A way to break bones? The weight of intuitiveness

D. Vettese[1,2] (Corresponding author), T. Stavrova[1], A. Borel[1,3], J. Marin[1], M.-H. Moncel[1], M. Arzarello[2], C. Daujeard[1].

[1]Histoire Naturelle de l'Homme Préhistorique (HNHP, UMR 7194), Sorbonne Universités, Muséum national d'Histoire naturelle (MNHN), Homme et Environnement, CNRS, Institut de Paléontologie Humaine, Paris, France
[2]Università degli Studi di Ferrara, Dipartimento degli Studi Umanistici, Sezione di Scienze Preistoriche e Antropologiche, Corso Ercole I d'Este 32, 44121 Ferrara, Italy
[3]Institute of Archaeological Sciences, Eötvös Loránd University, Múzeum krt. 4/b, 1088 Budapest, Hungary.

Delphine Vettese (corresponding author): delphine.vettese@mnhn.fr, https://orcid.org/0000-0001-6441-6054

Trajanka Stavrova: stavrovatrajanka@gmail.com

Antony Borel: antony.borel@mnhn.fr

Juan Marin: juan.marin.hernando@gmail.com, http://orcid.org/0000-0002-5698-602X

Marie-Hélène Moncel: marie-helene.moncel@mnhn.fr

Marta Arzarello: marta.arzarello@unife.it

Camille Daujeard: camille.daujeard@mnhn.fr, https://orcid.org/0000-0001-7489-8691

Abstract:

During the Middle Paleolithic period, bone marrow extraction was an essential source of fat
nutrients for hunter-gatherers especially throughout cold and dry seasons. This is attested by the
recurrent findings of percussion marks in osteological material from anthropized archaeological levels.
Among them some showed indicators that the marrow extraction process was part of a butchery cultural
practice, meaning that the inflicted fracturing gestures and techniques were recurrent, standardized and
counter-intuitive i.e. culturally influenced. In order to assess the weight of the counter-intuitive factor
in the percussion mark pattern distribution, we carried out an experiment that by contrast focuses on the
intuitive approach of fracturing bones to extract marrow, involving individual without experience in this
activity.

We wanted to evaluate the influence of bone morphology and the individuals' behaviour on the
distribution of percussion marks. Twelve experimenters broke 120 limb bones, a series of 10 bones per
individual. During the experiment, information concerning the fracture of the bones as well as individual
behaviour was collected and was subsequently compared to data from the laboratory study of the
remains. Then, we applied an innovative GIS (Geographic Information System) method to analyze the
distribution of percussion marks to highlight recurrent patterns. Results show that in spite of all the
variables there is a high similarity in the distribution of percussion marks which we consider as intuitive
patterns. The factor influenced the distribution for the humerus, radius-ulna and tibia series is the bone
morphology, while for the femur series individual behaviour seems to have more weight in the
distribution. To go further in the subject we need to compare the intuitive models with the distributions
of percussion marks registered in fossil assemblages. Thus, it would be possible to propose new
hypotheses on butchering practices based on the results presented in this work.

Keywords: GIS; bone breakage; experimental archaeology; intuitiveness; percussion marks; spatial
analyses

 Introduction

Lipids are essential nutrients for the human organism. During the Palaeolithic, fat was even
more important, due to its role in the gluconeogenesis metabolism, synthesizing glucose from non-
carbohydrate precursors (Costamagno and Rigaud 2014; Holt et al. 2018; Speth 2010). At that time,
various flora and fauna provided lipid resources but their accessibility was largely influenced by
environmental contexts. Lipids were found in oilseeds, such as hazelnuts. However, cold and dry periods
were characterized by a low vegetal biomass during which animal resources represented the most
important food resources (Binford 1981; Fiorenza et al. 2015; O'Connell and Marshall 1989; Speth
2010). The red marrow present in bone epiphyses and yellow marrow from the diaphysis were both
important resources of fat from animal carcasses. Marrow was even the most widely available fat
resource during the winter (Abe 2005; Binford 1978; Brugal and Defleur 1989; Costamagno and David
2009; Hardy et al. 2006; Kuntz et al. 2016; Speth and Spielman 1983). Indeed, the recovery of yellow
marrow was quasi-systematic among Palaeolithic hunter-gatherers, and particularly for Neanderthals
(Noe-Nygaard 1977; Patou-Mathis 1985; Speth 1989; Speth et al. 2012). Prehistorians noted widespread
evidence of marrow extraction through long bone breakage at Middle Palaeolithic sites (e.g.,: Blasco
and Fernández Peris 2012; Dart 1971; Daujeard et al. 2017; Gabucio et al. 2014; Marín et al. 2017;
Martin 1907; Romandini et al. 2014; Valensi et al. 2013; White 1992).

The traces left on bones by marrow recovery are mainly percussion marks. For several decades,
percussion marks and long bone breakage methods have been extensively studied (Boulestin 1999;
Capaldo and Blumenschine 1994; Fisher 1995; Gifford-Gonzalez 2018; Patou-Mathis 1985; White
1992). Since the beginning of the twentieth century and up until now, archaeological experiments on
long bone breakage have been carried out to characterize percussion marks and their location (e.g.,:
Blasco et al. 2014; Blumenschine and Selvaggio 1988; Brugal and Defleur 1989; Galán et al. 2009;
Martin 1910; Moclán and Domínguez-Rodrigo 2018; Noe-Nygaard 1977; Patou-Mathis 1985; Pickering
and Egeland 2006). Fracture patterns have also been extensively studied on different states of bone
preservation: fresh, dry, weathered, fossilized but also heated, frozen, boiled (Alhaique 1997; Karr and
Outram 2012; Lacroix 2011; Outram 2002; Villa and Mahieu 1991). The authors based their
experimental protocols on ethnographic works studying current populations of hunter-gatherers (e.g.,:
Abe 2005; Binford 1981; Costamagno and David 2009; Enloe 1993; Oliver 1993; Stiner et al. 2011;
Yravedra et al. 2017). More recently, the use of new methodological approaches (e.g., geometric
morphometrics, 3D modelling or GIS analyses of distribution along the diaphysis…) presents new
challenges for the experimental study of marrow recovery.

Some recent studies have focused on the distribution of percussion marks on long bones to
approach the subsistence behaviours of past hominins (Blasco et al. 2013; Masset et al. 2016; Moclán
and Domínguez-Rodrigo 2018; Vettese et al. 2017). One of these studies highlighted for the first time a
pattern of non-random bone breakage and interpreted it as a product of butchery traditions among

Neanderthals (Blasco et al. 2013). Through comparisons with an experiment performed by non-trained

individuals, the authors established that this systematic pattern differed from intuitive patterns. In

addition, Moclan and colleagues (2018) proved that the morphology of the skeletal element from which

marrow is extracted could influence the distribution of percussion marks.

86   We conducted a large-scale experiment to test how non-trained individuals recover marrow.

Indeed, in order to identify butchery traditions in Palaeolithic sites, it is necessary to differentiate know-

how from intuitiveness. The definition of a butchery tradition is a systematic and counterintuitive pattern

shared by a same group. The immediate apprehension of the non-trained butcher to break a bone is

influenced by numerous variables, including anatomical constraints (Blasco et al. 2013; Moclán and

Domínguez-Rodrigo 2018). Individuals who regularly break long bones acquire an empirical approach

and develop specific skills that enhance efficiency (Pickering and Egeland 2006). Hence, their skills

include habits and preferences gained by experience and/or group traditions. This know-how cannot be

assessed without differentiating between physical bone features and socio-cultural practices.

95   Thus, our aim is to experimentally test whether the inter-individual specificity of long bone

morphology may have some influence on the distribution of percussion marks. Based on (GIS) spatial

analysis of percussion marks on the bone surface, we also test the existence of a preferential pattern

regarding the intuitive breakage of long bones. Moreover, we intend to verify whether non-experienced

individuals develop their own method to improve efficiency, such as an auto-learning process. Finally,

it should be possible to grasp the influence of each bone structure by comparing the elements between

them and by assessing the performance of individuals on the same types of bones. This allows for the

comparison of behaviours and the influence of behaviour on the production of bone remains and the

marks recorded.

Material and Methods

Material

107   The studied sample includes 120 limb bones (humeri, radio-ulnas, femurs and tibias) from adult

cows at least 24 months old. A slaughterhouse supplied this experimental series, with 30 specimens of

each element, both left and right (Table 1). Professional butchers defleshed the carcasses. During the

process, they cut the metapodials with cutting pliers. Thus, these bones are absent from the present

experiment. After the reception of the bones, they were stored for less than a week in a fridge at 4 °C.

In addition, the elements broken by experimenters 11 and 12 were frozen for 40 days and thawed for

three days in the same fridge before the experiment (temperature: 4°C).

*Table 1: Data about the bone element broken by each individual (number of ID, element used, side). * These bones were frozen.*

Individual number	Element	Right	Left
1	Humerus	3	7
2	Radio-ulna	6	4
3	Femur	6	4
4	Tibia	2	8
5	Femur	4	6
6	Tibia	4	6
7	Humerus	5	5
8	Femur	6	4
9	Radio-ulna	6	4
10	Tibia	3	7
11	Humerus*	4	6
12	Radio-ulna*	3	7

We performed this experiment in a designated area in the property of the Museum national
d'Histoire naturelle (Paris). The experimental series involved 12 individuals (Table 2) without any
experience in bone breakage: eight men (mean age =34 years old; SD = 11.1 years) and four women
(mean age = 31 years old; SD = 7.1 years). Five individuals had theoretical knowledge of long bone
anatomy and one (individual number 1) was used to flaking lithic tools. Each experimenter broke a
series of 10 long bones of the same element. The number of tests is defined by the order in which each
bone is broken one after the other in a series of 10. To avoid selection biases, the bones were stacked
in a disordered pile when they were presented to the experimenters. Experimenters were isolated from
each other so they could not observe how the others broke the bones. No demonstration was performed.
Before the experiment, they only received one instruction: break the bone to extract the highest quantity
and quality of yellow marrow suitable for consumption. The breakage activity lasted for two to three
hours depending on the bone element and the individual. Experimenters had at their disposal a non-
retouched quartzite hammerstone weighing about 2 kg, a limestone anvil, a plate to deposit the marrow
and a wooden stick. The periosteum was not removed before breakage. Experimenters stopped once
they had extracted as much bone marrow as they could, and then collected it in the plate and weighed it
(Vettese 2020).

139 *Table 2: Data about the individuals for each series (W: Women; M: Men;,Stand. Dev.: Standard Deviation).*

Individual number	Age	Height (cm)	Weight (kg)	Sex	Used hand	Sport Practice	Bone knowledge	Broken bone element
1	29	167	53	W	Right	Yes	Yes	Humerus
2	40	174	82	M	Right	No	Yes	Radio-ulna
3	25	185	78	M	Right	Yes	Yes	Femur
4	30	186	80	M	Right	Yes	No	Tibia
5	51	160	59	W	Left	No	Yes	Femur
6	22	190	95	M	Left	Yes	No	Tibia
7	39	169	63	M	Right	Yes	No	Humerus
8	23	182	95	M	Right	No	Yes	Femur
9	23	167	58	W	Right	Yes	No	Radio-ulna
10	46	169	74	M	Right	Yes	No	Tibia
11	25	158	73	W	Right	Yes	No	Humerus
12	29	178	80	M	Right	Yes	No	Radio-ulna
Average	31.8	173.8	74.2					
Stand. dev.	9.8	10.4	13.7					
Maximum	51	190	95					
Minimum	22	158	53					

Our experiment involved individuals with no empirical knowledge of recovering marrow using
long bone breakage. Some were archaeologists or palaeontologists, the rest of the experimenters worked
in other fields with no relation to skeletal anatomy or bone tissue properties. The first category of
experimenters were familiar with bone morphology and structure as a study object only. Thus, the
approach to the long bones during marrow recovery was as intuitive as possible, independently of the
experimenters' age or sex. Besides, the experimenters broke a series of ten bones one after another. This
process allowed novice experimenters to self-assess and eventually learn from their mistakes.

After breakage of each element, all the fragmented remains were grouped together in a bag with
an identification code. The bones were boiled during 2 or 3 hours, to remove the grease and to soften
the flesh, and then placed in an oven at 40°C in a solution of water and Papain (papaya enzyme) for 48
hours. Then, they were soaked in a solution of water and sodium perborate for 24 hours, before being
air-dried. The material is currently kept in the Institut de Paléontologie Humaine (Paris).

Data acquisition

During the experiments, an observer recorded the following variables: the series number,
element laterality, position of the individual, the way the hammerstone was grasped, the position of the
bone, the use of the anvil and the number and location of blows. For these latter, we defined an area as
a long bone portion associated with a side (Figure 1). In order to evaluate the progression of individuals
during the experiment, from the first try to the end of the operation, we recorded task difficulty

evaluation and the number of blows. We asked the experimenters to auto-evaluate the difficulty encountered during marrow recovery on a scale from 1 to 5, (1 = very easy; 5= very hard), considering mainly the opening of the medullary cavity and marrow extraction. This auto-evaluation resulted in an empirical comparison between long bones. The aim of this auto-evaluation was to assess whether individuals progressed and developed an efficient method to extract marrow or if tiredness along with the long period of effort affected their capabilities. For quality control purposes, the experimenter and observer evaluated results on a scale from 1 to 5, 1 referring to contaminated marrow containing abundant splinters and 5 being very clean marrow.

Figure 1: Bone areas by portions and sides. Portion 1 (p1): proximal articular end; Portion 2 (p2): proximal diaphysis; Portion 3 (p3): medial diaphysis; Portion 4 (p4): distal diaphysis; Portion 5 (p5): distal articular end and Anterior (a), Posterior (p), Medial (m) and Lateral (l) side (adapted from Stavrova et al. 2019; https://doi.org/10.1371/journal.pone.0216733.s004).

The number of identified specimens (NISP), the number of undetermined specimens (NUSP) and the total number of specimens (NSP) were recorded. As the exact location of the percussion marks is necessary, only identified fragments for each bone element were included. Besides, throughout the treatment and the study of each bone, all the remains of one element were kept together, but the NUSP were splinters that we could not refit. Remains that could not be refitted were not considered in the following analysis. The width and length of fragments were measured using a digital caliper. The location of the percussion marks on the bone fragments was analysed according to the GIS protocol developed by Stavrova et al. (2019). The outline of the cortical surface of each identified fragment was

digitized using georeferenced photographic images representing the four-sided visualization of each
bone as a base.

Bone surface modifications were identified and recorded with the naked eye and 15-20x lens.
The centre of each percussion mark was recorded in ArcGIS with a point symbol, based on the
anatomical landmarks of each bone (e.g.,: foramen, crest)(Stavrova et al. 2019). In accordance with the
terminology of Vettese et al. (2020), five types of percussion marks were recorded: adhering flakes,
crushing marks, flakes, percussion notches, ovoid or triangular percussion pits and grooves. The number
of percussion marks (NPM) and the marks related to percussion (i.e., striations, microstriations and
pseudo-notches) were also noted. These last marks were not taken into account in the total number of
percussion marks. In addition, we excluded the flake because if we had taken into account the flake and
the notch, which were respectively positive and negative for the same percussion mark, we would have
counted the percussion point twice. In this analysis, percussion marks were divided into two groups: 1)
crushing marks, notches and adhering flakes (called "CNA") and 2) pits and grooves (called "Pits").
The "CNA" grouped the percussion marks related to cracking and fractures whereas "Pits" were only
produced by incipient percussion or the rebound effect.

Distances between percussion marks were calculated in ArcGIS and the "Optimized Hot Spot
analysis" tool was used to evaluate and highlight zones with high concentrations of percussion marks.
This tool identifies statistically significant spatial clusters of high concentrations (hot spots) and low
concentration of percussion marks (cold spots). The number of percussion marks recorded for each
series complies with the minimum of 30 features required to conduct the analysis (Stavrova et al. 2019).

The data collected during the experiment were compared to the data analysed in the laboratory
in order to understand which variables of the experimenters' behaviour could influence fragmentation
and the production of percussion marks. Spearman's rho was used to test the correlation between the
data collected during the experiment (i.e., number of blows and marrow weight) and the data recorded
on the bones after treatment (i.e., number of remains and number of percussion marks). Moreover, in
order to evaluate auto-learning and individual improvement, we tested the Spearman correlation of the
number of attempts according to the number of blows for each individual. We presented ρ of Spearman.
To highlight the evolution of the number of blows between the beginning and end of the experiment for
each of the individuals, we compared, on the one hand, the two averages of the first three and the last
three trials and, on the other hand, the first five and the last five trials using a non-parametric Mann-
Whitney U test. Indeed, the t-test hypothesis was never respected for our data (i.e., the normality and
equivalence of the variance of data). We propose an Efficiency Index calculated by dividing the mass
of marrow extracted from each bone by the number of blows. This Efficiency Index represents the
relationship between a form of expended energy and a form of recovered energy. The higher the
Efficiency Index, the more marrow was extracted with the least number of blows. To test a possible

evolution of the Efficiency Index between the three first and the last three tries and also between the

five first and the last five attempts, we compared the two averages using the Mann-Whitney U Test,

when assumptions for the t-test were not respected. We used the chi-squared test for the scale of auto-

assessment. As Cochran's rule was not respected, we used the Fisher exact test to check the independence

between the scale of marrow quality of the different elements. The correlation coefficients and the tests

were computed using R software (R Core Team 2019). We use an alpha of 0.05 for all the tests.

Results

Data recorded during the experiment

Bone and individual positions: different ways to break a bone

The position of the bone elements and the position of the individuals themselves varied during

the breaking process (Table 3 and SI 1). Five different positions chosen by the experimenters were

observed. The majority chose the squatting position (45%), followed by the kneeling position (43%).

The standing position was rarely selected. Only one individual broke bones in a seated cross-legged

position (individual n°2). A minority of the experimenters adjusted their positions during breakage (five

individuals: n°2, 4, 8, 9 and 11; 42%). Individual n°7 remained in the squatting position during the entire

experiment, but nonetheless tried the standing position once at the eighth try, without repeating it.

Individuals n°10 was the only one who did not adopt and maintain a position. By contrast, four

experimenters developed some kind of habit after a certain number of attempts. Individual n°3, after

three tries, chose the seated position and individual n°6 selected a kneeling position after two tries.

Individual n°5 developed a routine from the sixth attempt onwards. For each attempt, she started in a

standing position, then kneeled down, and she finished in a squatting position. Individual n°1 squatted

for all the attempts and half the time, she stood to hit the bone on the anvil. Only one individual tested

almost all the positions during the experiment (individual n°12). Nonetheless, after two tries, he

switched back to a squatting or kneeling position.

Regarding the choice of hand to grasp the hammerstone, only one individual used the left

hand, although two individuals were left-handed and one was ambidextrous. Therefore, the use of both

hands did not seem to be dependent on laterality. All the individuals breaking the radio-ulna used both

hands at least once to grasp the pebble, which was never the case for the individuals who broke the

tibias. The individuals who used both hands were both men and women.

We distinguished five ways to position the long bone in order to open its medullar (Table 3).

Ten of the individuals used the anvil at least once. Two experimenters (individuals n°8 and n°11) did

not use the anvil at all, but positioned the bones directly on the ground. One individual (individual n°7)

selected the batting technique. He hit the bone directly on the anvil with his hands grasping each

epiphysis in order to separate portion 1 from the diaphysis. Then, he laid the largest fragment on the anvil and pursued breakage using the pebble to recover as much marrow as possible. He began using this mixed technique after two tries and repeated it for the next attempts until the end of the series. He judged this mixed technique more efficient than only the hammerstone on anvil technique. One-third (33%) of the individuals always positioned their bones in the same way during the experiment. Therefore, the majority of the experimenters varied bone position during each try. We noticed that individuals experienced some difficulties in stabilizing the bone in order to hit it, especially for the radio-ulnas. Sometimes the bone slipped and the blow was less efficient. Some experimenters used an extra stone, their knees or the anvil to block the bone and keep it from slipping.

Table 3: Individual and bone position for each bone element, percentages were performed by 30 attempts by elements, and by the 120 attempts for the total.

	Squatting position	Standing position	Seated position	Two knees position	One knee position	Right grasp hand	Left grasp hand	Both grasp hand	On the anvil	One side on the anvil	Batting	On anvil and ground	On the ground
Femur	7 23.3%	4 13.3%	10 33.3%	13 43.3%	2 6.7%	20 66.7%	0 0%	12 40%	18 60%	0 0%	0 0%	1 3.3%	16 53.3%
3			10	3	2	10		2	10				
5	6	4						10	8			1	6
8	1			10		10							10
Humerus	20 66.7%	1 3.3%	3 10%	8 26.7%	0 0%	30 100%	0 0%	9 30%	20 66.7%	3 10%	9 30%	0 0%	10 33.3%
1	9	1	2			10			10				
7	10					10		8	10	3	9		
11	1		1	8		10		1					10
Radio-Ulna	16 53.3%	0 0%	11 36.7%	5 16.7%	3 10%	23 76.7%	0 0%	13 43.3%	16 53.3%	0 0%	0 0%	14 46.7%	10 33.3%
2			10			10		2	10				
9	10					3		10	1			5	10
12	6		1	5	3	10		1	5			9	
Tibia	20 66.7%	0 0%	0 0%	10 33.3%	6 20%	20 66.7%	10 33.3%	0 0%	26 86.7%	2 6.7%	0 0%	11 36.7%	0 0%
4	10					10			10	2			
6				10			10		10			1	
10	10				6	10			6			10	
Total	63 52.5%	5 4.2%	24 20%	36 30%	11 9.2%	93 77.5%	10 8.3%	34 28.3%	80 66.7%	5 4.2%	9 7.5%	26 21.7%	36 30.0%

The marrow quantity and quality

In order to extract marrow after breakage, most of the individuals used the wooden stick we provided, but some preferred selecting their own (SI 2). Only one individual (n°7) used the bone splinters created during breakage. Some also used their fingers. The radio-ulna was the long bone that yielded the smallest quantity of marrow; it represents 13% of the total collected marrow (Table 4). The quantity of marrow extracted from the femur is twice as high (26%). The individuals who broke the humeri and the tibias extracted a slightly higher quantity of marrow (around 30%). The total quantity of yellow marrow recovered represented 18 kg. We did not notice high variation among individuals, except for individual n°5 who recovered a significantly smaller quantity of marrow from the femurs than the other individuals did for the femur series (SI 3 4, 5, 6).

Element	N°indiv	MW (g)		NSP		NISP		NUSP		NB		NPM		Npit		NCNA	
Femur	3	1861	10.20%	259	9.50%	130	9.30%	129	9.80%	641	10.30%	263	5.30%	190	4.80%	73	7.40%
Femur	5	1030	5.60%	202	7.40%	137	9.80%	65	4.90%	496	7.90%	326	6.50%	282	7.10%	44	4.40%
Femur	8	1780	9.80%	222	8.20%	113	8.10%	109	8.20%	179	2.90%	221	4.40%	165	4.10%	56	5.70%
Femur	**Total**	**4671**	**25.60%**	**683**	**25.10%**	**380**	**27.10%**	**303**	**22.90%**	**1316**	**21.10%**	**810**	**16.20%**	**637**	**15.90%**	**173**	**17.50%**
Humerus	1	1710	9.40%	290	10.70%	109	7.80%	181	13.70%	573	9.20%	194	3.90%	72	1.80%	122	12.30%
Humerus	7	1737	9.50%	131	4.80%	77	5.50%	54	4.10%	244	3.90%	304	6.10%	217	5.40%	87	8.80%
Humerus	11	1941	10.60%	195	7.20%	84	6.00%	111	8.40%	377	6.00%	367	7.40%	301	7.50%	66	6.70%
Humerus	**Total**	**5388**	**29.50%**	**616**	**22.60%**	**270**	**19.30%**	**346**	**26.20%**	**1194**	**19.10%**	**865**	**17.30%**	**590**	**14.80%**	**275**	**27.80%**
Radio-ulna	2	799	4.40%	263	9.70%	119	8.50%	144	10.90%	808	12.90%	180	3.60%	49	1.20%	131	13.20%
Radio-ulna	9	775	4.20%	113	4.20%	80	5.70%	33	2.50%	873	14.00%	503	10.10%	428	10.70%	75	7.60%
Radio-ulna	12	811	4.40%	189	6.90%	73	5.20%	116	8.80%	822	13.20%	942	18.90%	894	22.40%	48	4.80%
Radio-ulna	**Total**	**2385**	**13.10%**	**565**	**20.80%**	**272**	**19.40%**	**293**	**22.20%**	**2503**	**40.10%**	**1625**	**32.60%**	**1371**	**34.30%**	**254**	**25.70%**
Tibia	4	1755	9.60%	439	16.10%	192	13.70%	247	18.70%	285	4.60%	280	5.60%	138	3.50%	144	14.50%
Tibia	6	1759	9.60%	175	6.40%	120	8.60%	55	4.20%	753	12.10%	738	14.80%	673	16.80%	65	6.60%
Tibia	10	2290	12.50%	244	9.00%	166	11.90%	78	5.90%	189	3.00%	669	13.40%	590	14.80%	79	8.00%
Tibia	**Total**	**5804**	**31.80%**	**858**	**31.50%**	**478**	**34.10%**	**380**	**28.70%**	**1227**	**19.70%**	**1687**	**33.80%**	**1401**	**35.00%**	**288**	**29.10%**
	Total	**18248**		**2722**		**1400**		**1322**		**6240**		**4987**		**3999**		**990**	

The evaluation of the quality of collected marrow shows a low proportion of bad or very bad assessments (10 %) (Table 5). In addition, most of those who made this judgment broke radio-ulnas. Most (2/3) of the recovered marrow was evaluated as good or very good to consume. None of the individuals who broke the humerus series evaluated the marrow as very consumable. However, more than a half of the individuals estimated the tibia marrow to be very good. The Fisher exact tests show that the scales of the tested elements were independent (df = 12, p<0.05).

We noted a significant negative correlation between the quantity of marrow and the number of blows. Indeed, the radio-ulna was the element that received the most blows during the experiment and where a smaller quantity of marrow was recovered. Concerning the radio-ulna, we observed a significant negative correlation between the NISP and the quantity of marrow collected. Besides, we noticed a significant positive correlation between the quantity of recovered marrow and both the NUSP and the number of CNA.

Felt scale	All		Femur		Humerus		Radio-ulna		Tibia	
Very easy	14	11.7%	4	13.3%	2	6.7%	1	3.3%	7	23.3%
Easy	53	44.2%	11	36.7%	21	70%	5	16.7%	16	53.3%
Moderate	28	23.3%	9	30%	4	13.3%	11	36.7%	4	13.3%
Hard	16	13.3%	4	13.3%	1	3.3%	8	26.7%	3	10%
Very hard	9	7.5%	2	6.7%	2	6.7%	5	16.7%	0	0%
Marrow quality scale	All		Femur		Humerus		Radio-ulna		Tibia	
Very bad	3	2.5%	0	0%	3	10%	0	0%	0	0.0%
Bad	9	7.5%	1	3.3%	0	0%	8	26.7%	0	0.0%
Neutral	29	24.2%	7	23%	15	50%	6	20%	1	3.3%
Good	55	45.8%	17	56.7%	12	40%	14	46.7%	12	40%
Very good	24	20%	5	16.7%	0	0%	2	6.7%	17	57%

The level of difficulty experienced during the experiment and the number of blows

In conclusion, the experimenters found the activity very easy 14 times (12%), easy 53 times
(44%), moderate 28 times (23%), difficult 16 times (13%) and very difficult only 9 times (13%).
individuals who broke the humeri did not encounter any difficulty in most cases (N=92; 77% easy or
very easy). The radio-ulna seems to be more difficult to break (N=52; 43% hard or very hard and N=44;
37% moderate). None of the individuals who broke a tibia estimated the task to be very difficult (Table
5). They did not find the task any easier or harder as they progressed. In other words, they did not find
the task harder at the beginning because they did not know how to go about it and then easier because
they found an efficient technique. Likewise, exhaustion from the activity did not influence whether
individuals experienced more difficulty at the end of the breakage series (SI 3, 4, 5). The difficulty
encountered according to the bone element varied significantly (Chi-squared test: df =12, X²=73.00362,
p<0.01).

The results of Spearman's correlation analysis between the number of attempts according to the
number of blows for each individual completed these observations. For five individuals (individuals n°1
and n°7 (humerus), individual n°3 (femur) and (individuals n°6 and n°9 (tibia)), the correlation analysis
shows a significant negative relationship (Spearman's rank correlation coefficient results respectively ρ
= -0.81; -0.77; -0.73; -0.93; -0.69). Regarding the other individual series, the results were not significant

(Spearman's rank correlation coefficient results respectively ρ: n°2 = -0.29; n°4 = 0.54; n°5 = -0.2; n°8

= 0.51; n°10 = 0.41; n°11 = -0.44; n°12 = 0.32).

The results of Spearman's correlation analysis between the number of attempts according to the

number of blows for each individual completed these observations. For five individuals (individuals n°1

and n°7 (humerus), individual n°3 (femur) and individuals n°6 and n°9 (tibia)), the correlation analysis

shows a significant negative relationship (Spearman's rank correlation coefficient results respectively ρ

= -0.81; -0.77; -0.73; -0.93; -0.69). Regarding the other individual series, the results were not significant

(Spearman's rank correlation coefficient results respectively ρ: n°2 = -0.29; n°4 = 0.54; n°5 = -0.2; n°8

= 0.51; n°10 = 0.41; n°11 = -0.44; n°12 = 0.32).

The proportion of the number of blows varied over the last attempts depending on individuals.

Experimenters n° 1, 3, 7, 9 reduced their number of hits by half, on average, n°6 by six between the

three first attempts and the last three (SI 3). Individual n°12 increased the average number of blows after

the three first tries. However, if we consider the same individual breaking the same element during the

whole experiment, we did not observe a significant difference between the five first and the last five

attempts. We found similar results between the three first and the last three, except for the humerus (p-

value = 0.0254) (SI 4).

The bi-plots showing the number of blows and marrow weight did not display any linearity in

the data regarding each element (SI 5). When we examine bone element type, the individuals who broke

the radio-ulnas reveal reduced Efficiency Index compared to the other elements (Figure 2; EFI all). On

the other hand, the experimenters who broke the tibias had a high average Efficient Index (>5). There

was marked variation between the individuals who broke the humeri and the femurs, especially for the

first three attempts (Figure 2; EFI 3). Between the first three and the last seven tries, the EFI of the

individuals who broke the tibias varied, like individual n°8. At the individual level, two-thirds of the

individuals increased their EFI between the first five and last five tries (n°3, 5, 1, 7, 11, 9, 6, 10).

Similarly, more than half the individuals augmented their EFI between the first and last three (n°3, 1, 7,

11, 2, 9, 6) (SI 6 and 7). However, regarding all the experimenters, the Wilcoxon signed rank test did

not show a significant difference between the first five and the last five tries or between the first three

and the last three tries (p-value > 0.05) (SI 8). We observed similar results regarding each element (p-

value >0.05).

Figure 2: Box plots of the Efficiency Index by Individual during the experiment regarding the third first tries, the last seventh tries and all tries.

Data recorded on the faunal assemblage obtained

Bone fragmentation

The experiments yielded 2,722 bone remains, among which 1,400 were refitted (Table 4). The NSP of the tibias (32%) is higher than for the other elements, which are almost identical (humerus: 25%, radio-ulnas: 23% and femurs: 21%). Likewise, the tibia NISP is the highest (34%) and the NISP of the humerus and radio-ulna are the lowest (19%). Nonetheless, we noticed that the average NUSP is around 25% for all the elements. We did not observe marked variability between individuals regarding the NISP, NUSP and the NSP SI 6). However, individual n°4 produced more numerous NUSP, which also affected the NSP. According to our application of the Spearman's rank correlation test, the NSP was positively correlated with both the NUSP and the NISP for all the elements (all elements: $\rho = 0.44$ and

$\rho = 0.76$; humerus: $\rho = 0.57$ and $\rho = 0.79$; radio-ulna: $\rho = 0.39$ and $\rho = 0.68$; femur: $\rho = 0.51$ and $\rho =$

0.8; tibia: $\rho = 0.51$ and $\rho = 0.8$). The NUSP and the NISP were also positively correlated between them

(all elements: $\rho = 0.91$; humerus: $\rho = 0.93$; radio-ulna: $\rho = 0.91$; femur: $\rho = 0.9$; tibia: $\rho = 0.9$) (SI 9).

Number of percussion marks

For one-third of the series, we recorded more percussion marks than the number of blows given

by the experimenters (Table 4). This anomaly was observed for one individual for each bone element.

Three of the individuals who produced a high number of percussion marks used an anvil, but this was

not the case for individual n°8. The tibia and radio-ulna series presented numerous percussion marks

(around 33%), corresponding to almost twice as many percussion marks as the femurs and humeri. We

recorded the most abundant pits and grooves on the tibia and the radio-ulna series. The other percussion

marks represented around 27% for all elements, with slightly less on the femurs (18%). Furthermore,

these marks display a lower standard deviation (SD = 33) than the pits (SD = 263) or all the percussion

marks combined (SD = 245). We observed an inter-individual difference, in particular for the first series

of the humeri, radio-ulnas and tibias. A different observer recorded these series.

According to our application of the Spearman's rank correlation test, we noted a significant

positive correlation between percussion marks with the number of pits and grooves on one hand, and

the number of crushing marks, adhering flakes and notches ("CNA"), on the other, for all the long bones

except the tibias ($\rho_{NPM/Npit}$: humerus = 0.95; radio-ulna = 0.96; femur = 0.95; tibia = 0.93; all elements =

0.93; $\rho_{NPM/NCAN}$: humerus = 0.35; radio-ulna = 0.7; femur = 0.35; tibia = -0.07; all elements =0.48). The

number of pits and grooves in the tibia series depended on the number of percussion marks, whereas the

other marks were independent. The number of percussion marks causing a fracture ("CNA") was not

significantly correlated with the other percussion marks (pits) ($\rho_{NCAN/Npit}$: humerus = 0.16; radio-ulna =

0.59; femur = 0.29; tibia = -0.32; all elements = 0.24). However, we observed a positive correlation for

the radio-ulna series and a negative one for the tibia series. The number of percussion marks producing

fractures ("CNA") was positively correlated with the NSP for the humerus series ($\rho = 0.34$). We

observed a significant positive correlation between the "CNA" percussion marks and the weight of

marrow and the NUSP for femurs and tibias (respectively, femurs: $\rho = 0.5$; $\rho = 0.38$ and tibias: $\rho = 0.62$;

$\rho = 0.92$). In addition, the number of pits and grooves on the tibias was correlated positively with the

NISP and the NSP (respectively: $\rho = 0.5$; $\rho = 0.38$). Finally, the radio-ulnas showed a positive correlation

between percussion marks causing breakage ("CNA") and the number of blows ($\rho = 0.39$) (SI 9).

The graph for the PCA analyses showed more differences between the tibia series on one hand

and the radio-ulna and humerus series, on the other, in particular on the second dimension (Figure 3).

These differences are mainly due to the NSP and the Efficiency Index. On the first dimension, radio-

ulnas and humeri are opposed by the number of blows and by the number of tries.

Figure 3: PCA regarding the number of remains (NSP), the number of percussion marks (NPM), the Efficiency Index and the number of attempts (Series Number) with ellipses at 0.95.

393
394

Percussion mark types and spatial distribution

We observed around 5,000 percussion marks on the identified fragments (Table 6). Pits were the most numerous (78%) and grooves were the least frequent marks (2.6%). Both triangular and ovoid pits were present, but the former were less numerous. We recorded more pits on the tibias and radio-ulnas than on the other bone types. Adhering flakes and notches represented 15% of all the percussion marks. Crushing marks were one of the least numerous traces (4.9 %), and were often located close to the articular portions. We recorded fewer crushing marks and more grooves on tibias compared to the other long bones. Conversely, we documented the most numerous crushing marks and adhering flakes on the humeri. We noted high variation in the number of pits and adhering flakes depending on the observers who documented the percussion marks. This difference in recording did not influence the number of other marks to the same degree. We also observed inter-individual variability, in particular for the radio-ulna series.

410 *Table 6: Type of percussion marks by individual and by element on NISP (Total PM: Percussion marks total).*

Element/ N°indiv	Percussion notch		Adhering flake		Triangular pit		Ovoid pit		Grooves		Crushing marks		Total PM
Femur	**85**	**10.5%**	**26**	**3.2%**	**41**	**5.1%**	**574**	**70.9%**	**22**	**2.7%**	**62**	**7.7%**	**810**
3	45	17.1%	10	3.8%	9	3.4%	173	65.8%	8	3.0%	18	6.8%	263
5	15	4.6%	5	1.5%	14	4.3%	257	78.8%	11	3.4%	24	7.4%	326
8	25	11.3%	11	5.0%	18	8.1%	144	65.2%	3	1.4%	20	9.0%	221
Humerus	**80**	**9.2%**	**93**	**10.8%**	**95**	**11.0%**	**462**	**53.4%**	**33**	**3.8%**	**102**	**11.8%**	**865**
1	39	20.1%	66	34.0%	8	4.1%	47	24.2%	17	8.8%	17	8.8%	194
7	21	6.9%	18	5.9%	44	14.5%	166	54.6%	7	2.3%	48	15.8%	304
11	20	5.4%	9	2.5%	43	11.7%	249	67.8%	9	2.5%	37	10.1%	367
Radio-ulna	**120**	**7.4%**	**77**	**4.7%**	**625**	**38.5%**	**717**	**44.1%**	**29**	**1.8%**	**57**	**3.5%**	**1625**
2	66	36.7%	39	21.7%	9	5.0%	33	18.3%	7	3.9%	26	14.4%	180
9	39	7.8%	17	3.4%	141	28.0%	275	54.7%	12	2.4%	19	3.8%	503
12	15	1.6%	21	2.2%	475	50.4%	409	43.4%	10	1.1%	12	1.3%	942
Tibia	**174**	**10.3%**	**86**	**5.1%**	**411**	**24.4%**	**943**	**55.9%**	**47**	**2.8%**	**24**	**1.4%**	**1687**
4	93	33.2%	42	15.0%	26	9.3%	93	33.2%	19	6.8%	5	1.8%	280
6	23	3.1%	31	4.2%	224	30.4%	435	58.9%	14	1.9%	11	1.5%	738
10	58	8.7%	13	1.9%	161	24.1%	415	62.0%	14	2.1%	8	1.2%	669
Total	**459**	**9.2%**	**282**	**5.7%**	**1172**	**23.5%**	**2696**	**54.1%**	**131**	**2.6%**	**245**	**4.9%**	**4987**

The crushing marks were often situated close to the articular portions (Figure 4, SI 10, 11, 12,
13). We documented some crushing marks on the medial diaphysis of the humeri and the radio-ulnas
and the tibias of individual n°4. Furthermore, we recorded most of the percussion marks on the
diaphysis; only the pits and grooves were located on the articular portions. It was difficult to observe a
concentrated area of percussion marks on the template on which we drew the marks. We merged all the
percussion marks derived from the 10 bone elements into a single series and performed an optimized
hot spot analysis to evaluate high concentration zones of percussion marks (Figure 5, 7, 9, 11). For each
long bone series, we tested all the percussion marks together; apart from percussion marks causing a
fracture ("CNA") and the others (pits and grooves) (Figure 6, 8, 10, 12).

Figure 4: Distribution of percussion marks of individual series merged along the humerus,the radio-ulnas, femurs and tibias. The type of percussion mark are red star: crushing marks, white triangle: pits and grooves and grey circle: notches and adhering flakes.

The analyses taking into consideration all the percussion marks generally highlighted cold spots on the articular portions, for all the studied long bones. Conversely, hot spots were generally highlighted on shaft portions. In particular, most of the time the ulna was a cold spot for all the series of radio-ulnas. Likewise, for almost all the series, the tibia crest was a cold spot. The majority of the cold spots were observed on proximal and distal articular portions with the exception of the proximal portion of the radius in the "CNA" analyses and on the tibia crest and ulnas (Figure 4, SI 10, 11, 12, 13).

The spatial analyses show that the majority of the high confidence hot spot zones for the humerus series were on the medial and lateral sides (Figure 5 A, B, D). Individual n°11 was an exception with hot spot areas on portion 4 of the posterior and anterior sides. The merged humerus series showed four high confidence hot spot zones on the medial and lateral sides. The hot spot zone on the medial side of portion 2 is the largest, and the one on the lateral side of portion 2 is the smallest. The last ones are

437   located on portion 4 (Figure 5 A, B, C, D). When we consider the two types of percussion marks, we

438   observe some differences in the location of the high confidence hot spot zone. The "CNA" were on the

439   lateral and medial sides of portion 2 , the pits and grooves were also on the medial side of portion 2, but

440   also on all sides of portion 4 (Figure 6).

Figure 5: Optimized Hot spot analysis of combined percussion marks from the humerus in each series: A- Individual n°1; B-
Individual n°7; C- Individual n°11; D- Merged humerus series (Ind.: Individual)

445
Figure 6: Optimized Hot spot analysis of combined percussion marks from the humerus in each series: A-E- Individual n°1;
B-F- Individual n°7; C-G- Individual n°11; D-H- Merged humerus series; A, B, C and D: "CNA" Percussion marks and E, F,
G and H: pits and grooves (Ind.: Individual)

For the radio-ulna series, we observed hot spot areas on the anterior and posterior sides for all
the individuals (Figure 7). Individual n°9 was the only one who frequently hit portion 3-4 on the medial
side. The analysis of the combined radius-ulna series highlighted a high confidence hot spot zone on the
posterior side of diaphysis portions 2 and 3. This result reflected the general tendencies observed for all
individuals. The analysis defined a reduced area, portion 2 on the anterior side, as a hot spot zone, which
was influenced by individual n°9. The results of the analysis showed that the high confidence hot spot
zone for the merged radio-ulna was on portion 3 of the posterior side, and on the anterior side portion
3-4 for "CNA" marks only (Figure 8 D).

Figure 7: Optimized Hot spot analysis of combined percussion marks from the radio-ulnas in each series: A- Individual n°2;
B- Individual n°9; C- Individual n°12; D- Merged radio-ulnas series (Ind.: Individual)

461
462 *Figure 8: Optimized Hot spot analysis of combined percussion marks from the radio-ulnas in each series: A-E- Individual n°2;*
463 *B-F- Individual n°9; C-G- Individual n°12; D-H- Merged radio-ulnas series; A, B, C and D: "CNA" Percussion marks and E,*
464 *F, G and H: pits and grooves (Ind.: Individual)*

The spatial analyses of the femur series show diversified distribution of the hot spot zones along the diaphysis, depending on the experimenter (Figure 9). For the three series, all sides were a hot spot area at least once, reflecting high inter-individual variability. However, the spatial analysis of the merged femur series shows three confident hot spot zones, quite similar to individual n°8 (Figure 9 C), on portions 2 and 4 on the anterior side, and on portions 3 and 4 on the medial side. High confidence hot spot zones for both percussion mark types are more numerous and dispersed on the merged femur than on the other bones (Figure 10 D, H).

Figure 9 Optimized Hot spot analysis of combined percussion marks from the femurs in each series: A- Individual n°3; B-Individual n°5; C- Individual n°8; D- Merged femurs series (Ind.: Individual)

Figure 10: Optimized Hot spot analysis of combined percussion marks from the femurs in each series: A-E- Individual n°3;
B-F- Individual n°5; C-G- Individual n°8; D-H- Merged femurs series; A, B, C and D: "CNA" Percussion marks and E, F, G
and H: pits and grooves (Ind.: Individual)

For the tibia series, we observed high confidence hot spot zones on anterior and posterior sides for each individual, in particular on portions 3 and 4 (Figure 11). The lateral side highlighted hot spot areas for individuals n°6 and n°10, especially on portion 2 (Figure 11 B, C). The merged tibia series reflects the dispersion of the high confidence hot spot zones observed for each series. The spatial analyses showed six hot spot zones, two of which very limited in size. However, the "CNA" area of high confidence hot spot was only on the posterior side of portions 3 and 4 (Figure 12 A, B, C, D). Zones with pits and grooves were numerous, distributed on the diaphysis of the lateral, anterior and posterior sides (Figure 12 E, F, G, H).

Figure 11: Optimized Hot spot analysis of combined percussion marks from the tibias in each series: A- Individual n°4; B- individual n°6; C- Individual n°10; D- Merged tibias series(Ind.: Individual)

494 *Figure 12: Optimized Hot spot analysis of combined percussion marks from the tibias in each series: A-E- Individual n°4; B-*
495 *F- Individual n°6; C-G- Individual n°10; D-H- Merged tibias series; A, B, C and D: "CNA" Percussion marks and E, F, G*
496 *and H: pits and grooves (Ind.: Individual)*

The most significant information highlighted by our experiments is related to the variability observed during the breakage process and that observed among the bone elements produced during experimentation. Indeed, some individual behavioural tendencies emerge during the breakage process, as well as patterns in the production and distribution of percussion marks. Our results also brought to light the influence of long bone morphology in an intuitiveness context.

How do individual gesture alternatives and bone elements influence breakage methods?

Some individuals could applied reduced force when they hit the bones, especially during the first attempts. Initially, experimenters tested the resistance of the bones and some were afraid of hurting themselves. In these cases, they hit the bones with less force. Likewise, at the end of the experiment, tiredness resulted in a reduction of the force applied. After the activity, most of the individuals had a shaking hand, and all the individuals had aches and pains the following days. This shows that marrow extraction requires a good physical condition and/or practice and/or good technical skill. During the activity, some individuals progressed or developed habits or a routine. However, they did not seem to become more efficient at optimally breaking bone to extract the maximum quantity of marrow. In addition, during the experiment, the individuals mentioned inter-element differences, whereby some bones were easier to break than others, irrespective of the experience acquired or the degree of exhaustion. Moreover, it is important to note that sometimes the hammerstone slipped on the cortical surface or slipped out of the experimenter's hand. This happened especially when the hammerstone and hands were covered in grease after several tries. When an individual broke several bones, he/she had to clean the tools or his/her hands to pursue marrow extraction.

One of the most significant results of our experiment was the variability of individual behaviour during marrow recovery. One individual adopted an unexpected posture: a cross-legged seated position. However, the observed variability only comprised five positions. In addition to these postural variations, we documented some non-linear behaviour during the experiment. Our experiment showed that non-trained individuals could change their position at different moments during both the marrow extraction from one bone and throughout the ten bone series. However, the majority of the individuals kept the same position or developed some kind of habit or routine in their practice. This routine concerned the individual and bone position and the way of grasping the bone and percussor. The auto-learning developed by experimenters to be as efficient as possible involved their position and their way of grasping the bone and percussor. The position influences the amplitude of the gestures, the force involved in breakage and the location of the blows. The standing position was only selected by women, and not by all of them. Sex did not influence the choice of posture or position of the bone. No link was found between the gender of the individual and how consistent they were in their posture choice or in

positioning the bone throughout the experiment. Likewise, sex did not seem to influence the way the percussor was grasped or the use of an anvil. These results could provide some evidence regarding the organization of butchery task distribution during the Palaeolithic. We noted that the posture adopted by individuals did not vary in relation to the skeletal element. However, the type of long bones seemed to affect practices and gestures, in particular for tibias and radio-ulnas. This observation could be related to the high number of blows applied to these two bone types by novice experimenters.

Although individuals proceeded differently depending on the elements, we noticed some divergences in our results, in particular between radio-ulnas and tibias. The experimenters who broke the radio-ulna series had a very low yield in relation to the number of blows. On the whole, the opposite is true for the tibia series, for which individuals had a high yield. Thus, individuals who broke radio-ulnas found the activity quite difficult and considered the marrow to be of relatively poor quality. On the other hand, the individual who broke tibias found the activity quite easy and the extracted marrow was considered to be of relatively good quality. Furthermore, experimenters used two hands at least once for breaking radio-ulnas, but never for tibias. The use of both hands to grasp the pebble could be an expression of the difficulty felt by the individuals breaking radio-ulnas. Regarding the NSP produced for both these elements, we recorded the highest number of blows and the lowest NSP for the radio-ulnas and the exact opposite for the tibias of our sample. This was why we observed a high number of pits and the lowest number of "CNA" percussion marks on radio-ulnas. Contrary to the tibia, the breakage of a radio-ulna of an animal such as a cow requires some mastery. Numerous studies have highlighted differences between the volumes of the medullar cavity of these two long bones. Tibias and femurs comprise large medullary cavities. On the other hand, medullar cavities are smaller for radio-ulnas and humeri and thus contain a lesser quantity of yellow marrow (Binford 1978, 1981; Jones and Metcalfe 1988). Therefore, the EFI (Efficiency Index) highlighted differences between the energy spent by counting the blows and the quantity of marrow extracted, in part because of the difference in volume between the radio-ulna and tibia series. However, medullary cavity volume alone cannot account for all the differences observed between bones. In terms of density, tibias and radiuses show the highest diaphysis density of all the long bones used in our experiment (Lam et al. 1999; Lyman 1994). However, they seem to react differently to blows. This difference was highlighted by the PCA, showing an opposition between the radio-ulna and the tibia series regarding the NSP, the Efficiency Index and the number of percussion marks.

What correlation can be made between blows and percussion marks?

Various researchers have stressed that the aim of mastering percussion breakage was to reduce the number of blows required to remove all the marrow from the medullary cavity (Capaldo and Blumenschine 1994; Galán et al. 2009; Moclán and Domínguez-Rodrigo 2018). For this reason, it was

expected that novice experimenters would average more blows than experts. Indeed, we noted a high number of blows, ranging from 6-279, with an average of 52, for the entire bone experiment. If we compare our experiment with previous works, we observe a huge difference between the average number and range of blows (Table 7). The average number of blows by individuals in our experiment was higher than the maximum in other studies. In particular, for their first tries, two individuals hit the bones more than 200 times. Besides, none of the experimenters stopped hitting the bones after the first fracture. The presence of the periosteum on the bones was not sufficient to explain the very high number of recorded blows. Indeed, during some experiments quoted in Table 7, the periosteum was not removed (Martin 1910; Pickering and Egeland 2006; Rovira Formento 2010). Novice experimenters tried to break the element further in order to recover the maximum amount of marrow. The weighing of the marrow after each attempt motivated some of them. At the individual scale, we observed some increase or decrease in the number of blows at the beginning and the end of the experiments, but overall, we did not highlight significant progress for all the individuals or a particular element. The only exception was the humerus, for which the number of blows decreased for all the individuals. The number of blows and the individuals' decision to pursue breakage after the first fracture can increase the NSP. Intensive breakage can reflect a high number of blows in order to access the yellow marrow. In addition, we observed the quasi-systematic conservation of the epiphyses in their entirety. Indeed, it is not necessary to break the articular portions for the extraction of yellow marrow only.

Table 7: Listing of different experiment of bone breakage for marrow extraction and comparing data (NME: Number of Minimum Element).

References	Species	Element	Range of number of blows	Mean of number of blows	NSP/NME=AverageNSP
(Rovira Formento 2010)	Horse	Femurs	1-27	8.8	-
(Galán et al. 2009)	Cow	Humerus Femurs Radio-ulna	- - -	- - -	145/22=6.5
(Pickering and Egeland 2006)	White deer	Radius Humerus	2-15 1-4	6 2	811/38=21.3 472/36=13.1
(Thiébaut et al. 2007)	Cow	Femurs	1-8	3.5	189/13=14.53
(De la Torre et al. 2013)	Cow	Limb bones	-	20	-
(Martin 1910)	Horse	Femurs Tibia	1-3 3	2 3	-

Interestingly, the number of blows is not correlated with the number of percussion marks. Comparing the two values, we noted that for two-thirds of the individual bone elements, the number of

blows is higher than the number of percussion marks and more surprisingly, for one-third of the bone
elements, the number of percussion marks is higher than the number of blows. For one individual per
bone element (humerus: n°7, radio-ulna: n°12, femur: n°8 and tibia: n°10), the number of blows was
inferior to the number of percussion marks.

In most cases (more than 95%), the number of pits and grooves or "CNA" marks was different
to the number of hits. In an archaeological context, it can thus be impossible to infer the number of
blows from the number of percussion marks. The use of an anvil or the hammerstone could explain the
differences recorded between the number of marks and blows. In their experiment, (Pickering and
Egeland 2006) also observed a majority of pits among the recorded percussion marks. They concluded
that most of these pits were produced by the counterblow of the anvil. In our experiment, pits were
generally more numerous than the other marks (67 %). Thus, pits and groves result from the rebound
effect caused by the hammerstone or by the anvil due to the counterblow. In this last case, the
counterblow is not a mark; but rather the result of the rebound effect of the bone on the anvil. The marks
due to counterblows include pits, microstriations, outer conchoidal scars or notches (e.g.,: Boulestin
1999; Capaldo and Blumenschine 1994; Gifford-Gonzalez 2018; Johnson 1985; Lyman 1994; Outram
2002), located on the opposite side to the blow. Due to the multiplicity of traces and the overlapping of
the traces produced by the hammerstone or when the long bone is batted against a hard surface, it is
difficult to discriminate blow marks from counterblow marks. In addition, we recorded some triangular
pits and grooves even though experimenters only used non-retouched pebbles. These marks,
characterized by straight and sharp edges, could be the product of the rebound effect from the anvil.
Therefore, the use of an anvil could create supplementary marks during the blow. Two individuals (n°8
and 11) did not use the anvil at all, but simply laid the bones on the ground. Nonetheless, some triangular
pits and grooves were identified (around 10 %). However, we sometimes recorded more marks than
blows. Thus, the counterblow is not the sole explanation for the difference between the number of blows
and the number of percussion marks. It is possible that the shape of the hammerstone, the impact with
the ground or the collision of bones between them during the activity created several pits. More than
one-quarter of the recorded pits were very small and superficial, with a diameter of 10 mm or less (28%).
In an archaeological context, the different taphonomical processes, such as trampling, for example
(Domínguez-Rodrigo et al. 2010), can erase these pits or produce others. For the remaining two-thirds
of the broken elements, we recorded a lower number of percussion marks than blows. In our protocol,
the periosteum was never removed. The presence of the periosteum, which absorbed some of the force
of the blows, could partly explain this difference.

Percussion marks distribution
The areas where the least percussion marks were observed, apart from the articular portions,
were the anterior crest of the tibia and the diaphysis of the ulna. The distribution of percussion marks
showed that novice experimenters avoided shape constraints, such as bone protuberances. They seemed

to prefer relatively flat areas. Thus, the spatial analyses for tibias and radio-ulnas highlighted significant

hot spot zones for the merged series, where bone shape was the flattest. Furthermore, for the "CNA"

marks, a single hot spot area was most impacted on the tibias. This area was located on the posterior

side of the distal diaphysis. It was one of the flattest and smoothest zones of the tibias with no muscular

lines. Moreover, the distal diaphysis is the least dense portion of the tibia (index density = 44) (Lyman

1994). Regarding the "CNA" marks, two specific high confidence hot spot zones were identified on the

radio-ulnas. They were located on relatively flat and easily accessible areas to blows. The first one is

situated on the anterior side of the medial shaft, which is relatively flat considering the globally very

arched shape of the radio-ulna. The radio-ulna section in this area is flattened compared with portions 3

and 4, which present a semi-lunar crescent shape. When the bone was placed on a small anvil, one of

the best ways to stabilize the radio-ulna was to place the medial shaft on the posterior side. Thus, the

anterior side was exposed and very accessible. The other hot spot area was on the exact opposite side

because, on the ground, it was easier to place the bone on the anterior side to avoid interference from

the ulna. In addition, when the radio-ulna was positioned on the anterior side, this area was located at

the inflexion point of the bone curve and was the most accessible to hit. Furthermore, these areas were

located on portions 2 and 3, which are the densest part of the radius (marrow index = 0.56 & 0.62).

Unlike zeugopods, the diaphyseal section of humeri and femurs is rather cylindrical. Their

diaphyses are relatively straight in comparison to the radio-ulna. The articular portion shows important

differences in terms of stability. The humerus is more stable on its lateral and medial sides whereas the

femur is relatively stable on all sides. Indeed, the very rounded shape of the humerus head causes

instability when placed on its posterior side. This is why the distribution of percussion marks,

highlighted by the hot spot analyses, is concentrated on the lateral and medial sides. Pit and groove

marks are generally distributed on the distal part of the bone. This part of the humerus diaphysis is

denser than the others (index density = 0.48). Conversely, the hot spot area of "CNA" marks is only

located on the proximal shaft portion, in the least dense part of the bone (index density = 0.25). These

"CNA" marks represent one of the most the numerous hot spot zones. In addition, these hot spot areas

were similar for the three merged spatial analyses of the femurs (all percussion marks, "CNA" marks

and pits and groove marks). All the diaphysis portions were hot spot zones at least once, independently

of density variability.

Thus, percussion mark location is influenced by the morphological constraints of bones, in

particular for the radio-ulna, the tibia and the humerus, in an intuitive context. Individuals were faced

with bone marrow extraction for the first time and were highly influenced by shape constraints. These

constraints were so strong that individuals continued hitting these zones after many attempts, as bone

stability and the accessibility of these areas facilitated breakage and therefore marrow recovery.

Nonetheless, the absence of morphological constraints seems to highlight novice choices. Therefore,

based on the results of our analyses, in an intuitive context, femurs seem to be the bones for which

individual variability was highest.

Archaeological, experimental and ethnographical data comparisons: some convergences

Binford (1981) described the practices and gestures of the Nunamiut in order to infer past

hunter-gatherer behaviour from archaeological assemblages. His observations describe systematic

breakage, hitting the same bone portions each time. Moreover, in the same way as experiments,

ethnographical studies highlight wide-ranging variables, which are hardly accessible in archaeological

contexts.

We highlighted the existence of a preferential pattern for the intuitive breakage of long bones,

using the spatial analysis of the location of percussion marks on radio-ulnas, tibias and humeri.

Archaeological and ethnographical studies focusing on the distribution of percussion marks along long

bone diaphyses have revealed the existence of preferentially impacted areas (Binford 1981; Blasco et

al. 2013; Masset et al. 2016; Moclán and Domínguez-Rodrigo 2018; Vettese et al. 2017). The

comparison between these results and ours underlines the influence of bone shape on some systematic

practices of extant and past hunter-gatherers (Figure 13). For radio-ulnas, archaeologists record

percussion marks on the anterior side, in particular in the faunal assemblages of level XVIIc of Bolomor

Cave (Blasco et al. 2013) and at the Abri du Maras layer 4.1 (Vettese et al. 2017), where some tendencies

were observed (Middle Palaeolithic), Combe Saunière and Cuzoul de Vers (Castel 1999) and the "upper

unit" of Saint-Germain-la-Rivière (Costamagno 1999; Masset et al. 2016) and the Rond du Barry (Upper

Palaeolithic), and also among the extant Nunamiut (Binford 1981). The areas highlighted by hot spot

analyses were similar to the areas of percussion mark concentrations in these comparative examples.

These faunal assemblages represent practically all the Palaeolithic sites for which percussion mark

distribution has been studied. The aforementioned faunal assemblages are dominated by different

species: reindeer, saiga antelope, ibex, red deer or larger-sized ungulates, like horse or bison, and in our

experiment, cow. This comparison suggests that the high shape constraints of the radio-ulnas strongly

influenced the distribution of percussion marks on whole long bone remains in an archaeological level

independently of ungulate size. Furthermore, for ungulates, the fact that the radius is merged with the

ulna constrains the morphological shape of the bones. However, level IV of Bolomor Cave and the

"lower unit" of Saint-Germain-la-Rivière showed different patterns on the lateral side for the former site

and both lateral and medial sides for the latter (Figure 13).

For the humerus, the distribution of percussion marks was quite similar to our observations.

Almost all the sites presented in the work of Masset et al. (2016) reveal a pattern on the medial and/or

lateral sides, with the sole exception of Bolomor Cave level IV. Compared to the other sites, we noted

variability in the exact area impacted by percussion marks, but we observed a preference for portion 4

in the majority of the assemblages. A minority of the studied tibia series showed a systematic pattern or

strong tendencies. The posterior sides displayed the most areas of percussion mark concentration.

Finally, most of the femur series from the archaeological sites studied do not seem to show systematic

distribution, with the exception of level VI of Bolomor, which indicates two preferentially impacted

zones, and the Abri du Maras, where one area was more impacted than the other. We observe similarities

with the Nunamiut for the anterior side.

Figure 13: Location of percussion marks (grey areas) for the humerus, radio-ulna, femur, and tibia for sites from the Middle
Paleolithic: Bolomor cave (ungulates: Blasco et al., 2013b.) and Abri du Maras (middle sized-ungulates: Vettese et al. 2017);
the Upper Paleolithic: Combe Saunière (reindeer, Solutrean: Castel, 1999), Cuzoul de Vers (reindeer, Badegoulian: Castel,
1999), Saint-Germain-la-Rivière, Lower Unit (saiga antelope, Lower Magdalenian: Costamagno, 1999) and Upper Unit (saiga
antelope: Masset et al. 2016), Rond du Barry (ibex, Upper Magdalenian: Costamagno, 1999), for these series the data
regarding the femurs were missing and most of the tibias did not present higher tendencies; Current: Nunamiut people of
Alaska (reindeer: Binford, 1981 and the results of our experiment regarding the CNA). Adapted according to Masset el al.
2016.

Conclusion

For the purposes of testing intuitiveness in bone breakage, we experimentally tested the

influence of individual behaviour and long bone morphology on percussion mark location. We observed

a high variability of gestures during the breakage process. Most of the individuals developed their own

routine. This routine included several positions and different techniques, involving variability in the

applied force and different bone element responses. Nevertheless, our results highlight similarities in

percussion mark distribution despite the specificity of each experimenter, for most of the long bones

studied here. Indeed, one of the most important results is the identification of specific hot spot areas,

regardless of the variability of experimenters' behaviour during the breakage process. These similarities

can be considered as intuitive patterns.

The location of percussion marks is influenced by numerous variables depending on: the
experimenters' behaviour and skills, the techniques employed, the location of the blows, the morphology
of the skeletal element and bone position during breakage. Nevertheless, our results highlighted the
predominance of two main factors in an intuitive context: long bone morphology for the humerus, radio-
ulna and tibia and experimenters' behaviour for the femur.

From a different perspective, we have shown that bone response can be very different in an
intuitive context, notably for the radio-ulna and the tibia. These differences can be observed at almost
all the different levels of the analysed data: experimenter behaviour or perception, marrow quality and
quantity, number of blows and remains produced and type of percussion marks recorded. These results
highlight the diametrically opposed differences between tibias and radio-ulnas in an intuitive context
Besides, to go further and specify our results, more experiment will be required with various cooking
methods, like roasting or boiling bones. Furthermore, the effect of prior periosteum removal will be
tested regarding the percussion marks formation or their number, and also, the breakage efficiency.

Our results shed new light on the importance of gestures in the breakage process. It could prove
interesting to model these results and evaluate their amplitude and the force involved. Numerous studies
focus on the force required to break long bones but generally concentrate on pressure methods.
Consequently, data pertaining to the percussion process are lacking. In addition to measuring the force
applied by the blow, it is essential to evaluate the force of the counter-blow caused by different materials,
such as stone, bone or wood. Moreover, it may be helpful to integrate our results regarding the whole
'chaine opératoire' of the bone breakage. Indeed, breakage function could be directed at grease
rendering, bone meal, bone fuel or craft activity instead of marrow recover alone.

Percussion marks are very instructive in an archaeological context to access past subsistence
behaviour. Our results show that percussion traces on fossil bones are only the tip of the iceberg. They
only reveal a small window into past bone marrow extraction practices. This experiment highlighted the
variability of experimenters' behaviour and the different bone responses during fracturing in order to
propose new hypotheses on past butchering practices based on the comparison of the different data
analysed in this work.

To conclude, the influence of bone morphology on the distribution of percussion marks in the
archaeological record should be taken into account before inferring cultural traditions. It is essential to
compare the intuitive patterns highlighted by spatial analysis with the patterns identified in
archaeological levels. It could also prove interesting in archaeological contexts to consider carefully
percussion mark distribution on the femur as this could be the most instructive bone element, in terms
of marrow recovery methods and group specificity.

Acknowledgements

We express our gratitude to CHARAL S.A.S. who kindly provided all the bones used in the preparation

of this experiment. We are grateful for the funding of the Fondation Nestlé France. We thank the

Museum national d'Histoire naturelle to permit the experiment and Philippe Meritte for his help. Many

thanks to Eric Pellé and Zoé Thalaud of the Service de Préparation Ostéologique et Taxidermique

(SPOT) from the Muséum national d'Histoire naturelle in Paris, for the osteological preparation of all

the long bones. We also wish to thank Laetitia Demay for her precious advice during proofreading and

her valuable help during the experiment. Laurent Crépin and colleagues provided helping logistical

advice and assistance during the experiment. We are deeply grateful to the volunteers for their selfless

efforts in helping with our experiment. We also thank Lou Albessard, Jeremy Duveau and Guilhem

Mauran for their precious advice. L. Byrne, an official translator and native English speaker, edited the

English manuscript. Finally, we thank A. Outram, T. O'Connor and the anonym reviewer for their

critical reviews that served to improve the paper.

Financial Disclosure

This project was supported by the Fondation Nestlé France (SJ 671-16) (https://fondation.nestle.fr/); the

Centre d'Information des Viandes – Viande, sciences et société (SJ 334-17); the Muséum national

d'Histoire naturelle.

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content

of this article. The authors reported no potential conflict of interest.

Marta Arzarello and Camille Daujeard are one of the *PCI Archaeology* recommenders

References

Abe, Y. (2005). *Hunting and Butchery Patterns of the Evenki in Northern Transbaikalia, Russia*. Stony brook University.

Alhaique, F. (1997). Do Patterns Of Bone Breakage Differ Between Cooked And Uncooked Bones ? An Experimental Approach. *Anthropozoologica*, *25–26*, 49–56.

Binford, L. R. (1978). *Nunamiut: Ethnoarchaeology* (Academic P.). New-York, Chicago, London.

Binford, L. R. (1981). *Bones: Ancient Men and Modern Myths.* (Academic P.). New-York.

Blasco, R., & Fernández Peris, J. (2012). Small and large game: Human use of diverse faunal resources at Level IV of Bolomor Cave (Valencia, Spain). *Comptes Rendus Palevol*, *11*(4), 265–282. doi:10.1016/j.crpv.2012.01.003

Blasco, R., Rosell, J., Domínguez-Rodrigo, M., Lozano, S., Pastó, I., Riba, D., et al. (2013). Learning by Heart: Cultural Patterns in the Faunal Processing Sequence during the Middle Pleistocene. *PLoS ONE*, *8*(2), e55863. doi:10.1371/journal.pone.0055863

Blasco, R., Domínguez-Rodrigo, M., Arilla, M., Camarós, E., & Rosell, J. (2014). Breaking Bones to Obtain Marrow: A Comparative Study between Percussion by Batting Bone on an Anvil and Hammerstone Percussion. *Archaeometry*, *56*(6), 1085–1104. doi:10.1111/arcm.12084

Blumenschine, R. J., & Selvaggio, M. M. (1988). Percussion marks on bone surfaces as a new diagnostic of hominid behaviour. *Nature*, *333*(23 June 1988), 763–765.

Boulestin, B. (1999). *Approche taphonomique des restes humains*. (Archaeopress, Ed.)*Bar International Series* (Vol. 776). Oxford: Publishers of British Archaeological Reports.

Brugal, J. P., & Defleur, A. (1989). Approche expérimentale de la fracturation des os des membres de grands mammifère. *Artefacts*, *7*, 14–20.

Capaldo, S. D., & Blumenschine, R. J. (1994). A Quantitative Diagnosis of Notches Made by Hammerstone Percussion and Carnivore Gnawing on Bovid Long Bones. *American Antiquity*, *59*(4), 724–748.

Castel, J.-C. (1999). *Comportements De Subsistance Au Solutreen Et Au Badegoulien D'Apres Les Faunes De Combe Sauniere (Dordogne) Et Du Cuzoul De Vers (Lot)*. Université Bordeaux I.

Costamagno, S. (1999). *Stratégies de Chasse et Fonction des Sites au Magdalénien dans le Sud de la France*. Unniversité Bordeaux I.

Costamagno, S., & David, F. (2009). Comparaison des pratiques bouchères et culinaires de différents groupes sibériens vivant de la renniculture. *Archaeofauna*, *18*, 9–25.

Costamagno, S., & Rigaud, J.-P. (2014). L' exploitation de la graisse au Paléolithique. In S. Costamagno (Ed.), *Histoire de l'alimentationn humaine : entre choix et contraintes* (Edition él.). Paris.

Dart, R. A. (1971). On the Osteodontokeratic Culture of the Australopithecinae Author (s): Raymond A . Dart and Donald L . Wolberg Published by : The University of Chicago Press on behalf of Wenner-Gren Foundation for Anthropological Research Stable URL : http://www.jstor. *Current Anthropology*, *12*(2), 233–236.

Daujeard, C., Vettese, D., Britton, K., Béarez, P., Boulbes, N., Crégut-Bonnoure, E., et al. (2017). Neanderthal selective hunting of reindeer? The case study of Abri du Maras (south-eastern France). *Archaeological and Anthropological Sciences*, 1–27. doi:10.1007/s12520-017-0580-8

De la Torre, I., Benito-calvo, A., Arroyo, A., Zupancich, A., & Proffitt, T. (2013). Experimental protocols for the study of battered stone anvils from Olduvai Gorge. *Journal of Archaeological Science*, *40*, 313–332. doi:10.1016/j.jas.2012.08.007

Domínguez-Rodrigo, M., Pickering, T. R., & Bunn, H. T. (2010). Configurational approach to
identifying the earliest hominin butchers. *Proceedings of the National Academy of Sciences*, (13
September 2010), 1–6. doi:10.1073/pnas.1013711107

Enloe, J. G. (1993). Ethnoarchaeology of marrow cracking: implications for the recognition of
prehistoric subsistence organization. In J. Hudson (Ed.), *From bones to behavior:*
ethnoarchaeological and experimental contributions to the interpretation of faunal remains (pp.
82–100). Carbondale: Southern Illinois University.

Fiorenza, L., Benazzi, S., Henry, A. G., Salazar-Garc??a, D. C., Blasco, R., Picin, A., et al. (2015). To
meat or not to meat? New perspectives on neanderthal ecology. *American Journal of Physical*
Anthropology, *156*(S59), 43–71. doi:10.1002/ajpa.22659

Fisher, J. W. (1995). Bone surface modifications in zooarchaeology. *Journal of Archaeological Method*
and Theory, *2*(1), 7–68. doi:10.1007/BF02228434

Gabucio, M. J., Cáceres, I., Rosell, J., Saladié, P., & Vallverdú, J. (2014). From small bone fragments
to Neanderthal activity areas: The case of Level O of the Abric Romaní (Capellades, Barcelona,
Spain). *Quaternary International*, *330*, 36–51. doi:10.1016/j.quaint.2013.12.015

Galán, B., Rodríguez, M., de Juana, S., & Domínguez-Rodrigo, M. (2009). A new experimental study
on percussion marks and notches and their bearing on the interpretation of hammerstone-broken
faunal   assemblages.   *Journal   of   Archaeological   Science*,   *36*,   776–784.
doi:10.1016/j.jas.2008.11.003

Gifford-Gonzalez, D. (2018). *An Introduction to Zooarchaeology*. Cham: Springer International
Publishing. doi:10.1007/978-3-319-65682-3

Hardy, K., Brand-miller, J., Brown, K. D., Thomas, M. G., & Copeland, L. (2006). The Importance of
Dietary Carbohydrate in Human Evolution. *Quaterly review of biology*, *90 (3)*, 251–268.

Holt, S. H., Eaton, S. B., Miller, J. B., Cordain, L., Speth, J. D., & Mann, N. (2018). Plant-animal
subsistence ratios and macronutrient energy estimations in worldwide hunter-gatherer diets. *The*
American Journal of Clinical Nutrition, *71*(3), 682–692. doi:10.1093/ajcn/71.3.682

Johnson, E. (1985). Current Developments in Bone Technology. In *Advances in Archaeological Method*
and Theory (Vol. 10, pp. 157–235). Elsevier. doi:10.1016/B978-0-12-003108-5.50010-5

Jones, K. T., & Metcalfe, D. (1988). Bare bones archaeology: Bone marrow indices and efficiency.
Journal of Archaeological Science, *15*(January), 415–423. doi:10.1016/0305-4403(88)90039-8

Karr, L. P., & Outram, A. K. (2012). Tracking changes in bone fracture morphology over time :
environment , taphonomy , and the archaeological record. *Journal of Archaeological Science*,
39(2), 555–559. doi:10.1016/j.jas.2011.10.016

Kuntz, D., Costamagno, S., Feyfant, L., & Martin, F. (2016). The exploitation of ungulates in the
Magdalenian in the Entre-Deux-Mers (Gironde, France). *Quaternary International*, *414*, 135–158.
doi:10.1016/j.quaint.2015.12.079

Lacroix, M. N. (2011). *A Study of the Impact of Weathering Upon the Minimal Force Required To*
Fracture Bone. Boston.

Lam, Y. M., Chen, X., & Pearson, O. M. (1999). Intertaxonomic Variability in Patterns of Bone Density
and the Differential Representation of Bovid , Cervid , and Equid Elements in the Archaeological
Record. *Society for American Archaeology*, *64*(2), 343–362. doi:10.2307/2694204

Lyman, R. L. (1994). *Vertebrate Taphonomy*. Cambridge: Cambridge University Press.

Marín, J., Saladié, P., Rodríguez-Hidalgo, A., & Carbonell, E. (2017). *Neanderthal hunting strategies*
inferred from mortality profiles within the Abric Romaní sequence. *Plos One* (Vol. 12).
doi:10.1371/journal.pone.0186970

Martin, H. (1907). Présentation d'ossements utilisés de l'époque moustérienne. *Bulletin de la Société*
préhistorique de France, *4*(5), 269–277. doi:10.3406/bspf.1907.11559

Martin, H. (1910). La Percussion osseuse et les esquilles qui en dérivent . Expérimentation. *Bulletin de*
la Societe Prehistorique Francaise, *5*(5), 299–304.

Masset, C., Costamagno, S., Cochard, D., & Laroulandie, V. (2016). La fracturation osseuse : du fait
technique à l'essai d'interprétation sociétale L'exemple de l'antilope saïga du gisement
magadelenien de Saint-Germain -la-Rivière (Gironde). *Bulletin de la Société préhistorique*
française, *113*(4), 691–712.

Moclán, A., & Domínguez-Rodrigo, M. (2018). An experimental study of the patterned nature of
anthropogenic bone breakage and its impact on bone surface modification frequencies. *Journal of*
Archaeological Science, *96*(May), 1–13. doi:10.1016/j.jas.2018.05.007

Noe-Nygaard, N. (1977). Butchering and Marrow Fracturing as a Taphonomic Factor in Archaeological.
Paleontological Society, *3*(2), 218–237.

O'Connell, J. F., & Marshall, B. (1989). Analysis of kangaroo body part transport among the alyawara
of central Australia. *Journal of Archaeological Science*, *16*(4), 393–405. doi:10.1016/0305-
4403(89)90014-9

Oliver, J. S. (1993). Carcass processing by the Hadza: bone breakage from butchery to consumption. In
J. Hudson (Ed.), *From bones to behavior: ethnoarchaeological and experimental contributions to*
the interpretation of faunal remains (pp. 200–227). Carbondale: Occasional Paper No. Center for
Archaeological Investigations, Southern Illinois University.

Outram, A. K. (2002). Bone Fracture and Within-bone Nutrients : an Experimentally Based Method for
Investigating Levels of Marrow Extraction. In P. Miracle & N. Milner (Eds.), *Consuming Passions*
and Patterns of Consumption. (McDonald I.). Cambridge.

Patou-Mathis, M. (1985). La fracturation des os longs de mammifères: élaboration d'un lexique et d'une
fiche type. *Artefacts*, *1*, 11–22.

Pickering, T. R., & Egeland, C. P. (2006). Experimental patterns of hammerstone percussion damage
on bones: Implications for inferences of carcass processing by humans. *Journal of Archaeological*
Science, *33*, 459–469. doi:10.1016/j.jas.2005.09.001

R Core Team. (2019). R: A language and environment for statistical computing. *Foundation for*
Statistical Computing, Vienna, Austria.

Romandini, M., Nannini, N., Tagliacozzo, A., & Peresani, M. (2014). The ungulate assemblage from
layer A9 at Grotta di Fumane, Italy: A zooarchaeological contribution to the reconstruction of
Neanderthal ecology. *Quaternary International*, *337*, 11–27. doi:10.1016/j.quaint.2014.03.027

Rovira Formento, M. (2010). *Aproximación experimental a la explotación de huesos largos de grandes*
animales para la recuperación de la médula ósea y su aplicación arqueológica al registro
faunístico del Nivel 3 colluvio de Isernia La Pineta (Molise , Italia). Arqueologia del Quaternari i
Evolució Humana.

Speth, J. D., & Spielman, K. A. (1983). Energy source, protein metabolism, and hunter-gatherer
subsistence strategies. *Journal of Anthropological Archaeology*, *2 (1)*, 1–31

Speth, J. D. (1989). Early homonid hunting and scavaging: the role of meat as an energy source. *Journal*
of Human Evolution, *18*(4), 329–343. doi:http://dx.doi.org/10.1016/0047-2484(89)90035-3

Speth, J. D. (2010). *The Paleoanthropology and Archaeology of Big-Game Hunting*. New York, NY:
Springer New York. doi:10.1007/978-1-4419-6733-6

Speth, J. D., Meignen, L., Bar-Yosef, O., & Goldberg, P. (2012). Spatial organization of Middle
Paleolithic occupation X in Kebara Cave (Israel): Concentrations of animal bones. *Quaternary*

International, *247*, 85–102. doi:10.1016/j.quaint.2011.03.001.

Stavrova, T., Borel, A., Daujeard, C., & Vettese, D. (2019). A GIS based approach to long bone breakage
patterns derived from marrow extraction. *Plos One*, *14*(5), e0216733.
doi:10.1371/journal.pone.0216733

Stiner, M. C., Gopher, A., & Barkai, R. (2011). Hearth-side socioeconomics, hunting and paleoecology
during the late Lower Paleolithic at Qesem Cave, Israel. *Journal of Human Evolution*, *60*(2), 213–
233. doi:10.1016/j.jhevol.2010.10.006

Thiébaut, C., Claud, É., Coudenneau, A., Coumont, M.-P., Asselin, G., Beauval, C., et al. (2007). *Des
traces et des Hommes*.

Valensi, P., Michel, V., El Guennouni, K., & Liouville, M. (2013). New data on human behavior from
a 160,000 year old Acheulean occupation level at Lazaret cave, south-east France: An
archaeozoological approach. *Quaternary International*, *316*, 123–139.
doi:10.1016/j.quaint.2013.10.034

Vettese, D., Daujeard, C., Blasco, R., Borel, A., Caceres, I., & Moncel, M. H. (2017). Neandertal long
bone breakage process: Standardized or random patterns? The example of Abri du Maras
(Southeastern France, MIS 3). *Journal of Archaeological Science: Reports*, *13*, 151–163.
doi:10.1016/j.jasrep.2017.03.029

Vettese, D., Blasco, R., Cáceres, I., Gaudzinski-Windheuser, S., Moncel, M., Hohenstein, U. T., &
Daujeard, C. (2020). Towards an understanding of hominin marrow extraction strategies: a
proposal for a percussion mark terminology. *Archaeological and Anthropological Sciences*, *12*(2),
48. doi:10.1007/s12520-019-00972-8

Delphine Vettese 2020. An archaeological experiment focused on the intuitive way of long bones
breakage to extract marrow. **protocols.io**dx.doi.org/10.17504/protocols.io.be7fjhjn

Villa, P., & Mahieu, E. (1991). Breakage patterns of human long bones. *Journal of Human Evolution*,
21(1910), 27–48.

White, T. (1992). *Prehistoric cannibalism at Mancos 5MTUMR-2346* (Princeton.). Oxford.

Yravedra, J., Maté-González, M. Á., Palomeque-González, J. F., Aramendi, J., Estaca-Gómez, V., San
Juan Blazquez, M., et al. (2017). A new approach to raw material use in the exploitation of animal
carcasses at BK (Upper Bed II, Olduvai Gorge, Tanzania): a micro-photogrammetric and
geometric morphometric analysis of fossil cut marks. *Boreas*. doi:10.1111/bor.12224

944