A common analgesic enhances the anti-tumour activity of 5-aza-2'-deoxycytidine through induction of oxidative stress

Hannah J. Gleneadie[1,10], Amy H. Baker[1], Nikolaos Batis[2], Jennifer Bryant[2], Yao Jiang[3], Samuel J.H. Clokie[4], Hisham Mehanna[2], Paloma Garcia[5], Deena M.A. Gendoo[6], Sally Roberts[5], Alfredo A. Molinolo[7], J. Silvio Gutkind[8], Ben A. Scheven[1], Paul R. Cooper[1], Farhat L. Khanim[9] and Malgorzata Wiench[1, 5,*].

[1]School of Dentistry, Institute of Clinical Studies, College of Medical and Dental Sciences, The University of Birmingham, Birmingham, B5 7EG, UK; [2]Institute of Head and Neck Studies and Education (InHANSE), The University of Birmingham, Birmingham, B15 2TT, UK; [3]School of Biosciences, The University of Birmingham, Birmingham, B15 2TT, UK; [4]West Midlands Regional Genetics Laboratory, Birmingham Women's and Children's Hospital, Birmingham, B15 2TG, UK; [5]Institute of Cancer and Genomic Sciences, College of Medical and Dental Sciences, The University of Birmingham, Birmingham, B15 2TT, UK; [6]Centre for Computational Biology, Institute of Cancer and Genomic Sciences, The University of Birmingham, Birmingham, B15 2TT, UK; [7]Moores Cancer Center and Department of Pathology, University of California San Diego, La Jolla, CA 92093, USA; [8]Department of Pharmacology and Moores Cancer Center, University of California San Diego, La Jolla, CA 92093, USA; [9]Institute of Clinical Studies, University of Birmingham, Birmingham B15 2TT, UK, [10]Present address: MRC London Institute of Medical Sciences, Imperial College London, London, W12 0NN, UK

*Correspondence: Malgorzata Wiench, School of Dentistry, Institute of Clinical Studies, Institute of Cancer and Genomic Sciences, The University of Birmingham, 5 Mill Pool Way, Birmingham B5 7EG, Tel: +44 (0) 1214665520, m.d.wiench@bham.ac.uk

Running title: Paracetamol sensitizes cancer cells to DAC

Financial support: This research was supported by grants from the FP7 Framework Marie Curie Actions CIG (methDRE) to M. Wiench and The Royal Society (RS-9-10-2012) to M. Wiench. H.J. Gleneadie received a joint University of Birmingham and Medical Research Council UK fellowship.

Character count (with spaces): 70,503

Abstract

The DNA demethylating agent 5-aza-2'-deoxycytidine (DAC) has anti-cancer therapeutic potential, but its clinical efficacy is hindered by DNA damage-related side effects and its use in solid tumours is debated. Here we describe how common analgesic paracetamol can augment the effects of DAC on cancer cell proliferation and differentiation, without enhancing DNA damage. Firstly, DAC treatment specifically upregulates cyclooxygenase-2-prostaglandin E_2 pathway, inadvertently providing cancer cells with survival potential, while the addition of paracetamol offsets this effect. Secondly, in the presence of paracetamol, DAC leads to glutathione depletion and ROS accumulation. Finally, this oxidative stress is further enhanced by DAC suppressing anti-oxidant and thioredoxin responses. Analyses of TCGA datasets show that the DAC/paracetamol combination targets pathways of clinical importance in many cancers. The benefits of combined treatment are demonstrated here in head and neck squamous cell carcinoma (HNSCC) and acute myeloid leukaemia cell lines and corroborated in a HNSCC xenograft mouse model. In summary, paracetamol increases DAC efficacy such that it could be used at clinically relevant doses; providing a strong rationale for consideration in cancer therapy.

Keywords: 5-aza-2'-deoxycytidine/acute myeloid leukemia/epigenetic therapies/head and neck squamous cell carcinoma/paracetamol

Introduction

Aberrant DNA methylation patterns are common in most cancers, arise early in tumour development and are potentially reversible by hypomethylating agents (Jones & Baylin, 2002). The DNA demethylating agent 5-aza-2'-deoxycytidine (Decitabine or DAC) is a nucleoside analogue that incorporates into replicating DNA in place of cytosine where it traps and promotes the degradation of DNA methyltransferases (DNMTs) (Stresemann & Lyko, 2008). This results in two anti-cancer activities: methyl marks cannot be copied during DNA replication causing widespread DNA demethylation and adducts are formed in the DNA leading to DNA damage and apoptosis (Stresemann & Lyko, 2008). DAC has been approved by the European Medicines Agency (EMA) for treatment of acute myeloid leukaemia (AML) (Kantarjian *et al*, 2012; Nieto *et al*, 2016). Pre-clinical studies suggest it might also be effective in solid tumours (Tsai *et al*, 2012), athough the outcomes of clinical trials are highly

variable (Linnekamp *et al*, 2017). DNA demethylating drugs are thought to de-repress epigenetically silenced tumour suppressor genes as well as demethylate endogenous retroviruses (ERVs), triggering an antiviral immune response and cancer cell death (Chiappinelli *et al*, 2015; Roulois *et al*, 2015; Stresemann & Lyko, 2008).

Head and neck squamous cell carcinoma (HNSCC), the 6th most common cancer worldwide, has 5-year survival rates of ≤40%, highlighting a pressing need for new therapies (Simard *et al*, 2014).The tumours originate from stratified squamous epithelium of the oral cavity and pharynx where the cells in the basal cell layer proliferate and replenish the suprabasal layers which undergo terminal differentiation (Squier & Kremer, 2001). Although DNA methylation aberrations are common in HNSCC (Steinmann *et al*, 2009) the clinical evaluation of DAC potential in HNSCC is very limited (Abele *et al*, 1987).

In solid tumours, DAC alone may not be curative, but favourable effects were observed for epigenetic agents combined with other chemo- and immune-therapies (Linnekamp *et al.*, 2017). So far it has not been explored whether the response to DAC could be enhanced by other compounds, not traditionally used in cancer treatment. In the current study, a custom-built library of 100-commonly used, cost-effective, off-patent drugs (Khanim *et al*, 2011) was investigated for their ability to sensitise HNSCC cells to DAC treatment. Of the drugs tested, paracetamol was identified to work in synergy with DAC.

Paracetamol (acetaminophen) is the most commonly used analgesic and antipyretic in both Europe and the United States, present on the market since the 1950s (Bertolini *et al*, 2006). Paracetamol affects the cyclooxygenase (COX) pathway wherein arachidonic acid (AA) is metabolized to prostaglandin H_2 (PGH_2) by either constitutively expressed COX-1 (PTGS1) or inducible COX-2 (PTGS2) (Graham *et al*, 2013). PGH_2 is then rapidly converted, by respective prostaglandin synthases, into effector prostanoids (prostaglandins PGE_2, PGF_2; PGI_2 and PGD_2 or thromboxane TXA) and these work, in part, through metabolite-specific G-protein coupled receptors to activate downstream pathways (Graham *et al.*, 2013). Additionally, other AA-derived metabolites are produced either through the lipoxygenase (LOX) or the monooxygenase (cytochrome P450) pathways (Wang & Dubois, 2010).

The COX-2-PGE_2 axis is associated with inflammation, growth and survival and is thought to contribute to the 'inflammogenesis of cancer' (Liu *et al*, 2015). Increased expression of COX-2 and production of PGE_2 are found in many solid tumours, including HNSCC, and correlate with tumour stage, metastasis and worse clinical outcome whilst low levels are

associated with better response to chemotherapy (Liu *et al.*, 2015; Saba *et al*, 2009). This pathway has also been implicated in the development of chemoresistance and linked to repopulation of cancer stem cells (Kurtova *et al*, 2015). Hence, COX-2 inhibitors have been tested for their therapeutic anti-cancer potential, showing protection against cancer development (Kim *et al*, 2010; Saba *et al.*, 2009). However, the only research that suggests paracetamol may have therapeutic potential in established tumours used overdose concentrations of the drug, either alone or in combination with other chemotherapeutics (Kobrinsky *et al*, 1996; Posadas *et al*, 2007; Wu *et al*, 2013).

Here we show that paracetamol can be used at clinically relevant concentrations to sensitize cancer cells (both HNSCC and AML) to DAC treatment, allowing for DAC dose-reduction.

Results

HNSCC cell lines show bimodal sensitivity to DAC treatment
To establish the potential of DAC as a therapeutic in HNSCC, relative cell viability was determined in four HNSCC cell lines and in normal human oral keratinocytes (HOK) after 96h of treatment (Fig 1A). HOK cells showed only moderate sensitivity to DAC treatment with no decrease in viability at clinically relevant concentrations and an IC_{50} of 8.93 µM. Interestingly, the four HNSCC cell lines could be divided into two distinct groups; DAC-sensitive (VU40T, IC_{50} of 2.17 µM and HN12, IC_{50} of 0.81 µM) and those with little (SCC040, IC_{50} of 10.61 µM) to no sensitivity (UDSCC2) (Fig 1A). This pattern was mirrored in the ability of DAC to demethylate DNA in the sensitive cell lines only (Fig 1B). These data suggest that the efficacy of DAC treatment is proportional to its ability to demethylate DNA, therefore likely dependent on cellular drug uptake, activation or retention.

Efficacy of DAC treatment can be synergistically increased with paracetamol
In patients with AML, a 5-day regimen of 20mg/m^2 DAC gave a maximum plasma concentration (Cmax) of 107 ng/ml, equivalent to 469 nM (European Medicines Agency, 2019; Nieto *et al.*, 2016), while all HNSCC cell lines have an IC_{50} value greater than 500 nM. Therefore, one sensitive (VU40T) and one resistant (SCC040) cell line were subjected to DAC sensitising screen to establish whether the efficacy of DAC could be increased (Fig 1C, Appendix Fig S1). The cells were treated with 500 nM DAC with or without one of a panel of 100 off-patent drugs (drug repurposing library FMC1(Khanim *et al.*, 2011)) and compared to the drug-only control. In the resistant SCC040 cells, none of the drugs were able to

sensitize the cells to DAC (Appendix Fig S1A). However, in the DAC-sensitive VU40T cell line, paracetamol, valproic acid (VPA) and zinc acetate further decreased cell viability (Fig 1C, Appendix Fig S1B). The paracetamol effect was replicated in the remaining cell lines: the efficacy of DAC was increased by paracetamol and zinc acetate in DAC-sensitive HN12 but not in DAC-resistant UDSCC2 cells (Fig 1D). Importantly, paracetamol alone did not alter the viability of HOK cells which were otherwise highly sensitive to both VPA and zinc acetate (Fig 1D). Therefore, DAC-paracetamol combination was further tested for synergy in VU40T and HN12 cells using the Chou-Talalay method (Chou, 2006). The cell viability was assessed in response to each drug separately and in combination across eight constant-ratio matched titration of Cmax (500 nM for DAC and 132 µM for paracetamol) (Fig 1E-F). This analysis showed combination index (CI) values less than 1 at all bar the lowest concentration, demonstrating synergy of supra-additive nature (Fig 1G). In VU40T cells, the dose reduction index (DRI) indicated that almost 5 times less of each drug could be used when applied in combination (Appendix Table S1), allowing DAC dose reduction from 2.26 µM to the clinically relevant 450 nM (European Medicines Agency, 2019; Nieto *et al.*, 2016). A similar reduction was observed for HN12 cells (Appendix Table S2).

Combined DAC-paracetamol treatment augments the effects of DAC on cell metabolism, proliferation and markers of basal epithelial cells

In DAC-responsive VU40T cells the combined DAC-paracetamol treatment decreased cell number compared to DAC alone (Fig 2A). It was next investigated whether the underlying mechanisms involved reduced proliferation, altered cell cycle progression or increased cell death, possibly due to enhanced DNA damage.

DAC is known to cause cell death and DNA damage (Stresemann & Lyko, 2008). As expected, 500 nM DAC increased both apoptosis and necrosis as shown by annexin V and propidium iodide staining but this was not altered by the addition of paracetamol (Fig 2A-B). Similarly, in VU40T cells, DAC treatment significantly increased DNA damage and double strand breaks as highlighted by an increase in the number of nuclei with γH2AX foci when compared to control; however, this too was not additionally enhanced by combined treatment (Fig 2C).

When compared to untreated cells none of the treatments resulted in change in cell cycle progression, however, the addition of paracetamol to DAC treatment led to fewer cells progressing to G2/M phase than for DAC alone (Fig 2D). This additional effect of

paracetamol was also observed for cell proliferation. Treatment with 500 nM DAC reduced the number of Ki67 positive nuclei and combined treatment further decreased their number, whilst paracetamol alone had no effect (Fig 2A and E). Furthermore, markers for the basal cell layer (the only cell layer in stratified epithelium that normally contains dividing cells), *TP63* and keratin 5 (*KRT5*), were down-regulated by DAC in a dose-dependent manner and this was enhanced by DAC-paracetamol combined treatment (Fig 2F). Notably, TP63 is a key transcription factor in stratified epithelium (Yoh & Prywes, 2015) and its overexpression and/or amplification is observed in ~ 30% of tumours in the TCGA HNSCC cohort (Appendix Fig S2). Conversely, involucrin (*IVL*), a marker for differentiated, suprabasal layers was upregulated upon DAC and even more so by DAC-paracetamol treatments (Fig 2G).

RNA sequencing performed on the DAC-sensitive VU40T cells demonstrated that DAC treatment substantially altered the transcriptome while paracetamol alone only resulted in modest changes (Fig 2H). However, addition of paracetamol to DAC treatment resulted in significantly greater changes in gene expression compared to DAC alone (Fig 2H). Six gene sets, up- and down-regulated by each treatment (Fig 2H, Appendix Table S3), were subjected to gene set enrichment analysis using GO Biological Processes (Ashburner *et al*, 2000) and consolidated in REVIGO (Supek *et al*, 2011) (Fig 2I, Appendix Table S4). Paracetamol treatment alone resulted in no significant enrichment for down-regulated gene sets; treatment primarily resulted in up-regulation of genes involved in respiratory electron transport chain. The profile upregulated by DAC was dominated by immune terms, especially interferon type I response. This was also evident in the combined treatment, although to a lesser extent. Interestingly, combined treatment was enriched for terms related to tissue development and differentiation (including 'pharyngeal system development' Appendix Table S4), confirming the possible change from basal cell-like to more differentiated epithelial cell phenotype. Genes down-regulated by both DAC and DAC+paracetamol showed similar ontology groupings related to DNA, protein and RNA metabolism but the enrichment was much stronger for the combined treatment (Fig 2I) due to higher number of differentially regulated genes (Fig 2H). This suggests that the addition of paracetamol to DAC significantly amplifies decreases in metabolism and DNA replication caused by DAC alone.

Therefore, although DAC alone has profound effects on proliferation, differentiation, cell death and DNA damage, the further reduction in viability observed after the addition of

paracetamol appears to be associated with decreased proliferation and divergence from the basal cell-like phenotype.

DAC treatment enhances the cyclooxygenase pathway, which is offset by paracetamol

Paracetamol is understood to act on the cyclooxygenase pathway, primarily through inhibition of COX-2 (PTGS2) (Graham *et al.*, 2013). Notably, a geneset for 'Prostanoid biosynthetic process' was enriched after both DAC and combined treatments (Appendix Table S4). Therefore, the effect of DAC on this pathway was further examined. In DAC-sensitive cell lines only, *PTGS2* RNA and protein levels were upregulated following DAC treatment and this was maintained after combined treatment (Fig 3A and B). A corresponding increase in the downstream product, prostaglandin E_2 (PGE_2), was observed in the DAC-sensitive VU40T cells but not in the resistant SCC040 cells (Fig 3C). PGE_2 returned to basal levels by the addition of 132 µM paracetamol (Fig 3C) which indicates that paracetamol dampens DAC-induced COX-2 pathway activation. In addition, expression of the PGE_2 receptors *PTGER1* and *PTGER2* increased in the DAC-sensitive cells VU40T and HN12, respectively, whilst no significant changes in *PTGER1-4* expression occurred in the DAC-resistant cells UDSCC2 and SCC040 (Fig 3D and Appendix Fig S3A). Transcriptional activation of the COX-2-PGE_2 pathway by DAC was evident in the RNA-seq data, together with COX-2-TXA_2, whilst the PGF_2, PGI_2 and PGD_2, pathways were not upregulated (Appendix Fig S3B and C). In summary, DAC treatment specifically upregulates many aspects of the COX-2-PGE_2 pathway, inadvertently providing the cancer cells with growth and survival potential, while the addition of paracetamol offsets this effect (Fig 3E).

There is a possibility that blocking the COX pathway with paracetamol could shunt AA towards the LOX pathway also leading to increased survival potential (Park *et al*, 2012). DAC treatment altered gene expression of enzymes involved in the LOX pathway by both up- (*ALOX5* and *ALOX15B*) and down- (*LT4H* and *ALOX12*) regulating them (Appendix Fig S3D). However, the secretion of cysteinyl leukotrienes and leukotriene B_4 remained below levels detectable by ELISA in all experimental conditions in VU40T and SCC040 cells. Therefore, there is no indication for LOX pathway compensation following COX-2 inhibition by paracetamol in the cell lines examined, although the involvement of other metabolites cannot be excluded (Appendix Fig S3E).

DAC treatment complements cancer-related activation of COX-2-PGE_2 pathway

Activation of the COX-2-PGE$_2$ pathway has been previously indicated in both HNSCC and other cancer types (Liu *et al.*, 2015; Saba *et al.*, 2009). In the most recent TCGA cohort, 29% of HNSCC tumours (150/520 patients) have at least one component of COX-2-PGE$_2$ pathway transcriptionally activated: mostly through upregulation of PGE$_2$ synthases or PGE$_2$ receptors (Fig 3F). Similar activation can be observed in other cancers (Appendix Table S5). However, over-expression of *PTGS2* itself is relatively rare (2.3% in HNSCC, Fig 3F), potentially serving as a limiting factor in the full activation of the pathway. Therefore, the DAC-induced upregulation of *PTGS2* could remove this limitation and counteract the anti-tumour effects of DAC. This mechanism could be applicable to other tumour types; *PTGS2* up-regulation by DAC was detected by Drug Perturbation Signatures (Fig 3G) from the cMAP dataset (Lamb *et al*, 2006) indicating the wider potential of DAC-paracetamol co-treatment.

Combined treatment mimics the effects of paracetamol overdose and depletes glutathione levels

If COX-2-PGE$_2$ pathway alterations were to be solely responsible for the synergy between DAC and paracetamol, other COX inhibitors should have a similar effect. However, neither the generic COX inhibitor ibuprofen, nor the COX-2 specific inhibitor valdecoxib sensitized HNSCC cells to DAC treatment (Fig 4A, Appendix Fig S4), despite valdecoxib being able to reduce DAC-stimulated PGE$_2$ production to a similar extent as paracetamol (Fig 4B). This suggests an additional, paracetamol-specific, mechanism accounts for the synergistic relationship between DAC and paracetamol.

Previous work on the anti-cancer therapeutic potential of paracetamol involved toxic doses of the drug (Kobrinsky *et al.*, 1996; Posadas *et al.*, 2007; Wu *et al.*, 2013). Efficacy was attributed to an accumulation of toxic metabolite of paracetamol, N-acetyl-p-benzoquinone-imine (NAPQI), resulting in glutathione depletion (Kobrinsky *et al.*, 1996; Posadas *et al.*, 2007; Wu *et al.*, 2013) (Fig 4C). By comparison, our current work was performed using a safe, clinically achievable serum concentration of paracetamol (132 µM) . However, in DAC-sensitive VU40T cells, DAC treatment caused an increase in the CYP450 enzyme, *CYP2E1*, thought to be primarily responsible for the conversion of paracetamol into NAPQI (Fig 4D-E). Assessing NAPQI levels is unreliable due to its highly reactive nature (Dahlin *et al*, 1984). However, combined treatment led to a dramatic reduction in GSH levels, a surrogate marker often used, and the effect was equivalent to a high dose (1mM) paracetamol (Fig 4F). Furthermore, N-acetyl cysteine (NAC) is clinically used as an antidote to treat paracetamol overdose by replenishing GSH and preventing NAPQI accumulation (Kobrinsky *et al.*, 1996).

In VU40T cells, 48h pre-treatment with 2.5 mM NAC led to increased cell viability in all conditions, but significantly and to the greatest extent in the DAC+paracetamol treated cells, where viability was restored to the control level (Fig 4G). Therefore, the combined treatment mimics the effects of paracetamol overdose at therapeutic concentrations and depletes GSH stores in tumour cells.

GSH is an intracellular antioxidant that acts primarily as a reactive oxygen species (ROS) scavenger (Forman *et al*, 2009). In agreement with this, DAC-paracetamol co-treatment significantly increased both intracellular ROS (Fig 4H) and mitochondrial superoxide (Fig 4I) as compared to DAC alone. This was specific for DAC co-treatment with paracetamol and not valdecoxib (Fig 4H-I). Accumulation of ROS typically triggers an anti-oxidant response which can protect cancer cells treated with chemotherapeutics (Ray *et al*, 2012). However, upon both DAC and DAC-paracetamol treatment the majority of genes described as direct anti-oxidant responders (Raghunath *et al*, 2018) are down-regulated (Fig 4J). Therefore, the transcriptional anti-ROS response is impaired by DAC treatment which leads to further exacerbation of oxidative stress and decreased cancer cell survival. Finally, it has been shown that keratinocytes can tolerate GSH depletion as long as the cysteine pools (uptake of cystine by SLC7A11 or cysteine by SLC1A4) and the thioredoxin reductase system (TXN/TXNRD) are functional (Telorack *et al*, 2016). Again, DAC downregulated the majority of these genes which would inhibit this compensating mechanism (Fig 4K).

To summarize, in cancer cells DAC-paracetamol co-treatment mimics the mechanisms of paracetamol overdose, whilst cell adaptation to oxidative stress is impaired by DAC (Fig 4L). The specificity of the DAC-paracetamol interaction is further supported through Drug Set Enrichment Analysis (DSEA). The DSEA algorithm aims to identify the mechanisms of action shared by a set of drugs (Napolitano *et al*, 2016). Indeed, DAC and paracetamol share significant enrichment for COX, cytochrome P450 and GSH metabolism pathways, while this was not observed for DAC and valdecoxib combination (Fig 4M, Appendix Fig S5). Interestingly, the analysis also indicated high enrichment for genes attributed to keratinocyte differentiation (Fig 4M).

DAC-paracetamol affected pathways contribute to HNSCC patients' survival

The mechanisms underlying the efficacy of DAC-paracetamol co-treatment identified so far are multifactorial (Fig 5A). Firstly, the COX-2-PGE$_2$ pathway is specifically upregulated by DAC, potentially improving cancer cells survival. Indeed, when upregulated, this pathway

negatively affects HNSCC patient survival (Fig 5B). This is specific to COX-2-PGE$_2$ but not to other COX pathways or to the LOX pathway (Appendix Fig S6A and B). The addition of paracetamol to DAC treatment counteracts the induction of this pro-survival mechanism. Secondly, combined treatment leads to depletion of glutathione stores followed by oxidative stress, both of which restrict tumour growth and improve the efficacy of anti-cancer drugs (Traverso *et al*, 2013). Therefore, in the proposed scenario, GSH is required to maintain cancer cell fitness. In agreement with this, genes involved in maintaining GSH stores are linked to poorer patients' survival in the TCGA cohort of HNSCC tumours (Fig 5C) as well as other cancers (Appendix Table S6). Thirdly, DAC treatment transcriptionally down-regulates anti-oxidant and thioredoxin responses known to prevent cellular adaptation to oxidative stress and protection from oxidative damage. Again, upregulation of these pathways, particularly the thioredoxin response, correlates with decreased survival of HNSCC patients (Fig 5D). These results suggest that the DAC-paracetamol combination targets pathways and genes of clinical importance in HNSCC patients.

In vivo potential of DAC-paracetamol combination treatment

An NSG mouse xenograft model using human HNSCC FaDu cells previously used in drug efficacy studies (Bryant *et al*, 2019) was utilized to assess the DAC-paracetamol treatment *in vivo*. Like HN12 and VU40T, FaDu cells showed high supra-additive effects for combined treatment (Fig 6A and B). Mice were injected with FaDu cells and the treatments (DAC, paracetamol, DAC+paracetamol or vehicle (PBS)) were administered 5 days a week (Fig 6C). After the first two weeks, tumours in the control and paracetamol-treated groups reached the maximum permissible size, while DAC alone and DAC+paracetamol groups were treated for another week (Fig 6C). Due to the strong initial response to DAC, the treatment was carried out with reduced DAC concentration (0.2 mg/kg). Although DAC alone showed a strong anti-tumour effect, the DAC-paracetamol treated tumours remained consistently smaller throughout the duration of the treatment (Fig 6C and D). No tumours from the DAC-paracetamol treated group exceeded 300 mm^3 (Fig 6E) and 5/6 animals survived until the end of the experiment at which point they started losing weight (Appendix Fig S7). Transcript analysis of RNA extracted from tumour tissue showed alterations consistent with the mechanism of DAC-paracetamol synergy: upregulation of *PTGS2* and *CYP2E1* (Fig 6F) and down-regulation of *TP63* and *KRT5* (Fig 6G).

Synergistic effects of DAC-paracetamol co-treatment are present in AML cell lines

Currently, DAC only has EMA approval in the treatment of AML (Nieto *et al.*, 2016). Therefore, combined treatment of DAC plus paracetamol was tested in two AML cell lines, SKM-1 and HL-60. The cells were treated for 72h and the drugs were found to work synergistically in both of them (Fig 7A-C). The combined effect was even more apparent after prolonged culture, which aimed to mimic the DAC treatment regimen in AML patients (European Medicines Agency, 2019) where the cells were treated for 72h followed by 21 days withdrawal over four cycles (Fig 7D). The benefits of DAC-paracetamol treatment were more pronounced with each consecutive cycle and by the time the cells reached the fourth cycle they were still responding well to combined treatment while becoming resistant to DAC alone (Fig 7D).

Similar to HNSCC cells, DAC treatment in AML cell lines activated several aspects of COX-2-PGE$_2$ pathway including upregulation of *PTGS2* and PGE$_2$ receptors' expression (Fig 7E). Upregulation of *CYP2E1* following DAC treatment was also noted for SKM-1 cells (Fig 7F). Ultimately, DAC increased ROS and mitochondrial superoxide production in these AML cells and this was further significantly enhanced by the addition of paracetamol but not valdecoxib (Fig 7G and H).

In further agreement with the results obtained for HNSCC cells, combined treatment did not add to the cell death caused by DAC alone as assessed by annexin V (Fig 7I). There was also little change to cell cycle progression after DAC when compared to control while combined treatment increased the number of cells retained in S phase when compared to DAC alone (Fig 7J). Following DAC and DAC-paracetamol treatments, AML cells displayed an increase of myeloid differentiation marker *CD11b* (*ITGAM*) (Fig 7K). Although this was not significantly different between combined treatment and DAC alone the effects of DAC+paracetamol were more apparent when cell morphology was assessed and showed a change towards myelocyte with increased cytoplasm and a high number of vacuoles (Fig 7L). These data indicate that DAC-paracetamol synergy and the effect it has on oxidative stress could be applicable to blood malignancies, where DAC is an accepted and widely used therapeutic option.

Discussion

The search for new drugs to improve the poor survival rate of HNSCC is ongoing. Our preliminary results using four HNSCC cell lines and an *in vivo* mouse model show that DAC alone has therapeutic potential in HNSCC which involves type I interferon activation.

However, the response is variable as it has been observed for other solid tumours (Linnekamp *et al.*, 2017) and patient stratification will be necessary to identify the DAC-responders. Future studies will look for predictive biomarkers, however the results shown here are clear that sensitivity to DAC is primarily dependent upon the ability of DAC to demethylate the DNA, and therefore likely due to incorporation, activation or retention of the drug, as suggested previously (Qin *et al*, 2009; Wu *et al*, 2016). Furthermore, an initial response to DAC treatment was a prerequisite for synergy with paracetamol, therefore the co-treatment is dependent upon the DNA demethylating capacity of DAC.

This is the first study to identify that paracetamol can enhance the anti-tumour activity of DAC. There are two main translational impacts of the DAC-paracetamol combination. Our data strongly indicate that adding paracetamol to DAC treatment could significantly lower the DAC dose needed to achieve therapeutic effects, broadening DAC application. In addition, it opens a potential for DAC dose reduction therefore potentially reducing DNA damage-related side effects.

Paracetamol is routinely prescribed as an analgesic, however it has not been considered so far whether its use may influence, positively or negatively, the efficacy of chemotherapy regimens. This study highlights that uncontrolled use of paracetamol during cancer therapies and clinical trials could affect the outcomes and interpretation of the results. Hence, further studies are required to look at the impact of supportive care medication in oncology.

Considering both DAC and paracetamol have broad effects with many potential interactions, we investigated and identified key mechanisms within the AA metabolism pathway that could underlie the synergy (summarized in Fig 5A). Both our data and analysis of public databases point towards DAC explicitly upregulating COX-2-PGE$_2$ pathway, therefore providing cancer cells with survival advantage. This also reflects the previously reported bias in cancer cells towards PGE$_2$ production to the detriment of other prostaglandins (Cebola *et al*, 2015; Cebola & Peinado, 2012). It remains to be established whether the COX-2-PGE$_2$ pathway activation is a direct result of DNA demethylation or is rather due to indirect mechanisms (e.g. response to dsRNA, cytokines or growth factors).

Surprisingly, the synergistic effect observed for paracetamol could not be reproduced using other COX-2 inhibitors. The data indicate an alternative paracetamol-specific mechanism; DAC-induced mimicry of paracetamol overdose leading to GSH depletion and exacerbated oxidative stress in cancer cells, both of which have the potential to restrict tumour growth and

improve patient survival (Traverso *et al.*, 2013). These cytoprotective pathways are also involved in development of drug resistance, hence their suppression might account for the extended drug sensitivity observed in AML cells after combined treatment.

Response to DAC has a profound effect on the transcriptional programme in HNSCC cells which is dominated by activation of type I interferon and anti-viral pathways, agreeing with recent reports on the role of 'viral mimicry' in cancer treatment with demethylating agents (Chiappinelli *et al.*, 2015; Roulois *et al.*, 2015). These effects are maintained but not increased by combined treatment. Instead, addition of paracetamol to DAC treatment led to a decrease in DNA, RNA and protein metabolism, together with reduced proliferation and enhanced differentiation. Concurrently, a strong enrichment for 'keratinocyte differentiation' was noted in the DSEA dataset shared by DAC and paracetamol signatures. Further experiments are required to establish the exact mechanisms leading to the changes in cell proliferation and differentiation upon combined treatment and whether they are linked to alterations in COX2-PGE$_2$ pathway and/or to 'mimicry of paracetamol overdose'. One possibility involves the effects of AA metabolism and ROS on PI3K/Akt and MAPK pathways, the main proliferation drivers in HNSCC (Cancer Genome Atlas, 2015). In HNSCC, mutations in the AA pathway have been reported to downregulate PI3K/Akt signalling (Cancer Genome Atlas, 2015), while patients with activating *PIK3CA* mutations specifically benefited from regular use of NSAIDs (Hedberg *et al*, 2019); confirming the significance of AA and COX pathways in HNSCC development and treatment.

Results obtained for AML cells confirm similar genes and pathways are being affected and suggest wider clinical potential for the DAC-paracetamol synergy. This is of importance because DAC is an accepted therapeutic option in AML patients which could directly benefit from the addition of paracetamol to their treatment regime. Paracetamol is one of the most commonly used drugs, both prescribed and self-medicated. Therefore, considering the mechanisms described here, its interaction with other drugs, especially chemotherapeutics, should be taken into consideration in cancer treatment.

Materials and methods

Cell lines and culture conditions

Five human HNSCC cell lines were used: SCC040 (German Culture Collection, DSMZ (#ACC660)), FaDu (ATCC (HTB-43)), VU40T (Prof H. Joenje (VU University Medical

Centre, Amsterdam)), HN12 (Dr J.F. Ensley (Wayne State University, Detroit, MI)) and UDSCC2 (Dr Henning Bier (University of Duesseldorf, Germany)). The cell lines were authenticated using STR profiling (NorthGene, UK). All HNSCC cell lines were maintained in DMEM (Sigma- Aldrich) supplemented with 10% fetal bovine serum (FBS) (Sigma-Aldrich), 1% penicillin-streptomycin (Gibco), 4 mM L-glutamine (Sigma-Aldrich), 1X non-essential amino acids (Life Technologies) and 1 mM sodium pyruvate (Life Technologies). Primary human oral keratinocyte (HOK) cells were purchased from Caltag Medsystems and cultured over Poly-L-Lysine in Oral Keratinocyte Medium supplemented with 1% oral keratinocyte growth supplement (all purchased from ScienceCell) and 1% penicillin-streptomycin. Two AML cell lines were used: SKM-1 (from Dr Stefan Heinrichs, University of Essen, Germany) and HL-60 (ATCC (CCL-240)). Cells were cultured in RPMI1640 medium (ThermoFisher Scientific), supplemented with 15% FBS (Sigma-Aldrich), 1% L-glutamine and 1% penicillin-streptomycin. All cell lines were regularly tested for mycoplasma using MycoAlert (Lonza) and experiments were performed within 18 passages after thawing.

Drugs treatments

All drugs were purchased from Sigma-Aldrich. 5-Aza-2'-deoxycytidine (DAC) was dissolved in either 50% acetic acid or in ≥99.9% dimethyl sulfoxide (DMSO), all other drugs were dissolved in DMSO. The treatments were carried out for 96h (HNSCC cells) or 72h (AML cells) using a relevant vehicle as control.

Viability assay

Relative viability was determined in 96 well plates using the CellTiter-Blue Cell Viability Assay (Promega), including a minimum of triplicate wells per sample, vehicle, high concentration vehicle and 'media only' controls. Samples were normalised to a vehicle-only control. Sigma plot software (Systat Software Inc.) was used to generate sigmoidal, 4-parameter dose response curves after drug titrations. IC_{50} values were calculated using MyCurveFit (MyAssays Ltd.).

DAC sensitising assay

A panel of 100 drugs (Drug Library FMC1) were administered at the reported peak serum concentrations (Cmax) (Khanim et al., 2011). Cells were seeded in 96-well plates and treated with each drug alone or in combination with 500 nM DAC. The assay was performed blind in biological triplicates with controls hidden within the panel.

Determining synergy

The Chou-Talalay method was employed to determine synergy (Chou, 2006) using constant-ratio matched titrations (0.125X, 0.25X, 0.5X, 1X, 2X, 3X, 4X, and 8X) of multiples of the Cmax (500 nM for DAC, 132 µM for paracetamol) and assessed by cell viability. The CompuSyn software (Chou, 2006) was used to calculate combination index (CI) and dose reduction index (DRI) values. CI values below 1 indicate synergy. DRI values indicate how many times less each drug can be used when in combination.

Long treatment of AML cells

SKM-1 and HL-60 cells were subjected to four cycles of 72h treatments followed by 21 days withdrawal period (details in Appendix Materials and Methods), the cells were counted at each passage using trypan blue (Sigma-Aldrich) staining and growth rates were calculated.

DNA dot blotting

DNA was extracted using the DNeasy Blood and Tissue Kit (Qiagen) and DNA dot blotting was performed as described in (Brown, 2001) using a titration of DNA; detailed protocol and antibodies are provided in Appendix Materials and Methods. Blots were normalized to methylene blue (Sigma-Aldrich) staining.

Western Blot analysis

WB analysis was performed as described previously (Khanim *et al.*, 2011), with some alterations; protocol details and antibodies are provided in Appendix Materials and Methods.

Giemsa-Jenner staining

VU40T cells were grown on coverslips for 96h, fixed in methanol and stained with Giemsa and Jenner (VWR) as previously described (Khanim *et al.*, 2011). SKM-1 cells were transferred to a glass slide with a Cytospin 3 (Thermo Shandon) before fixation. Microscope images were taken using EVOS XL Core Imaging System or BX-50 Olympus with a 100x oil immersion lens.

Immunofluorescence analysis

Immunofluorescence analysis was undertaken as described in (Roulois *et al.*, 2015) with details in Appendix Materials and Methods.

Apoptosis assay and cell cycle analysis

Apoptosis and necrosis were assessed using the Annexin V Apoptosis Detection APC kit (eBioscience, ThermoFisher Scientific) as the manufacturer recommends (see also Appendix

Materials and Methods). For cell cycle analysis, cells were fixed in 70% ice cold ethanol and resuspended in 200 µl 50 µg/ml propidium iodide solution (Sigma-Aldrich) and 50 µl of 100 µg/ml RNase solution (Roche). The solutions were analysed by flow cytometry on a Cyan B FACS analyser (Beckman Coulter).

ELISA

The levels of PGE$_2$, Leukotriene B$_4$ (LTB$_4$) and Cysteinyl leukotrienes (LTC$_4$, LTD$_4$, and LTE$_4$) were assessed in cell media using respective ELISA kits (Abcam). The results were normalised to the corresponding CellTiter-Blue viability results or cell count.

Determining glutathione concentration

The glutathione levels were measured using the GSH-Glo Glutathione Assay (Promega) with 1 mM paracetamol used as a positive control. The results were normalized against the corresponding CellTiter-Blue cell viability.

NAC rescue assay

N-acely-l-cysteine (NAC, Sigma-Aldrich) was dissolved in water and neutralised using 1 M sodium hydroxide (NaOH) (Sigma-Aldrich). VU40T cells were treated with 2.5 mM NAC or an equivalent volume of vehicle for 48h. Following this, wells were washed with fresh media and cells were treated with DAC, paracetamol or both for 96h and cell viability was determined.

Assessment of reactive oxygen species (ROS) and mitochondrial superoxide (mitosox)

MitoSOX Red (Cat# M36008, ThermoFisher Scientific) was used to assess mitosox, while ROS were measured using carboxy-H$_2$DCFDA (5-(and-6)-carboxy-2',7'-dichlorodihydrofluorescein diacetate, Cat# C369, Invitrogen), according to the manufacturers' protocols (details in Appendix Materials and Methods).

Real-Time Quantitative PCR (qRT-PCR)

Total RNA was extracted using RNeasy Mini Kit (Qiagen), including DNase I digestion step (RNase-Free DNase set, Qiagen). 1 µg of RNA was reverse transcribed (iScript cDNA Synthesis Kit (BioRad)) and the cDNA was purified with QiaQuick PCR Purification Kit (Qiagen). Purified cDNA was quantified using Qubit dsDNA High Sensitivity Assay kit (Thermo Fisher Scientific) to support normalization. qRT-PCRs were performed on a LightCycler 480 II using LightCycler 480 SYBR Green (Roche). Additional information and primers' sequences are provided in Appendix Materials and Methods.

RNA sequencing

For each sample three biological RNA replicates were pooled to make a library using the TruSeq Stranded mRNA Library Prep kit and sequenced using Illumina NextSeq 500 in paired-end mode at 2x75 bases in the Genomics Birmingham facility (Birmingham, UK). Reads were aligned to the genome (hg19) using HiSAT2 and processed with bedtools to generate normalised coverage plots. Details on quantification and data processing are in Appendix Materials and Methods.

Mouse xenograft study

The mouse xenograft study was performed as described previously (Bryant *et al.*, 2019) and details of the protocol are in Appendix Materials and Methods. To examine the toxicity and anti-tumour efficacy of DAC and paracetamol, we utilised male NOD/SCID/gamma (NSG) mice (Charles River) in accordance with the UK Home Office Animal (Scientific Procedures) Act 1986 and approved by the local University of Birmingham Ethical Review Committee. 24 males were implanted with 5×10^6 FaDu HNSCC cells suspended in serum-free medium and injected subcutaneously into the right flank. After three days the tumours established and mice were randomly allocated into four treatment groups on a 5 day on, 2 day off regimen: 0.4 mg/kg DAC in PBS via IP injection (or 0.2 mg/kg in the week three); 1000 mg/kg paracetamol in PBS via oral gavage; DAC plus paracetamol as above; control (PBS) given both through the oral gavage and IP injection. Animals were monitored daily for signs of ill health and the tumours were measured. Upon culling, the tumours were excised and snap frozen for RNA extraction and subsequent analysis.

Quantification and statistical analysis

Statistical analysis and graphs were performed using GraphPad PRISM 8 software, unless stated otherwise. Details of the statistical tests used for each experiment are given in the corresponding figure legends.

TCGA data

Genetic alterations, gene expression and patient survival data were retrieved from The Cancer Genome Atlas (TCGA) via cBioPortal for Cancer Genomics (Gao *et al*, 2013); the details are described in Appendix Materials and Methods.

Drug perturbation signatures

Drug perturbation signatures were downloaded for the BROAD Connectivity Map dataset (CMAP) using the PharmacoGx package (version 1.14.0) (Smirnov *et al*, 2016) in R (details in Appendix Materials and Methods).

Drug Set Enrichment Analysis

The effect of Decitabine, paracetamol and valdecoxib drug combinations on KEGG pathways and Gene Ontology Biological Processes was conducted using the Drug Set Enrichment Analysis (DSEA) server (Napolitano *et al.*, 2016). Significance of the perturbed pathways of interest was identified using the corresponding p-values per geneset; $\log_{10}$ (p-values) were plotted as heatmaps.

Data availability

RNA-Seq data: The data are deposited at the GEO repository, accession number GSE110045 and SRA, accession number SRP132039.

Acknowledgments

We would like to thank Dr Jo Parish (University of Birmingham, UK) for critically reviewing the manuscript. This research was supported by grants from the FP7 Framework Marie Curie Actions CIG (methDRE) to M. Wiench and The Royal Society (RS-9-10-2012) to M. Wiench. H.J. Gleneadie received a joint University of Birmingham and Medical Research Council fellowship.

Author contributions

H.J.G., M.W., F.L.K., B.A.S., and P.R.C designed the project. H.J.G., M.W., and F.L.K. analysed the data. H.J.G. and M.W. performed most of the experiments in HNSCC cell lines. A.B. and P.G. performed most of the experiments in AML cell lines. N.B. designed and analysed drug synergy experiments. J.B. and N.B. performed *in vivo* experiments. Y.J. performed and analysed ROS and mitosox assays. S.C. analysed RNA-seq data. D.M.A.G. performed DSEA analysis. S.R., J.S.G. and A.M. provided essential advice and material. M.W. and H.J.G. wrote the manuscript with input from S.R., F.L.K., S.C. and H.M.

Conflict of interest

The authors declare no conflict of interest.

The paper explained

Problem

Aberrant DNA methylation patterns are common in most cancers, arise early in tumour development and are potentially reversible by hypomethylating agents. 5-aza-2-deoxycytidine (DAC) is a DNA methylation inhibitor currently approved for treatment of some haematological malignancies, however the efficacy of DAC in solid tumours remains unclear. We do not fully understand why some cancers have limited sensitivity to DAC, why the response is highly heterogenous in solid tumours and how, after the initial response, the cancer cells develop resistance to DAC. In addition, the clinical application of DAC is limited by cytotoxic side effects, mostly due to DNA damage. Therefore, the ability to lower the DAC dose to clinically safe levels would be of great clinical significance. Here we hypothesized that anti-tumour DAC effects can be enhanced through interaction with other, non-oncological, compounds. The DAC potential was initially tested in HNSCC, a cancer which originates from squamous epithelial lining of oral cavity and pharynx. HNSCC has very poor survival rates despite improved surgical and chemotherapy-based regimes, therefore new therapies are being sought for.

Results

By screening a panel of 100 off-patent drugs, paracetamol (acetaminophen) was identified to synergistically increase DAC response in HNSCC cells. This finding has been corroborated in HNSCC xenograft mouse models and in AML cells. The mechanisms for such interaction are multifactorial. Firstly, DAC upregulates cyclooxygenase COX-2-PGE$_2$ pathway inadvertently increasing cancer cell survival. This is counteracted by paracetamol, a known COX-2 inhibitor. Secondly, the combined treatment mimics the effects of paracetamol overdose and depletes GSH stores, which is followed by an increase in oxidative stress, both of which have the potential to restrict tumour growth and improve efficacy of anti-cancer drugs. We further show that cells are unable to protect themselves from this increased oxidative stress as both anti-oxidant and thioredoxin responses are impaired by DAC treatment. Ultimately, the combined DAC/paracetamol treatment leads to decreased cell metabolism and proliferation, as well as enhanced differentiation of cancer cells.

Impact

The study provides evidence that the anti-tumour potential of DAC can be enhanced by the presence of paracetamol applied in non-toxic doses, commonly used to alleviate pain. This has several clinical implications. The addition of paracetamol could allow for DAC dose reduction hence reducing DNA damage-related side effects and widening its clinical usability. In addition, the impact of anti-pain implementation in cancer therapy has not been systematically investigated and is not considered when planning treatments and clinical trials. Here we highlight the fact that the commonly used drug, available as over-the-counter medicine and often self-medicated by patients, can change the cancer cell response to a chemotherapeutic. The molecular pathways affected by the DAC/paracetamol co-treatment are shown to target vulnerabilities present across many tumour types, indicating wider potential for the drug combination. In HNSCC, DAC/paracetamol combination could have therapeutic potential in tumours susceptible to DACinduced DNA demethylation.

Figure legends

Figure 1 - DAC therapeutic potential in HNSCC cell lines can be synergistically increased by co-treatment with paracetamol.

A. Dose dependent cell viability in response to 96h DAC treatment in four HNSCC cell lines and normal oral keratinocytes (HOK). Grey box indicates clinically relevant doses.

B. Global levels of DNA methylation (5mC) analysed by DNA dot blot in HNSCC cell lines +/- 500 nM DAC, 96h. The data were normalized to methylene blue staining and an example of 5mC dot blot is shown below. For each replicate the results for 1 and 0.5 µg of DNA were averaged.

C. Scatterplot of the results from DAC sensitizing screen (full data in Appendix Fig S1) using 100 off patent drugs (Drug Library FMC1). Y axis represents increase in sensitivity, grey area shows ± 2SD.

D. Cell viability in HNSCC cell lines and HOK cells treated for 96h with 132.3 µM paracetamol (Para), 601.7 µM valproic acid (VPA) and 323.4 µM zinc acetate (Zinc) +/- 500 nM DAC. The results are independent confirmation of data shown in Fig 1C and Appendix Fig S2. Dotted lines show the effect of 500 nM DAC alone.

E-G. Synergy was determined using Chou-Talalay method. Cell viability data for VU40T (**E**) and HN12 (**F**) cells treated with fixed Cmax titrations of DAC (500 nM), paracetamol (132 µM) or DAC + paracetamol were used to calculate combination index (CI) (**G**) and dose reduction index (DRI) (Appendix Tables S1 and S2). CI <1 denotes synergy.

DAC-sensitive cell lines are shown in reds, resistant – in greys, HOKs – in blue.

Data information: In A-C and E-G n=3, in D n=7. In B statistical analysis was performed by paired two-tailed *t* tests. In D, for each cell line a matched One-Way ANOVA with Dunnett's multiple comparison testing was used to compare DAC to DAC+drug. A separate paired *t* test was applied to compare no treatment with DAC. Values are displayed as means ± SEM. Significant p values are shown.

Source data are available for this figure.

Figure 2 - DAC-paracetamol combination enhances the effects of DAC on cell metabolism, proliferation and basal cell phenotype.

A. Response of VU40T cells to paracetamol, DAC or both as indicated (representative images from 3 experiments). Left: Giemsa-Jenner staining (4x magnification); middle: Ki67 immunostaining; right: Annexin V and propidium iodine (PI) FACS analysis (healthy cells (bottom left); early apoptotic cells (bottom right); late apoptotic cells (top right) and necrotic cells (top left)).

B. Proportion of VU40T cells undergoing cell death assessed by FACS following Annexin V and PI staining (examples shown in (A)).

C. Percentage of VU40T nuclei with indicated numbers of γH2AX foci following indicated treatments.

D. Cell cycle distribution in VU40T cells following indicated treatments.

E. Proportion of VU40T nuclei positive for Ki67 immunostaining after indicated treatments (examples in (A)).

F. qRT-PCR for *TP63* and *KRT5* normalized to *ACTB* in VU40T cells treated as indicated.

G. qRT-PCR for *IVL* normalized to *ACTB* in VU40T cells treated as indicated.

H. Venn diagrams showing overlap for the most upregulated (left, $\log_2$ fold change ≥1) and down-regulated (right, $\log_2$ fold change ≤-1) genes as compared to vehicle control.

I. REVIGO-consolidated GO-BP pathways enriched for each indicated treatment. Top 5 terms from each group are included. Heatmap displays category scores as $\log_{10}$ p-value.

Data information: Unless stated otherwise all treatments were for 96h with 500 nM DAC and 132.2 μM paracetamol. In B-D and F-G n=3, in E n=6. In B and D: Two-Way ANOVA with Dunnett's correction was preformed to compare Ctrl with treatments and a separate Two-Way ANOVA with Sidak's was performed to compare DAC with combined treatment. In C: matched Two-Way ANOVA with Tukey's multiple comparison test was used to compare the

distribution of foci number between each treatment group. In E-G: matched One-Way ANOVA with Dunnett's multiple comparison test was used to compare all treatments to Ctrl. A separate paired two-tailed *t* test was used to compare DAC to DAC+paracetamol. Values are displayed as means ± SEM.

Figure 3 - DAC treatment specifically activates COX-2-PGE$_2$ pathway.
A. qRT-PCR for *PTGS2* in HNSCC cell lines treated as indicated. Data are shown as relative to vehicle control (Ctrl=1).
B. PTGS2 protein levels in VU40T and SCC040 cells. Graph represents the data from three experiments normalized to Lamin A/C.
C. PGE$_2$ concentration assessed by ELISA in media of VU40T and SCC040 cells treated for 96h as indicated. The data are normalized to cell viability.
D. qRT-PCR for PGE$_2$ receptors, *PTGER1* and *PTGER2,* shown as in (A).
E. Schematic of COX-2-PGE$_2$ pathway with confirmed DAC effects shown in red. COX-2 catalyzes the conversion of arachidonic acid into prostaglandin H$_2$. This is then converted into prostaglandin E$_2$ which exerts its effects through G-protein coupled receptors. The full COX-2 pathway is shown in Appendix Fig S3B.
F. COX-2-PGE$_2$-related gene expression across 522 HNSCC tumour samples. Heatmap created in cBioPortal based on provisional HNSCC cohort (TCGA). % indicates fraction of tumours with alterations.
G. Drug Perturbation Signature for genes of the COX-2-PGE$_2$ pathway for Decitabine (DAC) and paracetamol (Para) using cMap dataset.
Data information: Unless stated otherwise all treatments were for 96h with 500 nM DAC and 132.2 µM paracetamol and performed in three biological replicates. In A, B and D: for each cell line a matched One-Way ANOVA with Dunnett's multiple comparison test was used to compare all treatments to Ctrl; a separate paired two-tailed *t* test was used to compare DAC to DAC+paracetamol. In C: for each cell line an ordinary One-Way ANOVA with Sidak's correction was applied to compare treated groups with Ctrl. Values are displayed as means ± SEM. Only significant p-values are shown.
Source data are available for this figure.

Figure 4 - Combined DAC-paracetamol treatment mimics the effects of paracetamol overdose, depletes glutathione levels and leads to oxidative stress.
A. Cell viability in VU40T and HN12 cells treated for 96h with 132.3 µM paracetamol, 10

µM valdecoxib or 193.9 µM ibuprofen +/- 500nM DAC. Dotted line shows the effect of DAC alone.

B. PGE_2 concentration in the media of VU40T cells treated with DAC with or without paracetamol or valdecoxib shown as relative to either vehicle control (left) or DAC only treatment control (right).

C. Schematic of paracetamol overdose. A fraction of paracetamol is converted into the toxic metabolite NAPQI, and detoxified by GSH. When paracetamol is taken in excess GSH stores deplete and NAPQI accumulates.

D. qRT-PCR for *CYP2E1* in HNSCC cell lines treated as indicated and shown relative to vehicle control (Ctrl=1).

E. CYP2E1 protein levels in VU40T and SCC040 cells. The graph represents the data from three experiments normalized to Lamin A/C.

F. Glutathione (GSH) levels in VU40T cells treated with DAC, paracetamol or both, normalized to cell viability. 1 mM paracetamol was used as a positive control.

G. NAC rescue experiment. VU40T cells were pre-treated for 48h with 2.5 mM NAC followed by 96h of DAC, paracetamol or combined treatments and cell viability was assessed in comparison to control without NAC (left panel).

H. Levels of intracellular ROS (DCFDA staining and FACS) in VU40T cells treated for 72h as indicated. Upper figure: representative FACS profile; lower graph: geometric mean normalized to vehicle control.

I. Levels of mitochondrial superoxide were assessed as in (H) by MitoSOX Red staining.

J. Gene expression (RNA-seq data) of direct responders to oxidative stress shown as heatmap of log_2 fold change values after indicated treatments.

K. Gene expression of cystin/TXN/TXNRD1 system components shown as in (J).

L. Schematic of proposed mechanism underlying DAC-paracetamol synergy: mimicry of paracetamol overdose leading to GSH depletion and oxidative stress. The effects of DAC are shown in red and paracetamol contribution – in blue.

M. Drug Set Enrichment Analysis for DAC, DAC + paracetamol, and DAC + valdecoxib combinations. Log_{10} p-values for selected pathways of interest (KEGG or GO BP genesets) are shown as heatmap.

Data information: Unless stated otherwise all treatments were for 96h with 500 nM DAC and 132.2 µM paracetamol, n=3 for A-B, D-F, H-I and n=4 for G. In A: for each cell line a mixed-effect analysis with Dunnett's correction was used to compare DAC to DAC+Vald and DAC+Ibup samples. In B, D-F: One-Way ANOVA with Dunnett's correction was used to

compare treatments with Ctrl. In D and F: a separate paired two-tailed *t* test was used to compare DAC to DAC+paracetamol. In G: for each group (+vehicle, +NAC) a matched One-Way ANOVA with Dunnett's was used to compare treatments with Ctrls; additionally, a paired two-tailed *t* test was used to compare Ctrl with NAC. In H-I: a matched One-Way ANOVA with Dunnett's correction was used to compare all groups to Ctrl; a separate ANOVA was used to compare DAC to DAC+Para and DAC+Vald. Values are displayed as means ± SEM. Only significant p values are shown.

Source data are available for this figure.

Figure 5 - Combined effects of DAC and paracetamol on AA metabolism and HNSCC patients' survival.

A. Arachidonic acid (AA) is metabolized to eicosanoids through COX, LOX, and cytochrome P450 monooxygenase pathways. In addition to DAC limiting cancer cell growth through either activation of tumour suppressor genes (TSGs) or viral mimicry (1), it inadvertently activates COX-2 PGE$_2$ pathway (2), which is contradicted by paracetamol. DAC also upregulates *CYP2E1* which, in the presence of paracetamol, leads to glutathione depletion and ROS accumulation, both enhanced by combined treatment (3). Simultaneously, DAC downregulates transcription of genes involved in antioxidant and thioredoxin responses, preventing cancer cells from developing adaptation to oxidative stress and protection from oxidative damage (4). GSH depletion has also potential to limit the production of LOX pathway metabolites dependent on GSH transferases.

B-D. Survival curves for 392 HNSCC patients with data available in TCGA provisional cohort with or without upregulation of genes (z-score ≥2.0) involved in indicated pathway. For gene sets see Appendix Materials and Methods. Kaplan Meier Estimates were plotted using cBioPortal. (**B**) Disease/ Progression-free survival curves for genes involved in COX-2-PGE$_2$ pathway. (**C**) Disease/Progression-free survival curves for genes involved in glutathione synthesis. (**D**) Disease/Progression-free survival curves for genes involved in antioxidant (left) and thioredoxin (right) responses.

Figure 6 - *In vivo* potential of DAC-paracetamol combination.

A-B. DAC-paracetamol synergy assessed in FaDu cells (as in Fig 1E-G); cell viability after treatment with fixed Cmax titrations of drugs (**A**) resulting in combination index (CI) shown in (**B**). CI value less than 1 denotes synergy.

C. Tumour growth of FaDu cells engrafted into NSG mice (Day 1, red arrow) treated as

indicated (black arrows).

D. Days to reach tumour size of 200 mm^3. Tumours below 200 mm^3 at the end of the experiment are counted as Day 25.

E. Tumour size at Day 15 and at the end of the study.

F. qRT-PCR for *PTGS2* and *CYP2E1* expression in tumour tissue.

G. qRT-PCR for *TP63* and *KRT5* expression in tumour tissue.

Data information: In A-B n=3. Mice groups: Ctrl (n=6), paracetamol (n=6), DAC (n=5), DAC+paracetamol (n=6). In D-G: a non-matched One-Way ANOVA with Dunnett's correction was used to compare treatments with control. To compare DAC and DAC+Para an unpaired two-tailed *t* test was used. Values are displayed as means ± SEM. Only significant p values are shown.

Figure. 7 - Potential of DAC-paracetamol combined treatment in AML.

A-C. DAC-paracetamol synergy determined in AML cell lines. Cell viability of SKM-1 (**A**) and HL-60 (**B**) treated as indicated for 72h. Combination index (CI) shown in (**C**).

D. Long term effects of DAC and DAC-paracetamol combined treatment on growth rates in SKM-1 (left) and HL-60 (right) cells: 4 cycles (C1-C4, 72h treatment as indicated followed by 21 day withdrawal). Vehicle and paracetamol controls are shown for the first 10 days.

E. qRT-PCR for *PTGS2*, *PTGER1*, *PTGER2* and *PTGER3* genes in SKM-1 and HL-60 cells after indicated treatments and shown relative to control.

F. qRT-PCR for *PTGS2* shown as in (E).

G. Levels of intracellular ROS measured by DCFDA staining and FACS in SKM-1 cells treated with indicated drugs and their combinations. Vald, 10 µM valdecoxib.

H. Levels of mitochondrial superoxide assessed by MitoSOX Red staining and FACS in SKM-1 cells as in (G).

I. Proportion of SKM-1 cells undergoing cell death assessed by FACS following Annexin V and PI staining.

J. Cell cycle distribution of SKM-1 cells after of DAC, paracetamol or combined treatment.

K. qRT-PCR for *CD11b* gene in SKM-1 and HL-60 cells after indicated treatments.

L. Giemsa-Jenner staining in SKM-1 cells. Upper: 100x magnification; lower: image zoomed to approximately one cell. Vacuole formation in cytoplasm indicated by black arrow.

Data information: Unless stated otherwise all treatments were for 72h with 500 nM DAC and 132.2 µM paracetamol and n=3. In E-H and K: a matched One-Way ANOVA with Dunnett's correction was applied to compare treatments with control; in G-H a separate ANOVA was

used to compare DAC to DAC+Para and DAC+Vald. In I-J: for each group a matched Two-Way ANOVA with Dunnett's was performed to compare treatments with control; a separate Two-Way ANOVA with Sidak's was used to compare DAC with DAC+Para. Values are displayed as means ± SEM. Only significant p values are shown.

References

Abele R, Clavel M, Dodion P, Bruntsch U, Gundersen S, Smyth J, Renard J, van Glabbeke M, Pinedo HM (1987) The EORTC Early Clinical Trials Cooperative Group experience with 5-aza-2'-deoxycytidine (NSC 127716) in patients with colo-rectal, head and neck, renal carcinomas and malignant melanomas. *Eur J Cancer Clin Oncol* 23: 1921-1924

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT *et al* (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 25: 25-29

Bertolini A, Ferrari A, Ottani A, Guerzoni S, Tacchi R, Leone S (2006) Paracetamol: new vistas of an old drug. *CNS Drug Rev* 12: 250-275

Brown T (2001) Dot and slot blotting of DNA. *Curr Protoc Mol Biol* Chapter 2: Unit2 9B

Bryant J, Batis N, Franke AC, Clancey G, Hartley M, Ryan G, Brooks J, Southam AD, Barnes N, Parish J *et al* (2019) Repurposed quinacrine synergizes with cisplatin, reducing the effective dose required for treatment of head and neck squamous cell carcinoma. *Oncotarget* 10: 5229-5244

Cancer Genome Atlas N (2015) Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature* 517: 576-582

Cebola I, Custodio J, Munoz M, Diez-Villanueva A, Pare L, Prieto P, Ausso S, Coll-Mulet L, Bosca L, Moreno V *et al* (2015) Epigenetics override pro-inflammatory PTGS transcriptomic signature towards selective hyperactivation of PGE2 in colorectal cancer. *Clin Epigenetics* 7: 74

Cebola I, Peinado MA (2012) Epigenetic deregulation of the COX pathway in cancer. *Prog Lipid Res* 51: 301-313

Chiappinelli KB, Strissel PL, Desrichard A, Li H, Henke C, Akman B, Hein A, Rote NS, Cope LM, Snyder A *et al* (2015) Inhibiting DNA Methylation Causes an Interferon Response in Cancer via dsRNA Including Endogenous Retroviruses. *Cell* 162: 974-986

Chou TC (2006) Theoretical basis, experimental design, and computerized simulation of synergism and antagonism in drug combination studies. *Pharmacol Rev* 58: 621-681

Dahlin DC, Miwa GT, Lu AY, Nelson SD (1984) N-acetyl-p-benzoquinone imine: a cytochrome P-450-mediated oxidation product of acetaminophen. *Proc Natl Acad Sci U S A* 81: 1327-1331

European Medicines Agency E, 2019. Dacogen, in: Agency E.M. (Ed.). https://www.ema.europa.eu/en/documents/product-information/dacogen-epar-product-information_en.pdf.

Forman HJ, Zhang H, Rinna A (2009) Glutathione: overview of its protective roles, measurement, and biosynthesis. *Mol Aspects Med* 30: 1-12

Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E *et al* (2013) Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* 6: pl1

Graham GG, Davies MJ, Day RO, Mohamudally A, Scott KF (2013) The modern pharmacology of paracetamol: therapeutic actions, mechanism of action, metabolism, toxicity and recent pharmacological findings. *Inflammopharmacology* 21: 201-232

Hedberg ML, Peyser ND, Bauman JE, Gooding WE, Li H, Bhola NE, Zhu TR, Zeng Y, Brand TM, Kim MO *et al* (2019) Use of nonsteroidal anti-inflammatory drugs predicts improved patient survival for PIK3CA-altered head and neck cancer. *J Exp Med* 216: 419-427

Jones PA, Baylin SB (2002) The fundamental role of epigenetic events in cancer. *Nat Rev Genet* 3: 415-428

Kantarjian HM, Thomas XG, Dmoszynska A, Wierzbowska A, Mazur G, Mayer J, Gau JP, Chou WC, Buckstein R, Cermak J *et al* (2012) Multicenter, randomized, open-label, phase III trial of decitabine versus patient choice, with physician advice, of either supportive care or low-dose cytarabine for the treatment of older patients with newly diagnosed acute myeloid leukemia. *J Clin Oncol* 30: 2670-2677

Khanim FL, Merrick BA, Giles HV, Jankute M, Jackson JB, Giles LJ, Birtwistle J, Bunce CM, Drayson MT (2011) Redeployment-based drug screening identifies the anti-helminthic niclosamide as anti-myeloma therapy that also reduces free light chain production. *Blood Cancer J* 1: e39

Kim YY, Lee EJ, Kim YK, Kim SM, Park JY, Myoung H, Kim MJ (2010) Anti-cancer effects of celecoxib in head and neck carcinoma. *Mol Cells* 29: 185-194

Kobrinsky NL, Hartfield D, Horner H, Maksymiuk A, Minuk GY, White DF, Feldstein TJ (1996) Treatment of advanced malignancies with high-dose acetaminophen and N-acetylcysteine rescue. *Cancer Invest* 14: 202-210

Kurtova AV, Xiao J, Mo Q, Pazhanisamy S, Krasnow R, Lerner SP, Chen F, Roh TT, Lay E, Ho PL *et al* (2015) Blocking PGE2-induced tumour repopulation abrogates bladder cancer chemoresistance. *Nature* 517: 209-213

Lamb J, Crawford ED, Peck D, Modell JW, Blat IC, Wrobel MJ, Lerner J, Brunet JP, Subramanian A, Ross KN *et al* (2006) The Connectivity Map: using gene-expression signatures to connect small molecules, genes, and disease. *Science* 313: 1929-1935

Linnekamp JF, Butter R, Spijker R, Medema JP, van Laarhoven HWM (2017) Clinical and biological effects of demethylating agents on solid tumours - A systematic review. *Cancer Treat Rev* 54: 10-23

Liu B, Qu L, Yan S (2015) Cyclooxygenase-2 promotes tumor growth and suppresses tumor immunity. *Cancer Cell Int* 15: 106

Napolitano F, Sirci F, Carrella D, di Bernardo D (2016) Drug-set enrichment analysis: a novel tool to investigate drug mode of action. *Bioinformatics* 32: 235-241

Nieto M, Demolis P, Behanzin E, Moreau A, Hudson I, Flores B, Stemplewski H, Salmonson T, Gisselbrecht C, Bowen D *et al* (2016) The European Medicines Agency Review of Decitabine (Dacogen) for the Treatment of Adult Patients With Acute Myeloid Leukemia:

Summary of the Scientific Assessment of the Committee for Medicinal Products for Human Use. *Oncologist* 21: 692-700

Park SW, Heo DS, Sung MW (2012) The shunting of arachidonic acid metabolism to 5-lipoxygenase and cytochrome p450 epoxygenase antagonizes the anti-cancer effect of cyclooxygenase-2 inhibition in head and neck cancer cells. *Cell Oncol (Dordr)* 35: 1-8

Posadas I, Vellecco V, Santos P, Prieto-Lloret J, Cena V (2007) Acetaminophen potentiates staurosporine-induced death in a human neuroblastoma cell line. *Br J Pharmacol* 150: 577-585

Qin T, Jelinek J, Si J, Shu J, Issa JP (2009) Mechanisms of resistance to 5-aza-2'-deoxycytidine in human cancer cell lines. *Blood* 113: 659-667

Raghunath A, Sundarraj K, Nagarajan R, Arfuso F, Bian J, Kumar AP, Sethi G, Perumal E (2018) Antioxidant response elements: Discovery, classes, regulation and potential applications. *Redox Biol* 17: 297-314

Ray PD, Huang BW, Tsuji Y (2012) Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cell Signal* 24: 981-990

Roulois D, Loo Yau H, Singhania R, Wang Y, Danesh A, Shen SY, Han H, Liang G, Jones PA, Pugh TJ *et al* (2015) DNA-Demethylating Agents Target Colorectal Cancer Cells by Inducing Viral Mimicry by Endogenous Transcripts. *Cell* 162: 961-973

Saba NF, Choi M, Muller S, Shin HJ, Tighiouart M, Papadimitrakopoulou VA, El-Naggar AK, Khuri FR, Chen ZG, Shin DM (2009) Role of cyclooxygenase-2 in tumor progression and survival of head and neck squamous cell carcinoma. *Cancer Prev Res (Phila)* 2: 823-829

Simard EP, Torre LA, Jemal A (2014) International trends in head and neck cancer incidence rates: differences by country, sex and anatomic site. *Oral Oncol* 50: 387-403

Smirnov P, Safikhani Z, El-Hachem N, Wang D, She A, Olsen C, Freeman M, Selby H, Gendoo DM, Grossmann P *et al* (2016) PharmacoGx: an R package for analysis of large pharmacogenomic datasets. *Bioinformatics* 32: 1244-1246

Squier CA, Kremer MJ (2001) Biology of oral mucosa and esophagus. *J Natl Cancer Inst Monogr*: 7-15

Steinmann K, Sandner A, Schagdarsurengin U, Dammann RH (2009) Frequent promoter hypermethylation of tumor-related genes in head and neck squamous cell carcinoma. *Oncol Rep* 22: 1519-1526

Stresemann C, Lyko F (2008) Modes of action of the DNA methyltransferase inhibitors azacytidine and decitabine. *Int J Cancer* 123: 8-13

Supek F, Bosnjak M, Skunca N, Smuc T (2011) REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One* 6: e21800

Telorack M, Meyer M, Ingold I, Conrad M, Bloch W, Werner S (2016) A Glutathione-Nrf2-Thioredoxin Cross-Talk Ensures Keratinocyte Survival and Efficient Wound Repair. *PLoS Genet* 12: e1005800

Traverso N, Ricciarelli R, Nitti M, Marengo B, Furfaro AL, Pronzato MA, Marinari UM, Domenicotti C (2013) Role of glutathione in cancer progression and chemoresistance. *Oxid Med Cell Longev* 2013: 972913

Tsai HC, Li H, Van Neste L, Cai Y, Robert C, Rassool FV, Shin JJ, Harbom KM, Beaty R, Pappou E *et al* (2012) Transient low doses of DNA-demethylating agents exert durable antitumor effects on hematological and epithelial tumor cells. *Cancer Cell* 21: 430-446

Wang D, Dubois RN (2010) Eicosanoids and cancer. *Nat Rev Cancer* 10: 181-193

Wu L, Shi W, Li X, Chang C, Xu F, He Q, Wu D, Su J, Zhou L, Song L *et al* (2016) High expression of the human equilibrative nucleoside transporter 1 gene predicts a good response to decitabine in patients with myelodysplastic syndrome. *J Transl Med* 14: 66

Wu YJ, Neuwelt AJ, Muldoon LL, Neuwelt EA (2013) Acetaminophen enhances cisplatin- and paclitaxel-mediated cytotoxicity to SKOV3 human ovarian carcinoma. *Anticancer Res* 33: 2391-2400

Yoh K, Prywes R (2015) Pathway Regulation of p63, a Director of Epithelial Cell Fate. *Front Endocrinol (Lausanne)* 6: 51

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

B

COX2 pathway

Logrank Test P-Value: 5.693e-3

Disease/Progression-free Survival (%)

n=274
Median: 76,15

n=118
Median: 35,61

Months Disease/Progression-free

C

Glutathione synthesis

Logrank Test P-Value: 2.763e-3

Disease/Progression-free Survival (%)

n=272
Median: 76,15

n=120
Median: 28,78

Months Disease/Progression-free

D

Antioxidant response
genes down-regulated by DAC

Logrank Test P-Value: 0.395

Disease/Progression-free Survival (%)

n=322
Median: 67,74

n=70
Median: 61,07

Months Disease/Progression-free

Thioredoxin pathway
genes down-regulated by DAC

Logrank Test P-Value: 3.639e-4

n=332
Median: 71,22

n=60
Median: 17,74

Months Disease/Progression-free

▪ HNSCC cases with genes upregulated   ▪ HNSCC cases without genes upregulation

Figure 6

Figure 7

Supplementary information for:

A common analgesic enhances the anti-tumour activity of 5-aza-2'-
deoxycytidine through induction of oxidative stress

Hannah J. Gleneadie, Amy Baker, Nikolaos Batis, Jennifer Bryant, Yao Jiang, Samuel J.H. Clokie, Hisham Mehanna, Paloma Garcia, Deena M.A. Gendoo, Sally Roberts, Alfredo A. Molinolo, J. Silvio Gutkind, Ben A. Scheven, Paul R. Cooper, Farhat L. Khanim, Malgorzata Wiench.

The Appendix contains the following supplementary information:

SI Materials and Methods

SI References

SI Tables S1 to S7

SI Figures S1 to S7

SI Materials and Methods

Detailed information is provided for the following methods and procedures: DNA dot blotting, Western blot analysis, immunofluorescence analysis, apoptosis assay, ROS and mitosox assessment, qRT-PCR, RNA-seq data analysis, mouse xenograft study, AML cells' growth rate assessment, TCGA data analysis and drug perturbation signatures.

Dot blotting of DNA

DNA was extracted using the DNeasy Blood and Tissue Kit (Qiagen) as described by the manufacturer. Dot blotting was performed as described in Current Protocols in Molecular Biology using a titration of DNA (Brown, 2001). Dried membranes (Nytran Supercharge Positively Charged Nylon Membrane (Fisher Scientific)) were blocked in 5% low-fat milk in PBS for 1h and incubated with primary antibody (anti-5mC, Cell Signalling (28692S), 1:750 dilution) in 1% bovine serum albumin (BSA) at 4°C overnight. Membranes were incubated with secondary antibody (Santa Cruz, Cat# sc-2004, 1:5000 dilution) for 2h at 4°C. Membranes were developed using 1:1 Amersham ECL western blotting reagent (GE Healthcare) to SuperSignal West Femto Chemiluminescent Substrate (Thermo Fisher Scientific) and detected by autoradiography. Blots were stained for total DNA by 30min incubation in 0.04% methylene blue (Sigma-Aldrich). ImageJ software (Schindelin *et al*, 2012) was used to quantify the intensity of the 1 μg and 0.5 μg dots and each 5-methylcytosine dot was normalised against its methylene blue counterpart. For each biological replicate an average was taken from the normalised 1 μg and 0.5 μg dots and the technical replicates for use in further analysis.

Western Blot analysis

Cell pellets were incubated with RIPA buffer (150mM sodium chloride, 0.5% sodium deoxycholate, 0.1% sodium dodecyl sulphate, 50mM Tris-HCl pH8) supplemented with protease inhibitors (Roche) for 45min then centrifuged at 17,000g for 10min and the supernatant was retained. Proteins were denatured in Laemmli buffer (4% SDS, 20% glycerol, 10% 2-mercaptoethanol, 0.004% bromophenol blue and 0.125 M Tris HCl), heated to 95°C for 5min and separated on a 10% SDS-PAGE gel before transferring onto a nitrocellulose membrane (Amersham Hybond ECL, GE Healthcare) using BioRad transfer apparatus. The protein transfer was confirmed with Ponceau S staining (SigmaAldrich).

Membranes were blocked in 20% milk in PBST for 1h following primary antibody incubation (COX-2: Abcam, Cat# ab15191, 1:1000 dilution, 1h at room temperature in 1%

BSA; CYP2E1: Abcam, Cat# ab28146, 1:2500 dilution, overnight 4°C in 5% milk; lamin A/C: Santa Cruz, Cat# sc-20681, 1:10,000, 1h at room temperature in 5% milk). Membranes were incubated with secondary antibody (anti-rabbit-IgG-HRP Santa Cruz, Cat# sc-2004, 1:5000 dilution) for 2h and developed as described for DNA dot blotting. ImageJ software was used to quantify the intensity of each band and bands of interest were normalised against the corresponding lamin A/C bands.

Immunofluorescence analysis

Immunofluorescence analysis was undertaken as described in (Roulois *et al*, 2015). For γH2AX staining, cells were first incubated with ice-cold pre-extraction buffer (10 mM PIPES pH 6.8, 300 mM sucrose, 20 mM NaCl, 3 mM $MgCl_2$ 0.5% Triton X-100) for 7min. For both γH2AX and Ki67 staining the cells were fixed in methanol for 15min, followed by 1min incubation with ice-cold acetate. Samples were blocked with 1% BSA in PBS for 1h at 4°C following incubation with the primary antibodies overnight (Ki67 (Abcam, Cat# ab15580) 1:1000; γH2AX (phospho S139) (Abcam, Cat# ab2893) 1:1000). Coverslips were incubated with secondary antibody (Jackson ImmunoResearch, Donkey anti-rabbit Alexa Fluor 488 cat# 711-545-152, 1:250) for 1h at room temperature, washed with PBS before drying and mounting using ProLong Gold Antifade Mountant with DAPI (Fisher Scientific). Cells were stored in the dark for 3 days prior to imaging on a Zeiss 780 Zen confocal microscope. For Ki67 staining, ImageJ was used to count percentage of positive cells in triplicate images from each biological replicate. For γH2AX staining, the number of γH2AX per nuclei was counted manually across all focal planes. 100 nuclei were analysed in each biological replicate.

Apoptosis assay

Following flow cytometry for cells stained with annexin V and propidium iodide (PI), the cells were gated into four populations and counted: healthy cells which were positive for neither PI nor annexin V; necrotic cells stained only with PI; early apoptotic cells stained only with annexin V and late apoptotic cells stained with both PI and annexin V. The proportion of cells in each quadrant after 48h, 72h, 96h of treatment was counted and compared to the vehicle control.

Assessment of reactive oxygen species (ROS) and mitochondrial superoxide (mitosox)

MitoSOX Red (Molecular Probes, Cat# M36008, ThermoFisher Scientific) was used to assess mitosox. PBS-washed cells were resuspended in 200 µl PBS (37°C) containing 5 µM

MitoSOX Red, incubated at 37°C for 10 min and then analysed by flow cytometry (emission wavelength 580 nm). Reactive oxygen species (ROS) were measured using carboxy-H_2DCFDA (5-(and-6)-carboxy-2',7'-dichlorodihydrofluorescein diacetate, Cat# C369, Invitrogen). 10 µM carboxy-H_2DCFDA was added to 0.5 ml cell suspension for 45 min, cells were washed with PBS and resuspended in 200 µl PBS (37°C). Resuspended cells were analysed by flow cytometry (emission wavelength 517-527 nm) with BD FACSCalibur and BD Cell Quest software. The geometric mean was calculated for each sample and experimental condition.

Real-Time quantitative PCR (qRT-PCR)

The primers used in qRT-PCR are listed in Appendix Table S7. Relative RNA values were normalised against the concentration of the purified cDNA as measured by Qubit dsDNA High Sensitivity Assay kit (Thermo Fisher Scientific). This method of normalization was treated as an additional step to the normalization to *ACTB* expression because the DAC treatments can lead to decrease of RNA yield and changes in housekeeping gene expression. Each sample was examined at least in triplicate. PCR product specificity was confirmed by a melting-curve analysis.

RNA sequencing

Quantification of the RNA sequencing data was performed according to the latest recommended pipeline as defined in the DeSeq2 software. A count for each gene was calculated using the reference-free aligner Salmon (Patro *et al*, 2017) and the resulting count table was processed using DeSeq2 (Love *et al*, 2014) to compare treatment groups. There were 1200 upregulated (log2 fold change ≥1) and 937 downregulated (log2 fold change ≤-1) genes, further divided into six groups: i) genes up-regulated by paracetamol (n=151); ii) genes down-regulated by paracetamol (n=127); iii) genes up-regulated by DAC (n=658); iv) genes down-regulated by DAC (n=293); v) genes up-regulated by combined treatment (n=915); and vi) genes down-regulated by combined treatment (n=721). Each group was subjected to gene ontology (GO) analysis using the Gene Ontology Consortium software (Ashburner *et al*, 2000) followed by removal of redundant terms using REVIGO (Supek *et al*, 2011). The top 5 terms with the Log_{10} p-value ≤3.0 from each group were collated; none of the GO terms from the gene set downregulated by paracetamol fulfilled this condition. The Log_{10} p-values were used to generate heatmap.

Mouse xenograft study – detailed protocol

To examine the toxicity and anti-tumour efficacy of DAC and paracetamol, we utilised male NOD/SCID/gamma (NSG) mice (Charles River) which were kept in 12 hour light and 12 hour dark cycles in individually ventilated cages. Mice were maintained on mouse feed and water, ad libitum, and were at least 6 weeks old at the start of the treatments.

Toxicity studies with 3 animals per single treatment and combination treatment were used to identify safe doses for use in subsequent efficacy study: 0.4 mg/kg DAC and 1000 mg/kg paracetamol. In the toxicity studies the drugs were given 5 days a week for three weeks, first for DAC and paracetamol separately and then in combination.

To assess the efficacy, 24 male NSG mice were implanted with $5x10^6$ FaDu HNSCC cells suspended in serum-free medium and injected subcutaneously into the right flank. Tumours were given three days to become established, at which point mice were randomly allocated into four treatment groups, each group containing 6 animals, as follows: (1) 0.4 mg/kg DAC dissolved in PBS via intraperitoneal (IP) injection on a 5 day on, 2 day off regimen from day 4 onwards; (2) 1000 mg/kg paracetamol dissolved in PBS given via oral gavage following the same regimen; (3) DAC plus paracetamol as above; (4) control (PBS) given both through the oral gavage and IP injection. Animals were monitored daily for signs of ill health and the tumours were measured. Tumour measurements were taken using callipers and volumes calculated using the formula L x W^2. Mice were culled once tumours reached a maximum of 1250 mm^3; tumours became ulcerated; when animals showed signs of ill health; or at the completion of the study. One mouse from group 1 had to be excluded on Day 10 due to dosing error. When displaying tumour growth over time, to gain an understanding of efficacy over a longer period of time, data were plotted until at least 4 animals per group remained; when fewer animals remained in a group, no more data were plotted. The tumour growth is also shown as time to reach volume of 200 mm^3; tumours smaller than 200 mm^3 at the end of the study (day 25) were counted as 'Day 25'. The difference in tumour size for each group was shown at day 15 (Control n=6, paracetamol n=6, DAC n=5, DAC+paracetamol n=6) and at the end of the study (day 25: DAC n=3 and DAC+paracetamol n=5).

Tumours were excised and snap frozen for subsequent analysis. Tumour tissue was available from 6 animals in control and paracetamol-treated groups, 4 animals from DAC-treated and 5 animals from DAC+paracetamol treated groups. Approximately 10-20mg tissue was pulverized and total RNA extracted as described before (see: Real-time quantitative PCR).

Long treatment of AML cells and growth rate assessment

SKM-1 and HL-60 cells were subjected to a repeated cycle of treatments. The cells were plated in 6-well plates at 0.5×10^6 cells/ml and 0.375×10^6 cells/ml for SKM-1 and HL-60, respectively, treated for 72h after which the cells were counted and re-plated without drugs at original concentrations. The withdrawal period lasted 21 days during which the cells were passaged twice a week. The treatment cycle was repeated four times and the cells were counted at each passage using trypan blue (Sigma-Aldrich) staining and haemocytometer cell counting. Growth rate was calculated using the following equation $Gr = \ln(N(t)/N(0))/t$, where Gr = growth rate, $N(t)$=number of cells at time t, $N(0)$=number of cells at time 0, and t=time(hours).

TCGA data

The following data were retrieved from The Cancer Genome Atlas (TCGA) via cBioPortal for Cancer Genomics (Cerami *et al*, 2012; Gao *et al*, 2013): (i) expression and genetic alterations in *TP63* gene available for 496 patients (Appendix Fig S2); (ii) percentage of samples for each cancer type with mRNA Expression z-score threshold ± 2.0 (RNA Seq V2 RSEM) for all cancers with available TCGA provisional data (Appendix Tables S5 and S6); (iii) clustered gene expression heatmap for genes in COX-2 pathway in HNSCC tumours (520 patients, Fig. 3F); (iv) Logrank test p-values for overall survival and/or disease/progression-free survival for all cancers with available TCGA provisional data in indicated set of genes (expression z-score threshold ± 2.0), (Appendix Tables S5 and S6, Fig. 5B-D, Appendix Fig. S6); (v) Kaplan-Meier curves for HNSCC (520 patients) disease-progression free survival based on gene expression alterations in indicated sets of genes. The following gene sets have been used: COX-2 pathway (*PTGS2*, *PTGES*, *PTGES2*, *PTGES3*, *PTGER1*, *PTGER2*, *PTGER3*, *PTGER4*; Fig. 5B); Glutathione synthesis (*GSS*, *GCLC*, *GCLM*, *GGCT*, *OPLAH*, *GSR*; Fig. 5C); Antioxidant response genes down-regulated by DAC (*PRDX1*, *PRDX6*, *SRXN1*, *UGT1A1*, *NQO1*, *FTL*, *AKR1C2*; Fig. 5D); Thioredoxin pathway genes down-regulated by DAC (*TXN2*, *SLC7A11*, *TXN*, *TXNRD1*; Fig. 5D); COX-2-PGF$_2$, -PGI$_2$, -PGD$_2$, -TBX$_2$ pathways (*PTGS2*, *PTGDR2*, *PRXL2B*, *PTGDS*, *PTGIS*, *PTGIR*, *PTGFR*, *TBXA2R*, *TBXAS1*; Appendix Fig. S6A); LOX pathway (*ALOX5*, *ALOX15*, *ALOX15B*, *ALOX12*, *LTA4H*; Appendix Fig. S6B).

Drug perturbation signatures

Drug perturbation signatures were downloaded for the BROAD Connectivity Map dataset (CMAP) using the PharmacoGx package (version 1.14.0) (Smirnov *et al*, 2016) in R. cMap drug perturbation signatures represent transcriptionally profiled, drugs-treated cancer cell

lines involving 11,833 genes and 1,309 drugs across 5 cancer cell lines (breast cancer MCF7 and ssMCF7, pancreatic cancer PC3, melanoma SKMEL5 and AML HL-60). Precomputed signatures for CMAP were available for 1288 drugs; the signatures are calculated using a linear regression model adjusted for treatment duration, cell line identity, and batch, as described previously (Smirnov *et al.*, 2016). Heatmaps of drug perturbation signatures (transcriptional profiles) for Decitabine (DAC), paracetamol (Para), and Valdecoxib were plotted using ggplots package (version 3.0.1.1). All analyses have been conducted using the R statistical software (version 3.6.0); listed software dependencies are available on the Comprehensive Repository R Archive Network (CRAN) or Bioconductor (BioC).

SI References

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT *et al* (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 25: 25-29

Brown T (2001) Dot and slot blotting of DNA. *Curr Protoc Mol Biol* Chapter 2: Unit2 9B

Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E *et al* (2012) The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov* 2: 401-404

Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E *et al* (2013) Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* 6: pl1

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15: 550

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C (2017) Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods* 14: 417-419

Roulois D, Loo Yau H, Singhania R, Wang Y, Danesh A, Shen SY, Han H, Liang G, Jones PA, Pugh TJ *et al* (2015) DNA-Demethylating Agents Target Colorectal Cancer Cells by Inducing Viral Mimicry by Endogenous Transcripts. *Cell* 162: 961-973

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B *et al* (2012) Fiji: an open-source platform for biological-image analysis. *Nat Methods* 9: 676-682

Smirnov P, Safikhani Z, El-Hachem N, Wang D, She A, Olsen C, Freeman M, Selby H, Gendoo DM, Grossmann P *et al* (2016) PharmacoGx: an R package for analysis of large pharmacogenomic datasets. *Bioinformatics* 32: 1244-1246

Supek F, Bosnjak M, Skunca N, Smuc T (2011) REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One* 6: e21800

A

B

Figure S1. Sensitivity of HNSCC cells to DAC treatment can be increased by drug combinations.

A-B. DAC sensitizing assay: SCC040 (**A**) and VU40T (**B**) cells were subjected to 96h treatment with one of a panel of 100 drugs, +/-500 nM DAC. Viability was recorded and is shown here relative to the vehicle only control cells (Drug #0 black bar). The horizontal white line shows the effect of 500 nM DAC alone. The bars far right (Drug #101) show the effect of 10 μM DAC. The sensitizing effect is observed when the combined effect of the two drugs is more effective than both DAC alone and the drug alone. The assay was performed in triplicate and error bars represent SEM.
Drug #28: zinc acetate, #29: valproic acid, #46: paracetamol.

Figure S2 - *TP63* **gene alterations in head and neck squamous cell carcinoma.**
Genomic and expression alterations in *TP63* gene in 496 tumours from the provisional TCGA cohort of
HNSCC patients. Upper graph shows genetic and expression (z-score => or =<2.0) alterations, explained in
the legend; lower graph depicts expression heatmap.

Figure S3 - COX-2 pathways are selectively affected by DAC treatment.

A. qRT-PCR for *PTGER3* and *PTGER4* genes in HNSCC cell lines treated for 96h with DAC and/or paracetamol as indicated. Data are shown as relative to vehicle control.

B. Schematic representation of the cyclooxygenase pathway. COX enzymes convert arachidonic acid (AA) into prostaglanding H_2 which is then converted to prostanoids (prostaglandins PGE_2, PGF_2, PGI_2, PGD_2 and thromboxane TXA_2) by respective synthases. Prostanoids are then recognized by G-protein coupled receptors in both autocrine and paracrine manner.

C. DAC-induced expression fold changes for genes encoding synthases and receptors involved in all cyclooxygenase pathways. Data obtained through RNA-seq in VU40T cells treated with 500 nM DAC for 96h. Black boxes indicate genes with highest upregulation. Colour-coded association with a specific pathway is shown to the right.

D. qRT-PCR for LOX pathway enzymes: *ALOX5*, *ALOX15B*, *ALOX12* and *LTA4H* in VU40T cells treated for 96h as indicated. Data are shown as relative to vehicle control.

E. Schematic of LOX pathway with confirmed DAC effects (up-regulation in red, down-regulation in green). Leukotrienes (LTs) shown in dotted boxes were below the detection levels of ELISA.

Data information: In A and D n=3, for each cell line a matched One-Way ANOVA with Dunnett's correction to compare all treatments to Ctrl. Values displayed as means +/-SEM. Only significant p-values are shown.

Figure S4. Valdecoxib does not sensitize HNSCC cells to DAC treatment.
VU40T and HN12 cell viability after 96h of treatment with indicated concentrations of Valdecoxib with or without 500 nM DAC. Dotted lines indicate viabillity at 500 nM DAC only. Mean ±SEM; n=3.

Figure S5 - DAC and paracetamol gene expression signatures share similar pathway enrichment. Drug perturbation signatures of Decitabine, paracetamol and valdecoxib, plotted for subsets of genes representing key pathways of interest that were identified from the DSEA analysis. Genes pertaining to each pathway were subsequently obtained from the corresponding genesets identified in MSigDB collection. Clustering of the drug perturbation profiles across these pathways indicates that Decitabine and paracetamol share similar drug perturbation profiles, compared to valdecoxib.

Figure S6 - COX-2 pathway alterations selectively affect HNSCC patients survival.
A. Disease/Progression-free survival curves for 392 HNSCC patients with data available in TCGA provisional cohort and gene upregulation for cyclooxygenase pathways other then PGE_2 (but including *PTGS2* gene).
B. Disease/Progression-free survival curves for 392 HNSCC patients with data available in TCGA provisional cohort and gene upregulation for LOX pathway.
Kaplan-Meier Estimates were obtained and plotted using cBioPortal.

Figure S7 - *In vivo* potential of DAC-paracetamol combination.
Mice weights recorded during the efficacy study comparing treatments with DAC alone, paracetamol alone and DAC+paracetamol combination.

SI Supplementary Tables

Table S1. DRI (Dose Reduction Index) for VU40T, combined treatment.

| Fa (%) | VU40T | | | |
| | Dose | | DRI | |
	DAC	Para	DAC	Para
10	0.01	105	0.30	10.18
25	0.16	251	1.22	7.11
50	2.26	595	**4.99**	**4.97**
75	31	1414	20.35	3.47
97	9300	9200	426	1.59

Table S2. DRI for HN12, combined treatment.

| Fa (%) | HN12 | | | |
| | Dose | | DRI | |
	DAC	Para	DAC	Para
10	0.004	186	0.49	80
25	0.06	1962	1.19	147
50	0.85	20653	**2.92**	**269**
75	11.92	217412	7.15	493
97	3630	3.55E+07	49.5	1827

Table S3. Differentially expressed genes in VU40T cells.

Increased by treatment			Decreased by treatment		
DAC	Para	DAC+Para	DAC	Para	DAC+Para
IFITM1	PTPRA	MT-ND1	STYXL1	FLOT1	MRPL15
OLR1	PHLPP1	MT-ND6	FAM172A	ZNF254	CDCA8
IFI6	INHBB	MT-ND4	ARHGEF10	AP002990.1	RRP8
LXN	TGM1	MT-ND5	MTO1	VAC14	TRAPPC5
PADI3	GPR37L1	TIMP3	PFAS	ALDH1A3	ADSL
FSTL1	GNL1	KLF15	PPIA	SBF1	ITPA
H1F0	SREK1	SERPINA1	TOX	HAUS5	RRP1
TNFSF10	MT-ATP6	MT-ATP6	PWP2	ASAH2	WIPF3
CLDN1	GPM6A	SLITRK6	MMP16	CCDC22	FLVCR1
STEAP4	ZNF302	SMARCA1	SBF1	PLD6	TMEM107
MTRNR2L1	BBS2	MUC16	EHMT2	TTC6	FABP5
GATA3	FGF23	GSN	RRM1	ERMAP	SLC7A5
FBLN1	NEGR1	OXTR	RPS21	ITPR1	CCNF
MMP13	HMGN5	CADM4	RPL22	AHNAK2	CA2
MUC20	GRAMD1C	MT-ATP8	RWDD2B	RABAC1	FAM57A
ICAM1	DHX37	MAML3	PLK1	ZNF688	LRRC45
GPRC5B	ANKRD44	MT-CYB	CHCHD4	TBC1D13	MCM5
CCL28	NDUFA13	UBALD2	POLR1B	TARBP2	SLC8A1
IGFBP7	RASSF5	MTRNR2L1	FASTKD2	WDR54	DOT1L
GNE	MT-ND4	EIF4E3	FANCD2	UBXN11	ZFAND2B
C3	AC138969.1	ALDH1A1	MVB12B	PCDH1	ALDH16A1
GBP2	MT-ATP8	MT-ND2	NME1-NME2	MAP4K2	ANKMY1
TNFAIP2	AGAP4	TP53INP2	C14ORF80	RHBDD3	TRMT112
PLA2G16	MT-ND5	GPC4	POLA2	NSL1	NDUFB6
C1S	NPHP3	MT-ND3	PHB2	PTPRE	ARHGEF10
SORBS2	DTNB	SORT1	WDR62	DSCAM	RPA1
MX1	AK9	FSTL1	THOC3	TRAF3IP3	DUSP7
AQP3	PEX5L	LMO7	UBE2C	PSD4	AP5Z1
RARRES3	PTN	TP53I11	NME1	NTPCR	FH
IGFBP4	KANK2	GATA3	MT1E	DDX39B	GINS4
GNL1	NDC1	GDPD3	PPIL1	MFSD13A	RPS6KA4
EPSTI1	AGAP2	NLRP10	PPIF	TRIM47	SNU13
IFIT1	PLAC8	PLA2G16	CASC4	XAB2	ACBD6
NLRP10	SCD	IFITM1	ZADH2	NAGS	DARS2
MT-ND1	INSIG1	IFI6	MARS2	HSPBAP1	PER2
TMOD1	MT-CYB	KLF9	RPS3A	CSNK2A3	GMNN
TP53INP2	PRRG3	ORAI2	MPDU1	ZNF235	FANCA
SMARCA1	IRX3	DNAH1	AURKB	ALOXE3	YRDC
MT-ND5	TCOF1	TIMP2	CDKN2C	SMYD3	TMPO
PTAFR	HAT1	LYPD3	MPP3	CYB561A3	IPO9
LMO7	SPATA17	KPNA7	DHRS4L2	SIK1	PA2G4
GCNT3	MT-ND2	MMP7	DLEU1	ABHD11	LAMTOR4
XAF1	ZNF571	PLSCR4	AFG1L	RAB6B	SLC2A4RG
MT-ND6	UTRN	MAF	PEF1	RFX7	HRAS
CCL5	EVI2B	ANGPT1	HERPUD1	SEC24D	C8ORF33
B3GNT7	MT-ND6	GBP2	PPIH	PHYH	KAT2A
SALL4	ARL6IP6	IL17RD	C12ORF73	ATP5J2-PTCD1	EIF3B
CADM4	CCDC191	PCDH12	MCM2	MAP3K6	RTN4IP1
TXNIP	NAV1	RARRES3	BIRC5	DUS1L	RAD18
GRHL1	KCNJ6	ABCG1	OSBPL6	ZNF514	KLHDC4
FKBP10	NEB	FMN1	CCNF	TRIM3	NCAPD2
KRTAP2-3	MT-CO3	DNAJC6	KIF23	LZTR1	DKC1
EVC2	YJEFN3	FGF23	FHOD3	PLEKHG4	TOMM22
ABCC2	PSIP1	ICAM1	MDP1	CEP170B	ECH1
DHX37	EDIL3	TMOD1	PUS1	TPRA1	PIGU
RHOF	WASHC2C	H1F0	DDT	SPHK1	CARS2
MT-ND4	GNAO1	HAP1	TUBA1C	AMTN	FUT10
LIPH	MT-ND1	MTRNR2L8	FAM173B	SQSTM1	FEZ1
PTPRR	TRAPPC2	CLDN1	DHODH	CERS2	ARMC5
ISG20	TRIM27	HEPHL1	TBCD	PI4KA	CAV1
DNMT3B	MT-CO2	RAET1E	KPTN	HAUS7	C14ORF159
ABCG1	TMEM123	PODXL	NHEJ1	HS3ST1	PDCD2L
INHBB	JDP2	PCSK5	NKX2-5	PLCD3	RNASEH2B
EFNA1	MDC1	CEACAM1	ACADVL	FASTK	DHX30
CORO1A	HEG1	EPSTI1	ADORA2B	NAT9	RAE1
LYPD3	RHBDL2	TTC9	TMEM11	MAMLD1	RPL11
CDK11B	MSMO1	TMOD2	RPS13	SKIV2L	AC015813.2
PHLPP1	C14ORF1	NPTXR	RPS4X	SERPINB2	LSM5
YJEFN3	EGLN3	YY2	EPN3	BACE2	UQCR10
IFI44L	CNTRL	LBH	PEMT	CNPY2	PPIH
ETNK2	SRRM5	EEF1A2	PMVK	MIB2	SULT1A1
CTSS	ESR1	TAGLN	RRP9	MYO18A	CWF19L1
KRT86	MNX1	PTAFR	RPA3	PWWP2B	RAB34
CXCL3	GALK1	CDK11B	RPLP0	FAXC	RRM2
FNBP1	HECTD1	LYPD5	SRM	SPOCK1	FANCD2
FGD3	CALB2	TNFSF10	RPP21	C16ORF91	POLR1A
SOX4	HMGCS1	PGM2L1	CHCHD5	MANBAL	TRAPPC12
TIMP2	SCEL	TFPI	RAN	FMNL2	MTERF3
PIGR	CLK1	TSPYL2	PRMT3	ADARB1	MRPL23
MINDY1	CYP51A1	AKAP8L	C19ORF48	UAP1L1	AKR7A2
CPED1	MED12L	CTSV	ZNF623	IMP4	DIP2A
FSBP	MMGT1	XAF1	MRPL13	CLCN5	CKS2
APOL1	CCDC14	CD74	MTERF3	FBXL15	DEF8

HEG1	CFI	CAMK2B	XAB2	MED17	MED16
MT-ATP6	ARPC4-TTLL3	GCNT3	TOMM40	SRSF8	MRPL20
MT-CYB	DMRT2	LGALS9	RPS7	OTUB2	NDUFAF1
TRIM27	MT-ND4L	IGFBP7	DNAAF5	TMEM8A	JAG2
COX6B2	MT-ND3	NAA30	SLC2A4RG	ADGRE2	GEMIN6
PFKFB4	LEAP2	LTBP2	DDB2	AC004832.3	SMPD4
HSD11B1	DZIP1L	MUC20	ATAD3B	DNER	KIF18A
SERPINA1	NEMF	TIAM2	COQ6	PORCN	WDR34
ANGPT1	BOC	TLR1	HADH	RCOR1	DNAAF5
TLR1	CFH	CCDC71L	LYRM7	ARMCX5	MACROD1
LTBP2	TSPYL2	YJEFN3	FAM207A	MRAS	XXYLT1
OAS1	ZNF821	ATP8B1	PRPF40A	SH3TC2	ESPL1
SCNN1A	ADD3	TUBA1A	BCKDHA	AC133555.5	BRCA1
AURKC	SH3YL1	ENO2	ATAD3A	ARHGEF40	ZNF202
SPP1	CBWD3	SATB1	POLD2	MIEF2	RABEPK
BATF2	C9ORF84	TPM1	RPSA	FEZ1	CRYBG3
CYB5R2	OTOGL	PBXIP1	PPAN	GPNMB	PRKAG1
ERV3-1	TMEM154	ANKRD24	POLR3H	ACADVL	LYRM4
DNAJB5	CDK11B	VPS13D	RPL6	KLHL21	CNOT6L
PCSK5	BLNK	BSPRY	ACAT1	S100A6	RACK1
APOLD1	KRI1	B4GALT6	MRPL53	RSRP1	MRPS24
MMP7	LRRCC1	MT-CO3	SLC25A14	ANKRD16	COG4
RDH10	MT-CO1	ERV3-1	DDX39A	FICD	MAD2L1
BSPRY	SLC25A27	MT-CO1	DYNC2H1	ZBTB17	NIP7
GPR160	ATP6V1G2-DDX39B	CMYA5	TRMT61A	ATG2A	DDX54
DUSP1	SCN3A	TXNDC15	DHRS4	HERPUD1	LSM6
CXCL11	FAM46A	SUGCT	HGH1	TBC1D30	COMMD4
LIPG	NRDC	ZNF708	NXT1	RAB11FIP4	ODC1
PLOD2	CCDC80	SLC16A13	ODC1	ST3GAL4	MRPS15
SORT1	FBN1	MT-CO2	SLFN11	AP5Z1	TMEM237
IFIT3	LAMC1	ZNF836	MRPL37	HYI	NME6
PLEKHA6	MPDZ	SHISA2	MRPL11	KCNMA1	DENND1B
PPM1M	FNBP1	CALB2	POLR2L	RBM15B	RPL27A
MTRNR2L8	ZCCHC8	SNED1	EEF2KMT	PLCE1	CENPM
HNF4G	MTRNR2L1	APOLD1	ADAMTS1	ANKRD28	NXT1
NECTIN4	CCDC77	ZNF506	ESPL1	IFI27L2	QSOX2
APOBEC3D	ATP8B3	LIMD2	UPP1	RAP1A	UQCRFS1
MITF	SLC6A9	NMNAT2	ATP5F1	BRD2	MINDY4
MGAT4A	NELFE	CDA	TYMS	SELENOT	POLR3B
NMNAT2	FADS1	PLOD2	TXNDC17	KIFC1	GSS
PRDM1	B4GALT6	PDGFB	RPL4	C8ORF33	ECE2
PLA2R1	FRYL	MFAP2	CTU2	EP400	GCSH
IL23A	USF3	BPGM	MRPL27	PTDSS1	ENTPD7
GAB2	RFTN2	RAB3B	CENPI	DUSP7	TRIP13
FMN1	KIAA0368	COL4A4	AC133555.5		NUDT8
SAMD9L	COL4A2	CELF6	MRTO4		CDCA5
MUC16	MEIS2	L1CAM	RFX7		AMER1
NEURL1	RASA2	LRRN1	TMEM241		DNAJA3
NFKBIZ	PIK3R1	CNGB3	BDKRB1		NCAPD3
MFAP2	COL12A1	OLFM2	TIMM10		CDT1
RND1	SEPT7	NMB	FAM185A		SSRP1
TTC9	TESMIN	TXNIP	SYBU		ZNF544
JPH2	TRAF3IP1	B3GNT7	THNSL1		TBC1D30
HERC5	KBTBD8	ATF3	RANBP1		MOCOS
NUPR1	BRSK1	PLEKHG2	DPH1		MRPS26
C15ORF48	PLEKHA4	TGOLN2	POLA1		ATP5F1
IFIT2	POLR3GL	TJP3	RTN4RL1		MARCH8
MT-ND3	TXNIP	SOX4	DDX49		ZNF138
CACNA1A	NFE2L3	SORBS2	ATP5G3		EEF2K
TDRD7	NUCB2	BVES	ZNF35		GNPDA1
PDGFB	TBC1D32	UPK2	AKAP8		CCDC150
TMEM25	MYO9B	EPPK1	CCNB1		SRA1
NTN4	LPAR6	GAB2	SIVA1		DPCD
S1PR1	GAS8	KIAA1551	AARSD1		GADD45GIP1
POU2F2	LENG8	KRT13	WDR4		MCM7
WFDC2	FAM184B	HIST1H2BD	PXMP2		PRDX2
CLIC3	FAM49A	NCCRP1	PTRH1		CPNE1
PLAC8	IQCK	MYO5C	GSTP1		RPS3
KPNA7		SREK1	AL591806.3		EPHX2
QPRT		ZNF518B	NUDT8		TONSL
GALNT6		GOLGA6L4	RPS12		MRPS27
ZNF750		TRAF1	VEPH1		GTF2I
EDN1		C3	ICOSLG		HSCB
TIMP3		MX1	PFKM		DHRS4
VASN		IL36RN	EMG1		BDKRB2
CALB2		ANOS1	BCL2		PMS2
SLC43A2		SEMA3F	HMGN2		BDKRB1
IL36RN		GABARAPL1	ING5		RPS14
TRPV4		C15ORF48	IMPDH2		NR2C2AP
VANGL2		GPM6A	NDUFA4		SRM
NMB		LIPH	MMS22L		GTF2H3
CLDN4		CCDC68	NRG1		PES1
ZEB2		GNAO1	MIPEP		ARL4D
USP18		PRRT2	RUVBL1		ALKBH3
SUN3		MARCH4	RAD51D		SLC43A3
TFPI		OLR1	TUBB		WRAP53
ABCD1		KLHL24	TLL1		DKK3
ST8SIA1		PARM1	METTL16		ZNF77

CMPK2	HSPE1-MOB4	TNPO3	RPL3
MPDZ	DUOXA1	H2AFZ	RPL7A
PAPSS2	TBX6	CARS2	LIG1
IGF2BP3	KCNN4	WDHD1	L3MBTL2
ARL11	AGO4	ATRIP	FPGT
MT-CO1	YPEL1	IL6R	CDC25A
DDX60	HOXC9	COMMD3	BOP1
TTLL3	RETREG1	POP5	POLR3E
FUT3	NIPSNAP2	ADSL	EIF3K
MT-ATP8	RNF32	RPAP1	COMMD9
MEX3A	SPP1	IAH1	WDR35
CFAP57	RUNX3	TDP1	RPL13
IL17RD	C10ORF10	TMEM161A	POLR2L
RIMS2	FBLN1	CDC25A	RPS29
PGM2L1	CTRC	POC1A	PTPRZ1
FGF11	ALDOC	EIF4A1	SLC25A20
CDA	VANGL2	POLE4	MRPS25
NXNL2	PGF	UROD	PHPT1
MT-ND2	ENPP1	ARMC6	BLM
FMNL1	VAMP2	CKMT1B	METTL16
LGALS9	BEX2	TMEM116	ATP23
YPEL3	MXRA5	TMEM14B	VPS51
INHBA	CD68	PI4KA	FAM173B
NDRG2	GPRC5B	ENOSF1	HINT2
TTN	PLA2G4C	MRPL40	MRPS34
CDH1	TMEM123	GEMIN6	MRPL43
LCN2	NECTIN4	RPTOR	DNA2
ZNF525	PTPRA	RBM14-RBM4	CHCHD6
MT-CO2	CCL5	ALOX12	ARHGAP11A
SLITRK6	DUSP1	GINS1	RPS10
SOD2	ECM1	CDH13	EME1
TSPAN13	LAT	PLA2G4A	H2AFX
SLC9A9	PARD6B	TMEM218	SKA3
VNN1	SERPING1	RNASEH2A	PPAN
TYMP	IL32	PBK	AVPI1
TNFSF13	BBS2	YBEY	PBK
PTPRA	MYO9B	PCNA	RRM1
HLA-B	STEAP4	GPR39	ATP5J
EPPK1	ITGB8	TIMM13	TRIP6
CNGB1	YPEL3	UMPS	MAN2C1
COL4A4	NUPR1	UQCC2	OIP5
IFITM2	ARGLU1	METTL8	SLC25A19
KLF9	HIST1H2BC	MRPS28	ATRAID
LOXL2	JUND	LEF1	DDX28
KRT13	MAP3K8	S100A2	RFC2
PEAR1	CAPS	STK11IP	PKP3
LRG1	ITM2C	CENPP	NDUFA9
TSPAN15	SEPT3	TRIM28	METTL8
AGO4	KIAA1462	PPA2	BCL2L13
PTPRJ	NFATC2	TUBA1B	TNPO3
IDO1	PRRG3	DCXR	H2AFV
CD38	STX11	TBC1D30	BABAM1
DDX58	CHD6	UNKL	ELOF1
ITGB8	REEP3	ARL4C	CKMT1B
LGMN	MAOA	SLC39A3	ATRIP
GSN	GPR132	ARHGEF1	ATP5O
ZNF780B	ATP12A	KIF26B	ZSCAN21
MOK	INHBA	CLUH	MAPK12
HEPHL1	ILDR1	GINS4	NME2
TVP23C-CDRT4	PAPSS2	RAB38	MYBL2
NOTCH3	FLNC	FLRT2	POC1A
ISG15	CCDC82	SLC37A2	POLD2
MTSS1	APH1B	CDK7	MLH3
LYPD5	RDH10	SNX5	C2ORF74
KIF13B	CCDC92	RRS1	RRP7A
MYO5C	MACF1	CENPO	KRT14
ARHGEF37	LIMCH1	TRAP1	MRPL13
MAP3K8	LGMN	SEPHS1	TMEM138
CYGB	AGPAT4	KITLG	PEX7
TRAF1	DYRK1B	IPO4	IFRD2
ZNF821	IFI44L	CCDC86	RRP1B
PRSS8	OAS1	NPR3	MUM1
ALDH3A1	PRSS8	UTP20	THOC1
TRANK1	SAMD9L	CDC20	NHEJ1
CDH5	LIPG	RING1	MECR
MYRF	MYH14	ZNF584	TELO2
SOCS3	IVL	PHF19	CDC25C
AC138811.2	STBD1	SLC25A5	GAPDH
ATF3	FAXDC2	BCS1L	KLHL22
TRIM38	ALDH3B2	PCDH7	LRR1
SAT1	PLEKHA6	UBE2G2	ATP5B
ITGA5	LY75	EIF3L	TMEM120B
OAS3	ZNF738	MRPS27	AGFG2
IGSF10	CACNA1A	HGSNAT	BUB1B
SHOX2	MAB21L3	PAAF1	MANEAL
LIMCH1	TMEM145	MRPL12	RPL22L1
NNAT	DLG4	ORC6	MMS19
FAXDC2	ARPC4-TTLL3	NCR3LG1	ZNF689
MT-CO3	FMNL1	EEF1AKMT1	WDR12

ZNF699	IGFL1	TTC26	FGFR2
SEMA3F	HIST2H2BE	PABPC4	EIF2B3
PPIC	ABAT	COPRS	NARS2
PXDN	NDRG2	SLC16A7	PEF1
A4GALT	NFKBIZ	XRCC3	URI1
ZNF738	KRT86	EARS2	NCAPH2
MMP10	PXDN	MFSD2A	ZNF25
MTRNR2L11	FCHSD1	ZRANB3	MCM10
CD74	SIX2	THADA	STARD3NL
A2ML1	FAM26E	DTYMK	ULK4
HAP1	IFIT1	DUS1L	KIF18B
WIPF1	DUSP5	CDK4	EXOSC8
DNAH2	A2ML1	TOE1	TMEM218
RUNX3	SHOX2	GINS2	MIPEP
TJP3	ZSCAN2	PLCE1	EXOC8
GDA	GATS	CCDC66	POLQ
ANKRD65	CORO1A	KIF4A	CCT7
AL136295.5	PRDM1	KRT5	NME1-NME2
YPEL2	TNFSF9	RBM15B	PMPCA
ENO2	MTRNR2L4	KCNJ15	PAAF1
CRABP2	HTR1D	ZNF26	HAUS4
CTSB	BHLHE41	ZBTB44	NDUFB2
ACVR2B	IL6	GSR	ATAD3A
SCARF1	A4GALT	C12ORF66	ZNF343
ZNF506	IGIP	IMPDH1	RNF123
EEF1A2	SUPT5H	CHTF18	DCUN1D2
SERPING1	SLC22A1	NCKAP5	EEF1AKMT1
GPC4	GCNT4	FAT4	ABLIM1
LRRN4	DBNDD1	NDUFAF8	SSSCA1
RAB11FIP1	CTGF	DUS3L	SRSF2
CHD6	ARID4A	WDR73	PPIA
IRAK2	N4BP3	SELENOT	RPUSD4
C10ORF10	PLD1	ERCC8	TUBB
NR4A1	LXN		REXO5
STBD1	SOX30		CIT
ZNF350	FBXO2		PTTG1
FOLR1	TTC39A		FIZ1
KBTBD8	CDH1		PABPC4
TP53I11	ZNF766		RTEL1
GPX7	HERC5		RIN1
NAA30	JDP2		CENPA
ACKR2	FLT4		PNPO
CFLAR	BNIPL		ARL4C
LBH	RSPH3		MRPL16
SGSM2	CLDN4		EMC9
CTH	KLK10		ZNF696
KANK2	ZFAT		DHRS4L2
ZNF518B	TM7SF2		DHX35
HOXA9	TNFSF13		HTD2
GMPR	RNF128		TTC19
CLSTN3	OSBP2		COPRS
PSCA	BATF2		SORD
TMEFF1	NNAT		PGAP2
PPFIA4	NEDD9		TMEM216
GIPR	ZNF681		FUS
BHLHE41	EVC2		MICB
RAB8B	CXCL14		NABP2
IL6	STIM1		ATP5SL
LLGL2	BMP4		POMK
POU2F3	ATP9B		BCL11B
ZNF365	ITSN2		SIVA1
IQCD	CYGB		CHCHD10
NEDD9	LRRC37A3		COX7B
ANGPT2	LMBR1L		CEP128
HOXC8	FGD3		PFAS
C16ORF54	INCA1		IDH3B
IGFN1	LRTOMT		UBE2T
CAVIN2	HHIPL1		METAP1D
TXNDC15	RPL39L		NOC2L
FAM26E	DNM1		AK2
LY75	CRYBG2		ITGB3BP
SIX2	SEMA5A		C16ORF59
TGM1	SULT1B1		RPS5
TRIB2	HSD11B1		L2HGDH
KLK10	MYO6		XAB2
NEBL	TNFAIP2		DTL
KCNN4	HMSD		RPL37A
KIAA1211L	MARCKSL1		SLC39A3
GDPD3	PIK3R3		CYP1A1
IL1R1	CNTN5		NDUFS2
DCAF4L1	CCDC73		TUBA1C
TIAM2	ERO1B		FAM98B
IFIH1	SMPD3		MPP3
ABCB5	KIAA1257		SUV39H1
RNFT2	EFNA1		STOML2
PARP14	MOSPD2		TACC3
NFKBIA	RORB		HS3ST3A1
OAS2	ID4		LRP8
GALC	SERPINF1		UCK2

CDKN2D		TSPAN2		RPS23
PCDHGA4		EDN1		TDP1
OXTR		RHOF		DPH1
TSPAN2		VMP1		ZNF655
DUSP5		STAT5A		CCDC86
UNC13D		MEIS3		PHF5A
ATL1		CCL28		HADH
CASC10		TDRKH		C11ORF1
RHOV		ITIH6		MED24
SLC2A12		PPP1R3B		MKS1
AGPAT4		TNFRSF10C		RPS16
FAM193B		IL2RG		NDUFAB1
APCDD1L		LGALSL		FAM173A
ERVW-1		PNRC1		ATP5G3
SYTL2		QPRT		EXD2
RNF157		TRPC1		RNF31
ZNF583		GOLM1		C6ORF141
DMPK		YPEL2		DSN1
MME		FARP1		MRPS28
AL365205.1		HHEX		TYMS
TMEM27		MME		ORC1
UBALD2		ADRA1B		NDUFS7
FGF13		ZNF468		COPS9
MT-ND4L		SERPINB7		RFC4
HYAL1		LRG1		FAM216A
DNAJC22		SIAE		RPL34
ARL14EPL		MLF1		TK1
THEMIS2		ARL14		PMF1
NFE2L3		KBTBD8		STUB1
TMEM98		TGFBR3		TRA2B
BCL2A1		RAB3IP		FAM196B
KIAA1257		TMEFF1		CCT4
BNIPL		VAV1		FEN1
PLAG1		GJB4		FAH
NPHP3		BTBD19		ZNF74
HELZ2		FNBP1L		SCLY
TMEM123		FGF11		TMEM192
CD68		FAM25A		FASTKD2
ADGRF1		PHYHIP		E2F1
IGFBP3		VGLL3		PRMT5
TNFSF9		LSM14A		MRPS17
PGPEP1		FSIP2		SMYD3
PNPLA3		IL23A		FBXO4
SLC25A37		WIPI1		NUDCD2
SLC16A4		FUT2		SLC25A15
PNRC1		SDR16C5		RPTOR
CD274		RGPD6		PSMA1
STAT1		ABCD1		DNAJC11
PKIB		TRPV4		ZNF511
ABAT		ARHGAP31		NCR3LG1
ALDOC		MMP10		ABCB6
ARHGAP27		SLC16A4		EEF1A1
HSPE1-MOB4		HOXB6		EXO1
CLDN7		TRIM38		ATP5G1
SYNGR3		SCEL		THOC3
DTNB		DNAJC22		C14ORF80
FGF14		ZNF677		SNRNP40
PLA2G4C		WDR45		FAAP24
RCAN1		VGLL1		TIMM44
GLRX		PIWIL2		NOC4L
COBL		HEG1		TBRG4
NDRG4		IL11		IPO11
CENPT		IGFBP4		HMGN2
CES3		SLCO1A2		NUBPL
RAET1L		ECE1		HSPD1
SLC12A7		TNFRSF6B		HIRIP3
GABRP		ZNF365		KIF26B
HTR1D		STAG3		RBM34
MAOA		PAPPA		RTN4RL1
ATF5		NPBWR1		LSM7
IFNLR1		TRIM2		CENPP
OSBP2		STX2		RPS27A
ATP12A		RAG1		DLST
IL24		PIK3IP1		TRAPPC9
SERPINF1		ATF7IP2		SLC16A14
VGLL3		PTTG1IP		LGALS1
LMBR1L		AC138969.1		RAD9A
C10ORF55		DDX60		IMPDH1
HOXB3		EFCAB13		TIMM8A
CXCL2		ATP2B4		SRSF7
LTBP1		MGAT4A		PIF1
LGI4		TMC8		PINX1
MYL9		AQP3		FOXM1
MAB21L3		C2CD2L		TOMM6
YY2		IRX5		NME1
FGF5		WFDC2		C7ORF50
BMP6		NTN4		AGPAT1
GOLM1		BMPR2		SETD6
PARP9		CYB5R2		TOMM40

GABARAPL1	TSPAN13	RPS2
ZC3H12A	KIAA1211L	DLEU1
CREG2	PSCA	NPRL2
DUOXA1	PDZK1	MAIP1
TMPRSS4	HOXB3	RPLP0
BPGM	SP140	TXNRD2
ALPP	PHLPP1	ZNF165
CASZ1	NEGR1	C12ORF45
AC138969.1	NREP	LPAR3
OASL	LZTS3	PHF19
PCED1B	SLC8B1	NOB1
PLSCR1	DNAJB2	PEMT
OSMR	ALPP	TMEM201
TRPC1	LRIF1	ALOX12
SWT1	MEX3A	USP47
GPR132	S1PR1	UTP20
MTRNR2L4	NEK11	CCNB2
PDZK1	PROS1	DUS3L
TNFRSF10C	ACKR2	DLX1
KRT6B	GRHL1	RPA3
FAM46B	LRRC4	TRIM7
SLC12A6	RABL2A	ACAT1
L1CAM	ZEB1	LIX1L
ELF3	SOD2	AGMAT
SPRR2E	MSC	RPL14
GATS	IFITM2	PTDSS1
TAGLN	AMZ1	CDCA2
HOXB6	POLR3GL	KRT10
PIPOX	GNE	ERVMER34-1
ACTBL2	CDC14A	USF1
AFF1	Z98749.3	RPL10A
PIK3IP1	MMAA	ARMCX5
GPR158	MYRF	GSDMD
BTBD19	HOXB8	GEMIN5
RABL2A	MYPN	PRPF19
TAB2	CCPG1	PTHLH
PIWIL2	STX3	TAF5
SYCP2L	APCDD1L	CYC1
ACE2	MAPK15	PAICS
PAPPA	RCAN1	CKMT1A
ERBB3	LCN2	SNRPD1
FBXW10	DBNDD2	NDUFAF6
SERPINB1	CPT1A	ADORA2B
IRF9	CTF1	KIF2C
ZNF223	LMBRD1	TTLL11
PARP12	PCED1B	POLR1C
PADI1	FILIP1L	NPM3
ALPPL2	TRIML2	MRPL13
RALGAPA2	HOXA9	GPSM2
HAPLN3	SSPN	EARS2
IL4I1	WISP3	RAD51
SLC28A3	ARL11	ACADVL
CRISPLD2	CT62	MRPL55
APOD	LEPR	RPL35A
CHD2	LIN7B	PRR5
RAB15	CASZ1	UBE2G2
LRRC4	SYTL2	NDUFC1
RBM38	GIPR	PEPD
IL32	ERBB3	CAD
COL28A1	GDNF	THNSL1
ADAP2	GCC2	ORC6
CERS4	KIF13B	MCM4
LRRC37A3	HYAL1	PPA2
IL36G	VLDLR	GSTO2
PLEKHA4	ITM2B	POLD1
WNT4	SNAP23	GCDH
DNAJC6	SLC12A7	SFXN2
PARD6B	CCDC62	MECOM
ECM1	ADA2	RPAP1
TMEM97	ADCY9	RPS3A
SPINK9	GGT1	RAB38
ARPC4-TTLL3	LEPROT	RPL6
PPP1R3B	CHST2	PIGL
CTC1	KLK7	HMBS
ZNF626	RNF19A	MT1E
CLK4	TNFRSF9	PCNA
SDR16C5	GALNT6	ADAMTS1
NPIPA5	USP53	WDR54
HOXD13	CDK11A	XRCC3
STX2	PPIC	TIMM13
FBXO39	ABCA13	SF3A3
SCARB1	ATG16L2	IMP4
IL1RL1	SLC2A12	BCS1L
PREX1	ITGA5	PRKAR1B
SKIL	CXORF36	PRMT1
DENND3	RBM5	SNF8
NEK5	PPM1H	EPN3
FBXO2	RAB11FIP1	POLE4
HIST1H2AC	AKAP13	CCDC51

ITSN2		HAT1		ERMAP
RILP		CLU		ECSIT
TCP11L2		CFAP53		EBNA1BP2
APOBR		ZNF432		IMP3
TMC8		TMEM52B		PHKA1
RTKN2		C1S		TUBE1
CEACAM1		PPP1R2		BLOC1S1
ASPHD1		CFLAR		NOP56
OPN3		MYL9		KIF22
LGALSL		RAB7A		CCT8
RPL39L		RAET1L		CCNA2
HIST1H2BC		PPFIA4		EIF2B5
ACPP		INHBB		DLL1
SLC16A13		KANK2		GLYR1
PGF		ABHD8		ITFG2
FNBP1L		COL5A1		DNPEP
VAV1		ATF5		C19ORF48
CCDC71L		COX6B2		AHSA1
ADAMTSL4		HLA-B		RNASEH2A
SAMHD1		FAM214B		S100A2
PTGS2		SPINK9		COQ3
ABHD8		FOXL2NB		RAD54L
DUOX2		ABCC2		EIF5A
ZFAT		KCNJ3		MRPS2
ECE1		KDM6B		UBE2C
KRT16		ZDHHC11		ERI3
TAGAP		GALC		GRK5
SMIM22		LENG8		MRPL37
ADCY9		LTBP1		WDHD1
IRF7		YPEL5		CDCA3
SAV1		IGFN1		POLR1E
GLIPR1L2		NEBL		H2AFZ
S100A4		STC1		UPP1
JUND		ATP13A2		RPS21
ARGLU1		CYTH1		PSMG3
HECTD1		CREBRF		TRIM34
PSG9		SLC4A5		C1ORF109
RFTN2		TFAP2E		RPLP2
CCDC144A		XKRX		TRMT1
UNC79		VASN		CIDEB
IL22RA1		CMPK2		FBL
RASA4		FAM49A		MT2A
CA13		CERCAM		RANBP1
TRIM2		ACAD11		NDUFAF5
VPS13D		CHIC2		WDR74
DOCK8		FGF14		FAAP20
SCHIP1		MFGE8		CDC45
PSTPIP2		GFPT2		CRADD
NCCRP1		ALDH3A1		LARS2
GALM		TAB3		NCAPH
MYPN		PPP1R3F		MRPL11
IFI44		DCAF4L1		TUBA1B
HBE1		IGF2BP3		PHKG2
SIRPB2		TNFSF15		SMYD5
CREB5		ADD3		NPR3
CTCFL		TDRD7		MTHFD1
NAA60		IFIH1		EIF4A1
IGFL1		F2R		WDR4
NLGN2		HIVEP1		C1QBP
CLDN16		ARL14EPL		POLR1B
SREK1		SCNN1A		CTPS1
TNS2		C9ORF72		MCM2
AMZ1		DTX3L		MTFP1
MAFF		UPK3B		DCTPP1
EFCAB10		MARCH3		CCNB1
GADL1		CSAG3		BIRC5
FSIP2		CDH5		TRAP1
ARRB2		BEND7		SKIV2L
MYADM		PTP4A3		ATAD3B
CDR2L		PADI1		RRS1
KLF15		ZFYVE1		RPL37
RABGAP1L		ZHX1-C8ORF76		PLK1
CPA4		DOK3		PCDH7
COL8A2		RHOV		RPL22
KIAA1551		COL4A2		KIAA0586
ZNF615		MLPH		PCLAF
PDE8A		ZNF615		NCLN
HIST1H2BD		USP2		GPAM
FUT6		MYADM		RPS7
ZNF714		NPTX1		NRG1
SDC4		APOBEC3D		RPS12
TMEM217		PDE1C		BUB1
BEX2		SWT1		PHB2
DPP3		UBL3		AURKB
GLIS3		PLAUR		CLUH
SHISA4		RAET1G		ZC3HC1
CREBRF		ATL1		NDUFS3
ZNF217		DHX37		MRPL22
LPCAT4		FAM102B		FJX1

HOXA5	IST1	DTYMK
ABCA1	HOXB2	PDSS1
EPHX3	MMP13	FARS2
UPK2	TYMP	TLL1
SH3D21	POM121	SLC37A2
ZSCAN2	PDZD7	PRMT7
ATF7IP2	SAT1	CNOT10
CDK11A	PTGS2	NDUFAF8
LEPR	XKR4	CENPO
CHMP5	FNDC3B	NOLC1
EPCAM	VSIG10	CENPX
DTX3L	EPN2	TRMT61B
RNF19A	PPM1N	BACE2
PLAUR	FOLR1	PSMB6
ZNF224	TMEM106B	GPHN
FRK	ABCA1	EMG1
SPATA20	PPM1J	DDX39A
SOWAHB	FOXN3	TADA1
ANKRD26	SLC16A3	SLC25A10
ZNF432	NEURL1B	EHBP1
RAG1	TEX35	CDCA7
FBP2	SERPINB1	SHCBP1
SP140	RASSF5	MCAT
SMPD3	HOXB4	UQCC2
VAMP2	N4BP2L1	SPC25
STEAP1B	CCDC186	PDF
TMCC3	PSTPIP2	HGH1
BRSK1	RHOQ	RPSA
TFAP2E	FLT1	KRBOX4
CCPG1	GPR89B	URB2
TNFAIP3	NXNL2	RRP9
TDRKH	HYAL4	PI4KA
RAB3B	MAFF	MCM6
ERO1A	EPB41	CC2D2A
ENKUR	SLCO4C1	GINS2
ZFP14	PPM1M	PLCE1
DLG4	DSC2	METTL17
MEIS3	LOXL3	CDK4
SAMD9	CD274	TRIM28
B4GALT1	GGT6	LYSMD2
CPEB3	SERPINB9	DHODH
DOK3	C7ORF57	MPHOSPH9
	MTRNR2L11	RPS4X
	RERE	MPDU1
	SLC12A2	KIF20A
	SYCP2	NUDT6
	DNAH5	SNX5
	SKIL	FAM58A
	HIST1H2AC	SLFN11
	HIST1H2BJ	TFAP4
	DOCK8	KITLG
	ZNF813	SLC16A7
	CPED1	MRPL40
	FAM214A	RBM15B
	KIF27	BLMH
	KATNAL1	MRTO4
	NPAP1	VOPP1
	ALOX5	KLHL13
	ZNF354A	FUNDC1
	PTPRJ	SLC25A5
	TMEM27	C20ORF27
	ZNF43	RFX7
	IFNLR1	MRM3
	C10ORF55	PUS1
	TRPA1	MRPL48
	SLC7A2	EEF2KMT
	LRRC56	KRT5
	VNN1	FANCB
	CALCOCO1	RPS13
	GRIK2	PLD6
	ZMYM5	GINS1
	SPECC1L-ADORA2A	NUP88
	RNF182	MRPL58
	ZSCAN30	CENPL
	ARRB2	NCEH1
	PKIA	PLA2G4A
	PARP9	METTL1
	TSPAN14	LMNB1
	L1TD1	MRPL12
	ZNF844	PFKM
	RBP7	GSR
	TMEM133	GTF2H2
	ZNF611	CDC20
	HEY1	GTPBP6
	DAPK1	MYBBP1A
	KLHL6	GAR1
	NDC1	MPV17
	FGF5	IMMP1L

		FUT4		POP4
		CPEB2		MRPS6
		OAS3		RFC5
		CLK1		PPM1G
		ZNF750		ANAPC5
		ZNF292		IPO4
		CAPN5		MYCBP2
		UBE2D1		AKR1C2
		ARHGEF37		COMMD3
		KDELR3		CENPW
		ISG20		EBPL
		ZFHX2		EIF3L
		SSC4D		TXNDC17
		PADI3		RPS17
		PLA2G7		RAN
		SLC44A2		SELENOT
		CLIC4		RPL4
		AC091167.3		
		TMEM38A		
		SERPINE2		
		NRDC		
		HOXC8		
		PLAG1		
		MDM2		
		HOXC6		
		TMPRSS4		
		STAT1		
		SLC28A3		
		PCDHGA11		
		MELTF		
		TSPAN6		
		CDKN2D		
		BTG1		
		ICA1		
		COL1A2		
		ENTPD2		
		CTTNBP2NL		
		GLRX		
		EPB41L5		
		TMEM97		
		MINDY1		
		FAM107B		
		ARHGEF28		
		ZEB2		
		ZNF563		
		FYN		
		SLCO2A1		
		CCDC17		
		HOXA13		
		PLAC8		
		BMP6		
		PARP14		
		KRT16		
		IFIT3		
		PLA2R1		
		HIC1		
		MAP1LC3B		
		CTSB		
		ULBP2		
		MT-ND4L		
		ANKIB1		
		DOPEY2		
		ARL8A		
		PRSS27		
		TTC28		
		AURKC		
		MCF2L2		
		KATNBL1		
		IL4I1		
		MLLT3		
		UBN1		
		UNC5B		
		GLDC		
		JPH2		
		ABCA7		
		TMEM45A		
		ASTN2		
		HAL		
		AC005324.3		
		APOD		
		PLPP6		
		PRODH		
		ZNF350		
		CLDN16		
		APOL1		
		KIF3C		
		EBLN2		
		TIMP1		
		USP18		

		ETV1			
		SMPDL3B			
		AZI2			
		ZNF185			
		VIM			
		BNIP3L			
		ZNF600			
		SH3BP2			
		IQCD			
		SLC43A2			
		KLRD1			
		OPN3			
		PDE8A			
		OSMR			
		PMEPA1			
		B3GALT5			
		TCP11L2			
		SLC44A3			
		SLC9A3R2			
		ANKRD65			
		PPARD			
		PLSCR1			
		SERPINI1			
		TMEM59			
		FAM46B			
		OCLN			
		NOTCH3			
		CAMK2N1			
		TBC1D8B			
		ZYG11A			
		ANO8			
		RP2			
		DAB2			
		TRIB2			
		GSTT2B			
		LGI4			
		SNX10			
		MOK			
		CPT1C			
		B3GNT3			
		SCARB1			
		MPZL3			
		IFITM10			
		CCDC39			
		CEP192			
		MICAL1			
		C1QTNF9			
		ADGRV1			
		MYO5B			
		CIR1			
		KCNJ2			
		APP			
		TMEM213			
		IL1RL1			
		NFKBIA			
		HBP1			
		CERS4			
		CREB5			
		ADAP2			
		TUSC3			
		CTSO			
		NPIPA2			
		REPS2			
		SV2B			
		AP3B2			
		SOHLH2			
		TMCC2			
		SPATA20			
		ZNF821			
		KRT9			
		S100A4			
		CLCN4			
		CDKL5			
		UBE2B			
		DTNA			
		DPYSL2			
		DHX57			
		GNL1			
		CASC10			
		SAV1			
		GLIS3			
		IL24			
		ZNF221			
		ZNF467			
		DYSF			
		TCF7L2			
		FLRT3			
		TMEM86A			
		MYRFL			

		COL5A3			
		ITGB7			
		VSIG10L			
		CPA4			
		ZFP28			
		TMEM159			
		KIAA1211			
		SYT11			
		NPIPA1			
		ZNF699			
		TP53INP1			
		RRM2B			
		ASPHD2			
		BEX5			
		RAP1GAP			
		LOXL2			
		CDKL2			
		FAM32A			
		MGLL			
		ZNF391			
		B4GALT1			
		EPCAM			
		KIAA1328			
		CTH			
		LLGL2			
		YIPF4			
		LRCH4			
		ACP7			
		ABCB5			
		MARK3			
		CHMP5			
		TRIM36			
		CLIC3			

Table S4. Full list of GO terms from REVIGO for differentially expressed groups.

Increased by Treatment								
DAC			**Para**			**DAC+ Para**		
term ID	description	log₁₀ p-value	term ID	description	log₁₀ p-value	term ID	description	log₁₀ p-value
GO:0060337	type I interferon signaling pathway	-13.79	GO:0022904	respiratory electron transport chain	-11.15	GO:0019221	cytokine-mediated signaling pathway	-6.73
GO:0045071	negative regulation of viral genome replication	-9.20	GO:0032981	mitochondrial respiratory chain complex I assembly	-6.23	GO:0052548	regulation of endopeptidase activity	-6.40
GO:0048660	regulation of smooth muscle cell proliferation	-4.88	GO:0006754	ATP biosynthetic process	-2.71	GO:0048523	negative regulation of cellular process	-4.50
GO:0032069	regulation of nuclease activity	-4.43	GO:1902930	regulation of alcohol biosynthetic process	-2.64	GO:0030198	extracellular matrix organization	-3.82
GO:0031325	positive regulation of cellular metabolic process	-4.27	GO:0015988	energy coupled proton transmembrane transport, against electrochemical gradient	-2.53	GO:0045601	regulation of endothelial cell differentiation	-3.14
GO:0042775	mitochondrial ATP synthesis coupled electron transport	-3.93	GO:0044782	cilium organization	-2.46	GO:0045953	negative regulation of natural killer cell mediated cytotoxicity	-3.14
GO:0017157	regulation of exocytosis	-3.56	GO:0045787	positive regulation of cell cycle	-2.44	GO:0043691	reverse cholesterol transport	-2.99
GO:0001818	negative regulation of cytokine production	-3.53	GO:0046578	regulation of Ras protein signal transduction	-2.30	GO:1905037	autophagosome organization	-2.98
GO:0010642	negative regulation of platelet-derived growth factor receptor signaling pathway	-2.81	GO:0048864	stem cell development	-2.25	GO:0007009	plasma membrane organization	-2.90
GO:0001990	regulation of systemic arterial blood pressure by hormone	-2.71	GO:0006636	unsaturated fatty acid biosynthetic process	-2.20	GO:1902003	regulation of beta-amyloid formation	-2.87
GO:0055091	phospholipid homeostasis	-2.59	GO:0052547	regulation of peptidase activity	-2.05	GO:0048678	response to axon injury	-2.79
GO:0001911	negative regulation of leukocyte mediated cytotoxicity	-2.59	GO:1901570	fatty acid derivative biosynthetic process	-1.82	GO:0042775	mitochondrial ATP synthesis coupled electron transport	-2.74
GO:0045216	cell-cell junction organization	-2.57	GO:0001501	skeletal system development	-1.59	GO:0044257	cellular protein catabolic process	-2.66
GO:0030865	cortical cytoskeleton organization	-2.56	GO:0070534	protein K63-linked ubiquitination	-1.48	GO:0006501	C-terminal protein lipidation	-2.63
GO:0033273	response to vitamin	-2.45	GO:0046782	regulation of viral transcription	-1.42	GO:0003180	aortic valve morphogenesis	-2.54
GO:0030198	extracellular matrix organization	-2.39				GO:0048485	sympathetic nervous system development	-2.19
GO:0016338	calcium-independent cell-cell adhesion via plasma membrane cell-adhesion molecules	-2.35				GO:0055091	phospholipid homeostasis	-2.19
GO:0015988	energy coupled proton transmembrane transport, against electrochemical gradient	-2.32				GO:1990000	amyloid fibril formation	-2.19
GO:0097067	cellular response to thyroid hormone stimulus	-2.32				GO:0032530	regulation of microvillus organization	-2.19
GO:0016042	lipid catabolic process	-2.28				GO:0043116	negative regulation of vascular permeability	-2.19
GO:0006006	glucose metabolic process	-2.27				GO:0050918	positive chemotaxis	-2.10
GO:0009435	NAD biosynthetic process	-2.20				GO:0060037	pharyngeal system development	-2.10
GO:0007009	plasma membrane organization	-2.16				GO:0033273	response to vitamin	-2.05
GO:0010745	negative regulation of macrophage derived foam cell differentiation	-2.01				GO:0044267	cellular protein metabolic process	-2.03
GO:0046457	prostanoid biosynthetic process	-2.01				GO:0018146	keratan sulfate biosynthetic process	-1.96
GO:0071985	multivesicular body sorting pathway	-2.01				GO:0030865	cortical cytoskeleton organization	-1.96
GO:0071360	cellular response to exogenous dsRNA	-2.01				GO:0015718	monocarboxylic acid transport	-1.95
GO:0036152	phosphatidylethanolamine acyl-chain remodeling	-1.95				GO:0015988	energy coupled proton transmembrane transport, against electrochemical gradient	-1.92
GO:0048485	sympathetic nervous system development	-1.77				GO:0010941	regulation of cell death	-1.88
GO:0051258	protein polymerization	-1.77				GO:0016050	vesicle organization	-1.84
GO:0044267	cellular protein metabolic process	-1.73				GO:0048169	regulation of long-term neuronal synaptic plasticity	-1.81
GO:0034375	high-density lipoprotein particle remodeling	-1.70				GO:0033013	tetrapyrrole metabolic process	-1.78
GO:0050918	positive chemotaxis	-1.70				GO:0009235	cobalamin metabolic process	-1.78
GO:0015919	peroxisomal membrane transport	-1.69				GO:0006892	post-Golgi vesicle-mediated transport	-1.75

term ID	description	value	term ID	description	value	term ID	description	value
GO:0017196	N-terminal peptidyl-methionine acetylation	-1.69				GO:0045216	cell-cell junction organization	-1.74
GO:0043471	regulation of cellular carbohydrate catabolic process	-1.69				GO:0009435	NAD biosynthetic process	-1.72
GO:0032347	regulation of aldosterone biosynthetic process	-1.69				GO:0090329	regulation of DNA-dependent DNA replication	-1.71
GO:0031397	negative regulation of protein ubiquitination	-1.59				GO:1901264	carbohydrate derivative transport	-1.71
GO:0030388	fructose 1,6-bisphosphate metabolic process	-1.58				GO:0014821	phasic smooth muscle contraction	-1.71
GO:0015677	copper ion import	-1.58				GO:0046457	prostanoid biosynthetic process	-1.62
GO:0006357	regulation of transcription from RNA polymerase II promoter	-1.54				GO:0010224	response to UV-B	-1.62
GO:0071493	cellular response to UV-B	-1.48				GO:0006706	steroid catabolic process	-1.62
GO:0035563	positive regulation of chromatin binding	-1.48				GO:0044272	sulfur compound biosynthetic process	-1.59
GO:0006501	C-terminal protein lipidation	-1.41				GO:0006357	regulation of transcription from RNA polymerase II promoter	-1.52
GO:0048015	phosphatidylinositol-mediated signaling	-1.41				GO:0010107	potassium ion import	-1.50
GO:0046323	glucose import	-1.39				GO:0043471	regulation of cellular carbohydrate catabolic process	-1.42
GO:0051180	vitamin transport	-1.37				GO:0034375	high-density lipoprotein particle remodeling	-1.33
GO:0006654	phosphatidic acid biosynthetic process	-1.35				GO:0051289	protein homotetramerization	-1.31
GO:0035418	protein localization to synapse	-1.31				GO:0030050	vesicle transport along actin filament	-1.31
						GO:2000427	positive regulation of apoptotic cell clearance	-1.31

Decreased by Treatment

DAC			Para			DAC+ Para		
term ID	description	$\log_{10}$ p-value	term ID	description	$\log_{10}$ p-value	term ID	description	$\log_{10}$ p-value
GO:0034645	cellular macromolecule biosynthetic process	-11.94	GO:0046512	sphingosine biosynthetic process	-2.85	GO:0042254	ribosome biogenesis	-35.37
GO:0042254	ribosome biogenesis	-9.79	GO:0042759	long-chain fatty acid biosynthetic process	-2.76	GO:0016072	rRNA metabolic process	-31.32
GO:0006259	DNA metabolic process	-9.53	GO:0043604	amide biosynthetic process	-2.53	GO:0010467	gene expression	-25.56
GO:0000082	G1/S transition of mitotic cell cycle	-8.29	GO:0046474	glycerophospholipid biosynthetic process	-2.39	GO:0006259	DNA metabolic process	-23.50
GO:0016072	rRNA metabolic process	-8.13	GO:1901888	regulation of cell junction assembly	-2.34	GO:0019083	viral transcription	-22.59
GO:0006261	DNA-dependent DNA replication	-8.04	GO:0034976	response to endoplasmic reticulum stress	-2.32	GO:0000956	nuclear-transcribed mRNA catabolic process	-18.64
GO:0016032	viral process	-6.75	GO:0000187	activation of MAPK activity	-2.17	GO:0006281	DNA repair	-18.24
GO:0006614	SRP-dependent cotranslational protein targeting to membrane	-6.10	GO:0006940	regulation of smooth muscle contraction	-2.00	GO:0006414	translational elongation	-17.28
GO:0022616	DNA strand elongation	-5.93	GO:0044271	cellular nitrogen compound biosynthetic process	-2.00	GO:0000082	G1/S transition of mitotic cell cycle	-8.95
GO:0007005	mitochondrion organization	-5.91	GO:0051128	regulation of cellular component organization	-1.90	GO:0015949	nucleobase-containing small molecule interconversion	-6.96
GO:0002181	cytoplasmic translation	-5.89	GO:0048146	positive regulation of fibroblast proliferation	-1.84	GO:0009168	purine ribonucleoside monophosphate biosynthetic process	-6.96
GO:0010467	gene expression	-5.49	GO:0045070	positive regulation of viral genome replication	-1.76	GO:1901796	regulation of signal transduction by p53 class mediator	-6.87
GO:0000956	nuclear-transcribed mRNA catabolic process	-4.96	GO:0034660	ncRNA metabolic process	-1.68	GO:0090305	nucleic acid phosphodiester bond hydrolysis	-6.47
GO:0006415	translational termination	-4.90	GO:0000422	mitophagy	-1.55	GO:0006839	mitochondrial transport	-6.36
GO:0009168	purine ribonucleoside monophosphate biosynthetic process	-4.63	GO:0001764	neuron migration	-1.55	GO:0061641	CENP-A containing chromatin organization	-6.01
GO:0006414	translational elongation	-4.58	GO:1901566	organonitrogen compound biosynthetic process	-1.54	GO:0000732	strand displacement	-4.44
GO:0015949	nucleobase-containing small molecule interconversion	-4.02	GO:0050708	regulation of protein secretion	-1.50	GO:0006479	protein methylation	-4.41
GO:0006283	transcription-coupled nucleotide-excision repair	-3.98	GO:0006526	arginine biosynthetic process	-1.36	GO:1904666	regulation of ubiquitin protein ligase activity	-4.38
GO:0006839	mitochondrial transport	-3.74	GO:0001765	membrane raft assembly	-1.36	GO:0009411	response to UV	-4.26
GO:1904666	regulation of ubiquitin protein ligase activity	-3.71	GO:0031076	embryonic camera-type eye development	-1.36	GO:0006362	transcription elongation from RNA polymerase I promoter	-4.22
GO:0006336	DNA replication-independent nucleosome assembly	-3.55	GO:0016311	dephosphorylation	-1.34	GO:0051302	regulation of cell division	-3.93

GO ID	Process	Value	GO ID	Process	Value	GO ID	Process	Value
GO:0044267	cellular protein metabolic process	-3.51	GO:0006470	protein dephosphorylation	-1.33	GO:0006360	transcription from RNA polymerase I promoter	-3.75
GO:0000731	DNA synthesis involved in DNA repair	-3.39	GO:0043484	regulation of RNA splicing	-1.31	GO:0034982	mitochondrial protein processing	-3.53
GO:0097033	mitochondrial respiratory chain complex III biogenesis	-3.33	GO:0035357	peroxisome proliferator activated receptor signaling pathway	-1.30	GO:0006353	DNA-templated transcription, termination	-3.32
GO:0009411	response to UV	-3.20	GO:0045714	regulation of low-density lipoprotein particle receptor biosynthetic process	-1.30	GO:0051188	cofactor biosynthetic process	-3.06
GO:0018022	peptidyl-lysine methylation	-2.74	GO:0042780	tRNA 3'-end processing	-1.30	GO:0040029	regulation of gene expression, epigenetic	-2.83
GO:0060249	anatomical structure homeostasis	-2.74	GO:0035280	miRNA loading onto RISC involved in gene silencing by miRNA	-1.30	GO:1904874	positive regulation of telomerase RNA localization to Cajal body	-2.67
GO:2001234	negative regulation of apoptotic signaling pathway	-2.55	GO:0010561	negative regulation of glycoprotein biosynthetic process	-1.30	GO:0001522	pseudouridine synthesis	-2.67
GO:0000959	mitochondrial RNA metabolic process	-2.47	GO:0010988	regulation of low-density lipoprotein particle clearance	-1.30	GO:0044806	G-quadruplex DNA unwinding	-2.64
GO:0042148	strand invasion	-2.25				GO:0006352	DNA-templated transcription, initiation	-2.49
GO:0043497	regulation of protein heterodimerization activity	-2.25				GO:0006396	RNA processing	-2.34
GO:0036297	interstrand cross-link repair	-2.24				GO:0006354	DNA-templated transcription, elongation	-2.22
GO:0019395	fatty acid oxidation	-2.18				GO:0000959	mitochondrial RNA metabolic process	-2.21
GO:0006362	transcription elongation from RNA polymerase I promoter	-2.07				GO:0048709	oligodendrocyte differentiation	-1.81
GO:0008213	protein alkylation	-2.03				GO:0032456	endocytic recycling	-1.76
GO:0045333	cellular respiration	-1.99				GO:0051181	cofactor transport	-1.74
GO:0006576	cellular biogenic amine metabolic process	-1.97				GO:0016236	macroautophagy	-1.72
GO:0072321	chaperone-mediated protein transport	-1.97				GO:0033688	regulation of osteoblast proliferation	-1.66
GO:0006360	transcription from RNA polymerase I promoter	-1.88				GO:0019218	regulation of steroid metabolic process	-1.66
GO:0032780	negative regulation of ATPase activity	-1.82				GO:0009749	response to glucose	-1.54
GO:0014032	neural crest cell development	-1.81				GO:0033119	negative regulation of RNA splicing	-1.53
GO:0051302	regulation of cell division	-1.80				GO:0015838	amino-acid betaine transport	-1.50
GO:0015985	energy coupled proton transport, down electrochemical gradient	-1.76				GO:0002903	negative regulation of B cell apoptotic process	-1.50
GO:0006084	acetyl-CoA metabolic process	-1.70				GO:0097384	cellular lipid biosynthetic process	-1.50
GO:0006352	DNA-templated transcription, initiation	-1.69				GO:0042148	strand invasion	-1.50
GO:0006353	DNA-templated transcription, termination	-1.68						
GO:0090305	nucleic acid phosphodiester bond hydrolysis	-1.63						
GO:0090181	regulation of cholesterol metabolic process	-1.63						
GO:0000377	RNA splicing, via transesterification reactions with bulged adenosine as nucleophile	-1.62						
GO:0006312	mitotic recombination	-1.51						
GO:0006266	DNA ligation	-1.51						
GO:0010501	RNA secondary structure unwinding	-1.50						
GO:0006301	postreplication repair	-1.48						
GO:0009081	branched-chain amino acid metabolic process	-1.35						
GO:0009451	RNA modification	-1.33						

Table S5. cBioPortal RNA-seq data (for cancers with provisional TCGA data): frequency of expression alterations (%) in genes from COX-2-PGE$_2$ pathway and correlation to survival (Logrank test p-value).

Cancer type	n	PTGS2	PTGES	PTGES2	PTGES3	PTGER1	PTGER2	PTGER3	PTGER4	OS	DFS
Adrenocortical ca.	79	1.3	4	5	19	9	1.3	5	6	0.181	0.130
Cholangiocarcinoma	36	6	6	6	2.8	8	2.8	2.8	2.8	0.322	
Bladder urothelial ca.	408	4	3	2.5	8	5	3	4	8	0.0702	
Colorectal adenoca.	379	3	6	1.6	7	2.4	4	4	4	0.0375	
Breast invasive ca.	1093	1.7	4	4	8	1.6	2.6	2.8	3	0.259	
Glioblastoma mult.	160	3	6	6	8	6	4	6	3		
Cervical SCC	304	2.6	6	4	9	5	4	8	6	0.17*	
Esophageal carcinoma	184	7	2.7	8	8	5	6	1.6	9		
Stomach adenoca.	415	5	4	9	9	4	0.7	4	7		0.09*
Uveal melanoma	80	5	2.5	6	6**	5	5	5	5		
HNSCC	520	2.3	4	7	7	2.5	4	3	6	0.090	5.693 e-3
Kidney renal clear cell ca	533	1.3	2.3	2.4	5	0.4	4	7	5	0.179	0.144
Kidney renal papillary cell ca.	290	2.4	4	4	9	3	4	2.4	4	0.034	0.307
Liver hepatocellular ca.	371	0.8	6	7	8	6	3	1.3	4	0.283	
Lung adenocarcinoma	515	4	4	3	11	2.1	5	3	4	0.039	
Lung SCC	501	4	6	3	5	2.4	1.2	1.4	7		0.202
AML	173	4	5	4	2.9	2.3	3	2.9	2.3		n/a
Ovarian serous cystadenoca.	307	0.7	2.6	7	4	2.3	2	5	5	2.305 e-3	0.286
Pancreatic adenoca.	178	4	4	1.7	10	3	4	6	4	0.089	0.080
Mesothelioma	87	3	1.1	1.1	10	8	1.1	3	5	n/a	n/a
Prostate adenoca.	497	2.8	5	6	5	4	4	5	3		0.262
Skin cutaneous melanoma	469	8	1.7	3	9	4	4	2.1	3	0.118	0.236
Sarcoma	259	5	10	6	8	5	8	4	5		
Testicular germ cell ca.	150	2	7	7	9	5	6	3	3		0.072
Thymoma	120	5	3	3	8**	4	4	3	4	0.017	0.290
Thyroid cancer	501	3	2.4	4	6	3	3	3	5		
Uterine corpus endothelial carc.	177	1.7	2.8	5	6	4	7	3	6		
Overall %		**3.4**	**4.3**	**4.7**	**7.7**	**4.1**	**3.7**	**3.7**	**4.8**		

OS, Overall Survival, DFS, Disease/Progression Free Survival. Only p-values < 0.35 are shown. OS and DFS values describe negative impact on survival, unless marked by * (correlation with better survival). Expression alterations are predominantly observed as overexpression unless marked by ** (where downregulation is observed). SCC, squamous cell carcinoma.

Table S6. cBioPortal RNA-seq data (for cancers with provisional TCGA data): frequency of expression alterations (%) in genes involved in glutathione synthesis and correlation to survival (Logrank test p-value).

Cancer type	n	GCLC	GCLM	GSS	GGCT	OPLAH	GSR	OS	DFS
Adrenocortical ca.	79	5	1.3	6	16	6	4	0.093	
Cholangiocarcinoma	36	8	2.8	2.8	11	8	6		0.296
Bladder urothelial ca.	408	4	4	18	16	11	2.9	6.327 e-3	0.231
Colorectal adenoca.	379	8	4	39	20	7	16**		
Breast invasive ca.	1093	5	7	10	8	15	4	0.017	0.339
Glioblastoma mult.	160	4	4	14	31	3	8	0.175	0.087
Cervical SCC	304	3	3	12	8	8	4		
Esophageal carcinoma	184	7	4	13	16	11	8		
Stomach adenoca.	415	7	6	14	13	15	9	0.04*	0.163*
Uveal melanoma	80	13	1.3	8	6	35	6		
HNSCC	520	6	6	10	8	12	6	0.019	2.763e-3
Kidney renal clear cell ca	533	5	4	6	8	5	5	0.185	0.070
Kidney renal papillary cell ca.	290	4	4	16	7	8	3	0.051	0.041
Liver hepatocellular ca.	371	6	3	5	7	20	2.7		
Lung adenocarcinoma	515	6	2.9	8	5	14	5	0.035	
Lung SCC	501	7	7	10	10	12	5		0.211*
AML	173	5	6	4	3	6	7		
Ovarian serous cystadenoca.	307	4	4	13	4	35	3		0.337*
Pancreatic adenoca.	178	4	2.2	8	7	7	5	0.15*	
Mesothelioma	87	7	3	6	3	9	6	n/a	n/a
Prostate adenoca.	497	6	4	4	8	11	6**	0.13*	7.141e-3
Skin cutaneous melanoma	469	13	5	11	18	8	8	2.585 e-3	0.184
Sarcoma	259	2.7	4	9	12	3	5	0.011	0.041
Testicular germ cell ca.	150	7	13	2.7	26	4	9 **	0.05	
Thymoma	120	6	3	7	6	4	5	1.111 e-4	
Thyroid cancer	501	2.4	1.8	6	4	1.4	1.6		
Uterine corpus endothelial carc.	177	5	4	6	6	12	3		
Overall %		**5.9**	**4.2**	**9.9**	**10.6**	**10.8**	**5.7**		

OS, Overall Survival, DFS, Disease/Progression Free Survival. Only p-values < 0.35 are shown. OS and DFS values describe negative impact on survival, unless marked by * (correlation with better survival). Expression alterations are predominantly observed as overexpression unless marked by ** (where downregulation is observed). SCC, squamous cell carcinoma.

Table S7. qRT-PCR primers' sequences.

Gene	Forward (5'-3')	Reverse (5'-3')
DNMT1	GAGCCACAGATGCTGACAAA	GACACAGGTGACCGTGCTTA
DNMT3A	AAGGAGGAGCGCCAAGAG	GGATGGGGACTTGGAGATCA
DNMT3B	GGGAGGTGTCCAGTCTGCTA	GGCTTTCTGAACGAGTCCTG
TP63	GTTTCGACGTGTCCTTCCAG	TCTGGATGGGGCATGTCTTT
KRT5	TGAGGTCAAGGCCCAGTATG	ATCTCATGCTTGGTGTTGCG
IVL	AACACAAAGGGATCAGCAGC	GCTCCAACAGTTGCTCTTTCT
PTGS2 (COX-2)	TCATCATCAGCGCCCTCAA	GCTCGTTCACAGCCTTCATG
PTGER1 (EP-1)	GCCAGCTTGTCGGTATCATG	CTGCAGGGAGGTAGAGCTC
PTGER2 (EP-2)	AAGCTGTGGTCAAGGCTACA	GCCAAGTACCATGCTCACTG
PTGER3 (EP-3)	GGATCATGTGCGTGCTGTC	TGTGTCTTGCAGTGCTCAAC
PTGER4 (EP-4)	TGCTCATCTGCTCCATCCC	ATTCGGATGGCCTGCAAATC
ALOX5	ACATCTACCTCAGCCTCGTG	AGTTCCTCGTCCACAGTCAC
ALOX15B	AAATCAAGGGGTTGCTGGAC	AACTGGGAGGCGAAGAAGG
ALOX12	CTTGCTGAACACTCACCTGG	ATGGTGTAGCGGATATGGGG
LTA4H	GACTTCTGGGAAGGAACACC	TGCCACCAGTTCTTTAGGGA
CYP2E1	CGGAACTATGGGATGGGGAA	CGGAAGAGGATGTCGGCTAT
ITGAM (CD11b)	TGTTTCACGGAACCTCAGGA	ATCCATTGTGAGGTCCTGGC
ACTB	AAAGACCTGTACGCCAACAC	GTCATACTCCTGCTTGCTGAT