Title: Polygenic adaptation of rosette growth variation in *Arabidopsis thaliana* populations

Authors: Benedict Wieters[1], Kim A. Steige[1], Fei He[1], Evan M. Koch[2,3] Sebastián E. Ramos-Onsins4, Hongya Gu5, Ya-Long Guo6, Shamil Sunyaev[2,3], Juliette de Meaux[1]

Address:

Institute of Botany, University of Cologne, Cologne, Germany

Genetics Division, Brigham & Women's Hospital and Harvard Medical School, Boston MA, United States

Department of Biomedical Informatics, Harvard Medical School, Boston MA, United States

Centre for Research in Agricultural Genomics (CRAG) CSIC/IRTA/UAB/UB, Bellaterra, Spain

State Key Laboratory for Protein and Plant Gene Research, College of Life Sciences, Peking University, Beijing, China

State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, Beijing, China

ORCID IDs:
Benedict Wieters https://orcid.org/0000-0001-6476-1587
Kim Steige
Fei He
Evan Koch https://orcid.org/0000-0002-1124-4559
Sebastián Ramos-Onsins https://orcid.org/0000-0002-1776-140X
Hongya Gu https://orcid.org/0000-0003-2619-1522
Ya-Long Guo https://orcid.org/0000-0002-4643-4889
Shamil Sunyaev https://orcid.org/0000-0001-5715-5677
Juliette de Meaux http://orcid.org/0000-0002-2942-4750

Abstract (193 words)

The rate at which plants grow is a major functional trait in plant ecology. However, little is known about its evolution in natural populations. Here, we investigate evolutionary and environmental factors shaping variation in the growth rate of *Arabidopsis thaliana*. We quantified variation in plant growth in two light environments, across a set of 278 genotypes sampled within four broad regions, including an outgroup set of genotypes from China. The observed variation was validated in a field experiment conducted under natural conditions. All genotypes grew a markedly larger rosette when the light supply was decreased, demonstrating that environmental plasticity is a predominant source of variation to adapt plant size to prevailing light conditions. Yet, we detected significant levels of genetic variation both in growth rate and growth plasticity. Genome-wide association studies revealed that only 2 single nucleotide polymorphisms associate with genetic variation for growth above Bonferroni confidence levels. However, marginally associated variants were strongly enriched among genes with an annotated role in growth and stress reactions. Polygenic scores computed from marginally associated variants confirmed the polygenic basis of growth variation. For both light regimes, phenotypic divergence of the most distantly related population, which originates from China, is smaller than the variation observed within Europe, indicating that plant growth is constrained by stabilizing selection. We observed that Spanish genotypes, however, reach a significantly larger size than Northern European genotypes. Tests of adaptive divergence and analysis of the individual burden of deleterious mutations reveal that adaptive processes have played a more important role in shaping polygenic regional differences in rosette growth than maladaptive evolution.

Introduction

Growth rate is a crucial component of individual fitness, as it reflects the capacity of the organism to acquire resources and conditions reproductive output [1,2]. In experimental evolutionary studies, relative growth rate provides a measure of microbial adaptation in response to selection [3]. In plants, however, little is known about the evolutionary processes that influence variation in plant growth rate, despite its cornerstone importance in plant ecology [4–6].

Three processes may contribute to variation in growth rate: plasticity, adaptation or maladaptation. Plasticity describes the immediate adjustment of plant growth rate in response to environmental modifications [7]. Such change may occur as a passive consequence of resource limitations. Plant growth, for example, is slower in drought conditions or at lower temperatures [8,9]. Plastic adjustments of plant growth, however, can also actively contribute to maintaining fitness under challenging conditions. For example, shade avoidance allows plants to outgrow neighbors competing for light [10]. Such reactions may allow the organism to maintain high fitness in diverse environments without having to evolve genetically.

As the distribution range of a species expands, plastic modifications may become insufficient to adjust fitness, and genetic variation may be required for local adaptation [7,11]. There is clear evidence that genetic variation in plastic life history traits such as flowering time or seed dormancy contributes to the evolution of life-history decisions that are tailored to the local optimal growth season [12–15]. Surprisingly, the extent to which genetic variation in plant growth rate itself contributes to local adaptation is not known.

Genetic variation in growth rate may also arise in the absence of a compelling environmental change, as a consequence of population genetics processes. In bottlenecked populations, or in the aftermath of rapid range expansion, increased drift hampers the efficient removal of deleterious mutations, and individuals may become less fit [16–20]. Because plant growth is a component of fitness, genotypes carrying a larger burden of deleterious mutations may show decreased growth. Genetic variation in growth rates may thus also reflect maladaptation resulting from decreased population size.

The annual species *A. thaliana* has become a model system for both molecular and evolutionary biology, and it is well suited for determining the ecological and evolutionary significance of plant growth rates [21,22]. *A. thaliana* individuals can adjust their growth rate plastically to maintain their fitness. Plant rosettes grow to a larger diameter when light becomes limited [10,23]. Ample genetic variation in plant growth rates has also been documented in this species [24–26]. In addition, there is evidence that the resources allocated to growth are not identical throughout the species' range, because trade-offs between growth rate and development change with latitude (reviewed in [11], [27,28]). Furthermore, traits related to how resources are allocated to growth, such as growth inhibition upon the activation of plant defense, or plant dwarfism, have been associated with adaptation [29–33]. In summary, adaptive variation in the rate of plant growth may have evolved in *A. thaliana*. At the same time, the maladaptive or neutral evolution of a decreased growth rate cannot be excluded *a priori*. Indeed, *A. thaliana* has experienced recent severe bottlenecks in parts of its range, such as in Northern European or Chinese populations, which locally increased the rate of genetic drift and led to an accumulation of deleterious genetic variants. [34–36]. Neutral evolutionary forces could therefore also have modified growth rate in these populations.

To determine the roles of deleterious variation, adaptive evolution and/or plasticity in the genetic variation among plant growth rates, we analyzed variation among rosette growth rates across genotypes sampled from four broad regions (China, Spain, Northern and Western Europe). To assess the relative roles of genetic and plastic variation, we measured growth rates under full versus limited light intensities and characterized genetic variation in growth plasticity. This analysis reveals extensive regional differences in growth dynamics, most of which have a polygenic basis. Population genetics analyses indicate that natural selection has helped shape this variation.

Material & Methods

Phenotypic analysis and estimation of growth rate parameters

We chose 278 genotypes of *Arabidopsis thaliana* originating from 220 locations distributed throughout 4 regions for phenotypic analyses of growth rate variation (Northern Europe, Western Europe, Spain and Central-Eastern China, Table S1, Fig S1). Seeds were stratified for 3 days at 4°C in the dark on wet paper, and six replicate seedlings per genotype were replanted, each in one 6x6 cm round pots containing soil ("Classic" from Einheitserde) mixed with perlite. Pots were randomized within three blocks of 8 trays with 7x5 pots. Replicates were split into two groups to measure growth under two light regimes, high light (HL) and low light (LL), for 8 and 13 weeks, respectively. Plants were grown in a temperature-controlled walk-in growth chamber (Dixell, Germany) set at 20 °C day and 18 °C night, and watered once a week. Trays were randomized and the rows in the trays were rotated every two to three days to account for variability within the chamber. The plants were exposed to light for 12 h with LEDs (LED Modul III DR-B-W-FR lights by dhlicht®) set to 100% intensity of blue (440nm), red (660nm) and white (HL conditions) or 30% of red and blue plus 100% of white light (LL conditions), followed by a 10 min far-red light pulse (40% intensity at 735nm). The total measured light intensity was 224 +/- 10 µmol/m²s in HL and 95 +/- 7 in LL.

Individual plants were photographed approximately bi-weekly with a Canon EOS 5D Mark III digital camera until days 46 and 89, for those grown under the HL and LL regimes, respectively. Flowering time was measured as days to first flower opening. For genotypes without a flowering individual by the end of the experiment, a flowering time value of 59 or 90 days (last date that flowering was scored) was assigned to HL and LL plants, respectively. Since only 37% of the plants in the experiment flowered, we also used flowering time data from the 1001 Genomes project, according to which flowering was scored at 10 and 16°C for 177 genotypes [37]. A measure of the diameter of each plant (defined as the longest distance between two leaves) was extracted at least once a week with ImageJ (v.1.50b, [38]). For day 18 in HL plants, both the diameter and the rosette area were estimated with *Rosette Tracker,* an ImageJ tool [39]. The resulting strong correlation confirmed that rosette diameter accurately predicts rosette area (r^2=0.929, p=<2.2e-16). Rosette diameter was therefore used to determine the increase in rosette size over time.

We conducted two additional experiments to test the ecological relevance of rosette growth variation measured under controlled conditions. First, all genotypes were grown under HL conditions, in 5 replicates. In this experiment, instead of rosette diameter, we measured hypocotyl length after 15 days, to quantify variation in seedling growth. We further weighted 3-week old plants with a precision balance (Sartorius AC 210 P with accuracy of 0.1 mg) to quantify variation in plant biomass. We also followed a similar

experimental design to measure the diameter of plants grown outdoor in 2 replicates in the
field of the Cologne Institute of Plant Sciences. Sand was used instead of soil in 9 cm
diameter pots. Seeds were sown in September 2016, which corresponds to the native
season in the area (G. Schmitz, unpublished), and put outside after a week.

All following analyses were conducted using R [40], and function names refer to those in
the back package unless otherwise noted. We extracted three parameters that together
provided a comprehensive description of individual rosette growth. For this, rosette
diameter measurements over time (our input phenotype) were modeled as a logistic
growth using the *drm* function from the *drc* package in R with [41]. The three following
growth parameters were extracted: final size (FS, largest estimated rosette diameter),
slope (factor of magnification in the linear phase) and t50 (inflection point; time at which
growth is maximum and half of FS has been reached). For each parameter, a genotypic
mean correcting for block, tray and position effect was computed with a generalized linear
model with a Gaussian error distribution and the following model:
parameter~accession+block+tray/(row+col)+error.

To quantify environmental plasticity, the interaction term (GxE) of the following generalized
linear model (GLM) was extracted: *parameter ~ G * E + error,* where the parameter was
the genotypic mean for FS, slope or t50, G the genotype and E the light regime (HL or LL)-
each genotype was grown in three replicates in each light regime. GxE estimates increase
with the magnitude of growth plasticity induced by a decrease in light intensity (Fig. S2).

To assess the correlation of phenotypic traits with climatic variables, we investigated solar
radiation estimates, temperature, precipitation, humidity and wind speed with 2.5-min grid
resolution (WorldClim2 database, [42], accessed on March 20, 2018) and soil water
content [43]. Following Dittberner et al. (2018) [44], we estimated the mean over the
putative growth season for each genotype in addition to the annual averages.

Because of the strong correlations between climatic variables, we conducted principal
component analyses (PCAs) to combine the data. We analyzed annual average radiation
separately and combined the other variables into the PCAs: growing season data,
variables related to precipitation and to temperature. Raw climatic data and the principle
components (PCs) are in Table S1.

Statistical analysis of genetic variance

Broad-sense heritability (H^2) was determined for each trait in each environment as
previously reported [44]. Genetic and environmental variances were estimated using the
lme function from the *nlme* package [45], with the block as fixed and genotype as random
effect. The heritability of GxE could not be estimated, because plasticity was calculated on
the basis of genotypic means, and the heritability calculation requires variation within a
genotype to estimate environmental variance. Yet, we were able to compute a pseudo-
heritability for all traits, including GxE effects, since the genome-wide association study

(GWAS) mixed model (see below) allowed us to estimate the proportion of the observed
phenotypic variance that is explained by genotypic relatedness [46].

Regional differences in mean growth parameters were tested with a multivariate analysis,
using the *manova* function, the matrix of growth parameters (genetic means) or plasticity
and the following model: *growth~ population*light regime*. Significance levels were
determined by the Pillai test.

For univariate analysis, we used GLMs to test the effect of population of origin on the
genotypic means. A Gaussian distribution was taken for error distribution, and the
dispersion parameter was estimated by the *glm* function. Group means were compared
with the *glht* function and plotted on boxplots using the *cld* function, both of the *multcomp*
package [47].

Pairwise trait correlations within and across populations were calculated with the *cor.test*
function, and p-values were established using the *lmekin* function in R*, which includes a
kinship matrix of individuals (see below) and thus corrects for population structure (after
[27]). We used the *corrplot* function from the *corrplot* package to plot correlations [48]).
Pairwise correlation coefficients were used for the pathway analysis, and significant
correlations were plotted using the *qgraph* function in the *qgraph* package (version 1.6.4,
[49]). Plots were modified using inkscape version 0.92.3 (inkscape.org, [50]). Significance
levels were adjusted for false-discovery rates with the function *p.adjust*.

Genome-wide association studies

Genomic data were available for 235 of the 278 genotypes included in the phenotypic
analysis, i.e. for 84 genotypes from Northern Europe (NE, predominantly Sweden), 3 from
Western Europe (WE), 119 from Spain (SP) and 15 from China (CH) [36,37]. Chinese
genotypes were excluded from the GWAS because of their limited number (15 genotypes)
and their strong genetic divergence ([36], Fig. S3). In total, the growth parameters of 203
and 201 genotypes grown under HL and LL, respectively, were used for the GWAS.
Genome-wide association studies were conducted using the method from [51]. The
corresponding   GWAS   package   was   downloaded   from
https://github.com/arthurkorte/GWAS [50]. Single Nucleotide Polymorphisms (SNPs) with
minor allele frequency below 5% or with more than 5% missing data were removed from
the genotype matrix, resulting in a matrix of 1,448,192 SNPs, produced with *vcftools* (--012
recode option) (version 0.1.15, [52]). A kinship matrix was computed with the
emma.kinship function. For each growth parameter, genotypic means were used as
phenotype measurements. We performed GWAS across the European sample of
genotypes but also within region (Spain and Northern Europe). For each SNP, the script
output delivered p-values, which were Bonferroni corrected for multiple testing, and effect
sizes. For each trait, we estimated pseudo-heritability, the proportion of the observed
phenotypic variance that is explained by the estimated relatedness (e.g. kinship matrix,

[46]). To identify candidate genes underpinning significant GWAS associations, we
calculated the linkage disequilibrium (LD) around the SNP of interest and selected all
genes that were in a genomic window with LD above 0.5 within a 250 kb window around
the SNP. The LD was calculated as the Pearson correlation between the frequencies of
allele pairs. Additionally, we downloaded an annotation of loss-of-function (LOF) variants
[53] and performed a GWAS association following the procedure described above except
that the SNP data set was replaced with the LOF data set.

Validation of the polygenic signal

In humans, where population structure and environmental variation are correlated,
insufficient correction of the genetic associations caused by shared ancestry has been
shown to create spurious associations [54]. Even though environmental variance is much
better controlled in common garden experiments including kinship as a covariate,
association tests can still be confounded by genetic relatedness [55]. This is of particular
concern when many trait/SNP associations are below the Bonferroni significance
threshold. The rate of false positive was not excessively inflated by genomic differentiation
between regions, because GWAS performed within regions (Northern Europe or Spain)
had similar p-value distributions than GWAS performed on the complete phenotypic
dataset (Figs. S4 & 5). We nevertheless used two additional approaches to confirm the
polygenic basis of traits. First, we examined whether phenotypic variation could be
predicted by polygenic scores derived from sub-significant GWAS hits, with $p<10^{-4}$. For this
we used two replicates, and, for each trait, we calculated genotypic means, performed
GWAS, as described above, and computed polygenic scores following Berg and Coop
(2014) [51]. SNPs with a significant association with the phenotype were pruned to remove
SNPs standing in strong linkage disequilibrium with *plink* version 1.90 [56], following Sohail
et al. (2019) [54]. The *plink -clump* function was set to select SNPs below a (GWAS) P-
value threshold of 0.0001, start clumps around these index SNPs in windows of 1 Mb, and
remove all SNPs with P < 0.01 that are in LD with the index SNPs. The SNP with the
lowest p-value in a clump was retained for further analysis. Briefly, input files, including
allele frequencies for all SNPs, all SNPs with GWAS p-values lower than 10^{-4} and their
effect size estimates were created with a custom R script. We defined each genotype as
its own population (genotypes from Spain and Northern Europe were grouped in regions).
Scripts were downloaded from *https://github.com/jjberg2/PolygenicAdaptationCode*. The
pipeline was run with default parameters, and polygenic scores (Z-scores) were estimated.
The correlation between polygenic scores and the phenotype of the third replicate was
calculated with the function *cor.test*.

Second, we investigated functional enrichment among all genes in *A. thaliana* with a
ranking based on the GWAS results of the different traits or the differentiation index (Fst)
between Northern Europe and Spain. If the polygenic signal was due to insufficient
correction for population structure, we hypothesized that similar functions would be
enriched among genes with high Fst and genes with low GWAS p-values. Fst-values for

each gene were calculated with the *F_ST.stats* function of the PopGenome library [57] for
genotypes from Spain and Northern Europe. Negative Fst values were set to zero. To
assign a single GWAS p-value for each gene, either we assigned for each trait the lowest
p-value of SNPs within the gene, or, if no SNP was within the gene, we assigned the p-
value from the physically closest SNP [46].

When there were GWAS hits in the vicinity of duplicated genes, we removed tandem
duplicated genes within a 10-gene sliding window. For this, we first aligned all TAIR10
genes against each other by using BLAST (version 2.9.0, available at
https://blast.ncbi.nlm.nih.gov). Then the duplicated genes were selected as genes with an
e-value <1e-30. Finally, tandem duplicated genes identified with gene distance <10-genes
were filtered out to avoid inflated functional enrichments.

Enrichments were tested as previously described [58]. To assess similarity between traits
in Gene Ontology (GO) enrichments, we calculated graph-based similarity with the
GOSemSim package [59]. A distance matrix was estimated with average connectivity
between the GO terms. The clustered GO categories were then plotted as a dendrogram
with the *plot.phylo* function from the *ape* package (version 5.3, [60]). GO categories
enriched at p-value below 0.001 were highlighted. The distribution of enriched GO
categories was evaluated by visual inspection.

Testing for adaptation or maladaptation

For population genetics analyses, we sampled one genotype at random whenever plants
were sampled in the same location, acquiring a total of 220 genotypes. As a proxy for the
genomic load imposed by deleterious mutations, the number of derived non-synonymous
mutations per haploid genome has been proposed [61]. This approach was not possible
here because the genomes of individuals from China and Europe were sequenced in
different labs, and the depth and quality of sequencing varied too much to make a fair
comparison. Instead, we used two data sets that together catalogued LOF alleles after
controlling as much as possible for heterogeneity in sequencing quality: one that included
European genotypes [53] and a more recent data set that included Chinese genotypes
[62]. As an estimate of the individual burden of deleterious mutations, we counted the
number of LOF alleles for each individual and tested whether individuals with a larger
number tended to have a lower growth rate using the Spearman rank correlation.

To search for footprints of adaptive evolution, we computed an Fst value between Spain
and Northern Europe for each SNP in the GWAS analysis using the R-package *hierfstat*
and the *basic.stats* function [63]. Negative Fst values were set to zero, and the *quantile*
function was used to calculate the 95th percentile. The Fst distribution of SNPs associated
with any GWAS (p<10e-4) was estimated, and the 95th percentile was compared to the
distribution of an equally large, randomly sampled set of SNPs in 10,000 permutations, to
determine the likelihood to detect a significant functional enrichment among random SNPs.
To compare the phenotypic differentiation of traits, Qst values for the phenotypic traits
were estimated as previously described [44]. Briefly, Qst was estimated as VarB / (VarW +

VarB), where VarW is the genotypic variance within and VarB between regions. These
variances were estimated with the *lme* function of the *nlme* package [45], with the block as
fixed and population/genotype as random effect. We extracted the intercept variance for
VarB and the residual variance for VarW. Since replicates were taken from the selfed
progeny of each genotype, VarB and VarW are broad-sense genetic variance components.
To reveal signatures of local adaptation, the Qst of each trait was compared to the 95[th]
percentile of the Fst distribution (between Spain and Northern Europe) [64,65]. To
consolidate evidence for a significant Qst, we permuted phenotypic data by randomizing
genotype labels and calculated the difference between the randomized Qst and 95[th]
percentile of Fst, following [44]. The permuted Qst distribution was compared to the true
Qst values to estimate p-values. In a second approach, we used a multivariate normal
distribution to generate phenotypic divergence based on the kinship matrix to generate an
expected Qst distribution [66]. Finally, we applied the over-dispersion test (Qx test), which
compares polygenic scores computed for associated versus random SNPs (null model), in
a process similar to a Qst/Fst comparison, but assuming that each population is composed
of the selfing progeny of one genotype [51]. A Qx significantly larger than the Qx computed
for the null model indicates that polygenic trait prediction is more differentiated than
expected from the kinship matrix and can be taken as an indication that the trait has
evolved under divergent selection, either within or between regions [51].

Results & Discussion

Ecological relevance of rosette growth variation

On the basis of the more than 15,000 rosette images we collected, we describe rosette growth variation with three parameters; each refers to the ways in which growth can differ among genotypes: i) the time lag until the exponential growth phase is reached (t50), ii) the speed of growth during the exponential growth phase (slope) and iii) the final size (FS) at which rosette diameter plateaus at the end of the rosette growth phase (Fig.1, Table S2-3). Of the parameters, FS displayed the highest broad-sense heritability, in plants grown under both regimes: high light (HL, H^2= 0.636) and low light (LL, H^2= 0.794, Table S4). Trait variation measured in controlled settings sometimes fails to reflect variation expressed in natural conditions [67,68]. This is not the case for rosette growth variation in *A. thaliana*. FS correlated positively with plant biomass (r^2= 0.230, p-value= 0.008) and seedling growth (r^2= 0.385, p-value= 8.5e-7) in the growth chamber. It also correlated positively with plant diameter measured in the field (r^2= 0.307, p-value= 0.0002). The variation in rosette growth we report is therefore ecologically relevant.

Environmental plasticity has the strongest impact on plant growth variation

Light regimes revealed that plasticity has the strongest impact on rosette growth (Fig. 1, MANOVA HL vs LL: F= 2529.20, df= 1, p-value= <2.2e-16, Table 1). In plants grown under LL, the maximum growth rate was delayed and rosette growth plateaued at a larger size (Table1, Fig. 2). This observation was in agreement with the reduced relative growth rate reported in many plant species when light supply decreases [69], whereas the larger FS reflected the expected shade avoidance reaction [70]. Estimate of GxE interaction effects quantified the growth plasticity of individual genotypes in relation to light regime (Fig. S6).

Table 1. Multi- and uni-variate analyses of growth variation in response to light regime, region and their interaction.

Response	df	Multivariate analysis (MANOVA)		Final Size		t50		Slope	
		F	p-value	F	p-value	F	p-value	F	p-value
Light regime	1	2529.2	**< 2.2E-16**	2638.5	**< 2.2E-16**	4961.5	**< 2.2E-16**	1.0	0.315
Region	3	8.4	**2.74E-12**	19.8	**4.57E-12**	6.9	**1.4e-5**	0.5	0.704
Light*Region	3	2.4	**0.009**	1.9	0.123	1.7	0.171	4.6	**0.004**

The multivariate analysis was conducted on a matrix of genotypic values for FS, t50 and slope for all four regions (Spain, Northern Europe, Western Europe, China).

Spanish genotypes show the most vigorous rosette growth

We found evidence for rosette growth variation across regions (MANOVA in Table 1, Figs. 1 & 2). Within Europe, Spanish genotypes reached the largest FS in both HL and LL plants (Tables 1 & S5, MANOVA: F=19.835, df= 3, p-value= 4.57e-12). Although the growth slope did not differ significantly across regions, we observed that, under HL conditions, Spanish genotypes reached 50% of their FS (=t50) significantly later than the genotypes originating from Northern Europe (t50= 15.16 vs 13.78, respectively, GLHT z= 3.567, p-value= 0.002, Fig. 2c). This effect was not significant in plants grown under LL conditions and was not

sufficiently different to contribute to GxE variation (Fig. 2d, Table 1). We verified that the larger rosette size observed in Spain was not due to a delay in flowering that would have extended the period of vegetative growth. FS variation among Northern European and Spanish genotypes did not correlate with the flowering time scored in our experiment (HL: r^2= -0.204, p-value= 0.146, LL: r^2= 0.147, p-value= 0.293). Published flowering time measures collected at 10°C and 16°C confirmed this finding (Fig. S7). Indeed, genotypes in these two regions are generally late flowering ([37,71,72], Fig. S8, GLHT z= -1.111, p-value= 0.267).

Within regions, patterns of co-variance between growth parameter and flowering time further showed that the evolution of plant growth cannot be explained by the evolution in flowering time. We observed that genetic variation in flowering time was not correlated with FS. Flowering time was indirectly correlated with growth via its co-variation with slope, but the sign of the correlation between the two regions reversed in LL plants (Fig. 3, correlation FT16 to slope in LL plants in Spain: 0.223, p-value= 0.041, in Northern Europe: -0.259, p-value= 0.049, Table S6). Within Spain, flowering time was also negatively correlated with the timing of growth (t50) in HL plants, indicating a trade-off, since early flowering lines had a larger time lag in their growth (Fig. 3, correlation t50 to FT16 in HL plants in Spain: -0.311, p-value= 0.0023, Table S6). This pattern confirmed a previous report that the correlation between flowering time and growth rate changes with latitude [28]. It also showed that the evolutionary history of rosette growth variation differs across regions.

Chinese genotypes show that growth rate variation is constrained in evolution

Despite a long history of population isolation [36], the growth rate of Chinese genotypes was comparable to that shown by most European genotypes (Table S5, Fig. 2 a-d). Under LL conditions, Chinese genotypes showed lower t50 and FS values only when compared to Spanish genotypes (Table S5, Fig. 2 a-d). Under HL, genotypes from China did not differ significantly from those from any other region (Fig. 2, Fig. S9). The Chinese genotypes indicate that the phenotypic evolution of rosette growth does not scale with the extent of genetic divergence. We therefore conclude that, in this outgroup lineage, it has been constrained by stabilizing selection.

The Chinese population was also the only one to show a difference in plasticity to the light regime (Fig. S6). Compared to Spanish genotypes, Chinese genotypes displayed a GxE that was lower for FS and higher for slope (FS: GLHT z= 2.729, p-value= 0.029; slope: GLHT z= 3.827, p-value= <0.001; Fig. S6). When grown under the LL regime, these genotypes displayed a lower FS than genotypes from Spain. In contrast, within Europe, we observed no significant difference in the growth plasticity of plants in relation to light regime, despite the fact that Northern populations are exposed to lower average light intensity (Fig. S10).

GWAS reveal only two SNPs significantly associated with rosette growth variation

We used GWAS to determine the genetic basis of variation in growth rate within Europe (Figs. 4 & S11-S13). The sample size (15) and strong population structure of Chinese genotypes precluded their inclusion in this analysis (Fig. S3). Overall, we found few significant genetic associations, indicating that genetic variance for growth rate is generally polygenic. One SNP (chromosome 1, position 24783843) associated with t50 variations in LL plants (effect size= -2.475, p-value= 2.6E-9, Fig. 4, Table S7). A second SNP (chromosome 3, position 951043) was associated significantly with the slope of rosette diameter growth in HL plants within Spain (effect size= 1.229, p-value= 8.4E-7, Fig. 4) and was polymorphic only in the Spanish set of genotypes. This SNP was within a 1Mb DNA fragment showing strong local LD and enclosing 21 genes. Two additional SNPs were associated with GxE for FS and t50 in HL plants in Northern Europe, respectively, with p-values just below the Bonferroni threshold (Table S7). Yet, we found no SNP significantly associated with FS above the Bonferroni threshold, although FS is the most heritable trait (Table S4). Local genetic variation in slope and t50, growth parameters which display moderate but significant genetic variance, appear to be controlled by low-frequency variants of comparatively larger effect, since some of them were associated above Bonferroni threshold. This genetic architecture resembles that reported in the same species for flowering time [73,74]. In contrast to slope and t50, variation in FS appeared more polygenic.

Polygenic scores and functional enrichments confirm the polygenic basis of growth variation

Traits with polygenic architecture are controlled by variation in many loci of low frequency and/or low effect sizes [75,76]. However, random SNPs with outlier frequency are not always sufficiently corrected for with the kinship matrix and these may give rise to spurious associations. Studies of polygenic traits such as human height have shown that residual effects of population structure can give signals of genetic association [54,77]. Similar effects were also encountered in studies of phenotypic variation in plant systems [55]. They are expected whenever environmental variance co-varies with population structure, as is likely the case in human studies, but can also persist in common garden studies if populations are geographically differentiated in the genetic component of the trait. To confirm the polygenic basis of growth variation, we evaluated the biological relevance of marginally significant genetic associations. For this, we first randomly chose two of the three replicate measures of growth and performed an independent GWAS analysis, extracting effect sizes of SNPs associated with a phenotype at a sub-significant p-value below 10^{-4}. The associated sets were composed of 22 to 32 unlinked SNPs. We used their effect size to compute polygenic scores for each parameter [51]. The polygenic scores computed for all growth parameters on two replicates correlated significantly with the phenotype measured independently in the third replicate (Table S8). The correlation was highest for FS measured in LL plants (Rho=0.567, p=<2.2e-16). In fact, FS, the most heritable trait, could be predicted with the highest accuracy in plants grown under both light regimes (Table S8). This confirms that genetic associations, despite their marginal

significance, can predict the observed phenotypes. This analysis did not entirely rule out the possibility that population structure contributes to the polygenic score, because the FS could also be predicted by random SNPs, but the correlation was much weaker (Table S11).

We thus further selected SNPs showing a sub-significant association ($p<0.001$) and investigated functional enrichment among genes that mapped within 10kb of the SNP but had been pruned for duplicates (Table S9). All traits showed functional enrichment within gene ontology (GO) categories related to growth, confirming that these genetic associations were biologically relevant (Table S9). For example, genes associated with variation in FS, the most polygenic trait, were enriched among genes involved in the growth-related functions "cotyledon development", "auxin polar transport", "response to mechanical stimulus" and "cell proliferation" (p-value= 0.0085 or lower). Interestingly, mechanical stimuli have been shown to strongly influence seedling growth, and we observed that FS correlated with hypocotyl length and biomass in 3-week-old plants (Fig. S14, [78]). Additionally, several categories related to defense and stress reactions, such as response to salt stress, response to chitin, regulation of defense response to fungus and negative regulation of defense response, were enriched. Variation in stress-related functions is known to have an impact on plant growth in *A. thaliana* [31]. Furthermore, we also found that SNPs associated with FS plasticity to light are enriched among genes involved in the shade avoidance response (p = 0.0023), by which plants exposed to limited light conditions increase stem elongation [10]. Associated genomic regions included, for example, PHY RAPIDLY REGULATED 2 (PAR2, AT3G58850), a negative regulator of shade avoidance [79] or LONG HYPOCOTYL UNDER SHADE (BBX21, AT1G75540), a regulator of de-etiolation and shade avoidance [80].

We reasoned that, if population structure outliers are driving the signal of association rather than variants of phenotypic relevance, we should observe similar functional enrichment among genes with elevated Fst and SNP associations. Indeed, genes with elevated Fst reflecting regional adaptation of traits could create spurious associations with traits that are also differentiated between regions. We thus ranked genes either by Fst or by the GWAS p-value of the closest SNP and determined enriched GO categories. We visualized overlaps in functional enrichment by clustering GO terms on the basis of the genes they shared (Fig. S15). The enrichment based on Fst revealed three strongly enriched GO terms: "organ morphogenesis", "circadian rhythm" and "virus-induced gene silencing" (p= 0.0009 or lower, Table S10). The enrichment in GO category "circadian rhythm" may reflect the local adaptation to Northern variations in day length [81,82]. Genes close to SNPs associated with the different growth parameters, however, had clearly distinctive patterns of functional enrichment (Fig. S15, Table S10). We therefore argue that even though population structure outliers may create some false-positive associations, the polygenic pattern of association that we observe at sub-significant level is clearly biologically relevant.

No association between per-individual burden and growth

In areas located at the edge of the distribution range of *A. thaliana*, populations may have accumulated an excess of deleterious mutations in the aftermath of their genetic isolation [37,83]. This could have resulted in a mutational load that would have decreased fitness components such as plant growth, because it influences the resources available for the production of progeny [1,18,20]. We thus hypothesized that the lower FS observed in Northern Europe may result from maladaptive forces associated with the demographic history of the region.

This hypothesis could not be supported. No significant difference was detected in total number of LOF mutations per genome in Northern Europe compared to Spain (GLHT: z-value= 0.634, p-value= 0.526, Fig. S16). This observation has been previously reported [53]. In addition, we detected no significant correlation between the number of LOF alleles per genome and the average final size in HL or LL plants within Europe (r^2 in HL= 0.079, p= 0.262, r^2 in LL= 0.029, p= 0.684). Furthermore, we observed no significant difference in growth between Northern European and Chinese populations, despite their significantly higher burden of LOF alleles per genome (GLHT China versus Northern Europe: z-value= -20.259, p-value= <1e-4, Fig. S17, [62]). Therefore, we conclude that the individual burden of LOF mutations is unrelated to rosette growth variation.

We reasoned that lower growth rate might also be associated with a small subset of LOF mutations. To test this hypothesis, we investigated genetic associations between LOF alleles and the three growth parameters (see Methods). We detected no association between LOF alleles and FS, yet there was a significant association of LOF variation at gene AT2G17750 with variation in both t50 in LL plants and t50 plasticity (Fig. 5, effect size= -3.542, p-value= 7.49e-6, gene-Fst= 0.113, and effect size= -4.470, p-value= 4.99e-6 for t50 and t50 plasticity, respectively). AT2G17750 encodes the NEP-interacting protein (NIP1) active in chloroplasts, which was reported to mediate intra-plastidial trafficking of an RNA polymerase encoded in the nucleus [84]. INIP1controls the transcription of the *rrn* operon in protoplasts or amyloplasts during seed germination and in chloroplasts during later developmental stages [84]. The LOF variant is present primarily in Northern Europe (MAF=16 and 0.8% in Northern Europe and Spain, respectively) but is unlikely to be deleterious: it correlates with a decrease of t50, which is a faster entry in the exponential growth phase indicative of increased growth vigor (Fig. 5). Taken together, this result does not support the hypothesis that decreased FS in Northern Europe or China is controlled by deleterious variation.

FS variation might reflect local adaptation at the regional scale
During the growth season, Northern European *A. thaliana* populations are exposed to lower average temperatures (Fig. S10). Smaller rosettes are more compact, and increased compactness is often observed in populations adapted to cold temperatures [30,85–87]. Freezing tolerance, which was indeed reported to be higher in Northern Europe, is associated with functions affecting rosette size [88]. We thus hypothesized that the

decreased FS and t50 observed for Northern European genotypes grown under both light
regimes is the result of polygenic adaptation to lower average temperatures. We used the
19 to 47 LD-pruned set of SNPs associating in GWAS at a sub-significant level to compute
polygenic scores for each genotype and each trait, and used Qx, a summary statistic that
quantifies their variance across locations of origin. A Qx value outside of neutral
expectations inferred from the kinship variance in the population, indicates excess
differentiation of polygenic scores, as expected if individual populations evolved under
divergent selection ([51]). We observed that all traits displayed a strongly significant Qx
(Table S11). The differentiation of polygenic scores between the individual populations of
origin suggests that divergent selection may be acting locally. Local adaptation has indeed
been reported at this scale in this species [89–91]. This result should however be taken
with caution, because, like the GWAS hits it is using, the Qx statistics is sensitive to
population structure outliers. but only final rosette size measured in HL plants and t50
measured in LL plants displayed polygenic scores that differed significantly between
regions (p-value= 0.0162 and 0.0309, respectively, Fig. S18).[89–91]

Interestingly, we observed that final rosette size measured in HL and t50 measured in LL
displayed polygenic scores that differed significantly between regions (p-value= 0.0162
and 0.0309, respectively, Fig. S18). We thus further tested whether, at the phenotypic
level, regional differentiation in growth rate departed from neutral expectations. We first
investigated whether variants associated with phenotypic variation in rosette diameter
showed increased genetic differentiation. The Fst distribution of the 1360 SNPs associated
with any of the growth parameters at $p<10^{-4}$ was significantly shifted towards higher values
(Kolmogorov-Smirnov test, $p<10^{-4}$). Thus, associated SNPs are collectively more likely to
be differentiated than the rest of the genome. This pattern is not caused by the
confounding effect of population structure, because the functional enrichments are mostly
specific to the phenotypes (Fig. S15). We note, however, that a few spurious genetic
associations could contribute to both higher Fst and over-dispersion of polygenic scores
[54,55,77]. Additional evidence based on approaches independent of GWAS is therefore
required to support the adaptive significance of regional differences in growth rate in
Europe. We first used the population kinship matrix to parameterize a multivariate normal
distribution and predict the amount of additive phenotypic divergence expected if the trait
evolves neutrally [66]. We observed that differentiation for FS measured in HL plants was
marginally more differentiated than predicted under neutral conditions (Qst=0.325, p-
value= 0.085, Fig. 6). The other parameters did not depart from neutrality (Qst ranging
from 0.029 to 0.27, min p= 0.11, Fig. 6). Since the divergent Chinese population revealed
that growth rate variation is constrained in evolution, this test is likely to be overly
conservative. In addition, it predicts the divergence in additive genetic variance, but in the
selfing species *A. thaliana*, the whole genetic variance, i.e. broad sense heritability, can
contribute to adaptation. We also compared the distribution of phenotypic variation within
and between regions to the SNP Fst-distribution [65]. We used the Fst between Northern
Europe and Spain as an estimate for nucleotide differentiation and compared it to
differentiation at the phenotypic level (Qst; [12,44,64]). For FS and t50, the Qst was
significantly greater than genetic differentiation at 95% of single nucleotide (Table 2). This

suggests that selective forces have contributed to the regional adaptation of final rosette
size in Europe. Nevertheless, we detected only weak correlations between growth
variation and temperature at the location of origin (Fig. S19), suggesting that growth rate
could be locally adapted to the conditions prevailing in each region. The environmental
factors contributing to adaptive divergence in plant growth thus remain to be determined in
this species.

Table 2. FS and t50 quantitative differentiation (Qst) exceed differentiation given by single
SNPs

Trait	Qst	Percentile of single SNP Fst
FSHL	**0.379**	**96.57**
FSLL	**0.282**	**95.80**
t50HL	**0.300**	**95.95**
t50LL	0.189	94.79
SLHL	0.081	93.23
SLLL	0.010	91.62

Legend. Qst for each trait measured in HL and LL plants. Linear mixed models were used
to quantify the ratio of genetic variation between versus within Spain and Northern Europe
(Qst). The 95th percentile of the distribution for single SNP Fst between regions was 0.205.
The 95th percentile of permuted Qsts can be used as an estimate of significance. HL:
plants grown under high light regime, LL: plants grown under low Light regime, FS: Final
Size, t50: time to maximum growth and SL: slope. Bold: significant Qst values (*p*<0.05).

Conclusion

Our comprehensive analysis of genetic diversity in rosette growth rate, within and between
four broad regions of the distribution area of *A. thaliana,* reveals the environmental and
evolutionary factors that control this complex trait, which is of central importance for plant
ecology. We show that plastic reactions to limited light conditions have the strongest
impact on variation in rosette growth rates. Yet, we also provide evidence for significant
genetic variation within and between regions. We observed that Spanish genotypes show
more vigorous rosette growth and reach the larger size, regardless of light conditions.
Although GWAS reveal very few associations that pass Bonferroni correction, analyses of
functional enrichments and polygenic scores demonstrate that the polygenic basis of trait
variation can also be explored in the presence of moderately significant genetic
associations. The greater phenotypic differentiation observed within Europe compared to
between Europe and China, which stands in contrast to the extent of genetic divergence,
demonstrates that stabilizing selective forces constrain the evolution of growth rate. The
analysis of polygenic scores and patterns of differentiation suggests that much of the
polygenic variation observed within Europe has been shaped by natural selection, rather

than by the burden imposed by deleterious mutations. Methodological developments that improve the use of polygenic associations for the study of local adaptation are needed to consolidate these conclusions. Leveraging polygenic associations in local adaptation studies remains challenging [74]. Methodological developments are needed to consolidate our conclusions. They will help better integrate complex traits in our understanding of the genetic processes underpinning ecological specialization [6].

Acknowledgments

This research was funded by the European Research Council (ERC) through the "AdaptoSCOPE" grant 648617, by the German American Fulbright Commission, and by grant AGL2016-78709-R (MEC, Spain) to S.E.R.-O. We also acknowledge the financial support of the Spanish Ministry of Economy and Competitivity for the Center of Excellence Severo Ochoa 2016–2019 (SEV-2015-0533) and by the CERCA Programme/Generalitat de Catalunya. We acknowledge grants R35 GM127131 and RO1 MH101244 from the National Institutes of Health (NIH) for S.S and E.K. We thank Prof. Andreas Beyer and Prof. Arthur Korte for advice regarding GWAS analyses and Emily Wheeler, Boston, for editorial assistance.

Figures

Fig. 1. Regional growth rate estimates in HL and LL. Estimated growth curves averaged over region (from drm function). The growth curves were estimated from diameter measurements at different time points. An illustration of the parameters that are estimated from these growth curves are included in the plot (Final Size is a diameter, t50 a time point and Slope the fold increase in the linear phase). HL (dashed line), LL (solid line), China (orange), Northern Europe (green), Spain (purple) and Western Europe (red).

Fig. 2. Significant regional differentiation of Final Size and t50 in HL and LL. *A.thaliana* genotypes are grouped based on geographical origin. Box plots show regional variation in Final Size (upper row) and t50 (lower row) for HL (left) and LL (right). Groups that do not share a letter are significantly different according to Tukey's HSD (p-value < 0.05). Region abbreviations: China (CH), Northern Europe (NE), Spain (SP) & Western Europe (WE).

Fig. 3. Trait network within SP and NE. Circles represent traits and lines correspond to significant pearson correlations (r^2) between traits for genotypes from Spain (SP, solid line) and Northern Europe (NE, line with dashes). The significance of interactions was estimated with a correction for population structure using *lmekin* and was corrected for multiple testing (FDR correction). Red lines represent positive and blue lines negative correlations. The traits are abbreviated: FT10&FT16 are flowering time at 10/16 °C from 1001 Genomes et al. (2016) and VERN is an estimate of vernalization requirement estimated as FT16-FT10. FSHL/FSLL are Final Size in HL/LL. t50H/t50L are t50 in HL/LL. SLHL&SLLL are Slope in HL/LL. Correlations and p-values in Table S6.

Fig. 4. GWAS-results for 4 phenotypes. Manhattan plots of GWAS of t50 in LL with all European genotypes (a), with a peak on Chromosome 1, GxE of Final Size with all European genotypes (b) with a peak on Chromosome 2, Slope in HL within Spain (c) with a peak on Chromsome 3 and t50 in HL in Northern Europe (d) with a peak on Chromosome 1. The dotted line shows the corresponding Bonferroni threshold adjusted for a p-value of 0.05.

Fig. 5. Loss-of-function association and phenotype of t50 in LL/GxE. Manhattan plot of a GWAS with loss-of-funcion alleles and t50LL (a) and t50GxE (c) as input phenotypes with the same association (AT2G17750) above the Bonferroni threshold (blue line). Boxplot of the phenotype of t50LL (b) and t50GxE (d) versus the allele state at AT2G17750 (0 means functional, 1 is a loss-of-function). The colors separate the populations into Spain (purple) and Northern Europe (green).

Fig. 6. Qst. Expected distribution for quantitative trait differentiation between the Spanish and Northern European population. The expectation is based on a multivariate normal distribution assuming a neutral trait with polygenic basis. Horizontal lines indicate observed Qst for the individual growth parameters, FS (Solid line), t50 (Dashed line), Slope (Dot line), in HL (orange) and LL (cyan). The red arrows show the 90^{th}, 95^{th} and 99^{th} percentiles of the distribution.

Supporting information

Fig. S1. Genotype origin map. Each dot represents the sampling point of a genotype. The genotypes where assigned to Northern Europe (green), Western Europe (red), Spain (purple) and China (orange).

Fig. S2. GxE for Final Size. GxE was estimated based on a glm(Final Size ~ genotype * environment) and is indicated by the color. Each dot corresponds to a genotype with its phenotype in HL (x-axis) and LL (y-axis). The black dot shows the average over all genotypes with standard deviation. The line shows a linear model for Final Size in LL ~ Finals Size HL.

Fig. S3. Principal component analysis of 227 genotypes. The PCA is based on 1.5 millions SNPs with a minor allele frequency larger than 0.05. The first two principle components explain about 16 % of the variance between the genotypes. Regions: China (orange), Northern Europe (green), Spain (purple), Western Europe (red).

Fig. S4. Regional comparison of the GWAS results within Spain to across Europe. qq-plots comparing the GWAS data in Fig. S10 (Spain, x-axis) to the data from Fig. S9 (Europe, y-axis). The traits are Final Size (upper row), t50 (2^{nd} row) and Slope (lower row) in HL (left column), LL (middle column) and their GxE (right column). The grey dotted line indicates the neutral expectation.

Fig. S5. Regional comparison of the GWAS results within Northern Europe to across Europe. qq-plots comparing the GWAS data in Fig. S11 (Northern Europe, x-axis) to the data from Fig. S9 (Europe, y-axis). The traits are Final Size (upper row), t50 (2^{nd} row) and Slope (lower row) in HL (left column), LL (middle column) and their GxE (right column). The grey dotted line indicates the neutral expectation.

Fig. S6. Regional differences for GxE for each trait. GxE for FS (left), t50 (center) and Slope (SL, right). The phenotypic values are based on 214 genotypes of *Arabidopsis thaliana*. Groups that do not share a letter are significantly different according to Tukey's HSD (p-value < 0.05). Population abbreviations: China (CH), Northern Europe (NE), Spain (SP) & Western Europe (WE).

Fig. S7. Flowering time from 1001 Genomes. Flowering time in 16°C conditions of each genotype plotted for Northern Europe (green) and Spain (purple, based on data from 1001Genomes, 2016). The regions showed no phenotypic difference, as indicated by the same letter (pairwise GLHT, p-value> 0.05).

Fig. S8. Flowering time in the experiment. Flowering time of each genotype in HL (left) and LL conditions(right). Missing values were replaced with 59 (HL) or 90 (LL) days after sowing. Boxplots with different letters are significantly different according to Tukey's HSD (p-value < 0.05). Population abbreviations: China (CH), Northern Europe (NE), Spain (SP) & Western Europe (WE).

Fig. S9. Regional differences in Slope. The phenotypic values are based on 220 genotypes of *Arabidopsis thaliana* in HL (left) and LL conditions (right). Groups that do not share a letter are significantly different according to Tukey's HSD (p-value < 0.05). Population abbreviations: China (CH), Northern Europe (NE), Spain (SP) & Western Europe (WE).

Fig. S10. Climatic variation between regions. Annual average of the monthly radiation (left), monthly average temperature (center) and monthly precipitation (right) estimates for the sampling location of each genotype from Worldclim2 data (estimate per ~1km²). Boxplots with different letters are significantly different according to Tukey's HSD (p-value < 0.05). Region abbreviations: China (CH), Northern Europe (NE), Spain (SP) & Western Europe (WE).

Fig. S11. GWAS results for all phenotypes across Europe. Manhattan plots using 201 (or more) genotypes from Europe (Spain and Northern Europe) as input. The traits are Final Size (upper row), t50 (2nd row) and Slope (lower row) in HL (left column), LL (middle column) and their GxE (right column). The dotted line denotes the 5% Bonferroni-corrected threshold.

Fig. S12. GWAS results for all phenotypes within Spain. Manhattan plots using 117 (or more) genotypes from Spain as input. The traits are Final Size (upper row), t50 (2nd row) and Slope (lower row) in HL (left column), LL (middle column) and their GxE (right column). The dotted line denotes the 5% Bonferroni-corrected threshold.

Fig. S13. GWAS results for all phenotypes within Northern Europe. Manhattan plots using 83 (or more) genotypes from Northern Europe as input. The traits are Final Size (upper row), t50 (2nd row) and Slope (lower row) in HL (left column), LL (middle column) and their GxE (right column). The dotted line denotes the 5% Bonferroni-corrected threshold.

Fig. S14. Correlation of phenotypic traits. Pearson correlations for each pair of traits. Colored boxes show significant correlations (p<0.05 after multiple testing correction (FDR correction) and correction for populations structure (lmekin)) for 195 genotypes across experiments. The significance is illustrated by box size (larger box represents lower p-values) and the color shows the direction and strength of correlation. Abbreviations are: HL= high light, GxE= Genome x Environment interaction, LL= low light, SL = Slope, FT= Flowering time, FS= Final Size, DiamFieldM2= Diameter in Field conditions after 2 Months, Biomass21d= Biomass in controlled (HL) conditions after 21 days.

Fig. S15. Functional enrichment dendrogram for GO enrichment. The enrichment is either based on ranking genes by p-value of the nearest SNP in GWAS (columns 1-9) or Fst of the gene (column 10). The GO terms are arranged into 9 clusters of similar function on the right side of the plot. Depicted are only enrichments with a p-value < 0.001.

Fig. S16. Loss-of-function alleles per population. Based on data from Monroe et al. (2018). The sum of LOF alleles per genotype for Northern Europe (green) and Spain (purple). The regions were not different from each other (GLHT: z-value= 0.634, p-value= 0.526, negative binomial distribution).

Fig. S17. Loss-of-function alleles per population. Based on data from Xu et al. (2019). Boxplot of the sum of LOF alleles per genotype for each region. Boxplots with different letters are significantly different according to Tukey's HSD (p-value < 0.05). Region abbreviations: China (CH), Northern Europe (NE) & Spain (SP).

Fig. S18. Polygenic Scores and regional differentiation for each trait. Summary results from the analysis after Berg and Coop (2014). Each boxplot depicts the polygenic scores of a trait for genotypes from Northern Europe (green) & Spain (purple). Boxplots with different letters are significantly different according to Tukey's HSD (p-value < 0.05). Furthermore, the plot contain information about the number of SNPs used as input, the Qx score for excess variance in SNPs associated with the trait and the p-value of the Qx-analysis. Traits: FS= Final Size, t50, SL= Slope, HL= High Light treatment, LL= Low Light treatment.

Fig. S19. Correlation of phenotypic traits and climate. Pearson correlations for each pair of traits/climatic variable. Colored boxes show significant correlations (p<0.05 after multiple testing correction (FDR correction) and correction for populations structure (lmekin)) for 195 genotypes across experiments. The significance is illustrated by box size (larger box represents lower p-values) and the color shows the direction and strength of correlation. Abbreviations are: HL= high light, GxE= Genome x Environment interaction, LL= low light, SL = Slope, FT= Flowering time, FS= Final Size, DiamFieldM2= Diameter in Field conditions after 2 Months, Biomass21d= Biomass in controlled (HL) conditions after 21 days, Radiation in kJ/m²/day, PC1/2_growS= Principle component 1 and 2 of all climatic data in the estimated growing Season, PC1/2_T= Principle component 1 and 2 for climatic variables related to Temperature, PC1/2_P= Principle component 1 and 2 for climatic variables related to Precipitation.

Table S1. Information on the genotypes used in this study, with their country of origin, assigned region, Genotype name and ID in 1001 Genomes, info on the sampling location and position (latitude and longitude) and the Collector. In the second part of the table the climatic information on the respective location is summarized with: Number of growing months; in the growing season: average Temperature [°C], Soil water content [%], Water vapor pressure [kPa], Wind speed [m s⁻¹], Radiation [kJ m⁻² day⁻¹], Rain [mm]. Afterwards the Bioclim variables 1 t 19 from the Worldclim database (http://worldclim.org/version2). After this the first 2 PCs for PCA on data based on growing season, Temperature variables from bioclim data and preciüitation variables from bioclim data.

Table S2. Raw phenotypic measurements for each plant In the experiment. Replicate is the block the plant was growing in with the corresponding tray number and row and

column for position on the tray (5 rows and 7 columns per tray). The "diam" measurements are diameter measurements where the number corresponds to days after sowing.

Table S3. Genotypic mean of each genotype after correction for positional effects. Information of the usage of genotypes: Phenotype_analysis is 1, if the genotype was used for phenotype-related analysis (regional differentiation, Qst) and GWAS is 1, if the genotype was used in GWAS and following analyses (also GO enrichment & polygenic scores). Additionally data from other experiments that was used for correlations: DiamFieldM2: Diameter in mm in the field in Cologne, after 2 months; Hypocotyllength: length of hypocotyls in mm in HL conditions, 15 days after sowing; Biomass21d: Plant dry weight in g after 21 days after sowing in HL conditions; FT_10/FT_16: flowering time in 10/16 °C from 1001 Genomes, 2016.

Table S4. Estimated heritabilities and pseudo-heritability from EMMAX. Rows contain the input sample size (N), heritability (H^2) and pseudo-heritability for each trait, treatment and population. The p-value of a heritabily is the genotype effect of the mixed linear model. Table S5. Pairwise comparisons of phenotypes for each trait and treatment. The mean difference between traits is given with Z- and p-value from a GLHT of a glm(parameter~population).

Table S6. Correlations between traits and p-values for the trait path analysis. Pearson correlation (r^2) is in the upper right triangle and the corresponding p-values from lmekin after fdr correction are in the lower left triangle. Significant correlations are in bold. The first matrix shows correlations for a dataset with genotypes from Spain (SP) and Northern Europe (NE), the second for Spain and the third for Northern Europe only.

Table S7. Associated SNPs for the different datasets, traits and environment. For each associated SNP the Chromosome, Base, minor allele frequency (MAF), -log10(P) and effect size are given. The LD for the focal SNP was estimated, with the number of SNPs and genes within the LD range. The p-value of two SNPs that exceeded the Bonferroni threshold are marked in bold, the others were just below threshold.

Table S8. Prediction accuracy testing of the polygenic adaptation test. For each phenotype, the number of input SNPs (Nr_SNPs) from GWAS for polygenic scores prediction is given. The input phenotype for the estimation is based on replicate 1 and 2 (Mean(Input)). The phenotype of the 3rd replicate is in Mean(Rep3). Rho and P(Prediction) are the spearman rank correlation and p-value for a correlation between predicted phenotype and phenotype of replicate 3. As a control of prediction accuracy the last 2 columns give Rho and P of the predicted polygenic scores to the input phenotype (replicate 1 and 2).

Table S9. GO-enrichment of genes in LD (within 10kb) to SNPs with p < 0.001 in a GWAS for the respective trait. Shown are terms with an enrichment < 0.001. GO.ID and term give information on the enriched GO term. Annotated states all genes that are in the term, Significant is the number of genes that are associated in the input data set and Expected the number of genes that are expected to be enriched by chance. The resultFisher gives the Fisher score for enrichment. We only report GO terms with >5 genes in them.

Table S10. GO-enrichment of all genes ranked by their Fst or p-value of the closest SNP in a GWAS of the respective trait. Shown are terms with an enrichment < 0.001. GO.ID and term give information on the enriched GO term. Nr_Genes is the number of genes in the respective term. The resultKS gives the Kolmogorov-Smirnov score for enrichment.

Table S11. Results from Polygenic adaptation test after Berg & Coop (2014). Qx is the test statistic for a signal of polygenic adaptation using all phenotypic data. Rho are the results from a spearman correlation of Z-scores predicted versus the phenotypes. The regional Z-values for Northern Europe and Spain are the effect of region on the trait. P-values from each test are in parentheses. The second half of the table contains results from the test without pruning for SNPs in LD.

Literature

1. Mitchell-olds T. Genetic constraints on life-history. Evolution (N Y). 1996;50: 140–145.

2. Wilson AJ, Pemberton JM, Pilkington JG, Clutton-Brock TH, Coltman DW, Kruuk LEB. Quantitative genetics of growth and cryptic evolution of body size in an island population. Evol Ecol. 2007;21: 337–356. doi:10.1007/s10682-006-9106-z

3. Elena SF, Lenski RE. Evolution experiments with microorganisms: The dynamics and genetic bases of adaptation. Nat Rev Genet. 2003;4: 457–469. doi:10.1038/nrg1088

4. Grime JP. Vegetation classification by reference to strategies. Nature. 1974;250: 26–31.

5. Grime JP. Evidence for the Existence of Three Primary Strategies in Plants and Its Relevance to Ecological and Evolutionary Theory. Am Nat. 1977;111: 1169–1194. doi:https://doi.org/10.1086/283244

6. Byers KJRP, Xu S, Schlüter PM. Molecular mechanisms of adaptation and speciation: why do we need an integrative approach? Mol Ecol. 2017;26: 277–290. doi:10.1111/mec.13678

7. Chevin LM, Lande R. When do adaptive plasticity and genetic evolution prevent extinction of a density-regulated population? Evolution (N Y). 2010;64: 1143–1150. doi:10.1111/j.1558-5646.2009.00875.x

8. Bac-Molenaar JA, Granier C, Keurentjes JJB, Vreugdenhil D. Genome wide association mapping of time-dependent growth responses to moderate drought stress in Arabidopsis. Plant Cell Environ. 2016;39: 88–102. doi:10.1111/pce.12595

9. Körner C. Paradigm shift in plant growth control. Curr Opin Plant Biol. 2015;25: 107–114. doi:10.1016/j.pbi.2015.05.003

10. Ballaré CL, Pierik R. The shade-avoidance syndrome: Multiple signals and ecological consequences. Plant Cell Environ. 2017;40: 2530–2543. doi:10.1111/pce.12914

11. Takou M, Wieters B, Kopriva S, Coupland G, Linstädter A. Linking genes with ecological strategies in Arabidopsis thaliana. J Exp Bot. 2019;70: 1141–1151. doi:10.1093/jxb/ery447

12. Kronholm I, Picó F, Alonso-Blanco C, Goudet J, de Meaux J. Genetic basis of adaptation in Arabidopsis thaliana: Local adaptation at the seed dormancy QTL DOG1. Evolution (N Y). 2012;66: 2287–2302. doi:10.5061/dryad.4rp76r87

13. Kerdaffrec E, Filiault DL, Korte A, Sasaki E, Nizhynska V, Seren Ü, et al. Multiple alleles at

a single locus control seed dormancy in Swedish *Arabidopsis*. Elife. 2016;5: e22502.
doi:10.7554/eLife.22502

14. Navarro JAR, Wilcox M, Burgueño J, Romay C, Swarts K, Trachsel S, et al. A study of
allelic diversity underlying flowering-time adaptation in maize landraces. Nat Genet.
2017;49: 476–480. doi:10.1038/ng.3784

15. Hughes PW, Soppe WJJ, Albani MC. Seed traits are pleiotropically regulated by the
flowering time gene PERPETUAL FLOWERING 1 (PEP1) in the perennial Arabis alpina.
Mol Ecol. 2019;28: 1183–1201. doi:10.1111/mec.15034

16. Hallatschek O, Hersen P, Ramanathan S, Nelson DR. Genetic drift at expanding frontiers
promotes gene segregation. Proc Natl Acad Sci U S A. 2007;104: 19926–19930.
doi:10.1073/pnas.0710150104

17. Excoffier L, Foll M, Petit RJ. Genetic Consequences of Range Expansions. Annu Rev Ecol
Evol Syst. 2009;40: 481–501. doi:10.1146/annurev.ecolsys.39.110707.173414

18. Willi Y, Fracassetti M, Zoller S, Van Buskirk J. Accumulation of Mutational Load at the
Edges of a Species Range. Mol Biol Evol. 2018;35: 781–791. doi:10.1093/molbev/msy003

19. Klopfstein S, Currat M, Excoffier L. The Fate of Mutations Surfing on the Wave of a Range
Expansion. Mol Biol Evol. 2006;23: 482–490. doi:10.1093/molbev/msj057

20. Takou M, Hämälä T, Steige KA, Koch E, Dittberner H, Yant L, et al. Maintenance of
adaptive dynamics in a bottlenecked range-edge population that retained out- crossing.
bioRxiv. 2019; 5–10. doi:http://dx.doi.org/10.1101/709873

21. Mitchell-Olds T, Schmitt J. Genetic mechanisms and evolutionary significance of natural
variation in Arabidopsis. Nature. 2006;441: 947–52. doi:10.1038/nature04878

22. Bergelson J, Roux F. Towards identifying genes underlying ecologically relevant traits in
Arabidopsis thaliana. Nat Rev Genet. 2010;11: 867–879. doi:10.1038/nrg2896

23. Hornitschek P, Kohnen M V., Lorrain S, Rougemont J, Ljung K, López-Vidriero I, et al.
Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions
by directly controlling auxin signaling. Plant J. 2012;71: 699–711. doi:10.1111/j.1365-
313X.2012.05033.x

24. Vasseur F, Exposito-Alonso M, Ayala-Garay OJ, Wang G, Enquist BJ, Vile D, et al. Adaptive
diversification of growth allometry in the plant Arabidopsis thaliana. Proc Natl Acad Sci.
2018;115: 3416–3421. doi:10.1073/pnas.1709141115

25. Bac-Molenaar JA, Vreugdenhil D, Granier C, Keurentjes JJB. Genome-wide association
mapping of growth dynamics detects time-specific and general quantitative trait loci. J Exp
Bot. 2015;66: 5567–5580. doi:10.1093/jxb/erv176

26. Hanemian M, Vasseur F, Marchadier E, Gilbault E, Bresson J, Gy I, et al. Transcriptional
natural variation at FLM induces synergistic pleiotropy in Arabidopsis thaliana. bioRxiv.
2019; 658013. doi:10.1101/658013

27. Glander S, He F, Schmitz G, Witten A, Telschow A, de Meaux J. Assortment of Flowering
Time and Immunity Alleles in Natural Arabidopsis thaliana Populations Suggests Immunity
and Vegetative Lifespan Strategies Coevolve. Genome Biol Evol. 2018;10: 2278–2291.
doi:10.1093/gbe/evy124

28. Debieu M, Tang C, Stich B, Sikosek T, Effgen S, Josephs E, et al. Co-Variation between Seed
Dormancy, Growth Rate and Flowering Time Changes with Latitude in Arabidopsis thaliana.
PLoS One. 2013;8: 1–12. doi:10.1371/journal.pone.0061075

29. Vetter MM, Kronholm I, He F, Häweker H, Reymond M, Bergelson J, et al. Flagellin

perception varies quantitatively in arabidopsis thaliana and its relatives. Mol Biol Evol.
2012;29: 1655–1667. doi:10.1093/molbev/mss011

30. Luo Y, Widmer A, Karrenberg S. The roles of genetic drift and natural selection in
quantitative trait divergence along an altitudinal gradient in Arabidopsis thaliana. Heredity
(Edinb). 2015;114: 220–8. doi:10.1038/hdy.2014.89

31. Todesco M, Balasubramanian S, Hu TT, Traw MB, Horton M, Epple P, et al. Natural allelic
variation underlying a major fitness trade-off in Arabidopsis thaliana. Nature. 2010;465:
632–636. doi:10.1038/nature09083

32. Züst T, Agrawal AA. Trade-Offs Between Plant Growth and Defense Against Insect
Herbivory: An Emerging Mechanistic Synthesis. Annu Rev Plant Biol. 2017;68: 513–534.
doi:10.1146/annurev-arplant-042916-040856

33. Davila Olivas NH, Frago E, Thoen MPM, Kloth KJ, Becker FFM, van Loon JJA, et al.
Natural variation in life history strategy of Arabidopsis thaliana determines stress responses
to drought and insects of different feeding guilds. Mol Ecol. 2017;26: 2959–2977.
doi:10.1111/mec.14100

34. Kronholm I, Loudet O, de Meaux J. Influence of mutation rate on estimators of genetic
differentiation--lessons from Arabidopsis thaliana. BMC Genet. 2010;11: 33–47.
doi:10.1186/1471-2156-11-33

35. Fulgione A, Koornneef M, Roux F, Hermisson J, Hancock AM. Madeiran Arabidopsis
thaliana Reveals Ancient Long-Range Colonization and Clarifies Demography in Eurasia.
Mol Biol Evol. 2017. doi:10.1093/molbev/msx300

36. Zou Y, Hou X, Wu Q, Chen J, Li Z, Han T, et al. Adaptation of Arabidopsis thaliana to the
Yangtze River basin. Genome Biol. 2017; 1–11. doi:10.1186/s13059-017-1378-9

37. 1001 Genomes Consortium T. 1,135 Genomes Reveal the Global Pattern of Polymorphism in
Arabidopsis thaliana. Cell. 2016;166: 1–11. doi:10.1016/j.cell.2016.05.063

38. Rasband W. ImageJ. U.S. National Institutes of Health. 2012.

39. De Vylder J, Vandenbussche F, Hu Y, Philips W, Van Der Straeten D. Rosette Tracker: an
open source image analysis tool for automatic quantification of genotype effects. 2012.

40. R-Development-Core-Team. R: A language and environment for statistical computing.
Vienna, Austria: R Foundation for Statistical Computing; 2008. Available: http://www.r-
project.org

41. Ritz C, Baty F, Streibig JC, Gerhard D. Dose-response analysis using R. PLoS One. 2015;10:
1–13. doi:10.1371/journal.pone.0146021

42. Fick SE, Hijmans RJ. Worldclim 2: New 1-km spatial resolution climate surfaces for global
land areas. International Journal of Climatology. 2017.

43. Trabucco A, Zomer R. Global soil water balance geospatial database. CGIAR Consortium for
Spatial Information. Publ Online. 2010. Available: http://www.cgiar-csi.org

44. Dittberner H, Korte A, Mettler-Altmann T, Weber APM, Monroe G, de Meaux J. Natural
variation in stomata size contributes to the local adaptation of water-use efficiency in
Arabidopsis thaliana. Mol Ecol. 2018; 4052–4065. doi:10.1111/mec.14838

45. Pinheiro J, Bates D, DebRoy S, Sarkar D, Team RC. nlme: linear and nonlinear mixed effects
models. 2018.

46. Korte A, Farlow A. The advantages and limitations of trait analysis with GWAS: a review.
Plant Methods. 2013;9: 29–38. doi:10.1186/1746-4811-9-29

47. Hothorn T, Bretz F, Westfall P, Heiberger RM, Schuetzenmeister A, Scheibe S. Package

‘multcomp’ - Simultaneous Inference in General Parametric Models. 2017.

48.   Wei T, Simko V. R package "corrplot": Visualisation of a Correlation Matrix (Version 0.84).
2017.

49.   Epskamp S, Costantini G, Haslbeck J, Isvoranu A, Cramer AOJ, Waldorp LJ, et al. Package
‘qgraph’: Graph Plotting Methods, Psychometric Data Visualization andGraphical Model
Estimation. 2019.

50.   The Inkscape Team. Inkscape. 2007. Available: inkscape.org

51.   Berg JJ, Coop G. A Population Genetic Signal of Polygenic Adaptation. PLoS Genet.
2014;10. doi:10.1371/journal.pgen.1004412

52.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call
format and VCFtools. Bioinformatics. 2011;27: 2156–2158.
doi:10.1093/bioinformatics/btr330

53.   Monroe J, Powell T, Price N, Mullen J., Howard A, Evans K, et al. Drought adaptation in
Arabidopsis thaliana by extensive genetic loss-of- function. Elife. 2018;7: 2–21.
doi:10.7554/eLife.41038

54.   Sohail M, Maier RM, Ganna A, Bloemendal A, Martin AR, Turchin MC, et al. Polygenic
adaptation on height is overestimated due to uncorrected stratification in genome-wide
association studies. Elife. 2019;8: 1–17. doi:10.7554/eLife.39702

55.   Fustier MA, Martínez-Ainsworth NE, Aguirre-Liguori JA, Venon A, Corti H, Rousselet A, et
al. Common gardens in teosintes reveal the establishment of a syndrome of adaptation to
altitude. PLoS Genetics. 2019. doi:10.1371/journal.pgen.1008512

56.   Purcell S, Chang C. PLINK 1.9. 2019. Available: www.cog-genomics.org/plink/1.9/

57.   Pfeifer B, Wittelsburger U, Ramos-Onsins SE, Lercher MJ, Wittelsbürger U, Ramos-Onsins
SE, et al. PopGenome: An Efficient Swiss Army Knife for Population Genomic Analyses in
R. Mol Biol Evol. 2014;31: 1929–1936. doi:10.1093/molbev/msu136

58.   He F, Arce AL, Schmitz G, Koornneef M, Novikova P, Beyer A, et al. The Footprint of
Polygenic Adaptation on Stress-Responsive Cis-Regulatory Divergence in the Arabidopsis
Genus. Mol Biol Evol. 2016;33: 2088–2101. doi:10.1093/molbev/msw096

59.   Yu G, Li F, Qin Y, Bo X, Wu Y, Wang S. GOSemSim: An R package for measuring semantic
similarity among GO terms and gene products. Bioinformatics. 2010;26: 976–978.
doi:10.1093/bioinformatics/btq064

60.   Paradis E, Blomberg S, Bolker B, Brown J, Claude J, Cuong HS, et al. Package ‘ape’:
Analyses of Phylogenetics and Evolution. 2019.

61.   Simons YB, Sella G. The impact of recent population history on the deleterious mutation
load in humans and close evolutionary relatives. Curr Opin Genet Dev. 2016;41: 150–158.
doi:10.1016/j.gde.2016.09.006

62.   Xu YC, Niu XM, Li XX, He W, Chen JF, Zou YP, et al. Adaptation and phenotypic
diversification in arabidopsis through loss-of-function mutations in protein-coding genes.
Plant Cell. 2019;31: 1012–1025. doi:10.1105/tpc.18.00791

63.   Goudet J. HIERFSTAT , a package for R to compute and test hierarchical F -statistics. Mol
Ecol Notes. 2005;5: 184–186. doi:10.1111/j.1471-8278

64.   Leinonen T, McCairns RJS, O'Hara RB, Merilä J. Q(ST)-F(ST) comparisons: evolutionary
and ecological insights from genomic heterogeneity. Nat Rev Genet. 2013;14: 179–90.
doi:10.1038/nrg3395

65.   Whitlock MC, Guillaume F. Testing for spatially divergent selection: Comparing QST to

1020   FST. Genetics. 2009;183: 1055–1063. doi:10.1534/genetics.108.099812

66. Koch EM. The effects of demography and genetics on the neutral distribution of quantitative
traits. Genetics. 2019;211: 1371–1394. doi:10.1534/genetics.118.301839

67. Brachi B, Faure N, Horton M, Flahauw E, Vazquez A, Nordborg M, et al. Linkage and
association mapping of Arabidopsis thaliana flowering time in nature. PLoS Genet. 2010;6:
40. doi:10.1371/journal.pgen.1000940

68. Külheim C, Ågren J, Jansson S. Rapid regulation of light harvesting and plant fitness in the
field. Science (80-). 2002;297: 91–93. doi:10.1126/science.1072359

69. Poorter H, Niinemets Ü, Ntagkas N, Siebenkäs A, Mäenpää M, Matsubara S, et al. A meta-
analysis of plant responses to light intensity for 70 traits ranging from molecules to whole
plant performance. New Phytol. 2019;223: 1073–1105. doi:10.1111/nph.15754

70. Franklin KA. Shade avoidance. New Phytol. 2008;179: 930–944. doi:10.1111/j.1469-
8137.2008.02507.x

71. Méndez-Vigo B, Gomaa NH, Alonso-Blanco C, Xavier Picó F. Among- and within-
population variation in flowering time of Iberian Arabidopsis thaliana estimated in field and
glasshouse conditions. New Phytol. 2013;197: 1332–1343. doi:10.1111/nph.12082

72. Shindo C, Aranzana MJ, Lister C, Baxter C, Nicholls C, Nordborg M, et al. Role of
FRIGIDA and FLOWERING LOCUS C in determining variation in flowering time of
Arabidopsis. Plant Physiol. 2005;138: 1163–1173. doi:10.1104/pp.105.061309.1

73. Zan Y, Carlborg Ö. A Polygenic Genetic Architecture of Flowering Time in the Worldwide
Arabidopsis thaliana Population. de Meaux J, editor. Mol Biol Evol. 2019;36: 141–154.
doi:10.1093/molbev/msy203

74. Price N, Lopez L, Platts AE, Lasky JR. In the presence of population structure: From
genomics to candidate genes underlying local adaptation. Ecol Evol. 2020; 1889–1904.
doi:10.1002/ece3.6002

75. Pritchard JK, Di Rienzo A. Adaptation - not by sweeps alone. Nat Rev Genet. 2010;11: 665–
667. doi:10.1038/nrg2880

76. Csilléry K, Rodríguez-Verdugo A, Rellstab C, Guillaume F. Detecting the genomic signal of
polygenic adaptation and the role of epistasis in evolution. Mol Ecol. 2018;27: 606–612.
doi:10.1111/mec.14499

77. Berg JJ, Harpak A, Sinnott-Armstrong N, Joergensen AM, Mostafavi H, Field Y, et al.
Reduced signal for polygenic adaptation of height in UK Biobank. Elife. 2019;8: 1–47.
doi:10.7554/eLife.39725

78. Braam J, Davis RW. Rain-, wind-, and touch-induced expression of calmodulin and
calmodulin-related genes in Arabidopsis. Cell. 1990. doi:10.1016/0092-8674(90)90587-5

79. Bou-Torrent J, Roig-Villanova I, Galstyan A, Martínez-García JF. PAR1 and PAR2 integrate
shade and hormone transcriptional networks. Plant Signal Behav. 2008;3: 453–454.
doi:10.4161/psb.3.7.5599

80. Holtan HE, Bandong S, Marion CM, Adam L, Tiwari S, Shen Y, et al. Bbx32, an arabidopsis
b-box protein, functions in light signaling by suppressing HY5-regulated gene expression
and interacting with STH2/BBX21. Plant Physiol. 2011;156: 2109–2123.
doi:10.1104/pp.111.177139

81. De Montaigu A, Giakountis A, Rubin M, Tóth R, Cremer F, Sokolova V, et al. Natural
diversity in daily rhythms of gene expression contributes to phenotypic variation. Proc Natl
Acad Sci U S A. 2015;112: 905–910. doi:10.1073/pnas.1422242112

82.   Salmela MJ, Weinig C. The fitness benefits of genetic variation in circadian clock regulation.
Curr Opin Plant Biol. 2019;49: 86–93. doi:10.1016/j.pbi.2019.06.003

83.   Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L. On the accumulation of deleterious
mutations during range expansions. Mol Ecol. 2013;22: 5972–5982. doi:10.1111/mec.12524

84.   Azevedo J, Courtois F, Hakimi M-A, Demarsy E, Lagrange T, Lerbs-mache S, et al.
Intraplastidial trafficking of a phage-type RNA polymerase is mediated by a thylakoid RING-
H2 protein. PNAS. 2008; 2–7.

85.   Jonas T, Rixen C, Sturm M, Stoeckli V. How alpine plant growth is linked to snow cover and
climate variability. J Geophys Res Biogeosciences. 2008;113: 1–10.
doi:10.1029/2007JG000680

86.   Byars SG, Papst W, Hoffmann AA. Local adaptation and cogradient selection in the alpine
plant, Poa hiemata, along a narrow altitudinal gradient. Evolution (N Y). 2007;61: 2925–
2941. doi:10.1111/j.1558-5646.2007.00248.x

87.   Li B, Suzuki J-I, Hara T. Latitudinal variation in plant size and relative growth rate in
Arabidopsis thaliana. Oecologia. 1998;115: 293–301. doi:10.1007/s004420050519

88.   Horton MW, Willems G, Sasaki E, Koornneef M, Nordborg M. The genetic architecture of
freezing tolerance varies across the range of Arabidopsis thaliana. Plant Cell Environ.
2016;39: 2570–2579. doi:10.1111/pce.12812

89.   Dubin MJ, Zhang P, Meng D, Remigereau M-S, Osborne EJ, Paolo Casale F, et al. DNA
methylation in Arabidopsis has a genetic basis and shows evidence of local adaptation. Elife.
2015;4: e05255. doi:10.7554/eLife.05255

90.   Vidigal DS, Marques ACSS, J.Willems LA, Buijs G, Méndez-Vigo B, Hilhorst HM, et al.
Altitudinal and climatic associations of seed dormancy and flowering traits evidence
adaptation of annual life cycle timing in Arabidopsis thaliana. Plant Cell Environ. 2016;39:
1737–1748. doi:10.1111/pce.12734

91.   Frachon L, Mayjonade B, Bartoli C, Hautekeete NC, Roux F. Adaptation to plant
communities across the genome of Arabidopsis thaliana . Mol Biol Evol. 2019.
doi:10.1093/molbev/msz078

1093

2.   Wilson AJ, Pemberton JM, Pilkington JG, Clutton-Brock TH, Coltman DW, Kruuk LEB.
Quantitative genetics of growth and cryptic evolution of body size in an island population.
Evol Ecol. 2007;21: 337–356. doi:10.1007/s10682-006-9106-z

3.   Elena SF, Lenski RE. Evolution experiments with microorganisms: The dynamics and
genetic bases of adaptation. Nat Rev Genet. 2003;4: 457–469. doi:10.1038/nrg1088

4.   Grime JP. Vegetation classification by reference to strategies. Nature. 1974;250: 26–31.

5.   Grime JP. Evidence for the Existence of Three Primary Strategies in Plants and Its Relevance
to Ecological and Evolutionary Theory. Am Nat. 1977;111: 1169–1194.
doi:https://doi.org/10.1086/283244

6.   Chevin LM, Lande R. When do adaptive plasticity and genetic evolution prevent extinction
of a density-regulated population? Evolution (N Y). 2010;64: 1143–1150.
doi:10.1111/j.1558-5646.2009.00875.x

7.   Bac-Molenaar JA, Granier C, Keurentjes JJB, Vreugdenhil D. Genome wide association
mapping of time-dependent growth responses to moderate drought stress in Arabidopsis.
Plant Cell Environ. 2016;39: 88–102. doi:10.1111/pce.12595

8.   Körner C. Paradigm shift in plant growth control. Curr Opin Plant Biol. 2015;25: 107–114.

doi:10.1016/j.pbi.2015.05.003

9.   Ballaré CL, Pierik R. The shade-avoidance syndrome: Multiple signals and ecological
consequences. Plant Cell Environ. 2017;40: 2530–2543. doi:10.1111/pce.12914

10.   Takou M, Wieters B, Kopriva S, Coupland G, Linstädter A. Linking genes with ecological
strategies in Arabidopsis thaliana. J Exp Bot. 2019;70: 1141–1151. doi:10.1093/jxb/ery447

11.   Kronholm I, Picó F, Alonso-Blanco C, Goudet J, de Meaux J. Genetic basis of adaptation in
Arabidopsis thaliana: Local adaptation at the seed dormancy QTL DOG1. Evolution (N Y).
2012;66: 2287–2302. doi:10.5061/dryad.4rp76r87

12.   Kerdaffrec E, Filiault DL, Korte A, Sasaki E, Nizhynska V, Seren Ü, et al. Multiple alleles at
a single locus control seed dormancy in Swedish *Arabidopsis*. Elife. 2016;5: e22502.
doi:10.7554/eLife.22502

13.   Navarro JAR, Wilcox M, Burgueño J, Romay C, Swarts K, Trachsel S, et al. A study of
allelic diversity underlying flowering-time adaptation in maize landraces. Nat Genet.
2017;49: 476–480. doi:10.1038/ng.3784

14.   Hughes PW, Soppe WJJ, Albani MC. Seed traits are pleiotropically regulated by the
flowering time gene PERPETUAL FLOWERING 1 (PEP1) in the perennial Arabis alpina.
Mol Ecol. 2019;28: 1183–1201. doi:10.1111/mec.15034

15.   Hallatschek O, Hersen P, Ramanathan S, Nelson DR. Genetic drift at expanding frontiers
promotes gene segregation. Proc Natl Acad Sci U S A. 2007;104: 19926–19930.
doi:10.1073/pnas.0710150104

16.   Excoffier L, Foll M, Petit RJ. Genetic Consequences of Range Expansions. Annu Rev Ecol
Evol Syst. 2009;40: 481–501. doi:10.1146/annurev.ecolsys.39.110707.173414

17.   Willi Y, Fracassetti M, Zoller S, Van Buskirk J. Accumulation of Mutational Load at the
Edges of a Species Range. Mol Biol Evol. 2018;35: 781–791. doi:10.1093/molbev/msy003

18.   Klopfstein S, Currat M, Excoffier L. The Fate of Mutations Surfing on the Wave of a Range
Expansion. Mol Biol Evol. 2006;23: 482–490. doi:10.1093/molbev/msj057

19.   Takou M, Hämälä T, Steige KA, Koch E, Dittberner H, Yant L, et al. Maintenance of
adaptive dynamics in a bottlenecked range-edge population that retained out- crossing.
bioRxiv. 2019; 5–10. doi:http://dx.doi.org/10.1101/709873

20.   Mitchell-Olds T, Schmitt J. Genetic mechanisms and evolutionary significance of natural
variation in Arabidopsis. Nature. 2006;441: 947–52. doi:10.1038/nature04878

21.   Bergelson J, Roux F. Towards identifying genes underlying ecologically relevant traits in
Arabidopsis thaliana. Nat Rev Genet. 2010;11: 867–879. doi:10.1038/nrg2896

22.   Hornitschek P, Kohnen M V., Lorrain S, Rougemont J, Ljung K, López-Vidriero I, et al.
Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions
by directly controlling auxin signaling. Plant J. 2012;71: 699–711. doi:10.1111/j.1365-
313X.2012.05033.x

23.   Vasseur F, Exposito-Alonso M, Ayala-Garay OJ, Wang G, Enquist BJ, Vile D, et al. Adaptive
diversification of growth allometry in the plant Arabidopsis thaliana. Proc Natl Acad Sci.
2018;115: 3416–3421. doi:10.1073/pnas.1709141115

24.   Bac-Molenaar JA, Vreugdenhil D, Granier C, Keurentjes JJB. Genome-wide association
mapping of growth dynamics detects time-specific and general quantitative trait loci. J Exp
Bot. 2015;66: 5567–5580. doi:10.1093/jxb/erv176

25.   Hanemian M, Vasseur F, Marchadier E, Gilbault E, Bresson J, Gy I, et al. Transcriptional
natural variation at FLM induces synergistic pleiotropy in Arabidopsis thaliana. bioRxiv.

2019; 658013. doi:10.1101/658013

26.   Glander S, He F, Schmitz G, Witten A, Telschow A, de Meaux J. Assortment of Flowering
1157   Time and Immunity Alleles in Natural Arabidopsis thaliana Populations Suggests Immunity
1158   and Vegetative Lifespan Strategies Coevolve. Genome Biol Evol. 2018;10: 2278–2291.
1159   doi:10.1093/gbe/evy124

1160   27.   Debieu M, Tang C, Stich B, Sikosek T, Effgen S, Josephs E, et al. Co-Variation between Seed
1161   Dormancy, Growth Rate and Flowering Time Changes with Latitude in Arabidopsis thaliana.
1162   PLoS One. 2013;8: 1–12. doi:10.1371/journal.pone.0061075

1163   28.   Vetter MM, Kronholm I, He F, Häweker H, Reymond M, Bergelson J, et al. Flagellin
1164   perception varies quantitatively in arabidopsis thaliana and its relatives. Mol Biol Evol.
1165   2012;29: 1655–1667. doi:10.1093/molbev/mss011

1166   29.   Luo Y, Widmer A, Karrenberg S. The roles of genetic drift and natural selection in
1167   quantitative trait divergence along an altitudinal gradient in Arabidopsis thaliana. Heredity
1168   (Edinb). 2015;114: 220–8. doi:10.1038/hdy.2014.89

1169   30.   Todesco M, Balasubramanian S, Hu TT, Traw MB, Horton M, Epple P, et al. Natural allelic
1170   variation underlying a major fitness trade-off in Arabidopsis thaliana. Nature. 2010;465:
1171   632–636. doi:10.1038/nature09083

1172   31.   Züst T, Agrawal AA. Trade-Offs Between Plant Growth and Defense Against Insect
1173   Herbivory: An Emerging Mechanistic Synthesis. Annu Rev Plant Biol. 2017;68: 513–534.
1174   doi:10.1146/annurev-arplant-042916-040856

1175   32.   Davila Olivas NH, Frago E, Thoen MPM, Kloth KJ, Becker FFM, van Loon JJA, et al.
1176   Natural variation in life history strategy of Arabidopsis thaliana determines stress responses
1177   to drought and insects of different feeding guilds. Mol Ecol. 2017;26: 2959–2977.
1178   doi:10.1111/mec.14100

1179   33.   Kronholm I, Loudet O, de Meaux J. Influence of mutation rate on estimators of genetic
1180   differentiation--lessons from Arabidopsis thaliana. BMC Genet. 2010;11: 33–47.
1181   doi:10.1186/1471-2156-11-33

1182   34.   Fulgione A, Koornneef M, Roux F, Hermisson J, Hancock AM. Madeiran Arabidopsis
1183   thaliana Reveals Ancient Long-Range Colonization and Clarifies Demography in Eurasia.
1184   Mol Biol Evol. 2017. doi:10.1093/molbev/msx300

1185   35.   Zou Y, Hou X, Wu Q, Chen J, Li Z, Han T, et al. Adaptation of Arabidopsis thaliana to the
1186   Yangtze River basin. Genome Biol. 2017; 1–11. doi:10.1186/s13059-017-1378-9

1187   36.   1001 Genomes Consortium T. 1,135 Genomes Reveal the Global Pattern of Polymorphism in
1188   Arabidopsis thaliana. Cell. 2016;166: 1–11. doi:10.1016/j.cell.2016.05.063

1189   37.   Rasband W. ImageJ. U.S. National Institutes of Health. 2012.

1190   38.   De Vylder J, Vandenbussche F, Hu Y, Philips W, Van Der Straeten D. Rosette Tracker: an
1191   open source image analysis tool for automatic quantification of genotype effects. 2012.

1192   39.   R-Development-Core-Team. R: A language and environment for statistical computing.
1193   Vienna, Austria: R Foundation for Statistical Computing; 2008. Available: http://www.r-
1194   project.org

1195   40.   Ritz C, Baty F, Streibig JC, Gerhard D. Dose-response analysis using R. PLoS One. 2015;10:
1196   1–13. doi:10.1371/journal.pone.0146021

1197   41.   Fick SE, Hijmans RJ. Worldclim 2: New 1-km spatial resolution climate surfaces for global
1198   land areas. International Journal of Climatology. 2017.

42.   Trabucco A, Zomer R. Global soil water balance geospatial database. CGIAR Consortium for

Spatial Information. Publ Online. 2010. Available: http://www.cgiar-csi.org

43.   Dittberner H, Korte A, Mettler-Altmann T, Weber APM, Monroe G, de Meaux J. Natural
variation in stomata size contributes to the local adaptation of water-use efficiency in
Arabidopsis thaliana. Mol Ecol. 2018; 4052–4065. doi:10.1111/mec.14838

44.   Pinheiro J, Bates D, DebRoy S, Sarkar D, Team RC. nlme: linear and nonlinear mixed effects
models. 2018.

45.   Korte A, Farlow A. The advantages and limitations of trait analysis with GWAS: a review.
Plant Methods. 2013;9: 29–38. doi:10.1186/1746-4811-9-29

46.   Hothorn T, Bretz F, Westfall P, Heiberger RM, Schuetzenmeister A, Scheibe S. Package
'multcomp' - Simultaneous Inference in General Parametric Models. 2017.

47.   Wei T, Simko V. R package "corrplot": Visualisation of a Correlation Matrix (Version 0.84).
2017.

48.   Epskamp S, Costantini G, Haslbeck J, Isvoranu A, Cramer AOJ, Waldorp LJ, et al. Package
'qgraph': Graph Plotting Methods, Psychometric Data Visualization andGraphical Model
Estimation. 2019.

49.   The Inkscape Team. Inkscape. 2007. Available: inkscape.org

50.   Berg JJ, Coop G. A Population Genetic Signal of Polygenic Adaptation. PLoS Genet.
2014;10. doi:10.1371/journal.pgen.1004412

51.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call
format and VCFtools. Bioinformatics. 2011;27: 2156–2158.
doi:10.1093/bioinformatics/btr330

52.   Monroe J, Powell T, Price N, Mullen J., Howard A, Evans K, et al. Drought adaptation in
nature by extensive genetic loss-of-function. Elife. 2018;7: 2–21. doi:10.7554/eLife.41038

53.   Sohail M, Maier RM, Ganna A, Bloemendal A, Martin AR, Turchin MC, et al. Polygenic
adaptation on height is overestimated due to uncorrected stratification in genome-wide
association studies. Elife. 2019;8: 1–17. doi:10.7554/eLife.39702

54.   Fustier MA, Martínez-Ainsworth NE, Aguirre-Liguori JA, Venon A, Corti H, Rousselet A, et
al. Common gardens in teosintes reveal the establishment of a syndrome of adaptation to
altitude. PLoS Genetics. 2019. doi:10.1371/journal.pgen.1008512

55.   Purcell S, Chang C. PLINK 1.9. 2019. Available: www.cog-genomics.org/plink/1.9/

56.   Pfeifer B, Wittelsburger U, Ramos-Onsins SE, Lercher MJ, Wittelsbürger U, Ramos-Onsins
SE, et al. PopGenome: An Efficient Swiss Army Knife for Population Genomic Analyses in
R. Mol Biol Evol. 2014;31: 1929–1936. doi:10.1093/molbev/msu136

57.   He F, Arce AL, Schmitz G, Koornneef M, Novikova P, Beyer A, et al. The Footprint of
Polygenic Adaptation on Stress-Responsive Cis-Regulatory Divergence in the Arabidopsis
Genus. Mol Biol Evol. 2016;33: 2088–2101. doi:10.1093/molbev/msw096

58.   Yu G, Li F, Qin Y, Bo X, Wu Y, Wang S. GOSemSim: An R package for measuring semantic
similarity among GO terms and gene products. Bioinformatics. 2010;26: 976–978.
doi:10.1093/bioinformatics/btq064

59.   Paradis E, Blomberg S, Bolker B, Brown J, Claude J, Cuong HS, et al. Package 'ape':
Analyses of Phylogenetics and Evolution. 2019.

60.   Simons YB, Sella G. The impact of recent population history on the deleterious mutation
load in humans and close evolutionary relatives. Curr Opin Genet Dev. 2016;41: 150–158.
doi:10.1016/j.gde.2016.09.006

61.   Xu YC, Niu XM, Li XX, He W, Chen JF, Zou YP, et al. Adaptation and phenotypic

diversification in arabidopsis through loss-of-function mutations in protein-coding genes.
Plant Cell. 2019;31: 1012–1025. doi:10.1105/tpc.18.00791

62.   Goudet J. HIERFSTAT , a package for R to compute and test hierarchical F -statistics. Mol
Ecol Notes. 2005;5: 184–186. doi:10.1111/j.1471-8278

63.   Leinonen T, McCairns RJS, O'Hara RB, Merilä J. Q(ST)-F(ST) comparisons: evolutionary
and ecological insights from genomic heterogeneity. Nat Rev Genet. 2013;14: 179–90.
doi:10.1038/nrg3395

64.   Whitlock MC, Guillaume F. Testing for spatially divergent selection: Comparing QST to
FST. Genetics. 2009;183: 1055–1063. doi:10.1534/genetics.108.099812

65.   Koch EM. The effects of demography and genetics on the neutral distribution of quantitative
traits. Genetics. 2019;211: 1371–1394. doi:10.1534/genetics.118.301839

66.   Brachi B, Faure N, Horton M, Flahauw E, Vazquez A, Nordborg M, et al. Linkage and
association mapping of Arabidopsis thaliana flowering time in nature. PLoS Genet. 2010;6:
40. doi:10.1371/journal.pgen.1000940

67.   Külheim C, Ågren J, Jansson S. Rapid regulation of light harvesting and plant fitness in the
field. Science (80-). 2002;297: 91–93. doi:10.1126/science.1072359

68.   Poorter H, Niinemets Ü, Ntagkas N, Siebenkäs A, Mäenpää M, Matsubara S, et al. A meta-
analysis of plant responses to light intensity for 70 traits ranging from molecules to whole
plant performance. New Phytol. 2019;223: 1073–1105. doi:10.1111/nph.15754

69.   Franklin KA. Shade avoidance. New Phytol. 2008;179: 930–944. doi:10.1111/j.1469-
8137.2008.02507.x

70.   Méndez-Vigo B, Gomaa NH, Alonso-Blanco C, Xavier Picó F. Among- and within-
population variation in flowering time of Iberian Arabidopsis thaliana estimated in field and
glasshouse conditions. New Phytol. 2013;197: 1332–1343. doi:10.1111/nph.12082

71.   Shindo C, Aranzana MJ, Lister C, Baxter C, Nicholls C, Nordborg M, et al. Role of
FRIGIDA and FLOWERING LOCUS C in determining variation in flowering time of
Arabidopsis. Plant Physiol. 2005;138: 1163–1173. doi:10.1104/pp.105.061309.1

72.   Zan Y, Carlborg Ö. A Polygenic Genetic Architecture of Flowering Time in the Worldwide
Arabidopsis thaliana Population. Mol Biol Evol. 2019;36: 141–154.
doi:10.1093/molbev/msy203

73.   Pritchard JK, Di Rienzo A. Adaptation - not by sweeps alone. Nat Rev Genet. 2010;11: 665–
667. doi:10.1038/nrg2880

74.   Csilléry K, Rodríguez-Verdugo A, Rellstab C, Guillaume F. Detecting the genomic signal of
polygenic adaptation and the role of epistasis in evolution. Mol Ecol. 2018;27: 606–612.
doi:10.1111/mec.14499

75.   Berg JJ, Harpak A, Sinnott-Armstrong N, Joergensen AM, Mostafavi H, Field Y, et al.
Reduced signal for polygenic adaptation of height in UK Biobank. Elife. 2019;8: 1–47.
doi:10.7554/eLife.39725

76.   Braam J, Davis RW. Rain-, wind-, and touch-induced expression of calmodulin and
calmodulin-related genes in Arabidopsis. Cell. 1990. doi:10.1016/0092-8674(90)90587-5

77.   Bou-Torrent J, Roig-Villanova I, Galstyan A, Martínez-García JF. PAR1 and PAR2 integrate
shade and hormone transcriptional networks. Plant Signal Behav. 2008;3: 453–454.
doi:10.4161/psb.3.7.5599

78.   Holtan HE, Bandong S, Marion CM, Adam L, Tiwari S, Shen Y, et al. Bbx32, an arabidopsis
b-box protein, functions in light signaling by suppressing HY5-regulated gene expression

and interacting with STH2/BBX21. Plant Physiol. 2011;156: 2109–2123.
doi:10.1104/pp.111.177139

79. De Montaigu A, Giakountis A, Rubin M, Tóth R, Cremer F, Sokolova V, et al. Natural
diversity in daily rhythms of gene expression contributes to phenotypic variation. Proc Natl
Acad Sci U S A. 2015;112: 905–910. doi:10.1073/pnas.1422242112

80. Salmela MJ, Weinig C. The fitness benefits of genetic variation in circadian clock regulation.
Curr Opin Plant Biol. 2019;49: 86–93. doi:10.1016/j.pbi.2019.06.003

81. Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L. On the accumulation of deleterious
mutations during range expansions. Mol Ecol. 2013;22: 5972–5982. doi:10.1111/mec.12524

82. Azevedo J, Courtois F, Hakimi M-A, Demarsy E, Lagrange T, Lerbs-mache S, et al.
Intraplastidial trafficking of a phage-type RNA polymerase is mediated by a thylakoid RING-
H2 protein. PNAS. 2008; 2–7.

83. Jonas T, Rixen C, Sturm M, Stoeckli V. How alpine plant growth is linked to snow cover and
climate variability. J Geophys Res Biogeosciences. 2008;113: 1–10.
doi:10.1029/2007JG000680

84. Byars SG, Papst W, Hoffmann AA. Local adaptation and cogradient selection in the alpine
plant, Poa hiemata, along a narrow altitudinal gradient. Evolution (N Y). 2007;61: 2925–
2941. doi:10.1111/j.1558-5646.2007.00248.x

85. Li B, Suzuki J-I, Hara T. Latitudinal variation in plant size and relative growth rate in
Arabidopsis thaliana. Oecologia. 1998;115: 293–301. doi:10.1007/s004420050519

86. Horton MW, Willems G, Sasaki E, Koornneef M, Nordborg M. The genetic architecture of
freezing tolerance varies across the range of Arabidopsis thaliana. Plant Cell Environ.
2016;39: 2570–2579. doi:10.1111/pce.12812

87. Dubin MJ, Zhang P, Meng D, Remigereau M-S, Osborne EJ, Paolo Casale F, et al. DNA
methylation in Arabidopsis has a genetic basis and shows evidence of local adaptation. Elife.
2015;4: e05255. doi:10.7554/eLife.05255

88. Vidigal DS, Marques ACSS, J.Willems LA, Buijs G, Méndez-Vigo B, Hilhorst HM, et al.
Altitudinal and climatic associations of seed dormancy and flowering traits evidence
adaptation of annual life cycle timing in Arabidopsis thaliana. Plant Cell Environ. 2016;39:
1737–1748. doi:10.1111/pce.12734

89. Frachon L, Mayjonade B, Bartoli C, Hautekeete NC, Roux F. Adaptation to plant
communities across the genome of Arabidopsis thaliana . Mol Biol Evol. 2019.
doi:10.1093/molbev/msz078

1323

1324
1325

Fig. 1. Regional growth rate estimates in HL and LL. Estimated growth curves averaged over region (from drm function). The growth curves were estimated from diameter measurements at different time points. An illustration of the parameters that are estimated from these growth curves are included in the plot (Final Size is a diameter, t50 a time point and Slope the fold increase in the linear phase). HL (dashed line), LL (solid line), China (orange), Northern Europe (green), Spain (purple) and Western Europe (red).

Fig. 2. Significant regional differentiation of Final Size and t50 in HL and LL. *A.thaliana* genotypes are grouped based on geographical origin. Box plots show regional variation in Final Size (upper row) and t50 (lower row) for HL (left) and LL (right). Groups that do not share a letter are significantly different according to Tukey's HSD (p-value < 0.05). Region abbreviations: China (CH), Northern Europe (NE), Spain (SP) & Western Europe (WE).

Fig. 3. Trait network within SP and NE. Circles represent traits and lines correspond to significant pearson correlations (r²) between traits for genotypes from Spain (SP, solid line) and Northern Europe (NE, line with dashes). The significance of interactions was estimated with a correction for population structure using *Imekin* and was corrected for multiple testing (FDR correction). Red lines represent positive and blue lines negative correlations. The traits are abbreviated: FT10&FT16 are flowering time at 10/16 °C from 1001 Genomes et al. (2016) and VERN is an estimate of vernalization requirement estimated as FT16-FT10. FSHL/FSLL are Final Size in HL/LL. t50H/t50L are t50 in HL/LL. SLHL&SLLL are Slope in HL/LL. Correlations and p-values in Table S6.

Fig. 4. GWAS-results for 4 phenotypes. Manhattan plots of GWAS of t50 in LL with all European genotypes (a), with a peak on Chromosome 1, GxE of Final Size with all European genotypes (b) with a peak on Chromosome 2, Slope in HL within Spain (c) with a peak on Chromsome 3 and t50 in HL in Northern Europe (d) with a peak on Chromosome 1. The dotted line shows the corresponding Bonferroni threshold adjusted for a p-value of 0.05.

Fig. 5. Loss-of-function association and phenotype of t50 in LL/GxE. Manhattan plot of a GWAS with loss-of-funcion alleles and t50LL (a) and t50GxE (c) as input phenotypes with the same association (AT2G17750) above the Bonferroni threshold (blue line). Boxplot of the phenotype of t50LL (b) and t50GxE (d) versus the allele state at AT2G17750 (0 means functional, 1 is a loss-of-function). The colors separate the populations into Spain (purple) and Northern Europe (green).

Fig. 6. Qst. Expected distribution for quantitative trait differentiation between the Spanish and Northern European population. The expectation is based on a multivariate normal distribution assuming a neutral trait with polygenic basis. Horizontal lines indicate observed Qst for the individual growth parameters, FS (Solid line), t50 (Dashed line), Slope (Dot line), in HL (orange) and LL (cyan). The red arrows show the 90th, 95th and 99th percentiles of the distribution.