Title: Conservation genomics reveals low genetic diversity and multiple parentage in the

threatened freshwater mussel, *Margaritifera hembeli*

Nicole L. Garrison[1]

Paul D. Johnson[2]

Nathan V. Whelan[1,3]

1: School of Fisheries, Aquaculture, and Aquatic Sciences, College of Agriculture, Auburn

University, 203 Swingle Hall, Auburn, Alabama 36849, USA

2: Alabama Aquatic Biodiversity Center, Alabama Department of Conservation and Natural

Resources, 2200 Highway 175, Marion, Alabama, USA

3: Southeast Conservation Genetics Lab, Warm Springs Fish Technology Center, US Fish and

Wildlife Service, 203 Swingle Hall, Auburn, Alabama, 36849

Abstract

Margaritifera hembeli is a federally threatened freshwater mussel species restricted to three central Louisiana drainages. Currently, management efforts are being formulated without an understanding of population-level genetic patterns, which could result in sub-optimal conservation outcomes. In particular, information about riverscape genetic patterns is needed to design effective propagation and reintroduction plans. We apply a genomic approach (RADseq) to assess genetic diversity and structure among four wild populations sampled from across the species range. We also assess the genetic diversity of a captively reared cohort produced from a single female. We recovered population differentiation between individuals sampled to the north and south of the Red River. All sites had similarly low levels of heterogeneity and other measures of genetic diversity. The captive cohort displayed higher levels of genetic diversity than expected and likely represents a case of multiple paternity. Future propagation efforts will likely be able to produce genetically diverse cohorts from a small number of wild-caught females, and we recommend future reintroduction efforts utilize brooders within the sub-drainage closest to the reintroduction effort.

Keywords: Margaritiferidae, RADseq, Louisiana Pearlshell, endangered species act, population genomics.

Introduction

In a world of ever-increasing anthropogenic encroachment and climatic change, effective

conservation decisions for species of concern must be made quickly. Regardless of taxon,

detailed genetic assessments clarify the pattern and process of diversity across a landscape,

diagnose specific conservation challenges, and answer crucial questions managers face during

the decision making process (DeSalle and Amato 2004; Segelbacher et al. 2010; Richardson et

al. 2016). Genetic information is invaluable to captive propagation programs, where data play a

role in the selection of broodstock and maintenance of genetic diversity over time (Witzenberger

and Hochkirch 2011). In cases where propagation methods are actively being developed and

reintroduction efforts planned, genetic analysis can assist production and brood stock selection,

better insure successful reintroduction outcomes, and enable continued monitoring of established

populations (Jones et al. 2006; Schwartz et al. 2007; McMurray and Roe 2017). Despite the

potential contribution of genetic information to successful conservation efforts, many species

have little or no genetic data available; this is particularly true of freshwater mussels(Haag and

Williams 2014; Strayer et al. 2019).

Conservation genetic information can now be inferred using thousands of markers from

across a genome, providing a high resolution alternative to historic standbys like single gene or

microsatellite analyses (Luikart et al. 2003; Davey and Blaxter 2010). Importantly, high

throughput methods can be employed in non-model systems with no pre-existing genomic

resources. Population genomic analyses can also directly aid in the development of markers for

continued monitoring of wild and captive populations (Schwartz et al. 2007; Karlsson et al.

2011; Hendricks et al. 2018) and they represent a powerful tool for conservation managers and

stakeholders overseeing long-term recovery programs (Witzenberger and Hochkirch 2011).

Leveraging the power and accessibility of current conservation genomic methods to examine

understudied groups such as freshwater mollusks should be a major focus of conservation

research as anthropogenic pressures increase.

Freshwater mussels are among the most critically imperiled aquatic organisms. A history

of overexploitation and environmental impacts including habitat fragmentation, channel

alterations, agricultural inputs, and pollutants has led to global endangerment of mussels. In the

last 100 years, an estimated 28 species have gone extinct in North America alone. Of the

remaining North American species, approximately 65% are currently at risk of extinction

(Williams et al. 1993; Haag and Williams 2014). The importance of mussels to freshwater

ecosystems globally cannot be overstated. Mussels are long-lived benthic invertebrates that

actively filter suspended particles from water, providing valuable ecosystem services (Vaughn

and Hakenkamp 2001). Furthermore, freshwater mussels are intimately linked to water quality,

and their shells can provide a record of environmental changes, at both recent (Pfister et al. 2011)

and historic scales (Fritts et al. 2017). As part of their complex life cycle, mussels produce

parasitic larvae (glochidia) that must attach and mature on the gills of a host fish. This

relationship can sometimes be very specific (i.e. a single host species) or more general (a

particular family of fishes), and is required for the maturation and dispersal of juvenile mussels

(Wächtler et al. 2001). A history of overexploitation and environmental impacts including habitat

fragmentation, channel alteration, agricultural inputs, and pollution has led to global

endangerment of mussels (Lopes-Lima et al. 2018b). Environmental degradation undoubtedly

impacts mussels directly through physiological stress (Strayer et al. 2004), but may also impair

recruitment via the reduction of host fish population densities (Bogan 1993; Haag 2012) and

their ability to migrate freely (Watters 1996).

Margaritifera hembeli, Louisiana Pearlshell (Fig. 1a), is federally threatened under the

U.S. Endangered Species Act and is restricted to three tributary drainages of the Red River in

central Louisiana (Smith 1988)(Fig. 2). Belonging to the family Margaritiferidae, *M. hembeli* is a

morphologically and phylogenetically distinct group of mussels with only five extant North

American representatives, much fewer than the number of mussel species representing the family

Unionidae (Bogan 2008; Lopes-Lima et al. 2018a). *Margaritifera hembeli* individuals are long

lived (~45-75 years) and inhabit shallow, high velocity stream reaches with relatively stable

substrates (Johnson and Brown 1998; Johnson and Brown 2000). Healthy occurrences of this

species are characterized by large, dense beds (Fig. 1b), sometimes exceeding 300 individuals/m^2

(Johnson and Brown 1998). In the wild, *M. hembeli* glochidia have been found on *Noturus*

phaeus (Brown Madtom), *Luxilus chrysocephalus* (Striped Shiner), *Lythrurus umbratilis* (Redfin

Shiner), and *Notemigonus crysoleucas* (Golden Shiner), but these may be spurious records and

overall host suitability has not been confirmed (Hill 1986; Johnson and Brown 1998). In

particular, host fish specificity in the wild remains unclear given that *M. hembeli* glochidia have

only been found in small numbers on implicated fish species, yet they transform particularly well

on *Esox* spp. species in captivity (see below). Many negative factors influencing mussels on a

global scale are also affecting *M. hembeli*. The 1988 United States Fish and Wildlife (USFWS)

status report for the species noted that *M. hembeli* populations face threats from poor land

management practices (silviculture, gravel mining), reservoir construction, and pollution runoff.

During this initial USFWS assessment of *M. hembeli*, it was known only from 11 headwater

streams in Bayou Boeuf (Rapides Parish, Louisiana), which contributed to the designation of

endangered status under the U.S. Endangered Species Act (USFWS 1988). Upon the discovery

of additional populations in Grant Parish, it was later down listed to threatened (USFWS 1993).

Nevertheless, *M. hembeli* remains in urgent need of conservation attention.

For *M. hembeli* and other mussels, there is considerable interest in captive propagation as

a management tool (Haag and Williams 2014). Currently, most mussel propagation programs

involve the capture of one or a few wild, gravid females followed by inoculation of glochidia on

the appropriate host fish in captivity. Progress has been made in technical aspects of rearing

certain species such as determining temperature thresholds (Steingraeber et al. 2007), appropriate

nutrition (Gatenby et al. 1996), sediment requirements (Jones et al. 2005), host fish identification

(Keller and Ruessler 1997; Hart et al. 2018), and creating capacity for large scale production

(Barnhart 2006). However, genetic impacts of propagation are less clear. Though many

propagation operations exist at federal and state government facilities across the southeast,

genetic consequences of in-use propagation protocols for freshwater mussels is an area of

research in its infancy. Recommendations exist (Jones et al. 2006), but few studies have

explicitly examined the genetic consequences of specific propagation protocols for freshwater

mussels.

An initial transformation and culture trial for *M. hembeli* producing individuals utilized in

this study began in 2016. Transformation of *M. hembeli* juveniles was completed on a single

Esox americanus americanus (Redfin Pickerel) by USFWS (Natchitoches National Fish

Hatchery in Louisiana and the Ecological Services Office in Louisiana). Infection rate was robust

and the effort resulted in nearly 9,000 juvenile *M. hembeli* transformed from the single *E.*

americanus americanus. Once transformation had been completed, culture protocols were

initiated at the Alabama Aquatic Biodiversity Center (AABC) in Marion, Alabama to evaluate

three different methods for rearing *M. hembeli* juveniles. As this initial effort was focused

primarily on determining basic culture methods for *M. hembeli*, transformation of juveniles from

a single gravid *M. hembeli* from Black Creek (Grant Parish, Louisiana) represented an

opportunity to examine genetic variation of progeny found in a single female. Past research has

indicated females of other *Margaritifera* species can mate with multiple males in the wild

(Wacker et al. 2018) and multiple paternity has been noted in several Unionid mussel species

(Christian et al. 2007; Bai et al. 2012; Ferguson et al. 2013). No study has examined this

potential mating system in *M. hembeli*. Given conservation challenges facing *M. hembeli*, the

prospect of reintroducing large numbers of low diversity individuals should be carefully

considered. Understanding of both the genetic structure of wild populations and the opportunity

to contrast against the diversity of a single-female brood could prove invaluable for informing

future conservation and management recommendations for *M. hembeli* reintroduction efforts.

Despite genetic concerns for the management and survival of *M. hemebli,* modern genetic

data has not been generated to inform management decisions. Previous genetic work on *M.*

hembeli found extreme monomorphism ($H_o = 0$) across 25 allozyme markers (Curole et al. 2004).

This observed near-absence of heterozygosity was hypothesized to result from the common (and

likely natural) stochastic extirpation of *M. hembeli* beds, coupled with high re-colonization rates.

A more recent study utilizing microsatellites (Roe 2009) found low heterozygosity at all sampled

populations and some genetic structure among *M. hembeli* populations, but conclusions were

drawn from only five loci. In total, previously generated genetic data offer scant information for

use in designing management efforts. Modern, genomic-level data are necessary to assess

population connectivity, measure genome-wide genetic diversity, and provide managers with

useful information that can be used to inform propagation and reintroduction efforts.

The goals of this study were to 1) assess the genetic diversity of the 2016 AABC captive

M. hembeli cohort 2) determine whether previous reports of low genetic diversity of wild *M.*

hembeli populations using allozymes is supported by genomic markers and 3) and characterize

patterns of gene flow or potential barriers to aid in reintroduction of future culture efforts. To

accomplish these objectives, a restriction enzyme associated high-throughput sequencing method

was employed (RADseq) to generate a dataset of thousands genomic loci. RADseq is capable of

utilizing genomic DNA samples collected non-lethally to generate millions of sequence reads for

each individual. These reads can be processed with existing tools to identify thousands of single

nucleotide polymorphisms (SNPs) from across the genome, providing high resolution data for

determining demographic and evolutionary dynamics within a species. Ultimately, these data

will inform recovery efforts of *M. hembeli* and provide the first genetic profile of a propagated

cohort for this threatened freshwater mussel.

Materials and Methods

Sample Collection and Sequencing

We sampled 20 individuals each from four sites in the Red River drainage in Louisiana,

and 20 captively reared individuals produced from a single wild caught female from Black

Creek. Two sampling localities were from Rapides Parish (Brown Creek – Bayou Rapides and

Loving Creek – Bayou Boeuf) and two were from Grant Parish (Jordan Creek and Black Creek –

Bayou Rigolette). Sampling sites were selected for ease of access, abundance of *M. hembeli*, and

because they encompass the major drainages from which *M. hembeli* is known to occur (Fig. 2).

Mussels were collected by hand and effort was taken to select a broad size range likely

representing individuals from multiple age classes. A sterile, individually wrapped buccal swab

was used to collect genetic material from the foot of each mussel and immediately placed in

swab stabilization buffer from the Buccal-PrepPlus DNA isolation kit (Isohelix™). Mussels were

photographed on site and returned immediately to their bed after swabbing. The captively reared

individuals were lethally sampled at the AABC and a clip of the foot was placed in 96-100%

ethanol until DNA extractions could be completed. The mother of the captive cohort was not

available for sampling and a tissue voucher/buccal swab was never taken. Shells of captively

reared individuals were deposited at the Auburn Museum of Natural History (AUM 45578-

45595).

DNA extractions of buccal swabs and tissue were completed with Isohelix Xtreme DNA

isolation kit and the Qiagen DNeasy blood and tissue kit respectively, following the

manufacturers' instructions. After DNA extraction, each sample was treated with RNase A at a

final concentration of 100 µg/ml and incubated at 37°C for 15 minutes to remove any co-purified

RNA. Each extraction was quantified with a Qubit Fluorometer and checked for integrity of high

molecular weight DNA through standard gel electrophoresis. Samples were standardized to a

concentration of 20 ng/µL and 50 µL of standardized DNA was sent to Floragenex Inc.

(Portland, OR) for RADseq library preparation using the SbfI restriction enzyme following Baird

et al. (2008). Samples were tagged with unique barcode identifiers, pooled, and sequenced in

three replicate lanes on the Illumina HiSeq 4000 platform using 100bp paired-end chemistry.

Sequence Processing and Variant Identification

Reads from all three sequenced lanes were combined for each individual and processed

with the STACKS v2.1 pipeline for population genomic analysis (Rochette et al. 2019).

STACKS demultiplexes raw sequencing data, aligns reads to form stacks of loci, identifies

variants (i.e. SNPs), and generates descriptive population statistics. Paired-end reads were

demultiplexed with the *process_radtags* command using default settings that allow for barcode

sequences to be rescued if the barcode varies by only one nucleotide from the expected sequence.

Input file preparation details and *process_radtags* settings can be found online

(https://github.com/nlgarrison/ConservationGenomics). The denovo_map.pl script was used to

automate the STACKS pipeline, as a reference genome is not available for *M. hembeli* or any

closely related species. Stack assembly required a minimum of five reads per locus (-m 5),

allowed for three mismatches within stacks of the same individual initially (-M 3) and two

mismatches between stacks of different individuals (-r 2). All other parameters were set to

default. To identify SNPs in stacked loci, a catalog of potential sites must be formed; in our

pipeline only individuals from the wild populations were used to generate the catalog. To do this,

at the *cstacks* step in the denovo_map.pl pipeline, a population map including only wild-caught

individuals was provided to the catalog building command. This approach was chosen to reduce

bias in called variants that might be introduced by uneven population sampling, and it allowed

for a true test of population assignment for the captive individuals.

We ran the *populations* command on two sets of samples: wild-caught individuals only

(NoCaptive) and all sampled individuals (WithCaptive). The captive cohort was excluded in

initial runs of *populations* to eliminate bias when estimating the baseline population parameters

for wild individuals. The NoCaptive dataset was analyzed with a minimum minor allele

frequency setting of 0.025, maximum heterozygosity setting of 0.50, and a requirement that a

variant be present in at least three sampling sites and at least 50% of individuals within each

sampling site; these settings were also used in the analysis of the WithCaptive dataset. For both

the NoCaptive and WithCaptive datasets, one *populations* analysis was generated allowing only

one random SNP per locus (denoted "S") and a second was produced allowing multiple SNPs

per locus (denoted "M"), resulting in a total of four datasets; NoCaptiveS, NoCaptiveM,

WithCaptiveS, WithCaptiveM. This was done because some downstream analyses assume

unlinked loci, whereas others can use multiple SNPs originating from the same locus. For

downstream analyses requiring subsets of individuals as input, the program VCFtools (Danecek

et al. 2011) was used to generate reduced datasets as needed from the WithCaptiveS/M vcf files.

Population Genomic Analyses

For each dataset, average heterozygosity, nucleotide diversity, pairwise F_{ST} among each

sampling site, and F_{IS} at each sampling site was reported by *populations*. The *basicStats* function

of the R (R CORE Team 2019) package diveRsity (Keenan et al. 2016) was used to calculate

allelic richness. We assessed population structure among sampling sites with an Analysis of

Molecular Variance (AMOVA; Excoffier, Smouse, and Quattro 1992) and a series of clustering

methods. AMOVA was performed using the *poppr.amova* command in the R package adegenet

(Kamvar et al. 2014; Jombart et al. 2018). Individuals were stratified by sample site and whether

the site was located north or south of the Red River to assess whether the Red River serves as a

barrier to gene flow or if genetic structure is better explained by local landscape characteristics.

Significance was tested with a 500 permutation randomization test.

Discriminant analysis of principal components (DAPC) was implemented using the R

package adegenet following Jombart and Collins (2015), using the NoCaptiveM and

WithCaptiveM datasets. The best-fit number of clusters (K) was assessed using *k*-means

clustering with Bayesian information criteria. The *snmf* function of the R package LEA (Frichot

and François 2015) was used to identify clusters of individuals and determine admixture

proportions, and the best-fit K was determined using the cross-entropy criterion. For these

analyses, only the unlinked SNPs (NoCaptiveS, WithCaptiveS) were used. Though similar in

function to STRUCTURE (Pritchard et al. 2000), LEA can be more accurate than STRUCTURE

in the face of inbreeding (Frichot et al. 2014) and handles genomic data more efficiently.

We also used the model-based method fineRADstructure (Malinsky et al. 2018) to

generate a summary of haplotype coancestry for all individuals. A major advancement provided

by fineRADstructure is its ability to utilize linkage and polymorphism in genomic data without a

reference genome, allowing fine scale patterns of relatedness among individuals to be examined.

This analysis was conducted with the NoCaptiveM, WithCaptiveM, and the captive cohort in

isolation. The captive cohort was examined independently in order to make more accurate

inferences about potential multiple paternity.

We tested for a signature of isolation-by-distance using a Mantel test of correlation

between geographical distance and pairwise F_{ST} values for the wild populations. Geographic

distances among sites were measured by plotting sample collection points in QGIS (QGIS

Development Team 2014) and hand tracing stream distance between sampling sites in a pairwise

fashion. As river connections were sometimes difficult to assess and to account for minor

idiosyncrasies associated with tracing river path, hand tracing was repeated three times for each

pair and an average of distances in meters was taken as the geographic stream distance. F_{ST}

values used for the Mantel test were those reported by STACKS (Table 1). The Mantel test was

done using the R package 'ade4' (Dray, Dufour, et. al 2007); significance was evaluated with

1,000 random permutations. As Mantel tests have received criticism for use as a measure of

isolation by distance (Legendre et al. 2015; Meirmans 2015), we also performed a multiple

regression on distance matrices with the MRM function of the Ecodist R package (Goslee and

Urban 2007); significance was tested with 10,000 permutations.

The programs diveRsity and Migrate-n version 4.2.14 (Beerli and Palczewski 2010) were

used to evaluate connectivity between wild populations of *M. hembeli*. The *divMigrate* function

of the R package diveRsity uses differences in allele frequencies to model asymmetric, relative

rates of migration between populations (Sundqvist et al. 2016). Though it was designed for use

with microsatellite data, *divMigrate* is capable of handling genomic SNP datasets and has been

shown to reflect biologically realistic scenarios of population connectivity in recent studies (e.g.

Woodings et al. 2018; Manuzzi et al. 2019). An analysis of the WithCaptiveS dataset was done

with the *divMigrateOnline* implementation (Keenan 2012) to calculate relative rates of migration

between wild populations using a cutoff value of 40, an alpha of 0.05, and the G_{ST} statistic (Nei

1973). Support for the asymmetry of migration rates was evaluated with 1000 bootstrap

replicates. Although the captive population was included in our initial *divMigrate* analysis, due

to its high similarity ($G_{ST} = 1.0$) with Black Creek individuals it was masked using the "exclude

population" option without recalibrating rates.

The *divMigrateOnline* analysis was complemented by the Bayesian population genetics

program migrate-n. Given computational demands of migrate-n, a subset of 100 polymorphic

loci were randomly selected from the NoCaptiveM dataset. Migrate-n analyses were done with

loci in their entirety, rather than individual SNPs, as the SNP model has not been thoroughly

tested (see migrate-n manual). Loci that appeared more than once in the random subset were

filtered out, leaving 95 polymorphic loci for inclusion in the migrate-n analysis. Geographic

distances between sites were calculated as described previously. Five migration models were

investigated as indicated by geography and other population genomic analyses 1) full migration

2) northern panmixia with unidirectional gene flow from Loving Creek to Brown Creek 3)

northern panmixia with bidirectional gene flow between Loving Creek and Brown Creek 4)

panmixia (Fig. 3). For each migration model, Bayesian inference was performed using the DNA

sequence model (sampling of 20,000,000 total steps, 10,000 steps discarded as burn-in, default

priors) and combined with thermodynamic integration (four parallel heated chains) at

temperatures 1, 1.5, 3.0, and 1×10^6. Log marginal likelihood values were calculated with Bezier

approximation within migrate-n and log Bayes factors were used to rank the models following

Beerli & Palczewski (2010).

Effective population size (N_e) and the number of effective breeders (N_b) was estimated

for each sampling site, geographically proximate wild populations combined, and the captive

cohort in isolation with NeEstimator2 (Do et al. 2014) using the NoCaptiveS and WithCaptiveS

datasets as input. Given results indicating that the Red River represents a barrier to gene flow

(see below), we combined sites to assess N_e of "northern" (Black Creek and Jordan Creek) and

"southern" (Loving Creek and Brown Creek) populations. The molecular coanscestry method of

Nomura (2008) and linkage disequilibrium method (Hill 1981; Waples 2006) as implemented in

NeEstimator2 were used to evaluate N_e and N_b without a genomic map.

The program COLONY (Jones and Wang 2010) was used to validate suspected multiple

paternity in the captive cohort. A captive-only subset of SNPs filtered from the WithCaptiveM

dataset using vcftools was used as input. Three different filtering strategies were tested, allowing

sites with missing data proportions of 0%, 25%, and 50%. Files were converted to COLONY

input format using tidy_vcf and the write_colony function of the R package radiator (Gosselin et

al. 2020). Long, full likelihood runs were performed with 10 replications for each filtered

dataset; male and female polygamy settings were used and half sibship of all individuals with an

unknown mother was specified in the input file.

Results

      Sequencing resulted in >460 million reads per lane and >1 million reads per individual

(n=94). Sequence reads filtered out due to barcode ambiguity, unclear restriction cut sites, and

potential Illumina adapter contamination comprised ~15% of the total raw reads; 1.4 billion

paired-end reads were included in the assembly process. We saw no evidence of contamination

in our samples and most loci present in Black Creek were also present in the Captive Cohort,

indicating that there was no bias associated with DNA sampling method in our study.

Furthermore, there were no detectable differences in the quality or quantity of sequences

obtained from the two sample collection methods (swab and tissue clips). The STACKS pipeline

identified 1,185,792 SNPs across 20,464 loci, further filtered by *populations* program constraints

to 2,563 putatively independent variant SNPs across the wild populations. When multiple SNPs

per locus were allowed, the number of variant SNPs recovered across all wild populations

increased to 7,601. After initial runs of the STACKS pipeline, four individuals having less than

half the identified SNPs found in the rest of the samples (Jordan18, Captive17, Loving20, and

Brown5) were removed, resulting in datasets generated for downstream analyses which included

90 individuals from the original set.

      The vast majority of markers sampled were not variable across the four wild populations;

out of ~20,000 potential markers (i.e. SNPs), about 10%, met the population parameters

specified and contained informative variation. Average nucleotide diversity (π) ranged from

0.21-0.23. Observed heterozygosity was considerably lower than expected heterozygosity across

all sites ($H_{obs} = 0.08\text{-}0.09$, $H_{exp} = 0.20\text{-}0.22$), likely indicating genetic bottlenecks at all sites.

Allelic richness was also similar across all sampled groups (1.16-1.51). Analysis of the captive

population generated 2,416 variable SNPs (8,069 when multiple SNPs per locus were allowed)

with an average observed H_{obs} of 0.10 and an average nucleotide diversity of 0.25. Pairwise F_{ST}

values were low, with the Brown Creek population presenting the highest differentiation from

other populations (0.063-0.082; see Table 1 for all pairwise F_{ST} values); the captive cohort and

the Black Creek population had the lowest pairwise F_{ST} (0.035). F_{IS} values for each collection site

ranged from 0.44 to 0.48 and was 0.39 for the captive cohort.

Genetic structure was seen among collection sites to varying degrees depending on

analytical method. AMOVA was significant at all hierarchical levels ($p = 0.001$), but only 3.89%

of genetic variation was explained by whether the sampling site was north or south of the Red

River. A further 6.7% of genetic variation was explained by collection sites within regions,

indicating landscape barriers to gene flow that are more complex than one major river. The

Mantel test and multiple regression for isolation-by-distance was significant ($p < 0.05$),

suggesting isolation by distance patterns in the data. DAPC analysis of wild individuals indicated

that a K of 4 most accurately captured the diversity of the samples, with individuals from Jordan

and Black (Grant Parish) overlapping and Brown and Loving (Rapides Parish) each forming

their own distinct cluster (Fig. 4a). When all populations were included, captive mussels

overlapped entirely with the Black Creek cluster (Fig. 4b). Notably, the captive mussels showed

similar spread in the DAPC analysis as wild individuals from Black Creek, indicating the captive

cohort possesses variation in genetic diversity comparable to wild populations.

Analysis with LEA suggested data were best explained by two genetic clusters when only

wild populations were included. LEA analysis with K = 2 showed that individuals from sites

north of the Red River (Grant Parish) had more similar admixture profiles to each other than

those from south of the Red River (Rapides Parish) and *vice versa* (Fig. 5). Hierarchical

clustering analysis with fineRADstructure, mirrored DAPC and LEA analyses (Fig. 6). Brown

Creek and Loving Creek formed groupings, to the exclusion of most other individuals, with

particularly high coancestry values detected within the Brown Creek population. Individuals

from Jordan Creek, Black Creek, and the Captive cohort are nearly indistinguishable from each

other with only minor internal structuring comprising a subset of Black Creek and Captive

individuals (Fig. 6). When examined in isolation, the captive cohort displayed structured

relationships at varying levels of coancestry; at the top of the coancestry value range a cluster of

individuals (Cap3, Cap10, and Cap4) may represent a grouping of full siblings (Fig. 7).

The program *divMigrateOnline* detected relatively high migration rates between the two

northern sampling sites, Jordan Creek and Black Creek. To a lesser degree, the analysis showed

migration from the northern populations to Loving Creek and unidirectional movement from

Loving Creek to Brown Creek (Fig. 8). The asymmetry of migration rates inferred by

divMigrateOnline were not supported by 1000 bootstrap replicates, meaning that while gene

flow was detected between Jordan Creek and Black Creek, no strong directionality could be

inferred. Of the models assessed by migrate-n, the most likely was model 3 (Table 3) with

populations occurring north of the Red River being panmictic, bidirectional gene flow from the

panmictic northern population to Loving Creek, and bidirectional migration between Loving

Creek and Brown Creek.

Estimates of effective population size for some populations in isolation resulted in

negative or infinite values, suggesting that the values were driven by sampling error as a result of

insufficient sample size or marker informativeness (Marandel et al. 2019). Combining

individuals from the northern sites (Jordan Creek, Black Creek) and southern sites (Brown Creek

and Loving Creek) allowed for more realistic estimation of effective population sizes (Table 2).

The captive cohort generated the lowest estimates of N_e (86 -111 individuals) but an N_b of 5.

This high estimate for the number of effective breeders within a cohort known to originate from

a single female is a strong indication of multiple paternity.

COLONY also indicated multiple paternity was present in the captive cohort samples, but

the point estimates of male input varied with the level of missing data allowed into the analysis.

Using the most complete dataset (sites every individual shared), COLONY output indicated

several full-sibling clusters and 7 fathers. The most relaxed filtering strategy (50% site coverage)

indicated a different father for each individual in the sample (n = 19). Estimates of N_e produced

by COLONY (N_e = 4, 95% CI = 2-12) were stable across analyses and reflected both the

NeEstimator2 values for the captive samples (Table 2) and the fineRADstructure clustering

analysis (Fig. 7).

Discussion

Population genomic data revealed low levels of heterozygosity across *M. hembeli*

populations and complex patterns of gene flow among sites. Although low levels of genetic

diversity in wild populations is concerning from a conservation standpoint, our results are similar

to past studies on *M. hembeli* (Curole et al. 2004; Roe 2009). Moreover, low genetic diversity

and a signature of bottlenecks at every sampling site may reflect natural processes such as rapid

colonization after the loss of a habitat that results from stream meanders and cutoffs that are

common in low-elevation, flat terrains like those in central Louisiana. Despite low genetic

diversity, high gene flow among some sample sites was observed. Genetic structure generally

followed a pattern of isolation-by-distance, but the Red River also appeared to be a factor in

observed genetic structure. This suggests movement between populations on the same side of the

Red River may occur during major flood events, thereby facilitating migration with infected

fishes moving to adjacent stream channels and does not require host-fish passage through the

Red River channel.

Our data showed a captive cohort of *M. hembeli* produced from a single female had

comparable genetic diversity to the wild population from which the female was sampled.

Broadly, analyses indicate evidence of multiple paternity, which has considerable implications

for propagation efforts of *M. hembeli*. Although multiple paternity has been known to occur in at

least some freshwater mussels (Christian et al. 2007; Bai et al. 2012; Ferguson et al. 2013;

Wacker et al. 2018), more recent studies have empirically demonstrated wild female mussels can

mate with multiple males to produce genetically diverse captive cohorts (Wacker et al. 2019).

Multiple paternity had not been empirically observed in *M. hembeli* prior to this study. Results

suggest progeny generated from a few (randomly selected) wild-fertilized females annually

would mirror the heterogenity found in the wild host populaton, likely resulting in low detectable

differences between propagated and wild mussels with a recovery effort that spans multiple

years.

Genetic Diversity and Effective Population Size

Low observed heterozygosity relative to expected heterozygosity (Table 2), suggests *M.*

hembeli populations are small and experiencing associated effects such as inbreeding and genetic

bottlenecks. This is also reflected by high F_{IS} values (Table 2). Given the threatened status of *M.*

hembeli under the U.S. Endangered Species Act, anthropogenic activities have clearly caused

severe population declines. However, life history, natural demographics, and stochastic habitat

disturbances probably also play a possibly an outsized role, in influencing low heterozygosity of

M. hembeli. For instance, a single beaver dam once led to the extirpation of an *M. hembeli* bed

(~1000 individuals) located on Forest Service land (Stewart 1990). Furthermore, Johnson and

Brown (2000) showed that channel stability of the study area on the time-scale of a single year

can be low, even though they noted that *M. hembeli* appeared to be associated with relatively

stable microhabitats. Assuming rapid colonization ability of *M. hembeli* (see Curole, Foltz, and

Brown 2004), low heterozygosity is likely the result of repeated bottlenecks caused by habitat

destruction followed by colonization of newly exposed suitable habitat. Although natural

destruction of mussel beds and rapid colonization of new habitats may be a normal part of *M.*

hembeli biology, habitat fragmentation caused by anthropogenic activity exacerbates

contemporary population decline by restricting recruitment across populations (Geist and

Auerswald 2007). The habitat fragmentation witnessed through modification of waterways or

changes in landuse are likely limiting re-colonization options for *M. hembeli*.

When considered in isolation, N_e estimates for all sampling sites except Black Creek

were inferred to be infinite, which is indicative of N_e estimates being driven by sampling error,

rather than large population sizes (Marandel et al. 2019). Pooling individuals from north and

south of the Red River, respectively, allowed for more precise estimates of N_e (Table 2).

Effective population size is difficult to accurately estimate even with genome-wide markers as

many methods make assumptions that are typically violated such as sampling of non-overlapping

generations (Waples et al. 2016; Wang et al. 2016; Marandel et al. 2019). The linkage

disequilibrium method applied here is known to be downwardly biased (as much as 30%) when

samples consist of different age classes (Waples et al. 2014). We anticipate that our estimates of

N_e for wild populations are much lower than reality, as multiple age classes were sampled. Thus,

N_e estimates reported here may be of value to resource managers, but they should be approached

with caution and not the sole genetically derived metric used for assessing populations of *M.*

hembeli. The temporal trajectory of N_e is more crucial to conservation managers than a point

estimate, and we recommend additional sampling of size/age class cohorts to reveal trends

through time.

Population Connectivity

In general, analyses determined genetic demarcation between populations occurring north

of the Red River in those occurring south of it. However, fine-scale relationships and migration

patterns among populations that our SNP-based approach illuminated are novel. Sampled sites

appeared to demonstrate an isolation-by-distance effect, which is a general pattern seen in many

freshwater organisms (Meffe and Vrijenhoek 1988; Whelan et al. 2019). Furthermore, AMOVA

indicated significant genetic structure with a genetic break occurring between collection sites

north and south of the Red River, with further significant genetic structure between populations

in the north and south. These patterns were recovered to varying degrees by clustering analyses,

with DAPC indicating at least some genetic distinctness among all four sites (Fig. 4).

Analyses that examined finer-scale gene-flow patterns indicated a high amount

connectivity between Grant Parish populations (Jordan Creek and Black Creek), with the best-fit

model as assessed by migrate-n having a panmictic Jordan Creek and Black Creek (Fig.

2).Whether the observed pattern is a result of active gene flow or a historical connection is

unknown. Black Creek has been putatively isolated from Jordan Creek for the past 60 years

behind Iatt Lake, but the long lifespan of *M. hembeli* may result in a longer time-period for

genomic signatures of isolation to be detectable. Documentation of Iatt Lake's management

history (Moses et al. 2016) indicates *Esox americanus americanus*, was not detected in fish

surveys but other *Esox* species (*Esox americanus vermiculitus*, *Esox niger*) have been collected

during Iatt Lake surveys. Iatt Lake is prone to flooding and has experienced several high water

events in recent history (Moses et al. 2016), which could facilitate fish passage between

tributaries. Assuming the presence of *Esox* spp. around Lake Iatt and periodic flooding it is

possible that a connection was recent and possibly intermittent between Black Creek and Jordan

Creek drainages.

Relative genetic homogeneity among northern sites contrasts with the relative isolation of

those sampled from south of the Red River. Loving Creek and Brown Creek shared similar

genomic backgrounds, as indicated by LEA (Fig. 4), but *divMigrate* and migrate-n analyses

indicated somewhat limited gene flow between the two southern sites. Several analyses indicated

that Brown Creek was the most isolated group of individuals sampled, which may be partially

explained by a higher stream distance from its Red River confluence compared to other sites.

Connectivity analyses appear to indicate that Loving Creek represents Brown Creek's only

connection to the rest of the species range (Fig. 6). Though there are sites not sampled during the

course of this study which occur in streams located between Loving and Brown Creek, our data

still indicates that movement of *M. hembeli* in the southern part of its range is relatively more

restricted than in the northern section of its range.

Further natural history work is needed to better understand the conditions required for

successful dispersal of *M. hembeli*. Although a detailed host fish trial has not been completed for

M. hembeli, formal trials have been completed for *M. marrianae*, the Alabama Pearlshell. These

trials indicate *Esox vermicularis vermicularis* is the primary host in Alabama. A minimal

transformation rate was also documented for *Noturus leptacanthus* (Speckled Madtom) but it's

not a primary host for *M. marrianae* (Fobian et al., *in prep.*). In contrast, *M. hembeli* readily

transforms transform on *Esox* spp. in captivity (Schmidt-Frater, *pers. comm.*) but a formal host

trial has not been completed. Additionally, several non-sampled populations of *M. hembeli* occur

in headwater streams isolated behind reservoirs – future studies including those sites will likely

provide further insights into the impact of such barriers on the dispersal of *M. hembeli* across its

range.

Captive Propagation and Reintroduction

Our analysis of captively reared individuals revealed a surprising amount of genetic

diversity given that the cohort was produced by a single female from Black Creek. Notably,

genetic diversity estimates for this cohort were virtually identical to estimates from the wild

population (Table 2). Clustering analyses provided evidence these data were more than sufficient

to assign captive individuals to their population of origin, always grouping them with wild Black

Creek individuals (Fig. 4b). Multiple paternity was also evident based on estimates of the

number of effective breeders for the captive cohort (N_b =5). Coupled with inferences from

COLONY and fineRADstructure, where multiple paternal genotypes (7-19) and multiple clusters

of individuals with high (but not identical) co-ancestry were observed, our study supports the

presence of a multiple paternity strategy within *M. hembeli*. Overall, this represents a best-case

scenario for managers as a limited number of wild-caught *M. hembeli* females can be brought

into captivity annually and produce a genetically diverse cohort for reintroduction efforts.

However, given the difficulty of *Margaritifera* spp. production in a hatchery setting (Paul

Johnson *pers. obs.*), any serious reintroduction effort would likely be a decades-long endeavor at

minimum.

The ability to produce genetically diverse individuals from a small number of females

should facilitate propagation programs; however, care must still be taken when choosing brood

stock and determining reintroduction sites. Managers should be guided by our findings of

population structure and isolation by distance. Broadly, brood stock should be as geographically

proximate to the chosen reintroduction site as possible, coming from one or a few sites within the

same drainage. At the very least, broodstock should come from the same side of the Red River as

the chosen reintroduction site. Furthermore, genetic diversity and number of effective breeders

for another Margaritiferid, *M. margaritifera*, was recently determined to be higher when females

were fertilized in the wild relative to those fertilized in captivity (Wacker et al. 2019). Our work

suggests multiple paternity is likely the case for *M. hembeli* as well. The best chance we have at

maintaining appropriate levels of diversity is to utilize wild-fertilized broodstock while it is still

available, rather than attempt to establish a captive breeding colony of *M. hembeli*.

Our data also give reasons to avoid augmentation (i.e., placing captively reared

individuals on top of a natural population) in favor of reintroductions (i.e., placing captively

reared individuals at a site from which *M. hembeli* has been extirpated). Each population

analyzed here was considered genetically distinct in at least one analysis, albeit with limited or

no genetic distinction between Black Creek and Jordan Creek in most analyses. That said,

consequences of outbreeding depression are impossible to predict at this time, and augmentation

violates recent recommendations for freshwater mussel propagation and release (Mobile River

Basin Mollusk Restoration Committee 2010; Cumberlandian Region Mollusk Restoration

Committee 2010; Strayer et al. 2019). Our data support such recommendations. If managers are

faced with no suitable sites for propagation and release other than sites with natural *M. hembeli*

population, then we argue that habitat restoration should be a higher priority than captive

propagation of *M. hembeli*. Good animals placed into poor habitat will not have the desired

outcome (Geist and Auerswald 2007).

Conclusions

This study provides information that can be used to facilitate successful propagation

efforts and a framework for studying the existing diversity in imperiled mussel species using

modern methods. We have demonstrated that genetically diverse cohorts of margaritiferids may

be produced from a small number of wild-caught, gravid females. Importantly, our findings also

indicate that the occurrence of multiple paternity in freshwater mussels may be more widespread

than the limited number of explicitly documented cases. Our findings can likely be generalized

to closely related species such as the federally endangered Alabama Pearlshell, *M. marrianae,*

the sister species to *M. hembeli*, which is currently the focus of intense propagation and

management efforts. More broadly, we have demonstrated the utility of RAD-seq approaches,

compared to older technologies, in providing fine-scale information for freshwater mussel

conservation. Although RAD-seq is now widely used for many conservation genetics studies of

non-model organisms, its use for freshwater mussel research is still rare.

In terms of Louisiana Pearlshell recovery, our recommendation is to continue propagation

efforts utilizing wild-fertilized females with a focus on habitat restoration and continued life

history research. Given the likelihood of multiple parentage, captive cohorts produced from a

single female will have more diversity than might have been previously expected. However,

efforts should be made to not re-use the same females over multiple propagation years, and the

use of multiple broodstock females per year is encouraged when possible. Broodstock selected

from populations north of the Red River should not be used to propagate individuals that are to

be released into localities south of the river. While our findings provide some hope for the

efficacy of propagating the Louisiana Pearlshell using existing populations as sources of diverse

brood stock, it also indicates that high levels of inbreeding and loss of population connectivity

may be a looming problem for long-term survival of the species. Indeed given the genetic

bottlenecks at multiple sites sampled in this study, further analyses might reveal hererogeneity of

a reintroduction effort could be improved by mixing propagules from multiple adjacent

populations, taking care to keep individual efforts within their respecitive subdrainages. More

work is needed to ensure that reintroduced and existing populations of this threatened species

form a connected, contemporarily recruiting network of individuals capable of sustaining itself if

true recovery is to be achieved.

Acknowledgements

We thank Jared Streeter (Louisiana Department of Wildlife and Fisheries), Ted Soileau, and

Steve Shively (US Forest Service) for assistance in the field. Nathan Johnson (US Geological

Survey) and Jürgen Geist (Technische Universität München) provided advice on swabbing and

extraction methods. A special thanks to the staff of the Alabama Aquatic Biodiversity Center for

maintaining captively reared *M. hembeli*, developing propagation techniques for the species, and

allowing access to tissues. We also thank the landowners who provided access to sampling sites

on privately owned land. The findings and conclusions in this article are those of the authors and

do not necessarily represent the views of the U.S. Fish and Wildlife Service.

Funding This work was funded by a 1311 system-wide funding grant from U.S. Fish and
Wildlife Service.
591

Availability of data and material

Demultiplexed Illumina sequence data have been uploaded to NCBI SRA (ascession numbers to

be provided upon manuscript acceptance). Processed datasets in various file formats and certain

program input and output files (e.g., colony) are available on FigShare (private link for

reviewers: https://figshare.com/s/2707ff9ee3050d6b10f6; *DOI to be provided upon acceptance*).

Additional code related to the execution of pipelines used are available online at

https://github.com/nlgarrison/ConservationGenomics.

References

Bai Z, Luo M, Zhu W, et al (2012) Multiple paternity in the freshwater pearl mussel Hyriopsis
cumingii (Lea, 1852). J Molluscan Stud 78:142–146.
https://doi.org/10.1093/mollus/eyr049

Baird NA, Etter PD, Atwood TS, et al (2008) Rapid SNP Discovery and Genetic Mapping Using
Sequenced RAD Markers. PLoS ONE 3:e3376.
https://doi.org/10.1371/journal.pone.0003376

Barnhart MC (2006) Buckets of muckets: A compact system for rearing juvenile freshwater
mussels. Aquaculture 254:227–233. https://doi.org/10.1016/j.aquaculture.2005.08.028

Beerli P, Palczewski M (2010) Unified Framework to Evaluate Panmixia and Migration
Direction Among Multiple Sampling Locations. Genetics 185:313–326.
https://doi.org/10.1534/genetics.109.112532

Bogan AE (1993) Freshwater Bivalve Extinctions (Mollusca: Unionoida): A Search for Causes.
Am Zool 33:599–609. https://doi.org/10.1093/icb/33.6.599

Bogan AE (2008) Global diversity of freshwater mussels (Mollusca, Bivalvia) in freshwater. In:
Balian EV, Lévêque C, Segers H, Martens K (eds) Freshwater Animal Diversity
Assessment. Springer Netherlands, Dordrecht, pp 139–147

Christian AD, Monroe EM, Asher AM, et al (2007) Methods of DNA extraction and PCR
amplification for individual freshwater mussel (Bivalvia: Unionidae) glochidia, with the
first report of multiple paternity in these organisms: TECHNICAL ARTICLE. Mol Ecol
Notes 7:570–573. https://doi.org/10.1111/j.1471-8286.2007.01743.x

Cumberlandian Region Mollusk Restoration Committee (2010) Plan for the Population
Restoration and Conservation of Imperiled Freshwater Mollusks of the Cumberlandian
Region

Curole JP, Foltz DW, Brown KM (2004) Extensive allozyme monomorphism in a threatened
species of freshwater mussel, Margaritifera hembeli Conrad (Bivalvia: Margaritiferidae).
Conserv Genet 5:271–278. https://doi.org/10.1023/B:COGE.0000030005.14751.76

Danecek P, Auton A, Abecasis G, et al (2011) The variant call format and VCFtools.
Bioinformatics 27:2156–2158

Davey JW, Blaxter ML (2010) RADSeq: next-generation population genetics. Brief Funct
Genomics 9:416–423. https://doi.org/10.1093/bfgp/elq031

DeSalle R, Amato G (2004) The expansion of conservation genetics. Nat Rev Genet 5:702.
https://doi.org/10.1038/nrg1425

Do C, Waples RS, Peel D, et al (2014) NeEstimator v2: re-implementation of software for the
estimation of contemporary effective population size (Ne) from genetic data. Mol Ecol
Resour 14:209–214

Dray S, Dufour A-B, others (2007) The ade4 package: implementing the duality diagram for
ecologists. J Stat Softw 22:1–20

Excoffier L, Smouse PE, Quattro JM (1992) Analysis of molecular variance inferred from metric
distances among DNA haplotypes: application to human mitochondrial DNA restriction
data. Genetics 131:479–491

Ferguson CD, Blum MJ, Raymer ML, et al (2013) Population structure, multiple paternity, and
long-distance transport of spermatozoa in the freshwater mussel Lampsilis cardium
(Bivalvia:Unionidae). Freshw Sci 32:267–282. https://doi.org/10.1899/12-028.1

Frichot E, François O (2015) LEA: an R package for landscape and ecological association
studies. Methods Ecol Evol 6:925–929

Frichot E, Mathieu F, Trouillon T, et al (2014) Fast and Efficient Estimation of Individual
Ancestry Coefficients. Genetics 196:973–983.
https://doi.org/10.1534/genetics.113.160572

Fritts AK, Fritts MW, Haag WR, et al (2017) Freshwater mussel shells (Unionidae) chronicle
changes in a North American river over the past 1000 years. Sci Total Environ 575:199–
206. https://doi.org/10.1016/j.scitotenv.2016.09.225

Gatenby CM, Neves RJ, Parker BC (1996) Influence of Sediment and Algal Food on Cultured
Juvenile Freshwater Mussels. J North Am Benthol Soc 15:597–609.
https://doi.org/10.2307/1467810

Geist J, Auerswald K (2007) Physicochemical stream bed characteristics and recruitment of the
freshwater pearl mussel (Margaritifera margaritifera). Freshw Biol 52:2299–2316

Goslee SC, Urban DL (2007) The ecodist package for dissimilarity-based analysis of ecological
data. J Stat Softw 22:1–19

Gosselin T, Lamothe M, Grewe P (2020) radiator: RADseq Data Exploration, Manipulation and
Visualization using R

Haag WR (2012) North American Freshwater Mussels: Natural History, Ecology, and
Conservation. Cambridge University Press

Haag WR, Williams JD (2014) Biodiversity on the brink: an assessment of conservation
strategies for North American freshwater mussels. Hydrobiologia 735:45–60.
https://doi.org/10.1007/s10750-013-1524-7

Hart MA, Haag WR, Bringolf R, Stoeckel JA (2018) Novel technique to identify large river host
fish for freshwater mussel propagation and conservation. Aquac Rep 9:10–17.
https://doi.org/10.1016/j.aqrep.2017.11.002

Hendricks SA, Schweizer RM, Wayne RK (2018) Conservation genomics illuminates the
adaptive uniqueness of North American gray wolves. Conserv Genet.
https://doi.org/10.1007/s10592-018-1118-z

Hill A (1986) The Glochidia (larvae) of the Freshwater Mussel Margaritifera hembeli
(Unionacea: Margaritiferidae) Fish Host Species, Morphology, and Period of Fish Host
Infection

Hill WG (1981) Estimation of effective population size from data on linkage disequilibrium.
Genet Res 38:209–216. https://doi.org/10.1017/S0016672300020553

Johnson PD, Brown KM (2000) The importance of microhabitat factors and habitat stability to
the threatened Louisiana pearl shell, Margaritifera hembeli (Conrad). 78:7

Johnson PD, Brown KM (1998) Intraspecific life history variation in the threatened Louisiana
pearlshell mussel, Margaritifera hembeli. Freshw Biol 40:317–329

Jombart T, Collins C (2015) A tutorial for discriminant analysis of principal components
(DAPC) using adegenet 2.0. 0. Imp Coll Lond-MRC Cent Outbreak Anal Model 43:

Jombart T, Kamvar ZN, Collins C, et al (2018) Package 'adegenet'

Jones J, Mair R, Neves R (2005) Factors Affecting Survival and Growth of Juvenile Freshwater
Mussels Cultured in Recirculating Aquaculture Systems. North Am J Aquac 67:210–220.
https://doi.org/10.1577/A04-055.1

Jones JW, Hallerman EM, Neves RJ (2006) Genetic management guidelines for captive
propagation of freshwater mussels (Unionoidea). J Shellfish Res 25:527–535.
https://doi.org/10.2983/0730-8000(2006)25[527:GMGFCP]2.0.CO;2

Jones OR, Wang J (2010) COLONY: a program for parentage and sibship inference from
multilocus genotype data. Mol Ecol Resour 10:551–555

Kamvar ZN, Tabima JF, Grünwald NJ (2014) Poppr: an R package for genetic analysis of
populations with clonal, partially clonal, and/or sexual reproduction. PeerJ 2:e281.
https://doi.org/10.7717/peerj.281

Karlsson S, Moen T, Lien S, et al (2011) Generic genetic differences between farmed and wild
Atlantic salmon identified from a 7K SNP-chip. Mol Ecol Resour 11:247–253.
https://doi.org/10.1111/j.1755-0998.2010.02959.x

Keenan K (2012) divMigrate-online: Visualise and test gene flow patterns among populations.
https://popgen.shinyapps.io/divMigrate-online/

Keenan K, McGinnity P, Cross TF, et al (2016) diveRsity: An R package for the estimation and
exploration of population genetics parameters and their associated errors. Methods Ecol
Evol 782–788. https://doi.org/10.1111/2041-210X.12067@10.1111/(ISSN)2041-
210X.ecologyandevolutionireland

Keller AE, Ruessler DS (1997) Determination or Verification of Host Fish for Nine Species of
Unionid Mussels. Am Midl Nat 138:402–407. https://doi.org/10.2307/2426831

Legendre P, Fortin M-J, Borcard D (2015) Should the Mantel test be used in spatial analysis?
Methods Ecol Evol 6:1239–1247

Lopes-Lima M, Bolotov IN, Do VT, et al (2018a) Expansion and systematics redefinition of the
most threatened freshwater mussel family, the Margaritiferidae. Mol Phylogenet Evol
127:98–118. https://doi.org/10.1016/j.ympev.2018.04.041

Lopes-Lima M, Burlakova LE, Karatayev AY, et al (2018b) Conservation of freshwater bivalves
at the global scale: diversity, threats and research needs. Hydrobiologia 810:1–14.
https://doi.org/10.1007/s10750-017-3486-7

Luikart G, England PR, Tallmon D, et al (2003) The power and promise of population genomics:
from genotyping to genome typing. Nat Rev Genet 4:981.
https://doi.org/10.1038/nrg1226

Malinsky M, Trucchi E, Lawson DJ, Falush D (2018) RADpainter and fineRADstructure:
population inference from RADseq data. Mol Biol Evol 35:1284–1290

Manuzzi A, Zane L, Muñoz-Merida A, et al (2019) Population genomics and phylogeography of
a benthic coastal shark (Scyliorhinus canicula) using 2b-RAD single nucleotide
polymorphisms. Biol J Linn Soc 126:289–303. https://doi.org/10.1093/biolinnean/bly185

Marandel F, Lorance P, Berthelé O, et al (2019) Estimating effective population size of large
marine populations, is it feasible? Fish Fish 20:189–198.
https://doi.org/10.1111/faf.12338

McMurray SE, Roe KJ (2017) Perspectives on the Controlled Propagation, Augmentation, and
Reintroduction of Freshwater Mussels (Mollusca: Bivalvia: Unionoida). Freshw Mollusk
Biol Conserv 20:1. https://doi.org/10.31931/fmbc.v20i1.2017.1-12

Meffe GK, Vrijenhoek RC (1988) Conservation genetics in the management of desert fishes.
Conserv Biol 2:157–169

Meirmans PG (2015) Seven common mistakes in population genetics and how to avoid them.
Mol Ecol 24:3223–3231

Mobile River Basin Mollusk Restoration Committee (2010) A Plan for the Population
Restoration and Conservation of Freshwater Mollusks of the Mobile River Basin

Moses RD, Mathews L, Richard S (2016) Iatt Lake; Lake History and Management Issues

Nei M (1973) Analysis of Gene Diversity in Subdivided Populations. Proc Natl Acad Sci
70:3321–3323. https://doi.org/10.1073/pnas.70.12.3321

Nomura T (2008) Estimation of effective number of breeders from molecular coancestry of
single cohort sample. Evol Appl 1:462–474

Pfister CA, McCoy SJ, Wootton JT, et al (2011) Rapid Environmental Change over the Past
Decade Revealed by Isotopic Analysis of the California Mussel in the Northeast Pacific.
PLOS ONE 6:e25766. https://doi.org/10.1371/journal.pone.0025766

Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus
genotype data. Genetics 155:945–959

QGIS Development Team (2014) QGIS Geographic Information System. Open Source
Geospatial Foundation

Richardson JL, Brady SP, Wang IJ, Spear SF (2016) Navigating the pitfalls and promise of
landscape genetics. Mol Ecol 25:849–863. https://doi.org/10.1111/mec.13527

Rochette NC, Rivera-Colón AG, Catchen JM (2019) Stacks 2: Analytical methods for paired-end
sequencing improve RADseq-based population genomics. Mol Ecol 28:4737–4754.
https://doi.org/10.1111/mec.15253

Roe K (2009) Conservation Genetics of the Freshwater Mussel Margaritifera hembeli (Bivalvia:
Margaritiferidae). Final report to USFWS Natchitoches National Fish Hatchery.
Natchitoches, LA.

Schwartz M, Luikart G, Waples R (2007) Genetic monitoring as a promising tool for
conservation and management. Trends Ecol Evol 22:25–33.
https://doi.org/10.1016/j.tree.2006.08.009

Segelbacher G, Cushman SA, Epperson BK, et al (2010) Applications of landscape genetics in
conservation biology: concepts and challenges. Conserv Genet 11:375–385.
https://doi.org/10.1007/s10592-009-0044-5

Smith DG (1988) Notes on the biology and morphology of Margaritifera hembeli(Conrad,
1838)(Unionacea: Margaritiferidae). Nautilus 102:159–163

Steingraeber MT, Bartsch MR, Kalas JE, Newton TJ (2007) Thermal Criteria for Early Life
Stage Development of the Winged Mapleleaf Mussel (Quadrula Fragosa). Am Midl Nat
157:297–311. https://doi.org/10.1674/0003-0031(2007)157[297:TCFELS]2.0.CO;2

Stewart JH (1990) Louisiana Pearlshell (Margaritifera hembeli) Recovery Plan

Strayer DL, Downing JA, Haag WR, et al (2004) Changing Perspectives on Pearly Mussels,
North America's Most Imperiled Animals. BioScience 54:429.
https://doi.org/10.1641/0006-3568(2004)054[0429:CPOPMN]2.0.CO;2

Strayer DL, Geist J, Haag WR, et al (2019) Essay: Making the most of recent advances in
freshwater mussel propagation and restoration. Conserv Sci Pract 1:e53

Sundqvist L, Keenan K, Zackrisson M, et al (2016) Directional genetic differentiation and
relative migration. Ecol Evol 6:3461–3475. https://doi.org/10.1002/ece3.2096

Team RC (2019) R: a language and environment for statistical computing, version 3.0. 2.
Vienna, Austria: R Foundation for Statistical Computing; 2013

U.S. Fish and Wildlife Service (USFWS) (1988) Endangered and threatened wildlife and plants;
final endangered status for the Louisiana pearlshell *Margaritifera hembeli*. Federal
Register 53(24): 3567-3569.
778
779 U.S. Fish and Wildlife Service (USFWS) (1993) Endangered and Threatened Wildlife and
780   Plants: Determination to Reclassify the Louisiana Pearlshell (*Margaritifera hembeli*)
781   from Endangered to Threatened. Federal Register 58(184): 49935-49937.
782
Vaughn CC, Hakenkamp CC (2001) The functional role of burrowing bivalves in freshwater
ecosystems. Freshw Biol 46:1431–1446. https://doi.org/10.1046/j.1365-
2427.2001.00771.x

Wächtler K, Dreher-Mansur MC, Richter T (2001) Larval Types and Early Postlarval Biology in
Naiads (Unionoida). In: Bauer G, Wächtler K (eds) Ecology and Evolution of the
Freshwater Mussels Unionoida. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 93–
125

Wacker S, Larsen BM, Jakobsen P, Karlsson S (2018) High levels of multiple paternity in a
spermcast mating freshwater mussel. Ecol Evol 8:8126–8134.
https://doi.org/10.1002/ece3.4201

Wacker S, Larsen BM, Jakobsen P, Karlsson S (2019) Multiple paternity promotes genetic
diversity in captive breeding of a freshwater mussel. Glob Ecol Conserv 17:e00564.
https://doi.org/10.1016/j.gecco.2019.e00564

Wang J, Santiago E, Caballero A (2016) Prediction and estimation of effective population size.
Heredity 117:193–206. https://doi.org/10.1038/hdy.2016.43

Waples RK, Larson WA, Waples RS (2016) Estimating contemporary effective population size
in non-model species using linkage disequilibrium across thousands of loci. Heredity
117:233–240. https://doi.org/10.1038/hdy.2016.60

Waples RS (2006) A bias correction for estimates of effective population size based on linkage
disequilibrium at unlinked gene loci*. Conserv Genet 7:167.
https://doi.org/10.1007/s10592-005-9100-y

Waples RS, Antao T, Luikart G (2014) Effects of Overlapping Generations on Linkage
Disequilibrium Estimates of Effective Population Size. Genetics 197:769–780.
https://doi.org/10.1534/genetics.114.164822

Watters GT (1996) Small dams as barriers to freshwater mussels (Bivalvia, Unionoida) and their
hosts. Biol Conserv 75:79–85. https://doi.org/10.1016/0006-3207(95)00034-8

Whelan NV, Galaska MP, Sipley BN, et al (2019) Riverscape genetic variation, migration
patterns, and morphological variation of the threatened Round Rocksnail, Leptoxis
ampla. Mol Ecol 28:1593–1610

Williams JD, Warren ML, Cummings KS, et al (1993) Conservation Status of Freshwater
Mussels of the United States and Canada. 18:17

Witzenberger KA, Hochkirch A (2011) Ex situ conservation genetics: a review of molecular
studies on the genetic consequences of captive breeding programmes for endangered
animal species. Biodivers Conserv 20:1843–1861. https://doi.org/10.1007/s10531-011-
0074-4

Woodings LN, Murphy NP, Doyle SR, et al (2018) Outlier SNPs detect weak regional structure
against a background of genetic homogeneity in the Eastern Rock Lobster, Sagmariasus
verreauxi. Mar Biol 165:185. https://doi.org/10.1007/s00227-018-3443-7

844
845
846

Table 1 Pairwise FST (top) and estimated in-stream distance in km (bottom) for all populations analyzed.

	Jordan	Black	Loving	Brown	Captive
Jordan	-	0.043	0.063	0.078	0.048
Black	36.5	-	0.064	0.082	0.035
Loving	77.1	91.4	-	0.063	0.071
Brown	87.8	102.5	41.2	-	0.086
Captive	36.5	0	91.4	102.5	-

888
889

Table 2 Population summary statistics (in part, full summary available online) from left to right; number of individuals (N), number of loci recovered (Loci), number of private alleles (PA), nucleotide diversity (Π) and standard deviation (SD), observed heterozygosity (H$_o$), expected heterozygosity (H$_e$), coefficient of inbreeding (F$_{IS}$), effective populaiton size (N$_e$) and 95% confidence interval (CI), number of effective breeders (N$_b$), and allelic richness (AR).

Population	N	Loci	PA	Π (SD)	H$_o$ (SD)	H$_e$ (SD)	F$_{IS}$ (SD)	N$_e$ (CI)	N$_b$	AR (CI)
Jordan	18	2134	91	0.25 (±0.04)	0.10 (±0.02)	0.24 (±0.03)	0.45 (±0.17)	INF	4.2	1.47 (1.41-1.51)
Black	20	2203	88	0.25 (±0.04)	0.10 (±0.02)	0.24 (±0.04)	0.43 (±0.16)	205 (188-225)	3.9	1.51 (1.46-1.55)
North	38	--	--	--	--	--	--	177 (170-183)	3.9	--
Loving	16	1672	40	0.27 (±0.04)	0.09 (±0.02)	0.25 (±0.04)	0.46 (±0.17)	INF	4.3	1.16 (1.12-1.19)
Brown	17	1941	142	0.27 (±0.04)	0.09 (±0.01)	0.26 (±0.03)	0.49 (±0.17)	INF	3.9	1.35 (1.30-1.38)
South	33	--	--	--	--	--	--	488 (429-566)	4.1	--
Captive	19	2024	41	0.25 (±0.04)	0.10 (±0.02)	0.24 (±0.04)	0.40 (±0.16)	70 (64-78)	5.0	1.37 (1.33-1.41)

Table 3 Description of models assessed by migrate-n, sorted by model rank; model numbers correspond to those in Fig. 2.

Number	Model Description	Log marginal likelihood value	Rank
3	**Northern panmixia, bidirectional South**	**-93595.56**	**1**
4	Northern panmixia, full migration	-93803.10	2
1	Full migration	-94306.55	3
5	Panmixia	-94030.00	4
2	Northern panmixia, unidirectional South	-94612.37	5

Fig. 1 Map depicting the extent of M. hembeli distribution in central Louisiana. Pictures illustrate mussel bed density at Black Creek site (upper) and individual shell (lower).

Fig. 2 Population migration models assessed using migrate-N.

Fig. 3 Discriminant analysis of principal components output showing clustering of a) all wild populations and b) wild + captive individuals using the multiple SNP per locus datasets.

Fig. 4 Individual admixtures for all wild individuals as inferred by LEA analysis, using the best-fit K of 2. Colors correspond to parish designations depicted in Fig. 1.

Fig. 5 Hierarchical heatmap generated by fineRADstructure, all individuals included. Letters correspond to population identifiers (A = Jordan Creek, B = Black Creek/Captive cohort, C = Brown Creek, D = Loving Creek). Boxes highlight patterns of population clustering, colors represent relative co-ancestry values averaged by population.

Fig. 6 Hierarchical heatmap generated by fineRADstructure, only captive individuals. Boxes highlight patterns of individual clustering, colors represent relative co-ancestry values averaged by cluster.

Fig. 7 Network generated by divMigrateOnline using the WithCaptiveS dataset; GST statistic, 0.40 filtering level, alpha 0.05. Colors correspond to those in Fig. 3.