The Na$^+$/K$^+$ ATPase Regulates Glycolysis and Modifies Immune Metabolism in Tumors

Sydney M. Sanderson[1#], Zhengtao Xiao[1#], Amy J. Wisdom[1, 2], Shree Bose[1], Maria V. Liberti[1], Michael A. Reid[1],

Emily Hocke[3], Simon G. Gregory[3], David G. Kirsch[2], Jason W. Locasale[1*]

[1]Department of Pharmacology and Cancer Biology, Duke University School of Medicine, Durham, NC, 27710, USA

[2]Department of Radiation Oncology, Duke University School of Medicine, Durham, NC, 27710, USA

[3]Duke Molecular Physiology Institute, Duke University School of Medicine, Durham, NC, 27710, USA

[#]These authors contributed equally to this work

[*]**Corresponding author: dr.jason.locasale@gmail.com**

Abstract

Cancer therapies targeting metabolism have been limited due to a lack of understanding of the controlling properties of vulnerable pathways. The Na$^+$/K$^+$ ATPase is responsible for a large portion of cellular energy demands but how these demands influence metabolism and create metabolic liabilities are not known. Using metabolomic approaches, we first show that digoxin, a cardiac glycoside widely used in humans, acts through disruption to central carbon metabolism via on target inhibition of the Na$^+$/K$^+$ ATPase that was fully recovered by expression of an allele resistant to digoxin. We further show in vivo that administration of digoxin inhibits glycolysis in both malignant and healthy cells, particularly within clinically relevant cardiac tissue, while exhibiting tumor-specific cytotoxic activity in an allografted soft tissue sarcoma. Single-cell expression analysis of over 31,000 cells within the sarcoma shows that acute Na$^+$/K$^+$ ATPase inhibition shifts the immune composition of the tumor microenvironment, leading to selective alterations to metabolic programs in specific immune cells thus acting both through tumor cell and microenvironmental (e.g. macrophage) cells. These results provide evidence that altering energy demands can be used to regulate glycolysis with cell-type specific consequences in a multicellular environment of biomedical interest.

Introduction

Cancer cells exhibit metabolic reprogramming to support their uncontrolled proliferation, most notably in their utilization of processes that ultimately generate energy including glycolysis and the tricarboxylic acid (TCA) cycle (collectedly referred to as central carbon metabolism). Energy metabolism is coupled to redox, biosynthesis, and signaling[1,2]. Altered metabolism such as that observed in cancer has been attributed to a myriad of factors, such as oncogene activation[3-5], dysregulation of mitochondria[6-9], adaptation to oxygen[10,11] and nutrient[12] scarcity. While many of these alterations have been shown to promote dependencies on central carbon metabolism, therapies targeting these metabolic processes are toxic, and the efficacy of those that are sufficiently tolerable have poorly understood specificities[13,14]. Therefore, novel mechanisms that reveal vulnerabilities associated with metabolic reprogramming remain highly desired.

Cardiac glycosides (CGs) are tolerated agents commonly prescribed for the treatment of cardiac arrythmias or congestive heart failure. They are largely believed to act as inhibitors of the sodium-potassium pump (also referred to as the Na^+/K^+ ATPase)[15]. This transmembrane enzyme imports two potassium ions while exporting three sodium ions in an ATP-dependent manner, thereby maintaining the electrochemical gradient across the cell membrane[16]. The activity of this pump additionally contributes to the regulation of intracellular pH[17], glucose uptake[18], and Ca^{2+} levels[19].

Interestingly, CGs have been shown to exhibit anticancer activity[20-22]. Nearly half a century ago, Efraim Racker postulated that a partially defective Na^+/K^+ ATPase could be a primary tumorigenic event by disrupting the cellular energetic state[23]. Given that the Na^+/K^+ ATPase accounts for roughly 20-70% of cellular ATP demand[24,25], it was suggested that lowered enzyme activity could induce dysregulation of ATP-producing processes and resulting cellular

programming towards an oncogenic state[23]. While this hypothesis has remained largely

unexplored, it suggests that the Na^+/K^+ ATPase could be a major mechanism of control over

central carbon metabolism.

In this study, we show that an immediate and direct consequence of Na^+/K^+ ATPase

inhibition is disruption of central carbon metabolism and then show how this reprograms nearly

all of metabolism. We demonstrate that the downstream metabolic consequences of digoxin

treatment are specifically mediated through on-target inhibition of the Na^+/K^+ ATPase (that

could be fully resotred by a resistant allele of Na^+/K^+ ATPase) and that digoxin treatment can

impact metabolic processes in both healthy and malignant tissues with differential requirements

in malignant cells. Furthermore, we use single-cell RNA sequencing to explore the metabolic

consequences of Na^+/K^+ ATPase inhibition to show that acute inhibition reveals specific

metabolic requirements in the tumor microenvironment and exerts intriguing metabolic changes

in different immune cell compartments leading to coordinated regulation of non-metabolic genes.

Results

Digoxin disrupts central carbon metabolism and related processes in a time- and dose-
dependent manner

To determine whether cell sensitivity to digoxin is associated with intrinsic metabolic state, we

compared basal metabolic uptake and excretion rates with digoxin IC_{50} values across a panel of

59 cancer cell lines. This analysis demonstrated that digoxin treatment correlated with the

metabolic flux of TCA intermediates (Figure 1A), including malate (Spearman correlation, $r = -$

0.38, $p = 0.0025$) and citrate (Spearman correlation, $r = -0.35$, $p = 0.0056$), as well as the ATP-

recycling metabolite creatine (Spearman correlation, $r = 0.29$, $p = 0.02$) (Figure S1A). This

initial finding suggested that Na^+/K^+ ATPase activity could be linked to central carbon metabolic

processes.

78   To assess the metabolic consequences of digoxin treatment, we generated metabolite

profiles of HCT-116 cells both temporally using the digoxin IC_{50} concentration of 100 nM

(Figure 1B), as well as acutely (3 hours) using increasing concentrations of digoxin (Figure S1B

and 1C). Pathway analyses demonstrated that central carbon metabolism, as well as processes

associated with its activity including aspartate/glutamate metabolism and taurine metabolism,

were among the most significantly impacted metabolic pathways by digoxin in both a time- and

dose-dependent manner (Figure 1D).

85   Closer examination of these metabolite profiles revealed an increase in upper glycolytic

intermediates as well as a decrease in TCA cycle metabolites in both a time- (Figure 1E and

S1C) and dose-dependent (Figure 1F and S1C) manner. Importantly, the accumulation of

fructose 1,6-bisphosphate (F1,6BP) has been shown to indicate changes to glycolytic flux [26],

indicating that the observed increases in F1,6BP levels (Figure 1E and 1F) are demonstrative of

disruption to central carbon metabolism. Additionally, we consistently observed decreases in the

levels of taurine (Figure 1F) and hypotaurine (Figure S1D), as well as creatine (Figure 1F) and

phosphocreatine (Figure S1E). Taurine has been suggested to partially regulate mitochondrial

electron transport chain (ETC) activity [27], while creatine enables the rapid anaerobic recycling of

ATP under high energetic demand [28], thus providing additional evidence of disrupted energy

status.

96   To study the effects of digoxin on downstream processes of central carbon metabolism,

we first measured steady-state glucose incorporation into glycolysis and the downstream TCA

cycle using media supplemented with uniformly labeled [^{13}C]-glucose ([U-^{13}C]-glucose) (Figure

1G and S1F). We found reductions in the labeling of both glycolytic and TCA intermediates

(Figure 1H and S1G). Additional assessment of the kinetics of glycolytic flux into the

mitochondria further revealed that the labeling of TCA intermediates [U-^{13}C]-glucose

administration was significantly reduced (Figure 1I and S1H). Finally, by measuring the steady-

state incorporation of glutamine into the TCA cycle via conversion to α-ketoglutarate (α-KG)

using [U-^{13}C]-glutamine (Figure 1J), we found significantly reduced labeling of TCA

intermediates (Figure 1J and S1I) further indicating alterations in mitochondrial activity.

Altogether, these findings demonstrate that digoxin substantially disrupts multiple nodes of

central carbon metabolism which extends to other intracellular energetic processes.

Digoxin exerts its metabolic effects via on-target inhibition of the Na$^+$/K$^+$ ATPase

Upon consideration of the metabolic reprogramming induced by digoxin treatment, we assessed

whether the cytotoxic effects of digoxin could be mitigated by nutrient supply. We cultured

HCT-116 cells in media containing 100 nM digoxin as well as supplementations of nutrients

related to central carbon metabolism or redox balance which is coupled to central carbon

metabolism; surprisingly, we found that the majority of these treatments were insufficient to

rescue cells from digoxin treatment (Figure 2A), although the supplementation of either

nucleosides or the antioxidant NAC were modestly cytoprotective (Figure 2B), in line with

previous reports [29]. These results suggest that the metabolic disruption induced by digoxin likely

exerts extensive downstream consequences on cellular homeostasis and function beyond the

targeting of a single pathway.

The possibility remained that the metabolic consequences of digoxin were due to factors

other than inhibition of Na$^+$/K$^+$ ATPase activity. It is established that murine Na$^+$/K$^+$ ATPase

enzymes are substantially less sensitive to CGs, with the mouse isoform exhibiting roughly

1000-fold lower affinity for CGs than the human counterpart [30,31]. Indeed, it has been shown that

ectopic expression of the α-subunit of the mouse ATPase (mATP1a1) is sufficient to rescue cell

viability upon digitoxin treatment [32]. Therefore, to determine whether the observed metabolic

effects were specifically attributable to disruption of Na^+/K^+ ATPase activity, we ectopically

expressed the mouse mATP1a1 subunit in HCT-116 cells and demonstrated a complete rescue of

cell viability after treatment with digoxin (Figure 2C). We then treated the mATP1a1-expressing

cells with digoxin at the IC_{50} concentration and compared the resulting metabolite profiles

(Figure 2D). We found that mATP1a1 expression completely blocked the metabolic

consequences of digoxin treatment (Figure 2E and 2F), thereby restoring the activity of central

carbon metabolism (Figure 2G and 2H) as well as taurine (Figure 2I) and creatine (Figure 2J)

levels. This finding establishes that disrupted central carbon metabolism is an intrinsic

component of digoxin-induced cytotoxicity, which is a direct result of Na^+/K^+ ATPase inhibition.

Digoxin treatment impacts energy metabolism in a tissue-specific and antineoplastic
manner

After defining the impact of digoxin on central carbon metabolism and related energetic

processes in cell culture, we next sought to determine whether these effects could be achieved in

a more complex *in vivo* setting. Numerous findings of CGs effectively inhibiting tumor growth

in xenograft studies have been reported [29,32,33]; however, given the extreme (greater than 1000-

fold) differential in CG-binding to the ATPase enzymes found in human-derived cells and the

murine host, these findings are confounded by the artificially high concentrations in xenograft

studies of digoxin that can be used with no effects on the mouse host due to their weak binding

affinity to murine Na^+/K^+ ATPases. Thus, a therapeutic window needed to achieve tumor

growth inhibition could be prohibitive in a setting where the host and tumor express the same

ATPase enzymes [34]. Additionally, it remains to be explored whether Na^+/K^+ ATPase inhibition

can impact metabolism in both healthy and malignant tissue.

To investigate these essential considerations, we cultured mouse sarcoma cells generated

from primary sarcoma tumors driven by expression of oncogenic $KRAS^{G12D}$ and p53 deletion

($Kras^{LSL-G12D/+};Trp53^{flox/flox}$, or KP) and found that they exhibited an IC_{50} of 100 μM, in line with

their murine isoform expression (Figure S2A). As proof of principle, we first verified that

administration of this concentration reliably impacted central carbon metabolism in cell culture

(Figure S2B and S2C). We then orthotopically injected these cells into the right gastrocnemius

muscle of syngeneic 129/SvJae mice. Upon tumor palpation (approximately 11 days after

injection), we treated mice with a previously reported dose[35] of 2 mg/kg digoxin every 24 hours

and collected tumors as well as healthy tissues after administration of the fourth dose (Figure

3A). Of note, while this treatment regimen appeared to trend towards tumor growth inhibition,

treated mice exhibited some weight loss upon daily digoxin treatment (Figure S2D and S2E)

although no other physiological or behavioral signs of toxicity were observed.

Metabolite profiling revealed that cardiac tissue was the most significantly impacted

tissue type (Figure 3B), with substantial alterations in glycolytic (Figure S2F) and TCA cycle

(Figure S2G) intermediates. These effects were even more pronounced in cardiac than tumor

tissue in this context (Figure 3B and 3C). A number of similar changes were also found in

muscle, brain, liver, and kidney tissue (Figure S2H), although to a much lesser extent than was

found in cardiac tissue (Figure S2F, S2G, S2I and S2J) or the cultured sarcoma cells (Figure S2B

and S2C).

To examine the antineoplastic potential of digoxin in this setting, we treated the second

cohort of orthotopically-engrafted syngeneic mice with digoxin (2mg/kg) or vehicle every 48

hours and monitored tumor growth (Figure 3D). There were no discernible histological

differences between vehicle- and digoxin-treated groups, with the majority of tumors exhibiting

regions of both low (Figure 3E, left) and high (Figure 3E, right) vascularity. Of note, this

heterogeneous landscape, likely resulting from the relatively large size of these KP allograft

tumors (Figure 3D), closely resembles the regional heterogeneity found in patient sarcomas[36].

Regression analyses demonstrated that digoxin treatment significantly delayed tumor growth as

measured by time to tumor quintupling (Figure 3F). Furthermore, the most prominent metabolic

alterations found in tumors exposed to chronic digoxin treatment were consistent with

dysregulation of energy-related metabolic processes (Figure 3G), with disruptions to

mitochondrial metabolism (Figure 3H) as well as taurine (Figure 3I) and creatine (Figure 3J)

levels. Collectively, these findings demonstrate that digoxin can impact metabolic processes in

both healthy and malignant tissues and that these metabolic perturbations lead to antineoplastic

effects in a physiological setting.

Acute digoxin treatment remodels the immune compartment of the tumor
microenvironment

An important feature of allograft models is the presence of an intact immune system[37,38]. Given

that the differential requirements of central carbon metabolism have become increasingly

appreciated in immune cell function[39-41], we performed single-cell RNA sequencing (scRNA-

seq) to explore the effects of digoxin treatment on the tumor microenvironment. Briefly, we

treated mice with vehicle or 2 mg/kg digoxin (using two mice per group) with the daily treatment

regimen described previously (Figure 4A) and harvested the tumors after the final fourth dose.

We then dissociated the tumors into single-cell suspensions and performed 10x scRNA-seq

without cell type sorting or purification using the Chromium drop-seq platform (10x Genomics)

(Figure 4A). We analyzed mRNA expression from >7,200 cells from each tumor after quality

controls (Figure 4A), with roughly 100,000 reads from each cell that covered nearly 5,000 genes

(Figure S3A—S3C), and used previously published scRNA-seq data of healthy murine muscle

tissue[42] as the reference to distinguish between malignant and non-malignant cells based on

relative gene copy number[43] (Figure 4B, S3D and S3E). Acute digoxin treatment trended

towards a reduction in the relative population of malignant cells (Figure 4C), consistent with

previous observations of early tumor growth inhibition (Figure S2D).

Upon further analysis of the single-cell transcriptomes, we identified fifteen immune cell

populations, including five distinct T-cell populations (Figure 4D, S4A and S4B). Myeloid cells

(i.e. type I and type II macrophages, M1 and M2 respectively) were the most abundant immune

cell population (Figure 4E), consistent with previous reports of autochthonous KP sarcomas [44].

While the relative populations of intratumoral B cells and T cells did not appear to be

appreciably skewed by this short-term digoxin treatment, the relative populations of neutrophil

and dendritic cell infiltrates were slightly increased and decreased, respectively (Figure 4F).

These results demonstrate that acute digoxin treatment induces an immediate shift in the tumor

microenvironment.

Targeting the Na^+/K^+ ATPase with digoxin alters metabolic processes in tumor cells and
immune infiltrates

To determine whether digoxin treatment exerts differential effects on metabolic programming

between cell types, we compared the scRNA-seq transcriptome of each tumor cell population

after digoxin treatment with vehicle treatment. Oxidative phosphorylation was the most

extensively altered metabolic program in malignant cells, with significant increases in expression

of *Atp5k* transcripts corresponding to the ATP synthase gene (Figure 5 A and B). ATP synthase

expression is linked to glycolytic activity, thereby providing additional evidence that central

carbon metabolism disruption is an immediate and direct consequence of digoxin treatment.

Further functional enrichment analysis of all significantly altered genes showed that the

apoptosis process was activated (Figure S5), which could be induced by mitochondrial OXPHOS

system[45,46].

Upon examination of the non-malignant cell populations, we found that different cell

populations were subject to various magnitude alterations on their metabolic genes after digoxin

treatment (Figure 5C). Pathway analysis of the significantly altered metabolic genes (|Cohens' d|

> 0.1 and p-value < 0.05) suggested that digoxin exposes diversity in metabolic plasticity among

different cell populations (Figure 5D and S6). The macrophage populations exhibited metabolic

reprogramming specifically within central carbon metabolism, with M1 macrophages

characterized by upregulation of glycolysis while M2 macrophages were characterized by an

upregulation in oxidative phosphorylation similar to malignant cells (Figure 5D). Interestingly,

previous studies of macrophage behavior have also observed similar metabolic shifts in these

two populations upon their activation[47,48]. To examine this possibility more closely, we

performed KEGG pathway analysis on their full transcriptomes and additionally found

transcriptional signatures (i.e. endoplasmic reticulum associated genes) consistent with their

polarization[49-51] (Figure S7 A-C). These observations illustrate that digoxin is associated with

broad transcriptional consequences, including within metabolic processes, across multiple tumor

cell populations.

Discussion

Metabolic programming, on both a cellular and physiological level, is known to be highly

regulated by environmental factors which are extrinsic to discrete metabolic reactions[52-56]. In line

with this, many conventional therapies have been shown to exert substantial metabolic effects

beyond their understood mechanism of action[57-59]. As many of the metabolic vulnerabilities

inherent to cancer cells are difficult to target without inducing toxic consequences on healthy

tissue, the identification of agents that can be repurposed to exploit these processes remains an

active area of investigation[60-63].

Cardiac glycosides have been shown to exhibit antineoplastic effects in numerous

settings[20,21,35], which have been attributed to a myriad of sources such disruption of proton

gradients[32] and activation of kinases that physically interact with the sodium-potassium pump

(i.e. Na^+/K^+ ATPase "signalosome")[64-66]. However, fluctuations in energetic demand from

membrane transport activity have been shown to impact glycolytic rate[67], and it has been

historically hypothesized that alterations in Na^+/K^+ ATPase activity may be a contributing factor

to enhanced dependence on glycolysis in cancer[23]. Our results demonstrate that digoxin, through

on-target inhibition of the Na^+/K^+ ATPase, induces broad metabolic disruptions in a time- and

dose-dependent manner; these disruptions were most prominent within central carbon and

energy-related metabolic processes, including taurine and creatine metabolism. Interestingly,

taurine has been suggested to act as a mild cardiac glycoside in cardiac tissue through its

modulation of mitochondrial ROS[27], and taurine loss has been associated with Na^+ efflux thereby

reducing the degree of Ca^{2+} overload[68]. It's therefore tempting to speculate that the dual loss of

taurine and creatine upon digoxin treatment are compensatory responses to ion imbalance and

reduced ATP production, respectively.

Our results further provide the first characterization of the global metabolic

consequences of digoxin on healthy tissues to allow direct comparison with metabolic profiles in

tumors. We found that acute digoxin treatment effectively impacted central carbon metabolism

or other energy-related metabolites (i.e. taurine and creatine) in diverse healthy tissues, with the

most observable metabolic consequences found in cardiac tissue. Although the higher degree of

central carbon metabolic disturbance in cardiac compared to tumor tissue after short-term

digoxin treatment was unexpected, the use of digoxin for treating cardiac dysfunction likely

indicates a tissue-specific affinity for CGs; indeed, it has been shown that the human cardiac-

specific alpha subunit isoform exhibits a higher affinity for CGs compared to the isoforms

present in most other tissues[69], thereby providing a potential mechanism for the observed effects.

Importantly, despite these prominent metabolic alterations in cardiac tissue, we did not observe

any signs of cardiotoxicity at the doses we used, and these alterations were considerably more

prominent in tumors with the antineoplastic long-term digoxin treatment, indicating the enhanced

dependency of tumor cells on these processes.

We have recently demonstrated that scRNA metabolic gene transcriptomes can be used

to investigate aspects of metabolic activity and plasticity within individual cells [46]. Our results

additionally contribute to our understanding of unique cell-autonomous responses within the

tumor microenvironment by using scRNA-seq to examine diverse cell populations and metabolic

programs within tumors following acute cytotoxic treatment. Our findings of both altered

representations of immune infiltrate, as well as transcriptional programs of metabolic processes

featuring clear distinctions between malignant and immune cell populations, in response to short-

term digoxin treatment provide a novel glimpse into cell-specific intratumoral heterogeneity.

It remains unclear whether the altered metabolic programs and possible activation of

different myeloid populations are a direct result of exposure to digoxin or indirect response to

signals from adjacent malignant cells; furthermore, the observed variations in cell populations

might become more pronounced with longer exposure to digoxin or at different stages of tumor

development. Indeed, the potential dual polarization of M1 and M2 macrophages (which are

commonly characterized as pro-inflammatory or anti-inflammatory, respectively)[70] is intriguing

and warrants further investigation. It will be interesting in future studies to determine the

functional consequences of these variations in metabolic reprogramming between immune cell

populations, especially upon consideration of previous studies reporting consequences of digoxin

on immune cell activity in various noncancerous settings [71,72]. Additional studies of these

interactions could illuminate possible synergistic mechanisms that may elicit enhanced clinical

efficacy of digoxin when combined with targeted immune therapies as has been found with other

compounds[44,73,74].

Methods

Cell Culture

294 Cells were cultured at 37°C, with 5% atmospheric CO_2 in RPMI-1640 (Gibco), 10% heat-

295 inactivated fetal bovine serum (FBS; F2442, Sigma), 100 U/mL penicillin (Gibco), and 100

296 mg/mL streptomycin (Gibco). HCT-116 cells were obtained from the American Tissue Culture

297 Collection (ATCC), and cultured murine sarcoma cells were generated from primary murine

298 sarcomas described previously [44] and below in "Allograft Mouse Studies".

Digoxin IC$_{50}$ measurements

Cells were plated at a density of 5.0×10^3 cells/well in triplicate in a 96-well plate and were

allowed to adhere in full RPMI 1640 media for 24 hours. Cells were briefly washed with PBS

and then incubated in medium containing vehicle (DMSO; #97061-250, VWR) or indicated

concentrations of digoxin (#D6003, Sigma Aldrich). After 48 hours, the media was aspirated and

replaced with 100 µl phenol-red free RPMI-1640 (Gibco) and 12 mM 3-[4,5-Dimethylthiazol-2-

yl]-2,5-diphenyltetrazolium (MTT) (Thermo Fisher Scientific, #M6494). After 4 hours, the

MTT-containing media was aspirated and 50 µl DMSO was added to dissolve the formazan.

After 5 minutes, absorbance was read at 540 nm. For all experiments, three technical replicates

per culture condition were used.

Nutrient Rescue Experiments

For all nutrient rescue experiments, cells were plated at a density of 5.0×10^5 cells/well in

triplicate in a 6-well plate and were allowed to adhere for 24 hours. Cells were briefly washed

with PBS and then incubated with vehicle (DMSO) or 100 nM digoxin, as well as one or more of

the following supplementations: 5mM N-acetyl-cysteine (NAC; A9165, Sigma), 2 µM Trolox

(238813, Sigma), 100 µM adenosine 5'-triphosphate disodium salt hydrate (ATP; A26209,

Sigma), 10mM creatine (C3630, Sigma), 10mM Taurine (T8691, Sigma), 2µM dimethyl-a-

ketoglutarate (cell-permeable a-KG; 349631, Sigma), 5mM sodium pyruvate (sc-208397A, Santa

Cruz Biotechnology), 500 µM nicotinamide (N0636, Sigma), 100µM beta-nicotinamide adenine

dinucleotide (NAD$^+$; 160047, MP Biomedical), and $1 \times$ Embryomax Nucleosides (ES-008-D,

Millipore). For glucose-restricted medium, RPMI 1640 glucose- and glutamine-free medium was

supplemented with 2 mM glutamine, 10% heat-inactivated FBS, 100 U/mL penicillin, and 100

mg/mL streptomycin; indicated concentrations of glucose were then serially added to the

medium. After 48 hours, an MTT cell viability assay was performed as described above. For all

experiments, three technical replicates per culture condition were used.

[U-^{13}C] glucose and [U-^{13}C] glutamine Tracing

[U-^{13}C] glucose (CLM-1396-10) and [U-^{13}C] glutamine (CLM-1822-H-PK) were purchased

from Cambridge Isotope Laboratories. [U-^{13}C] glucose was added to RPMI 1640 glucose-free

medium at a concentration of 11 mM, while [U-^{13}C] glutamine was added to glucose- and

glutamine-free RPMI-1640 medium (supplemented with 11 mM glucose) to a concentration of 2

mM. Cells were plated at a density of 3.0×10^5 cells/well in a 6-well plate, and after 24 hours

were treated with either vehicle (DMSO) or 100 nM digoxin. After 4 hours of drug treatment, the

medium was quickly aspirated and washed with glucose-free RPMI 1640 medium, and either [U-

^{13}C] glucose- or [U-^{13}C] glutamine-containing medium was added to each well. Cells were then

collected for metabolite extraction either 20 hours later or at the indicated time points. For all

experiments, three technical replicates per culture condition were used.

Lentiviral Transfection and Transduction

HEK-293T cells were plated at a density of 1.0×10^6 cells/10 cm plate in RPMI 1640 (Gibco)

supplemented with 10% heat-inactivated FBS, penicillin (100 U/ml), and streptomycin (100

mg/ml) and were allowed to adhere and reach 70% confluency. 15 μg of mATP1a1 (EX-

Mm01329-Lv105, GeneCopoeia) or GFP control (EX-EGFP-Lv105, GeneCopoeia) plasmid, 10

μg of PsPAX2 packaging vector (no. 12260, Addgene), and 5 μg of PMD2.G envelope–

expressing plasmids (no. 12259, Addgene) were diluted in 500 μl of jetPRIME buffer (no. 114-

07, Polyplus-transfection) and vortexed. Next, 60 μl of the jetPRIME transfection reagent (no.

114-07, Polyplus-transfection) was added to the mixture, vortexed for 10 seconds, and left to

incubate for 10 minutes at room temperature. The medium in the plate was replaced with fresh

medium, and the transfection mix was then added to the 10-cm plate dropwise. After 24 hours,

the transfection medium was replaced with fresh medium. After an additional 24 hours, the

medium was collected and filtered through a 0.45-μm filter for virus collection. HCT-116 cells

were plated in 10-cm plates, and when they reached 30-50% confluency, virus-containing

medium (1:1 with fresh RPMI 1640 medium) was added to the plates along with polybrene (4

μg/μl). After 24 hours, the virus-containing medium was removed and replaced with fresh RPMI

1640 medium. Cells were incubated with puromycin (1 µg/ml) for 48 hours for positive

selection.

Metabolite Extraction

Media was quickly aspirated and 1 mL of extraction solvent (80% methanol/water, cooled to -

80°C) was added to each well of the 6-well plates, and were then transferred to -80°C for 15

minutes. Plates were removed and cells scraped into the extraction solvent on dry ice. All

metabolite extracts were centrifuged at 20,000g at 4°C for 10 min. Finally, the solvent in each

sample was evaporated in a speed vacuum, and the resulting pellets were stored in -80°C until

resuspension. For polar metabolite analysis, the cell extract was dissolved in 15 µL water and 15

µL methanol/acetonitrile (1:1, v/v) (LC-MS optima grade, Thermo Fisher Scientific). Samples

were centrifuged at 20,000g for 2 minutes at 4°C, and the supernatants were transferred to liquid

chromatography (LC) vials.

Liquid Chromatography

Ultimate 3000 HPLC (Dionex) with an Xbridge amide column (100 x 2.1 mm i.d., 3.5 µm;

Waters) was coupled to Q Exactive-Mass spectrometer (QE-MS, Thermo Scientific) for

metabolite separation and detection at room temperature. The mobile phase A reagent was

composed of 20 mM ammonium acetate and 15 mM ammonium hydroxide in 3% acetonitrile in

HPLC-grade water (pH 9.0), while the mobile phase B reagent was acetonitrile. All solvents

were LC-MS grade and were purchased from Fischer Scientific. The flow rate used was 0.15

mL/min from 0-10 minutes and 15-20 minutes, and 0.3 mL/min from 10.5-14.5 minutes. The

linear gradient was as follows: 0 minutes 85% B, 1.5 minutes 85% B, 5.5 minutes 35% B, 10

minutes 35% B, 10.5 minutes 25% B, 14.5 minutes 35% B, 15 minutes 85% B, and 20 minutes

85% B.

Mass Spectrometry

The QE-MS is outfitted with a heated electrospray ionization probe (HESI) with the following

parameters: evaporation temperature, 120°C; sheath gas, 30; auxiliary gas, 10; sweep gas, 3;

spray voltage, 3.6 kV for positive mode and 2.5 kV for negative mode. Capillary temperature

was set at 320°C and S-lens was 55. A full scan range was set at 60 to 900 (m/z), with the

resolution set to 70,000. The maximum injection time (max IT) was 200 ms. Automated gain

control (AGC) was targeted at 3,000,000 ions.

Peak Extraction and Metabolomics Data Analysis

Data collected from LC-Q Exactive MS was processed using commercially available software

Sieve 2.0 (Thermo Scientific). For targeted metabolite analysis, the method "peak alignment and

frame extraction" was applied. An input file ("frame seed") of theoretical m/z (width set at 10

ppm) and retention time of ~260 known metabolites was used for positive mode analysis, while a

separate frame seed file of ~200 metabolites was used for negative mode analysis. To calculate

the fold changes between different experimental groups, integrated peak intensities generated

from the raw data were used. Hierarchical clustering and heatmaps were generated using

Morpheus software (The Broad Institute, https://software.broadinstitute.org/morpheus/). For

hierarchical clustering, Spearman correlation parameters were implemented for row and column

parameters. Pathway enrichment analysis was conducted by MetaboAnalyst 3.0 software

(http://www.metaboanalyst.ca/faces/home.xhtml) using HMDB IDs of the metabolites that were

significantly enriched ($p < 0.05$). The pathway library used was *Homo sapiens* and Fishers'

Exact test was employed for over-representation analysis. Other quantitation and statistics were

calculated using Graphpad Prism software.

Allograft Mouse Studies

All animal studies were performed following protocols approved by the Duke University

Institutional Animal Care and Use Committee (IACUC) and adhere to the NIH Guide for the

Care and Use of Laboratory Animals. KP sarcoma cells were obtained from a primary $Kras^{LSL\text{-}}$

$^{G12D/+}$;$Trp53^{flox/flox}$ sarcoma. The tumor was dissected from the hind limb and dissociated by

shaking for 45 minutes at 37°C in collagenase Type IV (Gibco), dispase (Gibco), and trypsin

(Gibco). Cell suspension was then strained through a 40 µm filter, washed in PBS, and plated for

culture. KP cells were maintained *in vitro* in DMEM (Gibco) containing 10% heat-inactivated

FBS (Gibco), and 1% Penicillin/Streptomycin (Gibco) for 8-10 passages before transplanting

into syngeneic mice. All mice were maintained on a pure 129/SvJae genetic background. For

allograft tumor initiation, cultured KP murine cells were suspended in DMEM medium at a

concentration of 5×10^6 cells/mL, and 5×10^4 cells were injected into the gastrocnemius muscle of

recipient mice. When tumors reached 70-150 mm^3 (as determined by caliper measurement in two

dimensions), the sarcomas were randomized to vehicle or digoxin groups. Mice were

administered an intraperitoneal (i.p.) injection of either vehicle (PBS) or 2 mg/kg digoxin

(prepared in DMSO and then diluted in PBS) with a volume not exceeding 250 µL. For short-

term treatments, injections were administered every 24 hours, until mice were euthanized via

cervical dislocation 3 hours after administration of the fourth dose. For long-term treatments,

injections were administered every 48 hours with tumor growth measured 3 times weekly until

sarcomas exceeded 13 mm in any dimension, at which point mice were euthanized via cervical

dislocation following IACUC guidelines at Duke University. Tumor, heart, kidney, liver, brain,

muscle, and plasma samples were collected and immediately snap-frozen in liquid nitrogen. For

the longitudinal treatment study, sections of tumors were preserved for histological analysis.

Histology and Microscopy

Fresh tumor samples were harvested after euthanasia, fixed in 4% PFA overnight, and preserved

in 70% ethanol until paraffin embedding. Immunohistochemistry was performed in order to stain

for CD31 as previously described[44]. Representative images of each H&E section were captured

using a Leica DM IL LED microscope equipped with a Leica MC170HD camera with a 20×

objective using LAS EZ software (Leica). Scale bars = 20 μm.

Single-cell RNA Sequencing

Tumors were dissected and minced following the manufacturer's protocol using MACS C tubes

and the mouse Tumor Dissociation Kit (Miltenyi Biotec). After tumor dissociation, the cells

were filtered through a 40 μM strainer. Red blood cells were lysed using ACK Lysing Buffer

(Lonza) and washed with flow buffer made of HBSS (cat 13175-095, Gibco), 5 mM EDTA

(E7899, Sigma-Aldrich), and 2.5% FBS (Gibco). Cells were washed twice more in 0.04% bovine

serum albumin (BSA) in PBS, then resuspended at 1000 cells per μL. Cell suspensions were

loaded on the 10x Genomics Chromium Controller Single-Cell Instrument (10x Genomics) using

the Chromium Single Cell 3' Reagent V3 Kit. Cells were mixed with reverse transcription

reagents, gel beads, and oil to generate single-cell gel beads in emulsions (GEM) for reverse

transcription (RT). After RT, GEMs were broken, and the single-stranded cDNA was purified

with DynaBeads. cDNA was amplified by PCR and the cDNA product was purified with the

SPRIselect Reagent Kit (Beckman Coulter). Sequencing libraries were constructed using the

reagents provided in the Chromium Single-Cell 3' Library Kit following the user guide.

Sequencing libraries were sequenced with the Illumina Novaseq 6000 platform at the Duke GCB

Sequencing and Genomic Technologies Core.

Single-cell RNA sequencing data processing

The single cell sequencing data were processed into the gene expression tables using the

pipelines from C*ell Ranger* v3.0.2 (https://support.10xgenomics.com/). Briefly, raw base call

files generated by Illumina sequencers were first demultiplexed into sample-specific FASTQ

files with the *cellranger mkfastq* pipeline. The FASTQ files for each sample were then aligned to

the mouse reference genome (mm10) using *STAR*[75]. The aligned reads for each gene were further

counted by the *cellranger count* pipeline. Quality control and filtering steps were performed to

remove the low-quality cells (with fewer than 1800 genes detected) and uninformative genes

(detected in fewer than 10 cells) for the downstream analyses.

Classification of single cells into malignant and non-malignant cells

Since malignant cells typically harbor large-scale copy number alteration (*i.e.* gains or deletions

of whole chromosomes or large chromosomal regions) that distinguish them from non-malignant

cells [76-79], we performed the copy number variations (CNVs) analysis for each sample to classify

the single cells into malignant and non-malignant cells. The copy number profiles were estimated

based on the average expression of large sets of genes in each chromosomal region using

inferCNV v0.99.7 (https://github.com/broadinstitute/inferCNV). A processed single-cell

transcriptomic dataset from limb muscles of two healthy mouse [42] was served as the reference for

CNVs calling. The same filtering procedures were applied to this reference dataset so that the

cells with less than 1800 expressed genes, and the genes expressed in fewer than 10 cells were

excluded. Genes with average expression values larger than 0.1 in reference cells were included

in the following analyses. The CNVs were estimated by sorting the genes according to their

chromosomal positions and using a moving average window with length 101 within each

chromosome. For each sample, the cells were separated into two clusters based on the

hierarchical clustering of CNV scores. We assigned the cluster with low-frequency CNVs like

the reference cells as the non-malignant, while the other cluster with high-frequency CNVs was

considered as the malignant.

Classification of cell populations within non-malignant cells

The Seurat v3.0.2 (http://satijalab.org/seurat/) R package was used for the identification of non-

malignant cell types. The gene expression data were firstly log-normalized and scaled with

default parameters. The top 2000 most variable genes selected by Seurat were used in the

principal component analysis (PCA). The first 85 principal components (PCs) selected based on

the built-in jackstraw analysis were used for downstream clustering analysis and t-SNE analysis.

Cell clusters were defined using FindClusters functions implemented in Seurat with default

parameters and resolution=0.15. The t-SNE analysis was used to visualize the clustering results

with perplexity setting to 1% of cell number whenever it was larger than 30 and learning rate

setting to 1/12 of cell number whenever it was above 200, as suggested by the previous study[80].

Each cluster was annotated by comparing it's specifically expressed genes with cell markers

reported in the literature[42] and CellMarker database[81] (http://biocc.hrbmu.edu.cn/CellMarker/). T

cells were further separated into different subtypes based on the following procedures: cells were

firstly classified as $CD8^+$ and $CD4^+$ based on the expression levels of gene *Cd8a* and *Cd4*. T

cells with *Cd8a* expression level larger than 0.1 were considered as the $CD8^+$. Similarly, those

with *Cd4* larger than 0.1 were considered as $CD4^+$ type. While the remaining cells with both *Cd4*

and *Cd8a* expression below than 0.1 were tentatively labeled as the double negative T cells.

$CD4^+$ T cells with the total expression level of *Foxp3* and *Il2ra* higher than 0.2 were further

labeled as Tregs, while other CD4$^+$ T cells were labeled as Ths. Hence the T cells were initially

classified as CD8$^+$, CD4$^+$ Tregs, CD4$^+$ Ths and double-negative T cells based on the expression

level of *Cd8, Cd4, Foxp3,* and *Il2ra*. We identified the differentially expressed genes

(Bonferroni-corrected p-value < 0.1) for these four groups of T cells using the FindAllMarkers

function in Seurat and then performed the PCA based on these identified differential genes. The

top 4 PCs selected based on jackstraw analysis were used for the next clustering analysis. We

then re-clustered these T cells into 5 groups using the FindClusters function with resolution=0.1

and annotated them as the CD8$^+$, Tregs, Ths, natural killer T cell (NKT) by comparing to known

cell markers, as well as an unknown cell group with no discernible markers.

Differential gene expression and pathway enrichment analysis

The Wilcoxon Rank Sum test was performed on metabolic genes to identify differences in

metabolism between single cells in vehicle and digoxin treatment groups. Cohen's d, a measure

of effect size, was calculated as below to estimate the magnitude of changes in gene expression

in response to Digoxin treatment.

$$Cohen's\ d = \frac{M_{dig} - M_{veh}}{Pooled\ SD} \tag{1}$$

500 Where M_{dig} and M_{veh} is the average of each gene's expression within a cell population from

501 digoxin and vehicle treatment groups, respectively. The *Pooled SD,* represents the population

502 standard deviation of a gene, is given by,

$$Pooled\ SD = \begin{cases} \sqrt{\dfrac{Sd_{dig}^2 + Sd_{veh}^2}{2}}, & n_{dig} = n_{veh} \\ \sqrt{\dfrac{(n_{dig}-1) \times Sd_{dig}^2 + (n_{veh}-1) \times Sd_{veh}^2}{n_{dig}+n_{veh}-2}}, & n_{dig} \neq n_{veh} \end{cases} \tag{2}$$

Where n_{dig} and n_{veh} are number of single cells in each population; the Sd_{dig} and S_{dveh} are standard

deviations. Lists of metabolic genes and pathways were obtained from the KEGG database

(https://www.kegg.jp/). The metabolic genes with p-value smaller than 0.01 (for malignant cells)

or 0.05 (for nonmalignant cells) and the absolute value of Cohen's d larger than 0.1 were

considered statistically significant and included in the pathway enrichment analysis. The one-

tailed Fisher's exact test was used to evaluate the enrichment significance of differential

metabolic genes in each metabolic pathway. GSEA analysis was performed using the javaGSEA

package available at https://www.gsea-msigdb.org/gsea/downloads.jsp with default parameters.

The completed differential gene sets (i.e. both metabolic and non-metabolic genes) were

searched against KEGG pathways using the Metascape (http://metascape.org).

Quantification and statistical analysis

All error bars were reported as +/- SEM with n=3 independent biological replicates and

statistical tests resulting in p-value computations were obtained using a two-tailed Student's t-

test unless otherwise noted. All statistics were computed using Graphpad Prism 6 (GraphPad,

http://graphpad.com/scientific-software/prism/) unless otherwise noted. * p < 0.05; **p < 0.01;

***p < 0.001.

Data Availability

The scRNA-seq data generated during this study are available in Gene Expression Omnibus

(GEO) under accession GSE149751 and can be visualized through the Single Cell Portal

(https://singlecell.broadinstitute.org/single_cell/study/SCP916).

Code Availability

Scripts reproducing the single-cell RNA-seq analysis are available at

https://github.com/LocasaleLab/mouse-sarcoma-scRNA

Author Contributions

Conceptualization, S.M.S. and J.W.L.; Computational Data Analysis, Z.X.; Animal Experiments, A.J.W., S.M.S., M.E.R., and D.G.K.; Single-cell RNA Sequencing, E.H. and S.G.G.; Metabolomics, S.M.S.; Isotope Tracing, S.M.S. and S.B.; Metabolic Flux and IC_{50} Correlations, M.V.L.; All Other Experiments, S.M.S.; Writing, S.M.S. with critical input from all authors; Supervision, S.M.S. and J.W.L.

Acknowledgments

We thank members of the Locasale laboratory for their helpful feedback and advice. We also thank members of the Kirsch laboratory for their assistance with the mouse studies. We would also like to thank the Duke GCB Sequencing and Genomic Technologies Core for their help with RNA sequencing. Work was supported by the National Institutes of Health (F31CA224973 to SMS, R01CA193256 to JWL, F30CA221268 to AJW, R35CA197616 to DGK). Support from the Duke Compute Cluster and Data Commons Storage for computational resources are gratefully acknowledged. J.W.L serves advisory roles in Nanocare Technologies, Raphael Pharmaceuticals, and Restoration Foodworks.

Figures

Figure 1 | Digoxin disrupts central carbon metabolism and related processes in a time- and dose-dependent manner.

(**A**) Spearman rank correlations of digoxin IC_{50} values and basal metabolic flux of individual metabolites in NCI-60 cancer cell panel. The metabolites are ranked based on the correlation coefficients.

(**B**) Experimental measurement of digoxin IC_{50} in HCT-116 cells after 48 hours.

(**C**) Clustered heatmaps of relative intensities of global metabolites upon digoxin treatment at indicated time points (left) and dose levels (right). Colored boxes highlight the most prominent metabolic changes.

(**D**) Enriched metabolic pathways determined from significantly altered metabolites ($p < 0.05$, Student's t-test) indicated in (C). The bar color matches box color in (C).

(**E and F**) Relative intensities of representative central carbon and energy metabolites at different time points (E) and dose levels (F).

(**G**) Diagram of isotopologue labeling of [U-^{13}C] glucose through glycolysis into the TCA cycle.

(**H**) Fractional abundance of each [U-^{13}C] glucose labeled isotopologue relative to the sum of all isotopologues of the glycolytic and TCA intermediates.

(**I**) Relative intensities of biologically relevant [U-^{13}C] glucose labeled isotopologues of lactate and succinate at different time points.

(**J**) Diagram of isotopologue labeling from [U-^{13}C] glutamine to the TCA cycle (left) and fractional abundance of each [U-^{13}C] glutamine-labeled isotopologue of TCA intermediates (right).

Figure 2 | Digoxin exerts its metabolic effects via on-target inhibition of the Na^+/K^+ ATPase.

(**A and B**) Relative cell viabilities of HCT-116 cells treated with vehicle or 100 nM digoxin in regulator media or with supplementation of the indicated nutrient for 72 hours. Nuc.: nucleosides.

(**C**) Relative cell viability of HCT-116 cells transfected with mATP1a1 digoxin-resistant subunit,

cultured in incremental concentrations of digoxin.

(**D**) Heatmap of fold changes in global metabolite levels with or without 48-hours 100 nM

digoxin treatment in HCT-116 transfected cells.

(**E and F**) Volcano plots displaying metabolites profiles of HCT-116 cells transfected with GFP

control (E) or mATP1a1 (F) vectors after treated with digoxin and vehicle.

(**G—J**) Relative intensities of TCA cycle metabolites (G), lactate (H), taurine (I) and creatine (J)

of HCT-116 cells transfected with GFP or mATP1a1 vectors upon digoxin and vehicle treatment.

Figure 3 | Digoxin treatment impacts energy metabolism in a tissue-specific and
antineoplastic manner.

(**A**) Diagram of digoxin treatment schedule in allograft sarcoma model. i.m., intramuscular; i.p.,

intraperitoneal injection.

(**B**) Volcano plots displaying metabolites profiles of cardiac tissue between vehicle and digoxin

treatment.

(**C**) Same as in (B) but for tumor tissue.

(**D**) Representative image of allograft sarcoma at growth endpoint. The tumor measured 12.8 mm

×13.4 mm at the time it was determined to have reached the endpoint.

(**E**) H&E sections of representative tumors in vehicle treatment, showing regions of both low

(left) and high (right) vascularity.

(**F**) Kaplan-Meier survival curve (left) and quantification of time for tumors to quintuple in

volume (right). Values in the right panel are represented as mean ± SD.

(**G**) Same as in (B) but for tumors exposed to chronic digoxin treatment.

(**H—J**) Relative intensities of TCA intermediates (H), taurine metabolites (I), and creatine (J) in

chronic vehicle- and digoxin-treated tumors. N=10.

Figure 4 | Acute digoxin treatment shifts the tumor microenvironment.

(**A**) Overview of single-cell RNA sequencing workflow, from tumor generation, cell harvest,

RNA sequencing, to gene expression analysis.

(**B**) Chromosomal landscape of large-scale CNVs for individual cells (rows) from normal mouse

muscle tissue and representative tumor vehicle replicate 1, allowing us to distinguish cells into

malignant and non-malignant. Amplifications (red) and deletions (blues) were inferred by

averaging gene expression over 100 genes stretch on each chromosome (columns).

(**C**) Relative proportion of malignant and non-malignant cells in each treatment.

(**D**) t-SNE plots show identified non-malignant cell populations (left) and T cell subpopulations

(right). M1: type I macrophages; M2: type II macrophages; M2α: highly proliferation M2; NK:

Natural killer cells; Tregs: Regulatory T cells; Th: T-helper cells; NKT: Natural killer T cells.

(**E**) Relative proportion of cell populations in the total non-malignant cell pool.

(**F**) Comparison of relative proportions of indicated cell populations in vehicle and digoxin

treatment.

Figure 5 | Digoxin treatment is associated with transcriptional reprogramming of metabolic
processes in malignant and non-malignant cells.

(**A**) Volcano plot of significance (measured by p value) against magnitude (measured by Cohen's

d) of metabolic gene expression differences between vehicle and digoxin treatment. The

differential genes above a significance threshold of p value < 0.01 and the absolute value of

Cohen's d > 0.1 are labeled in red.

(**B**) Metabolic pathways enriched in pathway analysis using differentially expressed metabolic

genes determined from (A). Rich factor is the ratio of the number of enriched genes (represented

by the size of the dots) to the number of background genes in the corresponding pathway.

(**C**) Distribution of Cohen's d for metabolic genes in each non-malignant cell population, with

significantly altered genes ($p < 0.05$ and | Cohens'd | > 0.1) marked in red (upregulated in

digoxin treatment) or blue (downregulated in digoxin treatment).

(**D**) Enriched metabolic pathways in non-malignant cell populations, determined by differentially

expressed metabolic genes in (C). For each cell population, the top 3 enriched pathways are

shown.

References
625

1   Pavlova, N. N. & Thompson, C. B. The Emerging Hallmarks of Cancer Metabolism. *Cell*
Metab **23**, 27-47 (2016).
2   Liberti, M. V. & Locasale, J. W. The Warburg Effect: How Does it Benefit Cancer Cells?
Trends Biochem Sci **41**, 211-218 (2016).
3   Miller, D. M., Thomas, S. D., Islam, A., Muench, D. & Sedoris, K. c-Myc and cancer
metabolism. *Clin Cancer Res* **18**, 5546-5553 (2012).
4   Papa, S., Choy, P. M. & Bubici, C. The ERK and JNK pathways in the regulation of
metabolic reprogramming. *Oncogene* **38**, 2223-2240 (2019).
5   Serna-Blasco, R., Sanz-Alvarez, M., Aguilera, O. & Garcia-Foncillas, J. Targeting the
RAS-dependent chemoresistance: The Warburg connection. *Semin Cancer Biol* **54**, 80-90
(2019).
6   Baysal, B. E. *et al.* Mutations in SDHD, a mitochondrial complex II gene, in hereditary
paraganglioma. *Science* **287**, 848-851 (2000).
7   Xu, W. *et al.* Oncometabolite 2-hydroxyglutarate is a competitive inhibitor of alpha-
ketoglutarate-dependent dioxygenases. *Cancer Cell* **19**, 17-30 (2011).
8   Selak, M. A. *et al.* Succinate links TCA cycle dysfunction to oncogenesis by inhibiting
HIF-alpha prolyl hydroxylase. *Cancer Cell* **7**, 77-85 (2005).
9   Gaude, E. & Frezza, C. Defects in mitochondrial metabolism and cancer. *Cancer Metab*
2, 10 (2014).
10   Gatenby, R. A. & Gillies, R. J. Why do cancers have high aerobic glycolysis? *Nat Rev*
Cancer **4**, 891-899 (2004).
11   Weljie, A. M. & Jirik, F. R. Hypoxia-induced metabolic shifts in cancer cells: moving
beyond the Warburg effect. *Int J Biochem Cell Biol* **43**, 981-989 (2011).
12   Chang, C. H. *et al.* Metabolic Competition in the Tumor Microenvironment Is a Driver of
Cancer Progression. *Cell* **162**, 1229-1241 (2015).
13   Hay, N. Reprogramming glucose metabolism in cancer: can it be exploited for cancer
therapy? *Nat Rev Cancer* **16**, 635-649 (2016).
14   Luengo, A., Gui, D. Y. & Vander Heiden, M. G. Targeting Metabolism for Cancer
Therapy. *Cell Chem Biol* **24**, 1161-1180 (2017).
15   Barry, W. H., Hasin, Y. & Smith, T. W. Sodium pump inhibition, enhanced calcium
influx via sodium-calcium exchange, and positive inotropic response in cultured heart
cells. *Circ Res* **56**, 231-241 (1985).
16   Kaplan, J. H. Biochemistry of Na,K-ATPase. *Annu Rev Biochem* **71**, 511-535 (2002).
17   Clausen, T. Potassium and sodium transport and pH regulation. *Can J Physiol Pharmacol*
70 Suppl, S219-222 (1992).
18   Wright, E. M., Hirayama, B. A. & Loo, D. F. Active sugar transport in health and
disease. *J Intern Med* **261**, 32-43 (2007).
19   Sibarov, D. A., Bolshakov, A. E., Abushik, P. A., Krivoi, II & Antonov, S. M. Na+,K+-
ATPase functionally interacts with the plasma membrane Na+,Ca2+ exchanger to prevent
Ca2+ overload and neuronal apoptosis in excitotoxic stress. *J Pharmacol Exp Ther* **343**,
596-607 (2012).
20   Kepp, O. *et al.* Anticancer activity of cardiac glycosides: At the frontier between cell-
autonomous and immunological effects. *Oncoimmunology* **1**, 1640-1642 (2012).
21   Slingerland, M., Cerella, C., Guchelaar, H. J., Diederich, M. & Gelderblom, H. Cardiac
glycosides in cancer therapy: from preclinical investigations towards clinical trials. *Invest*
New Drugs **31**, 1087-1094 (2013).

22   Stenkvist, B. *et al.* Evidence of a modifying influence of heart glucosides on the
673   development of breast cancer. *Anal Quant Cytol* **2**, 49-54 (1980).
674   23   Racker, E. Bioenergetics and the problem of tumor growth. *Am Sci* **60**, 56-63 (1972).
675   24   Howarth, C., Gleeson, P. & Attwell, D. Updated energy budgets for neural computation
676   in the neocortex and cerebellum. *J Cereb Blood Flow Metab* **32**, 1222-1232 (2012).
677   25   Milligan, L. P. & McBride, B. W. Energy costs of ion pumping by animal tissues. *J Nutr*
678   **115**, 1374-1382 (1985).
679   26   Shestov, A. A. *et al.* Quantitative determinants of aerobic glycolysis identify flux through
680   the enzyme GAPDH as a limiting step. *Elife* **3** (2014).
681   27   Jong, C. J., Azuma, J. & Schaffer, S. Mechanism underlying the antioxidant activity of
682   taurine: prevention of mitochondrial oxidant production. *Amino Acids* **42**, 2223-2232
683   (2012).
684   28   Kurosawa, Y. *et al.* Creatine supplementation enhances anaerobic ATP synthesis during a
685   single 10 sec maximal handgrip exercise. *Mol Cell Biochem* **244**, 105-112 (2003).
686   29   Kim, N. *et al.* Cardiac glycosides display selective efficacy for STK11 mutant lung
687   cancer. *Sci Rep* **6**, 29721 (2016).
688   30   Calderon-Montano, J. M., Burgos-Moron, E. & Lopez-Lazaro, M. The in vivo antitumor
689   activity of cardiac glycosides in mice xenografted with human cancer cells is probably an
690   experimental artifact. *Oncogene* **33**, 2947-2948 (2014).
691   31   Gupta, R. S., Chopra, A. & Stetsko, D. K. Cellular basis for the species differences in
692   sensitivity to cardiac glycosides (digitalis). *J Cell Physiol* **127**, 197-206 (1986).
693   32   Eskiocak, U. *et al.* Synergistic effects of ion transporter and MAP kinase pathway
694   inhibitors in melanoma. *Nat Commun* **7**, 12336 (2016).
695   33   Zhang, H. *et al.* Digoxin and other cardiac glycosides inhibit HIF-1alpha synthesis and
696   block tumor growth. *Proc Natl Acad Sci U S A* **105**, 19579-19586 (2008).
697   34   Calderon-Montano, J. M. *et al.* Evaluating the cancer therapeutic potential of cardiac
698   glycosides. *Biomed Res Int* **2014**, 794930 (2014).
699   35   Huang, L. *et al.* Systems biology-based drug repositioning identifies digoxin as a
700   potential therapy for groups 3 and 4 medulloblastoma. *Sci Transl Med* **10** (2018).
701   36   Brizel, D. M., Rosner, G. L., Prosnitz, L. R. & Dewhirst, M. W. Patterns and variability
702   of tumor oxygenation in human soft tissue sarcomas, cervical carcinomas, and lymph
703   node metastases. *Int J Radiat Oncol Biol Phys* **32**, 1121-1125 (1995).
704   37   Morton, C. L. & Houghton, P. J. Establishment of human tumor xenografts in
705   immunodeficient mice. *Nat Protoc* **2**, 247-250 (2007).
706   38   Buque, A. & Galluzzi, L. Modeling Tumor Immunology and Immunotherapy in Mice.
707   *Trends Cancer* **4**, 599-601 (2018).
708   39   Sukumar, M. *et al.* Inhibiting glycolytic metabolism enhances CD8+ T cell memory and
709   antitumor function. *J Clin Invest* **123**, 4479-4488 (2013).
710   40   Chang, C. H. *et al.* Posttranscriptional control of T cell effector function by aerobic
711   glycolysis. *Cell* **153**, 1239-1251 (2013).
712   41   Patel, C. H., Leone, R. D., Horton, M. R. & Powell, J. D. Targeting metabolism to
713   regulate immune responses in autoimmunity and cancer. *Nat Rev Drug Discov* **18**, 669-
714   688 (2019).
715   42   Tabula Muris, C. *et al.* Single-cell transcriptomics of 20 mouse organs creates a Tabula
716   Muris. *Nature* **562**, 367-372 (2018).

43 Lee, C. L. *et al.* Mutational landscape in genetically engineered, carcinogen-induced, and
radiation-induced mouse sarcoma. *JCI Insight* **4** (2019).

44 Wisdom, A. J. *et al.* Neutrophils promote tumor resistance to radiation therapy. *Proc Natl*
Acad Sci U S A (2019).

45 Yadav, N. *et al.* Oxidative phosphorylation-dependent regulation of cancer cell apoptosis
in response to anticancer agents. *Cell Death Dis* **6**, e1969 (2015).

46 Xiao, Z., Dai, Z. & Locasale, J. W. Metabolic landscape of the tumor microenvironment
at single cell resolution. *Nat Commun* **10**, 3763 (2019).

47 Newsholme, P., Curi, R., Gordon, S. & Newsholme, E. A. Metabolism of glucose,
glutamine, long-chain fatty acids and ketone bodies by murine macrophages. *Biochem J*
239, 121-125 (1986).

48 Newsholme, P., Gordon, S. & Newsholme, E. A. Rates of utilization and fates of glucose,
glutamine, pyruvate, fatty acids and ketone bodies by mouse macrophages. *Biochem J*
242, 631-636 (1987).

49 Gagnon, E. *et al.* Endoplasmic reticulum-mediated phagocytosis is a mechanism of entry
into macrophages. *Cell* **110**, 119-131 (2002).

50 Oh, J. *et al.* Endoplasmic reticulum stress controls M2 macrophage differentiation and
foam cell formation. *J Biol Chem* **287**, 11629-11641 (2012).

51 Canton, J. Phagosome maturation in polarized macrophages. *J Leukoc Biol* **96**, 729-738
(2014).

52 Pan, M. *et al.* Regional glutamine deficiency in tumours promotes dedifferentiation
through inhibition of histone demethylation. *Nat Cell Biol* **18**, 1090-1101 (2016).

53 Davidson, S. M. *et al.* Environment Impacts the Metabolic Dependencies of Ras-Driven
Non-Small Cell Lung Cancer. *Cell Metab* **23**, 517-528 (2016).

54 Erecinska, M. & Dagani, F. Relationships between the neuronal sodium/potassium pump
and energy metabolism. Effects of K+, Na+, and adenosine triphosphate in isolated brain
synaptosomes. *J Gen Physiol* **95**, 591-616 (1990).

55 Bensaad, K. & Harris, A. L. Hypoxia and metabolism in cancer. *Adv Exp Med Biol* **772**,
1-39 (2014).

56 Moghetti, P., Bacchi, E., Brangani, C., Dona, S. & Negri, C. Metabolic Effects of
Exercise. *Front Horm Res* **47**, 44-57 (2016).

57 Hennings, J. M. *et al.* Effect of mirtazapine on metabolism and energy substrate
partitioning in healthy men. *JCI Insight* **4** (2019).

58 Ballon, J. S. *et al.* Pathophysiology of drug induced weight and metabolic effects:
findings from an RCT in healthy volunteers treated with olanzapine, iloperidone, or
placebo. *J Psychopharmacol* **32**, 533-540 (2018).

59 Madiraju, A. K. *et al.* Metformin suppresses gluconeogenesis by inhibiting mitochondrial
glycerophosphate dehydrogenase. *Nature* **510**, 542-546 (2014).

60 Liu, X., Romero, I. L., Litchfield, L. M., Lengyel, E. & Locasale, J. W. Metformin
Targets Central Carbon Metabolism and Reveals Mitochondrial Requirements in Human
Cancers. *Cell Metab* **24**, 728-739 (2016).

61 Chan, K. K., Oza, A. M. & Siu, L. L. The statins as anticancer agents. *Clin Cancer Res* **9**,
10-19 (2003).

62 Mills, E. J. *et al.* Low-dose aspirin and cancer mortality: a meta-analysis of randomized
trials. *Am J Med* **125**, 560-567 (2012).

63   Wheaton, W. W. *et al.* Metformin inhibits mitochondrial complex I of cancer cells to
reduce tumorigenesis. *Elife* **3**, e02242 (2014).

64   Banerjee, M. *et al.* Na/K-ATPase Y260 Phosphorylation-mediated Src Regulation in
Control of Aerobic Glycolysis and Tumor Growth. *Sci Rep* **8**, 12322 (2018).

65   Felippe Goncalves-de-Albuquerque, C., Ribeiro Silva, A., Ignacio da Silva, C., Caire
Castro-Faria-Neto, H. & Burth, P. Na/K Pump and Beyond: Na/K-ATPase as a
Modulator of Apoptosis and Autophagy. *Molecules* **22** (2017).

66   Xie, Z. & Askari, A. Na(+)/K(+)-ATPase as a signal transducer. *Eur J Biochem* **269**,
2434-2439 (2002).

67   Epstein, T., Xu, L., Gillies, R. J. & Gatenby, R. A. Separation of metabolic supply and
demand: aerobic glycolysis as a normal physiological response to fluctuating energetic
demands in the membrane. *Cancer Metab* **2**, 7 (2014).

68   Schaffer, S. W., Solodushko, V. & Kakhniashvili, D. Beneficial effect of taurine
depletion on osmotic sodium and calcium loading during chemical hypoxia. *Am J Physiol
Cell Physiol* **282**, C1113-1120 (2002).

69   Katz, A. *et al.* Selectivity of digitalis glycosides for isoforms of human Na,K-ATPase. *J
Biol Chem* **285**, 19582-19592 (2010).

70   Najafi, M. *et al.* Macrophage polarity in cancer: A review. *J Cell Biochem* **120**, 2756-
2765 (2019).

71   Huh, J. R. *et al.* Digoxin and its derivatives suppress TH17 cell differentiation by
antagonizing RORgammat activity. *Nature* **472**, 486-490 (2011).

72   Zhyvoloup, A. *et al.* Digoxin reveals a functional connection between HIV-1 integration
preference and T-cell activation. *PLoS Pathog* **13**, e1006460 (2017).

73   Sen, T. *et al.* Targeting DNA Damage Response Promotes Antitumor Immunity through
STING-Mediated T-cell Activation in Small Cell Lung Cancer. *Cancer Discov* **9**, 646-
661 (2019).

74   Kuntz, E. M. *et al.* Targeting mitochondrial oxidative phosphorylation eradicates therapy-
resistant chronic myeloid leukemia stem cells. *Nat Med* **23**, 1234-1240 (2017).

75   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).

76   Puram, S. V. *et al.* Single-Cell Transcriptomic Analysis of Primary and Metastatic Tumor
Ecosystems in Head and Neck Cancer. *Cell* **171**, 1611-1624 e1624 (2017).

77   Patel, A. P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in primary
glioblastoma. *Science* **344**, 1396-1401 (2014).

78   Venteicher, A. S. *et al.* Decoupling genetics, lineages, and microenvironment in IDH-
mutant gliomas by single-cell RNA-seq. *Science* **355** (2017).

79   Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-
cell RNA-seq. *Science* **352**, 189-196 (2016).

80   Kobak, D. & Berens, P. The art of using t-SNE for single-cell transcriptomics. *bioRxiv*
(2019).

81   Zhang, X. *et al.* CellMarker: a manually curated resource of cell markers in human and
mouse. *Nucleic Acids Res* **47**, D721-D728 (2019).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5