1

HbtR, a heterofunctional homolog of the virulence regulator TcpP, facilitates the

transition between symbiotic and planktonic lifestyles in *Vibrio fischeri*.

Brittany D. Bennett, Tara Essock-Burns, and Edward G. Ruby

Pacific Biosciences Research Center, University of Hawaiʻi—Manoa, Honolulu, HI

96813

Running head: Regulation of lifestyle transition in *V. fischeri*

Address correspondence: eruby@hawaii.edu

Abstract

The bioluminescent bacterium *Vibrio fischeri* forms a mutually beneficial symbiosis with the Hawaiian bobtail squid, *Euprymna scolopes*, in which the bacteria, housed inside a specialized light organ, produce light used by the squid in its nocturnal activities. Upon hatching, *E. scolopes* juveniles acquire *V. fischeri* from the seawater through a complex process that requires, among other factors, chemotaxis by the bacteria along a gradient of *N*-acetylated sugars into the crypts of the light organ, the niche in which the bacteria reside. Once inside the light organ, *V. fischeri* transitions into a symbiotic, sessile state in which the quorum-signaling regulator LitR induces luminescence. In this work we show that expression of *litR* and luminescence are repressed by a homolog of the *V. cholerae* virulence factor TcpP, which we have named HbtR. Further, we demonstrate that LitR represses genes involved in motility and chemotaxis into the light organ and activates genes required for exopolysaccharide production.

Importance

TcpP homologs are widespread throughout the *Vibrio* genus; however, the only protein in this family described thus far is a *V. cholerae* virulence regulator. Here we show that HbtR, the TcpP homolog in *V. fischeri*, has both a biological role and regulatory pathway completely unlike that in *V. cholerae*. Through its repression of the quorum-signaling regulator LitR, HbtR affects the expression of genes important for colonization of the *E. scolopes* light organ. While LitR becomes activated within the crypts, and upregulates luminescence and exopolysaccharide genes and downregulates chemotaxis and motility genes, it appears that HbtR, upon expulsion of *V. fischeri* cells into seawater, reverses this process to aid the switch from a symbiotic to a planktonic state. The

40 possible importance of HbtR to the survival of *V. fischeri* outside of its animal host may

41 have broader implications for the ways in which bacteria transition between often vastly

42 different environmental niches.

Introduction

The Gram-negative bacterium *Vibrio (Aliivibrio) fischeri* is a model organism for the

study of biochemical processes underpinning bioluminescence, quorum sensing, and

bacterial-animal symbioses. Luminescence in *V. fischeri* is activated by a quorum-

sensing pathway that responds to high concentrations of two autoinducer molecules, *N*-

(3-oxohexanoyl)-L-homoserine lactone (3-oxo-C6-HSL (1)) and *N*-octanoyl-L-

homoserine lactone (C8-HSL (2)). When *V. fischeri* cells multiply to a certain density, 3-

oxo-C6-HSL and C8-HSL accumulate to high local concentrations, initiating a signaling

cascade that leads to upregulation of the regulator LuxR by the regulator LitR and

consequent activation of the luciferase operon, *luxICDABEG* (3–5); reviewed in (6).

Light production by *V. fischeri* is a crucial factor in the mutualistic symbiosis it

forms within the light-emitting organ of the Hawaiian bobtail squid, *Euprymna scolopes*

(7–9). Juvenile *E. scolopes* become colonized with *V. fischeri* shortly after hatching into

seawater containing this bacterium (10). *V. fischeri* cells initiate light-organ colonization

through a series of steps, including chemotaxis towards *N*-acetylated sugars released

by the squid (11, 12). After initial colonization of the squid light organ, the symbiosis

undergoes a daily cyclic rhythm of three basic stages for the remainder of the squid's

life: during the day, *V. fischeri* cells grow to a high density in the crypts on carbon

sources provided by the squid (13, 14); at night, the bacteria produce light that aids in

camouflage for the squid (15, 16); and at dawn, ~95% of the bacterial cells are expelled

from the light organ into the seawater, where they may initiate colonization of new squid

hatchlings, while the remaining ~5% repopulate the light organ (17, 18).

Previous work indicated that, in *V. fischeri* cells newly expelled from the light

organ, numerous genes are up- and downregulated relative to their level of expression

in cells that had been planktonic for some time (19). Two of the genes upregulated in

expelled cells were VF_A0473 and VF_A0474, comprising a small operon annotated as

encoding homologs of the genetic regulator TcpP and its chaperone, TcpH. TcpP was

first described in *V. cholerae* as a virulence gene regulator (20, 21). During the early

stages of *V. cholerae* infection, TcpP works synergistically with another transcriptional

regulator, ToxR, to upregulate expression of the virulence regulator *toxT* and thereby

activate expression of cholera toxin and the toxin-coregulated pilus in response to

changing environmental conditions (21, 22). *tcpPH* expression is itself induced by AphA

and AphB in response to low oxygen or acidic pH (23–25). In *V. cholerae* El Tor

biotypes, *aphA* expression is repressed by HapR (the *V. cholerae* ortholog of LitR) upon

initiation of quorum sensing at high cell densities (26, 27). Additionally, activation of

tcpPH expression by AphA and AphB is inhibited by the global metabolic regulator Crp

(28). This complex regulatory pathway serves to upregulate *tcpPH* and, consequently,

downstream virulence factors under conditions consistent with entry into the vertebrate

gut, a process reversed late in the infection cycle before the bacteria exit the host and

re-enter an aquatic environment (29, 30).

No published genomes of *V. fischeri* strains encode homologs of either *toxT* or

cholera toxin genes. The genome of the model strain *V. fischeri* ES114 does encode

annotated homologs of the toxin-coregulated pilus genes *tcpFETSD* and *tcpCQBA*, but

this strain is alone among the 66 currently available *V. fischeri* genomes to do so. The

V. fischeri genes regulated by the protein product of VF_A0473 are therefore unknown,

as are any upstream regulatory factors that determine under which conditions the

regulator is produced and active. Here we present evidence that VF_A0473 regulates

genes governing phenotypes relevant to the switch from symbiosis to a planktonic

lifestyle. We therefore rename VF_A0473 and VF_A0474, currently annotated as *tcpP*

and *tcpH*, as *hbtR* (*hab*itat *t*ransition *r*egulator) and *hbtC* (*hab*itat *t*ransition *c*haperone).

In this work, we identify the HbtR regulon and uncover aspects of *hbtRC* regulation.

Further, we determine that LitR is repressed by HbtR and describe LitR-regulated

phenotypes beyond luminescence that are important for light-organ colonization.

Results

<u>The HbtR regulon is distinct from that of TcpP.</u> Considering that *V. fischeri* and *V.*

cholerae have significantly different lifestyles, as well as the absence in 65 of 66

sequenced *V. fischeri* genomes of homologs of virulence-factor genes regulated by

TcpP in *V. cholerae*, we postulated that HbtR has a distinct function from that of TcpP.

To determine the regulon of HbtR, RNA-seq was performed using Δ*hbtRC* mutant

strains of *V. fischeri* ES114 carrying either empty vector or an inducible vector with

hbtRC under control of the *lac* promoter. Notable among the RNA-seq results (Table

S1) is that none of the *tcp* pilus gene homologs encoded in the *V. fischeri* ES114

genome were expressed at significantly different levels in *hbtRC* strains carrying either

empty vector or the inducible *hbtRC* vector (Table S1), ruling out the possibility that

HbtR regulates the same genes in *V. fischeri* ES114 as TcpP does in *V. cholerae*.

<u>HbtR represses *litR* expression and luminescence *in vitro*.</u> Among the genes

significantly downregulated in the RNA-seq results when *hbtRC* was expressed were

the quorum signaling-regulated genes *litR*, *luxR*, *qsrP*, and *rpoQ* (Table 1, Table S1).

Less strongly, but statistically significantly, downregulated genes included most of the

luciferase operon (Table 1, Table S1). To determine whether repression of *lux* genes by

HbtR affects bacterial light emission, luminescence and growth were monitored in

cultures of wild-type *V. fischeri* and its Δ*hbtRC* derivative carrying either empty vector or

vector constitutively expressing *hbtRC*. Deletion of *hbtRC* did not affect light production;

however, overexpression of *hbtRC* in either wild type or Δ*hbtRC* led to a delay in onset

and significant decrease in intensity of light production (Fig. 1A). There was no

difference in growth rate between the strains (Fig. 1B), eliminating the possibility that

the reduced luminescence by the *hbtRC*-overexpression strains was due to a growth

defect. To determine whether deletion of *hbtRC* would lead to a change in light

production by *V. fischeri* within the light organ, *E. scolopes* hatchlings were colonized

with wild-type *V. fischeri* or Δ*hbtRC*. There was no difference in the luminescence of

juvenile squid colonized by either *V. fischeri* strain at either 24 or 48 h (Fig. S1),

suggesting that HbtR is not active inside the light organ.

<u>HbtR activates and LitR represses chemosensory genes involved in *E. scolopes* light-</u>

<u>organ colonization</u>. Among the genes upregulated in the *hbtRC*-expressing strain in the

RNA-seq experiment were four genes annotated as encoding methyl-accepting

chemotaxis proteins (MCPs): VF_1133, VF_2042, VF_A0246, and VF_A0389 (Table 1,

Table S1). To confirm that the upregulation of these MCP genes by HbtR functions

through its repression of *litR* expression, RT-qPCR was performed on wild-type *V.*

fischeri and Δ*litR*, as well as Δ*hbtRC* carrying either empty vector or vector

constitutively expressing *hbtRC*. Expression of all four of these MCP genes was

significantly increased in Δ*litR* compared to wild-type *V. fischeri* (Table 2). Expression of

VF_2042 was significantly increased in Δ*hbtRC* constitutively expressing *hbtRC*

compared to Δ*hbtRC* carrying empty vector (Table 2), confirming the RNA-seq results

indicating that HbtR affects MCP gene regulation.

As chemotaxis is essential to colonization of the *E. scolopes* light organ by *V.*

fischeri (12), we asked whether any of the four MCPs repressed by LitR are involved in

colonization. Newly hatched *E. scolopes* juveniles were exposed to a 1:1 ratio of wild-

type *V. fischeri* and a quadruple mutant with in-frame deletions of all four of the LitR-

repressed MCP genes (ΔVF_1133ΔVF_2042ΔVF_A0246ΔVF_A0389; "ΔΔΔΔ").

Regardless of which strain carried a *gfp*-labeled plasmid, the squid light organ

populations tended to be dominated by wild-type *V. fischeri* (Fig. 2), indicating that HbtR

and LitR regulate chemotaxis genes involved in initiating symbiosis.

To determine whether the colonization defect displayed by ΔΔΔΔ is due to the

chemosensory function of any of the four MCP genes, we sought to identify the

chemoattractants recognized by these MCPs. Wild-type *V. fischeri* and MCP deletion

mutants were spotted onto minimal-medium, soft-agar plates containing 1 mM of three

known chemoattractants for *V. fischeri* (12): *N*-acetyl-D-glucosamine (GlcNAc), *N,N′*-

diacetylchitobiose ($(GlcNAc)_2$), or *N*-acetylneuraminic acid (Neu5Ac). The chemotactic

zone sizes for ΔVF_1133 and ΔVF_A0246 were significantly smaller than for wild-type

V. fischeri on each of the *N*-acetylated sugars (Table 3). Oddly, while the zone sizes of

ΔΔΔΔ on GlcNAc and $(GlcNAc)_2$ were similar to those of ΔVF_1133 and ΔVF_A0246,

on Neu5Ac the ΔΔΔΔ zone size was comparable to wild type. This is a repeatable

phenotype we have yet to explain. However, a ΔVF_1133 ΔVF_A0246 mutant produced

a similar zone size to the single mutants on Neu5Ac. There was no significant difference

in zone sizes between ΔΔΔΔ and wild-type *V. fischeri* spotted onto plates containing

either no chemoattractant or 1 mM glucose (Table 3), indicating these mutations do not

affect motility or chemotaxis in general. Because they grew equally well on either

GlcNAc or Neu5AC (Fig. S2), the differences in zone sizes of the MCP mutants and

wild-type *V. fischeri* were unlikely to be growth-related.

<u>HbtR activates and LitR represses motility.</u> Two flagellin genes, *flaC* and *flaF*, were

upregulated by *hbtRC* (Table 1, Table S1), indicating the possible activation of motility

by HbtR. An earlier study demonstrated that LitR represses motility in *V. fischeri* (31),

and microarrays performed previously (Studer, Schaefer, and Ruby; Schaefer and Ruby

unpublished data) indicated that AinS and LitR downregulate the expression of a

majority of the flagellin genes within the locus VF_1836–77, as well as *flaF*, *fliL2*, *motX*,

motA1B1, and *cheW*. To confirm the repression of motility genes by LitR, RT-qPCR was

employed to determine the level of expression of three flagellin or motor genes in

different genomic regions in wild-type *V. fischeri* and Δ*litR*. Expression of *cheW*, *motA1*,

flaC, and *flaF* was moderately, but statistically significantly, higher in Δ*litR* than in wild-

type *V. fischeri* (Table 2). However, we were not able to determine the mechanism by

which LitR represses multiple motility gene loci, as expression levels of the known

flagellin regulatory genes *rpoN*, *flrA*, *flrB*, *fliA*, and *flgM* were comparable between Δ*litR*

and wild-type *V. fischeri* strains (Table 2).

The sigma factor-like regulator RpoQ, which is upregulated by LitR, represses

motility and/or chemotaxis towards GlcNAc when overexpressed (32). Thus, we wanted

to determine whether the effect LitR has on motility and *N*-acetylated sugar chemotaxis

is mediated through its regulation of *rpoQ*. On soft-agar chemotaxis plates containing no

chemoattractant or 1 mM GlcNAc, we observed the same pattern regardless of the

presence of GlcNAc: either deletion of *litR* or overexpression of *hbtRC* led to larger

migration zones than for wild-type *V. fischeri*, while overexpression of *litR* essentially

eliminated bacterial migration (Table 3). There were no significant differences between

Δ*rpoQ* and wild-type *V. fischeri* strains carrying the same vector (Table 3), indicating

that the repression of motility by LitR is not mediated through RpoQ.

HbtR represses and LitR activates exopolysaccharide production. An approximately 25-

kb locus constituting the genes VF_0157–80 was downregulated when *hbtRC* was

expressed (Table 1, Table S1). Previously performed microarrays indicated that most of

the genes in this locus (VF_0159 and VF_0162–0180) are also upregulated by AinS

and LitR (Studer, Schaefer, and Ruby; Schaefer and Ruby, unpublished data). To

confirm the upregulation of this locus by LitR, RT-qPCR was used to evaluate the

expression of three genes, located in separate operons, in Δ*litR* and wild-type *V.*

fischeri. Expression of VF_0157 (*wbfB*), VF_0164 (*etp*), and VF_0165 (*wzc*) was

significantly higher in wild-type *V. fischeri* than in Δ*litR* (Table 2). Expression of *etp* was

significantly lower in Δ*hbtRC* constitutively expressing *hbtRC* compared to Δ*hbtRC*

carrying empty vector (Table 2), confirming that HbtR regulation of *litR* represses gene

expression in this locus.

Most of the genes in the locus VF_0157–80 are annotated as being involved in

production of extracellular polysaccharides. However, gene annotation alone could not

indicate which component(s) of the cell envelope (lipopolysaccharide O-antigen,

capsule, or exopolysaccharide (EPS)) might be affected by the genes in this locus. To

determine which polysaccharide type is produced by the enzymes encoded in

VF_0157–80, alcian blue staining was used to detect negatively charged

polysaccharides in the supernatants of liquid cultures of *V. fischeri* strains. Both Δ*litR*

and ΔVF0157–80 produced less alcian blue-staining EPS than wild-type *V. fischeri*, with

ΔVF_0157–80 producing even less EPS than Δ*litR* (Fig. 3). Overexpression of *hbtRC*

repressed EPS production in wild-type *V. fischeri*, but not in Δ*litR* or ΔVF_0157–80 (Fig.

3). Overexpression of *litR* in ΔVF_0157–80 did not complement the reduction in EPS

produced by that mutant (Fig. 3), indicating that essentially all EPS is produced by this

locus.

<u>*hbtRC* expression is regulated by Crp, but not by AphB or quorum sensing</u>. To

determine whether *hbtRC* expression in *V. fischeri* is affected by quorum sensing and/or

Crp, a regulator known to affect *tcpPH* expression in *V. cholerae*, RT-qPCR was

performed on wild-type *V. fischeri*, Δ*litR*, and Δ*crp* grown to OD_{600} ~0.3 and ~1.0.

Expression of *hbtRC* was lower at high cell density than at low cell density for all three

strains, with no difference between wild-type *V. fischeri* and Δ*litR* at either higher or

lower OD_{600} (Table 2). Expression of *hbtRC* in Δ*crp* was below that in wild-type *V.*

fischeri at low OD_{600}, but comparable at high OD_{600}. Thus, Crp appears to activate

hbtRC expression, the reverse of *tcpPH* regulation by Crp in *V. cholerae*.

Because Crp activates quorum-signaling genes in the *V. fischeri* genome (33,

34), we asked whether quorum signaling is responsible for the change in *hbtRC*

expression at different culture densities, and whether this regulation explains the lower

expression in the Δ*crp* mutant. The RT-qPCR experiment was repeated for wild-type *V.*

fischeri grown in SWT with or without added autoinducers. *hbtRC* expression remained

higher at lower cell density in the presence of both autoinducers (Table 2), indicating

that quorum signaling is not involved in Crp- and cell density-mediated *hbtRC*

regulation*.*

To determine whether the regulator AphAB and/or a change in pH affect *hbtRC*

expression in *V. fischeri*, RT-qPCR was performed on RNA extracted from cultures of

wild type or Δ*aphB V. fischeri* grown at pH 5.5 or 8.5. There was no significant

difference in *hbtRC* expression between any of the conditions (Table S2).

tcpPH homologs do not cross-complement between *V. fischeri* and *V. cholerae*. Δ*tcpPH*

and Δ*hbtRC* strains of *V. cholerae* and *V. fischeri*, respectively, were complemented

with empty vector, vector expressing *tcpPH*, or vector expressing *hbtRC*. RT-qPCR was

performed on RNA extracted from cultures of each strain to determine the level of *toxT*

expression in the *V. cholerae* strains or *litR* expression in the *V. fischeri* strains. While

expression of each *tcpPH* homolog complemented the respective deletion, cross-

complementation had no effect on output gene expression (Table S2), illustrating the

divergent activities of HbtR and TcpP.

In *V. cholerae*, ToxR cooperates with TcpP to activate *toxT* expression (35, 36),

begging the question of whether ToxR is also involved in controlling the *V. fischeri* HbtR

regulon. RT-qPCR performed on RNA extracted from cultures of Δ*hbtRC* and

Δ*hbtRC*Δ*toxRS* carrying either empty vector or vector expressing *hbtRC* showed that

litR expression was repressed to the same degree by HbtR regardless of the presence

or absence of *toxRS* (Table S2), further demonstrating that regulation by HbtR is

independent of ToxR and differs substantially from the TcpP system.

Δ*hbtRC* has no defect in colonizing juvenile *E. scolopes* light organs. Previous work

indicated that Δ*hbtRC* (then referred to as Δ*tcpPH*) had a light-organ colonization

defect, which appeared to increase through 96 h (19). We replicated this defect when

using the same conditions as the previous study (Fig. 4A); however, we observed an

advantage for Δ*hbtRC* when we reversed the fluorescent marker plasmids (Fig. 4A),

indicating the earlier results were due simply to experimental design. Co-colonization of

juvenile *E. scolopes* with chromosomally labeled strains resulted in no significant

colonization defect for either strain (Fig. 4B), demonstrating that HbtR is, in fact, not

required for entry into, or life inside, the *E. scolopes* light organ. Based on this

experience, we urge caution by other researchers using *rfp*-labeled pVSV208 in co-

colonization experiments.

Transcription of *hbtR* is activated as symbionts exit the light organ. Considering the

effects HbtR has on chemotaxis and luminescence gene expression (Tables 1, 2, and

S1), and that *hbtRC* has no effect on squid luminescence (Fig. S1) or colonization (Fig.

4B), we postulated that HbtR was likely to be involved in the transition into planktonic

life as *V. fischeri* exits the light organ. To establish at which point in the juvenile squid

diel cycle *V. fischeri hbtRC* expression is activated, we performed *in situ* hybridization

chain reaction (HCR) targeting *hbtR* mRNA to determine whether this gene was

expressed in the bacteria before, during, or after their expulsion from the squid light

organ into seawater. *hbtR* mRNA was not detected in *V. fischeri* cells within the crypts

or during expulsion along the path out of the light organ (Fig. 5A). Only in those cells

that had completely exited the light-organ pores was *hbtR* expression detected (Fig. 5B

and Movie S1). Within an hour of expulsion into seawater, ~25% of expelled *V. fischeri*

cells were expressing *hbtR* (Fig. 5C). As expected, transcripts of the quorum signaling-

induced gene *litR* were detected within *V. fischeri* cells inside the *E. scolopes* light-

organ crypts (Fig. S3); thus, it is unlikely that HCR was unable to detect *hbtR* transcripts

in colonizing bacteria due to failure of the reagents to penetrate the light organ.

Discussion

While much research has gone into understanding the mechanisms necessary for

colonization of the *E. scolopes* light organ by *V. fischeri* (reviewed most recently in

(37)), there has been little investigation into how the ~95% of symbiotic bacteria

expelled from the light organ at dawn each day transition back into life in seawater. In

this work we present HbtR, a heterofunctional homolog of the *V. cholerae* virulence

regulator TcpP that represses the quorum-signaling regulator LitR. While it is unclear at

this time whether HbtR represses *litR* expression directly or through some intermediate

regulator(s), the RNA-seq data indicate HbtR does not act through a step in the

quorum-signaling pathway upstream of LitR (Table S1). Through the discovery of

additional functions regulated by LitR, namely chemotaxis, motility, and EPS production,

we have begun to build a picture in which LitR aids the transition by *V. fischeri* into a

colonization lifestyle and, upon expulsion from the light organ, HbtR reverses that

process to help transition back into planktonic life (Fig. 6). This is a markedly different

role for HbtR from that of TcpP.

The regulatory mechanisms controlling *hbtRC* expression, as elucidated thus far,

are similar to the *tcpPH* regulatory pathway in *V. cholerae* only in that both operons are

regulated by the global regulator Crp, albeit in the opposite manner. Notably, the *hapR*

homolog in *V. fischeri*, *litR*, is itself repressed by HbtR (Tables 1, S1, S2), a reversal of

roles from those in *V. cholerae*. Considering the distinct regulatory pathways for HbtR

and TcpP, as well as the low percent identity (26.8%) of the two protein sequences, it

appears, evolutionarily, that either progenitors of *tcpPH* and *hbtRC* were acquired

independently after separation of the two species' ancestors, or the parent operon

evolved in dramatically different fashions as the two species evolved to inhabit disparate

environmental niches. Determining which of these histories is the more likely is beyond

the scope of this study, but in either case it is clear the two homologs have been

recruited into specialized roles befitting the two species' distinct lifestyles, i.e.,

pathogenesis versus beneficial symbiosis within animal hosts.

The data presented here suggest HbtR acts as a first-wave responder as *V.*

fischeri bacteria exit the light organ and re-enter the seawater (Fig. 5), a significant

change in environmental conditions and an imperative for survival. This role could

explain some of the discrepancies observed between the RNA-seq results in the

present study and those in Thompson 2017 (19). For example, the Thompson data

indicate that in addition to *hbtR* and *hbtC*, the luminescence genes *luxR*, *luxI*, and

luxCDABEG were also highly upregulated in cells recently expelled from the light

organs compared to planktonic cells, while *litR*, which activates the *lux* genes, was not

differentially regulated between the two cell populations. Meanwhile, the results of this

study indicate that HbtR represses LitR and consequently causes the downregulation of

lux and other downstream quorum-signaling genes (Tables 1, S1, S2). Similarly, the

gene most highly upregulated by HbtR in the RNA-seq data presented here was

VF_2042 (Table S1), one of the four MCP genes repressed by LitR (Table 2); in the

earlier study, VF_2042 was downregulated in cells released from the light organs (19).

These results are consistent with a model in which HbtR becomes active once *V.*

fischeri exits the light organ, with some of the earliest effects seen in its repression of

litR. In the earlier study, it may have been that in the ~25 minutes between initiation of

light-organ expulsion and harvesting the *V. fischeri* RNA, HbtR had become activated

(Fig. 5) and was already repressing *litR* expression, but the downstream effects (e.g.,

on VF_2042 and *lux* gene expression) were not yet apparent.

Previous studies have also demonstrated a repression of several flagellin genes

by AinS and increased motility in a *litR* mutant (31). Here we show that LitR represses

numerous flagellin, motor, and chemotaxis genes, including at least two MCP genes

involved in chemotaxis toward (GlcNAc)$_2$, an *N*-acetylated sugar reported to be an

important chemoattractant during light-organ colonization (12). Previous work indicated

that a *litR* mutant colonized *E. scolopes* to a lower level than wild-type *V. fischeri* (31),

which the authors suggested was due to an initiation delay due to hypermotility.

However, in the study in which LitR was first described (4), a *litR* mutant had an

advantage in initiating light-organ colonization over wild-type *V. fischeri.* Based on work

presented here, including a demonstration that MCP genes repressed by LitR are

involved in light-organ colonization (Fig. 2), we posit that the *litR*-mutant colonization

benefit (4) may have resulted from the increase in both motility and chemotaxis into the

light organ by that mutant. We hypothesize that the lower level of colonization observed

for the *litR* mutant in single-strain experiments (31) may have been due to dysregulation

of a separate symbiotic process, perhaps extracellular polysaccharide production.

LitR appears to activate, and HbtR to thus repress, a large locus of genes

(VF_0157–80) involved in extracellular polysaccharide production (Tables 1, S1, S2).

Fidopiastis et al. (4) noted that *litR*[-] colonies are less opaque than wild-type *V. fischeri*

and suggested this phenotype may be due to a difference in the extracellular

polysaccharides present in each strain. The nature of polysaccharide affected by

VF_0157–80 is not fully established, yet it seems most likely to be an EPS, or "slime

layer," due to the ease with which it separates from the bacterial cell during

centrifugation (Fig. 3A). While lipopolysaccharide and the *syp* polysaccharide have

been implicated in the initiation of host colonization by *V. fischeri* (38–40), little has been

determined regarding the possible involvement of extracellular polysaccharides in

colonization persistence. *V. fischeri*-colonized crypts, despite having no goblet cells

(Nyholm, personal communication), have been shown to stain alcian blue-positive (41),

indicating that the presumptive EPS made by VF_0157–80 may be a relevant factor in

E. scolopes-V. fischeri symbiosis. This aspect of the symbiotic relationship could have

broader implications for host-microbe interactions in general.

Materials and Methods

Bacterial strains and growth conditions. Bacterial strains and plasmids used in this

study are summarized in Table S3. *E. coli* and *V. cholerae* were grown in lysogeny

broth (LB; (42), (19)) at 37°C; *V. fischeri* was grown in LB-salt (LBS; (43)) at 28°C.

Overnight cultures were inoculated with single colonies from freshly streaked -80°C

stocks. Liquid cultures were shaken at 225 rpm. Unless otherwise noted, 15 g of agar

were added per liter for solid media. Antibiotics were added to overnight cultures, where

applicable, at the following concentrations: kanamycin (Km), 50 µg/mL; chloramphenicol

(Cm), 5 µg/mL (*V. fischeri*) or 25 µg/mL (*E. coli*). Experimental cultures were grown in

either seawater-tryptone (SWT) medium (44) or minimal-salts medium (MSM; per L: 8.8

g NaCl, 6.2 g $MgSO_4 \cdot 7H_2O$, 0.74 g $CaCl_2 \cdot 2H_2O$, 0.0204 g H_2NaPO_4, 0.37 g KCl, 8.66 g

PIPES disodium salt, 0.65 mg $FeSO_4 \cdot 7H_2O$; pH 7.5) supplemented with 5 mg/L

casamino acids and a carbon source as indicated. The results of all experiments are

reported as the means of three (or more, as indicated) biological replicates ± 1 standard

deviation (SD). Statistical analysis was performed using Welch's t-test.

Plasmid and mutant construction. Primers used to construct the deletion and

expression vectors in this study are listed in Table S4. In-frame deletion of genes from

the *V. fischeri* genome was performed as previously described (45, 46). Counter-

selection to remove the target gene and pSMV3 was performed on LB-sucrose (per L:

2.5 g NaCl, 10 g Bacto Tryptone, 5 g yeast extract, 100 g sucrose) for two days at room

temperature. Expression vectors were constructed by amplifying and ligating the target

gene into the multiple cloning site of pVSV105 (47). Inducible expression of *hbtRC* was

achieved by ligating *lacI^q* from pAKD601 to *hbtRC* cloned from the *V. fischeri* genome

and inserting the fusion into the multiple cloning site of pVSV105. Overexpression of

hbtRC brought the level of expression closer to that found upon expulsion from juvenile

squid (19), which was higher than we measured for wild-type *V. fischeri* in culture

conditions. Genomic insertion of *gfp* or *rfp* under control of the *lac* promoter at the

attTn7 site was performed using a mini-Tn7 vector as described (48). Previously

constructed Δ*hbtRC* (19) was used to recapitulate conditions for follow-up on previous

experiments (RNA-seq and squid co-colonization); a newly derived Δ*hbtRC* strain was

used for all other experiments so it would have the same parent wild-type *V. fischeri*

stock as other mutants created in this study.

Generation and analysis of RNAseq libraries. RNA extraction and RNAseq were

performed as previously described (19). Briefly, RNA was stabilized with RNAprotect

Bacterial Reagent (Qiagen) and extracted with an RNeasy Mini Kit (Qiagen) from 500

μL of cultures grown to mid-log phase in SWT with 1.75 mM IPTG. RNA was extracted

from biological triplicates for each strain. Contaminating DNA was degraded by

treatment with TURBO DNase (Invitrogen). Ribosomal reduction, strand-specific library

preparation, and paired-end 50-bp sequence analysis on an Illumina HiSeq 2500 on

high-output mode were performed at the University of Minnesota Genomics Center.

Sequences were processed on the open-source Galaxy server (usegalaxy.org; (49))

using the following workflow (default settings used unless otherwise indicated): reads

were trimmed with Trimmomatic (maximum mismatch:1, LEADING:3, TRAILING:3,

SLIDINGWINDOW:4,20, MINLEN:35), mapped to the *V. fischeri* chromosomes and

plasmid (accession nos. NC_006840.2, NC_006841.2, and NC_006842.2) using

Bowtie2 (-un-con, --sensitive), and counted with featureCounts (-p

disabled, -t=gene, -g=locus_tag, -Q=10, -s); differential expression was analyzed with

DESeq2 (outliers filtering and independent filtering turned on). Between 11.7 million and

12.7 million paired reads were mapped to the *V. fischeri* genome per biological

replicate. Raw and processed read files have been deposited into the NCBI Gene

Expression Omnibus server (GSE151621).

Reverse transcription and quantitative PCR. RNA from bacterial cultures was

extracted as described above and treated with TURBO DNase (Invitrogen). cDNA was

synthesized from DNase-treated RNA using SMART MMLV Reverse Transcriptase

(Clontech) and Random Hexamer Primer (Thermo Scientific). Gene expression was

measured by qPCR performed with LightCycler 480 SYBR Green I Master Mix (Roche)

under the following conditions: 95°C for 5 min; 45 cycles of 95°C for 10 sec, 60°C for 20

sec, and 72°C for 20 sec; melting curve acquisition from 65°C to 97°C. qPCR primer

pairs were designed for these conditions and confirmed to have efficiencies between 90

and 110%. Cycle thresholds (Ct) for each sample were normalized to those of the

reference gene *polA* (ΔCt = target gene-*polA*; *polA* expression primers provided

courtesy of Silvia Moriano-Gutierrez). $\Delta\Delta$Ct values were calculated by subtracting the

average ΔCt for the parent strain, at higher pH or lower OD_{600} when appropriate; fold

changes were calculated as $2^{(-\Delta\Delta Ct)}$.

Growth and luminescence curves. Overnight LBS cultures of each strain were

pelleted, washed once, and resuspended in 1 mL of the intended growth medium.

Luminescence growth curves were performed in 20 mL SWT in flasks shaking at 225

rpm; periodic 1 mL and 300 µL aliquots were taken for luminescence and OD_{600}

readings on a luminometer (Turner Designs) and Tecan Genios plate reader,

respectively. Growth curves were performed in 1.2 mL growth medium and monitored in

the Tecan Genios plate reader with continuous shaking at high speed.

Squid colonization assays. Juvenile squid colonization experiments were performed

as described previously (50), with juvenile squid exposed to *V. fischeri* strains at 1,000–

6,000 CFU/mL either for 3 h or overnight, as indicated, before being placed in fresh

sterile seawater until euthanizing by freezing. Colonization competition experiments

were performed by exposing juvenile squid to a 1:1 inoculum of each strain.

***In situ* hybridization chain reaction.** HCR (51) was performed on squid colonized with

wild-type *V. fischeri* carrying a *gfp*-labeled plasmid and on *V. fischeri* cells 1 h after

being expelled from squid light organs as previously described (52–54). Briefly, *E.*

scolopes juveniles and expelled *V. fischeri* cells were fixed in 4% paraformaldehyde in

marine phosphate-buffered saline either before or after a light cue stimulated bacterial

expulsion. HCR was performed on dissected light organs (52) and expelled *V. fischeri*

cells affixed to gelatin-coated slides (adapted from (53)) using HCR version 3.0

chemistry (54). Ten probes targeting *hbtR* mRNA and eleven probes targeting *litR*

mRNA (Table 3) were amplified with Alexa Fluor 546-labeled hairpins (Molecular

Instruments). Light organs were then counterstained overnight with a 1:750 dilution of

4′,6-diamidino-2-phenylindole (DAPI; Thermo Fisher Scientific) in 5x SSC-Tween before

mounting on slides with Vectashield (Vector Laboratories) and overlaid with a coverslip

(#1.5, Fisherbrand, Fisher Scientific). Imaging was done on an upright Zeiss LSM 710

laser-scanning confocal microscope at the University of Hawaiʻi Kewalo Marine

Laboratory; images were analyzed using FIJI (ImageJ) (55).

Motility and chemotaxis assays. Soft-agar motility assays were performed as

previously described (56). Briefly, the equivalent of 10 µL of a bacterial culture grown to

an OD_{600} of 0.5 in SWT were spotted onto minimal-medium plates containing 0.25%

agar and, when appropriate, 1 mM chemoattractant. Migration zones were measured

after 18–24 h of static incubation at 28°C.

Alcian blue detection of extracellular polysaccharides. EPS was detected by

staining with alcian blue essentially as previously described (57). Single colonies from

freshly streaked -80°C stocks were used to inoculate MSM supplemented with 6.5 mM

Neu5Ac and 0.05% (wt/vol) casamino acids and grown with shaking 18–22 h. EPS was

separated from cells by centrifugation of the equivalent of 2.5 mL at OD_{600}=1.0 for 15

min at 12,000 xg and 4°C. 250 µL of this supernatant was mixed with 1 mL alcian-blue

solution (per L: 0.5 g alcian blue, 30 mL glacial acetic acid, pH 2.5) and rocked for 1 h at

room temperature, then centrifuged for 10 min at 10,000 rpm and 4°C. Pellets were

resuspended in 1 mL 100% ethanol, then centrifuged for 10 min at 10,000 xg and 4°C.

Pellets were solubilized in 500 µL SDS (per L: 100 g sodium dodecyl sulfate, 50 mM

sodium acetate, pH 5.8), and the absorbance was read at OD_{620}.

Acknowledgments

The authors thank Silvia Moriano-Gutierrez and Alexandrea Duscher for advice

regarding RNA-seq, Elliott McCloskey and Roxane Gaedeke for help with experiment

execution, the University of Minnesota Genomics Center for RNA processing and next-

generation sequencing, and the Ruby and McFall-Ngai laboratories for their insights.

The research was supported by NIH grants R37 AI50661 (Margaret McFall-Ngai and

E.G.R.), R01 OD11024, and R01 GM135254 (E.G.R. and M.M.-N.) and NSF INSPIRE

Grant MCB1608744 (Eva Kanso and Scott Fraser, University of Southern California).

Acquisition of the Leica TCS SP8 X confocal was supported by NSF DBI 1828262

(Marilyn Dunlap, E.G.R., and M.M.-N.), and microscopy was performed in the MICRO

facility, supported by COBRE P20 GM125508 (M.M.-N. and E.G.R.).

References

1.   Eberhard A, Burlingame AL, Eberhard C, Kenyon GL, Nealson KH, Oppenheimer
NJ. 1981. Structural identification of autoinducer of *Photobacterium fischeri* luciferase.
Biochemistry 20:2444–2449.

2.   Kuo A, Blough NV, Dunlap PV. 1994. Multiple N-acyl-L-homoserine lactone
autoinducers of luminescence in the marine symbiotic bacterium *Vibrio fischeri*. J
Bacteriol 176:7558–7565.

3.   Eberhard A. 1972. Inhibition and activation of bacterial luciferase synthesis. J
Bacteriol 109:1101–1105.

4.   Fidopiastis PM, Miyamoto CM, Jobling MG, Meighen EA, Ruby EG. 2002. LitR, a
new transcriptional activator in *Vibrio fischeri*, regulates luminescence and symbiotic
light organ colonization. Mol Microbiol 45:131–143.

5.   Engebrecht J, Silverman M. 1984. Identification of genes and gene products
necessary for bacterial bioluminescence. Proc Natl Acad Sci U S A 81:4154–4158.

6.   Verma SC, Miyashiro T. 2013. Quorum Sensing in the squid-vibrio symbiosis. Int
J Mol Sci 14:16386–16401.

7.   Visick KL, Foster J, Doino J, McFall-Ngai M, Ruby EG. 2000. *Vibrio fischeri lux*
genes play an important role in colonization and development of the host light organ. J
Bacteriol 182:4578–4586.

8.   Koch EJ, Miyashiro T, McFall-Ngai MJ, Ruby EG. 2014. Features governing symbiont persistence in the squid-vibrio association. Mol Ecol 23:1624–1634.

9.   Moriano-Gutierrez S, Koch EJ, Bussan H, Romano K, Belcaid M, Rey FE, Ruby EG, McFall-Ngai MJ. 2019. Critical symbiont signals drive both local and systemic changes in diel and developmental host gene expression. Proc Natl Acad Sci U S A 116:7990–7999.

10.   Wei SL, Young RE. 1989. Development of symbiotic bacterial bioluminescence in a nearshore cephalopod, *Euprymna scolopes*. Mar Biol 103:541–546.

11.   Graf J, Dunlap PV, Ruby EG. 1994. Effect of transposon-induced motility mutations on colonization of the host light organ by *Vibrio fischeri*. J Bacteriol 176:6986–6991.

12.   Mandel MJ, Schaefer AL, Brennan CA, Heath-Heckman EAC, Deloney-Marino CR, McFall-Ngai MJ, Ruby EG. 2012. Squid-derived chitin oligosaccharides are a chemotactic signal during colonization by *Vibrio fischeri*. Appl Environ Microbiol 78:4620–4626.

13.   Ruby EG, Asato LM. 1993. Growth and flagellation of *Vibrio fischeri* during initiation of the sepiolid squid light organ symbiosis. Arch Microbiol 159:160–167.

14.   Graf J, Ruby EG. 1998. Host-derived amino acids support the proliferation of symbiotic bacteria. Proc Natl Acad Sci U S A 95:1818–1822.

15.   Jones BW, Nishiguchi MK. 2004. Counterillumination in the Hawaiian bobtail squid, *Euprymna scolopes* Berry (Mollusca: Cephalopoda). Mar Biol 144:1151–1155.

16.   Stabb EV, Millikan DS, Millikan DS. 2009. Is the *Vibrio fischeri–Euprymna scolopes* symbiosis a defensive mutualism? *In* Defensive Mutual Microb Symbiosis. CRC Press.

17.   Lee K-H, Ruby EG. 1994. Effect of the squid host on the abundance and distribution of symbiotic *Vibrio fischeri* in nature. Appl Environ Microbiol 60:1565–1571.

18.   Nyholm SV, McFall-Ngai MJ. 1998. Sampling the light-organ microenvironment of *Euprymna scolopes*: description of a population of host cells in association with the bacterial symbiont *Vibrio fischeri*. Biol Bull 195:89–97.

19.   Thompson LR, Nikolakakis K, Pan S, Reed J, Knight R, Ruby EG. 2017. Transcriptional characterization of *Vibrio fischeri* during colonization of juvenile *Euprymna scolopes*. Environ Microbiol 19:1845–1856.

20.   Thomas S, Williams SG, Manning PA. 1995. Regulation of *tcp* genes in classical and El Tor strains of *Vibrio cholerae* O1. Gene 166:43–48.

21.   Häse CC, Mekalanos JJ. 1998. TcpP protein is a positive regulator of virulence gene expression in *Vibrio cholerae*. Proc Natl Acad Sci U S A 95:730–734.

22.   Taylor RK, Miller VL, Furlong DB, Mekalanos JJ. 1987. Use of *phoA* gene fusions
to identify a pilus colonization factor coordinately regulated with cholera toxin. Proc Natl
Acad Sci U S A 84:2833–2837.

23.   Skorupski K, Taylor RK. 1999. A new level in the *Vibrio cholerae* ToxR virulence
cascade: AphA is required for transcriptional activation of the *tcpPH* operon. Mol
Microbiol 31:763–771.

24.   Kovacikova G, Skorupski K. 1999. A *Vibrio cholerae* LysR homolog, AphB,
cooperates with AphA at the *tcpPH* promoter to activate expression of the ToxR
virulence cascade. J Bacteriol 181:4250–4256.

25.   Kovacikova G, Lin W, Skorupski K. 2010. The LysR-type virulence activator
AphB regulates the expression of genes in *Vibrio cholerae* in response to low pH and
anaerobiosis. J Bacteriol 192:4181–4191.

26.   Zhu J, Miller MB, Vance RE, Dziejman M, Bassler BL, Mekalanos JJ. 2002.
Quorum-sensing regulators control virulence gene expression in *Vibrio cholerae*. Proc
Natl Acad Sci U S A 99:3129–3134.

27.   Kovacikova G, Skorupski K. 2002. Regulation of virulence gene expression in
Vibrio cholerae by quorum sensing: HapR functions at the *aphA* promoter. Mol Microbiol
46:1135–1147.

28.   Kovacikova G, Skorupski K. 2001. Overlapping binding sites for the virulence
gene regulators AphA, AphB and cAMP-CRP at the *Vibrio cholerae tcpPH* promoter.
Mol Microbiol 41:393–407.

29.   Larocque RC, Harris JB, Dziejman M, Li X, Khan AI, Faruque ASG, Faruque SM,
Nair GB, Ryan ET, Qadri F, Mekalanos JJ, Calderwood SB. 2005. Transcriptional
profiling of *Vibrio cholerae* recovered directly from patient specimens during early and
late stages of human infection. Infect Immun 73:4488–4493.

30.   Schild S, Tamayo R, Nelson EJ, Qadri F, Calderwood SB, Camilli A. 2007.
Genes induced late in infection increase fitness of *Vibrio cholerae* after release into the
environment. Cell Host Microbe 2:264–277.

31.   Lupp C, Ruby EG. 2005. *Vibrio fischeri* uses two quorum-sensing systems for the
regulation of early and late colonization factors. J Bacteriol 187:3620–3629.

32.   Cao X, Studer SV, Wassarman K, Zhang Y, Ruby EG, Miyashiro T. 2012. The
novel sigma factor-like regulator RpoQ controls luminescence, chitinase activity, and
motility in *Vibrio fischeri*. mBio e00285-11

33.   Dunlap PV, Greenberg EP. 1985. Control of *Vibrio fischeri* luminescence gene
expression in *Escherichia coli* by cyclic AMP and cyclic AMP receptor protein. J
Bacteriol 164:45–50.

34.   Lyell NL, Colton DM, Bose JL, Tumen-Velasquez MP, Kimbrough JH, Stabb EV.
2013. Cyclic AMP receptor protein regulates pheromone-mediated bioluminescence at
multiple levels in *Vibrio fischeri* ES114. J Bacteriol 195:5051–5063.

35.   Murley YM, Carroll PA, Skorupski K, Taylor RK, Calderwood SB. 1999.
Differential transcription of the *tcpPH* operon confers biotype-specific control of the
Vibrio cholerae ToxR virulence regulon. Infect Immun 67:5117–5123.

36.   Krukonis ES, Yu RR, Dirita VJ. 2000. The *Vibrio cholerae* ToxR/TcpP/ToxT
virulence cascade: distinct roles for two membrane-localized transcriptional activators
on a single promoter. Mol Microbiol 38:67–84.

37.   Norsworthy AN, Visick KL. 2013. Gimme shelter: how *Vibrio fischeri* successfully
navigates an animal's multiple environments. Front Microbiol 4:356.

38.   Yip ES, Grublesky BT, Hussa EA, Visick KL. 2005. A novel, conserved cluster of
genes promotes symbiotic colonization and σ^{54}-dependent biofilm formation by *Vibrio
fischeri*: *syp*, a novel symbiosis locus in *Vibrio fischeri*. Mol Microbiol 57:1485–1498.

39.   Post DM, Yu L, Krasity BC, Choudhury B, Mandel MJ, Brennan CA, Ruby EG,
McFall-Ngai MJ, Gibson BW, Apicella MA. 2012. O-antigen and core carbohydrate of
Vibrio fischeri lipopolysaccharide: composition and analysis of their role in *Euprymna
scolopes* light organ colonization. J Biol Chem 287:8515-30.

40.   Schwartzman JA, Lynch JB, Flores Ramos S, Zhou L, Apicella MA, Yew JY,
Ruby EG. 2019. Acidic pH promotes lipopolysaccharide modification and alters
colonization in a bacteria-animal mutualism. Mol Microbiol 112:1326–1338.

41.   Nyholm SV, Deplancke B, Gaskins HR, Apicella MA, McFall-Ngai MJ. 2002.
Roles of *Vibrio fischeri* and nonsymbiotic bacteria in the dynamics of mucus secretion
during symbiont colonization of the *Euprymna scolopes* light organ. Appl Environ
Microbiol 68:5113–5122.

42.   Bertani G. 1951. Studies on lysogenesis. I. The mode of phage liberation by
lysogenic *Escherichia coli*. J Bacteriol 62:293–300.

43.   Stabb EV, Reich KA, Ruby EG. 2001. *Vibrio fischeri* genes *hvnA* and *hvnB*
encode secreted NAD+-glycohydrolases. J Bacteriol 183:309–317.

44.   Boettcher KJ, Ruby EG. 1990. Depressed light emission by symbiotic *Vibrio
fischeri* of the sepiolid squid *Euprymna scolopes*. J Bacteriol 172:3701–3706.

45.   Saltikov CW, Newman DK. 2003. Genetic identification of a respiratory arsenate
reductase. Proc Natl Acad Sci U S A 100:10983–10988.

46.   Bennett BD, Redford KE, Gralnick JA. 2018. Survival of anaerobic Fe^{2+} stress
requires the ClpXP protease. J Bacteriol 200:e00671-17.

47.   Dunn AK, Millikan DS, Adin DM, Bose JL, Stabb EV. 2006. New *rfp-* and pES213-derived tools for analyzing symbiotic *Vibrio fischeri* reveal patterns of infection and *lux* expression *in situ*. Appl Environ Microbiol 72:802–810.

48.   McCann J, Stabb EV, Millikan DS, Ruby EG. 2003. Population dynamics of *Vibrio fischeri* during Infection of *Euprymna scolopes*. Appl Environ Microbiol 69:5928–5934.

49.   Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Cech M, Chilton J, Clements D, Coraor N, Grüning BA, Guerler A, Hillman-Jackson J, Hiltemann S, Jalili V, Rasche H, Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D. 2018. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Res 46:W537–W544.

50.   Bongrand C, Ruby EG. 2019. Achieving a multi-strain symbiosis: strain behavior and infection dynamics. ISME J 13:698-706.

51.   Choi HMT, Chang JY, Trinh LA, Padilla JE, Fraser SE, Pierce NA. 2010. Programmable *in situ* amplification for multiplexed imaging of mRNA expression. Nat Biotechnol 28:1208–1212.

52.   Nikolakakis K, Lehnert E, McFall-Ngai MJ, Ruby EG. 2015. Use of hybridization chain reaction-fluorescent *in situ* hybridization to track gene expression by both partners during initiation of symbiosis. Appl Environ Microbiol 81:4728–4735.

53.   Choi HMT, Calvert CR, Husain N, Huss D, Barsi JC, Deverman BE, Hunter RC, Kato M, Lee SM, Abelin ACT, Rosenthal AZ, Akbari OS, Li Y, Hay BA, Sternberg PW, Patterson PH, Davidson EH, Mazmanian SK, Prober DA, van de Rijn M, Leadbetter JR, Newman DK, Readhead C, Bronner ME, Wold B, Lansford R, Sauka-Spengler T, Fraser SE, Pierce NA. 2016. Mapping a multiplexed zoo of mRNA expression. Dev Camb Engl 143:3632–3637.

54.   Choi HMT, Schwarzkopf M, Fornace ME, Acharya A, Artavanis G, Stegmaier J, Cunha A, Pierce NA. 2018. Third-generation *in situ* hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. Development 145:dev165753.

55.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. Nat Methods 9:676–682.

56.   DeLoney-Marino CR. 2013. Observing chemotaxis in *Vibrio fischeri* using soft agar assays in an undergraduate microbiology laboratory. J Microbiol Biol Educ JMBE 14:271–272.

57.   Plante C. 2000. Role of bacterial exopolymeric capsules in protection from deposit-feeder digestion. Aquat Microb Ecol 21:211–219.

 FIGURE LEGENDS

FIG 1 Luminescence and growth of wild-type *V. fischeri* and Δ*hbtRC* strains. The rates

of (A) light production and (B) growth in SWT were measured for wild-type *V. fischeri*

(circles) and Δ*hbtRC* (squares) carrying empty pVSV105 (open symbols) or

pVSV105::*hbtRC* (closed symbols). Results represent the means of three biological

replicates ± one standard deviation. RLU, relative light units; Abs, absorbance.

FIG 2 Light-organ colonization competition between wild-type *V. fischeri* and a

chemotaxis mutant. Ratios of wild-type *V. fischeri* (WT) carrying a GFP-encoding

plasmid (pVSV102) vs. unlabeled ΔVF_1133 ΔVF_2042 ΔVF_A0246 ΔVF_A0389

("ΔΔΔΔ") (circles) or unlabeled wild-type *V. fischeri* vs. ΔΔΔΔ carrying pVSV102

(squares) colonizing juvenile *E. scolopes* light organs were measured after 3 h of

exposure to the inoculum followed by 21 h of incubation in sterile seawater. RCI,

relative competitive index of bacterial strains. Each point represents one animal. Empty

symbols represent the limit of detection for light organs in which only the wild-type strain

was found. Error bars represent ± one standard deviation.

FIG 3 EPS production in wild-type *V. fischeri*, Δ*litR*, and ΔVF_0157–80 strains. EPS in

supernatants of 24-h cultures grown in 0.05% casamino acids and 6.5 mM Neu5Ac was

stained with alcian blue for wild-type *V. fischeri* (circles), Δ*litR* (squares), and

ΔVF_0157–80 (triangles) carrying empty pVSV105 (open symbols), pVSV105::*hbtRC*

(half-filled symbols), or pVSV105::*litR* (closed symbols). Error bars represent ± one

standard deviation. Abs, absorbance.*, $p<0.05$; **, $p<0.005$; ***, $p<0.001$; n.s., not

statistically significant.

FIG 4 Light-organ colonization competition between wild-type *V. fischeri* and Δ*hbtRC*

strains. (A) Ratios of wild-type *V. fischeri* carrying a GFP-encoding plasmid (pVSV102)

vs. Δ*hbtRC* carrying an RFP-encoding plasmid (pVSV208) (circles, squares, and

triangles) or wild-type *V. fischeri* (WT) carrying pVSV208 vs. Δ*hbtRC* carrying pVSV102

(diamonds) colonizing juvenile *E. scolopes* light organs were measured after 1 to 10 d.

(B) Ratios of wild-type *V. fischeri* with chromosomal *gfp* vs. Δ*hbtRC* with chromosomal

rfp (squares) or wild-type *V. fischeri* with chromosomal *rfp* vs. Δ*hbtRC* with

chromosomal *gfp* (circles) colonizing juvenile *E. scolopes* light organs were measured

after 1 d. RCI, relative competitive index of bacterial strains. Each point represents one

animal. Empty symbols represent the limit of detection for light organs in which only the

gfp-carrying strain was found. Error bars represent ± one standard deviation. n.s., not

statistically significant.

FIG 5 Expression of *hbtR* in *V. fischeri* during expulsion from *E. scolopes* light organs.

Transcripts of *hbtR* (magenta, arrowheads) were detected using *in situ* HCR in *V.*

fischeri cells (green) during expulsion from *E. scolopes* juvenile light organ crypts. (A) *V.*

fischeri symbionts originate in the crypt, pass through the migration path (mp), and exit

through the external surface pore; (A) merged image, (A') GFP channel, (A") *hbtR*

transcript channel (Alexa 546). (B) *V. fischeri* cells (green) are expelled through the

external surface pore (asterisk) into seawater 1 h after a dawn light cue; (B) a single

optical slice of the superficial layer above the exit point from host and *V. fischeri* cells

outside of the pore expressing *hbtR*; (B'), the corresponding merged composite stack

(video in Movie S1). (C) One hour after expulsion into ambient seawater, vented cells

express *hbtR* (magenta); enlargement is of the area within the dashed square. (C')

Percentage and 95% confidence intervals of vented cells expressing *hbtR* calculated

from four samples of ventate on slides, from two experiments. Blue, *E. scolopes* nuclei

(DAPI).

FIG 6 Model proposing regulation by HbtR and LitR in *V. fischeri* during colonization of

and expulsion from the *E. scolopes* light organ. Upon colonization, LitR regulates

relevant physiological phenotypes, including activation of luminescence and EPS

production and repression of motility and chemotaxis. HbtR causes the repression of

litR expression upon expulsion of *V. fischeri* from the light organ, reversing this process

and leading to the upregulation of motility and chemotaxis and the downregulation of

luminescence and EPS production. Gray text, downregulated genes and phenotypes;

blue text, upregulated genes and phenotypes.

SUPPLEMENTAL MATERIAL LEGENDS

FIG S1 Luminescence of wild-type *V. fischeri* and Δ*hbtRC* strains during symbiosis.

Luminescence was measured in juvenile squid 24 h or 48 h after colonization with either

wild-type *V. fischeri* (●) or Δ*hbtRC* (■). Each point represents one animal. Mean values

indicated; no significant differences were noted between strains at either time-point.

RLU, relative light units.

FIG S2 Growth of wild-type *V. fischeri* and chemotaxis mutants on *N*-acetylated sugars.

(A) The rates of growth in MSM supplemented with 0.05% casamino acids and 6.5 mM

Neu5Ac were measured for wild-type *V. fischeri* (●), ΔΔΔΔ (■), and ΔVF_1133

ΔVF_A0246 (▲). (B) The rates of growth in MSM supplemented with 0.05% casamino

acids and 6.5 mM GlcNAc were measured for wild-type *V. fischeri* (●) and ΔΔΔΔ (■).

Results represent means of three biological replicates ± one standard deviation. Abs,

absorbance.

FIG S3 Expression of *litR* in *V. fischeri* when host-associated within the *E. scolopes* light

organ. Transcripts of *litR* (green) were detected using *in situ* HCR in *V. fischeri* cells

colonizing *E. scolopes* crypts (C) after traversing the tissues of the migration path (MP).

Blue, *E. scolopes* nuclei (DAPI); green, *litR* transcripts (Alexa 546). Bar 15 μm.

TABLE S1 RNA-seq data file Expression-level counts of all orfs in the *V. fischeri*

genome. Comparison of mapped read counts for Δ*hbtRC* and *hbtRC* (induced)

TABLE S2 RT-qPCR results

TABLE S3 Bacterial strains and plasmids used in this work

TABLE S4 Primers and probes used in this work

MOVIE S1 Expression of *hbtR* in *V. fischeri* cells during their expulsion from an *E.*

scolopes light organ. *hbtR* transcripts were detected using *in situ* HCR in *V. fischeri*

cells expelled from the external surface pore of the light organ 1 h after a dawn light

cue. Merged stack of images depicted in Fig 5B'. Blue, *E. scolopes* nuclei (DAPI);

green, *gfp*-labeled *V. fischeri*; magenta, *hbtR* transcripts (Alexa 546).

Table 1. RNA-seq results for genes analyzed in this study[a]

Quorum-sensing genes

Locus	Gene	Product description	log2(FC)[b]	*p*-adj[c]
VF_2177	*litR*	quorum sensing transcriptional regulator LitR	-1.09	1.30E-25
VF_A0919	*luxE*	long-chain-fatty-acid ligase	-0.56	0.0002
VF_A0920	*luxB*	luciferase beta chain	-0.39	0.007
VF_A0921	*luxA*	luciferase alpha chain	-0.42	0.001
VF_A0922	*luxD*	acyl transferase	-0.36	0.027
VF_A0923	*luxC*	acyl-CoA reductase	-0.58	0.0005
VF_A0924	*luxI*	3-oxo-C6-HSL autoinducer synthesis protein	-0.67	0.0005
VF_A0925	*luxR*	LuxR family transcriptional regulator	-1.29	1.78E-27
VF_A1015	*rpoQ*	sigma-Q factor RpoQ, quorum-sensing regulated RpoS-like sigma subunit	-1.25	2.01E-14
VF_A1058	*qsrP*	LuxR family transcriptional regulator	-1.38	1.41E-05

Extracellular polysaccharide genes

Locus	Gene	Product description	log2(FC)	*p*-adj
VF_0157	*wbfB*	WbfB protein	-1.36	6.75E-21
VF_0158	VF_0158	hypothetical protein	-1.61	5.16E-23
VF_0160	*wbfD*	WbfD protein	-1.73	4.19E-14
VF_0161	VF_0161	hypothetical protein	-1.23	4.30E-28
VF_0162	*gfcE*	exopolysaccharide export protein	-1.98	8.73E-43
VF_0163	VF_0163	hypothetical protein	-1.26	2.53E-17
VF_0164	*etp*	phosphotyrosine-protein phosphatase	-2.08	1.33E-37
VF_0165	*wzc*	protein-tyrosine kinase, chain length regulator in capsular polysaccharide biosynthesis	-1.81	2.47E-38
VF_0166	*rffG*	dTDP-glucose 4,6-dehydratase	-1.98	1.01E-45
VF_0167	*rffH*	glucose-1-phosphate thymidylyltransferase	-1.65	1.01E-37
VF_0168	*rfbC*	dTDP-4-deoxyrhamnose-3,5-epimerase	-1.80	2.75E-38
VF_0169	*rmlB*	dTDP-glucose-4,6-dehydratase	-1.64	3.03E-31
VF_0170	*rfbX*	polisoprenol-linked O-antigen transporter	-1.72	3.03E-24

VF_0171	VF_0171	hypothetical protein	-1.14	1.68E-09
VF_0172	VF_0172	O-acetyltransferase	-1.64	3.26E-53
VF_0173	VF_0173	hypothetical protein	-1.94	1.66E-33
VF_0174	VF_0174	beta-D-GlcNAc beta-1,3-galactosyltransferase	-1.72	2.02E-36
VF_0175	VF_0175	glycosyltransferase	-1.50	1.16E-42
VF_0176	VF_0176	3-deoxy-8-phosphooctulonate synthase	-1.34	5.18E-30
VF_0177	VF_0177	3-deoxy-manno-octulosonate-8-phosphatase	-1.56	1.57E-56
VF_0178	VF_0178	3-deoxy-manno-octulosonate cytidylyltransferase	-1.60	1.91E-60
VF_0179	*kpsF*	arabinose-5-phosphate isomerase	-1.80	3.50E-39
VF_0180	*rfe*	UDP-GlcNAc:undecaprenylphosphate GlcNAc-1-phosphate transferase	-1.78	2.39E-71

Chemotaxis and motility genes

Locus	Gene	Product description	log2(FC)	*p*-adj
VF_1133	VF_1133	methyl-accepting chemotaxis protein	1.55	4.08E-41
VF_1864	*flaC*	flagellin	1.03	1.80E-16
VF_2042	VF_2042	methyl-accepting chemotaxis protein	2.78	2.11E-87
VF_2079	*flaF*	flagellin	0.93	6.75E-16
VF_A0246	VF_A0246	methyl-accepting chemotaxis protein	1.04	8.31E-12
VF_A0389	VF_A0389	methyl-accepting chemotaxis protein	1.20	1.34E-09

[a] An abbreviated subset of data in Table S1
[b] Log 2 fold change (Δ*hbtRC*+*hbtRC* / Δ*hbtRC*)
[c] *p*-adj, Benjamini-Hochberg adjustment of Wald test *p*-value

Table 2. RT-qPCR results[a]

	Wild-type *V. fischeri*			ΔlitR			
Gene	**ΔCt[b] (gene-*polA*)**	**SD**		**ΔCt (gene-*polA*)**	**SD**	***p*-value[c]**	**Fold change**
VF1133	3.01	0.03		1.12	0.07	0.001	3.70
VF2042	4.64	0.15		-1.73	0.52	0.002	86.13
VFA0246	3.87	0.06		-1.47	0.12	0.000	40.32
VFA0389	0.52	0.15		-0.81	0.18	0.002	2.52
cheW	-1.55	0.05		-2.53	0.24	0.019	1.99
motA1	0.95	0.23		-0.16	0.30	0.015	2.17
flaC	-1.58	0.22		-4.17	0.51	0.015	6.27
flaF	1.31	0.14		-1.44	0.75	0.024	7.41
rpoN	-0.01	0.14		0.14	0.11	0.233	0.90
flrA	-0.74	0.04		-0.48	0.12	0.071	0.84
flrB	0.85	0.13		1.38	0.63	0.293	0.74
fliA	-1.37	0.11		-1.81	0.27	0.119	1.38
flgM	-0.53	0.14		-1.35	0.31	0.053	1.80
wbfB	1.56	0.12		3.72	0.24	0.005	0.23
etp	0.38	0.09		3.42	0.08	< 0.0001	0.12
wzc	0.35	0.20		3.46	0.17	0.000	0.12

	Δ*hbtRC* + pVSV105			Δ*hbtRC* + pVSV105::*hbtRC*			
Gene	**ΔCt (gene-*polA*)**	**SD**		**ΔCt (gene-*polA*)**	**SD**	***p*-value**	**Fold change**
VF_2042	4.53	0.37		2.10	0.06	0.008	5.39
etp	-1.23	0.08		-0.26	0.08	0.0007	0.51

	Low Cell Density				High Cell Density			
Strain	**ΔCt (*hbtRC-polA*)**	**SD**	***p*-value**	**Fold change**	**ΔCt (*hbtRC-polA*)**	**SD**	***p*-value[d]**	**Fold change[d]**
Wild-type *V. fischeri*	8.08	0.12	—	—	13.07	0.35	0.002	0.03
Δ*litR*	8.19	0.58	0.775	0.97	13.02	0.69	0.007	0.04
Δ*crp*	10.06	0.68	0.038	0.27	12.29	0.12	< 0.0001	0.05

	Low Cell Density				High Cell Density			
Condition	**ΔCt (*hbtRC-polA*)**	**SD**	***p*-value**	**Fold change**	**ΔCt (*hbtRC-polA*)**	**SD**	***p*-value[e]**	**Fold change[e]**
SWT Alone	6.00	0.49	—	—	10.04	0.53	0.002	0.06
+ C8-HSL	5.72	0.13	0.473	1.25	10.60	0.87	0.001	0.06
+ 3-oxo-C6-HSL	5.71	0.39	0.432	1.22	10.16	0.26	0.004	0.05

Table 3. Soft-agar migration zone sizes

Strain	GlcNAc			(GlcNAc)$_2$			Neu5Ac		
	Zone size (mm)[a]	SD	*p*-value[b]	Zone size (mm)	SD	*p*-value	Zone size (mm)	SD	*p*-value
Wild-type *V. fischeri*	25.7	0.6	—	21.7	0.6	—	24	1	—
ΔVF_1133	18.0	0.0	< 0.0001	14.0	0.0	< 0.0001	18	1	0.002
ΔVF_2042	26.3	0.6	0.230	21.3	0.6	0.519	25	1.7	0.450
ΔVF_A0246	19.7	0.6	0.0002	16.3	0.6	0.0003	19.7	0.6	0.007
ΔVF_A0389	26.0	1.0	0.651	22.0	1.7	0.782	24	0	0.651
ΔVF_1133 ΔVF_A0246	ND[c]	ND	ND	ND	ND	ND	17.7	0.6	0.003
ΔΔΔΔ	19.3	2.1	0.037	15.3	1.2	0.014	23.7	3.2	0.880

Strain	Glucose			No chemoattractant		
	Zone size (mm)	SD	*p*-value	Zone size (mm)	SD	*p*-value
Wild-type *V. fischeri*	18.2	0.8	—	17.0	0.0	—
ΔΔΔΔ	19.2	2.9	0.455	16.7	0.6	0.230

Strain	GlcNAc			No chemoattractant		
	Zone size (mm)	SD	*p*-value[d]	Zone size (mm)	SD	*p*-value[d]
Wild-type *V. fischeri* + empty vector	10.3	1.2	—	6.0	1.0	—
Wild-type *V. fischeri* + hbtRC	13.7	1.2	0.024	9.0	1.0	0.021
Wild-type *V. fischeri* + litR	5.3	0.6	0.022	5.3	0.6	0.391
Δ*litR* + empty vector	18.7	0.6	0.008	17.7	1.5	0.002
Δ*litR* + hbtRC	20.7	0.6	0.011	18.3	1.2	0.002
Δ*litR* + litR	6.0	1.0	0.391	7.7	0.6	0.008
Δ*rpoQ* + empty vector	10.7	0.6	0.699	5.0	0.0	0.139
Δ*rpoQ* + hbtRC	15.3	1.2	0.152	9.7	1.5	0.572
Δ*rpoQ* + litR	4.7	0.6	0.230	5.3	0.6	1.000

[a] Average of ≥ three biological replicates.
[b] *p*-value, Welch's t-test as compared to wild-type zones in that chemoattractant.
[c] ND, not done.
[d] *p*-value for wild-type strains, as compared to wild type carrying empty vector; for Δ*litR* and Δ*rpoQ* strains, as compared to wild type carrying the respective vector.

A

RLU

Abs (OD$_{600}$)

B

Abs (OD$_{600}$)

Time (h)