A hybrid cyt *c* maturation system enhances the bioelectrical performance of engineered *Escherichia coli* by improving the rate-limiting step

Author List:

Lin Su [1,2,3], Tatsuya Fukushima [3], Caroline M. Ajo-Franklin [2,3,4,5*]

Author Affiliations:

[1] State Key Laboratory of Bioelectronics, School of Biological Science and Medical Engineering, Southeast University, Nanjing 210018, China

[2] Department of BioSciences, Rice University, Houston, TX 77005, USA

[3] Molecular Foundry, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

[4] Molecular Biophysics and Integrated Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

[5] Institute for Biosciences and Bioengineering, Rice University, Houston, TX 77005, USA

* To whom correspondence should be addressed. Tel: 1-713-348-3497; Fax: 1-713-348-4790; Email: cajo-franklin@rice.edu

ABSTRACT

Bioelectronic devices can use electron flux to enable communication between biotic components

and abiotic electrodes. We have modified *Escherichia coli* to electrically interact with electrodes

by expressing the cytochrome *c* from *Shewanella oneidensis* MR-1. However, we observe

inefficient electrical performance, which we hypothesize is due to the limited compatibility of the

E. coli cytochrome *c* maturation (Ccm) systems with MR-1 cytochrome *c*. Here we test whether

the bioelectronic performance of *E. coli* can be improved by constructing hybrid Ccm systems

containing protein domains from both *E. coli* and *S. oneidensis* MR-1. The hybrid CcmH increased

cytochrome *c* expression by increasing the abundance of CymA 60%, while only slightly changing

the abundance of the other cytochromes *c*. Electrochemical measurements showed that the overall

current from the hybrid *ccm* strain increased 121% relative to the wildtype *ccm* strain, with an

electron flux per cell of 12.3 ± 0.3 fA·cell^{-1}. Additionally, the hybrid *ccm* strain doubled its

electrical response with the addition of exogenous flavin, and quantitative analysis of this

demonstrates CymA is the rate-limiting step in this electron conduit. These results demonstrate

that this hybrid Ccm system can enhance the bioelectrical performance of the cyt *c* expressing *E.*

coli, allowing the construction of more efficient bioelectronic devices.

KEYWORDS

Microbial electrochemistry, microbial fuel cell, bioelectronic sensor, synthetic biology,

cytochrome *c* maturation

1. INTRODUCTION

The flow of electrons can transfer both energy and information between living and non-living systems for bioelectronic and biosensing applications. Extracellular electron transfer (EET) pathways (Shi et al., 2016) can electrically connect living microorganisms with electrochemical cells, enabling applications such as power generation in microbial fuel cells (MFCs) (Gul and Ahmad, 2019; Logan, 2009; Santoro et al., 2017; Slate et al., 2019) and environmental sensing in bioelectronic sensing systems (Chouler et al., 2018; Wang et al., 2013; Zhou et al., 2017). Engineering EET in native electroactive microorganisms, such as *Shewanella oneidensis* MR-1 (Cheng et al., 2020; Min et al., 2017; West et al., 2017), and non-native hosts, such as *Escherichia coli* (Jensen et al., 2016; Sturm-Richter et al., 2015; Thirumurthy and Jones, 2020) or *Pseudomonas putida* (Schmitz et al., 2015), has expanded the range of potential applications (Harris et al., 2017; Su and Ajo-Franklin, 2019). However, a fundamental limitation in these bioelectronic systems is the low rate of EET, which causes these devices to have low power output, low sensing sensitivity, and limited robustness. For example, expressing the EET pathway from *Shewanella oneidensis* MR-1 into engineered *E. coli* only yields ~10% the current produced by MR-1 (TerAvest et al., 2014).

The molecular structure of these EET pathways is increasingly well-understood. The direct EET pathways (Mtr pathway) in *Shewanella oneidensis* MR-1 is a group of multiheme cytochrome *c* (cyt *c*) proteins embedded in the inner membrane (CymA), in the periplasm (Stc, FccA), and outer membrane (MtrC, MtrA) (Clarke et al., 2011; Okamoto et al., 2011; Rosenbaum et al., 2012). We and others (Jensen et al., 2016; Su et al., 2020; TerAvest et al., 2014) have found that expressing the CymA, MtrC, MtrB and MtrA proteins of the Mtr pathway enables *E. coli* to perform EET (**Figure 1A**).

Expressing the Mtr cyt *c* proteins requires the cytochrome *c* maturation (Ccm) system, which

covalently ligates heme *b* groups to the apo-cyt *c* protein to form the holo-cyt *c* (Sanders et al.,

2010; Verissimo et al., 2011). The Ccm systems of α- and β-proteobacteria contains three modules

(**Figure 1B**): the heme *b* translocation module that conveys heme *b* into the periplasm and to the

ligation module, the apo cyt *c* chaperone module that maintains apocyt *c* in the reduced state, and

the ligation module that catalyzes formation of a thioether bond between the heme b and apocyt *c*

(Kranz et al., 2009; Sanders et al., 2010; Thöny-Meyer, 2000). In both *S. oneidensis* and *E. coli*,

CcmA, B, C, D and E form the heme delivery module, and CcmG and DsbD compose the

chaperone module. While *E. coli* uses two proteins, CcmF and CcmH, as the ligation module, this

module in *S. oneidensis* MR-1 includes a third protein - CcmI (**Figure S1**). We recently found that

mutations in the C-terminal domain of CcmH can significantly change the EET efficiency by

changing the stoichiometry of the Mtr cyt *c (Su et al., 2020)*. This finding suggested that optimizing

the heme ligation process by redesigning CcmH of *E. coli* could remodel the stoichiometry of cyt

c and enhance the bioelectrical performance of Mtr-expressing *E. coli*.

Figure 1. A schematic of the Mtr pathway in engineered *E. coli* and the three modules of Ccm system in *E. coli*. (A) The Mtr pathway includes MtrC, MtrB, MtrA and CymA from *Shewanella oneidensis* MR-1, which enable engineered *E. coli* to perform extracellular electron transfer. (B) For cytochrome *c* (cyt *c*) such as MtrC, MtrA and CymA to mature, the Ccm system catalyzes ligation of heme with apo-cyt *c* to form covalent bonds with heme groups. In *E. coli*, the Ccm system is composed of eight different Ccm proteins, CcmA, B, C, D, E, F, G, H, and a DsbD (disulfide bond formation protein). These proteins form three functional modules: heme delivery (green and purple), apo cyt *c* chaperone (red), and ligation (orange).

The role of the C-terminal domain of CcmH in cyt *c* maturation is disputed. Fabianek et al. (Fabianek et al., 1999) and Robertson et al. (Robertson et al., 2008) demonstrated that *E. coli* strains with an incomplete CcmH lacking its C-terminal domain can still mature both the native diheme cyt *c* (NapB) and a heterologous monoheme cyt *c* (Cyt *c* C550). They contend that the C-terminal is not required for maturation because of functional redundancy between the two domains of CcmH. In contrast, Zheng et al. (Zheng et al., 2012) predicted the C-terminal of CcmH is a TRP (tetratricopeptide repeat) -like domain which recognizes apo-cyt *c* and concluded it is essential for cyt *c* expression. Our previous results suggest the C-terminal of CcmH is essential in expressing multi-heme heterologous cyt *c* (Su et al., 2020), agreeing more closely with Zheng et al. (Zheng et al., 2012).

To optimize the heme ligation process for Mtr cyt *c*, here we constructed a hybrid CcmH that uses protein domains from *E. coli* (the N-terminal of CcmH) and *S. oneidensis* MR-1 (the C-

96   terminal of CcmI). We then explored the effects on cyt *c* expression and EET efficiency to test if

97   this hybrid CcmH could boost the bioelectrical performance of Mtr-expressing *E. coli*.

99   2. METHODS

100   2.1   Construction of *E. coli* strains with *ccm* variants

101   We used Phusion DNA polymerase (NEB) for all the PCR amplification (please refer to

102   Supplementary Information for cloning procedures). We have listed all the primers, plasmids, and

103   strains we used in Table S1 and S2. We sequence-verified all plasmids used and transformed them

104   into the *E. coli* C43(DE3) background (Lucigen) to construct all the strains.

105   2.2   Cyt *c* expression and growth conditions for *E. coli* strains

To express cyt *c*, we first inoculated all the *E. coli* strains directly from -80 °C glycerol stocks in

5 mL LB media with antibiotics (50 μg/mL kanamycin and 34 μg/mL chloramphenicol) and grew

them at 37 °C overnight with 250 rpm sharking. Then we inoculated cell cultures (1:100 v/v ratio)

in 50 mL 2xYT media in foil-covered 250 mL flasks with the same antibiotic concentrations and

1 mM δ-ALA (aminolevulinic acid), and grew these cultures at 37 °C with 250 rpm sharking.

When the cell OD_{600} reached between 0.5 to 0.6, we added 4 μM IPTG to induce the expression

of cyt *c*, followed by culturing at 30 °C with 250 rpm shaking overnight. Then we washed cells

twice with M9 minimal medium (BD). All the chemicals and media are from Sigma-Aldrich unless

otherwise noted.

2.3   Redness screening of the *E. coli* strains with *ccm* variants

We used redness screening for detecting the overall expression level of cyt *c*, as previously described (Su et al., 2020). We first grew all the strains from -80 °C glycerol stocks, then inoculated cell cultures in 2xYT media with foil-covered flasks (see **section 2.2**). When the cell OD_{600} reached between 0.5 to 0.6, we split the cell cultures into 96-well deep-well plates (Polypropylene 2.2 mL, VWR), where each well contained 0.5 mL cell culture. We then induced the cells with a gradient concentration of IPTG (0, 0.5, 1, 2, 4, 8, 16, 32, 62.5, 125, 250, and 500 μM). After overnight incubation, we transferred the cell cultures into 96-well white round-bottom microplates (3789A, Corning), and centrifuged the plate to pellet the cells. After removing the media, we used a scanner to take pictures of the plates and measured the redness of the cell pellets with a MATLAB image processing program (Su et al., 2020).

2.4   Enhanced chemiluminescence measurement of cyt *c* expression

We used enhanced chemiluminescence (ECL) to detect the cyt *c* expression as previously described (Su et al., 2020). In brief, we mixed 20 μL of 10 OD_{600} washed cells (described in **2.1**) with an equal volume of Laemmli Sample Buffer (2x), and heated the mixture to 95°C for 15 min. Then we loaded the samples and protein ladder (Precision Plus Protein WesternC Standards) into an SDS-gel (4-20% Crit TGX Stain-Free Gel) to perform SDS-PAGE (160 V, 45 min). Next, we transferred proteins from the SDS-gel onto the nitrocellulose membrane (Trans-Blot Turbo Midi NC Transfer Packs) via the Trans-Blot Turbo Transfer System (Bio-Rad). We then used ECL kit (Pierce Pico West Enhanced Chemiluminescence substrate, ThermoFisher) to perform the cyt *c* detection assay. Finally, we imaged the ECL signals using the FluorChem E system (ProteinSimple). All the chemicals and supplies are from Bio-Rad unless otherwise indicated.

2.5   Bioelectrochemical measurement

We used two-chamber, three-electrode electrochemical reactors to perform the electrochemical

experiments as previously described (Su et al., 2020). The reactors contained three electrodes: a

working electrode (graphite felt, Alfa Aesar) and a reference electrode (CHI111, Ag/AgCl with 3

M KCl, CH Instruments) in the anodic chamber, and a counter electrode (titanium wire, Alfa

Aesar) in the cathodic chamber. We also used a cation exchange membrane (CMI-7000S,

Membranes International) to separate these two chambers. Each chamber contained 150 mL M9

minimal medium (BD) and 50 mM D,L-lactate (Sigma-Aldrich) as the electrolyte. During the test,

we continuously purged the anodic chamber with N_2 gas to maintain anaerobic conditions, and we

kept the temperature at 30 °C by placing all the reactors in an incubator. We used a potentiostat

(VSP-300, Bio-Logic Science Instruments) to perform all the electrochemical measurements. We

set the applied potential at 0.2 V versus the reference electrode to conduct chronoamperometry,

and the electrical current (averaged at every 36 s) is reported as a function of the geometric surface

area of the working electrode (0.0016 m^2, two sides, diameter of 32 mm). For the differential pulse

voltammetry (DPV) measurement (Xu et al., 2016), we scanned the potential from -0.4 V to 0.2 V

versus the reference electrode, with the pulse height set to 50 mV, pulse width set to 500 ms, step

height set to 1.0 mV, and step time set to 1000 ms.

3.   RESULTS AND DISCUSSION

3.1   Design and construction of a hybrid cyt _c_ maturation operon in _Escherichia coli_

To test our hypothesis that MR-1's ligation module is more efficient at maturing MR-1 cyt _c_ than

E. coli's ligation module, we made a series of modifications to the _ccmH_ gene in the plasmid which

carries the _ccm_ operon from _E. coli_ (**Figure 2**). We then co-transformed these plasmids with a

separate plasmid that expresses *cymAmtrCAB* from MR-1 into *E. coli* C43(DE3) to create the

wildtype *ccm*, *ccmHI*, *ccmH^N*, and hybrid *ccmH* strains (**Table 1**). Specifically, to test the effect

of altering the entire ligation module, we replaced *ccmH* from *E. coli* with *ccmHI* from MR-1,

creating the I5101 plasmid in the *ccmHI* strain. To test the impact of replacing only the apo cyt *c*

binding component, we fused the N-terminal region of the *E. coli ccmH* with the C terminal region

of MR-1 *ccmI*, notated as the I5099 plasmid in hybrid *ccmH* strain. As a control, we also made the

I5100 plasmid in *ccmH^N* strain from the I5099 plasmid, in which the C-terminal domain of *S.*

oneidensis CcmI was removed. We also constructed a strain with the pACYC184 empty vector

and *cymAmtrCAB* plasmid, dubbed the 'no *ccm*' strain, as a negative control.

Figure 2. A schematic illustration describing the two plasmids we used for cytochrome *c*

expression in engineered *E. coli*. For the Mtr conduit plasmid, *cymA*, *mtrC*, *mtrA*, *mtrB* are under

the same T7 lacO promoter (P_{T7}); for the Ccm plasmid, all the components are under a constitutive

promoter ($P_{FAB\#640}$). In the Ccm plasmid, we replaced the coding sequence of *ccmH* from the *E.*

coli ccm operon with three other variants. These plasmids coded for CcmHI from MR-1 (I5101),

a fusion between the N terminal region of *E. coli ccmH* and the C terminal region of CcmI from

MR-1 (I5099), and a control that removed the C terminal region of MR-1 CcmI (I5100). Gray and

red regions indicate DNA regions that code for proteins from *E. coli* and MR-1, respectively. (ori

origin of replication; ⌐ promoter; ⬤ ribosome-binding site, RBS; ➡ coding sequence, CDS; ⊤

terminator).

Table 1. Strains Used in This Study

Strain Name	Parental Strain	Plasmid(s)	Genes*	Source
wildtype *ccm*	C43(DE3)	M0640	*ccmABCDEFGH*	(Goldbeck et al., 2013)
		I5049	*cymA, mtrCAB*	(Jensen et al., 2016)
hybrid *ccm*	C43(DE3)	I5099	*ccmABCDEFG, hybrid ccmH*	this work
		I5049	*cymA, mtrCAB*	(Jensen et al., 2016)
ccmH^N	C43(DE3)	I5100	*ccmABCDEFG, ccmH (N-term.)*	this work
		I5049	*cymA, mtrCAB*	(Jensen et al., 2016)
ccmHI	C43(DE3)	I5101	*ccmABCDEFG, ccmHI_SO*	this work
		I5049	*cymA, mtrCAB*	(Jensen et al., 2016)
no *ccm*	C43(DE3)	pACYC184	Empty plasmid vector (no *ccm*)	

I5049	*cymA, mtrCAB*	(Jensen et al., 2016)

181   * Details of the genes are available in supporting information, Table S2.

3.2   The hybrid Ccm system can increase cyt *c* abundance in *E. coli*

To test if these *ccm* variants can mature and change the cyt *c* expression level, we first examined

the cyt *c* expression of these *E. coli* strains with a redness assay (Su et al., 2020) across different

IPTG induction levels (**Figure 3**). Since IPTG induction level is non-linear with protein

expression, we present this data in term of relative promoter strength, which is the GFP expression

per cell at a IPTG concentration normalized by the maximal GFP expression per cell (Goldbeck et

al., 2013). The redness value, which indirectly measures the cyt *c* per cell, peaked at the relative

promoter strength of ~ 0.1 (arb. strength units) for all the strains with a Ccm plasmid. As expected,

the no *ccm* strain has the lowest redness value because it cannot mature cyt *c,* and the wildtype

ccm strain has a high redness value, consistent with its ability to mature cyt *c*. Compared to the

wildtype *ccm* strain, the redness value of the *ccmHI* strain decreased by ~40% (0.15 ± 0.003 arb.

red units in *ccmHI* vs. 0.25 ± 0.003 arb. red units in wildtype). This decrease in cyt *c* expression

might be due to a poor ability of the *E. coli* and MR-1 proteins to form a functional complex

(Zheng et al., 2012). In contrast, the hybrid *ccmH* strain has a redness value (0.27 ± 0.002 arb. red

units) that is 8% greater than the wildtype *ccm* strain, indicating that it can mature cyt *c* slightly

better. This robust cyt *c* expression shows the C terminal region of CcmI from MR-1 can

functionally replace the C terminal region of *E. coli* CcmH. Overall, these data show that the hybrid

ccm system can increase cyt *c* abundance in *E. coli*, which potentially may increase its EET

efficiency.

Figure 3. The hybrid *ccmH* strain expresses more cytochromes *c*. Relative redness as a function of relative promoter strength, i.e., induction level, for *E. coli* expressing *cymAmtrCAB* with different *ccm* operons. The hybrid *ccmH* strain has a higher level of overall cyt *c* expression than wildtype *ccm* strain. Results are representative of three independent experiments.

3.3   The hybrid Ccm system changes Mtr cyt *c* stoichiometry

To further investigate how the hybrid *ccm* system alters the expression of cyt *c* in the Mtr pathway, we used enhanced chemiluminescence (ECL) to characterize the abundance of CymA, MtrC, and MtrA in hybrid *ccmH* strain, wildtype *ccm* strain, and no *ccm* strain (**Figure S2**). These strains were all induced at 0.068 arb. strength units relative promoter strength (4 μM IPTG) since the hybrid *ccm* strain has one of the highest cyt *c* expressions at this induction (**Figure 3**). As expected, there was no detectable amount of cyt *c* in the no *ccm* strain (**Figure S2**). While the MtrC abundance was very similar, the abundance of MtrA decreased ~15%, and CymA increased ~60% in the hybrid *ccm* strain relative to the wildtype *ccm* strain (**Figure 4**). These data show that

215   the hybrid *ccm* system increases the overall cyt *c* abundance by increasing the maturation of
216   CymA.

Figure 4. The hybrid *ccm* system increases the CymA expression in *E. coli*. The relative density
of enhanced chemiluminescence (ECL) bands arising from three cyt *c* heterologously expressed
in the hybrid *ccm* and wildtype *ccm* strains. The cyt *c* expression in negative control no *ccm* strain
was below the limit of detection. Results are representative of three independent experiments.

3.4   The hybrid Ccm system increases the current generation of Mtr-expressing *E. coli*

Altering the stoichiometry of Mtr cyt *c* in *E. coli* can negatively or positively affect its
bioelectrochemical performance (Su et al., 2020). To probe if the increase of CymA expression in
hybrid *ccmH* strain will also enhance its bioelectrical performance, we measured the current
generation (**Figure 5**) by the wildtype *ccm*, hybrid *ccmH*, *ccmH^N* strains in three-electrode

electrochemistry bioreactors. The wildtype *ccm* strain produced 11.47 ± 0.19 mA/m^2, in line with

our previous observations (Su et al., 2020). The hybrid *ccmH* strain produced 25.32 ± 3.61mA/m^2,

a 121% increase in current over the wildtype *ccm* strain. The *ccmH^N* strain produced only $3.42 \pm$

0.44 mA/m^2, about a 70% decrease relative to the wildtype *ccm* strain, similar to the no *mtr* strain

previously reported (TerAvest et al., 2014). Thus, we conclude that the hybrid *ccmH* strain matures

more CymA and produces higher current than the *E. coli* Ccm system.

Figure 5. The hybrid *ccmH* strain produces a higher current. The current generation measured

by chronoamperometry in the bioreactors showing hybrid *ccmH* strain produces more current than

the wildtype *ccm* strain, while the negative control strain *ccmH^N* produces less current. Results are

representative of three independent experiments. The dotted arrow shows the time the bacteria

were injected.

To probe the mechanism underlying the increase in current production by the hybrid *ccmH* strain,

we performed differential pulse voltammetry (DPV) (**Figure 6**) after strains reached the steady-

state current generation after 90 h chronoamperometry testing. We observed a single anodic peak

at redox potential (E_p) of +51 mV$_{SHE}$ in all three strains, which has been previously assigned to the

MtrCAB complex (Okamoto et al., 2013). This observation confirms that all three strains use the

MtrCAB conduit for the current production. The peak height in hybrid *ccmH* strain reached 8.31

μA, about 118% higher than the 3.82 μA in wildtype *ccm* strain, while in the *ccmH*^{N} strain, it was

only 0.09 μA, about 2% of the wildtype. The ratio of the peak height for the hybrid *ccm* strain

relative to the wildtype *ccm* strain (2.18) closely mirrors the ratio of steady-state current for these

strains (2.21), indicating the increase in steady-state current arises from an increase in the flux of

electrons through the MtrCAB complex. Thus, the hybrid *ccm* strain produces more current by

more effectively maturing CymA, boosting its relative abundance, and increasing the electron flux

through the MtrCAB conduit.

Figure 6. The hybrid *ccmH* strain shows higher redox activity through cyt *c*. Differential pulse

voltammetry (DPV) shows that the hybrid *ccmH* strain has a higher redox peak at the cyt *c* related

redox reaction potential (+51 mV vs. SHE). DPV signal smoothing and background deduction by

QSoas Version 2.0 (Fourmond et al., 2009).

3.5   Hybrid *ccmH* strain produces more current per cell via both direct and indirect EET

While DPV confirms that the increase in electron flux through the Mtr pathway increases the

current density, this increased flux could arise from an increase in the current per cell, an increase

in the number of viable cells, or an increase in both. To distinguish between these possibilities, we

measured both the current production and the number of viable cells using colony-forming unit

counting in the $ccmH^N$, hybrid *ccm*, and wildtype *ccm* strains (**Figure 7A**). The $ccmH^N$ strain

produced near baseline current on a per-cell basis (1.0 ± 0.005 fA·cell^{-1}), as expected. Interestingly,

the hybrid *ccmH* strains produced ~50% more current per cell than the wildtype *ccm* (12.3 ± 0.3

fA·cell^{-1} versus 8.3 ± 0.4 fA·cell^{-1}). These data confirm that the hybrid *ccmH* strain produces more

current at the single-cell level by direct EET.

In addition to direct EET via Mtr conduit, MR-1 (Brutinel and Gralnick, 2012; Coursolle et al.,

2010) can also perform EET with flavins as a bound cofactor to cyt *c* (Edwards et al., 2015;

Okamoto et al., 2014) or as a soluble mediator (Kotloski and Gralnick, 2013). To test if the hybrid

ccmH strain can also utilize flavin for EET, we added exogenous flavins and measured the current

density per cell. The addition of riboflavin to the $ccmH^N$ strain resulted in a meager current increase

(0.02 ± 0.004 fA·cell^{-1}·µM^{-1}). As expected, the current produced by both the wildtype *ccm* and

hybrid *ccm* strains linearly increased with the exogenous flavin concentration (**Figure 7A**).

However, the hybrid *ccm* strain produced significantly more current per µM exogenous flavin

(0.32 ± 0.06 fA·cell^{-1}· µM^{-1}) than the wildtype *ccm* strain (0.16 ± 0.01 fA·cell^{-1}·µM^{-1}) (**Figure**

7A). These data indicate that the hybrid *ccm* strain produces more current at the single-cell level

via a flavin-mediated EET as well as a direct EET mechanism.

(A)

Figure 7. Current production by the hybrid *ccmH* strain is more sensitive to exogenous

flavins. (A) The hybrid *ccmH* strain (red) has a more rapid and higher current response as a

function of extra riboflavin concentration, compared to the wildtype *ccm* strain (gray), and *ccmH^N*

strain (yellow). Results are representative of two independent experiments. Linear fit analysis

(with Y error) is performed by OriginPro 2019b (OriginLab Corporation). (B) A schematic

depicting the extracellular electron transfer via Mtr conduit. MtrCAB complex receives the

electrons from CymA, then delivers to the electrode (I_{strain}) through either MtrC (I_{Mtr}) or flavin

(I_{flavin}). (OM: outer membranes, IM: Inner membranes).

3.6   Direct and flavin-mediated EET in the hybrid *ccmH* strain is limited by CymA

Our data suggest that the CymA amount limits electron flux and that the hybrid *ccmH* strain

achieves higher current by enabling a greater electron flux from CymA into the same direct and

indirect EET routes found in the wildtype *ccm* strain. To quantitatively test these hypotheses, we

developed a quantitative model that relates EET in the hybrid *ccmH* strain to EET in the wildtype

ccm strain and then tested whether this model could explain the current per cell measurement in

the hybrid *ccmH* strain (**Figure 7**).

Explicitly, we modeled the overall EET current per cell (I_{strain}) as the sum of the direct EET via

Mtr conduit (I_{Mtr}) and flavin-mediated EET (I_{flavin}) (**Figure 7B**). We describe the flavin-mediated

EET as the product of the flavin concentration ($[flavin]$) and a constant of proportionality (k_{strain}),

(**Equation 1**):

$$I_{strain} = I_{strain,Mtr} + I_{strain,flavin} = I_{strain,Mtr} + k_{strain}[flavin] \qquad (1)$$

Assuming the amount of CymA is rate-limiting in our system (**Figure 4**), we can then calculate

EET current in the hybrid *ccmH* strain (I_{hybrid}) as a ratio of the CymA concentration in the two

strains and the current in the wildtype *ccm* strain ($I_{WT,Mtr}$) with the equation below (**Equation 2**):

$$I_{hybrid} = \frac{[CymA]_{hybrid}}{[CymA]_{WT}} I_{WT} = \frac{[CymA]_{hybrid}}{[CymA]_{WT}} I_{WT,Mtr} + \frac{[CymA]_{hybrid}}{[CymA]_{WT}} k_{WT}[flavin] \qquad (2)$$

Thus, both the intercept and slope of the best-fit line for the hybrid *ccm* strain should be equal to

the intercept and slope best-fit line of the wildtype *ccm* strain times the ratio of CymA

concentration in these two strains.

Using the experimentally-determined values of $I_{WT,Mtr}$ (8.3 ± 0.4 fA·cell^{-1}), k_{WT} (0.16 ± 0.01

fA·cell^{-1}·µM^{-1}), and $[CymA]_{hybrid}/[CymA]_{WT} = 1.6$, we can predict the values of I_{hybrid} and compare

it to our experimentally-determined values (**Figure 7A**). Both the predicted $I_{hybrid, Mtr}$ and $I_{hybrid,}$

flavin values, 13.28 ± 0.64 fA·cell^{-1} and 0.26 ± 0.02 fA·cell^{-1}·µM^{-1}, respectively, are quite close to

our observation of the direct EET (12.3 ± 0.3 fA·cell^{-1}) and mediated EET (0.32 ± 0.06 fA·cell^{-1}·µM^{-1}) in hybrid *ccmH* strain. This result provides strong evidence that the hybrid *ccmH* strain's

improved bioelectrochemical performance arises mainly from the CymA expression increase and

underscores the importance of CymA in both direct EET and indirect EET.

4. DISCUSSION

In this study, we have shown that the C-terminal region of CcmH can be complemented by the

homologous region of CcmI from *S. oneidensis* MR-1, which increases the bioelectronic

performance of Mtr-expressing *E. coli*. We found that the hybrid CcmH can increase the overall

cyt *c* expression (**Figure 3**), with a preferred 60% increase in CymA expression (**Figure 4**). This

increase in CymA expression led to a 48% more EET current per bacterium and doubled the

electrical response to exogenous flavin (**Figure 7**), consistent with CymA being rate-limiting in

EET. In the following, we discuss how these findings impact the development of biofuel cells and

biosensors.

First, our data show that the C-terminal region of CcmH is essential for cyt *c*-based EET.

Shewanella and *Geobacter* species are the dominant organisms used for EET applications, such as

MFCs. These microorganisms express a significant amount of cyt *c* to perform EET (Meyer et al.,

2004; Shi et al., 2009). As a result, there are increasing efforts focused on genetically introducing

cyt *c* to make new organisms capable of EET (Schuergers et al., 2017; Sekar et al., 2016;

Thirumurthy and Jones, 2020), overexpressing cyt *c* to understand the structural basis of EET

(Edwards et al., 2020, 2019; Wang et al., 2019), and making cyt *c* -based conductive biomaterials

for bioelectrical applications (Liu et al., 2020; Sun et al., 2018). Previously we used a random

mutagenesis approach to improve cyt *c* maturation, which yielded strains that both increased

current by up to 77% and decreased current production by as much as -66% (Su et al., 2020).

Leveraging understanding from our previous study, this work presents a rational approach that

increased current by 121%. Thus, understanding how cyt *c* mature can provide guidance on

manipulating cyt *c* expression, which will assist both fundamental studies and these applied efforts

to increase the bioelectrical performance of MFCs.

Second, we found that the hybrid *ccmH* strain generates higher electrical signals using both Mtr

and flavin-mediated EET. Increasing the EET efficiency of Mtr-expressing *E. coli* is a particular

interest for biosensing because it can increase the signal-to-noise ratio, enabling higher sensitivity

detection. In recent years, much progress has been made in engineering the electrical

communication between bacteria and electrodes, such as responding to specific chemicals using

chemical-sensitive promoters (Golitsch et al., 2013; Webster et al., 2014; Yang et al., 2018),

miniaturized reactors with optimized structure design (Zhou et al., 2017), and modified material-

bacteria interface (Zajdel et al., 2018). Enhancing the bioelectrical performance of Mtr-expression

E. coli offers a reliable chassis in biosensing applications.

Lastly, our work provides guidance us on how to build a more efficient EET conduit for

bioelectrical sensing. Our model for the current generation in hybrid *ccmH* strain together with the

fact that CymA can only carry 40% (fewer heme groups per protein) as many electrons as MtrA

or MtrC, strongly suggests that electron transfer through CymA is a bottleneck for EET in the Mtr-

expression *E. coli*. Future work will focus on increasing the electron flux between menaquinone

and the MtrCAB complex to overcome this rate-limiting step. Also, since the electrical response

is linear in both CymA amount and flavin concentration, this informs more information rich

sensing scenarios, including building an "AND" gate with dual inputs.

5. CONCLUSIONS

In summary, a hybrid Ccm system altered the stoichiometry of the expressed cyt *c,* thus accelerating the rate-limiting step in EET and enhancing the bioelectronic performance of *E. coli*. Our results suggest that increasing the electron flux from menaquinone to MtrCAB complex could further enhance the bioelectrical performance of engineered *E. coli*. More broadly, our work provides a new strategy to express cyt *c* that underlies microbial bioelectronics and opens new possibilities for engineering more sophisticated and sensitive biosensors.

SUPPLEMENTARY DATA

Supporting Information is available online.

AUTHOR CONTRIBUTIONS

L.S. contributed to strain construction, design of the study, conducting experiments, analysis of the data, and writing and revision of the manuscript. T.F. contributed to strain construction, design of the study, conducting experiments, analysis of the data, review and revision of the manuscript. C.M.A-F. contributed to the design of the study, analysis of the data, and writing of the manuscript.

ACKNOWLEDGMENT

We thank Dr. Shuai Xu and Prof. Mohamed Y. El-Naggar, the University of Southern California for discussions on the DPV experiments. Work at the Molecular Foundry was supported by the Office of Science, Office of Basic Energy Sciences, of the U.S. Department of Energy [DE-AC02-

377   05CH11231]. This work was supported by the Office of Naval Research, [N0001418IP00037];

378   and the China Scholarship Council [201606090098 to L.S.].

REFERENCES

Brutinel, E.D., Gralnick, J.A., 2012. Appl. Microbiol. Biotechnol. 93, 41–48.

   https://doi.org/10.1007/s00253-011-3653-0

Cheng, Z., Xiong, J., Min, D., Cheng, L., Liu, D., Li, W., Jin, F., Yang, M., Yu, H., 2020. Biotechnol.

   Bioeng. 117, 1294–1303. https://doi.org/10.1002/bit.27305

Chouler, J., Cruz-Izquierdo, Á., Rengaraj, S., Scott, J.L., Di Lorenzo, M., 2018. Biosens. Bioelectron.

   102, 49–56. https://doi.org/10.1016/j.bios.2017.11.018

Clarke, T.A., Edwards, M.J., Gates, A.J., Hall, A., White, G.F., Bradley, J., Reardon, C.L., Shi, L.,

   Beliaev, A.S., Marshall, M.J., Wang, Z., Watmough, N.J., Fredrickson, J.K., Zachara, J.M., Butt,

   J.N., Richardson, D.J., 2011. Proc. Natl. Acad. Sci. U. S. A. 108, 9384–9389.

   https://doi.org/10.1073/pnas.1017200108

Coursolle, D., Baron, D.B., Bond, D.R., Gralnick, J.A., 2010. J. Bacteriol. 192, 467–474.

   https://doi.org/10.1128/JB.00925-09

Edwards, M.J., Richardson, D.J., Paquete, C.M., Clarke, T.A., 2019. Protein Sci. 29, 830–842.

   https://doi.org/10.1002/pro.3787

Edwards, M.J., White, G.F., Butt, J.N., Richardson, D.J., Clarke, T.A., 2020. Cell. 181, 665–673.

   https://doi.org/10.1016/j.cell.2020.03.032

Edwards, M.J., White, G.F., Norman, M., Tome-Fernandez, A., Ainsworth, E., Shi, L., Fredrickson, J.K.,

   Zachara, J.M., Butt, J.N., Richardson, D.J., Clarke, T.A., 2015. Sci. Rep. 5, 11677.

   https://doi.org/10.1038/srep11677

Fabianek, R.A., Hofer, T., Thöny-Meyer, L., 1999. Arch. Microbiol. 171, 92–100.

   https://doi.org/10.1007/s002030050683

Fourmond, V., Hoke, K., Heering, H.A., Baffert, C., Leroux, F., Bertrand, P., Léger, C., 2009.

   Bioelectrochemistry 76, 141–147. https://doi.org/10.1016/j.bioelechem.2009.02.010

Goldbeck, C.P., Jensen, H.M., TerAvest, M.A., Beedle, N., Appling, Y., Hepler, M., Cambray, G.,

Mutalik, V., Angenent, L.T., Ajo-Franklin, C.M., 2013. ACS Synth. Biol. 2, 150–159.

https://doi.org/10.1021/sb300119v

Golitsch, F., Bücking, C., Gescher, J., 2013. Biosens. Bioelectron. 47, 285–291.

https://doi.org/10.1016/j.bios.2013.03.010

Gul, M.M., Ahmad, K.S., 2019. Biosens. Bioelectron. 142, 111576.

https://doi.org/10.1016/j.bios.2019.111576

Harris, H.W., Sánchez-Andrea, I., McLean, J.S., Salas, E.C., Tran, W., El-Naggar, M.Y., Nealson, K.H.,

2017. Front. Microbiol. 8, 2568. https://doi.org/10.3389/fmicb.2017.02568

Jensen, H.M., TerAvest, M.A., Kokish, M.G., Ajo-Franklin, C.M., 2016. ACS Synth. Biol. 5, 679–688.

https://doi.org/10.1021/acssynbio.5b00279

Kotloski, N.J., Gralnick, J.A., 2013. mBio. 4, e00553-12. https://doi.org/10.1128/mbio.00553-12

Kranz, R.G., Richard-Fogal, C., Taylor, J.-S., Frawley, E.R., 2009. Microbiol. Mol. Biol. Rev. 73, 510–

528. https://doi.org/10.1128/MMBR.00001-09

Liu, X., Gao, H., Ward, J.E., Liu, X., Yin, B., Fu, T., Chen, J., Lovley, D.R., Yao, J., 2020. Nature. 578,

550–554. https://doi.org/10.1038/s41586-020-2010-9

Logan, B.E., 2009. Nat. Rev. Microbiol. 7, 375–381. https://doi.org/10.1038/nrmicro2113

Meyer, T.E., Tsapin, A.I., Vandenberghe, I., de Smet, L., Frishman, D., Nealson, K.H., Cusanovich,

M.A., van Beeumen, J.J., 2004. OMICS 8, 57–77. https://doi.org/10.1089/153623104773547499

Min, D., Cheng, L., Zhang, F., Huang, X.-N., Li, D.-B., Liu, D.-F., Lau, T.-C., Mu, Y., Yu, H.-Q., 2017.

Environ. Sci. Technol. 51, 5082-5089. https://doi.org/10.1021/acs.est.6b04640

Okamoto, A., Hashimoto, K., Nealson, K.H., Nakamura, R., 2013. Proc. Natl. Acad. Sci. U. S. A. 110,

7856–7861. https://doi.org/10.1073/pnas.1220823110

Okamoto, A., Nakamura, R., Hashimoto, K., 2011. Electrochim. Acta 56, 5526–5531.

https://doi.org/10.1016/j.electacta.2011.03.076

Okamoto, A., Nakamura, R., Nealson, K.H., Hashimoto, K., 2014. ChemElectroChem 1, 1808–1812.

https://doi.org/10.1002/celc.201402151

Robertson, I.B., Stevens, J.M., Ferguson, S.J., 2008. FEBS Lett. 582, 3067–3072.

https://doi.org/10.1016/j.febslet.2008.07.052

Rosenbaum, M.A., Bar, H.Y., Beg, Q.K., Segrè, D., Booth, J., Cotta, M.A., Angenent, L.T., 2012. PLoS

One 7, e30827. https://doi.org/10.1371/journal.pone.0030827

Sanders, C., Turkarslan, S., Lee, D.-W., Daldal, F., 2010. Trends Microbiol. 18, 266–274.

https://doi.org/10.1016/j.tim.2010.03.006

Santoro, C., Arbizzani, C., Erable, B., Ieropoulos, I., 2017. J. Power Sources 356, 225–244.

https://doi.org/10.1016/j.jpowsour.2017.03.109

Schmitz, S., Nies, S., Wierckx, N., Blank, L.M., Rosenbaum, M.A., 2015. Front. Microbiol. 6, 284.

https://doi.org/10.3389/fmicb.2015.00284

Schuergers, N., Werlang, C., Ajo-Franklin, C.M., Boghossian, A.A., 2017. Energy Environ. Sci. 10,

1102–1115. https://doi.org/10.1039/C7EE00282C

Sekar, N., Jain, R., Yan, Y., Ramasamy, R.P., 2016. Biotechnol. Bioeng. 113, 675–679.

https://doi.org/10.1002/bit.25829

Shi, L., Dong, H., Reguera, G., Beyenal, H., Lu, A., Liu, J., Yu, H.-Q., Fredrickson, J.K., 2016. Nat. Rev.

Microbiol. 14, 651–662. https://doi.org/10.1038/nrmicro.2016.93

Shi, L., Richardson, D.J., Wang, Z., Kerisit, S.N., Rosso, K.M., Zachara, J.M., Fredrickson, J.K., 2009.

Environ. Microbiol. Rep. 1, 220–227. https://doi.org/10.1111/j.1758-2229.2009.00035.x

Slate, A.J., Whitehead, K.A., Brownson, D.A.C., Banks, C.E., 2019. Renewable Sustainable Energy Rev.

101, 60–81. https://doi.org/10.1016/j.rser.2018.09.044

Sturm-Richter, K., Golitsch, F., Sturm, G., Kipf, E., Dittrich, A., Beblawy, S., Kerzenmacher, S.,

Gescher, J., 2015. Bioresour. Technol. 186, 89–96. https://doi.org/10.1016/j.biortech.2015.02.116

Su, L., Ajo-Franklin, C.M., 2019. Curr. Opin. Biotechnol. 57, 66–72.

https://doi.org/10.1016/j.copbio.2019.01.018

Su, L., Fukushima, T., Prior, A., Baruch, M., Zajdel, T.J., Ajo-Franklin, C.M., 2020. ACS Synth. Biol. 9,

115–124. https://doi.org/10.1021/acssynbio.9b00379

Sun, Y.-L., Tang, H.-Y., Ribbe, A., Duzhko, V., Woodard, T.L., Ward, J.E., Bai, Y., Nevin, K.P.,

Nonnenmann, S.S., Russell, T., Emrick, T., Lovley, D.R., 2018. Small 14, e1802624.

https://doi.org/10.1002/smll.201802624

TerAvest, M.A., Zajdel, T.J., Ajo-Franklin, C.M., 2014. ChemElectroChem 1, 1874–1879.

https://doi.org/10.1002/celc.201402194

Thirumurthy, M.A., Jones, A.K., 2020. Nanotechnology 31, 124001. https://doi.org/10.1088/1361-

6528/ab5de6

Thöny-Meyer, L., 2000. Biochim. Biophys. Acta 1459, 316–324. https://doi.org/10.1016/s0005-

2728(00)00167-5

Verissimo, A.F., Yang, H., Wu, X., Sanders, C., Daldal, F., 2011. J. Biol. Chem. 286, 40452–40463.

https://doi.org/10.1074/jbc.M111.277764

Wang, F., Gu, Y., O'Brien, J.P., Yi, S.M., Yalcin, S.E., Srikanth, V., Shen, C., Vu, D., Ing, N.L.,

Hochbaum, A.I., Egelman, E.H., Malvankar, N.S., 2019. Cell 177, 361–369.

https://doi.org/10.1016/j.cell.2019.03.029

Wang, X., Gao, N., Zhou, Q., 2013. Biosens. Bioelectron. 43, 264–267.

https://doi.org/10.1016/j.bios.2012.12.029

Webster, D.P., TerAvest, M.A., Doud, D.F.R., Chakravorty, A., Holmes, E.C., Radens, C.M., Sureka, S.,

Gralnick, J.A., Angenent, L.T., 2014. Biosens. Bioelectron. 62, 320–324.

https://doi.org/10.1016/j.bios.2014.07.003

West, E.A., Jain, A., Gralnick, J.A., 2017. ACS Synth. Biol. 6, 1627–1634.

https://doi.org/10.1021/acssynbio.6b00349

Xu, S., Jangir, Y., El-Naggar, M.Y., 2016. Electrochim. Acta 198, 49–55.

https://doi.org/10.1016/j.electacta.2016.03.074

Yang, Y., Zhou, H., Mei, X., Liu, B., Xing, D., 2018. Front. Microbiol. 9, 1924.

https://doi.org/10.3389/fmicb.2018.01924

Zajdel, T.J., Baruch, M., Méhes, G., Stavrinidou, E., Berggren, M., Maharbiz, M.M., Simon, D.T., Ajo-

Franklin, C.M., 2018. Sci. Rep. 8, 15293. https://doi.org/10.1038/s41598-018-33521-9

Zheng, X.-M., Hong, J., Li, H.-Y., Lin, D.-H., Hu, H.-Y., 2012. Biochim. Biophys. Acta 1824, 1394–

1400. https://doi.org/10.1016/j.bbapap.2012.06.017

Zhou, A.Y., Baruch, M., Ajo-Franklin, C.M., Maharbiz, M.M., 2017. PLoS One 12, e0184994.

https://doi.org/10.1371/journal.pone.0184994

487