A hybrid cyt *c* maturation system enhances the bioelectrical performance of engineered *Escherichia coli* by improving the rate-limiting step

Author List:

Lin Su [1,2,3], Tatsuya Fukushima [3], Caroline M. Ajo-Franklin [2,3,4,5*]

Author Affiliations:

[1] State Key Laboratory of Bioelectronics, School of Biological Science and Medical Engineering, Southeast University, Nanjing 210018, China

[2] Department of BioSciences, Rice University, Houston, TX 77005, USA

[3] Molecular Foundry, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

[4] Molecular Biophysics and Integrated Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

[5] Synthetic Biology Institute, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

* To whom correspondence should be addressed. Tel: 1-713-348-3497; Fax: 1-713-348-4790; Email: cajo-franklin@rice.edu

ABSTRACT

Bioelectronic devices can use electron flux to enable communication between biotic components

and abiotic electrodes. We have modified *Escherichia coli* to electrically interact with electrodes

by expressing the cytochrome *c* from *Shewanella oneidensis* MR-1. However, we observe

inefficient electrical performance, which we hypothesize is due to the limited compatibility of the

E. coli cytochrome *c* maturation (*ccm*) systems with MR-1 cyt *c*. Here we test whether the

bioelectronic performance of *E. coli* can be improved by constructing hybrid *ccm* systems with

parts from both *E. coli* and *S. oneidensis MR-1*. Our results show that the hybrid CcmH can

increase cytochrome *c* expression and the stoichiometry of protein components also shifted.

Electrochemical measurements show that the electron flux per cell increased 48% with the hybrid

system. Additionally, the electrical response doubled with addition of exogenous flavin, which

gives us a quantitative understanding of the bottleneck step in this electron conduit. These results

demonstrate that this hybrid *ccm* system can enhance the bioelectrical performance of the cyt *c*

expressing *Escherichia coli*, allowing to build more efficient bioelectronic devices.

KEYWORDS

Microbial electrochemistry, microbial fuel cell, bioelectronic sensor, synthetic biology,

cytochrome *c* maturation

1. INTRODUCTION

The flow of electrons can transfer both energy and information between living and non-living systems for bioelectronic and biosensing applications. Extracellular electron transfer (EET) pathways (Shi et al., 2016) can electrically connect living microorganisms with electrochemical cells, enabling applications such as power generation in microbial fuel cells (MFCs) (Gul and Ahmad, 2019; Logan, 2009; Santoro et al., 2017; Slate et al., 2019) and environmental sensing in bioelectronic sensing systems (Chouler et al., 2018; Wang et al., 2013; Zhou et al., 2017). Engineering EET in native electroactive microorganisms, such as *Shewanella oneidensis* MR-1 (Cheng et al., 2020; Min et al., 2017; West et al., 2017), and non-native hosts, such as *Escherichia coli (Jensen et al., 2016; Sturm-Richter et al., 2015)*, has expanded the range of potential applications (Harris et al., 2017; Su and Ajo-Franklin, 2019). However, a fundamental limitation in these bioelectronic systems is the low rate of EET, which causes these devices to have low power output, low sensing sensitivity, and limited robustness. For example, expressing the EET pathway from *Shewanella oneidensis* MR-1 into engineered *E. coli* only yields ~10% the current produced by MR-1 (TerAvest et al., 2014).

Although the molecular structure of these EET pathways is increasingly well-understood, how these complex proteins are biosynthesized is far less understood. The EET pathways (Mtr pathway) in *Shewanella oneidensis* MR-1 is a group of multiheme cytochrome *c* (cyt *c*) proteins embedded in the inner membrane (CymA), periplasm (ScyA, CctA, etc.), and outer membrane (MtrCAB complex) (Clarke et al., 2011; Okamoto et al., 2011; Rosenbaum et al., 2012). Expressing these cyt *c* requires the cytochrome *c* maturation (Ccm) system, which covalently ligates heme groups to the apo-protein (Sanders et al., 2010; Verissimo et al., 2011). We and others have found that expressing components of the Mtr pathway and the Ccm system enables *E. coli* to perform EET

(Jensen et al., 2016; Su et al., 2020; TerAvest et al., 2014). In the Ccm systems of Gram-negative

bacteria, CcmH is essential to the heme ligation process (Sanders et al., 2010), which includes

translocating heme groups, and formatting a thioether bond between the heme group and the apo

cyt *c*. We recently found that mutations in the C-terminal domain of CcmH can significantly

change the EET efficiency by changing the stoichiometry of the Mtr cyt *c (Su et al., 2020)*. This

finding suggested that optimizing the heme ligation process by redesigning CcmH could remodel

the stoichiometry of cyt *c* and enhance the bioelectrical performance of Mtr-expressing *E. coli*.

The role of the C-terminal domain of CcmH in cyt *c* maturation is disputed. Fabianek et al.

(Fabianek et al., 1999) and Robertson et al. (Robertson et al., 2008) demonstrated that a native

diheme cyt *c* (NapB) and a heterologous monoheme cyt *c* (Cyt c C550) can be matured in *E. coli*

lacking the C-terminal of CcmH. They contend that the C-terminal is not required for maturation

because of functional redundancy between the two domains of CcmH. In contrast, Zheng et al.

(Zheng et al., 2012) predicted the C-terminal of CcmH is a TRP (tetratricopeptide repeat) -like

domain which for recognition of apo-cyt *c* and predicts it is essential for cyt *c* expression. Our

previous results suggest the C-terminal of CcmH is essential in expressing multi-heme

heterologous cyt *c (Su et al., 2020)*, agreeing more closely with Zheng et al. (Zheng et al., 2012).

To determine if MR-1's ligation module is more efficient at maturing cyt *c*, here we constructed

a hybrid CcmH that uses protein domains from *E. coli* and *S. oneidensis* MR-1. We then explored

the effects on cyt *c* expression and EET efficiency to test if this hybrid CcmH could boost the

bioelectrical performance of Mtr-expressing *E. coli*.

2. METHODS

2.1   Construction of *E. coli* strains with *ccm* variants

We used Phusion DNA polymerase (NEB) for all the PCR amplification (please refer to

Supplementary Information for cloning procedures). We have listed all the primers, plasmids, and

strains we used in Table S1 and S2. We sequence verified all plasmids used and transformed them

into the *E. coli* C43(DE3) background (Lucigen) to construct all the strains.

2.2   Cyt *c* expression and growth conditions for *E. coli* strains

To express cyt *c*, we first grew all the *E. coli* strains directly from -80 °C glycerol stores in 5 mL

LB media with antibiotics (50 µg/mL kanamycin and 34 µg/mL chloramphenicol), incubated at

37 °C overnight with 250 rpm sharking. Then we inoculated cell cultures (1:100 v/v ratio) in 50

mL 2xYT media in foil-covered 250 mL flasks with the same antibiotic concentrations and 1 mM

δ-ALA (aminolevulinic acid), and grew these cultures at 37 °C with 250 rpm sharking. When the

cell OD_{600} reached between 0.5 to 0.6, we added 4 µM IPTG to induce the expression of cyt *c*,

followed by culturing at 30 °C with 250 rpm shaking overnight. Then we washed cells twice with

M9 minimal medium (BD). All the chemicals and media are from Sigma-Aldrich unless otherwise

noted.

2.3   Redness screening of the *E. coli* strains with *ccm* variants

We used redness screening for detecting the overall expression level of cyt *c*, as previously

described (Su et al., 2020). We first grew all the strains from -80 °C glycerol stores, then inoculated

cell cultures in 2xYT media with foil-covered flasks (see **section 2.2**). When the cell OD_{600} reached

between 0.5 to 0.6, we split the cell cultures into 96-well deep-well plates (Polypropylene 2.2 mL,

VWR), where each well contained 0.5 mL cell culture. We then induced the cells with a gradient

concentration of IPTG (0, 0.5, 1, 2, 4, 8, 16, 32, 62.5, 125, 250, and 500 µM). After overnight

incubation, we transferred the cell cultures into 96-well white round-bottom microplates (3789A,

Corning), and centrifuged to pellet the cells. After removing the media, we used a scanner to take

pictures of the plates and measured the redness of the cell pellets with a MATLAB image

processing program (Su et al., 2020).

2.4   Enhanced chemiluminescence measurement of cyt *c* expression

We used enhanced chemiluminescence (ECL) to detect the cyt *c* expression as previously

described (Su et al., 2020). In brief, we mixed 20 μL of 10 OD_{600} washed cells (described in **2.1**)

with an equal volume of Laemmli Sample Buffer (2x), and heated the mixture to 95°C for 15 min.

Then we loaded the samples and protein ladder (Precision Plus Protein WesternC Standards) into

an SDS-gel (4-20% Crit TGX Stain-Free Gel) to perform SDS-PAGE (160 V, 45 min). Next, we

transferred proteins from the SDS-gel onto the nitrocellulose membrane (Trans-Blot Turbo Midi

NC Transfer Packs) via the Trans-Blot Turbo Transfer System (Bio-Rad). We then used ECL kit

(Pierce Pico West Enhanced Chemiluminescence substrate, ThermoFisher) to perform the cyt *c*

detection assay. Finally, we used the FluorChem E system (ProteinSimple) to take the ECL images.

All the chemicals and supplies are from Bio-Rad unless otherwise indicated.

2.5   Bioelectrochemical measurement

We used two-chamber, three-electrode electrochemical reactors to perform the electrochemical

experiments as previously described (Su et al., 2020). The reactors contained three electrodes: a

working electrode (graphite felt, Alfa Aesar) and a reference electrode (CHI111, Ag/AgCl with 3

M KCl, CH Instruments) in the anodic chamber, and a counter electrode (titanium wire, Alfa Aesar)

in the cathodic chamber. We also used a cation exchange membrane (CMI-7000S, Membranes

International) to separate these two chambers. The electrolyte contained M9 minimal medium (BD)

and 50 mM D,L-lactate (Sigma-Aldrich). During the test, we continuously purged the anodic

chamber with N_2 gas to maintain anaerobic conditions, and we kept the temperature at 30 °C by

placing all the reactors in an incubator. We used a potentiostat (VSP-300, Bio-Logic Science

Instruments) to perform all the electrochemical measurements. We set the applied potential at 0.2

V versus the reference electrode to conduct chronoamperometry, and the electrical current

(averaged at every 36 s) is reported as a function of the geometric surface area of the working

electrode. For the differential pulse voltammetry (DPV) measurement (Xu et al., 2016), we

scanned the potential from -0.4 V to 0.2 V versus the reference electrode, with the pulse height set

to 50 mV, pulse width set to 500 ms, step height set to 1.0 mV, and step time set to 1000 ms.

3. RESULTS AND DISCUSSION

3.1   Design and construction of a hybrid cyt *c* maturation operon in *Escherichia coli*

To test our hypothesis that MR-1's ligation module is more efficient at maturing MR-1 cyt *c* than

E. coli's ligation module, we made a series of modifications to the *ccmH* gene in the plasmid which

carries the *ccm* operon from *E. coli* (Figure 1). To test the effect of altering the entire ligation

module, we replaced *ccmH* from *E. coli* with *ccmHI* from MR-1, creating the *ccmHI* plasmid. To

test the impact of replacing only the apo cyt *c* binding component, we fused the N-terminal region

of the *E. coli ccmH* with the C terminal region of MR-1 *ccmI*, notated as the hybrid *ccmH* plasmid.

As a control, we also made a $ccmH^N$ plasmid from the hybrid *ccmH* plasmid, in which the C-

terminal domain of *S. oneidensis* CcmI was removed. We then co-transformed the *ccm*, *ccmHI*,

hybrid *ccmH* and $ccmH^N$ plasmids with a separate plasmid that expresses *cymAmtrCAB* from MR-

1 (Figure S1) into *E. coli* C43(DE3) to create the wildtype *ccm*, *ccmHI*, $ccmH^N$, and hybrid *ccmH*

strains. We also constructed a strain with the pACYC184 empty vector and *cymAmtrCAB* plasmid,

dubbed the 'no *ccm'* strain, as a negative control.

3.2   The hybrid *ccm* system can increase cyt *c* abundance in *E. coli*

To test if these *ccm* variants can mature and change the cyt *c* expression level, we first examined

the cyt *c* expression of these *E. coli* strains with a redness assay (Su et al., 2020) across different

IPTG inducing levels (Figure 2), i.e., different promoter strength (Goldbeck et al., 2013). The

redness value, which indirectly measures the cyt *c* abundance per cell, peaked at the relative

promoter strength region around 0.1 intensity (arb. units) for all the strains with a *ccm* plasmid. As

expected, the no *ccm* strain has the lowest redness value because it cannot mature cyt *c,* and the

wildtype *ccm* has a high redness value, consistent with its ability to mature cyt *c*. Compared to the

wildtype *ccm* strain, the redness value of *ccmHI* strain decreased by ~40% (0.15 ± 0.003 in *ccmHI*

vs. 0.25 ± 0.003 in wildtype). This decrease in cyt *c* expression might be due to a poor ability of

the *E. coli* and MR-1 proteins to form a functional complex (Zheng et al., 2012). In contrast, the

hybrid *ccmH* strain has redness value (0.27 ± 0.002) that is 8% greater than the wildtype *ccm*,

indicating that this strain can mature cyt *c* slightly better than the wildtype. Increased cyt c

expression in hybrid *ccmH* strain shows the C terminal region of CcmI from MR-1 can functionally

replace the C terminal region of *E. coli ccmH*. Overall, these data show that the hybrid *ccm* system

can increase cyt *c* abundance in *E. coli*, which potentially may increase its EET efficiency.

3.3   The hybrid *ccm* system changes Mtr cyt *c* stoichiometry

To further investigate how the hybrid *ccm* system alters the expression of cyt *c* in the Mtr pathway,

we used enhanced chemiluminescence (ECL) to characterize the abundance of CymA, MtrC, and

MtrA in hybrid *ccmH* strain, wildtype *ccm* strain, and no *ccm* strain (**Figure S2**). These strains

were all induced at 0.068 relative promoter strength since the hybrid *ccm* strain has the highest cyt

c expression at this induction (Figure 2). As expected, there was no detectable amount of cyt *c* in

the no *ccm* strain (**Figure S2**). While the MtrC abundance was very similar, the abundance of

MtrA decreased ~15%, and CymA increased ~60% in the hybrid *ccm* strain relative to the wildtype

ccm strain ([Figure 3](#)). These data show that the hybrid *ccm* system increases the overall cyt *c*

abundance by increasing the maturation of CymA.

3.4   The hybrid *ccm* system increases the current generation of Mtr-expressing *E. coli*

Altering the stoichiometry of Mtr cyt *c* in *E. coli* can negatively or positively affect its

bioelectrochemical performance (Su et al., 2020). To probe if the increase of CymA expression in

hybrid *ccmH* strain will also enhance its bioelectrical performance, we measured the current

generation ([Figure 4](#)) by the wildtype *ccm*, hybrid *ccmH*, *ccmH^N* strains in three-electrode

electrochemistry bioreactors. The wildtype *ccm* strain produced 11.47 ± 0.19 mA/m^2, in line with

our previous observations (Su et al., 2020). The hybrid *ccmH* produced 25.32 ± 3.61mA/m^2, a 121%

increase in current over the wildtype *ccm*. The *ccmH^N* produced only 3.42 ± 0.44 mA/m^2, about a

70% decrease relative to the wildtype *ccm*, similar to the no *mtr* strain previously reported

(TerAvest et al., 2014). Thus, we conclude that the hybrid *ccmH* strain matures more CymA and

produces higher current than the *E. coli ccm* system.

To probe the mechanism underlying the increase in current production by the hybrid *ccmH* strain,

we performed differential pulse voltammetry (DPV) ([Figure 5](#)) after strains reached the steady-

state current generation after 90 h chronoamperometry testing. We observed a single anodic peak

at redox potential (E_p) of +51 mV$_{SHE}$ in all three strains, which has been previously assigned to the

MtrCAB complex *(Okamoto et al., 2013)*. This observation confirms that all three strains use the

MtrCAB conduit for the current production. The peak height in hybrid *ccmH* strain reached 8.31

µA, about 118% higher than the 3.82 µA in wildtype *ccm* strain, while in the *ccmH^N* strain, it was

only 0.09 μA, about 2% of the wildtype. The ratio of the peak height for the hybrid *ccm* strain

relative to the wildtype *ccm* strain (2.18) closely mirrors the ratio of steady-state current for these

strains (2.21), indicating the increase in steady-state current arises from an increase in the flux of

electrons through the MtrCAB complex. Thus, the hybrid *ccm* strain produces more current by

more effectively maturing CymA, boosting its relative abundance, and increasing the electron flux

through the MtrCAB conduit.

3.5   Hybrid *ccmH* strain produces more current per cell via both direct and indirect EET

While DPV confirms that the increase in electron flux through the Mtr pathway increases the

current density, this increased flux could arise from an increase in the current per cell, an increase

in the number of viable cells, or an increase in both. To distinguish between these possibilities, we

measured both the current production and the number of viable cells using colony-forming unit

counting in the *ccmH^N*, hybrid *ccm*, and wildtype *ccm* strains ([Figure 6](#)A). The *ccmH^N* strain

produced near baseline current on a per cell basis (1.0 ± 0.005 fA·cell^{-1}), as expected. Interestingly,

the hybrid *ccmH* strains produced ~50% more current per cell than the wildtype *ccm* (12.3 ± 0.3

fA·cell^{-1} versus 8.3 ± 0.4 fA·cell^{-1}). These data confirm that the hybrid *ccmH* strain produces more

current at the single-cell level by direct EET.

In addition to direct EET, MR-1 (Brutinel and Gralnick, 2012; Coursolle et al., 2010) as a bound

cofactor (Edwards et al., 2015; Okamoto et al., 2014)or as a soluble mediator (Kotloski and

Gralnick, 2013). To test if the hybrid *ccmH* strain can also utilize flavin for EET, we added

exogenous flavins and measured the current density per cell. The addition of riboflavin to the

ccmH^N strain resulted in a meager current increase (0.02 ± 0.004 fA·cell^{-1}·μM^{-1}). As expected, the

current produced by both the wildtype *ccm* and hybrid *ccm* linearly increased with the exogenous

flavin concentration (Figure 6A). However, the hybrid *ccm* strain produced significantly more

current per µM exogenous flavin (0.32 ± 0.06 fA·cell^{-1}· µM^{-1}) than the wildtype *ccm* (0.16 ± 0.01

fA·cell^{-1}·µM^{-1}) (Figure 6A). These data indicate that the hybrid *ccm* strain produces more current

at the single-cell level via a flavin-mediated EET as well as a direct EET mechanism.

3.6   Direct and flavin-mediated EET in the hybrid *ccmH* strain is limited by CymA

Our data suggest that CymA amount limits electron flux and that the hybrid *ccm* strain

achieves higher current by enabling a greater electron flux from CymA into the same direct and

indirect EET routes found in the wildtype *ccm* strain. To quantitatively test these hypotheses, we

developed a quantitative model that relates EET in the hybrid *ccm* strain to EET in the wildtype

ccm strain and then tested whether this model could explain the current per cell measurement in

the hybrid *ccm* strain (Figure 6).

Explicitly, we modeled the overall EET current per cell (I_{strain}) as the sum of the direct EET

via Mtr conduit (I_{Mtr}) and flavin-mediated EET (I_{flavin}) (Figure 6B). We describe the flavin-

mediated EET as the product of the flavin concentration ($[flavin]$) and a constant of

proportionality (k_{strain}), (**Equation 1**):

$$I_{strain} = I_{strain,Mtr} + I_{strain,flavin} = I_{strain,Mtr} + k_{strain}[flavin] \qquad (1)$$

Assuming the concentration of CymA is rate-limiting in our system (Figure 3), both I_{Mtr} and I_{flavin}

should be dependent on the CymA concentration (**Equation 2**).

$$I_{strain,Mtr} = a[CymA]_{strain}; \; I_{strain,flavin} = b[CymA]_{strain}k_{strain}[flavin] \qquad (2)$$

By substituting and rearranging, we can then calculate EET current in the hybrid *ccmH* strain

(I_{hybrid}) as a ratio of the CymA concentration in the two strains and the current in the wildtype *ccm*

strain ($I_{WT,Mtr}$) with the equation below (**Equation 3**).

236
$$I_{hybrid} = \frac{[CymA]_{hybrid}}{[CymA]_{WT}} I_{WT,Mtr} + \frac{[CymA]_{hybrid}}{[CymA]_{WT}} k_{WT}[flavin] \qquad (3)$$

Thus, both the intercept and slope of the best-fit line for the hybrid *ccm* strain should be equal to

the intercept and slope best-fit line of the wildtype *ccm* strain times the ratio of CymA

concentration in these two strains.

Using the experimentally-determined values of $I_{WT,Mtr}$ (8.3 ± 0.4 fA·cell^{-1}), k_{WT} (0.16 ± 0.01

fA·cell^{-1}·µM^{-1}), and *[CymA]$_{hybrid}$/[CymA]$_{WT}$* = 1.6, we can predict the values of I_{hybrid} and compare

it to our experimentally-determined values (Figure 6A). Both the predicted $I_{hybrid, Mtr}$ and $I_{hybrid, flavin}$

values, 13.28 ± 0.64 fA·cell^{-1} and 0.26 ± 0.02 fA·cell^{-1}·µM^{-1}, respectively, are quite close to our

observation of the direct EET (12.3 ± 0.3 fA·cell^{-1}) and mediated EET (0.32 ± 0.06 fA·cell^{-1}·µM^{-}

^{1}) in hybrid *ccmH* strain. This result provides strong evidence that the hybrid *ccmH* strain's

improved bioelectrochemical performance arises mainly from the CymA expression increase and

underscores the importance of CymA in both direct EET and indirect EET.

4. DISCUSSION

In this study, we have shown that the C-terminal region of CcmH can be complemented by the

homologous region of CcmI from *S. oneidens* MR-1, which increases the bioelectronic

performance of Mtr-expressing *E. coli*. We found that the hybrid *ccm* can increase the overall cyt

c expression (Figure 2), with a preferred 60% increase in CymA expression (Figure 3). This

increase in CymA expression led to a 48% more EET current per bacterium and doubled the

electrical response to exogenous flavin ([Figure 5](#)), consistent with CymA being rate limiting in

EET. In the following, we discuss how these findings impact the development of biofuel cells and

biosensors.

First, our data show that the C-terminal region of CcmH is essential for cyt c-based EET.

Shewanella and *Geobacter* species are the dominant organisms used for EET applications, such as

MFCs. These microorganisms express a significant amount of cyt c to perform EET (Meyer et al.,

2004; Shi et al., 2009). As a result, there are increasing efforts focused on genetically introducing

cyt c to make new organisms capable of EET ((Schuergers et al., 2017; Sekar et al., 2016;

Thirumurthy and Jones, 2020), overexpressing cyt c to understand the structural basis of EET (M.

Edwards et al., 2019; M. J. Edwards et al., 2019; Wang et al., 2019), and making cyt c -based

conductive biomaterials for bioelectrical applications (Liu et al., 2020; Sun et al., 2018).

Understanding how cyt c mature can provide guidance on manipulating cyt c expression, which

will assist both fundamental studies and these applied efforts to increase the bioelectrical

performance of MFCs.

Second, we found that the hybrid *ccmH* strain generates higher electrical signals using both Mtr

and flavin-mediated EET. Increasing the EET efficiency of Mtr-expressing *E. coli* is of particular

interest for biosensing because it can increase the signal-to-noise ratio, enabling higher sensitivity

detection. In recent years, much progress has been made in engineering the electrical

communication between bacteria and electrodes, such as responding to specific chemicals using

chemical-sensitive promoters (Golitsch et al., 2013; Webster et al., 2014; Yang et al., 2018),

miniaturized reactors with optimized structure design (Zhou et al., 2017), and modified material-

bacteria interface (Zajdel et al., 2018). Enhancing the bioelectrical performance of Mtr-expression

E. coli offers a reliable chassis in biosensing applications.

Last, the cyt *c* stoichiometry shifts to more abundant in CymA, which guides us on how to build

a more efficient EET conduit for bioelectrical sensing. Our analysis of modeling the current

generation in hybrid *ccmH* strain, with and without exogenous flavin, together with the fact that

CymA can only carry 40% (fewer heme groups per protein) as many electrons as MtrA or MtrC,

highly suggests that electron transfer through CymA is a bottleneck for EET in the Mtr-expression

E. coli. Future work will focus on increasing the electron flux between menaquinone and the

MtrCAB complex to overcome this rate-limiting step. Also, the electrical response we observed

which is linear to both CymA amount and flavin concentration, could expand the application

scenarios such as building an "AND" gate with dual inputs.

5. CONCLUSIONS

In summary, a hybrid *ccm* system altered the stoichiometry of the expressed cyt *c,* thus

accelerating the rate limiting step in EET and enhancing the bioelectronic performance of *E. coli*.

Our results suggest that increasing the electron flux from menaquinone to MtrCAB complex could

further enhance the bioelectrical performance of engineered *E. coli*. More broadly, our work

provides a new strategy to express cyt *c* that underlie microbial bioelectronics and opens new

possibilities for engineering more sophisticated and sensitive biosensors.

SUPPLEMENTARY DATA

The Supporting Information is available online.

ABBREVIATIONS

cyt *c*, EET, ccm, MFC.

AUTHOR CONTRIBUTIONS

L.S. contributed to strains construction, design of the study, conducting experiments, analysis of the data, and writing of the manuscript. T.F. contributed to strain construction, design of the study, conducting experiments, analysis of the data, review and editing the manuscript. C.M.A-F. contributed to the design of the study, analysis of the data, and writing of the manuscript.

ACKNOWLEDGMENT

We thank Dr. Shuai Xu and Prof. Mohamed Y. El-Naggar, the University of Southern California for discussions on the DPV experiments. Work at the Molecular Foundry was supported by the Office of Science, Office of Basic Energy Sciences, of the U.S. Department of Energy [DE-AC02-05CH11231]. This work was supported by the Office of Naval Research, [N0001418IP00037]; and the China Scholarship Council [201606090098 to L.S.].

REFERENCES

Brutinel, E.D., Gralnick, J.A., 2012. Shuttling happens: soluble flavin mediators of extracellular electron transfer in Shewanella. Appl. Microbiol. Biotechnol. 93, 41–48. https://doi.org/10.1007/s00253-011-3653-0

Cheng, Z., Xiong, J., Min, D., Cheng, L., Liu, D., Li, W., Jin, F., Yang, M., Yu, H., 2020. Promoting bidirectional extracellular electron transfer of Shewanella oneidensis MR-1 for hexavalent chromium reduction via elevating intracellular cAMP level. Biotechnol. Bioeng. https://doi.org/10.1002/bit.27305

Chouler, J., Cruz-Izquierdo, Á., Rengaraj, S., Scott, J.L., Di Lorenzo, M., 2018. A screen-printed paper microbial fuel cell biosensor for detection of toxic compounds in water. Biosens. Bioelectron. 102, 49–56. https://doi.org/10.1016/j.bios.2017.11.018

Clarke, T.A., Edwards, M.J., Gates, A.J., Hall, A., White, G.F., Bradley, J., Reardon, C.L., Shi, L., Beliaev, A.S., Marshall, M.J., Wang, Z., Watmough, N.J., Fredrickson, J.K., Zachara, J.M., Butt, J.N., Richardson, D.J., 2011. Structure of a bacterial cell surface decaheme electron conduit. Proc. Natl. Acad. Sci. U. S. A. 108, 9384–9389. https://doi.org/10.1073/pnas.1017200108

Coursolle, D., Baron, D.B., Bond, D.R., Gralnick, J.A., 2010. The Mtr respiratory pathway is essential for reducing flavins and electrodes in Shewanella oneidensis. J. Bacteriol. 192, 467–474. https://doi.org/10.1128/JB.00925-09

Edwards, M.J., Richardson, D.J., Paquete, C.M., Clarke, T.A., 2019. Role of multiheme cytochromes involved in extracellular anaerobic respiration in bacteria. Protein Sci. https://doi.org/10.1002/pro.3787

Edwards, M.J., White, G.F., Norman, M., Tome-Fernandez, A., Ainsworth, E., Shi, L., Fredrickson, J.K., Zachara, J.M., Butt, J.N., Richardson, D.J., Clarke, T.A., 2015. Redox Linked Flavin Sites in Extracellular Decaheme Proteins Involved in Microbe-Mineral Electron Transfer. Sci. Rep. 5, 11677. https://doi.org/10.1038/srep11677

Edwards, M., White, G., Butt, J., Richardson, D.J., Clarke, T.A., 2019. The Structure of a Biological

Insulated Transmembrane Molecular Wire. https://doi.org/10.2139/ssrn.3445677

Fabianek, R.A., Hofer, T., Thöny-Meyer, L., 1999. Characterization of the Escherichia coli CcmH protein

reveals new insights into the redox pathway required for cytochrome c maturation. Arch. Microbiol.

171, 92–100. https://doi.org/10.1007/s002030050683

Fourmond, V., Hoke, K., Heering, H.A., Baffert, C., Leroux, F., Bertrand, P., Léger, C., 2009. SOAS: a

free program to analyze electrochemical data and other one-dimensional signals.

Bioelectrochemistry 76, 141–147. https://doi.org/10.1016/j.bioelechem.2009.02.010

Goldbeck, C.P., Jensen, H.M., TerAvest, M.A., Beedle, N., Appling, Y., Hepler, M., Cambray, G.,

Mutalik, V., Angenent, L.T., Ajo-Franklin, C.M., 2013. Tuning promoter strengths for improved

synthesis and function of electron conduits in Escherichia coli. ACS Synth. Biol. 2, 150–159.

https://doi.org/10.1021/sb300119v

Golitsch, F., Bücking, C., Gescher, J., 2013. Proof of principle for an engineered microbial biosensor

based on Shewanella oneidensis outer membrane protein complexes. Biosens. Bioelectron. 47, 285–

291. https://doi.org/10.1016/j.bios.2013.03.010

Gul, M.M., Ahmad, K.S., 2019. Bioelectrochemical systems: Sustainable bio-energy powerhouses.

Biosens. Bioelectron. 142, 111576. https://doi.org/10.1016/j.bios.2019.111576

Harris, H.W., Sánchez-Andrea, I., McLean, J.S., Salas, E.C., Tran, W., El-Naggar, M.Y., Nealson, K.H.,

2017. Redox Sensing within the Genus Shewanella. Front. Microbiol. 8, 2568.

https://doi.org/10.3389/fmicb.2017.02568

Jensen, H.M., TerAvest, M.A., Kokish, M.G., Ajo-Franklin, C.M., 2016. CymA and Exogenous Flavins

Improve Extracellular Electron Transfer and Couple It to Cell Growth in Mtr-Expressing Escherichia

coli. ACS Synth. Biol. 5, 679–688. https://doi.org/10.1021/acssynbio.5b00279

Kotloski, N.J., Gralnick, J.A., 2013. Flavin Electron Shuttles Dominate Extracellular Electron Transfer by

Shewanella oneidensis. mBio. https://doi.org/10.1128/mbio.00553-12

Liu, X., Gao, H., Ward, J.E., Liu, X., Yin, B., Fu, T., Chen, J., Lovley, D.R., Yao, J., 2020. Power

generation from ambient humidity using protein nanowires. Nature. https://doi.org/10.1038/s41586-

020-2010-9

Logan, B.E., 2009. Exoelectrogenic bacteria that power microbial fuel cells. Nature Reviews

Microbiology. https://doi.org/10.1038/nrmicro2113

Meyer, T.E., Tsapin, A.I., Vandenberghe, I., de Smet, L., Frishman, D., Nealson, K.H., Cusanovich,

M.A., van Beeumen, J.J., 2004. Identification of 42 possible cytochrome C genes in the Shewanella

oneidensis genome and characterization of six soluble cytochromes. OMICS 8, 57–77.

https://doi.org/10.1089/153623104773547499

Min, D., Cheng, L., Zhang, F., Huang, X.-N., Li, D.-B., Liu, D.-F., Lau, T.-C., Mu, Y., Yu, H.-Q., 2017.

Enhancing Extracellular Electron Transfer of Shewanella oneidensis MR-1 through Coupling

Improved Flavin Synthesis and Metal-Reducing Conduit for Pollutant Degradation. Environmental

Science & Technology. https://doi.org/10.1021/acs.est.6b04640

Okamoto, A., Hashimoto, K., Nealson, K.H., Nakamura, R., 2013. Rate enhancement of bacterial

extracellular electron transport involves bound flavin semiquinones. Proc. Natl. Acad. Sci. U. S. A.

110, 7856–7861. https://doi.org/10.1073/pnas.1220823110

Okamoto, A., Nakamura, R., Hashimoto, K., 2011. In-vivo identification of direct electron transfer from

Shewanella oneidensis MR-1 to electrodes via outer-membrane OmcA–MtrCAB protein complexes.

Electrochim. Acta 56, 5526–5531. https://doi.org/10.1016/j.electacta.2011.03.076

Okamoto, A., Nakamura, R., Nealson, K.H., Hashimoto, K., 2014. Bound Flavin Model Suggests Similar

Electron-Transfer Mechanisms in Shewanella and Geobacter. CHEMELECTROCHEM 1, 1808–

1812. https://doi.org/10.1002/celc.201402151

Robertson, I.B., Stevens, J.M., Ferguson, S.J., 2008. Dispensable residues in the active site of the

cytochrome c biogenesis protein CcmH. FEBS Lett. 582, 3067–3072.

https://doi.org/10.1016/j.febslet.2008.07.052

Rosenbaum, M.A., Bar, H.Y., Beg, Q.K., Segrè, D., Booth, J., Cotta, M.A., Angenent, L.T., 2012.

Transcriptional analysis of Shewanella oneidensis MR-1 with an electrode compared to Fe(III)citrate

or oxygen as terminal electron acceptor. PLoS One 7, e30827.

https://doi.org/10.1371/journal.pone.0030827

Sanders, C., Turkarslan, S., Lee, D.-W., Daldal, F., 2010. Cytochrome c biogenesis: the Ccm system.

Trends Microbiol. 18, 266–274. https://doi.org/10.1016/j.tim.2010.03.006

Santoro, C., Arbizzani, C., Erable, B., Ieropoulos, I., 2017. Microbial fuel cells: From fundamentals to

applications. A review. J. Power Sources 356, 225–244.

https://doi.org/10.1016/j.jpowsour.2017.03.109

Schuergers, N., Werlang, C., Ajo-Franklin, C.M., Boghossian, A.A., 2017. A Synthetic Biology

Approach to Engineering Living Photovoltaics. Energy Environ. Sci. 10, 1102–1115.

https://doi.org/10.1039/C7EE00282C

Sekar, N., Jain, R., Yan, Y., Ramasamy, R.P., 2016. Enhanced photo-bioelectrochemical energy

conversion by genetically engineered cyanobacteria. Biotechnol. Bioeng. 113, 675–679.

https://doi.org/10.1002/bit.25829

Shi, L., Dong, H., Reguera, G., Beyenal, H., Lu, A., Liu, J., Yu, H.-Q., Fredrickson, J.K., 2016.

Extracellular electron transfer mechanisms between microorganisms and minerals. Nat. Rev.

Microbiol. 14, 651–662. https://doi.org/10.1038/nrmicro.2016.93

Shi, L., Richardson, D.J., Wang, Z., Kerisit, S.N., Rosso, K.M., Zachara, J.M., Fredrickson, J.K., 2009.

The roles of outer membrane cytochromes of Shewanella and Geobacter in extracellular electron

transfer. Environ. Microbiol. Rep. 1, 220–227. https://doi.org/10.1111/j.1758-2229.2009.00035.x

Slate, A.J., Whitehead, K.A., Brownson, D.A.C., Banks, C.E., 2019. Microbial fuel cells: An overview of

current technology. Renewable Sustainable Energy Rev. 101, 60–81.

https://doi.org/10.1016/j.rser.2018.09.044

Sturm-Richter, K., Golitsch, F., Sturm, G., Kipf, E., Dittrich, A., Beblawy, S., Kerzenmacher, S.,

Gescher, J., 2015. Unbalanced fermentation of glycerol in Escherichia coli via heterologous

production of an electron transport chain and electrode interaction in microbial electrochemical cells.

Bioresour. Technol. 186, 89–96. https://doi.org/10.1016/j.biortech.2015.02.116

Su, L., Ajo-Franklin, C.M., 2019. Reaching full potential: bioelectrochemical systems for storing

renewable energy in chemical bonds. Curr. Opin. Biotechnol. 57, 66–72.

https://doi.org/10.1016/j.copbio.2019.01.018

Su, L., Fukushima, T., Prior, A., Baruch, M., Zajdel, T.J., Ajo-Franklin, C.M., 2020. Modifying

cytochrome c maturation can increase bioelectronic performance of engineered Escherichia coli.

ACS Synth. Biol. 9, 115–124. https://doi.org/10.1021/acssynbio.9b00379

Sun, Y.-L., Tang, H.-Y., Ribbe, A., Duzhko, V., Woodard, T.L., Ward, J.E., Bai, Y., Nevin, K.P.,

Nonnenmann, S.S., Russell, T., Emrick, T., Lovley, D.R., 2018. Conductive Composite Materials

Fabricated from Microbially Produced Protein Nanowires. Small 14, e1802624.

https://doi.org/10.1002/smll.201802624

TerAvest, M.A., Zajdel, T.J., Ajo-Franklin, C.M., 2014. The Mtr pathway of shewanella oneidensis MR-1

couples substrate utilization to current production in escherichia coli. ChemElectroChem 1, 1874–

1879.

Thirumurthy, M.A., Jones, A.K., 2020. Geobacter cytochrome OmcZs binds riboflavin: implications for

extracellular electron transfer. Nanotechnology 31, 124001. https://doi.org/10.1088/1361-

6528/ab5de6

Verissimo, A.F., Yang, H., Wu, X., Sanders, C., Daldal, F., 2011. CcmI subunit of CcmFHI heme ligation

complex functions as an apocytochrome c chaperone during c-type cytochrome maturation. J. Biol.

Chem. 286, 40452–40463. https://doi.org/10.1074/jbc.M111.277764

Wang, F., Gu, Y., O'Brien, J.P., Yi, S.M., Yalcin, S.E., Srikanth, V., Shen, C., Vu, D., Ing, N.L.,

Hochbaum, A.I., Egelman, E.H., Malvankar, N.S., 2019. Structure of Microbial Nanowires Reveals

Stacked Hemes that Transport Electrons over Micrometers. Cell 177, 361–369.e10.

https://doi.org/10.1016/j.cell.2019.03.029

Wang, X., Gao, N., Zhou, Q., 2013. Concentration responses of toxicity sensor with Shewanella

oneidensis MR-1 growing in bioelectrochemical systems. Biosens. Bioelectron. 43, 264–267.

https://doi.org/10.1016/j.bios.2012.12.029

Webster, D.P., TerAvest, M.A., Doud, D.F.R., Chakravorty, A., Holmes, E.C., Radens, C.M., Sureka, S.,

Gralnick, J.A., Angenent, L.T., 2014. An arsenic-specific biosensor with genetically engineered

Shewanella oneidensis in a bioelectrochemical system. Biosens. Bioelectron. 62, 320–324.

https://doi.org/10.1016/j.bios.2014.07.003

West, E.A., Jain, A., Gralnick, J.A., 2017. Engineering a Native Inducible Expression System in

Shewanella oneidensis to Control Extracellular Electron Transfer. ACS Synth. Biol. 6, 1627–1634.

https://doi.org/10.1021/acssynbio.6b00349

Xu, S., Jangir, Y., El-Naggar, M.Y., 2016. Disentangling the roles of free and cytochrome-bound flavins

in extracellular electron transport from Shewanella oneidensis MR-1. Electrochim. Acta 198, 49–55.

https://doi.org/10.1016/j.electacta.2016.03.074

Yang, Y., Zhou, H., Mei, X., Liu, B., Xing, D., 2018. Dual-Edged Character of Quorum Sensing

Signaling Molecules in Microbial Extracellular Electron Transfer. Front. Microbiol. 9, 1924.

https://doi.org/10.3389/fmicb.2018.01924

Zajdel, T.J., Baruch, M., Méhes, G., Stavrinidou, E., Berggren, M., Maharbiz, M.M., Simon, D.T., Ajo-

Franklin, C.M., 2018. PEDOT:PSS-based Multilayer Bacterial-Composite Films for Bioelectronics.

Sci. Rep. 8, 15293. https://doi.org/10.1038/s41598-018-33521-9

Zheng, X.-M., Hong, J., Li, H.-Y., Lin, D.-H., Hu, H.-Y., 2012. Biochemical properties and catalytic

domain structure of the CcmH protein from Escherichia coli. Biochim. Biophys. Acta 1824, 1394–

1400. https://doi.org/10.1016/j.bbapap.2012.06.017

Zhou, A.Y., Baruch, M., Ajo-Franklin, C.M., Maharbiz, M.M., 2017. A portable bioelectronic sensing

system (BESSY) for environmental deployment incorporating differential microbial sensing in

miniaturized reactors. PLoS One 12, e0184994. https://doi.org/10.1371/journal.pone.0184994

 FIGURE CAPTIONS

Figure 1. A schematic illustration depicting the different *ccmH* genes for cytochrome *c*

expression in engineered *E. coli*. We replaced the coding sequence of *ccmH* from the *E. coli*

ccmABCDEFGH operon with three other variants. These variants coded for CcmHI from MR-1

(*ccmHI*), a fusion between the N terminal region of *E. coli ccmH* and the C terminal region of

CcmI from MR-1 (hybrid *ccmH*), and a control that removed the C terminal region of MR-1 CcmI

(*ccmH^N*). Black and red regions indicate DNA regions that code for proteins from *E. coli* and MR-

1, respectively.

Figure 2. The hybrid *ccmH* strain expresses more cytochromes *c*. Relative redness as a function

of relative promoter strength, i.e., induction level, for *E. coli* expressing *cymAmtrCAB* with

different *ccm* operons. The hybrid *ccmH* strain has a higher level of overall cyt *c* expression than

wildtype *ccm*. Results are representative of three independent experiments.

Figure 3. The hybrid *ccm* system increases the CymA expression in *E. coli*. The relative density

of enhanced chemiluminescence (ECL) bands arising from three cyt *c* heterologously expressed

in the hybrid *ccm* and wildtype *ccm* strains. The cyt *c* expression in negative control no *ccm* strain

was below the limit of detection. Results are representative of three independent experiments.

Figure 4. The hybrid *ccmH* strain produces a higher current. The current generation measured

by chronoamperometry in the bioreactors showing hybrid *ccmH* strain produces more current than

the wildtype *ccm*, while the negative control strain *ccmH^N* produces less current. Results are

representative of three independent experiments. The dotted arrow shows the injecting time of

bacteria.

Figure 5. The hybrid *ccmH* strain shows higher redox activity through cyt *c*. Differential pulse

voltammetry (DPV) shows that the hybrid *ccmH* strain has a higher redox peak at the cyt *c* related

redox reaction potential (+51 mV vs. SHE). DPV signal smoothing and background deduction by

QSoas Version 2.0 (Fourmond et al., 2009).

Figure 6. Current production by the hybrid *ccmH* strain is more sensitive to exogenous

flavins. (A) The hybrid *ccmH* strain (red) has a more rapid and higher current response as a

function of extra riboflavin concentration, compared to the wildtype *ccm* strain (gray), and *ccmH^N*

strain (yellow). Results are representative of two independent experiments. Linear fit analysis

(with Y error) is performed by OriginPro 2019b (OriginLab Corporation). (B) A schematic

depicting the extracellular electron transfer via Mtr conduit. MtrCAB complex receives the

electrons from CymA (I_{strain}), then delivers to the electrode through either MtrC (I_{Mtr}) or flavin

(I_{flavin}). (OM: outer membranes, IM: Inner membranes).

Figure1

Figure2

Figure3

Figure4

Figure6

(A)

(B)