Structural basis to design multi-epitope vaccines against Novel Coronavirus 19 (COVID19) infection, the ongoing pandemic emergency: an in silico approach

Sukrit Srivastava*[a,b], Sonia Verma[c], Mohit Kamthania[d], Rupinder Kaur[e], Ruchi Kiran Badyal[f], Ajay Kumar Saxena[b], Ho-Joon Shin[g], Kailash Pandey[c], Michael Kolbe[h,i]

a   Department of Biotechnology, Institute of Bio-Medical Education and
        Research, Mangalayatan University, 202146 Aligarh, India.
b   Molecular Medicine Lab., School of Life Science, Jawaharlal Nehru
        University, 110067 New Delhi, India.
c   National Institute of Malaria Research-Indian Council of Medical Research,
        New Delhi, India
d   Department of Biotechnology, Institute of Applied Medicines and Research,
        201206 Ghaziabad, India.
e   Department of Chemistry, Guru Nanak Dev University, Amritsar, India
f   Department of economics, Mangalayata University, India
g   TIDCL, Ajou University, South Korea
h   Department for Structural Infection Biology, Centre for Structural Systems
        Biology (CSSB) & Helmholtz-Centre for Infection Research, Notkestraße
        85, 22607 Hamburg, Germany.
i   Faculty of Mathematics, Informatics and Natural Sciences, University of
        Hamburg, Rothenbaumchaussee 19, 20148 Hamburg, Germany.

*Correspondence should be addressed to S.S.
(sukrit.srivastav@mangalayatan.edu.in; srivastav.sukrit@gmail.com)

Abstract

The 2019 novel coronavirus (COVID19 / Wuhan coronavirus), officially named as Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), is a positive-sense single-stranded RNA coronavirus. SARS-CoV-2 causes the contagious COVID19 disease also known as 2019-nCoV acute respiratory disease and has led to the ongoing 2019–20 pandemic COVID19 outbreak. The effective

countermeasures against SARS-CoV-2 infection requires the design and

development of specific and effective vaccine candidate. In the present study we

have screened 38 CTL, 42 HTL and 12 B cell epitopes by utilizing different in

silico tools. The screened epitopes were further validated for their binding with

their respective HLA allele binders and TAP (Transporter associated with antigen

processing) molecule by molecular docking. The screened epitopes were further

utilized to design novel two multi-epitope vaccines (MEVs) composed of CTL,

HTL and B cell epitopes overlaps with potential to elicit humoral as well as

cellular immune response against SARS-CoV-2. To enhance the immune

response for our vaccine design, truncated (residues 10-153) Onchocerca

volvulus activation-associated secreted protein-1 (Ov-ASP-1) has been utilized

as an adjuvant at N terminal of both the MEVs. Further molecular models for both

the MEVs were prepared and validated for their stable molecular interaction with

Toll-Like Receptor 3 (TLR 3). The codon-optimized cDNA of both the MEVs were

further analyzed for their potential of high level of expression in Human cell line.

The present study proposes *in-silico* validated design of two MEVs against

SARS-CoV-2 to elicit humoral as well as cellular immune response.

CTL Multi-epitope COVID19 vaccine - Toll Like Receptor 3 complex

CTL Multi-epitope COVID19 vaccine - Toll Like Receptor 3 complex

Graphical abstract: The designed CTL (Cytotoxic T lymphocyte) and HTL (Helper T lymphocyte) multi-epitope vaccine (MEV) against COVID19 infection. Both the CTL and HTL MEVs models show a very stable and very fitting conformational complex formation tendency with the Toll like receptor 3. CTL and HTL MEVs: ribbon; Toll like receptor 3: gray cartoon; Adjuvant [truncated (residues 10-153) Onchocerca volvulus activation-associated secreted protein-1]: orange ribbon; Epitopes: cyan ribbons regions; 6xHis Tag: magenta ribbon region

Abbreviations: Antigen-presenting cell (APCs), Codon Adaptation Index (CAI), Coronavirus ID 19 (COVID19), Coronavirus (CoV), Coverage (Cov), Cryo-Electron Microscopy (Cryo-EM), Cytotoxic T lymphocyte (CTL), ectodomain (ECD), Envelope Protein (E Protein), endoplasmic reticulum (ER), European Bioinformatics Institute (EBI), Nucleocapsid Protein (N Protein), Ope reading Frame (ORF), global distance test (GDT), Grand average of hydropathicity (GRAVY), Surface protein (S Protein), half maximal inhibitory concentration (IC50), Helper T lymphocyte (HTL), Human Leukocyte Antigen (HLA), Immune Epitope Database (IEDB), Immuno Polymorphism Database (IPD), Interferon Gama (IFN-γ), International ImMunoGeneTics project (IMGT), major histocompatibility complex (MHC), Membrane Protein (M Protein), Motif-EmeRging with ClassesIdentification (MERCI), Multi-epitope Vaccine (MEV), Multiple Sequence Alignment (MSA), non-structural protein (nsp), National Center for Biotechnology Information (NCBI), peptide-MHC (pMHC), Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), Stabilization Matrix alignment method (SMM), Transmembrane (TM), Protein Data Bank (PDB), Qualitative Model Energy ANalysis (QMEAN), root mean square deviation (RMSD), Support Vector Machine (SVM), Toll-Like Receptor 3 (TLR3), Transporter associated with antigen processing (TAP), un-normalized global distance test (uGDT)

Introduction

The novel coronavirus (COVID19), officially named as Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has caused the ongoing outbreak of

severe form of flu leading to death with mortality rate of 3.4 %. The SARS-CoV-2

is a novel coronavirus associated with a respiratory disease initiated from the

Wuhan of Hubei province, China. The disease is highly contagious and has

spread to 182 countries/ Territories since its outbreak in China in December 2019

till 21st of march 2020. Worldwide, as of 21st March 2020 the total confirmed

cases has been reported to be 2,66,073 and total death count reported is 11,184

(WHO Situation report 21st March 2020). Over all the SARS-CoV-2 infection has

put a global emergency condition. The economic impact of COVID-19 is even

more harsh and has put the world on economic risk. As of 9th March, the

downside scenario sees a \$2 trillion shortfall in global income with a \$220 billion

hit to developing countries. The COVID-19 shock will cause a recession in

several countries and depress global annual growth this year to 2.5 per cent

lower, the recessionary threshold for the world economy (UNCTAD report 9th

March 2020).

The infection mechanism and pathogenesis of SARS-CoV-2 is largely

unknown yet. The proteome of SARS-CoV-2 is composed of 11 structural and

non-structural proteins. These includes polyprotein (ORF1ab), Surface protein (S

Protein), ORF3, Envelope Protein (E Protein), Membrane Protein (M Protein),

ORF6, ORF7a, ORF7b, ORF8, Nucleocapsid Protein (N Protein), and ORF10

(NCBI protein sequence database). The actual function and pathogenic or

proliferative role of these SARS-CoV-2 coronavirus proteins are largely not

known yet.

The SARS-CoV-2 coronavirus polyprotein (ORF1ab) with length of 7,096 amino acid (aa) is composed of 16 different expressed protein viz. leader protein (nsp1, location: 1-180 aa); nsp2 (location: 181-818); nsp3 (former nsp1, carry conserved domains - N-terminal acidic, predicted phosphoesterase, papain-like proteinase, Y-domain, transmembrane domain 1 (TM1) and adenosine diphosphate-ribose 1"-phosphatase, location: 819-2,763); nsp4 (contains transmembrane domain 2 (TM2), location: 2,764-3,263); 3C-like proteinase (nsp5, main proteinase, mediates cleavages downstream of nsp4, location: 3,264-3,569); nsp6 (putative transmembrane domain, location: 3,570-3,859); nsp7 (location: 3,860-3,942); nsp8 (location: 3,943-4,140); nsp9 (ssRNA-binding protein, location: 4,141-4,253); nsp10 (formerly known as growth-factor-like protein, location: 4,254-4,392); nsp11 (location: 4,393-4,405); RNA-dependent RNA polymerase (nsp12, location: 4,393-5,324); Helicase (nsp13; zinc-binding domain, NTPase/helicase domain, RNA 5'-triphosphatase, location: 5,325-5,925); 3'-to-5' exonuclease (nsp14, location: 5,926-6,452); endo RNAse (nsp15, location: 6,453-6,798); and 2'-O-ribose methyltransferase (nsp16; location: 6,799-7,096).

The SARS-CoV-2 coronavirus surface glycoprotein (S Protein) is a structural protein and acts as spike protein, its location is 21,563-25,384 aa, and length is 1273 aa); The SARS-CoV-2 coronavirus ORF3a protein has location 25,393-26,220 aa and length 275 aa. The SARS-CoV-2 coronavirus envelope protein (E Protein) (ORF4) is a structural protein and has location 26,245-26,472 aa and length 75 aa. The SARS-CoV-2 coronavirus membrane glycoprotein (M

Protein) (ORF5) is a structural protein and has location 26,523-27,191aa and

length 222 aa. The SARS-CoV-2 coronavirus ORF6 protein has location 27,202-

27,387 aa and length 61 aa. The SARS-CoV-2 coronavirus ORF7a protein has

location 27,394-27,759 aa and length 121 aa. The SARS-CoV-2 coronavirus

ORF7b protein has location 27,756-27,887 aa and length 43 aa. The SARS-CoV-

2 coronavirus ORF8 protein has location 27,894-28,259 aa and length 121 aa.

The SARS-CoV-2 coronavirus nucleocapsid phosphoprotein (N Protein) (ORF9)

is a structural protein with location 28,274-29,533 aa and has length of 419 aa.

The SARS-CoV-2 coronavirus ORF10 protein has location 29,558-29,674 aa and

length 38 aa (NCBI protein sequence database).

Although the exact mechanism and roles of all the above-mentioned

proteins of SARS-CoV-2 coronavirus proteome is not well known yet, but these

proteins could act as potential vaccine candidate against the SARS-CoV-2

coronavirus infection. In the present study we have screened highly potential

epitopes from all the above-mentioned protein and further we have also designed

and proposed CTL (Cytotoxic T lymphocyte) and HTL (Helper T lymphocyte)

multi-epitope based vaccine candidates against the SARS-CoV-2 coronavirus

infection.

Methodology

In the present study on SARS-CoV-2 coronavirus we have screened potential epitopes and have designed and proposed two multi-epitope vaccines (MEVs) composed of screened CTL (Cytotoxic T lymphocyte) and HTL (Helper T lymphocyte) epitopes with overlapping regions of B cell epitopes. Hence the proposed MEVs are supposed to have potential to elicit both the humoral as well as cellular immune response. To enhance immune response, truncated (residues 10-153) Onchocerca volvulus activation-associated secreted protein-1 (Ov-ASP-1) has been utilized as adjuvant at N-terminal of both the MEVs. The truncated Ov-ASP-1 was chosen due to its potential to activate antigen-processing cells (APCs) (MacDonald et al., 2005; Guo et al., 2015; He et al., 2009). All SARS-CoV-2 coronavirus proteins mentioned in the introduction, were utilized to screen the potential CTL, HTL and B cell epitopes. The screened epitopes were further studied for overlapping consensus regions amongst them. The epitopes showing overlapping regions in partial or complete were chosen for detailed further studies.

The chosen CTL and HTL epitopes were analyzed for their molecular interaction with their respective HLA allele binders. Moreover, the chosen CTL epitopes were also analyzed for their molecular interaction with TAP (Transporter associated with antigen processing) transporter cavity to observe their smooth passage from cytoplasm to endoplasmic reticulum lumen (Oldham et al., 2016; Abele et al., 2004). The tertiary model for both MEVs were generated and refined.

Both the MEVs models were further utilized to screened B Cell linear and

discontinuous epitopes as well as IFN-γ inducing epitopes.

The molecular signaling by multiple TLRs, is an essential component of

the innate immune response against SARS-CoV-2 coronavirus. Since the rOv-

ASP-1 primarily binds APCs among human PBMCs and trigger pro-inflammatory

cytokine production via Toll-like receptor 3 (TLR3), hence both the CTL and HTL

MEV models were further analyzed for their molecular interaction with the TLR-3

by molecular docking studies (Antoniou et al., 2003; Delneste et al., 2007; Totura

et al., 2015; Farina et al., 2005). Further the codon optimized cDNA of both the

MEVs were analyzed to have high level of expression in mammalian cell line

(human), which would facilitate *in-vivo* expression, experimentation and trials

(Fig.1).

Figure 1. Workflow chart.

Screening of Potential Epitopes

T cell Epitope Prediction

Screening of Cytotoxic T lymphocyte (CTL) Epitope. The screening of Cytotoxic T lymphocyte epitopes was performed by the IEDB (Immune Epitope Database) tools "MHC-I Binding Predictions" (http://tools.iedb.org/mhci/) and "MHC-I Processing Predictions" (http://tools.iedb.org/processing/) (Tenzer et al., 2005; Peters et al., 2003; Hoof et al., 2009). Both the tools use six different methods (viz. Consensus, NN-align, SMM-align, Combinatorial library, Sturniolo and NetMHCIIpan) and generate "Percentile rank" and a "total score" respectively.

The screening is based on the total amount of cleavage site in the protein. TAP score estimates an effective log -(IC50) values (half maximal inhibitory concentration (IC50)) for binding to TAP of a peptide or its N-terminal prolonged precursors. The MHC binding prediction score is -log(IC50) values for binding to MHC of a peptide (Calis et al., 2013). The IC(50) (nM) value for each epitope and MHC allele binding pairs were also obtained by this IEDB tool. Epitopes having high, intermediate, and least affinity of binding to their HLA allele binders have IC50 values < 50 nM, < 500 nM and < 5000 nM respectively.

Immunogenicity of all the screened CTL epitopes was also obtained by using "MHC I Immunogenicity" tool of IEDB (http://tools.iedb.org/immunogenicity/) with all the parameters set to default analyzing 1st, 2nd, and C-terminus amino acids of the given screened epitope (Calis et al., 2013). The tool predicts the immunogenicity of a given peptide-MHC (pMHC) complex on the basis of the

physiochemical properties of constituting amino acid and their position within the

peptide sequence.

Screening of Helper T lymphocyte (HTL) Epitopes. To screen out the Helper T

lymphocyte epitopes from SARS-CoV-2 proteins, the IEDB tool "MHC-II Binding

Predictions" (http://tools.iedb.org/mhcii/) was used. The tool generates

"Percentile rank" for each potential peptide. The lower the value of percentile,

higher would be the affinity. This percentile rank is generated by the combination

of three different methods viz. combinatorial library, SMM_align & Sturniolo; and

by comparing the score of the peptide against the scores of other random five

million 15-mer peptides of SWISSPROT database (Wang et al., 2010; Sidney et

al., 2008; Nielsen et al., 2007; Sturniolo et al., 1999). The rank from the

consensus of all three methods was generated by the median percentile rank of

the three methods.

Population Coverage by CTL and HTL epitopes. The "Population Coverage" tool

of IEDB (http://tools.iedb.org/population/) was used to elucidate the world human

population coverage by the shortlisted 38 CTL and 42 HTL epitopes derived from

nine SARS-CoV-2 proteins (Bui et al., 2006). The T cells recognize complex

between a specific major MHC molecule and a particular pathogen-derived

epitope. The given epitope will elicit a response only in an individual that express

an MHC molecule, which is capable of binding that particular epitope. This

denominated MHC restriction of T cell responses and the MHC polymorphism

provide basis for population coverage study. The MHC types are expressed at

dramatically different frequencies in different ethnicities. Hence a vaccine with

larger population coverage could be of greater importance (Sturniolo et al., 1999).

Clinical administration of multiple-epitopes involving both the CTL and the HTL

epitopes are predicted here to have a greater probability of larger human

population coverage worldwide.

B Cell Epitope Prediction

Sequence-based B Cell epitope prediction. Protein sequence-based method

"Bepipred Linear Epitope Prediction" was utilized to screen linear B cell epitopes

from eleven different SARS-CoV-2 proteins. The tool "B Cell Epitope Prediction

Tools" of IEDB server (http://tools.iedb.org/bcell/) was utilized. In this screening

the parameters such as hydrophilicity, flexibility, accessibility, turns, exposed

surface, polarity and antigenic propensity of the polypeptides is correlated with its

location in the protein. This allows the search for continuous epitopes prediction

from protein sequence. The prediction is base on the propensity scales for each

of the 20 amino acids. For a window size n, the i - (n-1)/2 neighboring residues

on each side of residue i are used to compute the score for the residue i. The

method "Bepipred Linear Epitope Prediction" utilized here is based on the

propensity scale method as well as the physiochemical properties of the given

antigenic sequence to screen potential epitopes (Larsen et al., 2006).

Characterisation of potential epitopes

Epitope conservation analysis. The shortlisted CTL, HTL and B cell epitopes screened from eleven SARS-CoV-2 proteins were analysed for the conservancy of their amino acid sequence by "Epitope Conservancy Analysis" tool (http://tools.iedb.org/conservancy/) of IEDB. The epitope conservancy is the number of protein sequences (retrieved from NCBI) that contain that particular epitope. The analysis was done against their entire respective source protein sequences of SARS-CoV-2 proteins retrieved from the NCBI protein database (Bui et al., 2007).

Epitope Toxicity prediction. The tool ToxinPred (http://crdd.osdd.net/raghava/toxinpred/multi_submit.php) was used to analyse the toxicity of shortlisted CTL, HTL and B cell epitopes. The tool allows to identifying highly toxic or non-toxic short peptides. The toxicity check analysis was done by the "SVM (Swiss-Prot) based" (support vector machine) method utilizing dataset of 1805 sequences as positive, 3593 negative sequences from Swissprot as well as an alternative dataset comprises the same 1805 positive sequences and 12541 negative sequences from TrEMBLE (Gupta et al., 2013).

Overlapping residue analysis. The overlapping residue analysis for the shortlisted 38 CTL, 42 HTL and the 12 B cell linear epitopes was performed by the Multiple Sequence Alignment (MSA) analysis by Clustal Omega tool (https://www.ebi.ac.uk/Tools/msa/clustalo/) of EBI (European Bioinformatics

Institute) (Sievers et al., 2011). The Clustal Omega multiple sequence alignment

tool virtually aligns any number of protein sequences and delivers an accurate

alignment.

Epitope selected for molecular interaction study with HLA allele and TAP

transporter. On the basis of the overlapping residue analysis of shortlisted CTL,

HTL and linear B cell epitopes few numbers of CTL and HTL epitopes were

chosen for further analysis. The chosen epitopes are encircled shown in Fig.2.

These epitopes were chosen on the basis of them having partial or full

overlapping sequence region amongst all three types of epitopes (CTL, HTL and

B Cell). The chosen epitopes were further analyzed for their interaction with their

respective HLA allele binders and TAP cavity interaction.

Molecular interaction analysis of selected epitopes with HLA allele and TAP

transporter.

Tertiary structure modelling of HLA alleles and selected T cell epitopes. The

Swiss-model was used for homology modelling of the HLA class I and II allele

binders of the chosen epitopes (Arnold et al., 2006). The amino acid sequence of

the HLA allele binders were retrieved from Immuno Polymorphism Database

(IPD-IMGT/HLA) (https://www.ebi.ac.uk/ipd/imgt/hla/allele.html). Templates for

homology modelling were chosen on the basis of highest amino acid sequence

similarity. All the generated HLA allele models had acceptable QMEAN value

(cutoff -4.0) (Supplementary table S1). The QMEAN value gives a composite

quality estimate involving both global as well as local analysis of the model

(Benkert et al., 2008).

The PEP-FOLD 2.0 a de novo structure prediction tool at RPBS Web

Portal   (https://mobyle.rpbs.univ-paris-diderot.fr/cgi-bin/portal.py#forms::PEP-

FOLD) was utilized to generate tertiary structures for the chosen CTL and HTL

epitopes (Shen et al., 2014).

Molecular interaction analysis of chosen CTL and HTL epitopes with HLA alleles.

The PatchDock tool (http://bioinfo3d.cs.tau.ac.il/PatchDock/) was utilized for *in-*

silico molecular docking study of the selected CTL and HTL epitopes with their

respective HLA class I and II allele binders (Bell et al., 2005; Duhovny et al.,

2002; Schneidman-Duhovny et al., 2005). PatchDock utilizes algorithm for

unbound (real life) docking of molecules for protein-protein complex formation.

The algorithm carries out the rigid docking, with the surface variability/flexibility

implicitly addressed through liberal intermolecular penetration. The algorithm

focuses on the (i) initial molecular surface fitting on localized, curvature based

surface patches (ii) use of Geometric Hashing and Pose Clustering for initial

transformation detection (iii) computation of shape complementarity utilizing the

Distance Transform (iv) efficient steric clash detection and geometric fit scoring

based on a multi-resolution shape representation and (v) utilization of biological

information by focusing on hot spot rich surface patches (Bell et al., 2005;

Duhovny et al., 2002; Schneidman-Duhovny et al., 2005).

Molecular interaction analysis of selected CTL epitopes with TAP transporter.

TAP transporter plays an important role in the presentation of CTL epitope. From

the cytosol after proteasome processing, the fragmented peptide of foreign

protein gets transported to endoplasmic reticulum (ER) through the TAP

transporter. From the ER these short peptides reach to the golgi bodies and then

get presented on the cell surface (Abele et al., 2004). Molecular interaction study

of the chosen CTL epitopes within the TAP transporter cavity was performed by

molecular docking study utilizing the PatchDock tool. For accurate prediction, the

cryo-EM structure of TAP transporter (PDB ID: 5u1d) was used by removing the

antigen from TAP cavity of the original structure (Oldham et al., 2016).

Design, characterisation and molecular interaction analysis of Multi-

Epitope Vaccines with immunological receptor.

Design of Multi-Epitope Vaccines. The screened and shortlisted high scoring 38

CTL and 42 HTL epitopes were utilized to design CTL and HTL Multi-Epitope

vaccines (Table 1 and 2). Short peptide EAAAK and GGGGS were used as rigid

and flexible linkers respectively (Fig.2). The GGGGS linker provides proper

conformational flexibility to the vaccine tertiary structure and hence facilitates

stable conformation to the vaccine. The EAAAK linker facilitates in domain

formation and hence facilitates the vaccine to obtain its final stable structure.

Truncated (residues 10-153) Onchocerca volvulus activation-associated secreted

protein-1 (Ov-ASP-1) has been utilized as an adjuvant at N terminal of both the

CTL and HTL MEVs (MacDonald et al., 2005; Guo et al., 2015; He et al., 2009;

Hu et al., 2004; Hajighahramani et al., 2017; Chen et al., 2013; Srivastava et al.,

2019; Srivastava et al., 2018).

B. HTL epitopes | **A. CTL epitopes**

Ov-ASP-1 (truncated (10-153)):
IVVAVTGYNCPGGKLTALERKKIVGQNNKYRSDLINGKL
KNRNGTYMPRGKNMLELTWDCKLESSAQRWANQCIFG
HSPRQQREGVGENVYAYWSSVSVEGLKKTAGTDAGKS
WWSKLPKLYENNPSNNMTWKVAGQGVLHFTQ

B column (HTL epitopes) — ORF1ab:
EAAAK
WFLAYILFTRFFYVL
GGGGS
FLAYILFTRFFYVLG
GGGGS
LAYILFTRFFYVLGL
GGGGS
AYILFTRFFYVLGLA
GGGGS
RAMPNMLRIMASLVL
GGGGS
AMPNMLRIMASLVLA
GGGGS
MPNMLRIMASLVLAR
GGGGS
PNMLRIMASLVLARK
GGGGS
QESPFVMMSAPPAQY
GGGGS
NMLRIMASLVLARKH
GGGGS
ESPFVMMSAPPAQYE
GGGGS
SPFVMMSAPPAQYEL
GGGGS
AIILASFSASTSAFV
GGGGS
IILASFSASTSAFVE
GGGGS
FAWWTAFVTNVNASS
GGGGS

Surface protein:
KTQSLLIVNNATNVV
GGGGS
TQSLLIVNNATNVVI
GGGGS
QSLLIVNNATNVVIK
GGGGS
SLLIVNNATNVVIKV
GGGGS
LLIVNNATNVVIKVC
GGGGS

ORF3a:
LLFVTVYSHLLLVAA
GGGGS
FVRATATIPIQASLP
GGGGS

Envelope protein:
VLLFLAFVVFLLVTL
GGGGS
LLFLAFVVFLLVTLA
GGGGS

Membrane protein:
GLMWLSYFIASFRLF
GGGGS
LMWLSYFIASFRLFA
GGGGS
LSYYKLGASQRVAGD
GGGGS

ORF6:
TFKVSIWNLDYIINL
GGGGS
FKVSIWNLDYIINLI
GGGGS
KVSIWNLDYIINLII
GGGGS

ORF7a:
IILFLALITLATCEL
GGGGS
ILFLALITLATCELY
GGGGS

ORF7b:
YLCFLAFLLFLVLIM
GGGGS
LCFLAFLLFLVLIML
GGGGS
CFLAFLLFLVLIMLI
GGGGS

ORF8:
CTQHQPYVVDDPCPI
GGGGS
HQPYVVDDPCPIHFY
GGGGS
QPYVVDDPCPIHFYS
GGGGS

Nucleocapsid phosphoprotein:
PQIAQFAPSASAFFG
GGGGS
IAQFAPSASAFFGMS
GGGGS
AQFAPSASAFFGMSR

ORF10:
YINVFAFPFTIYSLL
HHHHHH

A column (CTL epitopes):
EAAAK
SEMVMCGGSLY
GGGGS
FYWFFSNYLKR
GGGGS
ISNSWLMW
GGGGS
ETISLAGSYK
GGGGS
QEILGTVSW
GGGGS
STFNVPMEK
GGGGS
RMYIFFASFY
GGGGS
FLFVAAIFYL
GGGGS
RYFRLTLGVY
GGGGS
FLNGSCGSV
GGGGS
CTDDNALAY
GGGGS
CTDDNALAYY
GGGGS
MYKGLPWNVVR
GGGGS
SIINNTVYTK
GGGGS
LPVNVAFELW
GGGGS
DEWSMATYY
GGGGS
YILFTRFFYV
GGGGS
YIFFASFYYV
GGGGS
WTAGAAAYYV
GGGGS
FPNITNLCPF
GGGGS
NYNYLYRLFR
GGGGS
NYLYRLFR
GGGGS
YLYALVYFL
GGGGS
IPYNSVTSSI
GGGGS
LLFLAFVVF
GGGGS
LTALRLCAY
GGGGS
YFIASFRLFAR
GGGGS
ATSRTLSYYK
GGGGS
RTFKVSIW
GGGGS
AEILLIIMRTF
GGGGS
RARSVSPK
GGGGS
QLRARSVSPK
GGGGS
FLAFLLFLV
GGGGS
HFYSKWYIR
GGGGS
MEVTPSGTW
GGGGS
KPRQKRTAT
GGGGS
MGYINVFAF
GGGGS
GYINVFAFPF
HHHHHH

Figure 2. Design of SARS-CoV-2 Multi-Epitope Vaccine (MEVs). (A) CTL and

(B) HTL epitopes were linked by the short peptide linker 'GGGGS'. Truncated

(residues 10-153) Onchocerca volvulus activation-associated secreted protein-1

(Ov-ASP-1) has been utilized at the N terminal of both the MEVs. The short

peptide EAAAK was used to link Ov-ASP-1 at N terminal. Epitopes from different

proteins were colored in different colors. C terminal 6xHis is designed as His tag.

Characterisation of designed Multi-Epitope Vaccines

Physicochemical property analysis of designed MEVs. The ProtParam

(https://web.expasy.org/protparam/) tool was utilized to analyse the

physiochemical properties of the amino acid sequence of the designed CTL and

HTL MEVs (Gasteiger et al., 2005). The ProtParam analysis performs an

empirical investigation for the given query amino acid sequence. ProtParam

computes various physicochemical properties derived from a given protein

sequence.

Interferon gama inducing epitope prediction. From the designed amino acid

sequence of both the MEVs potential interferon gama (IFN-γ) epitopes were

screened by "IFNepitope" server

(http://crdd.osdd.net/raghava/ifnepitope/scan.php) using "Motif and SVM hybrid",

(MERCI: Motif-EmeRging and with Classes-Identification, and SVM: support

vector machine) method. The tool predicts peptides from protein sequences

having the capacity to induce IFN-gamma release from CD4+ T cells. This

module generates overlapping peptides from query sequence and predicts IFN-
gamma inducing peptides. For the screening, IEDB database with 3705 IFN-
gamma inducing and 6728 non-inducing MHC class II binders is utilized (Nagpal
et al., 2015; Dhanda et al., 2013).

MEVs allergenicity and antigenicity prediction. Both the designed MEVs were
further analysed for allergenicity and antigenicity prediction by utilizing the
AlgPred (http://crdd.osdd.net/raghava/algpred/submission.html) and the Vaxigen
(http://www.ddg-pharmfac.net/vaxijen/VaxiJen/VaxiJen.html) tools respectively
(Saha et al., 2006; Doytchinova and Flower 2007). The AlgPred prediction is
based on the similarity of already known epitope with any region of the submitted
protein. For the screening of allergenicity, the Swiss-prot dataset consisting of
101725 non-allergens and 323 allergens is utilized. The VaxiJen utilizes an
alignment-free approach, solely based on the physicochemical properties of the
query amino acid sequence. For prediction of antigenicity, the Bacterial, viral and
the tumour protein datasets are used to derive models for the prediction of whole
protein antigenicity. Every set consisted of known 100 antigens and 100 non-
antigens.

Tertiary structure modelling and refinement of MEVs. The tertiary structure of
both the designed CTL and HTL MEVs were generated by homology modelling
utilizing the I-TASSER modeling tool (https://zhanglab.ccmb.med.umich.edu/I-
TASSER/). The I-TASSER is a tool for protein structure prediction based on the

sequence-to-structure-to-function paradigm (Roy et al., 2010). The tool

generates three-dimensional (3D) atomic models from multiple threading

alignments and iterative structural assembly simulations for a submitted amino

acid sequence. I-TASSER works on the basis of the structure templates

identified by LOMETS, a meta-server, from the PDB library. I-TASSER only uses

the templates of the highest Z-score which is the difference between the raw and

average scores in the unit of standard deviation. For each target model the I-

TASSER simulations generate a large ensemble of structural conformations,

called decoys. To select the final models, I-TASSER uses the SPICKER program

to cluster all the decoys based on the pair-wise structure similarity, and reports

up to five models. The Normalized Z-score >1 mean a good alignment and vice

versa. The Cov represents the coverage of the threading alignment and is equal

to the number of aligned residues divided by the length of query protein. Ranking

of templet proteins is based on TM-score of the structural alignment between the

query structure model and known structures. The RMSD is the RMSD between

templet residues and query residues that are structurally aligned by TM-align.

The refinement of both the generated MEV models was performed by

ModRefiner   (https://zhanglab.ccmb.med.umich.edu/ModRefiner/)   and

GalaxyRefine   tool   (http://galaxy.seoklab.org/cgi-bin/submit.cgi?type=REFINE)

(Dong et al., 2011). TM-score generated by ModRefiner indicates the structural

similarity of the refined model with the original input model. Closer the TM-Score

to 1, higher would be the similarity of original and the refined model. RMSD value

of the refined model shows the conformational deviation from the initial input

models.

The GalaxyRefine tool refines the query tertiary structure by repeated

structure perturbation as well as by utilizing the subsequent structural relaxation

by the molecular dynamics simulation. The tool GalaxyRefine generates reliable

core structures from multiple templates and then re-builds unreliable loops or

termini by using an optimization-based refinement method (Ko et al., 2012; Wang

et al., 2013; Shin et al., 2014). To avoid any breaks in the 3D model

GalaxyRefine uses triaxial loop closure method. The MolProbity score generated

for a given refined model indicates the log-weighted combination of the clash

score, the percentage of Ramachandran not favoured residues and the

percentage of bad side-chain rotamers.

Validation of CTL and HTL MEVs refined models. Both the refined CTL and HTL

MEV 3D models were further validated by RAMPAGE analysis tool

(http://mordred.bioc.cam.ac.uk/~rapper/rampage.php) (S.C. Lovell et al., 2008;

Ramakrishnan, C. et al., 1965). The generated Ramachandran plots for the MEV

models show the sterically allowed and disallowed residues along with their

dihedral psi (ψ) and phi (φ) angles.

Linear and Discontinuous B-cell epitope prediction from MEVs. The Ellipro

(ElliPro: Antibody Epitope Prediction tool; http://tools.iedb.org/ellipro/) method

available at IEDB, was used to screen the linear and the discontinuous B cell

epitopes from the MEVs vaccine models. The ElliPro method analyses on the basis of the location of residue in the protein's 3D structure. The residues lying outside of an ellipsoid covering 90% of the inner core protein residues score highest Protrusion Index (PI) of 0.9; and so on. The discontinuous epitopes predicted by the ElliPro tool are clustered on the basis of the distance "R" in Å between two residue's centers of mass lying outside of the largest possible ellipsoid. The larger value of R indicates larger distant residues (residue discontinuity) are screened in the epitopes (Kringelum et al., 2012; Ponomarenko et al., 2008).

Molecular interaction analysis of MEVs with immunological receptor.

Molecular docking and dynamics simulation study of MEVs and TLR-3. Molecular interaction analysis of both the designed MEVs with Toll-Like receptor-3 (TLR-3), was performed by molecular docking and molecular dynamics simulation. Molecular docking was performed by PatchDock server (http://bioinfo3d.cs.tau.ac.il/PatchDock/) (Schneidman-Duhovny et al., 2005). PatchDock utilizes algorithm for unbound (real life) docking of molecules for protein-protein complex formation as explained earlier (Bell et al., 2005; Duhovny et al., 2002). For molecular docking, the 3D structure of human TLR-3 ectodomain (ECD) was retrieved from PDB databank (PDB ID: 2A0Z). The study provides dynamical properties of the designed system with MEVs-TLR3 complexes with a guess at the interactions between the molecules, and also it

gives 'exact' predictions of bulk properties including the hydrogen bond formation

and conformation of the molecules forming the complex.

In-silico analysis of MEVs for cloning and expression potency

Analysis of cDNA of both the MEVs for cloning and expression in mammalian

host cell line. Complementary DNA of both the MEVs, codon optimized for

expression in Mammalian cell line (Human) were generated by Java Codon

Adaptation Tool (http://www.jcat.de/). The generated cDNA of both the MEVs

were further analysed by GenScript Rare Codon Analysis Tool

(https://www.genscript.com/tools/rare-codon-analysis). The tool analyses the GC

content, Codon Adaptation Index (CAI) and the Tandem rare codon frequency for

a given cDNA (Morla et al., 2016; Wu et al., 2010). The CAI indicates the

possibility of cDNA expression in a chosen expression system. The tandem rare

codon frequency indicates the presence of low-frequency codons in the given

cDNA.

RESULTS & DISCUSSION

Screening of potential epitopes

T cell Epitope Prediction

Screening of Cytotoxic T lymphocyte (CTL) Epitope. Cytotoxic T lymphocyte

(CTL) epitopes were screened by "MHC-I Binding Predictions" and "MHC-I

Binding Predictions" IEDB tools. These epitopes are shortlisted on the basis of

total amount of cleavage site in the protein, low IC(50) (nM) value for epitope-

HLA class I allele pairs, and for binding to TAP cavity.

The 38 epitopes predicted by "MHC-I Binding Predictions" tool with

highest "Percentile Rank" were shortlisted for multi-epitope vaccine design and

are listed in table 1. Rest 101 epitopes are listed in Supplementary table S8. The

67 epitopes predicted by "MHC-I Processing Predictions" tool with highest "Total

score" are listed in Supplementary table 9.

The immunogenicity of the shortlisted CTL epitopes was also determined

and are mentioned in table 1, Supplementary table S8, and S9. The higher

immunogenicity score indicates the greater immunogenic potential of the given

epitope.

Table 1. Shortlisted high percentile ranking SARS-CoV-2 CTL epitopes. Selected
high percentile CTL epitopes and their respective HLA alleles binders are listed. *In-silico*
analysis have shown all the selected epitopes to be non-toxic (Non-Toxin) as well as they
show significant conservancy and high immunogenicity. Six epitope shown with ** show
exact match with the epitopes published by Grifoni et al., 2020, indicate consensus in
results.
576
577

SARS-CoV-19 Proteins	Peptide	Conservancy	Immunogenicity	Toxicity	Allele	Position	Length	Method used	Percentile Rank
E Protein	LLFLAFVVF	100.00% (35/35)	0.2341	Non-Toxin	B*15:01	18-26	9	Consensus (ann/comblib_sidney2008/smm)	0.1
E Protein	LTALRLCAY	97.14% (34/35)	0.01886	Non-Toxin	A*01:01	34-42	9	Consensus (ann/smm)	0.12
M Protein	ATSRTLSYYK**	100.00% (41/41)	-0.13563	Non-Toxin	A*11:01	31-40	10		0.06
M Protein	YFIASFRLFAR	100.00% (41/41)	0.19709	Non-Toxin	A*33:01	25-35	11	ann	0.03
N protein	KPRQKRTAT	100.00% (40/40)	-0.20542	Non-Toxin	B*07:02	47-55	9	Consensus (ann/comblib_sidney2008/smm)	0.1
N protein	MEVTPSGTW	100.00% (40/40)	-0.06279	Non-Toxin	B*44:02	42-50	9	Consensus (ann/smm)	0.06
ORF-1ab	GYINVFAFPF**	100.00% (28/28)	0.32004	Non-Toxin	A*23:01	2-11	10		0.11
ORF-1ab	MGYINVFAF	100.00% (28/28)	0.28694	Non-Toxin	B*35:01	1-9	9	Consensus (ann/comblib_sidney2008/smm)	0.1
ORF-1ab	CTDDNALAY	100.00% (36/36)	0.07355	Non-Toxin	A*01:01	33-41	9	Consensus (ann/smm)	0.06
ORF-1ab	CTDDNALAYY**	100.00% (36/36)	0.08174	Non-Toxin	A*01:01	33-42	10		0.06

ORF-1ab	DEWSMATYY**	97.22% (35/36)	-0.19814	Non-Toxin	B*44:03	58-66	9		0.07
ORF-1ab	ETISLAGSYK	100.00% (36/36)	-0.20585	Non-Toxin	A*68:01	33-42	10		0.06
ORF-1ab	FLFVAAIFYL	100.00% (36/36)	0.37766	Non-Toxin	A*02:01	50-59	10		0.06
ORF-1ab	FLNGSCGSV	100.00% (36/36)	-0.24791	Non-Toxin	A*02:03	43-51	9		0.06
ORF-1ab	FYWFFSNYLKR	100.00% (36/36)	-0.00254	Non-Toxin	A*33:01	3-13	11	ann	0.04
ORF-1ab	ISNSWLMW	100.00% (36/36)	-0.11151	Non-Toxin	B*58:01	49-56	8		0.05
ORF-1ab	LPVNAFELW	97.22% (35/36)	0.27341	Non-Toxin	B*53:01	61-70	10	Consensus (ann/smm)	0.06
ORF-1ab	MYKGLPWNVVR	94.44% (34/36)	0.21107	Non-Toxin	A*33:01	58-68	11	ann	0.06
ORF-1ab	QEILGTVSW	100.00% (36/36)	0.03976	Non-Toxin	B*44:02	35-43	9		0.06
ORF-1ab	RMYIFFASFY	100.00% (36/36)	0.32633	Non-Toxin	A*30:02	2-11	10	Consensus (ann/smm)	0.06
ORF-1ab	RYFRLTLGVY	100.00% (36/36)	0.15936	Non-Toxin	A*30:02	22-31	10		0.06
ORF-1ab	SEMVMCGGSLY	97.22% (35/36)	-0.32016	Non-Toxin	B*44:02	16-26	11	ann	0.03
ORF-1ab	SIINNTVYTK**	97.22% (35/36)	0.12661	Non-Toxin	A*11:01	37-46	10		0.06
ORF-1ab	STFNVPMEK	100.00% (36/36)	-0.02845	Non-Toxin	A*11:01	10-18	9	Consensus (ann/smm)	0.06
ORF-1ab	YIFFASFYYV	100.00% (36/36)	0.13772	Non-Toxin	A*02:06	4-13	10		0.07
ORF-1ab	YILFTRFFYV	100.00% (36/36)	0.40924	Non-Toxin	A*02:06	21-30	10		0.07
ORF3a	IPYNSVTSSI	100.00% (28/28)	-0.32835	Non-Toxin	B*51:01	18-27	10		0.11
ORF3a	YLYALVYFL**	100.00% (28/28)	0.13151	Non-Toxin	A*02:01	37-45	9	Consensus (ann/comblib_sidney2008/smm)	0.1
ORF6	AEILLIIMRTF	100.00% (34/34)	0.1815	Non-Toxin	B*44:02	12-22	11	ann	0.06
ORF6	RTFKVSIW	100.00% (34/34)	-0.18221	Non-Toxin	B*57:01	20-27	8		0.05
ORF7a	QLRARSVSPK	100.00% (28/28)	-0.16177	Non-Toxin	A*03:01	6-15	10	Consensus (ann/smm)	0.16
ORF7a	RARSVSPK	100.00% (28/28)	-0.27456	Non-Toxin	A*30:01	8-15	8	ann	0.11
ORF7b	FLAFLLFLV	100.00% (2/2)	0.20158	Non-Toxin	A*02:03	13-21	9		0.07
ORF8	HFYSKWYIR	100.00% (34/34)	-0.09452	Non-Toxin	A*31:01	40-48	9	Consensus (ann/smm)	0.11
S Protein	FPNITNLCPF	100.00% (41/41)	0.1009	Non-Toxin	B*53:01	49-58	10		0.06
S Protein	NYLYRLFR	100.00% (41/41)	0.13144	Non-Toxin	A*33:01	30-37	8	ann	0.07
S Protein	NYNYLYRLFR	100.00% (41/41)	0.08754	Non-Toxin	A*33:01	28-37	10	Consensus (ann/smm)	0.07
S Protein	WTAGAAAYYV	100.00% (41/41)	0.15455	Non-Toxin	A*68:02	48-57	10		0.06

578
579
580

Screening of Helper T lymphocyte (HTL) epitopes. The screening of helper T lymphocyte (HTL) epitopes from eleven different proteins of SARS-CoV-2 was performed on the basis of "Percentile rank". The smaller the value of percentile rank the higher would be the affinity of the peptide with its respective HLA allele binders. The 42 epitopes with high percentile ranking were shortlisted (Table 2). Another 171 potential HTL cell epitopes with high Percentile rank, screened in our study are listed in Supplementary table S10.

589 **Table 2. Shortlisted high scoring SARS-CoV-2 HTL epitopes.** Selected high scoring
590 HTL epitopes and their respective HLA alleles binders are listed. *In-silico* analysis have
591 shown all the selected epitopes to be non-toxic (Non-Toxin) as well as they show
592 significant conservancy. Six epitope shown with ** show exact match with the epitopes
593 published by Grifoni et al., 2020, indicate consensus in results.
594
595

SARS-CoV- 19 Proteins	Peptide	Conservancy	Non-Toxin	Allele	Position	Method used	Percentile Rank
E Protein	VLLFLAFVVFLLVTL	100.00% (35/35)	Non-Toxin	DPA1*03:01/DPB1*04:02	17-31		0.02
E Protein	LLFLAFVVFLLVTLA	100.00% (35/35)	Non-Toxin	DPA1*03:01/DPB1*04:02	18-32		0.02
M Protein	GLMWLSYFIASFRLF	100.00% (41/41)	Non-Toxin	DPA1*01:03/DPB1*02:01	19-33		0.05
M Protein	LMWLSYFIASFRLFA	100.00% (41/41)	Non-Toxin	DPA1*01:03/DPB1*02:01	20-34		0.05
M Protein	LSYYKLGASQRVAGD**	100.00% (41/41)	Non-Toxin	DRB1*09:01	36-50		0.06
N Protein	PQIAQFAPSASAFFG	100.00% (40/40)	Non-Toxin	DRB1*09:01	22-36		0.01
N Protein	IAQFAPSASAFFGMS	100.00% (40/40)	Non-Toxin	DRB1*09:01	24-38		0.01
N Protein	AQFAPSASAFFGMSR	100.00% (40/40)	Non-Toxin	DRB1*09:01	25-39		0.01
ORF1ab	WFLAYILFTRFFYVL	100.00% (36/36)	Non-Toxin	DPA1*01:03/DPB1*02:01	17-31		0.01
ORF1ab	FLAYILFTRFFYVLG	100.00% (36/36)	Non-Toxin	DPA1*01:03/DPB1*02:01	18-32		0.01
ORF1ab	LAYILFTRFFYVLGL	100.00% (36/36)	Non-Toxin	DPA1*01:03/DPB1*02:01	19-33	Consensus (comb.lib./smm/nn)	0.01
ORF1ab	AYILFTRFFYVLGLA	100.00% (36/36)	Non-Toxin	DPA1*01:03/DPB1*02:01	20-34		0.01
ORF1ab	RAMPNMLRIMASLVL**	97.22% (35/36)	Non-Toxin	DRB1*15:01	46-60		0.01
ORF1ab	AMPNMLRIMASLVLA	97.22% (35/36)	Non-Toxin	DRB1*15:01	47-61		0.01
ORF1ab	MPNMLRIMASLVLAR	97.22% (35/36)	Non-Toxin	DRB1*15:01	48-62		0.01
ORF1ab	PNMLRIMASLVLARK	97.22% (35/36)	Non-Toxin	DRB1*15:01	49-63		0.01
ORF1ab	QESPFVMMSAPPAQY	100.00% (36/36)	Non-Toxin	DRB1*01:01	50-64		0.01
ORF1ab	NMLRIMASLVLARKH	97.22% (35/36)	Non-Toxin	DRB1*15:01	50-64		0.01
ORF1ab	ESPFVMMSAPPAQYE**	100.00% (36/36)	Non-Toxin	DRB1*01:01	51-65		0.01
ORF1ab	SPFVMMSAPPAQYEL	100.00% (36/36)	Non-Toxin	DRB1*01:01	52-66		0.01
ORF1ab	AIILASFSASTSAFV	100.00% (36/36)	Non-Toxin	DRB1*09:01	54-68		0.01
ORF1ab	IILASFSASTSAFVE	100.00% (36/36)	Non-Toxin	DRB1*09:01	55-69		0.01
ORF1ab	FAWWTAFVTNVNASS	97.22% (35/36)	Non-Toxin	DRB3*02:02	55-69	NetMHCIIpan	0.01
ORF3a	FVRATATIPIQASLP	100.00% (28/28)	Non-Toxin	DPA1*02:01/DPB1*14:01	28-42		0.12
ORF3a	LLFVTVYSHLLLVAA	100.00% (28/28)	Non-Toxin	DRB1*01:01	15-29		0.1
ORF6	TFKVSIWNLDYIINL**	100.00% (34/34)	Non-Toxin	DQA1*01:01/DQB1*05:01	21-35		0.02
ORF6	FKVSIWNLDYIINLI	100.00% (34/34)	Non-Toxin	DQA1*01:01/DQB1*05:01	22-36		0.02
ORF6	KVSIWNLDYIINLII	100.00% (34/34)	Non-Toxin	DQA1*01:01/DQB1*05:01	23-37	Consensus (comb.lib./smm/nn)	0.02
ORF7a	IILFLALITLATCEL	100.00% (28/28)	Non-Toxin	DRB1*01:01	3-17		0.16
ORF7a	ILFLALITLATCELY	100.00% (28/28)	Non-Toxin	DRB1*01:01	4-18		0.16
ORF7b	YLCFLAFLLFLVLIM	100.00% (2/2)	Non-Toxin	DPA1*03:01/DPB1*04:02	10-24		0.02
ORF7b	LCFLAFLLFLVLIML	100.00% (2/2)	Non-Toxin	DPA1*03:01/DPB1*04:02	11-25		0.02
ORF7b	CFLAFLLFLVLIMLI	100.00% (2/2)	Non-Toxin	DPA1*03:01/DPB1*04:02	12-26		0.03

ORF8	CTQHQPYVVDDPCPI	100.00% (34/34)	Non-Toxin	DRB3*01:01	25-39		0.08
ORF8	HQPYVVDDPCPIHFY	100.00% (34/34)	Non-Toxin	DRB3*01:01	28-42		0.08
ORF8	QPYVVDDPCPIHFYS	100.00% (34/34)	Non-Toxin	DRB3*01:01	29-43		0.07
ORF10	YINVFAFPFTIYSLL	100.00% (28/28)	Non-Toxin	DPA1*01:03/DPB1*02:01	3-17		0.29
S protein	KTQSLLIVNNATNVV	100.00% (41/41)	Non-Toxin	DRB1*13:02	43-57		0.01
S protein	TQSLLIVNNATNVVI	100.00% (41/41)	Non-Toxin	DRB1*13:02	44-58		0.01
S protein	QSLLIVNNATNVVIK	100.00% (41/41)	Non-Toxin	DRB1*13:02	45-59		0.01
S protein	SLLIVNNATNVVIKV**	100.00% (41/41)	Non-Toxin	DRB1*13:02	46-60		0.01
S protein	LLIVNNATNVVIKVC	100.00% (41/41)	Non-Toxin	DRB1*13:02	47-61		0.01

596
597
598

Population Coverage by CTL and HTL epitopes. The population coverage by the

shortlisted epitopes was also studied, in particular involving China, France, Italy,

United States of America, South Asia, East Asia, Northeast Asia and the Middle

East. From this study, we may conclude that the combined use of all the

shortlisted CTL and HTL epitopes would have an average worldwide population

coverage as high as 96.10%, with a standard deviation of 31.17 (Table 3).

Table 3. World population coverage by the shortlisted SARS-CoV-2 CTL and HTL epitopes combined. With a standard deviation of 31.17 on an average 96.10 % of world population could be covered by the joint administration of selected CTL and HTL epitopes (given in Table 1 and 2) as vaccine candidate.

a projected population coverage
b average number of epitope hits / HLA combinations recognized by the population
c minimum number of epitope hits / HLA combinations recognized by 90% of the population

Population/area	Class combined		
	% coverage[a]	average_hit[b]	pc90[c]
Algeria	27.85	1.51	0.69
American Samoa	70.35	1.93	0.34
Argentina	96.78	3.56	1.35
Australia	79.62	2.73	0.49
Austria	99.28	7.17	2.74
Belgium	97.21	4.63	1.71
Bolivia	35.81	2.14	0.77
Borneo	30.61	1.53	0.72
Brazil	92.28	3.69	1.15
Bulgaria	95.49	4.48	1.64

Burkina Faso	61.26	1.3	0.26
Cameroon	88.9	2.71	0.9
Canada	8.75	0.46	0.54
Cape Verde	98.02	5.71	2.07
Central Africa	87.66	3.35	0.81
Central African Republic	36.4	1.25	0.16
Central America	16.77	0.65	0.12
Chile	90.76	4.36	1.06
China	90.43	4.74	1.05
Colombia	24.9	1.18	0.27
Congo	29.65	1.58	0.64
Cook Islands	22.52	1.33	0.65
Croatia	97.46	5.28	1.97
Cuba	97.36	6.08	2.03
Czech Republic	98.68	5.99	2.31
Denmark	59.3	3.61	1.23
East Africa	90.29	3.46	1.02
East Asia	93.23	5.49	1.29
Ecuador	51.11	2.55	0.2
England	99.45	7.37	3.14
Equatorial Guinea	15.47	0.81	0.58
Ethiopia	34.14	1.72	0.71
Europe	98.35	6.12	2.32
Fiji	23.54	1.33	0.59
Finland	99.58	5.27	2.46
France	98.2	6.39	2.31
Gabon	7.02	0.42	0.64
Georgia	93.92	4.08	1.31
Germany	99.34	7.17	2.92
Greece	26.34	1.51	0.68
Guatemala	17.11	0.67	0.12
Guinea-Bissau	95.03	4.49	1.37
Hong Kong	84.82	2.7	0.66
India	86.82	3.86	0.76
Indonesia	78.44	2.93	0.46
Iran	94.14	4.47	1.45
Ireland Northern	99.55	7.7	3.3
Ireland South	99.28	7.31	3.16
Israel	84.33	3.49	0.64
Italy	96.7	4.11	1.74
Ivory Coast	65.09	1.16	0.29
Jamaica	4.55	0.23	0.52
Japan	92.78	5.3	1.23
Jordan	85.1	3.54	0.67
Kenya	86.65	2.27	0.75
Kiribati	6.3	0.31	0.53
Korea; South	95.79	6.18	1.72
Lebanon	14.26	0.8	0.58
Macedonia	43.69	2.18	0.18
Malaysia	80.94	3.79	0.52
Mali	95.75	2.67	1.3
Martinique	43.47	1.8	0.18
Mexico	93.68	3.6	1.35

Mongolia	87.03	4.25	0.77
Morocco	97.32	5.55	2.01
Nauru	18.44	0.93	0.61
Netherlands	40.49	2.35	0.84
New Caledonia	61.85	2.73	0.26
New Zealand	24.11	1.39	0.66
Niue	20.26	1.2	0.63
North Africa	94.14	4.19	1.3
North America	97.36	6	2.09
Northeast Asia	90.3	4.71	1.04
Norway	52.39	3.08	1.05
Oceania	75.64	3.29	0.41
Oman	94.27	3.2	1.35
Pakistan	85	2.54	0.67
Papua New Guinea	71.23	3.29	0.35
Paraguay	0	0	0.6
Peru	95.12	3.38	1.52
Philippines	67.23	1.92	0.61
Poland	98.46	6.18	2.36
Portugal	96.27	5.25	1.75
Romania	95.9	3.54	1.56
Russia	94.88	5.26	1.49
Rwanda	45.12	1.85	0.18
Samoa	43.64	2.65	0.89
Sao Tome and Principe	96.32	4.15	1.54
Saudi Arabia	96.53	4.87	1.65
Scotland	73.06	4.19	0.37
Senegal	94.11	3.54	1.24
Serbia	62.91	1.07	0.27
Singapore	86.92	4.37	0.76
Slovenia	43.37	2.58	0.88
South Africa	88.81	2.6	0.89
South America	81.68	2.95	0.55
South Asia	90.37	4.25	1.03
Southeast Asia	82.33	3.79	0.57
Southwest Asia	86.38	3.13	0.73
Spain	87.57	3.8	0.8
Sri Lanka	52.39	1.59	0.42
Sudan	87.98	2.69	0.83
Sweden	99.29	7.32	2.59
Taiwan	84.93	4.28	0.66
Thailand	91.26	4.81	1.17
Tokelau	7.84	0.47	0.65
Tonga	33.43	1.98	0.69
Tunisia	94.64	4.17	1.38
Turkey	61.05	2.54	0.51
Uganda	92	2.99	1.13
United Arab Emirates	2.19	0.02	0.1
United States	97.45	6.04	2.11
Venezuela	69.56	1.16	0.33
Vietnam	86.19	4.11	0.72
Wales	1	0.01	0.1
West Africa	94.34	4.44	1.32

West Indies	96.71	5.04	1.74
Zambia	97.5	3.1	1.67
Zimbabwe	93.6	3.4	1.21
World	**96.10**	**5.55**	**1.8**
Average	**70.54**	**3.35**	**1.07**
Standard deviation	**31.17**	**1.87**	**0.73**

B Cell epitope prediction

Sequence-based B Cell epitope prediction. To screen B cell epitopes we utilized the Bepipred Linear Epitope Prediction method. In our study, we screened 12 B cell epitopes, from eleven SARS-CoV-2 proteins, which show partial or complete overlap with shortlisted CTL and HTL epitope (Table 4). Another 206 B Cell epitope, with the epitope length of at least four amino acids and maximum 20 amino acid were screened and are listed in Supplementary table S11.

Table 4. Shortlisted SARS-CoV-2 B Cell epitopes. BepiPred Linear B Cell epitopes showing sequence overlap with CTL and HTL epitopes are shortlisted. *In-silico* analysis have shown all the selected epitopes to be non-toxic (Non-Toxin) as well as they show significant amino acid sequence conservancy.

S.No	SARS-2-CoV-Proteins	Overlapping B Cell Epitope (BepiPred method)	Conservancy	Length	Position	Toxicity
1	Membrane Protein	KLGASQRVAGDS	100.00% (41/41)	12	180-191	Non-Toxin
2	N protein	RLNQLESKMSGKGQQQQGQTVTKKSAAEASK KPRQKRTATKA	100.00% (40/40)	42	226-267	Non-Toxin
3	ORF1ab	GTTQTACTDDNALAYYNTTK	100.00% (36/36)	20	4157-4176	Non-Toxin
4	ORF3a	QGEIKDATPSDF	100.00% (28/28)	12	17-28	Non-Toxin
5	ORF3a	PYNSVT	100.00% (28/28)	6	159-164	Non-Toxin
6	ORF7a	LYHYQECVR	100.00% (28/28)	9	17-25	Non-Toxin
7	ORF7a	VKHVYQLRARSVSPKLFIRQEEVQEL	100.00% (28/28)	26	71-96	Non-Toxin
8	ORF8	QSCTQHQPYVVDDPCPIHFYSKW	100.00% (34/34)	23	23-45	Non-Toxin
9	ORF8	RVGARKSAP	100.00% (34/34)	9	48-56	Non-Toxin
10	S protein	TPGDSSSGWTA	100.00% (41/41)	11	250-260	Non-Toxin
11	S protein	FPNITNLCPFGEVFNATRFASVYAWNRKRISNCVA	100.00% (41/41)	35	329-363	Non-Toxin
12	S protein	NLDSKVGGNYNYLYRLFRKSNLKPFERDISTEIY QAGSTPCNGVEGFNCYFPLQSYGFQPTN	100.00% (41/41)	62	440-501	Non-Toxin

643
Characterisation of potential epitopes

Epitope conservation analysis. Sequence conservation analysis of the screened

CTL, HTL and B cell epitopes have shown highly conserved nature of the

shortlisted epitopes. Both the CTL epitopes and the HTL epitope were found to

be significantly conserved with their 100% amino acid sequence amongst the

NCBI retrieved protein sequences of SARS-CoV-2 (CTL epitopes 94.44% to

100% conserved and HTL epitopes were 97.22% to 100% conserved (Table 1, 2

4, Supplementary table S8, S9, S10 & S11).

Epitope toxicity prediction. Toxicity analysis of all the screened CTL, HTL and B

Cell epitopes was also performed. The ToxinPred study of all the shortlisted

epitopes shows that they all are non-toxic in nature (Table 1, 2 4, Supplementary

table S8, S9, S10 & S11).

Overlapping residue analysis. Amino acid sequence overlap analysis amongst

the shortlisted CTL, HTL and B cell epitopes from eleven SARS-CoV-2 proteins

was performed by the Multiple Sequence Alignment (MSA) analysis tool Clustal

Omega. The analysis has shown that several epitopes of CTL, HTL and B cell

were having amino acid sequence overlap. The CTL, HTL and B cell epitopes

having two or more than two amino acid residues overlap are shown in Fig.2.

Overlapping T & B cell epitopes across proteome of SARS-CoV-2

Figure 3. Overlapping SARS-CoV-2 CTL, HTL and B cell epitopes. Multiple

sequence alignment performed by Clustal Omega at EBI to identify the

consensus overlapping regions of CTL (red), HTL (blue) and B cell epitopes

(green) amongst shortlisted epitopes. Epitopes with overlapping region amongst

all the three types of epitopes (CTL, HTL and B Cell epitopes) were chosen for

further studies (encircled).

Epitope selected for molecular interaction study with HLA allele and TAP

transporter. The epitopes showing overlap amongst all the three type of epitopes

i.e CTL, HTL and B cell epitopes have been encircled in Fig.2 and are chosen for

further study for their interaction with HLA allele and TAP (Transporter

Associated with Antigen Processing) transporter.

Molecular interaction analysis of selected epitopes with HLA allele and TAP

transporter.

Molecular interaction analysis of chosen CTL and HTL epitopes with HLA alleles.

The molecular docking study of chosen CTL and HTL epitopes with their

respective HLA class I and II allele binders was performed by PatchDock tool.

The study revealed a significant molecular interaction between all the chosen

epitopes and their HLA allele binders showing multiple hydrogen bond formations

(Fig.3). Furthermore, the B-factor analysis of all the epitope-HLA allele

complexes has also shown the epitope ligand to have stable (blue) binding

conformation in complex with the HLA allele molecule (VIBGYOR colour

presentation, blue being very stable) Fig.4.

1. ATSRTLSYYK-HLA-A*11:01

2. HFYSKWYIR-HLA-A*31:01

CTL epitopes - HLA Class I allele complexes

3. CTQHQPYVVDDPCPI-HLA-DRB3*01:01

4. HQPYVVDDPCPIHFY-HLA-DRB3*01:01

5. LSYYKLGASQRVAGD-HLA-DRB1*09:01

6. QPYVVDDPCPIHFYS-HLA-DRB3*01:01

HTL epitopes - HLA Class II allele complexes

Figure 4. Molecular Docking analysis of SARS-CoV-2 CTL epitopes and

HLA alleles. Molecular docking of chosen CTL and HTL epitopes (cyan sticks)

with binding amino acid residues of their respective HLA class I and class II allele

binders (magenta sticks). The study shows the docked complexes to form stable

complex with multiple hydrogen bonds (green dots) formation. Images were

generated by the PyMOL Molecular Graphics System, Version 2.0 Schrödinger,

LLC.

1. ATSRTLSYYK-HLA-A*11:01   **2. HFYSKWYIR-HLA-A*31:01**

CTL epitopes - HLA Class I allele complexes

3. CTQHQPYVVDDPCPI-HLA-DRB3*01:01   **4. HQPYVVDDPCPIHFY-HLA-DRB3*01:01**

5. LSYYKLGASQRVAGD-HLA-DRB1*09:01   **6. QPYVVDDPCPIHFYS-HLA-DRB3*01:01**

HTL epitopes - HLA Class II allele complexes

Figure 5. B-Factor of CTL and HTL epitope in complex with HLA class I and II

allele. CTL and HTL Epitopes are shown in sticks and HLA Class I and II alleles are

shown in ribbon. The regions of the epitope in complex are shown in rainbow

(VIBGYOR), the regions in blue being very stable and the region towards red being

relatively unstable. In the complexes shown above, most of the regions of epitopes are in

blue indicating the complexes to be highly stable.

Molecular interaction analysis of selected CTL epitopes with TAP transporter.

The molecular docking interaction analysis of the chosen CTL epitopes with the

TAP transporter cavity has shown a significantly strong molecular interaction with

several hydrogen bonds formation at different sites of the TAP transporter cavity.

Two sites of interaction were of particular interest, one closer to the cytoplasmic

end and another closer to the ER lumen (Fig.5). This study confirms the

feasibility of transportation of chosen CTL epitopes from the cytoplasm to the ER

lumen which is an essential event for the representation of epitope by the HLA

allele molecules on the surface of antigen presenting cells.

Epitope ATSRTLSYYK through TAP cavity

Epitope HFYSKWYIR through TAP cavity

Figure 6. Molecular docking analysis of CTL epitopes within the TAP transporter cavity. Molecular interaction of CTL epitopes (cyan sticks) within the TAP cavity (gray ribbon/sticks) is shown. Detailed interaction between the residues of epitopes and TAP transporter residues have been shown with hydrogen formation shown with green dots.

Characterisation and molecular interaction analysis of designed Multi-Epitope Vaccines with immunological receptor

Characterisation of designed Multi-Epitope Vaccines

Physicochemical property analysis of designed MEVs. ProtParam analysis for both the CTL and HTL MEVs was performed to analyse their physiochemical properties. The empirical physiochemical properties of the CTL and HTL MEVs are given in the table 5. The aliphatic index and grand average of hydropathicity (GRAVY) of both the MEVs indicate globular and hydrophilic nature of the both the MEVs. The instability index score of both the MEVs indicate stable nature of the protein molecules (Table 5).

872 **Table 5. Physicochemical property analysis based on amino acid sequence**

873 **of designed CTL and HTL multi-epitope vaccine.**

MEVs	Amino acid length	Molecular Weight	Theoretical pI	Expected half-life	Aliphatic index	Grand average of hydropathicity (GRAVY)	Instability index score
CTL MEV	704 aa	72.62 kDa	9.70	E.Coli: 10 Hr Yeast: 20 Hr Mammalian cell: 30 Hr	61.09	-0.090	44.31
HTL MEV	989 aa	101.31 kDa	9.39	Coli: 10 Hr Yeast: 20 Hr Mammalian cell: 30 Hr	96.35	0.539	43.31

874

875

876 *Interferon-gamma inducing epitope prediction.* Interferon-gamma (IFN-γ)

877 inducing epitopes are involved in both the adaptive as well as in the innate

878 immune response. The IFN-γ inducing 15 mer peptide epitopes were screened

879 from the amino acid sequence of CTL and HTL MEVs by utilizing the IFNepitope

880 server. A total of 20 CTL MEV and 20 HTL MEV INF-γ inducing POSITIVE

881 epitopes with a score of 1 or more than 1 were shortlisted (Supplementary table

882 S2).

883

884

885

886

887

888

889

A. CTL Multi-epitope vaccine model

B. HTL Multi-epitope vaccine model

Figure 7. Tertiary structure modelling of CTL and HTL Multi-Epitope Vaccines. Tertiary structural models of CTL and HTL MEVs have been shown. The epitopes are shown in cyan. The adjuvant (Ov-ASP-1) is shown in orange. The linkers are shown in gray. Both the cartoon and surface presentation of both the MEVs are shown.

A. CTL Multi-epitope vaccine RAMPAGE analysis

Number of residues in favoured region (~98.0% expected) : 602 (85.8%)
Number of residues in allowed region (~2.0% expected) : 79 (11.3%)
Number of residues in outlier region : 21 (3.0%)

B. HTL Multi-epitope vaccine model RAMPAGE analysis

Number of residues in favoured region (~98.0% expected) : 718 (88.9%)
Number of residues in allowed region (~2.0% expected) : 72 (8.9%)
Number of residues in outlier region : 18 (2.2%)

Figure 8. RAMPAGE analysis of CTL and HTL MEVs. The RAMPAGE analysis

of both the CTL and HTL MEVs have been done and shown here.

MEVs allergenicity and antigenicity prediction. Both the CTL and HTL MEVs

were found to be NON-ALLERGEN by the AlgPred analysis (scoring -

0.95185601 and -1.1293352 respectively; threshold being -0.4). The CTL and

HTL MEVs were also analyzed by VaxiJen to be probable ANTIGENS (prediction

score 0.4485 and 0.4215 respectively; default threshold being 0.4). Hence with

the mentioned analysis tools both the CTL and HTL MEVs are predicted to be

non-allergic as well as potentially antigenic in nature.

Tertiary structure modelling and refinement of MEVs. 3D homology models were

generated for both the CTL and HTL MEVs by utilizing the I-TASSER modeling

tool (Fig.7). The models were generated for CTL Multi-epitope vaccine (PDB hit

5n8pA, Norm. Z-score of 1.49, Cov of 0.92, TM-score of 0.916 and RMSD of

1.04 Å) and HTL Multi-epitope vaccine (PDB hit 5n8pA, Norm. Z-score of 1.52,

Cov of 0.97, TM-score of 0.916 and RMSD of 1.04 Å).

Both the generated CTL and HTL 3D models were further refined by

ModRefiner to fix any gaps and then followed by GalaxyRefine refinement. The

refinement by ModRefiner showed the TM-score of 0.9189 and 0.9498 for the

CTL and HTL models respectively, hence being close to 1, the initial and the

refined models were structurally similar. After refinement, the RMSD for CTL and

HTL models with respect to the initial model was 3.367Å and 2.318Å respectively.

Further, both the CTL and HTL MEVs models were refined by GalaxyRefine and

model 1 was chosen on the basis of best scorings parameters. The CTL MEV

model refinement output model (Rama favoured was 83.6%, GDT-HA was

0.9371, RMSD was 0.459, MolProbity was 2.539, Clash score was 23.2, and

Poor rotamers was 1.8) and the HTL MEV model refinement output model (Rama

favoured was 87.7%, GDT-HA was 0.9552, RMSD was 0.402, MolProbity was

2.537, Clash score was 27.9, and Poor rotamers was 1.6) show a well refined

and acceptable models for both of the MEVs. After refinement, all the mentioned

parameter have found to be improved significantly in comparison to the initial

CTL and HTL MEV models (Supplementary table S3).

Validation of CTL and HTL MEVs refined models. Both the CTL and HTL model

were analyzed by the RAMPAGE analysis tool after refinement. The refined CTL

MEV model was found to have 85.8% residues in favored region, 11.3% residues

in allowed region, and only 3.0% residues in the outlier region; while the refined

HTL MEV model was found to have 88.9% of residues in favored region, 8.9%

residues in allowed region, and only 2.2% residues in the outlier region (Fig.8)

Linear and Discontinuous B-cell epitope prediction from MEVs. The linear and

discontinuous B-cell epitope prediction was performed to enlist potential linear

and discontinuous epitopes from the refined 3D models of CTL and HTL MEVs

utilizing the ElliPro tool available on IEDB server. The screening revealed that the

CTL MEV carries 17 linear and 2 potential discontinuous B cell epitopes and that

of HTL MEV carries 17 linear and 4 potential discontinuous epitopes. The high range of the PI (Protrusion Index) score of the linear and discontinuous epitopes from CTL and HTL MEV s show high potential of the epitopes to cause humoral immune response (PI score: CTL MEV linear & discontinuous B cell epitopes - 0.511 to 0.828 & 0.664 to 0.767 respectively; HTL MEV linear and discontinuous B cell epitopes - 0.518 to 0.831 & 0.53 to 0.776 respectively) (Supplementary table S4,S5, S6, S7).

Molecular interaction analysis of MEVs with immunological receptor.

Molecular docking and dynamics simulation study of MEVs and TLR-3. Both the refined models of CTL and HTL MEVs were further studied for their molecular interaction with the ectodomain (ECD) of human TLR-3. Therefore, molecular docking of CTL and HTL MEVs model with the TLR-3 crystal structure model (PDB ID: 2A0Z) was performed utilizing the PatchDock tool. Generated docking conformation with highest scores of 20776 and 20350 for CTL and HTL MEVs respectively were chosen for further study. The highest docking score indicates the best geometric shape complementarity fitting conformation of MEVs and the TLR-3 receptor as predicted by the PatchDock tool. Both the CTL and HTL MEVs were fitting into the ectodomain region of TLR-3 after docking (Fig.9A, 9C). The CTL and HTL MEVs have shown to form multiple hydrogen bonds within the ectodomain cavity region of TLR-3.

The B-factor analysis of MEVs-TLR3 complexes was also performed. The B-factor indicates the displacement of the atomic positions from an average

(mean) value i.e. the more flexible an atom is the larger the displacement from

the mean position will be (mean-squares displacement) (Fig.9B, 9D). The B-

factor analysis of the CTL and HTL MEVs bound to the TLR3 receptor shows that

most of the regions of MEVs bound to TLR3 are stable nature in nature. The B-

Factor analysis has been represented by the VIBGYOR color presentation with

blue represents low B-factor and red represents high B-factor. Hence, the results

suggest a stable complex formation tendency for both the CTL and HTL MEVs

with the ectodomain of the human TLR-3 receptor (Fig.9B, 9D).

CTL Multi-epitope vaccine model in complex with TLR3

HTL Multi-epitope vaccine model in complex with TLR3

Figure 9. Molecular Docking study of CTL and HTL MEVs with TLR-3

receptor. (A), (C) The docking complex of CTL-TLR and HTL-TLR3 have been

shown. The TLR3 is shown in cartoon, and the MEVs are shown in ribbon. The

Hydrogen bond formation is shown by cyan lines. **(B), (D)** B-factor presentation

for the docked MEVs to the TLR3 receptor. The presentation is in VIBGYOR

color, with blue showing low B-factor and red show high B-factor. Here most of

the MEV regions are in blue showing low B-factor and hence indicate stable

complex formation with TLR3 receptor. Images were generated by PyMol and

UCSF Chimera (Pettersen et al., 2004).

1037

In-silico **analysis of MEVs for cloning and expression potency**

Analysis of cDNA of both the MEVs for cloning and expression in mammalian

host cell line. Complementary DNA optimized for CTL and HTL expression in the

mammalian host cell line (Human) was generated by utilizing the Java Codon

Adaptation Tool. Further, the generated optimized cDNA's for both the MEVs

were analyzed by utilizing the GenScript Rare Codon Analysis Tool. The analysis

revealed that the codon optimized cDNA of both the CTL and HTL MEVs bear all

the crucial and favorable compositions for high level expression in mammalian

cell line (CTL-MEV: GC content 70.40%, CAI (Codon Adaptation Index) score

1.00 and 0% tandem rare codons; HTL MEV: GC content 69.26%, CAI score

1.00 and 0% tandem rare codons). Ideally, the GC content of a cDNA should be

30% to 70%, CAI score that indicates the possibility of cDNA expression in a

chosen expression system should be between 0.8-1.0, and the tandem rare

codon frequency that indicates the presence of low-frequency codons in cDNA

should be <30%. The tandem rare codons may hinder the proper expression of

the cDNA or even interrupt the translational machinery of the chosen expression

system. Hence as per the GenScript Rare Codon Analysis, the cDNA of both the

MEVs satisfy all the mentioned parameters and are predicted to have high

expression in the mammalian host cell line (Human).

1057

CONCLUSION

In the present study, we have designed and proposed two multi-epitope

vaccines derived from multiple CTL and HTL epitopes against SARS-CoV-2

(COVID19). The chosen CTL and HTL epitopes show significant sequence

overlap with screened linear B cell epitopes. The shortlisted CTL and HTL

epitopes were utilized to design CTI and HTL multi-epitope vaccine. Both the

generated CTL and HTL multi-epitope vaccine tertiary models have shown to

carry potential linear and discontinuous B cell epitopes as well as potential INF-γ

epitopes. Hence the designed MEVs are predicted to have the potential to elicit

humoral as well as cellular immune responses. Since Onchocerca volvulus

activation-associated secreted protein-1 (Ov-ASP-1) binds to the APCs and

trigger pro-inflammatory cytokine production via Toll-like receptor 3 (TLR3), the

truncated (residues 10-153) Ov-ASP-1 has been utilized as an adjuvant at N

terminal of both the CTL and HTL MEVs models. Chosen overlapping clustering

epitopes were validated for their molecular interaction with their respective HLA

allele binders by molecular docking studies. Molecular interaction of the chosen

CTL epitopes within the TAP transporter cavity was also analyzed. Analysis of

the average world population coverage by both the shortlisted CTL and HTL

epitopes combined revealed coverage of 96.10% world population. The

molecular interaction analysis of both the CTL and HTL MEVs with the

immunoreceptor TLR3 have shown very convincing structural fitting of the MEVs

into the ectodomain of TLR3 receptor cavity. The cDNA for both the MEVs were

generated considering codon-biasing for expression in the mammalian host cell

line (Human). Both the cDNA has been optimized in respect of their GC content

and zero tandem rare codons for the cDNA to have high expression possibility in

the mammalian host cell line (Human). Hence for further studies, both the design

of CTL and HTL MEVs could be cloned, expressed and tested for *in-vivo*

validations and animal trials as potential vaccine candidates against SARS-CoV-

2 infection.

1087

1088 **Supplementary table S1. Homology modeling for HLA alleles.** Tertiary
1089 structures of HLA alleles were modeled by homology modeling using SwissModel
1090 server. Templates were chosen with highest sequence identity. Generated
1091 models with acceptable QMEAN value were chosen for further studies.
1092
1093 **Supplementary table S2. INF-γ epitopes from CTL and HTL MEVs.** INF-γ
1094 inducing (POSITIVE) epitopes from CTL and HTL MEVs were screened by using
1095 "Motif and SVM hybrid" (MERCI & SVM) approaches.
1096
1097 **Supplementary table S3. Refinement models of CTL and HTL MEVs.** Both
1098 the CTL and HTL MEVs models were refined by GalaxyWEB server. After
1099 refinement in particular Rama favored residues increased significantly.
1100
1101 **Supplementary table S4. B Cell linear epitopes of CTL MEVs.** Linear B Cell
1102 epitopes predicted by ElliPro (IEDB) from CTL MEVs.
1103
1104 **Supplementary table S5. B Cell discontinuous epitopes of CTL MEVs.**
1105 Discontinuous B Cell epitopes predicted by ElliPro (IEDB) from CTL MEVs.

1106
1107 **Supplementary table S6. B Cell linear epitopes of HTL MEVs.** Linear B Cell
1108 epitopes predicted by ElliPro (IEDB) from HTL MEVs.
1109
1110 **Supplementary table S7. B Cell discontinuous epitopes of HTL MEVs.**
1111 Discontinuous B Cell epitopes predicted by ElliPro (IEDB) from HTL MEVs.
1112
1113 **Supplementary table S8. Shortlisted high scoring CTL epitopes (MHC-I**
1114 **Binding Predictions).** Selected high scoring CTL epitopes and their respective
1115 HLA alleles binders predicted by "MHC-I Binding Predictions" IEDB tool. *In-silico*
1116 analysis have shown all the selected epitopes to be non-toxic (Non-Toxin) as well
1117 as they show significant conservancy and high immunogenicity.
1118
Supplementary table S9. Shortlisted high scoring CTL epitopes (MHC-I
Processing Predictions). Selected high scoring CTL epitopes and their
respective HLA alleles binders predicted by "MHC-I Processing Predictions"
IEDB tool. The screening gives detailed and combined scoring "Total score" for
Proteasomal cleavage/TAP transport/MHC class I combined. *In-silico* analysis
have shown all the selected epitopes to be non-toxic (Non-Toxin) as well as they
show significant conservancy with high immunogenicity.
1126
Supplementary table S10. Shortlisted high scoring HTL epitopes (MHC-II
Binding Predictions). Selected high "Percentile rank" HTL epitopes with their
respective HLA class II alleles binders predicted by the "**MHC-II Binding**
Predictions" tool of IEDB are listed. *In-silico* analysis have shown all the
selected epitopes to be non-toxic (Non-Toxin) as well as they show significant
conservancy.
1133
Supplementary table S11. Shortlisted B Cell epitopes (BepiPred Linear B
Cell Prediction). B cell linear epitopes with length of 4 to 20 amino acids,
predicted by the "BepiPred Linear B Cell Prediction" IEDB tool, from eleven
SARS-CoV-2 proteins, are listed here. *In-silico* analysis have shown all the
selected epitopes to be non-toxic (Non-Toxin) as well as they show significant
amino acid sequence conservancy.
1140

1141 **AUTHOR CONTRIBUTION:**

1142 Protocol design: S.S., M.K.; Methodology performed by S.S., S.V., M.K., R.K.;

1143 Global Economic risk analysis: R.K.B.; Data analysis, scientific writing and

1144 revising the article: S.S., S.V., M.K., R.K., R.K.B., A.K.S., H.J.S., K.P., M.Kolbe.

1145

ADDITIONAL INFORMATION:

Authors declare to have no competing interests.

REFERENCES:

Abele, R. and Tampé, R., 2004. The ABCs of immunology: structure and function of TAP, the transporter associated with antigen processing. Physiology, 19(4), pp.216-224.

Antoniou, A.N., Powis, S.J. and Elliott, T., 2003. Assembly and export of MHC class I peptide ligands. Current opinion in immunology, 15(1), pp.75-81.

Arnold K, Bordoli L, Kopp J, and Schwede T (2006). The SWISS-MODEL Workspace: A web-based environment for protein structure homology modelling. Bioinformatics.,22,195-201.

Bell, J.K., Botos, I., Hall, P.R., Askins, J., Shiloach, J., Segal, D.M. and Davies, D.R., 2005. The molecular structure of the Toll-like receptor 3 ligand-binding domain. Proceedings of the National Academy of Sciences of the United States of America, 102(31), pp.10976-10980.

Benkert, P., Tosatto, S.C. and Schomburg, D., 2008. QMEAN: A comprehensive scoring function for model quality assessment. Proteins: Structure, Function, and Bioinformatics, 71(1), pp.261-277.

Bui H. H, Sidney J, Dinh K, Southwood S, Newman M. J, Sette A. 2006. Predicting population coverage of T-cell epitope-based diagnostics and vaccines. BMC Bioinformatics 17:153.

Bui HH, Sidney J, Li W, Fusseder N, Sette A. 2007. Development of an epitope conservancy analysis tool to facilitate the design of epitope-based diagnostics and vaccines. BMC Bioinformatics 8:361.

Calis JJA, Maybeno M, Greenbaum JA, Weiskopf D, De Silva AD, Sette A, Kesmir C, Peters B. 2013. Properties of MHC class I presented peptides that enhance immunogenicity. PloS Comp. Biol. 8(1):361.

Chen, X., Zaro, J.L. and Shen, W.C., 2013. Fusion protein linkers: property, design and functionality. Advanced drug delivery reviews, 65(10), pp.1357-1369.

Delneste, Y., Beauvillain, C. and Jeannin, P., 2007. Innate immunity: structure and function of TLRs. Medecine sciences: M/S, 23(1), pp.67-73.

Dhanda, S. K., Vir, P. & Raghava, G. P. Designing of interferon-gamma inducing MHC class-II binders. Biol. Direct. 8, 30 (2013).

Dong Xu and Yang Zhang. Improving the Physical Realism and Structural Accuracy of Protein Models by a Two-step Atomic-level Energy Minimization. Biophysical Journal, vol 101, 2525-2534 (2011).

Duhovny D, Nussinov R, Wolfson HJ. Efficient Unbound Docking of Rigid
Molecules. In Gusfield et al., Ed. Proceedings of the 2'nd Workshop on
Algorithms in Bioinformatics(WABI) Rome, Italy, Lecture Notes in
Computer Science 2452, pp. 185-200, Springer Verlag, 2002

Farina, C., Krumbholz, M., Giese, T., Hartmann, G., Aloisi, F. and Meinl, E., 2005.
Preferential expression and function of Toll-like receptor 3 in human
astrocytes. Journal of neuroimmunology, 159(1-2), pp.12-19.

Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S.E., Wilkins, M.R., Appel,
R.D. and Bairoch, A., 2005. Protein identification and analysis tools on the
ExPASy server (pp. 571-607). Humana Press.

Grifoni, A., Sidney, J., Zhang, Y., Scheuermann, R.H., Peters, B. and Sette, A.,
2020. A Sequence Homology and Bioinformatic Approach Can Predict
Candidate Targets for Immune Responses to SARS-CoV-2. Cell Host &
Microbe.

Guo, J., Yang, Y., Xiao, W., Sun, W., Yu, H., Du, L., Lustigman, S., Jiang, S.,
Kou, Z. and Zhou, Y., 2015. A truncated fragment of Ov-ASP-1 consisting
of the core pathogenesis-related-1 (PR-1) domain maintains adjuvanticity
as the full-length protein. Vaccine, 33(16), pp.1974-1980.

Gupta, S., Kapoor, P., Chaudhary, K., Gautam, A., Kumar, R., Raghava, G.P.
and Open Source Drug Discovery Consortium, 2013. In silico approach for
predicting toxicity of peptides and proteins. PLoS One, 8(9), p.e73957.

Hajighahramani, N., Nezafat, N., Eslami, M., Negahdaripour, M., Rahmatabadi,
S.S. and Ghasemi, Y., 2017. Immunoinformatics analysis and in silico
designing of a novel multi-epitope peptide vaccine against Staphylococcus
aureus. Infection, Genetics and Evolution, 48, pp.83-94.

He, Y., Barker, S.J., MacDonald, A.J., Yu, Y., Cao, L., Li, J., Parhar, R., Heck, S.,
Hartmann, S., Golenbock, D.T. and Jiang, S., 2009. Recombinant Ov-
ASP-1, a Th1-biased protein adjuvant derived from the helminth
Onchocerca volvulus, can directly bind and activate antigen-presenting
cells. The Journal of Immunology, 182(7), pp.4005-4016.

Hoof, I., Peters, B., Sidney, J., Pedersen, L.E., Sette, A., Lund, O., Buus, S. and
Nielsen, M., 2009. NetMHCpan, a method for MHC class I binding
prediction beyond humans. Immunogenetics, 61(1), p.1.

Hu W, Li F, Yang X, Li Z, Xia H, Li G, Wang Y, Zhang Z. A flexible peptide linker
enhances the immunoreactivity of two copies HBsAg preS1 (21-47) fusion
protein. J Biotechnol. 2004;107:83–90.

Irini A Doytchinova and Darren R Flower. VaxiJen: a server for prediction of
protective antigens, tumour antigens and subunit vaccines. BMC
Bioinformatics. 2007 8:4.

Ko, J., Park, H., Heo, L., and Seok, C., GalaxyWEB server for protein structure
prediction and refinement, Nucleic Acids Res. 40 (W1), W294-W297
(2012).

Kringelum, J.V., Lundegaard, C., Lund, O. and Nielsen, M., 2012. Reliable B cell
epitope predictions: impacts of method development and improved
benchmarking. PLoS computational biology, 8(12).

Larsen JE, Lund O, Nielsen M. 2006. Improved method for predicting linear B-
cell epitopes. Immunome Res 2:2.

MacDonald, A.J., Cao, L., He, Y., Zhao, Q., Jiang, S. and Lustigman, S., 2005.
rOv-ASP-1, a recombinant secreted protein of the helminth Onchocerca
volvulus, is a potent adjuvant for inducing antibodies to ovalbumin, HIV-1
polypeptide and SARS-CoV peptide antigens. Vaccine, 23(26), pp.3446-
3452.

Morla, S., Makhija, A. & Kumar, S. Synonymous codon usage pattern in
glycoprotein gene of rabies virus. Gene. 584, 1–6 (2016).

Nagpal, G., Gupta, S., Chaudhary, K., Dhanda, S.K., Prakash, S. and Raghava,
G.P., 2015. VaccineDA: Prediction, design and genome-wide screening of
oligodeoxynucleotide-based vaccine adjuvants. Scientific reports, 5,
p.12478.

NCBI protein sequence database: SARS-CoV-2 (Severe acute respiratory
syndrome coronavirus 2) Sequences
(https://www.ncbi.nlm.nih.gov/genbank/sars-cov-2-seqs/)

Nielsen, M., Lundegaard, C. and Lund, O., 2007. Prediction of MHC class II
binding affinity using SMM-align, a novel stabilization matrix alignment
method. BMC bioinformatics, 8(1), p.238.

Oldham, M.L., Grigorieff, N. and Chen, J., 2016. Structure of the Transporter
associated with antigen processing trapped by herpes simplex virus. eLife,
5, p.e21829.

Peters B, Bulik S, Tampe R, Van Endert PM, Holzhutter HG. 2003. Identifying
MHC class I epitopes by predicting the TAP transport efficiency of epitope
precursors. J Immunol171:1741-1749.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M.,
Meng, E.C. and Ferrin, T.E., 2004. UCSF Chimera—a visualization
system for exploratory research and analysis. Journal of computational
chemistry, 25(13), pp.1605-1612.

Ponomarenko JV, Bui H, Li W, Fusseder N, Bourne PE, Sette A, Peters B. 2008.
ElliPro: a new structure-based tool for the prediction of antibody epitopes.
BMC Bioinformatics 9:514.

PyMOL: The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.

Ramakrishnan, C. and Ramachandran, G.N., 1965. Stereochemical criteria for
polypeptide and protein chain conformations: II. Allowed conformations for
a pair of peptide units. Biophysical journal, 5(6), pp.909-933.
Roy, A., Kucukural, A. and Zhang, Y., 2010. I-TASSER: a unified platform for
automated protein structure and function prediction. Nature protocols, 5(4),
p.725.
S.C. Lovell, I.W. Davis, W.B. Arendall III, P.I.W. de Bakker, J.M. Word, M.G.
Prisant, J.S. Richardson and D.C. Richardson (2002) Structure validation
by Calpha geometry: phi,psi and Cbeta deviation. Proteins: Structure,
Function & Genetics. 50: 437-450.
Saha, S. & Raghava, G. AlgPred: prediction of allergenic proteins and mapping
of IgE epitopes. Nucleic. Acids. Res. 34, W202–W209 (2006).
Schneidman-Duhovny D, Inbar Y, Nussinov R, Wolfson HJ. PatchDock and
SymmDock: servers for rigid and symmetric docking. Nucl. Acids. Res. 33:
W363-367, 2005.
Shen, Y., Maupetit, J., Derreumaux, P. and Tufféry, P., 2014. Improved PEP-
FOLD approach for peptide and miniprotein structure prediction. Journal of
chemical theory and computation, 10(10), pp.4745-4758.
Shin, W.H., Lee, G.R., Heo, L., Lee, H. and Seok, C., 2014. Prediction of protein
structure and interaction by GALAXY protein modeling programs. Bio
Design, 2(1), pp.1-11.
Sidney, J., Assarsson, E., Moore, C., Ngo, S., Pinilla, C., Sette, A. and Peters, B.,
2008. Quantitative peptide binding motifs for 19 human and mouse MHC
class I molecules derived using positional scanning combinatorial peptide
libraries. Immunome research, 4(1), p.2.
Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R.,
McWilliam, H., Remmert, M., Söding, J. and Thompson, J.D., and Higgins
D.G., 2011. Fast, scalable generation of high-quality protein multiple
sequence alignments using Clustal Omega. Molecular systems biology,
7(1), p.539.
Srivastava, S., Kamthania, M., Kumar Pandey, R., Kumar Saxena, A., Saxena,
V., Kumar Singh, S., Kumar Sharma, R. and Sharma, N., 2019. Design of
novel multi-epitope vaccines against severe acute respiratory syndrome
validated through multistage molecular interaction and dynamics. Journal
of Biomolecular Structure and Dynamics, 37(16), pp.4345-4360.
Srivastava, S., Kamthania, M., Singh, S., Saxena, A.K. and Sharma, N., 2018.
Structural basis of development of multi-epitope vaccine against middle
east respiratory syndrome using in silico approach. Infection and drug
resistance, 11, p.2377.

Sturniolo, T., Bono, E., Ding, J., Raddrizzani, L., Tuereci, O., Sahin, U., Braxenthaler, M., Gallazzi, F., Protti, M.P., Sinigaglia, F. and Hammer, J., 1999. Generation of tissue-specific and promiscuous HLA ligand databases using DNA microarrays and virtual HLA class II matrices. Nature biotechnology, 17(6), p.555.

Tenzer S, Peters B, Bulik S, Schoor O, Lemmel C, Schatz MM, Kloetzel PM, Rammensee HG, Schild H, Holzhutter HG. 2005. Modeling the MHC class I pathway by combining predictions of proteasomal cleavage, TAP transport and MHC class I binding. Cell Mol Life Sci 62:1025-1037.

Totura, A.L., Whitmore, A., Agnihothram, S., Schäfer, A., Katze, M.G., Heise, M.T. and Baric, R.S., 2015. Toll-like receptor 3 signaling via TRIF contributes to a protective innate immune response to severe acute respiratory syndrome coronavirus infection. MBio, 6(3), pp.e00638-15.

United nations conference on trade and development report 9[th] March 2020 (https://unctad.org/en/pages/PressRelease.aspx?OriginalVersionID=548)

Wang, P., Sidney, J., Kim, Y., Sette, A., Lund, O., Nielsen, M. and Peters, B., 2010. Peptide binding predictions for HLA DR, DP and DQ molecules. BMC bioinformatics, 11(1), p.568.

Wang, Z. and Xu, J., 2013. Predicting protein contact map using evolutionary and physical constraints by integer programming. Bioinformatics, 29(13), pp.i266-i273.

World Health Organization report, Coronavirus disease 2019 (COVID-19) Situation Report – 61, 21[st] March 2020 (https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200321-sitrep-61-covid-19.pdf?sfvrsn=6aa18912_2)

Wu, X., Wu, S., Li, D., Zhang, J., Hou, L., Ma, J., Liu, W., Ren, D., Zhu, Y. and He, F., 2010. Computational identification of rare codons of Escherichia coli based on codon pairs preference. Bmc Bioinformatics, 11(1), p.61.

SUPPLEMENTARY:

Structural basis to design multi-epitope vaccines against Novel Coronavirus 19 (COVID19) infection, the ongoing pandemic emergency: an in silico approach

Sukrit Srivastava*[a,b], Sonia Verma[c], Mohit Kamthania[d], Rupinder Kaur[e], Ruchi Kiran Badyal[f], Ajay Kumar Saxena[b], Ho-Joon Shin[g], Kailash Pandey[c], Michael Kolbe[h,i]

a   Department of Biotechnology, Institute of Bio-Medical Education and Research, Mangalayatan University, 202146 Aligarh, India.
b   Molecular Medicine Lab., School of Life Science, Jawaharlal Nehru University, 110067 New Delhi, India.
c   National Institute of Malaria Research-Indian Council of Medical Research, New Delhi, India
d   Department of Biotechnology, Institute of Applied Medicines and Research, 201206 Ghaziabad, India.
e   Department of Chemistry, Guru Nanak Dev University, Amritsar, India
f   Department of economics, Mangalayata University, India
g   TIDCL, Ajou University, South Korea
h   Department for Structural Infection Biology, Centre for Structural Systems Biology (CSSB) & Helmholtz-Centre for Infection Research, Notkestraße 85, 22607 Hamburg, Germany.
i   Faculty of Mathematics, Informatics and Natural Sciences, University of Hamburg, Rothenbaumchaussee 19, 20148 Hamburg, Germany.

*Correspondence should be addressed to S.S.
(sukrit.srivastav@mangalayatan.edu.in; srivastav.sukrit@gmail.com)

Supplementary table S1. Homology modeling for HLA alleles. Tertiary structures of HLA alleles were modeled by homology modeling using SwissModel server. Templates were chosen with highest sequence identity. Generated models with acceptable QMEAN value were chosen for further studies.

S.No	HLA Class I Allele	Template used for modeling	% sequence identity	QMEAN
1	A*31:01	6o9b.1.A	94.29%	0.73
2	A*11:01	6o9b.1.A	97.50%	1.14
S.No	HLA Class II Allele	Template used for modeling	% sequence identity	QMEAN
3	DRB3*01:01	4is6.1.B	90.10%	(-)1.84
4	DRB1*09:01	1bx2.1.B	87.43%	(-)0.46

Supplementary table S2. INF-γ epitopes from CTL and HTL MEVs. INF-γ inducing (POSITIVE) epitopes from CTL and HTL MEVs were screened by using "Motif and SVM hybrid" (MERCI & SVM) approaches.

S.No	Start-END	Sequence	Method	Result	Score
colspan	CLT Epitopes also predicted to be IFN-gamma epitopes				

colspan CLT Epitopes also predicted to be IFN-gamma epitopes					
S.No	Start-END	Sequence	Method	Result	Score
1	67-82	RWANQCIFGHSPRQQ	MERCI	POSITIVE	2
2	68-83	WANQCIFGHSPRQQR	MERCI	POSITIVE	2
3	69-84	ANQCIFGHSPRQQRE	MERCI	POSITIVE	2
4	70-85	NQCIFGHSPRQQREG	MERCI	POSITIVE	2
5	71-86	QCIFGHSPRQQREGV	MERCI	POSITIVE	2
6	83-98	EGVGENVYAYWSSVS	MERCI	POSITIVE	1
7	84-99	GVGENVYAYWSSVSV	MERCI	POSITIVE	2
8	120-135	LYENNPSNNMTWKVA	MERCI	POSITIVE	1
9	121-136	YENNPSNNMTWKVAG	MERCI	POSITIVE	1
10	122-137	ENNPSNNMTWKVAGQ	MERCI	POSITIVE	2
11	413-428	CTDDNALAYYGGGGS	MERCI	POSITIVE	1
12	428-443	TDDNALAYYGGGGSM	MERCI	POSITIVE	1
13	429-444	DDNALAYYGGGGSMY	MERCI	POSITIVE	1
14	439-454	GGGGSSIINNTVYTK	MERCI	POSITIVE	1
15	440-455	GGGSSIINNTVYTKG	MERCI	POSITIVE	2
16	441-456	GGSSIINNTVYTKGG	MERCI	POSITIVE	3
17	442-457	GSSIINNTVYTKGGG	MERCI	POSITIVE	3
18	443-458	SSIINNTVYTKGGGG	MERCI	POSITIVE	3
19	444-459	SIINNTVYTKGGGGS	MERCI	POSITIVE	3
20	445-460	IINNTVYTKGGGGSL	MERCI	POSITIVE	3
HLT Epitopes also predicted to be IFN-gamma epitopes					
S.No	Start-END	Sequence	Method	Result	Score
1	67-82	RWANQCIFGHSPRQQ	MERCI	POSITIVE	2
2	68-83	WANQCIFGHSPRQQR	MERCI	POSITIVE	2
3	69-84	ANQCIFGHSPRQQRE	MERCI	POSITIVE	2
4	70-85	NQCIFGHSPRQQREG	MERCI	POSITIVE	2
5	71-86	QCIFGHSPRQQREGV	MERCI	POSITIVE	2
6	83-98	EGVGENVYAYWSSVS	MERCI	POSITIVE	1
7	84-99	GVGENVYAYWSSVSV	MERCI	POSITIVE	2
8	120-135	LYENNPSNNMTWKVA	MERCI	POSITIVE	1
9	121-136	YENNPSNNMTWKVAG	MERCI	POSITIVE	1
10	122-137	ENNPSNNMTWKVAGQ	MERCI	POSITIVE	2
11	585-600	GGGGSFAWWTAFVTN	MERCI	POSITIVE	1
12	586-601	GGGSFAWWTAFVTNV	MERCI	POSITIVE	1
13	587-602	GGSFAWWTAFVTNVN	MERCI	POSITIVE	1
14	588-603	GSFAWWTAFVTNVNA	MERCI	POSITIVE	1
15	589-604	SFAWWTAFVTNVNAS	MERCI	POSITIVE	1
16	560-575	FAWWTAFVTNVNASS	MERCI	POSITIVE	1
17	561-576	AWWTAFVTNVNASSG	MERCI	POSITIVE	1
18	806-821	GGGSCTQHQPYVVDD	MERCI	POSITIVE	1
19	807-822	GGSCTQHQPYVVDDP	MERCI	POSITIVE	1
20	808-823	GSCTQHQPYVVDDPC	MERCI	POSITIVE	1

Supplementary table S3. Refinement models of CTL and HTL MEVs. Both the CTL and HTL MEVs models were refined by GalaxyWEB server. After refinement in particular Rama favored residues increased significantly.

Galaxy Refinement for CTL MEV					
Model	GDT-HA	RMSD	Mol Probity	Clash score	Poor rotamers	Rama favored
Initial	1.0000	0.000	3.749	100.5	4.5	75.8
MODEL 1	0.9371	0.459	2.539	23.2	0.4	83.6
Galaxy Refinement for HTL MEV						
Initial	1.0000	0.000	3.554	131.2	2.2	81.3
MODEL 1	0.9552	0.402	2.537	27.9	0.7	87.7

Supplementary table S4. B Cell linear epitopes of CTL MEVs. Linear B Cell epitopes predicted by ElliPro (IEDB) from CTL MEVs.

S.No.	Start	End	Peptide	Number of residues	Score
1	1	97	IVVAVTGYNCPGGKLTALERKKIVGQNNKYRSDLINGKLKNRNGTY MPRGKNMLELTWDCKLESSAQRWANQCIFGHSPRQQREGVGENVYAYWSSV	97	0.828
2	604	704	LRARSVSPKGGGGSFLAFLLFLVGGGGSHFYSKWYIRGGGGSWTAG AAAYYVGGGGSFPNITNLCPFGGGGSNYNYLYRLFRGGGGSNYLYRLFRHHHHHH	101	0.812
3	400	415	LNGSCGSVGGGGSCTD	16	0.693
4	265	277	SSEMVMCGGSLYG	13	0.678
5	422	438	GGGGSCTDDNALAYYGG	17	0.662
6	584	596	FGGGGSRARSVSP	13	0.619
7	219	231	GGGSKPRQKRTAT	13	0.618
8	490	502	WSMATYYGGGGSY	13	0.618
9	294	304	GGGSISNSWLM	11	0.607
10	362	372	FYGGGGSFLFV	11	0.599
11	380	393	GGGSRYFRLTLGVY	14	0.599
12	320	327	KGGGGSQE	8	0.55
13	445	456	GLPWNVVRGGGG	12	0.548
14	99	109	VEGLKKTAGTD	11	0.547
15	342	349	FNVPMEKG	8	0.534
16	521	524	ASFY	4	0.528
17	464	477	VYTKGGGGSLPVNV	14	0.511

The header of the first table reads: CTL Multi-Epitope vaccine model linear epitopes

Supplementary table S5. B Cell discontinuous epitopes of CTL MEVs. Discontinuous B Cell epitopes predicted by ElliPro (IEDB) from CTL MEVs.

S.No.	Residues	Number of residues	Score
1	I1, V2, V3, A4, V5, T6, G7, Y8, N9, C10, P11, G12, G13, K14, L15, T16, A17, L18, E19, R20, K21, K22, I23, V24, G25, Q26, N27, N28, K29, Y30, R31, S32, D33, L34, I35, N36, G37, K38, L39, K40, N41, R42, N43, G44, T45, Y46, M47, P48, R49, G50, K51, N52, M53, L54, E55, L56, T57, W58, D59, C60, K61, L62, E63, S64, S65, A66, Q67, R68, W69, A70, N71, Q72, C73, I74, F75, G76, H77, S78, P79, R80, Q81, Q82, R83, E84, G85, V86, G87, E88, N89, V90, Y91, A92, Y93, W94, S95, S96, V97, S98, V99, E100, G101, L102, K103, K104, T105, G107, T108, D109, K112, W114, W115, N125, P126, S127, N128, N129, K133, V134, A135, G136	120	0.767
2	G160, A187, R188, G189, G219, G220, G221, S222, K223, P224, R225, Q226, R228, T229, A230, T231, G246, G247, G248, G249, S250, G264, S265, S266, E267, M268, V269, M270, C271, G272, G273, S274, L275, Y276, G277, G294, G295, G296, S297, I298, S299, N300, S301, W302, L303, M304, K320, G321, G322, G323, G324, S325, Q326, E327, I328, T341, F342, N343, V344, P345, M346, E347, K348, G349, Y363, G364, G365, G366, G367, S368, F369, L370, F371, V372, A374, G380, G381, G382, S383, R384, Y385, F386, R387, L388, T389, L390, G391, V392, L400, N401, G402, S403, C404, G405, S406, V407, G408, G409, G410, G411, S412, C413, T414, D415, G422, G423, G424, G425, S426, C427, T428, D429, D430, N431, A432, L433, A434, Y435, Y436, G445, L446, P447, W448, N449, V450, V451, R452, G453, G454, G455, G456, G469, G470, G471, S472, L473, P474, V475, N476, V477, W490, M492, A493, T494, Y495, Y496, G497, G498, G499, G500, S501, Y502, A521, S522, F523, Y524, I546, P547, Y548, N549, S550, K564, V565, S566, I567, W568, F584, G585, G586, G587, G588, S589, R590, A591, R592, S593, V594, S595, P596, G598, G599, G600, G601, Q603, L604, R605, A606, R607, S608, V609, S610, P611, K612, G613, G614, G615, G616, S617, F618, L619, A620, F621, L622, L623, F624, L625, V626, G627, G628, G629, G630, S631, H632, F633, Y634, S635, K636, W637, Y638, I639, R640, G641, G642, G643, G644, S645, W646, T647, A648, G649, A650, A651, A652, Y653, Y654, V655, G656, G657, G658, G659, S660, F661, P662, N663, I664, T665, N666, L667, C668, P669, F670, G671, G672, G673, G674, S675, N676, Y677, N678, Y679, L680, Y681, R682, L683, F684, G686, G687, G688, G689, S690, N691, Y692, L693, Y694, R695, R698, H699, H700, H701, H702, H703, H704	282	0.664

The header of the second table reads: CTL Multi-Epitope vaccine model discontinuous epitopes

Supplementary table S6. B Cell linear epitopes of HTL MEVs. Linear B Cell epitopes predicted by ElliPro (IEDB) from HTL MEVs.

S.No.	Start	End	Peptide	Number of residues	Score
1	691	810	INVFAFPFTIYSLLGGGGSKTQSLLIVNNATNVVGGGGSLLIVNNATNVVIK VCGGGGSQSLLIVNNATNVVIKGGGGSSLLIVNNATNVVIKVGGGGSTQ SLLIVNNATNVVIHHHHHH	120	0.831
2	1	138	IVVAVTGYNCPGGKLTALERKKIVGQNNKYRSDLINGKLKNRNGTYMPR GKNMLELTWDCKLESSAQRWANQCIFGHSPRQQREGVGENVYAYWSS VSVEGLKKTAGTDAGKSWWSKLPKLYENNPSNNMTWKVAGQG	138	0.777
3	452	468	VSIWNLDYIINLIGGGG	17	0.713
4	323	333	FVGGGGSESPF	11	0.69
5	474	489	WNLDYIINLIIGGGGS	16	0.685
6	434	444	TVYSHLLLVAA	11	0.679
7	543	551	LYGGGGSCF	9	0.651
8	416	424	TIPIQASLP	9	0.63
9	278	287	SAFFGMSGGG	10	0.608
10	351	360	ILASFSASTS	10	0.597
11	496	508	WNLDYIINLGGGG	13	0.586
12	399	404	PAQYEL	6	0.572
13	378	383	SAPPAQ	6	0.565
14	673	684	FAFPFTIYSLLL	12	0.556
15	515	527	ALITLATCELGGG	13	0.544
16	570	573	LCFL	4	0.54
17	655	660	DDPCPI	6	0.518

Supplementary table S7. B Cell discontinuous epitopes of HTL MEVs. Discontinuous B Cell epitopes predicted by ElliPro (IEDB) from HTL MEVs.

S. No.	Residues	Number of residues	Score
1	C639, P640, I641, D655, D656, P657, C658, P659, I660, H661, F662, F673, A674, F675, P676, F677, T678, I679, Y680, S681, L682, L683, L684, G687, G688, I691, N692, V693, F694, A695, F696, P697, F698, T699, I700, Y701, S702, L703, L704, G705, G706, G707, G708, S709, K710, T711, Q712, S713, L714, L715, I716, V717, N718, N719, A720, T721, N722, V723, V724, G725, G726, G727, G728, S729, L730, L731, I732, V733, N734, N735, A736, T737, N738, V739, V740, I741, K742, V743, C744, G745, G746, G747, G748, S749, Q750, S751, L752, L753, I754, V755, N756, N757, A758, T759, N760, V761, V762, I763, K764, G765, G766, G767, G768, S769, S770, L771, L772, I773, V774, N775, N776, A777, T778, N779, V780, V781, I782, K783, V784, G785, G786, G787, G788, S789, T790, Q791, S792, L793, L794, I795, V796, N797, N798, A799, T800, N801, V802, V803, I804, H805, H806, H807, H808, H809, H810	145	0.776
2	I1, V2, V3, A4, V5, T6, G7, Y8, N9, C10, P11, G12, G13, K14, L15, T16, A17, L18, E19, R20, K21, K22, I23, V24, G25, Q26, N27, N28, K29, Y30, R31, S32, D33, L34, I35, N36, G37, K38, L39, K40, N41, R42, N43, G44, T45, Y46, M47, P48, R49, G50, K51, N52, M53, L54, E55, L56, T57, W58, D59, C60, K61, L62, E63, S64, S65, A66, Q67, R68, W69, A70, N71, Q72, C73, I74, F75, G76, H77, S78, P79, R80, Q81, Q82, R83, E84, G85, V86, G87, E88, N89, V90, Y91, A92, Y93, W94, S95, S96, V97, S98, V99, E100, G101, L102, K103, K104, T105, A106, G107, T108, D109, A110, G111, K112, S113, W114, W115, S116, K117, L118, P119, K120, L121, Y122, E123, N124, N125, P126, S127, N128, N129, M130, T131, W132, K133, V134, A135, G136, Q137, G138, E145, A146, A147, A148, L150, F152, A154, V156, L163, A164, G165, G168, S169	151	0.748
3	S278, A279, F280, F281, G282, M283, S284, G285, G286, G287, F303, G304, G305, A322, F323, V324, G325, G326, G327, G328, S329, E330, S331, P332, F333, I350, I351, L352, A353, S354, F355, S356, A357, S358, T359, S360, S378, A379, P380, P381, A382, Q383, P399, A400, Q401, Y402, E403, L404, G405, G406, G407, T416, I417, P418, I419, Q420, A421, S422, L423, P424, T434, V435, Y436, S437, H438, L439, L440, L441, V442, A444, G446, V452, S453, I454, W455, N456, L457, D458, Y459, I460, I461, N462, L463, I464, G465, G466, G467, G468, W474, N475, L476, D477, Y478, I479, I480, N481, L482, I483, I484, G485, G486, G487, G488, S489, I495, W496, N497, L498, D499, Y500, I501, I502, N503, L504, G505, G506, G507, G508, A515, L516, I517, T518, A520, T521, C522, E523, L538, A539, L543, Y544, G545, G546, G547, G548, S549, C550, F551, L570, C571, F572, L573, F596, L597, L598, F599, P615, Y616, V617, V618, D619	150	0.609
4	L524, G525, G526, G527, G528	5	0.53

Supplementary table S8. Shortlisted high scoring CTL epitopes (MHC-I Binding Predictions). Selected high scoring CTL epitopes and their respective HLA alleles binders predicted by "MHC-I Binding Predictions" IEDB tool. *In-silico* analysis have shown all the selected epitopes to be non-toxic (Non-Toxin) as well as they show significant conservancy and high immunogenicity.

S.No.	Proteins	Peptide	Allele	Start	End	Length	Immunogenicity	Conservancy	Toxicity	Method used	Percentile Rank
							SARS-CoV-2 CTL epitopes (MHC-I Binding Predictions)				
1	E Protein	FLAFVVFLL	A*02:01	20	28	9	0.30188	100.00% (35/35)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.2
2	E Protein	FLAFVVFLL	A*02:03	20	28	9	0.30188	100.00% (35/35)	Non-Toxin	Consensus (ann/comblib)	0.25
3	E Protein	FLAFVVFLLV	A*02:01	20	29	10	0.30526	100.00% (35/35)	Non-Toxin	Consensus (ann/smm)	0.15
4	E Protein	FLAFVVFLLV	A*02:03	20	29	10	0.30526	100.00% (35/35)	Non-Toxin	Consensus (ann/smm)	0.23
5	E Protein	LAFVVFLLV	B*51:01	21	29	9	0.2141	100.00% (35/35)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.2
6	E Protein	LVKPSFYVY	B*15:01	51	59	9	-0.11106	100.00% (35/35)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.2
7	E Protein	RVKNLNSSR	A*31:01	61	69	9	-0.32968	100.00% (35/35)	Non-Toxin	Consensus (ann/smm)	0.16
8	E Protein	SLVKPSFYVY	B*15:01	50	59	10	-0.2443	100.00% (35/35)	Non-Toxin	Consensus (ann/smm)	0.28
9	E Protein	SVLLFLAFV	A*02:06	16	24	9	0.19022	100.00% (35/35)	Non-Toxin	Consensus (ann/smm)	0.33
10	E Protein	TLAILTALR	A*68:01	30	38	9	0.1989	97.14% (34/35)	Non-Toxin	Consensus (ann/smm)	0.21
11	E Protein	VLLFLAFVV	A*02:01	17	25	9	0.26315	100.00% (35/35)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.3
12	E Protein	VSLVKPSFY	A*30:02	49	57	9	-0.25372	100.00% (35/35)	Non-Toxin	Consensus (ann/smm)	0.33
13	M Protein	FIASFRLFAR	A*68:01	26	35	10	0.12185	100.00% (41/41)	Non-Toxin	Consensus (ann/smm)	0.12
14	M Protein	LIFLWLLWPV	A*02:06	51	60	10	0.40176	100.00% (41/41)	Non-Toxin	Consensus (ann/smm)	0.08
15	M Protein	LSYFIASFR	A*68:01	23	31	9	0.21181	100.00% (41/41)	Non-Toxin	Consensus (ann/smm)	0.11
16	M Protein	MWLSYFIASF	A*23:01	21	30	10	0.00197	100.00% (41/41)	Non-Toxin	Consensus (ann/smm)	0.11
17	M Protein	MWLSYFIASFR	A*68:01	21	31	11	0.03554	100.00% (41/41)	Non-Toxin	ann	0.09
18	M Protein	MWLSYFIASFR	A*33:01	21	31	11	0.03554	100.00% (41/41)	Non-Toxin	ann	0.1
19	M Protein	QWNLVIGFLF	A*23:01	19	28	10	0.28076	100.00% (41/41)	Non-Toxin	Consensus (ann/smm)	0.12
20	M Protein	RFLYIIKLIF	A*23:01	44	53	10	0.11728	100.00% (41/41)	Non-Toxin	Consensus (ann/smm)	0.12
21	M Protein	RTRSMWSF	A*32:01	35	42	8	-0.25178	100.00% (41/41)	Non-Toxin	ann	0.1
22	N Protein	AQFAPSASAF	B*15:01	25	34	10	-0.17446	100.00% (40/40)	Non-Toxin	Consensus (ann/smm)	0.16
23	N Protein	AQFAPSASAFF	B*15:01	25	35	11	-0.11074	100.00% (40/40)	Non-Toxin	ann	0.12
24	N Protein	FTALTQHGK	A*68:01	53	61	9	-0.0226	100.00% (40/40)	Non-Toxin	Consensus (ann/smm)	0.18
25	N Protein	KMKDLSPR	A*31:01	30	37	8	-0.22357	100.00% (40/40)	Non-Toxin	ann	0.15
26	N Protein	KTFPPTEPK	A*11:01	11	19	9	0.1306	100.00% (40/40)	Non-Toxin	Consensus (ann/smm)	0.11
27	N Protein	KTFPPTEPK	A*30:01	11	19	9	0.1306	100.00% (40/40)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.2
28	N Protein	RQKRTATKAY	B*15:01	49	58	10	-0.06462	100.00% (40/40)	Non-Toxin	Consensus (ann/smm)	0.2
29	N Protein	SSPDDQIGYY	A*01:01	8	17	10	0.07924	100.00% (40/40)	Non-Toxin	Consensus (ann/smm)	0.2
30	ORF-1ab	ASFYYVWKSY	A*30:02	8	17	10	0.00073	100.00% (36/36)	Non-Toxin	Consensus (ann/smm)	0.07
31	ORF-1ab	CLAYYFMRFRR	A*33:01	49	59	11	0.12614	100.00% (36/36)	Non-Toxin	ann	0.07
32	ORF-1ab	ETISLAGSY	A*26:01	33	41	9	-0.1653	100.00% (36/36)	Non-Toxin	Consensus (ann/smm)	0.1
33	ORF-1ab	IPLMYKGLPW	B*53:01	55	64	10	-0.37784	94.44% (34/36)	Non-Toxin	Consensus (ann/smm)	0.07
34	ORF-1ab	IPLMYKGLPW	B*53:01	55	64	10	-0.37784	94.44% (34/36)	Non-Toxin	Consensus (ann/smm)	0.07
35	ORF-1ab	KMNYQVNGY	A*30:02	36	44	9	-0.06542	97.22% (35/36)	Non-Toxin	Consensus (ann/smm)	0.07
36	ORF-1ab	KMNYQVNGY	A*30:02	36	44	9	-0.06542	97.22% (35/36)	Non-Toxin	Consensus (ann/smm)	0.07
37	ORF-1ab	STNVTIATY	A*30:02	33	41	9	0.25822	100.00% (36/36)	Non-Toxin	Consensus (ann/smm)	0.09

38	ORF-1ab	YAYLRKHFSM	B*08:01	28	37	10	-0.13937	97.22% (35/36)	Non-Toxin	Consensus (ann/smm)	0.07
39	ORF3a	FMRIFTIGTV	A*02:03	4	13	10	0.47908	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.17
40	ORF3a	FTIGTVTLK	A*68:01	8	16	9	0.18024	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.16
41	ORF3a	HVTFFIYNK	A*68:01	17	25	9	0.36278	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.16
42	ORF3a	IIMRLWLCWK	A*03:01	53	62	10	0.27346	96.43% (27/28)	Non-Toxin	Consensus (ann/smm)	0.13
43	ORF3a	VEHVTFFIY	B*44:03	15	23	9	0.3766	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.18
44	ORF3a	YFLQSINFVR	A*33:01	43	52	10	-0.03483	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.2
45	ORF6	AEILLIIM	B*40:01	12	19	8	0.25884	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.41
46	ORF6	FQVTIAEIL	A*02:06	7	15	9	0.38115	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.52
47	ORF6	ILLIIMRTFK	A*03:01	14	23	10	0.2388	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.17
48	ORF6	IMRTFKVSI	A*32:01	18	26	9	-0.09496	100.00% (34/34)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.44
49	ORF6	KVSIWNLDY	A*30:02	23	31	9	0.29343	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.41
50	ORF6	LLIIMRTFK	A*03:01	15	23	9	0.156	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.36
51	ORF6	LSKSLTENKY	A*30:02	40	49	10	-0.24668	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.48
52	ORF6	SIWNLDYII	A*32:01	25	33	9	0.15011	100.00% (34/34)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.2
53	ORF6	VTIAEILLI	B*58:01	9	17	9	0.28951	100.00% (34/34)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.34
54	ORF7a	CVRGTTVLLK	A*03:01	23	32	10	0.14952	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.17
55	ORF7a	EVQELYSPI	A*26:01	22	30	9	-0.09723	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.21
56	ORF7a	EVQELYSPIF	A*26:01	22	31	10	-0.03858	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.22
57	ORF7a	FALTCFSTQF	B*53:01	54	63	10	-0.09369	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.22
58	ORF7a	FLIVAAIVF	B*15:01	31	39	9	0.29611	100.00% (28/28)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.2
59	ORF7a	FLIVAAIVFI	A*02:01	31	40	10	0.38946	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.21
60	ORF7a	ITLCFTLKRK	A*03:01	40	49	10	-0.06825	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.17
61	ORF7a	QELYSPIFLI	B*44:02	24	33	10	0.03838	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.22
62	ORF7a	QELYSPIFLI	B*44:03	24	33	10	0.03838	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.25
63	ORF7a	RSVSPKLFI	B*58:01	10	18	9	-0.30783	100.00% (28/28)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.2
64	ORF7b	DFYLCFLAF	A*23:01	8	16	9	0.05884	100.00% (2/2)	Non-Toxin	Consensus (ann/smm)	0.29
65	ORF7b	FLAFLLFLVL	A*02:01	13	22	10	0.23386	100.00% (2/2)	Non-Toxin	Consensus (ann/smm)	0.26
66	ORF7b	FLLFLVLIML	A*02:01	16	25	10	0.14288	100.00% (2/2)	Non-Toxin	Consensus (ann/smm)	0.3
67	ORF7b	FYLCFLAFLL	A*23:01	9	18	10	0.1745	100.00% (2/2)	Non-Toxin	Consensus (ann/smm)	0.13
68	ORF7b	FYLCFLAFLL	A*24:02	9	18	10	0.1745	100.00% (2/2)	Non-Toxin	Consensus (ann/smm)	0.28
69	ORF7b	IELSLIDFY	B*44:03	2	10	9	0.03153	100.00% (2/2)	Non-Toxin	Consensus (ann/smm)	0.19
70	ORF7b	LVLIMLIIFW	B*53:01	20	29	10	0.25452	100.00% (2/2)	Non-Toxin	Consensus (ann/smm)	0.32
71	ORF7b	MIELSLIDFY	A*01:01	1	10	10	0.06184	100.00% (2/2)	Non-Toxin	Consensus (ann/smm)	0.35
72	ORF7b	MLIIFWFSL	A*02:01	24	32	9	0.50177	100.00% (2/2)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.2
73	ORF7b	SLIDFYLCFL	A*02:01	5	14	10	0.18838	100.00% (2/2)	Non-Toxin	Consensus (ann/smm)	0.14
74	ORF7b	YLCFLAFLL	A*02:01	10	18	9	0.21865	100.00% (2/2)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.3
75	ORF8	CPIHFYSKW	B*53:01	37	45	9	-0.07935	100.00% (34/34)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.2
76	ORF8	CPIHFYSKWY	B*53:01	37	46	10	-0.02216	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.2
77	ORF8	CSFYEDFLEY	A*01:01	32	41	10	0.31272	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.24
78	ORF8	DFLEYHDVR	A*33:01	37	45	9	0.16684	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.13
79	ORF8	FTINCQEPK	A*68:01	16	24	9	-0.04683	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.18
80	ORF8	GARKSAPLI	A*30:01	50	58	9	-0.34031	100.00% (34/34)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.3
81	ORF8	GIITTVAAF	B*15:01	8	16	9	0.2148	100.00% (34/34)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.3
82	ORF8	GSLVVRCSFY	A*30:02	26	35	10	0.00657	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.24

83	ORF8	IQYIDIGNY	A*30:02	1	9	9	0.30442	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.12
84	ORF8	LGIITTVAAF	B*15:01	7	16	10	0.34746	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.31
85	ORF8	LQSCTQHQPY	B*15:01	22	31	10	-0.25674	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.17
86	ORF8	NYTVSCLPF	A*23:01	8	16	9	-0.17355	58.82% (20/34)	Non-Toxin	Consensus (ann/smm)	0.23
87	ORF8	QSCTQHQPY	A*01:01	23	31	9	-0.16503	100.00% (34/34)	Non-Toxin	Consensus (ann/smm)	0.28
88	ORF10	FAFPFTIYSL	A*02:06	7	16	10	0.20414	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.48
89	ORF10	FPFTIYSL	B*51:01	9	16	8	0.06356	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.39
90	ORF10	FPFTIYSLL	B*51:01	9	17	9	0.05708	100.00% (28/28)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.3
91	ORF10	FPFTIYSLLL	B*53:01	9	18	10	0.03149	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.13
92	ORF10	FPFTIYSLLL	B*51:01	9	18	10	0.03149	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.14
93	ORF10	FTIYSLLLCR	A*68:01	11	20	10	-0.18372	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.12
94	ORF10	FTIYSLLLCR	A*33:01	11	20	10	-0.18372	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.35
95	ORF10	MGYINVFAFPF	A*23:01	1	11	11	0.40977	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.41
96	ORF10	TIYSLLLCR	A*68:01	12	20	9	-0.22977	100.00% (28/28)	Non-Toxin	Consensus (ann/smm)	0.47
97	ORF10	YINVFAFPF	B*53:01	3	11	9	0.28259	100.00% (28/28)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.35
98	ORF10	YINVFAFPF	A*32:01	3	11	9	0.28259	100.00% (28/28)	Non-Toxin	Consensus (ann/comblib_sidney2008/smm)	0.4
99	S Protein	FVFLVLLPLV	A*02:06	2	11	10	0.02996	100.00% (41/41)	Non-Toxin	Consensus (ann/smm)	0.1
100	S Protein	GNYNYLYRLFR	A*33:01	27	37	11	0.08205	100.00% (41/41)	Non-Toxin	ann	0.11
101	S Protein	KSFTVEKGIY	A*30:02	24	33	10	0.11812	100.00% (41/41)	Non-Toxin	Consensus (ann/smm)	0.11

Supplementary table S9. Shortlisted high scoring CTL epitopes (MHC-I Processing Predictions). Selected high scoring CTL epitopes and their respective HLA alleles binders predicted by "MHC-I Processing Predictions" IEDB tool. The screening gives detailed and combined scoring "Total score" for Proteasomal cleavage/TAP transport/MHC class I combined. *In-silico* analysis have shown all the selected epitopes to be non-toxic (Non-Toxin) as well as they show significant conservancy with high immunogenicity.

S.No.	Proteins	Allele	Start	End	Peptide Length	Peptide	Immunogenicity	Conservancy	Toxicity	Proteasome Score	TAP Score	MHC Score	Processing Score	Total Score	MHC IC50[nM]
							SARS-CoV-2 CTL epitopes (MHC-I Processing Predictions)								
1	E Protein	B*15:01	50	59	10	SLVKPSFYVY	-0.2443	100.00% (35/35)	Non-Toxin	1.51	1.36	-1.39	2.86	1.47	24.6
2	E Protein	B*15:01	51	59	9	LVKPSFYVY	-0.11106	100.00% (35/35)	Non-Toxin	1.51	1.35	-1.42	2.86	1.44	26.3
3	E Protein	A*30:02	51	59	9	LVKPSFYVY	-0.11106	100.00% (35/35)	Non-Toxin	1.51	1.35	-1.61	2.86	1.25	40.4
4	E Protein	A*02:01	20	28	9	FLAFVVFLL	0.30188	100.00% (35/35)	Non-Toxin	1.45	0.41	-0.81	1.86	1.05	6.5
5	E Protein	B*15:01	34	42	9	LTALRLCAY	0.01886	97.14% (34/35)	Non-Toxin	1.42	1.27	-1.69	2.69	1	49.5
6	M Protein	A*30:02	31	39	9	ATSRTLSYY	-0.11604	100.00% (41/41)	Non-Toxin	1.26	1.34	-1.12	2.6	1.48	13.3
7	M Protein	A*11:01	31	39	9	ATSRTLSYY	-0.11604	100.00% (41/41)	Non-Toxin	1.26	1.34	-1.52	2.6	1.08	32.9
8	M Protein	B*08:01	33	42	10	FARTRSMWSF	-0.12986	100.00% (41/41)	Non-Toxin	1.41	1.12	-1.34	2.53	1.19	22.1
9	M Protein	A*02:01	53	62	10	FLWLLWPVTL	0.31272	100.00% (41/41)	Non-Toxin	1.85	0.46	-1.16	2.31	1.16	14.3
10	M Protein	A*68:01	23	31	9	LSYFIASFR	0.21181	100.00% (41/41)	Non-Toxin	0.84	0.72	-0.46	1.56	1.1	2.9
11	M Protein	A*23:01	23	33	11	LSYFIASFRLF	0.2706	100.00% (41/41)	Non-Toxin	1.25	1.18	-1.36	2.43	1.07	23
12	M Protein	A*23:01	21	30	10	MWLSYFIASF	0.00197	100.00% (41/41)	Non-Toxin	1.38	1.26	-0.91	2.63	1.73	8.1
13	M Protein	A*24:02	21	30	10	MWLSYFIASF	0.00197	100.00% (41/41)	Non-Toxin	1.38	1.26	-1.27	2.63	1.36	18.7
14	M Protein	B*15:01	21	30	10	MWLSYFIASF	0.00197	100.00% (41/41)	Non-Toxin	1.38	1.26	-1.33	2.63	1.3	21.6

15	M Protein	A*23:01	44	53	10	RFLYIIKLIF	0.11728	100.00% (41/41)	Non-Toxin	1.25	1.35	-1.33	2.6	1.27	21.3
16	M Protein	A*30:01	35	42	8	RTRSMWSF	-0.25178	100.00% (41/41)	Non-Toxin	1.41	1.28	-1.51	2.69	1.18	32.7
17	M Protein	A*30:02	51	59	9	SGFAAYSRY	0.00261	100.00% (41/41)	Non-Toxin	1.53	1.17	-1.4	2.69	1.3	25
18	M Protein	B*15:01	44	53	10	SQRVAGDSGF	0.0305	100.00% (41/41)	Non-Toxin	1.33	1.24	-1.35	2.58	1.23	22.4
19	M Protein	A*23:01	24	32	9	SYFIASFRL	0.18333	100.00% (41/41)	Non-Toxin	1.45	0.62	-0.99	2.08	1.09	9.8
20	M Protein	B*35:01	30	38	9	VATSRTLSY	-0.17295	100.00% (41/41)	Non-Toxin	1.34	1.31	-1.38	2.65	1.27	23.9
21	M Protein	A*30:02	56	64	9	YSRYRIGNY	0.21358	100.00% (41/41)	Non-Toxin	1.36	1.37	-1.61	2.73	1.12	40.6
22	N Protein	B*15:01	25	34	10	AQFAPSASAF	-0.17446	100.00% (40/40)	Non-Toxin	1.23	1.25	-0.7	2.48	1.78	5
23	N Protein	B*15:01	25	35	11	AQFAPSASAFF	-0.11074	100.00% (40/40)	Non-Toxin	1.3	1.25	-1.46	2.55	1.08	29
24	N Protein	A*30:02	32	41	10	KDLSPRWYFY	0.14332	100.00% (40/40)	Non-Toxin	1.58	1.21	-1.68	2.79	1.1	48.3
25	N Protein	A*30:02	30	39	10	KMKDLSPRWY	-0.05692	100.00% (40/40)	Non-Toxin	1.31	1.35	-1.56	2.66	1.1	36.2
26	N Protein	A*68:02	48	56	9	NTASWFTAL	0.22775	100.00% (40/40)	Non-Toxin	1.42	0.49	-0.79	1.91	1.12	6.2
27	N Protein	B*15:01	49	58	10	RQKRTATKAY	-0.06462	100.00% (40/40)	Non-Toxin	1.51	1.34	-1.55	2.85	1.3	35.5
28	N Protein	A*01:01	8	17	10	SSPDDQIGYY	0.07924	100.00% (40/40)	Non-Toxin	1.24	1.36	-1.16	2.6	1.45	14.3
29	ORF1ab	A*01:01	32	41	10	ACTDDNALAY	0.10055	100.00% (36/36)	Non-Toxin	1.49	1.33	-0.82	2.82	2	6.6
30	ORF1ab	A*01:01	33	41	9	CTDDNALAY	0.07355	100.00% (36/36)	Non-Toxin	1.49	1.23	-0.45	2.72	2.27	2.8
31	ORF1ab	A*26:01	33	41	9	ETISLAGSY	-0.1653	100.00% (36/36)	Non-Toxin	1.4	1.21	-0.53	2.61	2.08	3.4
32	ORF1ab	B*35:01	2	10	9	FAVDAAKAY	-0.04849	100.00% (36/36)	Non-Toxin	1.73	1.35	-0.46	3.08	2.62	2.9
33	ORF1ab	B*15:01	53	61	9	ILMTARTVY	0.12576	100.00% (36/36)	Non-Toxin	1.65	1.34	-1.1	2.99	1.89	12.5
34	ORF1ab	B*15:01	52	61	10	LILMTARTVY	0.0012	100.00% (36/36)	Non-Toxin	1.65	1.36	-1.23	3.01	1.78	16.9
35	ORF1ab	B*15:01	13	22	10	LMSNLGMPSY	-0.30933	100.00% (36/36)	Non-Toxin	1.55	1.34	-1	2.9	1.89	10.1
36	ORF1ab	B*35:01	1	9	9	LPSLATVAY	0.06748	100.00% (36/36)	Non-Toxin	1.47	1.15	-0.34	2.62	2.28	2.2
37	ORF1ab	A*32:01	2	10	9	RMYIFFASF	0.29328	100.00% (36/36)	Non-Toxin	1.32	1.36	-0.82	2.68	1.86	6.6
38	ORF1ab	A*30:02	2	11	10	RMYIFFASFY	0.32633	100.00% (36/36)	Non-Toxin	1.28	1.53	-0.97	2.81	1.83	9.4
39	ORF1ab	A*03:01	2	11	10	RMYIFFASFY	0.32633	100.00% (36/36)	Non-Toxin	1.28	1.53	-1	2.81	1.8	10.1
40	ORF1ab	B*44:03	26	34	9	SEFSSLPSY	-0.40603	100.00% (36/36)	Non-Toxin	1.49	1.35	-1.03	2.84	1.82	10.6
41	ORF1ab	B*15:01	18	26	9	VMYMGTLSY	-0.21438	100.00% (36/36)	Non-Toxin	1.3	1.5	-0.8	2.79	1.99	6.3
42	ORF1ab	A*03:01	18	26	9	VMYMGTLSY	-0.21438	100.00% (36/36)	Non-Toxin	1.3	1.5	-0.98	2.79	1.82	9.5
43	ORF3a	B*44:03	29	39	11	AGLEAPFLYLY	0.21841	100.00% (28/28)	Non-Toxin	1.51	1.29	-1.63	2.8	1.17	42.9
44	ORF3a	A*23:01	41	50	10	LVYFLQSINF	-0.05419	100.00% (28/28)	Non-Toxin	1.3	1.31	-1.39	2.61	1.23	24.3
45	ORF6	A*30:02	23	31	9	KVSIWNLDY	0.29343	100.00% (34/34)	Non-Toxin	1.21	1.33	-1.46	2.54	1.08	28.8
46	ORF7a	B*35:01	67	75	9	CPDGVKHVY	-0.07008	100.00% (28/28)	Non-Toxin	1.43	1.11	-1.34	2.54	1.2	22
47	ORF7b	A*02:01	26	34	9	IIFWFSLEL	0.2683	100.00% (2/2)	Non-Toxin	1.65	0.55	-1.2	2.21	1.01	15.7
48	ORF7b	A*02:01	5	14	10	SLIDFYLCFL	0.18838	100.00% (2/2)	Non-Toxin	1.7	0.5	-0.74	2.2	1.46	5.5
49	ORF7b	A*02:03	5	14	10	SLIDFYLCFL	0.18838	100.00% (2/2)	Non-Toxin	1.7	0.5	-0.79	2.2	1.41	6.2
50	ORF7b	A*02:06	5	14	10	SLIDFYLCFL	0.18838	100.00% (2/2)	Non-Toxin	1.7	0.5	-0.93	2.2	1.27	8.6
51	ORF8	B*15:01	8	16	9	GIITTVAAF	0.2148	100.00% (34/34)	Non-Toxin	1.27	1.13	-1.26	2.4	1.14	18.1
52	ORF8	A*33:01	40	48	9	HFYSKWYIR	-0.09452	100.00% (34/34)	Non-Toxin	1.08	0.76	-0.68	1.84	1.16	4.8
53	ORF8	A*31:01	40	48	9	HFYSKWYIR	-0.09452	100.00% (34/34)	Non-Toxin	1.08	0.76	-0.79	1.84	1.05	6.2
54	ORF8	B*15:01	7	16	10	LGIITTVAAF	0.34746	100.00% (34/34)	Non-Toxin	1.27	0.97	-1.1	2.24	1.14	12.7
55	ORF10	A*68:01	5	14	10	NVFAFPFTIY	0.40129	100.00% (28/28)	Non-Toxin	1.47	1.4	-1.84	2.86	1.02	69.7
56	S Protein	B*35:01	11	19	9	CVADYSVLY	-0.09595	100.00% (41/41)	Non-Toxin	1.51	1.38	-1.39	2.89	1.5	24.4
57	S Protein	B*35:01	58	66	9	FAMQMAYRF	-0.28061	100.00% (41/41)	Non-Toxin	1.45	1.05	-0.8	2.5	1.7	6.3
58	S Protein	B*58:01	44	53	10	KRSFIEDLLF	0.29624	100.00% (41/41)	Non-Toxin	1.23	1.26	-1.11	2.49	1.37	12.9
59	S Protein	A*23:01	21	30	10	KWPWYIWLGF	0.56424	100.00% (41/41)	Non-Toxin	1.26	1.25	-1.04	2.51	1.47	11
60	S Protein	B*35:01	54	64	11	LQIPFAMQMAY	-0.22124	100.00% (41/41)	Non-Toxin	1.42	1.4	-1.25	2.82	1.57	17.8

61	S Protein	A*01:01	25	33	9	LTDEMIAQY	0.02757	100.00% (41/41)	Non-Toxin	1.21	1.21	-0.72	2.42	1.71	5.2
62	S Protein	B*35:01	55	64	10	QIPFAMQMAY	-0.25308	100.00% (41/41)	Non-Toxin	1.42	1.35	-1.16	2.77	1.61	14.5
63	S Protein	A*30:02	7	15	9	RISNCVADY	-0.02787	100.00% (41/41)	Non-Toxin	1.16	1.47	-1.26	2.63	1.37	18.2
64	S Protein	B*58:01	45	53	9	RSFIEDLLF	0.27446	100.00% (41/41)	Non-Toxin	1.23	1.32	-0.72	2.54	1.82	5.3
65	S Protein	B*15:01	4	13	10	RVYSTGSNVF	-0.23394	100.00% (41/41)	Non-Toxin	1.51	1.32	-1.02	2.83	1.81	10.5
66	S Protein	B*35:01	22	30	9	SANNCTFEY	0.13273	100.00% (41/41)	Non-Toxin	1.18	1.3	-1.11	2.48	1.37	12.8
67	S Protein	A*30:02	28	37	10	YTNSFTRGVY	0.08467	95.12% (39/41)	Non-Toxin	1.36	1.28	-1.26	2.64	1.38	18.2

Supplementary table S10. Shortlisted high scoring HTL epitopes (MHC-II Binding Predictions). Selected high "Percentile rank" HTL epitopes with their respective HLA class II alleles binders predicted by the "**MHC-II Binding Predictions**" tool of IEDB are listed. *In-silico* analysis have shown all the selected epitopes to be non-toxic (Non-Toxin) as well as they show significant conservancy.

SARS-CoV-2 HTL epitopes (MHC-II Binding Predictions)									
S.No	Proteins	Allele	Start	End	Peptide	Conservancy	Toxicity	Method used	Percentile Rank
1	E Protein	DPA1*03:01/DPB1*04:02	20	34	FLAFVVFLLVTLAIL	100.00% (35/35)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.06
2	E Protein	DPA1*03:01/DPB1*04:02	19	33	LFLAFVVFLLVTLAI	100.00% (35/35)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
3	E Protein	DPA1*01:03/DPB1*02:01	19	33	LFLAFVVFLLVTLAI	100.00% (35/35)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
4	E Protein	DPA1*01/DPB1*04:01	19	33	LFLAFVVFLLVTLAI	100.00% (35/35)	Non-Toxin	Consensus (comb.lib./smm)	0.08
5	E Protein	DPA1*02:01/DPB1*01:01	19	33	LFLAFVVFLLVTLAI	100.00% (35/35)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.1
6	E Protein	DPA1*01:03/DPB1*02:01	15	29	NSVLLFLAFVVFLLV	100.00% (35/35)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
7	E Protein	DPA1*03:01/DPB1*04:02	15	29	NSVLLFLAFVVFLLV	100.00% (35/35)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.06
8	E Protein	DPA1*01/DPB1*04:01	15	29	NSVLLFLAFVVFLLV	100.00% (35/35)	Non-Toxin	Consensus (comb.lib./smm)	0.07
9	E Protein	DPA1*02:01/DPB1*01:01	15	29	NSVLLFLAFVVFLLV	100.00% (35/35)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.1
10	E Protein	DPA1*02:01/DPB1*01:01	14	28	VNSVLLFLAFVVFLL	100.00% (35/35)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.1
11	M Protein	DQA1*01:01/DQB1*05:01	49	63	IKLIFLWLLWPVTLA	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.07
12	M Protein	DQA1*01:01/DQB1*05:01	50	64	KLIFLWLLWPVTLAC	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.11
13	M Protein	DRB1*09:01	34	48	RTLSYYKLGASQRVA	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.06
14	M Protein	DRB1*01:01	34	48	RTLSYYKLGASQRVA	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.67
15	M Protein	DRB1*09:01	33	47	SRTLSYYKLGASQRV	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.07
16	M Protein	DRB1*09:01	37	51	SYYKLGASQRVAGDS	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.48
17	M Protein	DRB1*09:01	35	49	TLSYYKLGASQRVAG	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.07
18	M Protein	DRB1*01:01	35	49	TLSYYKLGASQRVAG	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.67
19	M Protein	DPA1*01:03/DPB1*02:01	18	32	VGLMWLSYFIASFRL	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.07
20	N Protein	DQA1*01:02/DQB1*06:02	12	26	ANNAAIVLQLPQGTT	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.2
21	N Protein	DRB1*11:01	12	26	DQIGYYRRATRRIRG	100.00% (40/40)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.42
22	N Protein	DRB5*01:01	12	26	DQIGYYRRATRRIRG	100.00% (40/40)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.58
23	N Protein	DQA1*01:02/DQB1*06:02	7	21	GTRNPANNAAIVLQL	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
24	N Protein	DRB1*07:01	48	62	GTWLTYTGAIKLDDK	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.58
25	N Protein	DRB1*11:01	15	29	GYYRRATRRIRGGDG	100.00% (40/40)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.42
26	N Protein	DRB1*11:01	14	28	IGYYRRATRRIRGGD	100.00% (40/40)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.42
27	N Protein	DQA1*01:02/DQB1*06:02	13	27	NNAAIVLQLPQGTTL	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.42
28	N Protein	DQA1*01:02/DQB1*06:02	10	24	NPANNAAIVLQLPQG	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03

29	N Protein	DQA1*01:02/DQB1*06:02	11	25	PANNAAIVLQLPQGT	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
30	N Protein	DRB1*09:01	23	37	QIAQFAPSASAFFGM	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.01
31	N Protein	DRB1*11:01	13	27	QIGYYRRATRRIRGG	100.00% (40/40)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.39
32	N Protein	DRB5*01:01	13	27	QIGYYRRATRRIRGG	100.00% (40/40)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.58
33	N Protein	DQA1*01:02/DQB1*06:02	9	23	RNPANNAAIVLQLPQ	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
34	N Protein	DRB1*07:01	47	61	SGTWLTYTGAIKLDD	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.58
35	N Protein	DRB1*07:01	45	59	TPSGTWLTYTGAIKL	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.58
36	N Protein	DQA1*01:02/DQB1*06:02	8	22	TRNPANNAAIVLQLP	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
37	N Protein	DRB1*07:01	49	63	TWLTYTGAIKLDDKD	100.00% (40/40)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.58
38	ORF1ab	DPA1*01:03/DPB1*02:01	34	48	AAIMQLFFSYFAVHF	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
39	ORF1ab	DQA1*01:02/DQB1*06:02	45	59	AFASEAARVVRSIFS	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.08
40	ORF1ab	DPA1*01:03/DPB1*02:01	35	49	AIMQLFFSYFAVHFI	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
41	ORF1ab	DRB3*02:02	24	38	ANYIFWRNTNPIQLS	97.22% (35/36)	Non-Toxin	NetMHCIIpan	0.05
42	ORF1ab	DQA1*05:01/DQB1*03:01	35	49	ASIVAGGIVAIVVTC	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
43	ORF1ab	DRB3*02:02	56	70	AWWTAFVTNVNASSS	97.22% (35/36)	Non-Toxin	NetMHCIIpan	0.01
44	ORF1ab	DQA1*05:01/DQB1*03:01	32	46	DISASIVAGGIVAIV	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
45	ORF1ab	DPA1*03:01/DPB1*04:02	54	68	EETKFLTENLLLYID	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
46	ORF1ab	DQA1*05:01/DQB1*02:01	12	26	EIDFLELAMDEFIER	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
47	ORF1ab	DPA1*03:01/DPB1*04:02	55	69	ETKFLTENLLLYIDI	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
48	ORF1ab	DQA1*01:01/DQB1*05:01	48	62	FISNSWLMWLIINLV	94.44% (34/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
49	ORF1ab	DRB1*07:01	19	33	FSAVGNICYTPSKLI	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.06
50	ORF1ab	DRB1*07:01	52	66	FTPLVPFWITIAYII	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
51	ORF1ab	DQA1*01:01/DQB1*05:01	47	61	HFISNSWLMWLIINL	94.44% (34/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
52	ORF1ab	DQA1*05:01/DQB1*02:01	13	27	IDFLELAMDEFIERY	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
53	ORF1ab	DQA1*05:01/DQB1*03:01	33	47	ISASIVAGGIVAIVV	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
54	ORF1ab	DQA1*01:01/DQB1*05:01	49	63	ISNSWLMWLIINLVQ	94.44% (34/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.07
55	ORF1ab	DPA1*03:01/DPB1*04:02	55	69	KLINIIWFLLLSVC	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.09
56	ORF1ab	DPA1*02:01/DPB1*05:01	45	59	KVTLVFLFVAAIFYL	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
57	ORF1ab	DPA1*01:03/DPB1*02:01	33	47	LAAIMQLFFSYFAVH	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.07
58	ORF1ab	DPA1*03:01/DPB1*04:02	53	67	LEETKFLTENLLLYI	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
59	ORF1ab	DPA1*03:01/DPB1*04:02	56	70	LINIIWFLLLSVCL	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.09
60	ORF1ab	DRB1*07:01	55	69	LVPFWITIAYIICIS	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
61	ORF1ab	DQA1*05:01/DQB1*02:01	11	25	MEIDFLELAMDEFIE	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
62	ORF1ab	DPA1*02:01/DPB1*14:01	34	48	MNLKYAISAKNRART	97.22% (35/36)	Non-Toxin	NetMHCIIpan	0.08
63	ORF1ab	DPA1*01:03/DPB1*02:01	37	51	MQLFFSYFAVHFISN	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
64	ORF1ab	DPA1*01:03/DPB1*02:01	3	17	MYIFFASFYYVWKSY	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
65	ORF1ab	DRB1*11:01	25	39	NEFYAYLRKHFSMMI	97.22% (35/36)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.02
66	ORF1ab	DRB3*02:02	25	39	NYIFWRNTNPIQLSS	97.22% (35/36)	Non-Toxin	NetMHCIIpan	0.04
67	ORF1ab	DRB1*07:01	54	68	PLVPFWITIAYIICI	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.02
68	ORF1ab	DQA1*05:01/DQB1*02:01	10	24	QMEIDFLELAMDEFI	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
69	ORF1ab	DPA1*02:01/DPB1*14:01	33	47	QMNLKYAISAKNRAR	97.22% (35/36)	Non-Toxin	NetMHCIIpan	0.07
70	ORF1ab	DRB1*01:01	49	63	QQESPFVMMSAPPAQ	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.1
71	ORF1ab	DQA1*01:01/DQB1*05:01	26	40	QSTQWSLFFFLYENA	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.1
72	ORF1ab	DPA1*01:03/DPB1*02:01	29	43	QWSLFFFLYENAFLP	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
73	ORF1ab	DPA1*01:03/DPB1*02:01	2	16	RMYIFFASFYYVWKS	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
74	ORF1ab	DQA1*05:01/DQB1*03:01	34	48	SASIVAGGIVAIVVT	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
75	ORF1ab	DRB1*07:01	20	34	SAVGNICYTPSKLIE	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.09
76	ORF1ab	DQA1*05:01/DQB1*03:01	36	50	SIVAGGIVAIVVTCL	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03

77	ORF1ab	DPA1*03:01/DPB1*04:02	54	68	SKLINIIIWFLLLSV	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.09
78	ORF1ab	DQA1*01:01/DQB1*05:01	50	64	SNSWLMWLIINLVQM	94.44% (34/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.07
79	ORF1ab	DQA1*05:01/DQB1*02:01	9	23	SQMEIDFLELAMDEF	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
80	ORF1ab	DPA1*01:03/DPB1*02:01	27	41	STQWSLFFFLYENAF	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
81	ORF1ab	DQA1*01:01/DQB1*05:01	27	41	STQWSLFFFLYENAF	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.1
82	ORF1ab	DPA1*03:01/DPB1*04:02	56	70	TKFLTENLLLYIDIN	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
83	ORF1ab	DPA1*03:01/DPB1*04:02	52	66	TLEETKFLTENLLLY	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
84	ORF1ab	DRB1*07:01	53	67	TPLVPFWITIAYIIC	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
85	ORF1ab	DPA1*01:03/DPB1*02:01	28	42	TQWSLFFFLYENAFL	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.04
86	ORF1ab	DQA1*01:01/DQB1*05:01	28	42	TQWSLFFFLYENAFL	97.22% (35/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.09
87	ORF1ab	DRB1*11:01	24	38	VNEFYAYLRKHFSMM	97.22% (35/36)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.05
88	ORF1ab	DRB1*01:01	48	62	VQQESPFVMMSAPPA	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.1
89	ORF1ab	DQA1*01:02/DQB1*06:02	44	58	YAFASEAARVVRSIF	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.08
90	ORF1ab	DPA1*01:03/DPB1*02:01	4	18	YIFFASFYYVWKSYV	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
91	ORF1ab	DRB3*02:02	26	40	YIFWRNTNPIQLSSY	97.22% (35/36)	Non-Toxin	NetMHCIIpan	0.05
92	ORF1ab	DPA1*01:03/DPB1*02:01	21	35	YILFTRFFYVLGLAA	100.00% (36/36)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.08
93	ORF3a	DPA1*01/DPB1*04:01	33	47	APFLYLYALVYFLQS	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm)	0.12
94	ORF3a	DPA1*02:01/DPB1*14:01	27	41	DFVRATATIPIQASL	100.00% (28/28)	Non-Toxin	NetMHCIIpan	0.12
95	ORF3a	DPA1*01:03/DPB1*02:01	12	26	DTGVEHVTFFIYNKI	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	1.1
96	ORF3a	DPA1*02:01/DPB1*05:01	12	26	DTGVEHVTFFIYNKI	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	1.6
97	ORF3a	DRB1*04:05	20	34	FFIYNKIVDEPEEHV	100.00% (28/28)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.94
98	ORF3a	DPA1*01/DPB1*04:01	35	49	FLYLYALVYFLQSIN	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm)	0.12
99	ORF3a	DPA1*01:03/DPB1*02:01	14	28	GVEHVTFFIYNKIVD	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	1.2
100	ORF3a	DRB1*04:05	17	31	HVTFFIYNKIVDEPE	100.00% (28/28)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.84
101	ORF3a	DPA1*01/DPB1*04:01	34	48	PFLYLYALVYFLQSI	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm)	0.12
102	ORF3a	DRB1*04:05	19	33	TFFIYNKIVDEPEEH	100.00% (28/28)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.81
103	ORF3a	DPA1*01:03/DPB1*02:01	13	27	TGVEHVTFFIYNKIV	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.93
104	ORF3a	DPA1*02:01/DPB1*05:01	13	27	TGVEHVTFFIYNKIV	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	1.4
105	ORF3a	DRB1*04:05	18	32	VTFFIYNKIVDEPEE	100.00% (28/28)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.81
106	ORF3a	DPA1*01/DPB1*04:01	37	51	YLYALVYFLQSINFV	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm)	0.14
107	ORF6	DRB1*15:01	13	27	EILLIIMRTFKVSIW	100.00% (34/34)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.1
108	ORF6	DRB1*15:01	14	28	ILLIIMRTFKVSIWN	100.00% (34/34)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.1
109	ORF6	DQA1*01:01/DQB1*05:01	24	38	VSIWNLDYIINLIIK	100.00% (34/34)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
110	ORF7a	DRB1*01:01	2	16	KIILFLALITLATCE	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.16
111	ORF7b	DRB4*01:01	15	29	AFLLFLVLIMLIIFW	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.14
112	ORF7b	DPA1*03:01/DPB1*04:02	8	22	DFYLCFLAFLLFLVL	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
113	ORF7b	DPA1*01:03/DPB1*02:01	8	22	DFYLCFLAFLLFLVL	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.08
114	ORF7b	DPA1*01/DPB1*04:01	8	22	DFYLCFLAFLLFLVL	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm)	0.09
115	ORF7b	DPA1*03:01/DPB1*04:02	13	27	FLAFLLFLVLIMLII	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.06
116	ORF7b	DRB4*01:01	13	27	FLAFLLFLVLIMLII	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.14
117	ORF7b	DRB4*01:01	16	30	FLLFLVLIMLIIFWF	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.19
118	ORF7b	DPA1*03:01/DPB1*04:02	9	23	FYLCFLAFLLFLVLI	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
119	ORF7b	DPA1*01:03/DPB1*02:01	9	23	FYLCFLAFLLFLVLI	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.06
120	ORF7b	DPA1*01/DPB1*04:01	9	23	FYLCFLAFLLFLVLI	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm)	0.09
121	ORF7b	DPA1*01:03/DPB1*02:01	7	21	IDFYLCFLAFLLFLV	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.06
122	ORF7b	DPA1*03:01/DPB1*04:02	7	21	IDFYLCFLAFLLFLV	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.16
123	ORF7b	DPA1*03:01/DPB1*04:02	14	28	LAFLLFLVLIMLIIF	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.08
124	ORF7b	DRB4*01:01	14	28	LAFLLFLVLIMLIIF	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.12

125	ORF7b	DPA1*01:03/DPB1*02:01	6	20	LIDFYLCFLAFLLFL	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.05
126	ORF7b	DPA1*01:03/DPB1*02:01	4	18	LSLIDFYLCFLAFLL	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.15
127	ORF7b	DQA1*01:01/DQB1*05:01	20	34	LVLIMLIIFWFSLEL	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.14
128	ORF7b	DPA1*01:03/DPB1*02:01	5	19	SLIDFYLCFLAFLLF	100.00% (2/2)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.07
129	ORF8	DQA1*01:01/DQB1*05:01	98	112	LVVRCSFYEDFLEYH	100.00% (34/34)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.45
130	ORF8	DRB3*01:01	27	41	QHQPYVVDDPCPIHF	100.00% (34/34)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.08
131	ORF8	DRB3*01:01	26	40	TQHQPYVVDDPCPIH	100.00% (34/34)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.08
132	ORF10	DPA1*01/DPB1*04:01	7	21	FAFPFTIYSLLLCRM	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm)	0.56
133	ORF10	DPA1*02:01/DPB1*01:01	7	21	FAFPFTIYSLLLCRM	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	1.1
134	ORF10	DPA1*03:01/DPB1*04:02	7	21	FAFPFTIYSLLLCRM	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	1.1
135	ORF10	DPA1*01:03/DPB1*02:01	4	18	INVFAFPFTIYSLLL	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.29
136	ORF10	DPA1*01/DPB1*04:01	4	18	INVFAFPFTIYSLLL	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm)	0.46
137	ORF10	DPA1*02:01/DPB1*01:01	4	18	INVFAFPFTIYSLLL	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.72
138	ORF10	DPA1*01:03/DPB1*02:01	5	19	NVFAFPFTIYSLLLC	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.4
139	ORF10	DPA1*01/DPB1*04:01	5	19	NVFAFPFTIYSLLLC	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm)	0.56
140	ORF10	DPA1*02:01/DPB1*01:01	5	19	NVFAFPFTIYSLLLC	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.69
141	ORF10	DPA1*01/DPB1*04:01	6	20	VFAFPFTIYSLLLCR	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm)	0.56
142	ORF10	DPA1*02:01/DPB1*01:01	6	20	VFAFPFTIYSLLLCR	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.66
143	ORF10	DPA1*01:03/DPB1*02:01	6	20	VFAFPFTIYSLLLCR	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.71
144	ORF10	DPA1*03:01/DPB1*04:02	6	20	VFAFPFTIYSLLLCR	100.00% (28/28)	Non-Toxin	Consensus (comb.lib./smm/nn)	1.1
145	S protein	DRB3*01:01	47	61	ADSFVIRGDEVRQIA	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.49
146	S protein	DRB1*07:01	13	27	AIPTNFTISVTTEIL	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.4
147	S protein	DRB3*01:01	48	62	DSFVIRGDEVRQIAP	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.51
148	S protein	DRB3*02:02	42	56	EGVFVSNGTHWFVTQ	100.00% (41/41)	Non-Toxin	NetMHCIIpan	0.21
149	S protein	DRB5*01:01	52	66	FVFKNIDGYFKIYSK	100.00% (41/41)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.17
150	S protein	DRB1*11:01	27	41	GNYNYLYRLFRKSNL	100.00% (41/41)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.22
151	S protein	DRB1*07:01	12	26	IAIPTNFTISVTTEI	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.47
152	S protein	DRB5*01:01	23	37	INITRFQTLLALHRS	97.56% (40/41)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.32
153	S protein	DRB5*01:01	25	39	ITRFQTLLALHRSYL	97.56% (40/41)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.26
154	S protein	DPA1*02:01/DPB1*14:01	25	39	ITRFQTLLALHRSYL	97.56% (40/41)	Non-Toxin	NetMHCIIpan	0.43
155	S protein	DRB1*13:02	48	62	LIVNNATNVVIKVCE	100.00% (41/41)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.03
156	S protein	DRB1*01:01	23	37	LSFELLHAPATVCGP	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
157	S protein	DRB5*01:01	24	38	NITRFQTLLALHRSY	97.56% (40/41)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.32
158	S protein	DPA1*02:01/DPB1*14:01	24	38	NITRFQTLLALHRSY	97.56% (40/41)	Non-Toxin	NetMHCIIpan	0.45
159	S protein	DRB1*07:01	15	29	PTNFTISVTTEILPV	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.51
160	S protein	DPA1*02:01/DPB1*14:01	31	45	QLIRAAEIRASANLA	100.00% (41/41)	Non-Toxin	NetMHCIIpan	0.31
161	S protein	DPA1*02:01/DPB1*14:01	30	44	QQLIRAAEIRASANL	100.00% (41/41)	Non-Toxin	NetMHCIIpan	0.2
162	S protein	DRB5*01:01	50	64	REFVFKNIDGYFKIY	100.00% (41/41)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.17
163	S protein	DRB3*02:02	41	55	REGVFVSNGTHWFVT	100.00% (41/41)	Non-Toxin	NetMHCIIpan	0.2
164	S protein	DRB1*01:01	24	38	SFELLHAPATVCGPK	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.09
165	S protein	DRB3*01:01	49	63	SFVIRGDEVRQIAPG	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.51
166	S protein	DRB1*13:02	42	56	SKTQSLLIVNNATNV	100.00% (41/41)	Non-Toxin	Consensus (smm/nn/sturniolo)	0.03
167	S protein	DRB1*01:01	22	36	VLSFELLHAPATVCG	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
168	S protein	DRB1*01:01	21	35	VVLSFELLHAPATVC	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.03
169	S protein	DRB1*01:01	20	34	VVVLSFELLHAPATV	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.09
170	S protein	DRB3*01:01	46	60	YADSFVIRGDEVRQI	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.49
171	S protein	DQA1*05:01/DQB1*03:01	25	39	YIWLGFIAGLIAIVM	100.00% (41/41)	Non-Toxin	Consensus (comb.lib./smm/nn)	0.51

Supplementary table S11. Shortlisted B Cell epitopes (BepiPred Linear B Cell Prediction). B cell linear epitopes with length of 4 to 20 amino acids, predicted by the "BepiPred Linear B Cell Prediction" IEDB tool, from eleven SARS-CoV-2 proteins, are listed here. *In-silico* analysis have shown all the selected epitopes to be non-toxic (Non-Toxin) as well as they show significant amino acid sequence conservancy.

S.No.	Proteins	Start	End	Peptide	Conservancy	Toxicity	Length
				SARS-CoV-2 B Cell epitopes (Bepipred Linear Epitope Prediction)			
1	E protein	6	9	SEET	100.00% (35/35)	Non-Toxin	4
2	E protein	57	71	YVYSRVKNLNSSRVP	100.00% (35/35)	Non-Toxin	15
3	M protein	132	137	PLLESE	100.00% (41/41)	Non-Toxin	6
4	M protein	180	191	KLGASQRVAGDS	100.00% (41/41)	Non-Toxin	12
5	M protein	5	20	NGTITVEELKKLLEQW	100.00% (41/41)	Non-Toxin	16
6	M protein	199	218	YRIGNYKLNTDHSSSSDNIA	100.00% (41/41)	Non-Toxin	20
7	N protein	343	348	DPNFKD	97.50% (39/40)	Non-Toxin	6
8	N protein	119	127	AGLPYGANK	100.00% (40/40)	Non-Toxin	9
9	N protein	4	15	NGPQNQRNAPRI	100.00% (40/40)	Non-Toxin	12
10	N protein	404	416	SKQLQQSMSSADS	100.00% (40/40)	Non-Toxin	13
11	ORF1ab	95	98	IQYG	100.00% (36/36)	Non-Toxin	4
12	ORF1ab	1064	1067	YLKH	100.00% (36/36)	Non-Toxin	4
13	ORF1ab	1318	1321	RKVP	100.00% (36/36)	Non-Toxin	4
14	ORF1ab	1831	1834	YQCG	100.00% (36/36)	Non-Toxin	4
15	ORF1ab	2112	2115	ELSR	100.00% (36/36)	Non-Toxin	4
16	ORF1ab	2457	2460	TFCA	100.00% (36/36)	Non-Toxin	4
17	ORF1ab	2763	2766	GKIV	100.00% (36/36)	Non-Toxin	4
18	ORF1ab	4528	4531	EGNC	97.22% (35/36)	Non-Toxin	4
19	ORF1ab	5221	5224	LPYP	97.22% (35/36)	Non-Toxin	4
20	ORF1ab	5337	5340	SLRC	97.22% (35/36)	Non-Toxin	4
21	ORF1ab	5516	5519	KVQI	97.22% (35/36)	Non-Toxin	4
22	ORF1ab	6255	6258	HDIG	97.22% (35/36)	Non-Toxin	4
23	ORF1ab	6974	6977	WNAD	97.22% (35/36)	Non-Toxin	4
24	ORF1ab	113	117	EIPVA	97.22% (35/36)	Non-Toxin	5
25	ORF1ab	1124	1128	KGEDI	100.00% (36/36)	Non-Toxin	5
26	ORF1ab	1181	1185	NLYDK	100.00% (36/36)	Non-Toxin	5
27	ORF1ab	1689	1693	KFNPP	100.00% (36/36)	Non-Toxin	5
28	ORF1ab	1756	1760	GQQQT	100.00% (36/36)	Non-Toxin	5
29	ORF1ab	1936	1940	KFADD	100.00% (36/36)	Non-Toxin	5
30	ORF1ab	2128	2132	AAVNS	100.00% (36/36)	Non-Toxin	5
31	ORF1ab	2547	2551	SKCEE	100.00% (36/36)	Non-Toxin	5
32	ORF1ab	2903	2907	IEYTD	100.00% (36/36)	Non-Toxin	5
33	ORF1ab	3035	3039	IQPIG	100.00% (36/36)	Non-Toxin	5
34	ORF1ab	3243	3247	NSGSD	100.00% (36/36)	Non-Toxin	5
35	ORF1ab	3553	3557	EFTPF	100.00% (36/36)	Non-Toxin	5
36	ORF1ab	3899	3903	LLAKD	100.00% (36/36)	Non-Toxin	5
37	ORF1ab	3932	3936	EEMLD	100.00% (36/36)	Non-Toxin	5

38	ORF1ab	4017	4021	RSEDK	100.00% (36/36)	Non-Toxin	5
39	ORF1ab	5288	5292	TGHML	97.22% (35/36)	Non-Toxin	5
40	ORF1ab	5877	5881	AHSCN	97.22% (35/36)	Non-Toxin	5
41	ORF1ab	6736	6740	QTGSS	97.22% (35/36)	Non-Toxin	5
42	ORF1ab	18	23	LPVLQV	100.00% (36/36)	Non-Toxin	6
43	ORF1ab	31	36	FGDSVE	97.22% (35/36)	Non-Toxin	6
44	ORF1ab	76	81	ARTAPH	100.00% (36/36)	Non-Toxin	6
45	ORF1ab	208	213	AGKASC	100.00% (36/36)	Non-Toxin	6
46	ORF1ab	220	225	DFIDTK	100.00% (36/36)	Non-Toxin	6
47	ORF1ab	1813	1818	QYELKH	100.00% (36/36)	Non-Toxin	6
48	ORF1ab	1960	1965	FPDLNG	100.00% (36/36)	Non-Toxin	6
49	ORF1ab	2297	2302	SLDTYP	100.00% (36/36)	Non-Toxin	6
50	ORF1ab	2312	2317	SSFKWD	100.00% (36/36)	Non-Toxin	6
51	ORF1ab	2414	2419	KRNRAT	100.00% (36/36)	Non-Toxin	6
52	ORF1ab	2442	2447	KGFCKL	100.00% (36/36)	Non-Toxin	6
53	ORF1ab	2701	2706	AQVAKS	100.00% (36/36)	Non-Toxin	6
54	ORF1ab	3881	3886	VESSSK	100.00% (36/36)	Non-Toxin	6
55	ORF1ab	5905	5910	LYDKLQ	97.22% (35/36)	Non-Toxin	6
56	ORF1ab	6496	6501	ENKTTL	97.22% (35/36)	Non-Toxin	6
57	ORF1ab	6873	6878	DKGVAP	97.22% (35/36)	Non-Toxin	6
58	ORF1ab	1092	1098	ATNGPLK	100.00% (36/36)	Non-Toxin	7
59	ORF1ab	2678	2684	NKVENMT	100.00% (36/36)	Non-Toxin	7
60	ORF1ab	2747	2753	TRQVVNV	100.00% (36/36)	Non-Toxin	7
61	ORF1ab	2861	2867	EVGFVVP	100.00% (36/36)	Non-Toxin	7
62	ORF1ab	2924	2930	DASGKPV	100.00% (36/36)	Non-Toxin	7
63	ORF1ab	3009	3015	DYYRSLP	100.00% (36/36)	Non-Toxin	7
64	ORF1ab	3169	3175	GVSFSTF	100.00% (36/36)	Non-Toxin	7
65	ORF1ab	3269	3275	MAFPSGK	100.00% (36/36)	Non-Toxin	7
66	ORF1ab	3825	3831	SQGLLPP	100.00% (36/36)	Non-Toxin	7
67	ORF1ab	3858	3864	VQSKMSD	100.00% (36/36)	Non-Toxin	7
68	ORF1ab	3924	3930	AVDINKL	100.00% (36/36)	Non-Toxin	7
69	ORF1ab	3992	3998	DRDAAMQ	100.00% (36/36)	Non-Toxin	7
70	ORF1ab	4196	4202	FPKSDGT	100.00% (36/36)	Non-Toxin	7
71	ORF1ab	4336	4342	HPNPKGF	100.00% (36/36)	Non-Toxin	7
72	ORF1ab	5104	5110	GNKIADK	97.22% (35/36)	Non-Toxin	7
73	ORF1ab	5317	5323	YTPHTVL	97.22% (35/36)	Non-Toxin	7
74	ORF1ab	5524	5530	FEKGDYG	97.22% (35/36)	Non-Toxin	7
75	ORF1ab	5857	5863	VDSSQGS	97.22% (35/36)	Non-Toxin	7
76	ORF1ab	6078	6084	MYKGLPW	94.44% (34/36)	Non-Toxin	7
77	ORF1ab	6223	6229	IIGDELK	97.22% (35/36)	Non-Toxin	7
78	ORF1ab	6801	6807	QAWQPGV	97.22% (35/36)	Non-Toxin	7
79	ORF1ab	7012	7018	KPREQID	94.44% (34/36)	Non-Toxin	7
80	ORF1ab	5	12	VPGFNEKT	100.00% (36/36)	Non-Toxin	8
81	ORF1ab	175	182	RELNGGAY	100.00% (36/36)	Non-Toxin	8
82	ORF1ab	813	820	FTLKGGAP	100.00% (36/36)	Non-Toxin	8
83	ORF1ab	1720	1727	KTVGELGD	100.00% (36/36)	Non-Toxin	8

84	ORF1ab	2603	2610	NVPMEKLK	100.00% (36/36)	Non-Toxin	8
85	ORF1ab	4255	4262	GNATEVPA	100.00% (36/36)	Non-Toxin	8
86	ORF1ab	4286	4293	SGGQPITN	100.00% (36/36)	Non-Toxin	8
87	ORF1ab	6756	6763	EIIKSQDL	97.22% (35/36)	Non-Toxin	8
88	ORF1ab	860	868	YTVELGTEV	100.00% (36/36)	Non-Toxin	9
89	ORF1ab	1301	1309	IPTKKAGGT	100.00% (36/36)	Non-Toxin	9
90	ORF1ab	1333	1341	GLNGYTVEE	100.00% (36/36)	Non-Toxin	9
91	ORF1ab	1423	1431	YFYTSKTTV	100.00% (36/36)	Non-Toxin	9
92	ORF1ab	1586	1594	MSMTYGQQF	100.00% (36/36)	Non-Toxin	9
93	ORF1ab	2621	2629	AKNVSLDNV	100.00% (36/36)	Non-Toxin	9
94	ORF1ab	2814	2822	GGVTRDIAS	100.00% (36/36)	Non-Toxin	9
95	ORF1ab	4751	4759	VNLHSSRLS	97.22% (35/36)	Non-Toxin	9
96	ORF1ab	5125	5133	RNRDVDTDF	97.22% (35/36)	Non-Toxin	9
97	ORF1ab	5261	5269	LTKHPNQEY	97.22% (35/36)	Non-Toxin	9
98	ORF1ab	5465	5473	TEETFKLSY	97.22% (35/36)	Non-Toxin	9
99	ORF1ab	5967	5975	IPGIPKDMT	97.22% (35/36)	Non-Toxin	9
100	ORF1ab	6351	6359	FHTPAFDKS	97.22% (35/36)	Non-Toxin	9
101	ORF1ab	6464	6472	KGHFDGQQG	97.22% (35/36)	Non-Toxin	9
102	ORF1ab	6636	6644	GVVQQLPET	97.22% (35/36)	Non-Toxin	9
103	ORF1ab	6674	6682	IERYKLEGY	94.44% (34/36)	Non-Toxin	9
104	ORF1ab	56	65	VEKGVLPQLE	100.00% (36/36)	Non-Toxin	10
105	ORF1ab	245	254	SEKSYELQTP	100.00% (36/36)	Non-Toxin	10
106	ORF1ab	373	382	CHNSEVGPEH	100.00% (36/36)	Non-Toxin	10
107	ORF1ab	720	729	CVKSREETGL	100.00% (36/36)	Non-Toxin	10
108	ORF1ab	1510	1519	SYKDWSYSGQ	100.00% (36/36)	Non-Toxin	10
109	ORF1ab	1851	1860	ALLTKSSEYK	100.00% (36/36)	Non-Toxin	10
110	ORF1ab	1943	1952	QLTGYKKPAS	100.00% (36/36)	Non-Toxin	10
111	ORF1ab	2076	2085	ILKPANNSLK	100.00% (36/36)	Non-Toxin	10
112	ORF1ab	3500	3509	YNYEPLTQDH	100.00% (36/36)	Non-Toxin	10
113	ORF1ab	3965	3974	EQAVANGDSE	100.00% (36/36)	Non-Toxin	10
114	ORF1ab	4312	4321	PEANMDQESF	100.00% (36/36)	Non-Toxin	10
115	ORF1ab	4413	4422	PCGTGTSTDV	97.22% (35/36)	Non-Toxin	10
116	ORF1ab	4448	4457	FQEKDEDDNL	97.22% (35/36)	Non-Toxin	10
117	ORF1ab	4468	4477	TFSNYQHEET	97.22% (35/36)	Non-Toxin	10
118	ORF1ab	4493	4502	FFKFRIDGDM	97.22% (35/36)	Non-Toxin	10
119	ORF1ab	4936	4945	LKYAISAKNR	97.22% (35/36)	Non-Toxin	10
120	ORF1ab	5162	5171	YASQGLVASI	97.22% (35/36)	Non-Toxin	10
121	ORF1ab	5192	5201	WTETDLTKGP	97.22% (35/36)	Non-Toxin	10
122	ORF1ab	5990	5999	QVNGYPNMFI	97.22% (35/36)	Non-Toxin	10
123	ORF1ab	6540	6549	YKRDAPAHIS	97.22% (35/36)	Non-Toxin	10
124	ORF1ab	6578	6587	RVDGQVDLFR	97.22% (35/36)	Non-Toxin	10
125	ORF1ab	1652	1662	HTKKWKYPQVN	100.00% (36/36)	Non-Toxin	11
126	ORF1ab	2478	2488	KRPINPTDQSS	100.00% (36/36)	Non-Toxin	11
127	ORF1ab	4708	4718	LFSTVFPPTSF	97.22% (35/36)	Non-Toxin	11
128	ORF1ab	5493	5503	VGKPRPPLNRN	97.22% (35/36)	Non-Toxin	11
129	ORF1ab	6329	6339	RVLSNLNLPGC	97.22% (35/36)	Non-Toxin	11

130	ORF1ab	6558	6568	DIAKKPTETIC	97.22% (35/36)	Non-Toxin	11
131	ORF1ab	387	398	YHNESGLKTILR	100.00% (36/36)	Non-Toxin	12
132	ORF1ab	743	754	EGETLPTEVLTE	100.00% (36/36)	Non-Toxin	12
133	ORF1ab	1973	1984	KHYTPSFKKGAK	100.00% (36/36)	Non-Toxin	12
134	ORF1ab	2638	2649	RQGFVDSDVETK	100.00% (36/36)	Non-Toxin	12
135	ORF1ab	3195	3206	SDVLLPLTQYNR	100.00% (36/36)	Non-Toxin	12
136	ORF1ab	4138	4149	KLQNNELSPVAL	100.00% (36/36)	Non-Toxin	12
137	ORF1ab	4772	4783	MHAASGNLLLDK	97.22% (35/36)	Non-Toxin	12
138	ORF1ab	5068	5079	KPGGTSSGDATT	97.22% (35/36)	Non-Toxin	12
139	ORF1ab	5300	5311	TNDNTSRYWEPE	97.22% (35/36)	Non-Toxin	12
140	ORF1ab	5371	5382	PYVCNAPGCDVT	97.22% (35/36)	Non-Toxin	12
141	ORF1ab	6018	6029	GCHATREAVGTN	97.22% (35/36)	Non-Toxin	12
142	ORF1ab	6599	6610	SVKGLQPSVGPK	97.22% (35/36)	Non-Toxin	12
143	ORF1ab	6898	6909	LNDFVSDADSTL	97.22% (35/36)	Non-Toxin	12
144	ORF1ab	1354	1366	YILPSIISNEKQE	100.00% (36/36)	Non-Toxin	13
145	ORF1ab	1456	1468	GLNLEEAARYMRS	100.00% (36/36)	Non-Toxin	13
146	ORF1ab	1553	1565	FDNLKTLLSLREV	100.00% (36/36)	Non-Toxin	13
147	ORF1ab	1777	1789	EQFKKGVQIPCTC	100.00% (36/36)	Non-Toxin	13
148	ORF1ab	3310	3322	EDMLNPNYEDLLI	100.00% (36/36)	Non-Toxin	13
149	ORF1ab	5919	5931	RNVATLQAENVTG	97.22% (35/36)	Non-Toxin	13
150	ORF1ab	6932	6944	PKTKNVTKENDSK	97.22% (35/36)	Non-Toxin	13
151	ORF1ab	3211	3224	YNKYKYFSGAMDTT	100.00% (36/36)	Non-Toxin	14
152	ORF1ab	6002	6015	EEAIRHVRAWIGFD	97.22% (35/36)	Non-Toxin	14
153	ORF1ab	326	340	CGETSWQTGDFVKAT	100.00% (36/36)	Non-Toxin	15
154	ORF1ab	763	777	LQPLEQPTSEAVEAP	100.00% (36/36)	Non-Toxin	15
155	ORF1ab	886	900	VSELLTPLGIDLDEW	100.00% (36/36)	Non-Toxin	15
156	ORF1ab	2797	2811	SKHTDFSSEIIGYKA	100.00% (36/36)	Non-Toxin	15
157	ORF1ab	5730	5744	PAPRTLLTKGTLEPE	97.22% (35/36)	Non-Toxin	15
158	ORF1ab	6512	6526	KRNIKPVPEVKILNN	97.22% (35/36)	Non-Toxin	15
159	ORF1ab	6709	6723	RFKESPFELEDFIPM	97.22% (35/36)	Non-Toxin	15
160	ORF1ab	345	361	GTENLTKEGATTCGYLP	100.00% (36/36)	Non-Toxin	17
161	ORF1ab	2833	2849	DFDTWFSQRGGSYTNDK	100.00% (36/36)	Non-Toxin	17
162	ORF1ab	3356	3372	TANPKTPKYKFVRIQPG	100.00% (36/36)	Non-Toxin	17
163	ORF1ab	4546	4562	DDYFNKKDWYDFVENPD	97.22% (35/36)	Non-Toxin	17
164	ORF1ab	454	471	LQKEKVNINIVGDFKLNE	100.00% (36/36)	Non-Toxin	18
165	ORF1ab	2135	2152	WDTIANYAKPFLNKVVST	100.00% (36/36)	Non-Toxin	18
166	ORF1ab	1482	1500	AVTAYNGYLTSSSKTPEEH	100.00% (36/36)	Non-Toxin	19
167	ORF1ab	4380	4398	GCSCDQLREPMLQSADAQS	100.00% (36/36)	Non-Toxin	19
168	ORF1ab	6053	6071	PNNTDFSRVSAKPPPGDQF	97.22% (35/36)	Non-Toxin	19
169	ORF1ab	6817	6835	RMLLEKCDLQNYGDSATLP	97.22% (35/36)	Non-Toxin	19
170	ORF1ab	1398	1417	IVSTIQRKYKGIKIQEGVVD	100.00% (36/36)	Non-Toxin	20
171	ORF1ab	4157	4176	GTTQTACTDDNALAYYNTTK	100.00% (36/36)	Non-Toxin	20
172	ORF1ab	7035	7054	IQLSSYSLFDMSKFPLKLRG	97.22% (35/36)	Non-Toxin	20
173	ORF3a	135	140	SKNPLL	92.86% (26/28)	Non-Toxin	6
174	ORF3a	159	164	PYNSVT	100.00% (28/28)	Non-Toxin	6
175	ORF3a	216	225	STQLSTDTGV	100.00% (28/28)	Non-Toxin	10

176	ORF3a	61	71	KIITLKKRWQL	100.00% (28/28)	Non-Toxin	11
177	ORF3a	17	28	QGEIKDATPSDF	100.00% (28/28)	Non-Toxin	12
178	ORF6	44	57	LTENKYSQLDEEQP	100.00% (34/34)	Non-Toxin	14
179	ORF7a	17	25	LYHYQECVR	100.00% (28/28)	Non-Toxin	9
180	ORF7b	33	39	ELQDHNE	100.00% (2/2)	Non-Toxin	7
181	ORF8	91	95	QEPKL	100.00% (34/34)	Non-Toxin	5
182	ORF8	106	111	EDFLEY	100.00% (34/34)	Non-Toxin	6
183	ORF8	48	56	RVGARKSAP	100.00% (34/34)	Non-Toxin	9
184	ORF8	63	78	DEAGSKSPIQYIDIGN	100.00% (34/34)	Non-Toxin	16
185	S protein	293	296	LDPL	100.00% (41/41)	Non-Toxin	4
186	S protein	580	583	QTLE	100.00% (41/41)	Non-Toxin	4
187	S protein	1203	1206	LGKY	100.00% (41/41)	Non-Toxin	4
188	S protein	602	606	TNTSN	100.00% (41/41)	Non-Toxin	5
189	S protein	988	992	EAEVQ	100.00% (41/41)	Non-Toxin	5
190	S protein	773	779	EQDKNTQ	100.00% (41/41)	Non-Toxin	7
191	S protein	555	562	SNKKFLPF	100.00% (41/41)	Non-Toxin	8
192	S protein	807	814	PDPSKPSK	100.00% (41/41)	Non-Toxin	8
193	S protein	1035	1043	GQSKRVDFC	100.00% (41/41)	Non-Toxin	9
194	S protein	250	260	TPGDSSSGWTA	100.00% (41/41)	Non-Toxin	11
195	S protein	634	644	RVYSTGSNVFQ	100.00% (41/41)	Non-Toxin	11
196	S protein	656	666	VNNSYECDIPI	100.00% (41/41)	Non-Toxin	11
197	S protein	1107	1118	RNFYEPQIITTD	100.00% (41/41)	Non-Toxin	12
198	S protein	177	189	MDLEGKQGNFKNL	100.00% (41/41)	Non-Toxin	13
199	S protein	786	800	KQIYKTPPIKDFGGF	100.00% (41/41)	Non-Toxin	15
200	S protein	828	842	LADAGFIKQYGDCLG	100.00% (41/41)	Non-Toxin	15
201	S protein	206	221	KHTPINLVRDLPQGFS	100.00% (41/41)	Non-Toxin	16
202	S protein	695	710	YTMSLGAENSVAYSNN	100.00% (41/41)	Non-Toxin	16
203	S protein	138	154	DPFLGVYYHKNNKSWME	100.00% (41/41)	Non-Toxin	17
204	S protein	616	632	NCTEVPVAIHADQLTPT	100.00% (41/41)	Non-Toxin	17
205	S protein	304	322	KSFTVEKGIYQTSNFRVQP	100.00% (41/41)	Non-Toxin	19
206	S protein	672	690	ASYQTQTNSPRRARSVASQ	100.00% (41/41)	Non-Toxin	19

Eleven SARS-CoV-2 proteins were chosen
Protein sequences retrieved from NCBI

Screening of potential CTL, HTL and B Cell epitopes

Screening of CTL epitope
(IEDB "MHC-I Binding Predictions" and "MHC-I Processing Predictions")

Screening of HTL epitopes
(IEDB "MHC-II Binding Predictions")

Screening of protein sequence based B Cell epitope
(IEDB "B Cell Epitope Prediction Tools")

Population coverage by screened
CTL and HTL epitopes *(IEDB "Population Coverage")*

Characterization of screened potential epitopes

Epitope conservation analysis
(IEDB "Epitope Conservancy Analysis")

Epitope toxicity prediction
(ToxinPred)

Overlapping residue analysis for
CTL, HTL & B Cell epitopes *(Clustal Omega)*

Molecular interaction study of selected
overlapping CTL & HTL epitopes with HLA allele and TAP transporter

3D structure modelling of HLA alleles
and selected CTL & HTL epitopes *(Swiss-model & PEP-FOLD2.0)*

Molecular interaction analysis of selected
overlapping Ctl and HTL epitopes with HLA allele
(PatchDock Tool)

Molecular interaction analysis of selected
CTL epitopes with TAP transporter
(PatchDock Tool)

Design and characterisation of
CTL & HTL Multi-Epitope Vaccines

Interferon gama inducing epitope
prediction from designed MEVs *("IFNepitope")*

MEVs allergenicity and
antigenicity prediction *(AlgPred & VaxiJen)*

Physicochemical
analysis of MEVs *(ProtParam)*

3D structure modelling and refinement of CTL &
HTL MEVs *(I-TASSER, ModRefiner, GalaxyWEB)*

In-silico **validation of refined CTL**
& HTL MEV models *(RAMPAGE)*

Discontinuous B-cell epitope prediction
from CTL & HTL MEVs *(IEDB "ElliPro")*

Molecular interaction analysis of
CTL & HTL MEVs with TLR3
(PatchDock)

***In-silico* analysis of cDNA of CTL & HTL MEVs for**
cloning and expression in mammalian cell line (Human)
(Codon Usage Wrangler Tool & GenScript Rare Codon Analysis Tool)

Ov-ASP-1
(truncated (10-153))

IVVAVTGYNCPGGKLTALERKKIVGQNNKYRSDLINGKL
KNRNGTYMPRGKNMLELTWDCKLESSAQRWANQCIFG
HSPRQQREGVGENVYAYWSSVSVEGLKKTAGTDAGKS
WWSKLPKLYENNPSNNMTWKVAGQGVLHFTQ

B. HTL epitopes	A. CTL epitopes
EAAAK	EAAAK
WFLAYILFTRFFYVL	SEMVMCGGSLY
GGGGS	GGGGS
FLAYILFTRFFYVLG	FYWFFSNYLKR
GGGGS	GGGGS
LAYILFTRFFYVLGL	ISNSWLMW
GGGGS	GGGGS
AYILFTRFFYVLGLA	ETISLAGSYK
GGGGS	GGGGS
RAMPNMLRIMASLVL	QEILGTVSW
GGGGS	GGGGS
AMPNMLRIMASLVLA	STFNVPMEK
GGGGS	GGGGS
MPNMLRIMASLVLAR	RMYIFFASFY
GGGGS	GGGGS
PNMLRIMASLVLARK	FLFVAAIFYL
GGGGS	GGGGS
QESPFVMMSAPPAQY	RYFRLTLGVY
GGGGS	GGGGS
NMLRIMASLVLARKH	FLNGSCGSV
GGGGS	GGGGS
ESPFVMMSAPPAQYE	CTDDNALAY
GGGGS	GGGGS
SPFVMMSAPPAQYEL	CTDDNALAYY
GGGGS	GGGGS
AIILASFSASTSAFV	MYKGLPWNVVR
GGGGS	GGGGS
IILASFSASTSAFVE	SIINNTVYTK
GGGGS	GGGGS
FAWWTAFVTNVNASS	LPVNVAFELW
GGGGS	GGGGS
KTQSLLIVNNATNVV	DEWSMATYY
GGGGS	GGGGS
TQSLLIVNNATNVVI	YILFTRFFYV
GGGGS	GGGGS
QSLLIVNNATNVVIK	YIFFASFYYV
GGGGS	GGGGS
SLLIVNNATNVVIKV	WTAGAAAYYV
GGGGS	GGGGS
LLIVNNATNVVIKVC	FPNITNLCPF
GGGGS	GGGGS
LLFVTVYSHLLLVAA	NYNYLYRLFR
GGGGS	GGGGS
FVRATATIPIQASLP	NYLYRLFR
GGGGS	GGGGS
VLLFLAFVVFLLVTL	YLYALVYFL
GGGGS	GGGGS
LLFLAFVVFLLVTLA	IPYNSVTSSI
GGGGS	GGGGS
GLMWLSYFIASFRLF	LLFLAFVVF
GGGGS	GGGGS
LMWLSYFIASFRLFA	LTALRLCAY
GGGGS	GGGGS
LSYYKLGASQRVAGD	YFIASFRLFAR
GGGGS	GGGGS
TFKVSIWNLDYIINL	ATSRTLSYYK
GGGGS	GGGGS
FKVSIWNLDYIINLI	RTFKVSIW
GGGGS	GGGGS
KVSIWNLDYIINLII	AEILLIMRTF
GGGGS	GGGGS
IILFLALITLATCEL	RARSVSPK
GGGGS	GGGGS
ILFLALITLATCELY	QLRARSVSPK
GGGGS	GGGGS
YLCFLAFLLFLVLIM	FLAFLLFLV
GGGGS	GGGGS
LCFLAFLLFLVLIML	HFYSKWYIR
GGGGS	GGGGS
CFLAFLLFLVLIMLI	MEVTPSGTW
GGGGS	GGGGS
CTQHQPYVVDDPCPI	KPRQKRTAT
GGGGS	GGGGS
HQPYVVDDPCPIHFY	MGYINVFAF
GGGGS	GGGGS
QPYVVDDPCPIHFYS	GYINVFAFPF
GGGGS	
PQIAQFAPSASAFFG	
GGGGS	
IAQFAPSASAFFGMS	
GGGGS	
AQFAPSASAFFGMSR	
GGGGS	
YINVFAFPFTIYSLL	
HHHHHH	HHHHHH

Protein labels (left margin):

ORF1ab
Surface protein
ORF3a
Envelope protein
Membrane protein
ORF6
ORF7a
ORF7b
ORF8
Nucleocapsid phosphoprotein
ORF10

Overlapping T & B cell epitopes across proteome of SARS-CoV-2

1. ATSRTLSYYK-HLA-A*11:01

2. HFYSKWYIR-HLA-A*31:01

CTL epitopes - HLA Class I allele complexes

3. CTQHQPYVVDDPCPI-HLA-DRB3*01:01

4. HQPYVVDDPCPIHFY-HLA-DRB3*01:01

5. LSYYKLGASQRVAGD-HLA-DRB1*09:01

6. QPYVVDDPCPIHFYS-HLA-DRB3*01:01

HTL epitopes - HLA Class II allele complexes

1. ATSRTLSYYK-HLA-A*11:01

2. HFYSKWYIR-HLA-A*31:01

CTL epitopes - HLA Class I allele complexes

3. CTQHQPYVVDDPCPI-HLA-DRB3*01:01

4. HQPYVVDDPCPIHFY-HLA-DRB3*01:01

5. LSYYKLGASQRVAGD-HLA-DRB1*09:01

6. QPYVVDDPCPIHFYS-HLA-DRB3*01:01

HTL epitopes - HLA Class II allele complexes

Epitope ATSRTLSYYK through TAP cavity   **Epitope HFYSKWYIR through TAP cavity**

Number of residues in favoured region (~98.0% expected)   : 602 (85.8%)
Number of residues in allowed region (~2.0% expected)   : 79 (11.3%)
Number of residues in outlier region   : 21 (3.0%)

A. CTL Multi-epitope vaccine RAMPAGE analysis

Number of residues in favoured region (~98.0% expected)   : 718 (88.9%)
Number of residues in allowed region (~2.0% expected)   : 72 (8.9%)
Number of residues in outlier region   : 18 (2.2%)

B. HTL Multi-epitope vaccine model RAMPAGE analysis

CTL Multi-epitope vaccine model in complex with TLR3

HTL Multi-epitope vaccine model in complex with TLR3

CTL Multi-epitope COVID19 vaccine - Toll Like Receptor 3 complex   **CTL Multi-epitope COVID19 vaccine - Toll Like Receptor 3 complex**

Graphical abstract: The designed CTL (Cytotoxic T lymphocyte) and HTL (Helper T lymphocyte) multi-epitope vaccine (MEV) against COVID19 infection. Both the CTL and HTL MEVs models show a very stable and very fitting conformational complex formation tendency with the Toll like receptor 3. CTL and HTL MEVs: ribbon; Toll like receptor 3: gray cartoon; Adjuvant [truncated (residues 10-153) Onchocerca volvulus activation-associated secreted protein-1]: orange ribbon; Epitopes: cyan ribbons regions; 6xHis Tag: magenta ribbon region

A. CTL Multi-epitope vaccine model

B. HTL Multi-epitope vaccine model