Title: A microbial perspective on balancing trade-offs in ecosystem functions in a constructed

stormwater wetland

Author: Regina B. Bledsoe[a]*, Eban Z. Bean[b], Samuel S. Austin[a], Ariane L. Peralta[a]

*Corresponding author

Email: ginabbledsoe@gmail.com, ezbean@ufl.edu, austins15@ecualumni.ecu.edu,

peraltaa@ecu.edu

Affiliations:

[a]East Carolina University, Department of Biology, Greenville, NC, 27858, USA

[b]University of Florida, Department of Agricultural and Biological Engineering, Gainesville, FL,

32611, USA

Abbreviations: corrected Akaike information criteria = AICc, NH_4^+ = ammonium, C = carbon,

CO_2 = carbon dioxide, CW = constructed wetland, N_2 = dinitrogen, GWP = global warming

potential, GHG = greenhouse gas, CH_4 = methane, N= nitrogen, NO_3^- = nitrate, NO = nitric

oxide, NO_2^- = nitrite, N_2O = nitrous oxide, OTU = operational taxonomic unit, PCR =

polymerase chain reaction, PERMANOVA = permutated analysis of variance, PCoA = principal

coordinates analysis, SCM = stormwater control measure

Abstract

Green stormwater infrastructure, such as constructed wetlands (CWs), is a type of stormwater control measure that can decrease nutrient and pollutant loads from urban stormwater runoff. Wetland soil microorganisms provide nutrient and pollutant removal benefits which can also result in ecosystem disservices such as greenhouse gas (GHG) emissions and can inadvertently exacerbate climate change. Microbial respiration by facultative anaerobes in anoxic conditions is the primary pathway for nitrogen removal (benefit). Similar anoxic conditions that support denitrifying microorganisms can also support obligate anaerobes that produce methane (CH_4) via methanogenesis (disservice). We examined nitrogen removal potential, GHG production, and microbial community structure within permanently flooded and shallow land or temporarily-flooded areas of a stormwater CW to identify zones for CW design optimization. Results indicate that permanently flooded zones compared to shallow land zones are greater sources of CH_4 emissions (80.80 ± 118.31, 2.32 ± 9.33 mg CH_4-C m^{-2} hr^{-1}, respectively) and emit more carbon to the atmosphere (7161.27 kg CO_2, 93.20 kg CO_2 equivalents, respectively). However, nitrogen removal potential rates were similar across both flooded and shallow land zones (24.45 ± 20.18, 20.29 ± 15.14 ng N_2O-N hr^{-1} g^{-1} dry soil, respectively). At this particular CW, reduction of permanently flooded zones within the wetland could decrease GHG emissions (disservice) without limiting nitrogen removal (benefit) potential of the wetland. Holistic development and design of stormwater control measures, which account for microbial activity, provides the opportunity to maximize benefits (i.e., nutrient and pollutant removal) and reduce disservices (i.e., GHG emissions) of green stormwater infrastructure.

Keywords: Greenhouse gases, Denitrification, Methane, Stormwater, Urban ecosystems

Introduction

In urban areas, impervious surfaces increase nutrient and pollutant loads in stormwater

runoff, thereby decreasing water quality within neighboring watersheds (Bell et al., 2019; Line

and White, 2007; O'Driscoll et al., 2010). Nutrient loadings are of particular concern due to

adverse effects (e.g., harmful algal blooms and hypoxic zones) on downstream water quality

(Paerl et al., 2014; Rabalais et al., 2009). Green stormwater infrastructure, such as constructed

wetlands (CWs), are stormwater control measures (SCMs) that are designed to mimic natural

ecosystems to provide relief from stormwater runoff and improve water quality in and around

urban areas. The physical, chemical, and biological treatment processes that naturally occur in

wetlands make CWs an effective approach to reduce nutrient loads in stormwater runoff (Koch et

al., 2014; Mitsch et al., 2013; Payne et al., 2014; Reisinger et al., 2016). Thus, entities such as

the North Carolina Department of Environmental Quality encourage use of SCMs as a way to

reduce nutrient loadings to waterways by offering nutrient reduction credits. CWs that have a

ponding depth of less than 12 inches and a hydraulic residence time of two to five days are

credited with 40% removal of nitrogen (N) from stormwater runoff (NCDEQ, 2018). However,

this nutrient reduction policy and similar policies outside of North Carolina do not consider the

potential trade-off between water quality (e.g., N removal) and air quality (e.g., greenhouse gas

(GHG) emissions). The unintended GHG production from stormwater CWs could negatively

affect climate change mitigation and adaptation by reducing the carbon storage capacity typically

provided by wetlands (Demuzere et al., 2014). Therefore, in order to meet management goals in

a more holistic way (i.e., enhancing total N removal and reducing GHG emissions), a deeper

understanding of microbial processes occurring within CWs are needed to optimize wetland

design.

Plants and soil microorganisms carry out nutrient and pollutant removal processes that occur

within wetlands. While plants and microorganisms can assimilate ammonium (NH_4^+) and nitrate

(NO_3^-) ions into biomass, the primary pathway of N removal from a wetland ecosystem is

microbial respiration via denitrification and to a lesser degree anaerobic ammonium oxidation

(Lee et al., 2009). Denitrification is typically coupled with nitrification in which NH_4^+ is first

transformed to nitrite (NO_2^-) and then NO_3^- by aerobic nitrifying microorganisms (Lee et al.,

2009). The process of denitrification occurs when NO_3^- is reduced to nitric oxide (NO), then to

nitrous oxide (N_2O), a potent GHG, and finally to dinitrogen (N_2), an inert and abundant

atmospheric gas (Knowles, 1982; Lee et al., 2009; Smith and Tiedje, 1979). Denitrification is an

anaerobic microbial process that requires anoxic or low-oxygen conditions such as those found

in saturated and flooded soils within wetlands (Marton et al., 2015; Smith and Tiedje, 1979).

Many denitrifying microorganisms are facultatively anaerobic, meaning these microorganisms

can perform aerobic respiration when oxygen is available but can use other alternate electron

acceptors (i.e., nitrate) when anoxic soil conditions arise (Tiedje, 1988). Due to this facultative

anaerobic metabolism, these microorganisms respond to hydrologic change with varying degrees

depending on past environmental conditions (Peralta et al., 2014, 2013).

Similar anoxic conditions that support nitrogen removal by denitrification (transformation of

NO_3^- to N_2) can also support production of methane (CH_4), a GHG, which is considered an

ecosystem disservice (Demuzere et al., 2014). Methanogens are obligate anaerobes that produce

CH_4 via methanogenesis but only under anoxic conditions when the environment is depleted of

other electron acceptors (e.g., nitrate, sulfate, and ferric iron) (Fetzer and Conrad, 1993; Liu et

al., 2008). When organic matter is decomposed by fermentation, methanogens can use the

byproducts of acetate, hydrogen, and CO_2 for energy and produce CH_4 (Conrad, 2007). CH_4

production can also support the consumption of CH_4 by aerobic methanotrophs, which have been

reported to consume 45-90% of CH_4 produced (Brindha and Vasudevan, 2018; Conrad, 2007; Le

Mer and Roger, 2001). Systems in which the population of methanogens (i.e., producers of CH_4)

is greater than methanotrophs (i.e., consumers of CH_4) can lead an ecosystem to be a CH_4 source

(Altshuler et al., 2019; Rey-Sanchez et al., 2019; Wen et al., 2018). Since methanogens are

obligate anaerobes, even low amounts of oxygen can suppress CH_4 production (Fetzer and

Conrad, 1993; Le Mer and Roger, 2001). However, methanotrophs are aerobes, and there is

typically greater CH_4 oxidation in dynamic, wet-dry conditions that are not permanently

inundated (Chowdhury and Dick, 2013). Therefore, anoxic conditions that can support

denitrification can also result in GHG emissions, especially when nitrate is limiting (Burgin et

al., 2011).

The three major biogenic GHGs of concern, CO_2, CH_4, and N_2O, are known to contribute

to increased atmospheric warming and thereby influence global climate change. While CO_2

(416.21 ppm) is the most abundant GHG in the atmosphere, CH_4 (1.87 ppm) and N_2O (0.33

ppm) are found in lower atmospheric concentrations but have a greater global warming potential

(GWP) per molecule than CO_2 (IPCC, 2014) (https://www.esrl.noaa.gov/gmd/ccgg/trends/).

Specifically, CH_4 and N_2O have GWP of 28 and 265, respectively, over a 100 year time-scale

when compared to CO_2 which is equal to 1 (IPCC, 2014).Therefore, smaller amounts of CH_4 or

N_2O can contribute more to increasing GWP than equal amounts of CO_2.

To improve water quality management of CWs in the urban landscape, both ecosystem

services (e.g., nitrogen removal by denitrification) and disservices (e.g., GHG production) should

be considered during the planning and design phase (Demuzere et al., 2014). To accommodate

stormwater runoff and water quality functions, CWs generally have four main zones: upland

(rarely submerged except during large runoff events), shallow land (soil is saturated but only

temporarily inundated following runoff events), shallow pools (designed to be permanently

inundated except during droughts) and deep pools (permanently inundated) (NCDEQ, 2019). In

CWs, ammonification and nitrification occur in the shallow land zones as stormwater percolates

into aerated soils. If soils in shallow lands are saturated for sufficient durations, denitrification is

expected to eventually occur as well. It is in the shallow water zones where denitrification is

conventionally thought to occur due to saturated, anoxic conditions. In this study we focus on

denitrification as the ecosystem service since this process typically accounts for the most N

removal in CWs (Rahman et al., 2019).

The goal of this study is to identify areas within a stormwater CW that support

denitrification and reduce GHG

emissions, especially due to methanogenesis. We hypothesize that the dynamic wet-dry

conditions occurring in shallow land and flooded zones will both support the anaerobic process

of denitrification, while only the flooded zones will be sources of CH_4 gas. If production of CH_4

is greater than CO_2 production, the overall GWP of the wetland will increase since CH_4 has a

higher GWP than CO_2. However, due to the facultative nature of microbes residing in the

shallow land zones, denitrification will still occur in shallow land zones after a short period of

flooding if NO_3^- is not limiting. Further, we expect microbial community composition to differ

between land types such that flooded zones will be represented by a greater relative abundance

of microbes involved in CH_4 cycling while shallow land zones will have a greater proportion of

microbes involved in nitrogen cycling.

2. Materials and methods

To test these hypotheses, we monitored GHG fluxes monthly for one year and sampled

sediments and surface water for chemical analyses seasonally. To determine microbial (bacterial

and archaeal) community structure in sediments, we carried out 16s rRNA V4 amplicon

sequencing. We also determined potential denitrification rates across the wetland using a short-

term incubation experiment. By combining GHG rates with potential denitrification rates,

sediment and water chemistry parameters along with microbial community composition, we

examined the potential benefits and disservices occurring within this CW.

2.1 *Study site*

The study location is a recently constructed (2015) stormwater wetland located on East Carolina

University's main campus in Greenville, NC (N 35°35'22.4", W 77°22'13.4"). Prior to

construction, the area had served as a dry detention basin since at least 1998, based on historical

aerial imagery analysis. The 0.13 acre CW receives runoff from 5.7 acre parking lot in a highly

urbanized area of the city. To create the additional storage volume for runoff and permanent

ponding, the basin surface was excavated to create landform features. Soils were characterized

by orange and gray mottled loamy clay, indicative of reducing, low-oxygen soil conditions.

Infiltration rates were measured to be < 0.1 in/h, making for a suitable substrate to maintain the

permanent water levels in the CW. The temporary ponding depth (30.5 cm) of the wetland is

equal to 0.51 cm of runoff within the impervious drainage area (Fig. 1), undersized relative to

permitting requirements at the time (NCDEQ, 2019). The total surface area is divided among

main zones (% surface area): shallow land (35%), shallow water (35%), deep pools (10%),

forebay (10%), and outlet pool (10%)

(NCDEQ, 2019).The dominant plants

within saturated areas of the wetland are

Typha spp., *Pontederia cordata*, and *Iris*

versicolor L. The shallow land area along

the water's edge is dominated by *Carex*

lupulina Muhl. ex Willd. 2.2 *GHG*

monitoring and flux calculations

We used a static chamber method to

measure soil respiration as GHG fluxes

(Parkin et al., 2010). At the CW, we

established 13 GHG sampling plots along

two main transects that follow the central

channel within the wetland that flows from

inlet to outlet (Fig. 1). Of those 13 plots,

chambers 1-5 were placed in shallow water

and deep pools (flooded zones), and

chambers (6-13) were placed in shallow

land zones along the outside edge of the

main channel. Each sampling plot in

shallow land zones (n = 10) was fitted with

a GHG chamber bottom installed 6 cm

belowground. During sampling events,

Figure 1. Top: Landscape photograph of wetland. Center: Google Earth image of parking lot and stormwater CW outlined in red. Bottom: Google Earth image of wetland with blue circles (1-5) representing flooded plots and white circles (6-13) representing shallow land plots sampled for GHG and sediments.

chamber tops (30 cm height, 15 cm diameter) were attached to an aboveground groove, and the

groove was filled with water to create a gas tight seal. GHG chamber tops in flooded zones were

fitted with a Styrofoam float allowing 2-3 cm of the chamber top below the water surface.

We collected GHG samples monthly for one year. Sampling events occurred at least 48

hours after a rain event of > 0.51 cm. Gas samples were collected at 0, 12, 24, and 36 minutes

using 20 mL syringes. Each gas sample was divided between two 3.7 mL Exetainers® (Labco,

Lampeter, Wales, UK) with dual septa and stored upside at room temperature until analyzed. We

measure GHG concentrations using a Shimadzu 2014 Gas Chromatograph (Shimadzu Scientific

Instruments, Columbia, Maryland) fitted with an electron capture detector to detect N_2O and

flame ionization detector to measure CH_4 and CO_2. Monthly GHG fluxes were calculated using

the linear change in concentration and the ideal gas law (Millar et al. 2018). Seasonal estimates

were calculated by averaging monthly flux measurements: spring = March-May, summer = June-

August, fall = September-November and winter = December-February. In addition, the

equivalent CO_2 flux for CH_4 was calculated by multiplying CH_4 flux by 28, which is equal to

the increase in warming potential in one molecule of CH_4 versus one molecule of CO_2.

2.4 *Soil collection and chemical properties*

We characterized soil physicochemical properties seasonally (May, July, October, and

December). Each composite soil sample represented three cores (3.1 cm diameter, 12 cm depth)

that were adjacent to each GHG sampling chamber. We homogenized each composite sample by

passing the sample through a 4 mm sieve, and then processed subsamples for soil gravimetric

moisture, pH, extractable ammonium and nitrate, total carbon, and total nitrogen. We combined

5 g of field moist soil and 45 mL of 2M potassium chloride and shook samples for 1 hour before

gravity filtration to collect soil extracts. NH_4^+ and NO_3^- concentrations were colorimetrically

measured using a SmartChem 200 auto analyzer (Unity Scientific Milford, Massachusetts, USA)

at the East Carolina University Environmental Research Laboratory. Using a 20-30 g subsample,

we measured field-moist and oven-dry (dried 24 hours at 105°C) soil weights to determine

moisture content by dividing the mass of water by the mass of oven-dried soil. Using a mortar

and pestle, we coarsely ground a subsample of soil and combined 10 g of soil with 10 mL of

Nanopure® water to measure soil pH. On another subsample, we finely ground the soil for

elemental carbon and nitrogen analysis using an elemental analyzer (2400 CHNS Analyzer;

Perkin Elmer; Waltham, Massachusetts, USA) at the Environmental and Agricultural Testing

Service laboratory (Department of Crop and Soil Sciences at NC State). Finally, we measured

organic matter content based on the loss on ignition method by ashing 20 g coarsely ground

oven-dried soil at 550 °C (Hoogsteen et al., 2015) for 2 hours since the majority of OM is ashed

within the first 2 hours of combustion (Heiri et al., 2001). However, up to 4 hours may be

required to completely ash all organic matter from soils with high OM content (Heiri et al.,

2001). In addition, a subsample of field moist soil was stored at 4 °C until we performed

denitrification potential assays. Finally, a subsample of soil was stored at -80 °C until DNA

extraction for bacterial amplicon sequencing.

2.3 *Seasonal denitrification potential*

Within 72 hours of soil collection, we measured potential denitrification rates using the

acetylene block denitrification enzyme assay method (Schaller et al., 2004; Tiedje et al., 1989;

Wall et al., 2005). We weighed triplicate soil subsamples of 25 g and transferred each subsample

into separate 125 mL Wheaton bottles fitted with a phenolic cap and butyl septa. In each bottle,

we added 75 mL of 1 mM potassium chloride (KNO_3) to provide an abundant nitrogen supply

and 1.3 mL of chloramphenicol (100 mg/mL) to inhibit new denitrification enzyme production.

Bottles were sealed with septa-centered caps, shook, and purged with helium for 5 minutes to

create an anoxic environment. We removed 15 mL of helium from the headspace and replaced

with 15 mL of pure acetylene gas to prevent the conversion of N_2O to N_2. At each hour for 3

hours, we collected a 10 mL gas sample (time point T0, T1, T2, T3) from each bottle and stored

the gas sample in a 3.7 mL Exetainer®. We shook each bottle to equilibrate N_2O in aqueous and

sediment phases before each sample collection. We added 10 mL of 10% acetylene mixture to

the headspace after each sample collection to maintain acetylene concentrations within the bottle.

N_2O fluxes were calculated as described for GHG fluxes (Millar et al., 2018).

2.5 *Microbial Community Analysis*

We examined the microbial community from soils collected in May (spring) and October

(fall) from flooded plots (1-5) and shallow land plots (7-9, 11-13) (Fig. 1). First, we extracted

genomic DNA from soils using the Qiagen DNeasy Powerlyzer PowerSoil Kit. Genomic DNA

was amplified using barcoded primers 515FB/806R primer set, originally developed by the Earth

Microbiome Project (earthmicrobiome.org) to target the V4-V5 region of the bacterial 16S

subunit of the ribosomal RNA gene (Apprill et al., 2015; Caporaso et al., 2010; Parada et al.,

2016). Samples were prepared for sequencing by using polymerase chain reaction (PCR)

chemistry to amplify the 16S rRNA DNA with a unique sample identifier to each sequence

within each individual sample. For each sample, three 50 µL PCR libraries were prepared by

combining 38.35 µL molecular grade water, 5 µL Amplitaq Gold 360 10x buffer, 2.4 µL MgCl$_2$

(25 mM), 1 µL dNTPs (40mM total, 10mM each), 0.25 µL Amplitaq Gold 360 polymerase, 1 µL

515 forward barcoded primer (10 µM), 1 µL 806 reverse primer (10 µM), and 1 µL DNA

template (10 ng µL^{-1}) following the manufacturer's protocol (Applied Biosystems, Foster City,

CA, USA . Thermocycler conditions for PCR reactions were as follows: initial denaturation (94

°C, 3 minutes); 30 cycles of 94°C for 45 seconds, 50 °C for 30 seconds, 72 °C for 90 seconds;

final elongation (72 °C, 10 minutes). The triplicate 50 µL PCR libraries were combined and then

amplified DNA sequences separated from PCR reagents using the AMPure XP magnetic bead

protocol (Axygen, Union City, California, USA). Cleaned PCR products were quantified using

QuantIT dsDNA broad range assay (Thermo Scientific, Waltham, Massachusetts, USA) and

diluted to a concentration of 10 ng µL^{-1}. We combined barcoded PCR samples (5 ng µL^{-1}) in

equimolar concentration and sequenced the pooled library using the Illumina MiSeq platform

paired end read approach (Illumina Reagent Kit v2, 500 reaction kit) at the Indiana University

Center for Genomics and Bioinformatics Sequencing Facility.

Sequences were processed using a standard mothur pipeline (v1.40.1)(Kozich et al.,

2013; Schloss et al., 2009). We assembled contigs from the paired end reads, quality trimmed

using a moving average quality score (minimum quality score 35), aligned sequences to the Silva

Database (Quast et al. 2013; SSURef v123), and removed chimeric sequences using the

VSEARCH algorithm (Rognes et al., 2016). We created operational taxonomic units (OTUs) by

first splitting sequences based on taxonomic class and then binning into OTUs based on 97%

sequence similarity. The Silva database was used to assign taxonomy to each OTU.

2.6 *Statistical Analysis*

All statistical analyses were completed in the R statistical environment (RStudio

v1.1.383, Rv3.4.0) (R Core Team, 2019). We constructed linear mixed effects models with

sampling plot as a random effect using the *lme4* R package (Bates et al., 2015) to determine the

importance of hydrology, season, and distance from the inlet, and the interactions on GHG fluxes

and denitrification potential fluxes within the constructed wetland. We then conducted model

comparisons using sample size corrected Akaike information criteria (AICc) model comparisons,

which adjust for small sample size, against the null model to determine which fixed effects

(hydrology,season, and distance from inlet) to determine which combinations had the most

explanatory power (Gorsky et al., 2019; Hurvich and Tsai, 1993). Estimated marginal means

with Tukey post-hoc adjustment in the R package *emmeans* (Lenth, 2017) was used to determine

significance ($p<0.05$) in gas flux rates between treatment groups and the *multcomp* R package

(Hothorn et al., 2020) was used to assign Tukey post-hoc groups. Lastly, we used the *MuMIn* R

package to determine the proportion of variance explained by fixed effects (marginal) and the

complete model (conditional) (Barton, 2019; Gorsky et al., 2019). The CH_4, CO_2, and N_2O

fluxes were transformed to the log of the cube root and denitrification potential rates were log

transformed to meet normality assumptions of the statistical models.

Finally, we examined microbial community diversity and how that relates to soil

chemistry and GHG fluxes. We standardized read counts before calculating diversity metrics for

sediment microbial communities by removing singletons and doubletons, rarifying all sample to

the sample with the lowest read count, and resampling. Principal Coordinates of Analysis

(PCoA) based on Bray-Curtis dissimilarity was used to visualize patterns of bacterial diversity

among treatments and sampling dates. Permuted analysis of variance (PERMANOVA) was run

using the *adonis* function in the *vegan* package (Oksanen, 2015) to determine the importance of

hydrology and season in shaping overall microbial community composition. We performed a

Dufrene-Legendre indicator species analysis using the *labdsv* R package (Roberts, 2016) to

identify specific microbial community members that represented each hydrologic treatment.

Next, we examined the relationship between individual GHG fluxes and microbial community

Bray-Curtis dissimilarity matrix using distance based partial least square regression in the *dbstats*

R package (Boj et al., 2017). Finally, we used Mantel R statistic function in the *vegan* R package

(Oksanen, 2015) to examine the relationship between microbial community composition and all

soil properties (moisture, pH, organic matter, NH_4^+ and NO_3^- concentrations, total C, total N, and

C:N).

3. Results

3.1 Sediments and water properties

Over the one year study period in 2017, there were 268 days without precipitation.

Within flooded plots, water depth ranged from 8-28 cm in shallow water areas and 25-53 cm in

deep pools. Temperatures varied seasonally with July having the highest average ($\pm$ SD)

temperature (26.5 $\pm$ 1.8°C) and December having the lowest average temperature (4.9 $\pm$ 1.3°C)

(Table 1). Sediments in the wetland were acidic (4.6 $\pm$ 0.3 pH) ranging from pH 4.08 to 5.28.

Extractable NH_4^+ and NO_3^- concentrations were generally low in sediments (0.089 $\pm$ 0.066 mg

NH_4^+-N g^{-1} dry mass and 0.023 $\pm$ 0.008 mg NO_3^--N g^{-1} dry mass). In the water column, NH_4^+

concentrations were higher (2.57 $\pm$ 19.6 mg NH_4^+ L^{-1}) than NO_3^- concentrations (0.328 $\pm$ 0.803

mg NO_3^--N L^{-1}) throughout the wetland, especially during winter months (9.97 $\pm$ 39.4 mg NH_4^+

L^{-1} and 0.769 $\pm$ 1.27 mg NO_3^- L^{-1}). Sediment C:N ratios along with other sediment parameters

(total C, total N, and C:N) and water column phosphate (PO_4^{3-}) concentrations were similar

across space (hydrologic zone) and time (season) within the wetland (Table 1A&B).

3.2 GHG flux rates

3.2.1 Methane (CH_4)

Across the wetland, CH_4 fluxes differed by hydrology and season (Fig. 2A). Average ($\pm$

SD) CH_4 fluxes across all seasons were lowest in shallow land plots (2.32 $\pm$ 9.33 mg CH_4-C m^{-2}

hr^{-1}) and highest in flooded plots (80.80 $\pm$ 118.31 mg CH_4-C m^{-2} hr^{-1}). Within flooded plots, the

highest mean CH_4 fluxes (160.24 $\pm$ 156.28 mg CH_4-C m^{-2} hr^{-1}) were detected during summer

months, and the lowest CH_4 fluxes were detected in winter and spring months (45.02 $\pm$ 88.74 and

46.98 $\pm$ 59.40 mg CH_4-C m^{-2} hr^{-1}, respectively). Within shallow land plots, the highest mean CH_4

fluxes (6.38 $\pm$ 16.44 mg CH_4-C m^{-2} hr^{-1}) were detected during spring months and the lowest CH_4

fluxes were detected in winter (0.09 $\pm$ 0.42 mg CH_4-C m^{-2} hr^{-1}). The model that included

hydrology and seasonal interaction explained the most variation in CH_4 fluxes based on a ΔAICc

score of 0 and an AICc weight of 1 (Table S1.1). In this model the interaction of hydrology and

season and random effects accounted for 73% variation in CH_4 fluxes, while the interaction

alone explained 63% of that variation (Table S1.1).

3.2.2 Carbon dioxide (CO_2)

The wetland CO_2 fluxes varied seasonally but were generally highest in flooded plots

compared to shallow land plots (Fig. 2B). Across all seasons CO_2 fluxes were lowest in shallow

land plots (3.95 ± 43.96 mg CO_2-C m^{-2} hr^{-1}) and highest in flooded plots (73.52 ± 100.24 mg

CO_2-C m^{-2} hr^{-1}). We measured similar CO_2 flux rates (20.82 ± 46.36, 14.37 ± 44.68, and $18.88 \pm$

24.23 mg CO_2-C m^{-2} hr^{-1}) during spring, fall, and winter seasons, while CO_2 fluxes ($63.13 \pm$

133.21 mg CO_2-C m^{-2} hr^{-1}) were highest during the summer. The model with season only

explained the most variation in CO_2 fluxes based on a ΔAICc score of 0 and AICc weight of 0.93

(Table S1.1). The fixed effect of season explained 26% of the variation and the full model

explained 50% of the variation in CO_2 fluxes (Table S1.1).

Table 1A & B. Summary of mean (range) of soil and water properties measured at the constructed wetland. Annual wetland values are

averages of all samples for the entire sampling period. (1A) Hydrology (flooded vs. shallow land) values are averages of samples

collected in May (spring), July (summer), October (fall), and December (winter) for each hydrology type. (1B) Seasonal sediment

values are averages from of all plots based on samples collected in May (spring), July (summer), October (fall), and December

(winter). Seasonal water values are averages of monthly samples from flooded plots where spring represents months May-June,

summer represents months July-September, fall represents months October-November, and winter represents months December-

February. (Abbreviations: DM = dry mass. MDL = below detection limit).

A - Hydrology	Whole Wetland		Flooded		Shallow land	
	Mean	Range	Mean	Range	Mean	Range
Soil						
temperature (°C)	18.6	3.0-30.0				
moisture (%)	33.3	18.3-51.9	36.2	23.2-51.9	31.5	18.3-46.2
pH	4.6	4.1-5.3	4.7	4.2-5.3	4.6	4.1-5.2
organic matter (%)	6.20	3.40-10.17	6.54	3.41-10.17	6.01	3.56-8.42
mg NH_4^+-N g^{-1} DM	0.089	0.023-0.457	0.129	0.049-0.457	0.066	0.023-0.148
mg NO_3^--N g^{-1} DM	0.023	0.012-0.050	0.027	0.14-0.050	0.021	0.012-0.038
Total C (%)	1.66	0.27-3.61	1.78	0.27-3.61	1.60	0.57-2.81
Total N (%)	0.10	0.02-0.20	0.11	0.02-0.20	0.10	0.04-0.16
C:N (wt: wt)	15.78	11.00-19.00	15.88	11.00-19.00	15.73	13.88-18.44

B - Seasonal	Spring		Summer		Fall		Winter	
	Mean	Range	Mean	Range	Mean	Range	Mean	Range
Soil								
temperature (°C)	19.6	16.8-23.0	26.5	25.0-30.0	21.3	20.0-24.0	4.9	3.0-7.0
moisture (%)	32.1	26.6-42.9	35.3	27.0-48.1	35.6	26.6-51.9	29.3	18.3-44.1
pH	4.7	4.4-5.2	4.6	4.2-5.1	4.5	4.1-5.2	4.7	4.2-5.3
organic matter (%)	6.32	5.12-8.83	6.26	4.13-10.09	6.68	3.75-10.17	5.39	3.41-8.39
mg NH_4^+-N g-1 DM	0.085	0.023-0.148	0.089	0.036-0.204	0.098	0.028-0.457	0.082	0.040-0.201
mg NO_3^--N g-1 DM	0.020	0.016-0.027	0.025	0.017-0.038	0.023	0.014-0.034	0.026	0.012-0.050
Total C (%)	1.57	0.62-2.74	1.78	0.27-3.61	1.80	0.48-2.81	1.46	0.33-3.49
Total N (%)	0.10	0.04-0.17	0.11	0.02-0.19	0.11	0.03-0.16	0.09	0.03-0.20
C:N (wt: wt)	16.0	15.0-18.4	16.2	13.5-19.0	15.9	13.9-17.6	14.9	11.0-17.5
Water								
mg NH_4^+-N L-1	0.101	MDL-0.474	.206	0.019-0.923	0.263	0.068-1.065	9.972	MDL-197.441
mg NO_3^--N L-1	0.173	MDL-1.118	0.007	MDL-0.020	0.169	MDL-0.413	0.769	MDL-4.055
Total Dissolved	0.035	MDL-0.087	0.088	0.27-0.286	0.072	0.025-0.176	0.075	MDL-0.245

3.2.3 Nitrous oxide (N₂O)

The wetland N_2O fluxes

were near zero in all plots across all

seasons Fig. 2C). Specifically, N_2O

fluxes were highest in shallow land

plots (-0.006 ± 0.117 mg N_2O-N m^{-2}

hr^{-1}) and lowest in flooded plots (-

0.083 ± 0.243 mg N_2O-N m^{-2} hr^{-1}).

Average N_2O flux rates across

seasons ranged from -0.05 ± 0.20 to

-0.02 ± 0.15 mg N_2O-N m^{-2} hr^{-1}.

Figure 2. Boxplots of seasonal CH_4 (A), CO_2 (B), and N_2O (C) fluxes for flooded (blue circles) and shallow land (green triangles) plots. Spring = Mar-May, Summer = Jun-Aug, Fall = Sep-Nov, Winter = Dec-Feb. Letters indicate significantly different groups based on estimated marginal means with Tukey post-hoc assessment at p<0.05 using 95% confidence levels (Table S2).

Unlike CH_4 and CO_2 fluxes, hydrology and season did not influence N_2O fluxes. In this case, the

null model had a ΔAICc score of 0 and the greatest AICc weight (0.69) (Table S1.1). However,

the models with hydrology or season had similar ΔAICc scores (2.98, 3.98) but lower AICc

weights (0.15, 0.11) when compared to the null model. Comparison of the parameters hydrology

and season using estimated marginal means at 95% confidence level suggests season has the

greatest impact on N_2O fluxes (Table S1.2).

3.3 Global warming potential

In terms of GWP, flooded plots have a two-fold increase in magnitude of total warming

potential than shallow land plots especially during warmer months (2418.63, 58.46 CO_2

equivalents (mg m^{-2} hr^{-1}), respectively) (Fig. 3). This is primarily due to increases in CH_4

concentrations, which make up 99% of the CO_2 equivalents in both flooded and shallow land

plots (Fig. 3). Over one year, flooded areas within the wetland had the potential to produce about

7161.27 kg CO_2 equivalents, while shallow land areas produced about 93.20 kg CO_2 equivalents.

Figure 3. Monthly estimated global warming potentials of methane and carbon dioxide evaluated as carbon dioxide equivalents (+ standard error) measured over a year at a constructed wetland. Proportion of grey for each bar represents mg CH_4-C m^{-2} hr^{-1} converted to CO_2 equivalents and proportion of black for each bar represents mg CO_2-C m^{-2} hr^{-1}. Average mg CH_4-C m^{-2} hr^{-1} was multiplied by 28, which is the estimated increased radiative force of methane when compared to CO_2.

3.4 Denitrification potential

Figure 4. Boxplots of potential denitrification rates according to hydrology and season based on denitrification enzyme assay using the acetylene block method. Spring = Mar-May, Summer = Jun-Aug, Fall=Sep-Nov, Winter=Dec-Feb. Different letters indicate significantly different groups by Tukey adjusted estimated marginal means p<0.05. Blue circles indicate flooded plots and green triangles shallow land.

When nitrate is readily available and anoxic conditions are present, sediments from flooded and shallow land zones have similar denitrification potential (Fig. 4). In flooded and shallow land plots across all seasons, potential denitrification rates were 24.45 ± 20.18 and 20.29 ± 15.14 ng N_2O-N hr^{-1} g^{-1} dry mass, respectively. However, denitrification potential varied by season; we measured higher rates in spring and summer $(34.12 \pm 21.09$ and 25.16 ± 13.51 ng N_2O N hr^{-1} g^{-1} dry mass) than in fall and winter $(10.55 \pm 6.17$ and 16.45 ± 13.72 N_2O ng N hr^{-1} g^{-1} dry mass) months. The model with season only explained the most variation in potential denitrification rates based on a ΔAICc score of 0 and AICc weight of 0.78 (Table S1.3). The fixed effect of season explained 18% of the variation; however, the null model explained the majority of variation (42%) (Table S1.3). We selected the season only model even though the model with hydrology and season had a similar ΔAICc (1.27) and AICc

weight of 0.21, but there was little improvement in marginal and conditional R^2 values (Table

S1.3).

3.5 Microbial community analysis

Hydrology, and to a lesser extent season, influenced microbial community composition.

Illumina amplicon sequencing of the 16S rRNA V4 region resulted in 454,709 reads before and

435,331 reads after removing singletons and doubletons. Each sample (n=22) then comprised a

minimum of 12,510 reads and all samples were rarified to this minimum and resampled. The

final dataset contained a total of 14,563 OTUs among all rarified samples. Microbial community

composition in flooded plots were distinct from shallow land plots (PERMANOVA, R^2= 0.0886,

p=0.007; Fig. 5). Indicator species analysis identified one Operational Taxonomic Unit (OTU;

microbial taxon defined at 97% sequence similarly), in the family Syntrophaceae, that was

unique to flooded plots and identified eight OTUs that represented the shallow land plots. The

shallow land indicator taxa included OTUs from two unclassified bacteria, two

Betaproteobacteria, one Acidobacteria Gp6, and one *Geobacter* (Table S5). Further examination

of taxa with >0.6% relative abundance of the total community suggests that flooded zones are

dominated by taxa reported to be involved in methanogenic degradation of hydrocarbons (e.g.,

Methanomicrobia, Methylocystis, and Syntrophaceae), while shallow land zones have fewer

methanogens in comparison to flooded zones (Fig. 6). We chose to look at the top 0.6% of

microbial groups within the community because this threshold allowed us to further examine

changes in relative abundance of the indicator species Syntrophaceae; this group is implicated in

methanogenic hydrocarbon degradation and has greater relative abundance in flooded compared

to shallow land areas. In addition, shallow land zones are enriched with taxa known to be

involved in nitrogen cycle transformations (e.g., Bradyrhizobium, Nitrososphaera,

Hyphomicrobium) (Fig. 6).

Figure 5. Ordination plot based on Principal Coordinates Analysis depicting sediment microbial community composition. Each point represents the centroid and range across season and sampling location. Symbols represent sampling period (circle = spring, square = fall). Spring samples were collected in May and fall samples were collected in October. Colors represent sampling plots (blue = flooded, white = shallow land).

3.6 Analyses of relationship between GHGs, sediment chemistry, and microbial community

composition

Sediment chemical

properties shape microbial

community composition and

microbial community

composition influences GHG

fluxes more than sediment

chemistry properties.

Distance-based partial least

square regression analysis

suggests that microbial

community composition

explained 82.4%, 78.6%, and

79.1%, of variation in CH_4,

CO_2, and N_2O fluxes,

respectively, by components 1 and 2. Sediment chemistry components 1 and 2 best explained

variation in N_2O production (51.4%) but less so for CH_4 or CO_2 production (21.0%, 16.6%).

Mantel correlation analyses revealed a positive relationship between patterns in microbial

community composition and soil properties (r = 0.52, p = 0.003).

450 **4. Discussion**

Figure 6. Heat map of bacterial and archaeal taxa observed at >0.6% relative abundance. Color gradient (cool to warm, green to red) represents microbial relative abundance ~0.6 to 1.5%.

Examination of

microbial processes occurring at permanently and periodically inundated locations within a

stormwater CW revealed that modifications to wetland design could enhance water quality (via

denitrification) and reduce GHG production. Other studies comparing different types of CWs

indicate that free water surface (FWS) wetlands, like this study site, are known to produce

copious amounts of GHGs compared to vertical or horizontal subsurface flow CWs (Mander et

al., 2014; McPhillips and Walter, 2015). While vertical or subsurface CWs may reduce GHG

production, this type of installation is more costly than FWS wetlands and may not always be

practical. We recognize that this study focuses on a single CW; however, our results demonstrate

the need for more comprehensive studies exploring CW design features that can be modified to

enhance beneficial microbial ecosystem services (i.e., nitrogen removal) while reducing

ecosystem disservices (i.e., GHG emissions).

At our focal CW, CH_4 was the dominant GHG produced in flooded plots across all

seasons. Hydrologic conditions accounted for the most variation in CH_4 fluxes and microbial

community composition. Since high NO_3^- concentrations suppress methanogenesis (Kim et al.,

2015), it is likely the combination of anoxic conditions (McPhillips and Walter, 2015), low NO_3^-

concentrations in sediments and water, and availability of organic carbon in flooded plots

provided optimal conditions for CH_4 production (Rahman et al., 2019). However, season and

microbial community composition were stronger predicators than hydrology and distance from

inlet of CO_2 and N_2O rates within the wetland. We propose that season and hydrology played a

stronger role than distance from inlet in GHG fluxes and denitrification potential due to the small

size of the study CW. Seasonal differences in temperature and vegetation status (i.e., organic

carbon inputs) within this study influenced CO_2 and N_2O rates to a greater degree than

hydrology. In other studies, temperature and substrate availability strongly determined rates of

organic decomposition and denitrification (Davidson and Janssens, 2006; Knowles, 1982;

Moinet et al., 2018).

Overall, compared to wastewater treatment wetlands, flooded plots within our study

wetland represent a considerable source of GWP primarily due to CH_4 emissions. In order to

simultaneously compare the biogenic GHGs within the CW, we quantified GWP associated with

flooded and shallow land zones. In terms of GWP, CH_4 from flooded plots was the primary

carbon source within the wetland. While both CO_2 and CH_4 production was greatest in flooded

zones compared to shallow land zones, CH_4 production greatly exceeded CO_2 production leading

to high GWP within the wetland. Compared to surface flow wastewater treatment wetlands

receiving municipal or agricultural runoff, CO_2 emissions at our study wetland in flooded plots

were similar but approximately 95% lower in shallow land plots (Jahangir et al., 2016; Mander et

al., 2014). In contrast, CH_4 emissions at our study wetland had similar CH_4 rates in shallow land

plots when compared to wastewater treatment wetlands, but approximately 94% greater

emissions in flooded plots compared to wastewater treatment wetlands (Jahangir et al., 2016;

Mander et al., 2014). In terms of GWP, flooded plots in our study wetland have a GWP that is

98% greater than in wastewater treatment wetlands. The increased CH_4 emission in our study

wetland compared to wastewater wetlands is likely due to the lower availability of nutrients in

surface water and sediments.

Further, GHG emissions could be partially explained by microbial community structure.

This is due to hydrologic conditions in flooded zones providing optimal habitat for obligate

anaerobes that participate in methanogenesis. Methanogenesis is suppressed by even low oxygen

concentrations (Fetzer et al., 1993; Liu et al., 2008) and we could not detect methanogens in

shallow land areas where conditions can fluctuate between oxic and anoxic environments.

Additionally, analysis of microbial community structure suggests that hydrocarbon degradation,

a potential microbial ecosystem benefit, may be occurring in flooded plots. Hydrocarbons are

becoming recognized as a common contaminant in urban wetlands with the potential to reduce

and even harm wildlife taking refuge in these habitats (Clevenot et al., 2018; Mahler et al.,

2014). This finding demonstrates the value of examining the microbial community composition

when evaluating beneficial wetland ecosystem functions.

In retrospect of this study, we highlight three CW features for consideration to reduce

ecosystem function disservices: (i) surface area of hydrologic zones, (ii) incoming nutrient and

pollutant loads, and (iii) the anticipated number of rain days. Results from this study and others

indicate that areas that fluctuate between oxic and anoxic conditions, which occur in CW shallow

land areas in this study, are sites of reduced CH_4 production. This is due to the inhibition of

methanogenesis and a greater abundance of methanotrophs which consume methane

(Chowdhury and Dick, 2013; Lew and Glińska-Lewczuk, 2018). Additionally, even in CWs with

low nitrate availability, shallow land areas could potentially support enhanced denitrification due

to coupling of nitrification, which occurs during oxic periods and with steady organic carbon

inputs from vegetation (Mander et al., 2014; Rahman et al., 2019). Flooded zones increase water

storage capacity and provide habitat for predator species that control nuisance species; therefore,

elimination of flooded zones may not be desirable. However, an increase in shallow land area

and a reduction in surface area of flooded zones could reduce GHG emissions without sacrificing

benefits of flooded zones.

Secondly, substrate availability can directly impact microbial process rates. Compared to

the neighboring Tar River (0.52 mg NO_3-N L^{-1}), seasonal NO_3^- concentrations within the surface

water of the CW were much lower (0.007-0.173 mg NO_3-N L^{-1}) except during winter (0.769 mg

NO_3-N L^{-1}) (Humphrey et al., 2019). This increase in nitrate during winter is likely due to the

senescence of *Typha* spp. and *P. cordata* vegetation (Bachand and Horne, 2000), which are the

main plants growing in and around the main channel. Considering that NO_3^- availability limits

denitrification, ecosystems that receive runoff from agriculture, livestock, or are adjacent to a

high density of septic systems are areas of high denitrification potential and low methane

production (Naylor et al., 2018). Recent studies show that iron (II) (Fe^{2+}) additions can increase

N removal from vertical and horizontal subsurface CWs (Song et al., 2016; Zhang et al., 2019).

In the case of existing wetlands, further investigation into amendments as a way to increase

denitrification potential is warranted. Additionally, depending on the upstream runoff source,

pollutants such as hydrocarbons may be better targets for remediation within intentionally

designed CWs.

Lastly, we suggest that the frequency of rain days or inter-event duration is taken into

account during SCM planning (Andersen et al., 2017). Our results represent ambient conditions,

at least 48 hours after a storm event, which represent more than half the year for this CW. Within

this particular wetland, reducing the permanently flooded surface area could reduce GHGs while

still providing N removal benefits. In areas where rain events are less frequent (i.e., greater inter-

event durations) and more intense, SCMs that are not permanently flooded, such as dry

infiltration basins or dry detention basins, could provide relief for intense runoff events with

reduced GHG production (McPhillips and Walter, 2015; Morse et al., 2017). Therefore, we

suggest frequency of rain events as another factor to consider when planning SCM design.

5. Conclusion

Context matters when designing CWs in urban watersheds. This study demonstrates the

importance of considering microbial controls on biogeochemical processes within SCMs in order

to reduce tradeoffs between water quality and GHG production. In the case of this urban

stormwater CW, which is located in a parking lot of a highly urbanized area, we suggest that

reducing flooded areas and increasing shallow land area would reduce GHG emissions while still

providing nutrient removal benefit. However, our analysis of the microbial community also

indicates that potential hydrocarbon decomposition could be supported in flooded zones and

warrants a deeper investigation of this potential service. This study also highlights the need for a

comprehensive study of CW design features to better understand which design features and when

those features provide the most ecosystem benefits. By examining both microbial process rates

and the microbial community composition, we can approach SCM design and implementation in

a holistic way that accounts for ecosystem services and disservices.

Declaration of Competing Interest

None.

Acknowledgements

We thank B. Hinckley, A. Garcìa, and L. Armstrong for assistance with field research

and sample processing. We also thank Sound Rivers for installation of the constructed wetland

and ECU Grounds personnel and manager J. Gill for site maintenance. We also thank E. Field, L.

Kinsman-Costello, M. McCoy, and anonymous reviewers for feedback on earlier versions of this

manuscript. This work was supported by East Carolina University and the North Carolina Sea

Grant/Water Resources Research Institute Graduate Student Research Fellowship and the

National Science Foundation Graduate Research Fellowship to RBB. All code and data used

uyhin this study can be found in a public GitHub repository (DOI: 10.5281/zenodo.3924697)

and the NCBI SRA (BioProject PRJNA613095).

Appendix A. Supplementary data

Supplementary material related to this article can be found at: ###

References

Altshuler, I., Hamel, J., Turney, S., Magnuson, E., Lévesque, R., Greer, C.W., Whyte, L.G.,

2019. Species interactions and distinct microbial communities in high Arctic permafrost

affected cryosols are associated with the CH4 and CO2 gas fluxes. Environ. Microbiol. 21,

3711–3727. https://doi.org/10.1111/1462-2920.14715

Andersen, J.S., Lerer, S.M., Backhaus, A., Jensen, M.B., Danielsen Sørup, H.J., 2017.

Characteristic rain events: A methodology for improving the amenity value of stormwater

control measures. Sustain. 9. https://doi.org/10.3390/su9101793

Apprill, A., Mcnally, S., Parsons, R., Weber, L., 2015. Minor revision to V4 region SSU rRNA

806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat. Microb.

Ecol. 75, 129–137. https://doi.org/10.3354/ame01753

Bachand, P.A.M., Horne, A.J., 2000. Denitrification in contructed free-water surface wetlands:

II. Effects of vegetation and temperature. Ecol. Eng. 14, 17–32.

Barton, K., 2019. Package "MuMIn."

Bates, D., Machler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using

lme4. J. Stat. Softw. 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Bell, C.D., Tague, C.L., McMillan, S.K., 2019. Modeling Runoff and Nitrogen Loads From a

Watershed at Different Levels of Impervious Surface Coverage and Connectivity to Storm

Water Control Measures. Water Resour. Res. 55, 2690–2707.

https://doi.org/10.1029/2018WR023006

Boj, E., Caballe, A., Delicado, P., Fortiana, J., 2017. Package ' dbstats .'

Brindha, R.K., Vasudevan, N., 2018. Methane oxidation capacity of methanotrophs isolated from

different soil ecosystems. Int. J. Environ. Sci. Technol. 15, 1931–1940.

https://doi.org/10.1007/s13762-017-1546-1

Burgin, A.J., Yang, W.H., Hamilton, S.K., Silver, W.L., 2011. Beyond carbon and nitrogen: how

the microbial energy economy couples elemental cycles in diverse ecosystems. Front. Ecol.

Environ. 9, 44–52. https://doi.org/10.1890/090227

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-lyons, D., Lozupone, C.A., Turnbaugh, P.J.,

Fierer, N., Knight, R., 2010. <Global patterns of 16S rRNA diversity at a depth of millions

of sequences per sample.pdf>. Proc. Natl. Acad. Sci. U. S. A. 108, 4516–4522.

https://doi.org/10.1073/pnas.1000080107/-

/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1000080107

Chowdhury, T.R., Dick, R.P., 2013. Ecology of aerobic methanotrophs in controlling methane

fluxes from wetlands. Appl. Soil Ecol. 65, 8–22.

https://doi.org/10.1016/j.apsoil.2012.12.014

Clevenot, L., Carré, C., Pech, P., 2018. A Review of the factors that determine whether

stormwater ponds are ecological traps and/or high-quality breeding sites for amphibians.

Front. Ecol. Evol. 6, 1–12. https://doi.org/10.3389/fevo.2018.00040

Conrad, R., 2007. Microbial Ecology of Methanogens and Methanotrophs. Adv. Agron. 96, 1–

63. https://doi.org/10.1016/S0065-2113(07)96005-8

Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and

feedbacks to climate change. Nature 440, 165–173. https://doi.org/10.1038/nature04514

Demuzere, M., Orru, K., Heidrich, O., Olazabal, E., Geneletti, D., Orru, H., Bhave, A.G., Mittal,

N., Feliu, E., Faehnle, M., 2014. Mitigating and adapting to climate change: Multi-

functional and multi-scale assessment of green urban infrastructure. J. Environ. Manage.

146, 107–115. https://doi.org/10.1016/j.jenvman.2014.07.025

Fetzer, S., Bak, F., Conrad, R., 1993. Sensitivity of methanogenic bacteria from paddy soil to

oxygen and desiccation. FEMS Microbiol. Ecol. 12, 107–115.

https://doi.org/10.1111/j.1574-6941.1993.tb00022.x

Fetzer, S., Conrad, R., 1993. Effect of redox potential on methanogenesis by Methanosarcina

barkeri. Arch. Microbiol. 160, 108–113. https://doi.org/10.1007/BF00288711

Gorsky, A.L., Racanelli, G.A., Belvin, A.C., Chambers, R.M., 2019. Greenhouse gas flux from

stormwater ponds in southeastern Virginia (USA). Anthropocene 28.

https://doi.org/10.1016/j.ancene.2019.100218

Heiri, O., Lotter, A., Lemcke, G., 2001. Loss on ignition as a method for estimating organic and

carbonate content in sediments. J. Paleolimnol. 25, 101–110.

https://doi.org/10.1017/CBO9781107415324.004

Hoogsteen, M.J.J., Lantinga, E.A., Bakker, E.J., Groot, J.C.J., Tittonell, P.A., 2015. Estimating

soil organic carbon through loss on ignition: Effects of ignition conditions and structural

water loss. Eur. J. Soil Sci. 66, 320–328. https://doi.org/10.1111/ejss.12224

Hothorn, T., Bretz, F., Westfall, P., Heiberger, R.M., Schuetzenmeister, A., Scheibe, S., 2020.

Package ' multcomp .'

Humphrey, C., Iverson, G., Skibiel, C., Sanderford, C., Blackmon, J., 2019. Geochemistry of

Flood Waters from the Tar River, North Carolina Associated with Hurricane Matthew.

Resources 8, 48. https://doi.org/10.3390/resources8010048

Hurvich, C., Tsai, C., 1993. A CORRECTED AKAIKE INFORMATIONON CRITERION FOR

VECTOR AUTOREGRESSIVE MODEL SELECTION. J. Time Ser. Anal. 14, 271–279.

IPCC, 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and

III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Ipcc.

IPCC - Intergovernmental Panel on Climate Change, 2014. Climate Change 2014 Synthesis

Report (Unedited Version).

Jahangir, M.M.R., Richards, K.G., Healy, M.G., Gill, L., Müller, C., Johnston, P., Fenton, O.,

2016. Carbon and nitrogen dynamics and greenhouse gas emissions in constructed wetlands

treating wastewater: A review. Hydrol. Earth Syst. Sci. 20, 109–123.

https://doi.org/10.5194/hess-20-109-2016

Kim, S.Y., Veraart, A.J., Meima-Franke, M., Bodelier, P.L.E., 2015. Combined effects of

carbon, nitrogen and phosphorus on CH4 production and denitrification in wetland

sediments. Geoderma 259–260, 354–361. https://doi.org/10.1016/j.geoderma.2015.03.015

Knowles, R., 1982. Denitrification. Microbiol. Rev. 46, 43–70.

Koch, B.J., Febria, C.M., Gevrey, M., Wainger, L.A., Palmer, M.A., 2014. Nitrogen Removal by

Stormwater Management Structures: A Data Synthesis. J. Am. Water Resour. Assoc. 50,

1594–1607. https://doi.org/10.1111/jawr.12223

Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D., 2013. Development of

a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence

data on the miseq illumina sequencing platform. Appl. Environ. Microbiol. 79, 5112–5120.

https://doi.org/10.1128/AEM.01043-13

Le Mer, J., Roger, P., 2001. Production, oxidation, emission and consumption of methane by

soils: A review. Eur. J. Soil Biol. 37, 25–50. https://doi.org/10.1016/S1164-5563(01)01067-

6

Lee, C.G., Fletcher, T.D., Sun, G., 2009. Nitrogen removal in constructed wetland systems. Eng.

Life Sci. 9, 11–22. https://doi.org/10.1002/elsc.200800049

Lenth, R., 2017. Package ' lsmeans .' https://doi.org/10.1080/00031305.1980.10483031>.NOTE

Lew, S., Glińska-Lewczuk, K., 2018. Environmental controls on the abundance of

methanotrophs and methanogens in peat bog lakes. Sci. Total Environ. 645, 1201–1211.

https://doi.org/10.1016/j.scitotenv.2018.07.141

Line, D.E., White, N.M., 2007. Effects of Development on Runoff and Pollutant Export. Water

Environ. Res. 79, 185–190. https://doi.org/10.2175/106143006x111736

Liu, C. Te, Miyaki, T., Aono, T., Oyaizu, H., 2008. Evaluation of methanogenic strains and their

ability to endure aeration and water stress. Curr. Microbiol. 56, 214–218.

https://doi.org/10.1007/s00284-007-9059-7

Mahler, B.J., Van Metre, P.C., Foreman, W.T., 2014. Concentrations of polycyclic aromatic

hydrocarbons (PAHs) and azaarenes in runoff from coal-tar- and asphalt-sealcoated

pavement. Environ. Pollut. 188, 81–87. https://doi.org/10.1016/j.envpol.2014.01.008

Mander, Ü., Dotro, G., Ebie, Y., Towprayoon, S., Chiemchaisri, C., Nogueira, S.F., Jamsranjav,

B., Kasak, K., Truu, J., Tournebize, J., Mitsch, W.J., 2014. Greenhouse gas emission in

constructed wetlands for wastewater treatment: A review. Ecol. Eng. 66, 19–35.

https://doi.org/10.1016/j.ecoleng.2013.12.006

Marton, J.M., Creed, I.F., Lewis, D.B., Lane, C.R., Basu, N.B., Cohen, M.J., Craft, C.B., 2015.

Geographically isolated wetlands are important biogeochemical reactors on the landscape.

Bioscience 65, 408–418. https://doi.org/10.1093/biosci/biv009

McPhillips, L., Walter, M.T., 2015. Hydrologic conditions drive denitrification and greenhouse

gas emissions in stormwater detention basins. Ecol. Eng. 85, 67–75.

https://doi.org/10.1016/j.ecoleng.2015.10.018

Millar, N., Urrea, A., Kahmark, K., Shcherbak, I., Robertson, G.P., Ortiz-Monasterio, I., 2018.

Nitrous oxide (N2O) flux responds exponentially to nitrogen fertilizer in irrigated wheat in

the Yaqui Valley, Mexico. Agric. Ecosyst. Environ. 261, 125–132.

https://doi.org/10.1016/j.agee.2018.04.003

Mitsch, W.J., Bernal, B., Nahlik, A.M., Mander, Ü., Zhang, L., Anderson, C.J., Jørgensen, S.E.,

Brix, H., 2013. Wetlands, carbon, and climate change. Landsc. Ecol. 28, 583–597.

https://doi.org/10.1007/s10980-012-9758-8

Moinet, G.Y.K., Hunt, J.E., Kirschbaum, M.U.F., Morcom, C.P., Midwood, A.J., Millard, P.,

2018. The temperature sensitivity of soil organic matter decomposition is constrained by

microbial access to substrates. Soil Biol. Biochem. 116, 333–339.

https://doi.org/10.1016/j.soilbio.2017.10.031

Morse, N.R., McPhillips, L.E., Shapleigh, J.P., Walter, M.T., 2017. The Role of Denitrification

in Stormwater Detention Basin Treatment of Nitrogen. Environ. Sci. Technol. 51, 7928–

7935. https://doi.org/10.1021/acs.est.7b01813

Naylor, E., Humphrey, C., Kelley, T., Easter, L., Iverson, G., 2018. Evaluation of nitrate

concentrations and potential sources of nitrate in private water supply wells in North

Carolina. J. Environ. Health 80, 16–23.

NCDEQ, 2019. Stormwater Design Manual: C-4 Stormwater Wetland [WWW Document].

North Carolina Dep. Environ. Qual. URL https://deq.nc.gov/sw-bmp-manual

NCDEQ, 2018. Stormwater Control Measure Credit Document [WWW Document]. North

Carolina Dep. Environ. Qual. URL https://deq.nc.gov/sw-bmp-manual

O'Driscoll, M., Clinton, S., Jefferson, A., Manda, A., McMillan, S., 2010. Urbanization effects

on watershed hydrology and in-stream processes in the southern United States. Water

(Switzerland) 2, 605–648. https://doi.org/10.3390/w2030605

Oksanen, J., 2015. Vegan□: ecological diversity 1, 1–12. https://doi.org/10.1029/2006JF000545

Paerl, H.W., Hall, N.S., Peierls, B.L., Rossignol, K.L., 2014. Evolving Paradigms and

Challenges in Estuarine and Coastal Eutrophication Dynamics in a Culturally and

Climatically Stressed World. Estuaries and Coasts 37, 243–258.

https://doi.org/10.1007/s12237-014-9773-x

Parada, A.E., Needham, D.M., Fuhrman, J.A., 2016. Every base matters: Assessing small subunit

rRNA primers for marine microbiomes with mock communities, time series and global field

samples. Environ. Microbiol. 18, 1403–1414. https://doi.org/10.1111/1462-2920.13023

Parkin, T.B., Venterea, R.T., Follett, R.F. (editor), 2010. Sampling Protocols. Chapter 3.

Chamber-Based Trace Gas Flux Measurements. In Sampling Protocols. Available at:

www.ars.usda.gov/research/GRACEnet 2010, 3–1 to 3–39.

Payne, E.G.I., Fletcher, T.D., Cook, P.L.M., Deletic, A., Hatt, B.E., 2014. Processes and drivers

of nitrogen removal in stormwater biofiltration. Crit. Rev. Environ. Sci. Technol. 44, 796–

846. https://doi.org/10.1080/10643389.2012.741310

Peralta, A.L., Ludmer, S., Kent, A.D., 2013. Hydrologic history influences microbial community

composition and nitrogen cycling under experimental drying/wetting treatments. Soil Biol.

Biochem. 66, 29–37. https://doi.org/10.1016/j.soilbio.2013.06.019

Peralta, A.L., Ludmer, S., Matthews, J.W., Kent, A.D., 2014. Bacterial community response to

changes in soil redox potential along a moisture gradient in restored wetlands. Ecol. Eng.

73, 246–253. https://doi.org/10.1016/j.ecoleng.2014.09.047

R Core Team, 2019. R: A language and environment for statistical computing. R Found. Stat.

Comput. Vienna, Austria.

Rabalais, N.N., Turner, R.E., Díaz, R.J., Justić, D., 2009. Global change and eutrophication of

coastal waters. ICES J. Mar. Sci. 66, 1528–1537. https://doi.org/10.1093/icesjms/fsp047

Rahman, M.M., Roberts, K.L., Warry, F., Grace, M.R., Cook, P.L.M., 2019. Factors controlling

dissimilatory nitrate reduction processes in constructed stormwater urban wetlands.

Biogeochemistry 142, 375–393. https://doi.org/10.1007/s10533-019-00541-0

Reisinger, A.J., Groffman, P.M., Rosi-Marshall, E.J., 2016. Nitrogen-cycling process rates

across urban ecosystems. FEMS Microbiol. Ecol. 92, fiw198.

https://doi.org/10.1093/femsec/fiw198

Rey-Sanchez, C., Bohrer, G., Slater, J., Li, Y.F., Grau-Andrés, R., Hao, Y., Rich, V.I., Davies,

G.M., 2019. The ratio of methanogens to methanotrophs and water-level dynamics drive

methane transfer velocity in a temperate kettle-hole peat bog. Biogeosciences 16, 3207–

3231. https://doi.org/10.5194/bg-16-3207-2019

Roberts, D., 2016. Package ' labdsv .'

Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open

source tool for metagenomics. PeerJ 4, e2584. https://doi.org/10.7717/peerj.2584

Schaller, J.L., Royer, T. V., David, M.B., Tank, J.L., 2004. Denitrification associated with plants

and sediments in an agricultural stream. J. North Am. Benthol. Soc. 23, 667–676.

https://doi.org/10.1899/0887-3593(2004)023<0667:DAWPAS>2.0.CO;2

Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski,

R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B., Thallinger, G.G., Van

Horn, D.J., Weber, C.F., 2009. Introducing mothur: Open-source, platform-independent,

community-supported software for describing and comparing microbial communities. Appl.

Environ. Microbiol. 75, 7537–7541. https://doi.org/10.1128/AEM.01541-09

Smith, M.S., Tiedje, J.M., 1979. Phases of denitrification following oxygen depletion in soil.

Soil Biol. Biochem. 11, 261–267. https://doi.org/10.1016/0038-0717(79)90071-3

Song, X., Wang, S., Wang, Y., Zhao, Z., Yan, D., 2016. Addition of Fe2+ increase nitrate

removal in vertical subsurface flow constructed wetlands. Ecol. Eng. 91, 487–494.

https://doi.org/10.1016/j.ecoleng.2016.03.013

Tiedje, J.M., 1988. Ecology of denitrification and dissimilatory nitrate reduction to ammonium.

Environ. Microbiol. Anaerobes 179–244.

Tiedje, J.M., Simkins, S., Groffman, P.M., 1989. Perspectives on measurement of denitrification

in the field including recommended protocols for acetylene based methods. Plant Soil 115,

261–284. https://doi.org/10.1007/BF02202594

Wall, L.G., Tank, J.L., Royer, T. V., Bernot, M.J., 2005. Spatial and temporal variability in

sediment denitrification within an agriculturally influenced reservoir. Biogeochemistry 76,

85–111. https://doi.org/10.1007/s10533-005-2199-6

Wen, X., Unger, V., Jurasinski, G., Koebsch, F., Horn, F., Rehder, G., Sachs, T., Zak, D.,

Lischeid, G., Knorr, K.H., Böttcher, M.E., Winkel, M., Bodelier, P.L.E., Liebner, S., 2018.

Predominance of methanogens over methanotrophs in rewetted fens characterized by high

methane emissions. Biogeosciences 15, 6519–6536. https://doi.org/10.5194/bg-15-6519-

2018

Zhang, Y., Liu, X., Fu, C., Li, X., Yan, B., Shi, T., 2019. Effect of Fe2+ addition on chemical

oxygen demand and nitrogen removal in horizontal subsurface flow constructed wetlands.

Chemosphere 220, 259–265. https://doi.org/10.1016/j.chemosphere.2018.12.144

Figure Captions

Figure 1. Top: Landscape photograph of wetland. Center: Google Earth image of parking lot and

stormwater CW outlined in red. Bottom: Google Earth image of wetland with blue circles (1-5)

representing flooded plots and white circles (6-13) representing shallow land plots sampled for

GHG and sediments.

Table 1A & B. Summary of mean (range) of soil and water properties measured at the

constructed wetland. Annual wetland values are averages of all samples for the entire sampling

period. (1A) Hydrology (flooded vs. shallow land) values are averages of samples collected in

May (spring), July (summer), October (fall), and December (winter) for each hydrology type.

(1B) Seasonal sediment values are averages from of all plots based on samples collected in May

(spring), July (summer), October (fall), and December (winter). Seasonal water values are

averages of monthly samples from flooded plots where spring represents months May-June,

summer represents months July-September, fall represents months October-November, and

winter represents months December-February. (Abbreviations: DM = dry mass. MDL = below

detection limit).

Figure 2. Boxplots of seasonal CH_4 (A), CO_2 (B), and N_2O (C) fluxes for flooded (blue circles)

and shallow land (green triangles) plots. Spring = Mar-May, Summer = Jun-Aug, Fall = Sep-

Nov, Winter = Dec-Feb. Letters indicate significantly different groups based on estimated

marginal means with Tukey post-hoc assessment at p<0.05 using 95% confidence levels (Table

S2).

Figure 3. Monthly estimated global warming potentials of methane and carbon dioxide

evaluated as carbon dioxide equivalents (+ standard error) measured over a year at a constructed

wetland. Proportion of grey for each bar represents mg CH4-C m-2 hr-1 converted to CO2

equivalents and proportion of black for each bar represents mg CO2-C m-2 hr-1. Average mg

CH4-C m-2 hr-1 was multiplied by 28, which is the estimated increased radiative force of

methane when compared to CO_2.

Figure 4. Boxplots of potential denitrification rates according to hydrology and season based on

denitrification enzyme assay using the acetylene block method. Spring = Mar-May, Summer =

Jun-Aug, Fall=Sep-Nov, Winter=Dec-Feb. Different letters indicate significantly different

groups by Tukey adjusted estimated marginal means p<0.05. Blue circles indicate flooded plots

and green triangles shallow land.

Figure 5. Ordination plot based on Principal Coordinates Analysis depicting sediment microbial

community composition. Each point represents the centroid and range across season and

sampling location. Symbols represent sampling period (circle = spring, square = fall). Spring

samples were collected in May and fall samples were collected in October. Colors represent

sampling plots (blue = flooded, white = shallow land).

Figure 6. Heat map of bacterial and archaeal taxa observed at >0.6% relative abundance. Color

gradient (cool to warm, green to red) represents microbial relative abundance ~0.6 to 1.5%.