Title: A microbial perspective on balancing trade-offs in ecosystem functions in a constructed

stormwater wetland

Author: Regina B. Bledsoe[a]*, Samuel S. Austin[a], Eban Z. Bean[b], Ariane L. Peralta[a]

*Corresponding author

Email: ginabbledsoe@gmail.com, austins15@ecualumni.ecu.edu, ezbean@ufl.edu,

peraltaa@ecu.edu

Affiliations:

[a]East Carolina University, Department of Biology, Greenville, NC, 27858, USA

[b]University of Florida, Department of Agricultural and Biological Engineering, Gainesville, FL,

32611, USA

Abbreviations: corrected Akaike information criteria = AICc, NH_4^+ = ammonium, C = carbon,

CO_2 = carbon dioxide, CW = constructed wetland, N_2 = dinitrogen, GWP = global warming

potential, GHG = greenhouse gas, CH_4 = methane, N= nitrogen, NO_3^- = nitrate, NO = nitric

oxide, NO_2^- = nitrite, N_2O = nitrous oxide, OTU = operational taxonomic unit, PCR =

polymerase chain reaction, PERMANOVA = permutated analysis of variance, PCoA = principal

coordinates analysis, SCM = stormwater control measure

Abstract

Green stormwater infrastructure, such as constructed wetlands (CWs), is a type of

stormwater control measure that can reduce nutrient and pollutant loads in urban stormwater

runoff. Nutrient and pollutant removal processes within wetlands are primarily driven by

microbial functions that can also result in disservices such as greenhouse gas (GHG) emissions,

which can negate climate change mitigation. Specifically, microbial respiration by facultative

anaerobes in anoxic conditions is the primary pathway for nitrogen removal. Similar anoxic

conditions that support denitrifiers can also support obligate anaerobes that produce methane via

methanogenesis. In this study, we examined nitrogen removal potential, GHG production, and

microbial community structure within flooded and shallow land areas of a stormwater CW to

identify zones for CW design optimization. Our results indicate that permanently flooded zones

are sources of methane emissions and have the greatest contribution to climate change. However,

denitrification potential rates were similar across both flooded and shallow land zones. This

suggests that shallow land areas can provide nitrogen removal services with reduced GHG

emissions compared to flooded zones. In the case of this particular CW, a reduction of

permanently flooded zones within the wetland could decrease GHG emissions (i.e., disservice)

without limiting denitrification (i.e., benefit) potential of the wetland. We conclude that holistic

development and design of stormwater control measures, which account for microbial functions,

provides the opportunity to maximize benefits (i.e., nutrient and pollutant removal) and reduce

disservices (i.e., greenhouse gas emissions) of green stormwater infrastructure.

Keywords: Greenhouse gases, Denitrification, Methane, Stormwater, Urban ecosystems

Introduction

In urban areas, impervious surfaces increase nutrient and pollutant loads in stormwater runoff, thereby reducing water quality within neighboring watersheds (Bell et al., 2019; Line and White, 2007; O'Driscoll et al., 2010). Nutrient loadings are of particular concern due to adverse effects in downstream water quality (Paerl et al., 2014; Rabalais et al., 2009). Green stormwater infrastructure, such as constructed wetlands (CWs), are stormwater control measures (SCMs) that are designed to mimic natural ecosystems; SCMs provide relief from stormwater runoff and improve water quality in and around urban areas. The physical, chemical, and biological treatment processes that naturally occur in wetlands make CWs an effective approach to reduce nutrient loads in stormwater runoff (Koch et al., 2014; Mitsch et al., 2013; Payne et al., 2014; Reisinger et al., 2016). Thus, entities such as the North Carolina Department of Environmental Quality encourage use of SCMs as a way to reduce nutrient loadings to nutrient sensitive waterways by offering nutrient reduction credits. CWs that have a ponding depth of less than 12 inches and a hydraulic residence time of two to five days are credited with 40% removal of nitrogen (N) from stormwater runoff (NCDEQ, 2018) However, this nutrient reduction policy and similar policies outside of North Carolina do not consider the potential trade-off between water quality (e.g., N removal) and air quality (e.g., greenhouse gas (GHG) emissions). The unintended GHG production from stormwater CWs could negatively affect climate change mitigation and adaptation by reducing the carbon storage capacity typically provided by wetlands (Demuzere et al., 2014). Therefore, in order to meet management goals in a more holistic way (i.e., enhancing total N removal and reducing greenhouse gas emissions), a deeper understanding of microbial processes occurring within CWs are needed to optimize wetland design.

Plants and microorganisms carry out nutrient and pollutant removal processes that occur

within wetlands. While plants and microorganisms can assimilate ammonium (NH_4^+) and nitrate

(NO_3^-) ions into biomass, the primary pathway of complete N removal from the system is

microbial respiration via denitrification and to a lesser degree anaerobic ammonium oxidation

(Lee et al., 2009). Denitrification is typically coupled with nitrification in which NH_4^+ is first

transformed to nitrite (NO_2^-) and then NO_3^- by aerobic nitrifying microorganisms (Lee et al.,

2009). The process of complete denitrification occurs when NO_3^- is reduced to nitric oxide (NO),

then to nitrous oxide (N_2O), a potent GHG, and finally to dinitrogen (N_2), an inert and abundant

atmospheric gas (Knowles, 1982; Lee et al., 2009; Smith and Tiedje, 1979). Denitrification is an

anaerobic microbial process that requires anoxic conditions such as those found in saturated and

flooded soils within wetlands (Marton et al., 2015; Smith and Tiedje, 1979). Many denitrifying

microorganisms are facultatively anaerobic. These microorganisms can perform aerobic

respiration when oxygen is available but can use other alternate electron acceptors (i.e., nitrate)

when anoxic soil conditions arise (Tiedje, 1988). Due to this facultative anaerobic metabolism,

these microorganisms respond to hydrologic change with varying degrees depending on past

environmental conditions (Peralta et al., 2014, 2013).

Similar anoxic conditions that support the beneficial process of complete denitrification

(transformation of NO_3^- to N_2) can also support production of methane (CH_4), a GHG, which is

considered an ecosystem disservice (Demuzere et al., 2014). Methanogens are obligate anaerobes

that produce CH_4 via methanogenesis but only under anoxic conditions when the environment is

depleted of other electron acceptors (e.g., nitrate, sulfate, and ferric iron) (Fetzer and Conrad,

1993; Liu et al., 2008). When organic matter is decomposed by fermentation, methanogens can

use the byproducts of acetate, hydrogen, and CO_2 for energy and produce CH_4 (Conrad, 2007).

Methane production can also support the consumption of CH_4 by aerobic methanotrophs, which

have been reported to consume 45-90% of CH_4 produced (Brindha and Vasudevan, 2018;

Conrad, 2007; Le Mer and Roger, 2001). Systems in which the ratio of methanogens is greater

than methanotrophs are typically sources of methane, resulting in the ecosystem serving as a

methane source (Altshuler et al., 2019; Rey-Sanchez et al., 2019; Wen et al., 2018). Since

methanogens are obligate anaerobes, even low amounts of oxygen can suppress methanogenesis

(Fetzer and Conrad, 1993; Le Mer and Roger, 2001). However, methanotrophs are aerobes, and

there is typically greater CH_4 oxidation in pulsing systems that are not permanently inundated

(Chowdhury and Dick, 2013). Therefore, anoxic conditions that can support complete

denitrification can also result in GHG emissions, especially when nitrate is limiting (Burgin et

al., 2011).

The three major biogenic GHGs of concern, CO_2, CH_4, and N_2O, are known to contribute

to increased atmospheric warming and thereby influence global climate change. While CO_2 is the

most abundant GHG and is therefore most concerning, CH_4 and N_2O gases have a greater global

warming potential (GWP) than CO_2 (IPCC, 2014). Specifically, per molecule, CH_4 and N_2O

have GWP of 28 and 265, respectively, over a 100 year time-scale when compared to CO_2 which

is equal to 1 (IPCC, 2014). Generally, atmospheric quantities of CH_4 are lower than CO_2 and

N_2O is typically measured in the lowest atmospheric quantities, but the increased GWP of these

gases contribute disproportionately to increasing atmospheric GHG concentrations.

To improve water quality management of CWs in the urban landscape, both ecosystem

services (e.g., complete denitrification) and disservices (e.g., GHG production) should

considered during the planning and design phase (Demuzere et al., 2014). To accommodate

stormwater runoff and water quality functions, CWs generally have four main zones: upland

(rarely submerged except during large runoff events), shallow land (soil is saturated but only

temporarily inundated following runoff events), shallow pools (designed to be permanently

inundated except during droughts) and deep pools (permanently inundated) (NCDEQ, 2019). In

CWs, ammonification and nitrification are optimized to occur in the shallow land zones as

stormwater percolates into aerated soils as runoff enters. If soils in shallow lands are saturated

for sufficient durations, denitrification is expected to eventually occur as well. It is in the shallow

water zones where denitrification is conventionally thought to occur due to saturated, anoxic

conditions. In this study we focus on complete denitrification as the ecosystem service since this

process typically accounts for the most N removal in CWs (Rahman et al., 2019).

The goal of this study is to identify areas within a stormwater CW that support

denitrification and reduce GHG emissions, especially due to methanogenesis. We hypothesize

that the redox shifts occurring in shallow land and flooded zones will both support the anaerobic

process of denitrification, while only the flooded zones will be a source of methane gas. If

methane is greater than CO_2 production, the overall GWP of the wetland will increase since

methane has a higher GWP than CO_2. However, due to the facultative nature of microbes

residing in the shallow land zones, denitrification will still occur in shallow land zones after a

short period of flooding if nitrate is not limiting. To test these hypotheses, we monitored GHG

fluxes monthly for one year and sampled sediments and surface water for chemical analyses

seasonally. To determine microbial (bacterial and archaeal) community structure in sediments,

we carried out 16s rRNA V4 amplicon sequencing. We also determined potential denitrification

rates across the wetland using a short-term incubation experiment. By combining GHG rates

with potential denitrification rates, sediment and water chemistry parameters along with

microbial community composition, we examined the potential benefits and disservices occurring

within this CW.

2. Materials and methods

2.1 *Study site*

The study location is a recently constructed (2015) stormwater wetland located on East

Carolina University's main campus in Greenville, NC (N 35°35'22.4", W 77°22'13.4"). The 0.13

acre CW receives runoff from 5.7 acre parking lot in a highly urbanized area of the city. The

temporary ponding depth (30.5 cm) of the wetland is equal to 0.51 cm of runoff within the

impervious drainage area (Fig. 1). The total surface area is divided among main zones (% surface

area): shallow land (35%), shallow water (35%), deep pools (10%), forebay (10%), and outlet

pool (10%) (NCDEQ, 2019).The dominant plants within saturated areas of the wetland are

Typha latifolia L., *Pontederia cordata, and* L., and *Iris versicolor* L. The shallow land area along

the water's edge is dominated by *Carex lupulina* Muhl. ex Willd.

Figure 1. Field site and sampling design. Left: Real picture of wetland. Center: Google Earth

image of parking lot and adjacent stormwater CW. Right: Google Earth image of wetland with

white circles representing GHG and sediment sampling locations.

2.2 Greenhouse gas monitoring and flux calculations

We used a static chamber method to measure greenhouse gas (GHG) fluxes. At the CW,

we established 13 GHG sampling plots along two main transects that follow the central channel

within the wetland and flows from inlet to outlet (Fig. 1). Of those 13 plots, five chambers were

placed in shallow water and deep pools (flooded zones), and 10 chambers were placed in shallow

land zones along the outside edge of the main channel. Each sampling plot in shallow land zones

(n=10) was fitted with a GHG chamber bottom installed 6 cm belowground. During sampling

events, chamber tops (height = 30 cm and diameter = 15 cm) were attached to an aboveground

groove, and the groove was filled with water to create a gas tight seal. GHG chamber tops in

flooded zones were fitted with a Styrofoam float allowing 2-3 cm of the chamber top below the

surface water.

We collected GHG samples monthly for one year. Sampling events occurred at least 48

hours after a rain event of > 0.51 cm. Gas samples were collected at 0, 12, 24, and 36 minutes

using 20 mL syringes. Each gas sample was divided between two 3.7 mL exetainers with dual

septa and stored upside at room temperature until analyzed. We measure GHG concentrations of

using a Shimazdu 2014 Gas Chromatograph (Shimadzu Scientific Instruments, Columbia,

Maryland) fitted with an electron capture detector to detect N_2O and flame ionization detector to

measure CH_4 and CO_2. Monthly GHG fluxes were calculated using the linear change in

concentration and the ideal gas law (Millar et al. 2018). Seasonal estimates were calculated by

averaging monthly flux measurements: spring = March-May, summer = June-August,

fall=September-November and winter=December-February. In addition, the equivalent CO_2 flux

(CO$_2$e) for methane was calculated by multiplying CH$_4$ flux by 28, which is equal to the increase

in warming potential in methane versus carbon dioxide.

2.4 *Soil collection and chemical properties*

We characterize soil physicochemical properties seasonally (May, July, October,

December). Each CW composite soil sample represented three cores (3.1 cm diameter, 12 cm

depth) that were adjacent to each GHG sampling chamber. We homogenized each composite

sample by passing the sample through a 4 mm sieve, and then processed subsamples for soil

gravimetric moisture, pH, extractable ammonium and nitrate, total carbon, and total nitrogen.

Briefly, we combined 5 g of field moist soil and 45 mL of 2M potassium chloride and shook

samples for 1 hour before gravity filtration to collect soil extracts. Ammonium and nitrate

concentrations were colorimetrically measured using a SmartChem 200 auto analyzer (Unity

Scientific Milford, Massachusetts, USA) at the East Carolina University Environmental Research

Laboratory. Using a 20-30 g subsample, we measured field-moist and oven-dry (dried 24 hours

at 105°C) soil weights to determine moisture content by measuring the mass of water divided by

the mass of oven-dried soil. Using a mortar and pestle, we coarsely ground a subsample of soil

and combined 10 g of soil with 10 mL of Nanopure® water to measure soil pH. On another

subsample, we finely ground the soil for elemental carbon and nitrogen analysis using an

elemental analyzer (2400 CHNS Analyzer; Perkin Elmer; Waltham, Massachusetts, USA) at the

Environmental and Agricultural Testing Service laboratory (Department of Crop and Soil

Sciences at NC State). Finally, we measured organic matter content based on the loss on ignition

method by ashing 20 g oven-dried soil at 550 °C (Hoogsteen et al., 2015) for 2 hours, this

duration may underestimate organic carbon content (Heiri et al., 2001). In addition, a subsample

of field moist soil was stored at 4 °C until we performed denitrification potential assays. Another

subsample of soil stored at -80 °C until DNA extraction for bacterial amplicon sequencing.

2.3 *Seasonal denitrification potential*

200   Within 72 hours of soil collection, we measured potential denitrification rates using the

acetylene block denitrification enzyme assay (Schaller et al., 2004; Tiedje et al., 1989; Wall et

al., 2005). We weighed triplicate subsamples of 25 g and transferred samples into 125 mL

Wheaton bottles fitted with a phenolic cap and butyl septa. In each bottle, we added 75 mL of 1

mM potassium chloride (KNO_3) to provide an abundant nitrogen supply and 1.3 mL of

chloramphenicol (100 mg mL^{-1}) to inhibit new denitrification enzyme production. Bottles were

sealed with septa-centered caps, shook, and purged with helium for 5 minutes. We removed 15

mL of helium from the headspace and replaced with 15 mL of pure acetylene gas. At each hour

for 3 hours, we collected a 10 mL gas sample (time point T0, T1, T2, T3) from each bottle and

stored gas sample in a 3.7 mL exetainer. We shook each bottle to equilibrate N_2O in aqueous and

sediment phases before each sample collection. We added 10 mL of 10% acetylene mixture to

the headspace after each sample collection. N_2O fluxes were calculated as described for GHG

fluxes.

2.5 *Microbial Community Analysis*

We extracted DNA from soils using the Qiagen DNeasy Powerlyzer PowerSoil Kit.

Genomic DNA was amplified using barcoded primers 515FB/806R primer set, originally

developed by the Earth Microbiome Project to target the V4-V5 region of the bacterial 16S

subunit of the ribosomal RNA gene (Apprill et al., 2015; Caporaso et al., 2010; Parada et al.,

2016). For each sample, three 50 µL PCR libraries were prepared by combining 38.35 µL

molecular grade water, 5 µL Amplitaq Gold 360 10x buffer, 2.4 µL $MgCl_2$ (25 mM), 1 µL

dNTPs (40mM total, 10mM each), 0.25 µL Amplitaq Gold 360 polymerase, 1 µL 515 forward

barcoded primer (10 µM), 1 µL 806 reverse primer (10 µM), and 1 µL DNA template (10 ng µL^{-}

^{1}). Thermocycler conditions for PCR reactions were as follows: initial denaturation (94 °C, 3

minutes); 30 cycles of 94°C for 45 seconds, 50 °C for 30 seconds, 72 °C for 90 seconds; final

elongation (72 °C, 10 minutes). The triplicate 50 µL PCR libraries were combined and then

cleaned using the AMPure XP magnetic bead protocol (Axygen, Union City, California, USA).

Cleaned PCR products were quantified using QuantIT dsDNA broad range assay (Thermo

Scientific, Waltham, Massachusetts, USA) and diluted to a concentration of 10 ng µL^{-1}. We

combined barcoded PCR samples (5 ng µL^{-1}) in equimolar concentration and sequenced the

pooled library using the Illumina MiSeq platform paired end read approach (Illumina Reagent

Kit v2, 500 reaction kit) at the Indiana University Center for Genomics and Bioinformatics

Sequencing Facility.

Sequences were processed using a standard mothur pipeline (v1.40.1)(Kozich et al.,

2013; Schloss et al., 2009). We assembled contigs from the paired end reads, quality trimmed

using a moving average quality score (minimum quality score 35), aligned sequences to the Silva

Database (Quast et al. 2013; SSURef v123), and removed chimeric sequences using the

VSEARCH algorithm (Rognes et al., 2016). We created operational taxonomic units (OTUs) by

first splitting sequences based on taxonomic class and then binning into OTUs based on 97%

sequence similarity. The Silva database was used to assign taxonomy to each OTU.

2.6 *Statistical Analysis*

All statistical analyses were completed in the R statistical environment (RStudio

v1.1.383, Rv3.4.0) (R Core Team, 2019). We constructed linear mixed effects models with

sampling plot as a random effect using the *lme4* R package (Bates et al., 2015) to determine the

importance of hydrology, season, and distance from the inlet on GHG fluxes and denitrification

potential fluxes within the constructed wetland. We then used corrected Akaike information

criteria (AICc) model comparisons, which adjust for small sample size, against the null model to

determine which fixed effects (hydrology and season) had the most explanatory power (Gorsky

et al., 2019; Hurvich and Tsai, 1993). Estimated marginal means with Tukey post-hoc

adjustment in the R package *emmeans* (Lenth, 2017) was used to determine significance

($p < 0.05$) in gas flux rates between treatment groups and the *multcomp* R package (Hothorn et al.,

2020) was used to assign Tukey post-hoc groups. Lastly, we used the *MuMIn* R package to

determine the proportion of variance explained by fixed effects (marginal) and the complete

model (conditional) (Barton, 2019; Gorsky et al., 2019). The CH_4, CO_2, and N_2O fluxes were

transformed to the log of the cube root and denitrification potential rates were log transformed to

meet normality assumptions of the statistical models.

Principal Coordinates of Analysis (PCoA) based on Bray-Curtis dissimilarity was used to

visualize patterns of bacterial diversity among treatments and sampling dates. Permuted analysis

of variance (PERMANOVA) was run using the *adonis* function in the *vegan* package (Oksanen,

2015) to test clustering significance. We performed a Dufrene-Legendre indicator species

analysis using the *labdsv* R package (Roberts, 2016) to identify specific microbial community

members that represented each hydrologic treatment. Next, we examined the relationship

between individual GHG fluxes microbial community Bray-Curtis dissimilarity matrix using

distance based partial least square regression in the *dbstats* R package (Boj et al., 2017). Finally,

we used Mantel R statistic function in the *vegan* R package (Oksanen, 2015) to examine the

relationship between all soil properties (moisture, pH, organic matter, ammonium and nitrate

concentrations, total C, total N, and C:N) microbial composition.

3. Results

3.1 Sediments and water properties

Over the one year study period in 2017, there were 268 days without precipitation.

Within flooded plots, water depth ranged from 8-28 cm in shallow water areas and 25-53 cm in

deep pools. Temperatures varied seasonally with July having the highest average ($\pm$ SD)

temperature (26.5 ± 1.8°C) and December having the lowest average temperature (4.9 ± 1.3°C)

(Table 1). Sediments in the wetland were acidic (4.6 ± 0.3 pH) ranging from pH 4.08 to 5.28.

Ammonium and nitrate concentrations were generally lower in sediments (0.089 ± 0.066 mg

NH_4^+-N g^{-1} dry mass and 0.023 ± 0.008 mg NO_3^--N g^{-1} dry mass) than in the water column (2.57

$\pm$ 19.6 mg NH_4^+ L^{-1} and 0.328 ± 0.803 mg NO_3^--N g^{-1} dry mass) throughout the wetland,

especially during winter months (9.97 ± 39.4 mg NH_4^+ L^{-1} and 0.769 ± 1.27 mg NO_3^- L^{-1}).

Sediment C:N ratios along with other sediment parameters (total C, total N, and C:N) and water

column phosphate (PO_4^{3-}) concentrations were similar across space (hydrologic zone) and time

(season) within the wetland (Table 1).

3.2 Greenhouse gas flux rates

3.2.1 Methane

Across the wetland, CH_4 fluxes differed by hydrology and season (Fig. 2A). Average ($\pm$

SD) CH_4 fluxes across all seasons were lowest in shallow land plots (2.32 ± 9.33 mg CH_4-C m^{-2}

hr^{-1}) and highest in flooded plots (80.80 ± 118.31 mg CH_4-C m^{-2} hr^{-1}). Within flooded plots, the

highest mean CH_4 fluxes (160.24 ± 156.28 mg CH_4-C m^{-2} hr^{-1}) were detected during summer

months, and the lowest CH_4 fluxes were detected in winter and spring months (45.02 ± 88.74 and

46.98 ± 59.40 mg CH_4-C m^{-2} hr^{-1}, respectively). Within shallow land plots, the highest mean CH_4

fluxes (6.38 ± 16.44 mg CH_4-C m^{-2} hr^{-1}) were detected during spring months and the lowest CH_4

fluxes were detected in winter (0.09 ± 0.42 mg CH_4-C m^{-2} hr^{-1}). Based on the mixed effects

relative to the null model, the model with hydrology and seasonal interaction explained the most

variation in CH_4 fluxes. This model had the lowest AICc score (313.70) and carried 100% of the

AICc weight (Table S1). In this model the interaction of hydrology and season and random

effects accounted for 73% variation in CH_4 fluxes, while the interaction alone explained 63% of

that variation (Table S1).

3.2.2 Carbon dioxide

Carbon dioxide fluxes varied seasonally but were generally highest in flooded plots

compared to shallow land plots (Fig. 2B). Across all seasons CO_2 fluxes were lowest in shallow

land plots (3.95 ± 43.96 mg CO_2-C m^{-2} hr^{-1}) and highest in flooded plots (73.52 ± 100.24 mg

CO_2-C m^{-2} hr^{-1}). We measured similar CO_2 flux rates (20.82 ± 46.36, 14.37 ± 44.68, and $18.88 \pm$

24.23 mg CO_2-C m^{-2} hr^{-1}) during spring, fall, and winter seasons, while CO_2 fluxes ($63.13 \pm$

133.21 mg CO_2-C m^{-2} hr^{-1}) were highest during the summer. According to the mixed effects

models relative to the null model, the model with season only explained the most variation in

CO_2 fluxes. This model had the lowest AICc score (45.54) and carried 93% of the AICc weight

(Table S1). The fixed effect of season explained 26% of the variation and the full model

explained 50% of the variation in CO_2 fluxes (Table S1).

Table 1. Summary of mean (range) of soil and water properties measured at the constructed wetland. Annual wetland values are

averages of all samples for the entire sampling period. Hydrology values are averages of samples collected in May (spring), July

(summer), October (fall), and December (winter) for each hydrology type. Seasonal sediment values are averages from of all plots

based on samples collected in May (spring), July (summer), October (fall), and December (winter). Seasonal water values are averages

of monthly samples from flooded plots where spring represents months May-June, summer represents months July-September, fall

represents months October-November, and winter represents months December-February. (Abbreviations: DM = dry mass. MDL =

below detection limit).

	Whole	Hydrology		Season			
	Wetland	Flooded	Shallow land	Spring	Summer	Fall	Winter
Soil							
temperature (°C)	18.6 (3.0-30.0)	-	-	19.6 (16.8-23.0)	26.5 (25.0-30.0)	21.3 (20.0-24.0)	4.9 (3.0-7.0)
moisture (%)	33.3 (18.3-51.9)	36.2 (23.2-51.9)	31.5 (18.3-46.2)	32.1 (26.6-42.9)	35.3 (27.0-48.1)	35.6 (26.6-51.9)	29.3 (18.3-44.1)
pH	4.6 (4.1-5.3)	4.7 (4.2-5.3)	4.6 (4.1-5.2)	4.7 (4.4-5.2)	4.6 (4.2-5.1)	4.5 (4.1-5.2)	4.7 (4.2-5.3)
organic matter (%)	6.20 (3.40-10.17)	6.54 (3.41-10.17)	6.01 (3.56-8.42)	6.32 (5.12-8.83)	6.26 (4.13-10.09)	6.68 (3.75-10.17)	5.39 (3.41-8.39)
mg NH_4^+-N g^{-1} DM	0.089 (0.023-0.457)	0.129 (0.049-0.457)	0.066 (0.023-0.148)	0.085 (0.023-0.148)	0.089 (0.036-0.204)	0.098 (0.028-0.457)	0.082 (0.040-0.201)
mg NO_3^--N g^{-1} DM	0.023 (0.012-0.050)	0.027 (0.14-0.050)	0.021 (0.012-0.038)	0.020 (0.016-0.027)	0.025 (0.017-0.038)	0.023 (0.014-0.034)	0.026 (0.012-0.050)
Total C (%)	1.66 (0.27-3.61)	1.78 (0.27-3.61)	1.60 (0.57-2.81)	1.57 (0.62-2.74)	1.78 (0.27-3.61)	1.80 (0.48-2.81)	1.46 (0.33-3.49)
Total N (%)	0.10 (0.02-0.20)	0.11 (0.02-0.20)	0.10 (0.04-0.16)	0.10 (0.04-0.17)	0.11 (0.02-0.19)	0.11 (0.03-0.16)	0.09 (0.03-0.20)
C:N (wt: wt)	15.78 (11.00-19.00)	15.88 (11.00-19.00)	15.73 (13.88-18.44)	16.0 (15.0-18.4)	16.2 (13.5-19.0)	15.9 (13.9-17.6)	14.9 (11.0-17.5)
Water							
mg NH_4^+-N L^{-1}	-	-	-	0.101 (MDL-0.474)	.206 (0.019-0.923)	0.263 (0.068-1.065)	9.972 (MDL-197.441)
mg NO_3^--N L^{-1}	-	-	-	0.173 (MDL-1.118)	0.007 (MDL-0.020)	0.169 (MDL-0.413)	0.769 (MDL-4.055)
Total Dissolved mg PO_4^{3-} L^{-1}	-	-	-	0.035 (MDL-0.087)	0.088 (0.27-0.286)	0.072 (0.025-0.176)	0.075 (MDL-0.245)

3.2.3 Nitrous oxide

Nitrous oxide fluxes were near

zero in all plots across all seasons Fig.

2C). Specifically, N_2O fluxes were

highest in shallow land plots (-0.006 ±

0.117 mg N_2O-N m^{-2} hr^{-1}) and lowest

in flooded plots (-0.083 ± 0.243 mg

N_2O-N m^{-2} hr^{-1}). Average N_2O flux

rates across seasons ranged from -0.05

± 0.20 to -0.02 ± 0.15 mg N_2O-N m^{-2}

hr^{-1}. Unlike CH_4 and CO_2 fluxes,

Figure 2. Boxplots of seasonal methane (A), carbon dioxide (B), and nitrous oxide (C) fluxes for flooded (blue circles) and shallow land (green triangles) plots. Spring = Mar-May, Summer = Jun-Aug, Fall=Sep-Nov, Winter=Dec-Feb. Letters indicate significantly different groups based on estimated marginal means with Tukey post-hoc assessment at p<0.05 using 95% confidence levels (Table S2).

hydrology and season did not influence N_2O fluxes. In this case, the null model had the lowest

AICc score (264.19) and the greatest AICc weight (69%) (Table S1). However, the

parameterized models with hydrology or season had similar AICc scores (267.17, 267.88) but

lower AICc weights (0.15, 0.11) when compared to the null model. Comparison of parameters

hydrology and season using estimated marginal means at 95% confidence level suggest season

has the greatest impact on N_2O fluxes (Table S2).

3.3 Global warming potential

In terms of GWP, flooded plots have a two-fold increase in magnitude of total warming

potential than shallow land plots especially during warmer months (2418.63, 58.46 CO_2

equivalents (mg m^{-2} hr^{-1}), respectively) (Fig. 3). This is primarily due to increases in CH_4

concentrations, which make up 99% of the CO_2 equivalents in both flooded and shallow land

plots (Fig. 3). Over one year, flooded areas within the wetland had the potential to produce about

7161.27 kg CO_2 equivalents, while shallow land areas produced about 93.20 kg CO_2 equivalents.

Figure 3. Monthly estimated global warming potentials of methane and carbon dioxide evaluated as carbon dioxide equivalents measured over a year at a constructed wetland. Proportion of grey for each bar represents mg CH_4-C m^{-2} hr^{-1} converted to CO_2 equivalents and proportion of black for each bar represents mg CO_2-C m^{-2} hr^{-1}. Average mg CH_4-C m^{-2} hr^{-1} was multiplied by 28, which is the estimated increased radiative force of methane when compared to carbon dioxide.

3.4 Denitrification potential

When nitrate limitation is

relieved and anoxic conditions are

present, sediments from flooded and

shallow land zones have similar

denitrification potential (Fig. 4). In

flooded and shallow land plots across

all seasons, potential denitrification

rates were 24.45 ± 20.18 and $20.29 \pm$

15.14 ng N_2O-N hr^{-1} g^{-1} dry mass,

respectively. However, denitrification

potential varied by season; we measured

higher rates in spring and summer

$(34.12 \pm 21.09$ and 25.16 ± 13.51 ng

N_2O N hr^{-1} g^{-1} dry mass) than in fall and

winter $(10.55 \pm 6.17$ and 16.45 ± 13.72

N_2O ng N hr^{-1} g^{-1} dry mass) months.

Based on the mixed effects models

Figure 4. Boxplots of potential denitrification rates according to hydrology and season based on denitrification enzyme assay using the acetylene block method. Spring = Mar-May, Summer = Jun-Aug, Fall=Sep-Nov, Winter=Dec-Feb. Different letters indicate significantly different groups by Tukey adjusted estimated marginal means p<0.05. Blue circles indicate flooded plots and green triangles shallow land.

relative to the null model, the model with season only explained the most variation in potential

denitrification rates (Table S3). This model had the lowest AICc score (480.01) and carried 78%

of the AICc weight (Table S3). The fixed effect of season explained 18% of the variation;

however, the null model explained the majority of variation (42%) (Table S3). We selected the

season only model even though the model with the fixed effects of hydrology and season had a

similar AICc (482.61) and AICc weight (21%), but there was little improvement in marginal and

conditional R^2 values (Table S3).

3.5 Microbial community analysis

Hydrology, and to a lesser extent season, influenced microbial community composition.

Illumina amplicon sequencing of the 16S rRNA V4 region resulted in 454,709 reads, where each

sample (n=22) contained between 13,235 and 33,678 reads before removing singletons and

doubletons. Microbial community composition in flooded plots were distinct from shallow land

plots (PERMANOVA, R^2= 0.0886, p=0.007; Fig. 5). Indicator species analysis identified one

Operational Taxonomic Unit (OTU; microbial taxon defined at 97% sequence similarly), in the

family Syntrophaceae, that was unique to flooded plots and identified eight OTUs that

represented the shallow land plots. The shallow land indicator taxa included OTUs from two

unclassified bacteria, two Betaproteobacteria, one Acidobacteria Gp6, and one *Geobacter* (Table

S5). Further examination of taxa with >0.6% relative abundance of the total community suggests

that flooded zones are dominated by taxa putatively involved in methanogenic degradation of

hydrocarbons (e.g., Methanomicrobia, Methylocystis, and Syntrophaceae), while shallow land

zones have fewer methanogens in comparison to flooded zones (Fig. 6). In addition, shallow land

zones are enriched with taxa putatively involved in nitrogen cycle transformations (e.g.,

Bradyrhizobium, Nitrososphaera, Rhizobiales) (Fig. 6).

Figure 5. Ordination plot based on Principal Coordinates Analysis depicting sediment
microbial community composition. Each point represents the centroid and range across
season and sampling location. Symbols represent sampling period (circle = spring, square
= fall). Spring samples were collected in May and fall samples were collected in October.
Colors represent sampling plots (blue = flooded, green = shallow land).

3.6 Analyses of relationship between GHGs, sediment chemistry, and microbial community composition

Distance-based partial least square regression analysis suggests that microbial community composition explained 82.4%, 78.6%, and 79.1%, of variation in CH_4, CO_2, and N_2O, respectively, by components 1 and 2. Sediment chemistry components 1 and 2 best explained variation in N_2O production (51.4%) but less so for CH_4 or CO_2 production (21.0%, 16.6%). Mantel

Figure 6. Heat map of bacterial and archaeal taxa observed at >0.6% relative abundance. Color gradient (cool to warm, green to red) represents microbial relative abundance ~0.6 to 1.5%.

correlation analyses revealed a positive relationship between patterns in microbial community composition and soil properties (r = 0.52, p = 0.003).

4. Discussion

After examining microbial processes occurring at permanently and periodically inundated locations within a stormwater CW, results revealed that modifications to wetland design could enhance water quality (via denitrification) and reduce GHG production. Other studies comparing different types of CWs indicate that free water surface (FWS) wetlands, like this study site, are known to produce copious amounts of GHGs compared to vertical or horizontal subsurface flow CWs (Mander et al., 2014; McPhillips and Walter, 2015). While vertical or subsurface CWs may reduce GHG production, this type of installation is more costly that FWS wetlands and may not always be practical. We recognize that this study focuses on a single CW; however, our results demonstrate the need for more comprehensive studies exploring CW design features that can be modified to enhance beneficial microbial ecosystem services (i.e., complete denitrification) while reducing ecosystem disservices (i.e., GHG emissions).

At our focal CW, CH_4 was the dominant GHG produced in flooded plots across all seasons. Hydrologic conditions accounted for the most variation in CH_4 fluxes and microbial community composition. Since high NO_3^- concentrations suppress methanogenesis (Kim et al., 2015), it is likely the combination of anoxic conditions (McPhillips and Walter, 2015), low NO_3^- concentrations in sediments and water, and availability of organic carbon in flooded plots, which provided optimal conditions for CH_4 production (Rahman et al., 2019). However, season and microbial community composition were stronger predicators of CO_2 and N_2O rates within the wetland. In other studies, temperature and substrate availability strongly determined rates of organic decomposition and denitrification (Davidson and Janssens, 2006; Knowles, 1982; Moinet et al., 2018). Seasonal differences in temperature and vegetation status (i.e., organic

carbon inputs) within this study influenced CO_2 and N_2O rates to a greater degree than

hydrology.

In order to simultaneously compare the biogenic GHGs within the CW, we quantified

GWP associated with flooded and shallow land zones. In terms of GWP, CH_4 from flooded plots

was the primary carbon source within the wetland. While both CO_2 and CH_4 production was

greatest in flooded zones compared to shallow land zones, CH_4 production greatly exceeded CO_2

production leading to high GWP within the wetland. Compared to surface flow wastewater

treatment wetlands receiving municipal or agricultural runoff, CO_2 emissions at our study

wetland in flooded plots were similar but approximately 95% lower in shallow land plots

(Jahangir et al., 2016; Mander et al., 2014). In contrast, CH_4 emissions at our study wetland had

similar CH_4 rates in shallow land plots when compared to wastewater treatment wetlands, but

approximately 94% greater emissions in flooded plots compared to wastewater treatment

wetlands (Jahangir et al., 2016; Mander et al., 2014). In terms of GWP, flooded plots in our

study wetland have a GWP that is 98% greater than in wastewater treatment wetlands. Overall,

compared to wastewater treatment wetlands, flooded plots within our study wetland represent a

considerable source of GWP primarily due to CH_4 emissions.

Further, GHG emissions could be partially explained by microbial community structure.

This is due to hydrologic conditions in flooded zones providing optimal habitat for obligate

anaerobes that participate in methanogenesis. Since methanogenesis is suppressed by even low

oxygen concentrations (Fetzer et al., 1993; Liu et al., 2008), relative methanogen signal was not

detectable at shallow land areas which can fluctuate between oxic and anoxic conditions.

Additionally, analysis of microbial community structure suggests that hydrocarbon degradation,

a potential microbial ecosystem benefit, may be occurring in flooded plots. This finding

demonstrates the value of examining the microbial community composition when evaluating

beneficial wetland ecosystem functions.

From this study we highlight three CW features for consideration to reduce ecosystem

function disservices: (i) surface area of hydrologic zones, (ii) incoming nutrient and pollutant

loads, and (iii) the anticipated number of rain days. Results from this study and others indicate

that areas that fluctuate between oxic and anoxic conditions, which occur in CW shallow land

areas in this study, are sites of reduced CH_4 production. This is due to the inhibition of

methanogenesis and a greater abundance of methanotrophs which consume methane

(Chowdhury and Dick, 2013; Lew and Glińska-Lewczuk, 2018). Additionally, even in CWs with

low nitrate availability, shallow land areas could potentially support enhanced denitrification due

to coupling of nitrification, which occurs during oxic periods and steady organic carbon inputs

from vegetation (Mander et al., 2014; Rahman et al., 2019). Flooded zones increase water

storage capacity and provide habitat for predator species that control nuisance species; therefore,

elimination of flooded zones may not be desirable. However, an increase in shallow land area

and a reduction in surface area of flooded zones could reduce GHG emissions without sacrificing

benefits of flooded zones.

Secondly, substrate availability can directly impact microbial process rates. Compared to

the neighboring Tar River (0.52 mg NO_3-N L^{-1}), seasonal NO_3^- concentrations within the surface

water of the CW were much lower (0.007-0.173 mg NO_3-N L^{-1}) except during winter (0.769 mg

NO_3-N L^{-1}) (Humphrey et al., 2019). Considering that NO_3^- availability limits denitrification,

ecosystems that receive runoff from agriculture, livestock, or are adjacent to a high density of

septic systems are areas of high denitrification potential and low methane production (Naylor et

al., 2018). Recent studies show that iron (II) (Fe^{2+}) additions can increase N removal from

vertical and horizontal subsurface CWs (Song et al., 2016; Zhang et al., 2019). In the case of

existing wetlands, further investigation into amendments as an alternative to increasing

denitrification potential is warranted. Additionally, depending on the upstream runoff source,

pollutants such as hydrocarbons may be better targets for remediation within intentionally

designed CWs.

Lastly, we suggest that the frequency of rain days or inter-event duration is taken into

account during SCM planning (Andersen et al., 2017). Our results represent ambient conditions,

at least 48 hours after a storm event, which represent more than half the year for this CW. Within

this particular wetland, reducing the permanently flooded surface area could reduce GHGs while

still providing N removal benefits. In areas where rain events are less frequent (i.e., greater inter-

event durations) and more intense, SCMs that are not permanently flooded, such as dry

infiltration basins or dry detention basins, could provide relief for intense runoff events with

reduced GHG production (McPhillips and Walter, 2015; Morse et al., 2017). Therefore, we

suggest frequency of rain events as another factor to consider when planning SCM design.

5. Conclusion

This study demonstrates the importance of considering microbial controls on

biogeochemical processes within SCMs in order to reduce tradeoffs between water quality and

greenhouse gas production. Context matters when designing CWs in urban watersheds. In the

case of this urban stormwater CW, which is located in a parking lot of a highly urbanized area,

we suggest that reducing flooded areas and increasing shallow land area would reduce GHG

emissions while still providing nutrient removal benefit. However, our analysis of the microbial

community also indicates that potential hydrocarbon decomposition could be supported in

flooded zones and warrants a deeper investigation of this potential service. This study also

highlights the need for a comprehensive study of CW design features to better understand which

design features and when those features provide the most ecosystem benefits. By examining both

microbial process rates and the microbial community composition, we can approach SCM design

and implementation in a holistic way that accounts for ecosystem services and disservices.

Declaration of Competing Interest

None.

Acknowledgements

We thank B. Hinckley, A. Garcìa, and L. Armstrong for assistance with field research

and sample processing. We also thank Sound Rivers for installation of the constructed wetland

and ECU Grounds personnel and manager J. Gill for site maintenance. This work was supported

by East Carolina University and the North Carolina Sea Grant/Water Resources Research

Institute Graduate Student Research Fellowship and the National Science Foundation Graduate

Research Fellowship to RBB. All code and data used in this study can be found in a public

GitHub repository (https://github.com/PeraltaLab/ConstructedStormwaterWetland) and the

NCBI SRA (BioProject PRJNA613095).

Appendix A. Supplementary data

Supplementary material related to this article can be found at: ###

References

Altshuler, I., Hamel, J., Turney, S., Magnuson, E., Lévesque, R., Greer, C.W., Whyte, L.G.,

2019. Species interactions and distinct microbial communities in high Arctic permafrost

affected cryosols are associated with the CH_4 and CO_2 gas fluxes. Environ. Microbiol. 21,

3711–3727. https://doi.org/10.1111/1462-2920.14715

Andersen, J.S., Lerer, S.M., Backhaus, A., Jensen, M.B., Danielsen Sørup, H.J., 2017.

Characteristic rain events: A methodology for improving the amenity value of stormwater

control measures. Sustain. 9. https://doi.org/10.3390/su9101793

Apprill, A., Mcnally, S., Parsons, R., Weber, L., 2015. Minor revision to V4 region SSU rRNA

806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat. Microb.

Ecol. 75, 129–137. https://doi.org/10.3354/ame01753

Barton, K., 2019. Package "MuMIn."

Bates, D., Machler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using

lme4. J. Stat. Softw. 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Bell, C.D., Tague, C.L., McMillan, S.K., 2019. Modeling Runoff and Nitrogen Loads From a

Watershed at Different Levels of Impervious Surface Coverage and Connectivity to Storm

Water Control Measures. Water Resour. Res. 55, 2690–2707.

https://doi.org/10.1029/2018WR023006

Boj, E., Caballe, A., Delicado, P., Fortiana, J., 2017. Package ' dbstats .'

Brindha, R.K., Vasudevan, N., 2018. Methane oxidation capacity of methanotrophs isolated from

different soil ecosystems. Int. J. Environ. Sci. Technol. 15, 1931–1940.

https://doi.org/10.1007/s13762-017-1546-1

Burgin, A.J., Yang, W.H., Hamilton, S.K., Silver, W.L., 2011. Beyond carbon and nitrogen: how

the microbial energy economy couples elemental cycles in diverse ecosystems. Front. Ecol.

Environ. 9, 44–52. https://doi.org/10.1890/090227

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-lyons, D., Lozupone, C.A., Turnbaugh, P.J.,

Fierer, N., Knight, R., 2010. <Global patterns of 16S rRNA diversity at a depth of millions

of sequences per sample.pdf>. Proc. Natl. Acad. Sci. U. S. A. 108, 4516–4522.

https://doi.org/10.1073/pnas.1000080107/-

/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1000080107

Chowdhury, T.R., Dick, R.P., 2013. Ecology of aerobic methanotrophs in controlling methane

fluxes from wetlands. Appl. Soil Ecol. 65, 8–22.

https://doi.org/10.1016/j.apsoil.2012.12.014

Conrad, R., 2007. Microbial Ecology of Methanogens and Methanotrophs. Adv. Agron. 96, 1–

63. https://doi.org/10.1016/S0065-2113(07)96005-8

Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and

feedbacks to climate change. Nature 440, 165–173. https://doi.org/10.1038/nature04514

Demuzere, M., Orru, K., Heidrich, O., Olazabal, E., Geneletti, D., Orru, H., Bhave, A.G., Mittal,

N., Feliu, E., Faehnle, M., 2014. Mitigating and adapting to climate change: Multi-

functional and multi-scale assessment of green urban infrastructure. J. Environ. Manage.

146, 107–115. https://doi.org/10.1016/j.jenvman.2014.07.025

Fetzer, S., Bak, F., Conrad, R., 1993. Sensitivity of methanogenic bacteria from paddy soil to

oxygen and desiccation. FEMS Microbiol. Ecol. 12, 107–115.

https://doi.org/10.1111/j.1574-6941.1993.tb00022.x

Fetzer, S., Conrad, R., 1993. Effect of redox potential on methanogenesis by Methanosarcina

barkeri. Arch. Microbiol. 160, 108–113. https://doi.org/10.1007/BF00288711

Gorsky, A.L., Racanelli, G.A., Belvin, A.C., Chambers, R.M., 2019. Greenhouse gas flux from

stormwater ponds in southeastern Virginia (USA). Anthropocene 28.

https://doi.org/10.1016/j.ancene.2019.100218

Heiri, O., Lotter, A., Lemcke, G., 2001. Loss on ignition as a method for estimating organic and

carbonate content in sediments. J. Paleolimnol. 25, 101–110.

https://doi.org/10.1017/CBO9781107415324.004

Hoogsteen, M.J.J., Lantinga, E.A., Bakker, E.J., Groot, J.C.J., Tittonell, P.A., 2015. Estimating

soil organic carbon through loss on ignition: Effects of ignition conditions and structural

water loss. Eur. J. Soil Sci. 66, 320–328. https://doi.org/10.1111/ejss.12224

Hothorn, T., Bretz, F., Westfall, P., Heiberger, R.M., Schuetzenmeister, A., Scheibe, S., 2020.

Package ' multcomp .'

Humphrey, C., Iverson, G., Skibiel, C., Sanderford, C., Blackmon, J., 2019. Geochemistry of

Flood Waters from the Tar River, North Carolina Associated with Hurricane Matthew.

Resources 8, 48. https://doi.org/10.3390/resources8010048

Hurvich, C., Tsai, C., 1993. A CORRECTED AKAIKE INFORMATIONON CRITERION FOR

VECTOR AUTOREGRESSIVE MODEL SELECTION. J. Time Ser. Anal. 14, 271–279.

IPCC, 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and

III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Ipcc.

IPCC - Intergovernmental Panel on Climate Change, 2014. Climate Change 2014 Synthesis

Report (Unedited Version).

Jahangir, M.M.R., Richards, K.G., Healy, M.G., Gill, L., Müller, C., Johnston, P., Fenton, O.,

2016. Carbon and nitrogen dynamics and greenhouse gas emissions in constructed wetlands

treating wastewater: A review. Hydrol. Earth Syst. Sci. 20, 109–123.

https://doi.org/10.5194/hess-20-109-2016

Kim, S.Y., Veraart, A.J., Meima-Franke, M., Bodelier, P.L.E., 2015. Combined effects of

carbon, nitrogen and phosphorus on CH4 production and denitrification in wetland

sediments. Geoderma 259–260, 354–361. https://doi.org/10.1016/j.geoderma.2015.03.015

Knowles, R., 1982. Denitrification. Microbiol. Rev. 46, 43–70.

Koch, B.J., Febria, C.M., Gevrey, M., Wainger, L.A., Palmer, M.A., 2014. Nitrogen Removal by

Stormwater Management Structures: A Data Synthesis. J. Am. Water Resour. Assoc. 50,

1594–1607. https://doi.org/10.1111/jawr.12223

Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D., 2013. Development of

a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence

data on the miseq illumina sequencing platform. Appl. Environ. Microbiol. 79, 5112–5120.

https://doi.org/10.1128/AEM.01043-13

Le Mer, J., Roger, P., 2001. Production, oxidation, emission and consumption of methane by

soils: A review. Eur. J. Soil Biol. 37, 25–50. https://doi.org/10.1016/S1164-5563(01)01067-

6

Lee, C.G., Fletcher, T.D., Sun, G., 2009. Nitrogen removal in constructed wetland systems. Eng.

Life Sci. 9, 11–22. https://doi.org/10.1002/elsc.200800049

Lenth, R., 2017. Package ' lsmeans .' https://doi.org/10.1080/00031305.1980.10483031>.NOTE

Lew, S., Glińska-Lewczuk, K., 2018. Environmental controls on the abundance of

methanotrophs and methanogens in peat bog lakes. Sci. Total Environ. 645, 1201–1211.

https://doi.org/10.1016/j.scitotenv.2018.07.141

Line, D.E., White, N.M., 2007. Effects of Development on Runoff and Pollutant Export. Water

Environ. Res. 79, 185–190. https://doi.org/10.2175/106143006x111736

Liu, C. Te, Miyaki, T., Aono, T., Oyaizu, H., 2008. Evaluation of methanogenic strains and their

ability to endure aeration and water stress. Curr. Microbiol. 56, 214–218.

https://doi.org/10.1007/s00284-007-9059-7

Mander, Ü., Dotro, G., Ebie, Y., Towprayoon, S., Chiemchaisri, C., Nogueira, S.F., Jamsranjav,

B., Kasak, K., Truu, J., Tournebize, J., Mitsch, W.J., 2014. Greenhouse gas emission in

constructed wetlands for wastewater treatment: A review. Ecol. Eng. 66, 19–35.

https://doi.org/10.1016/j.ecoleng.2013.12.006

Marton, J.M., Creed, I.F., Lewis, D.B., Lane, C.R., Basu, N.B., Cohen, M.J., Craft, C.B., 2015.

Geographically isolated wetlands are important biogeochemical reactors on the landscape.

Bioscience 65, 408–418. https://doi.org/10.1093/biosci/biv009

McPhillips, L., Walter, M.T., 2015. Hydrologic conditions drive denitrification and greenhouse

gas emissions in stormwater detention basins. Ecol. Eng. 85, 67–75.

https://doi.org/10.1016/j.ecoleng.2015.10.018

Mitsch, W.J., Bernal, B., Nahlik, A.M., Mander, Ü., Zhang, L., Anderson, C.J., Jørgensen, S.E.,

638   Brix, H., 2013. Wetlands, carbon, and climate change. Landsc. Ecol. 28, 583–597.

639   https://doi.org/10.1007/s10980-012-9758-8

640   Moinet, G.Y.K., Hunt, J.E., Kirschbaum, M.U.F., Morcom, C.P., Midwood, A.J., Millard, P.,

641   2018. The temperature sensitivity of soil organic matter decomposition is constrained by

642   microbial access to substrates. Soil Biol. Biochem. 116, 333–339.

643   https://doi.org/10.1016/j.soilbio.2017.10.031

644   Morse, N.R., McPhillips, L.E., Shapleigh, J.P., Walter, M.T., 2017. The Role of Denitrification

645   in Stormwater Detention Basin Treatment of Nitrogen. Environ. Sci. Technol. 51, 7928–

646   7935. https://doi.org/10.1021/acs.est.7b01813

647   Naylor, E., Humphrey, C., Kelley, T., Easter, L., Iverson, G., 2018. Evaluation of nitrate

648   concentrations and potential sources of nitrate in private water supply wells in North

649   Carolina. J. Environ. Health 80, 16–23.

650   NCDEQ, 2019. Stormwater Design Manual: C-4 Stormwater Wetland [WWW Document].

651   North Carolina Dep. Environ. Qual. URL https://deq.nc.gov/sw-bmp-manual

652   NCDEQ, 2018. Stormwater Control Measure Credit Document [WWW Document]. North

653   Carolina Dep. Environ. Qual. URL https://deq.nc.gov/sw-bmp-manual

654   O'Driscoll, M., Clinton, S., Jefferson, A., Manda, A., McMillan, S., 2010. Urbanization effects

655   on watershed hydrology and in-stream processes in the southern United States. Water

656   (Switzerland) 2, 605–648. https://doi.org/10.3390/w2030605

Oksanen, J., 2015. Vegan□: ecological diversity 1, 1–12. https://doi.org/10.1029/2006JF000545

Paerl, H.W., Hall, N.S., Peierls, B.L., Rossignol, K.L., 2014. Evolving Paradigms and

Challenges in Estuarine and Coastal Eutrophication Dynamics in a Culturally and

Climatically Stressed World. Estuaries and Coasts 37, 243–258.

https://doi.org/10.1007/s12237-014-9773-x

Parada, A.E., Needham, D.M., Fuhrman, J.A., 2016. Every base matters: Assessing small subunit

rRNA primers for marine microbiomes with mock communities, time series and global field

samples. Environ. Microbiol. 18, 1403–1414. https://doi.org/10.1111/1462-2920.13023

Payne, E.G.I., Fletcher, T.D., Cook, P.L.M., Deletic, A., Hatt, B.E., 2014. Processes and drivers

of nitrogen removal in stormwater biofiltration. Crit. Rev. Environ. Sci. Technol. 44, 796–

846. https://doi.org/10.1080/10643389.2012.741310

Peralta, A.L., Ludmer, S., Kent, A.D., 2013. Hydrologic history influences microbial community

composition and nitrogen cycling under experimental drying/wetting treatments. Soil Biol.

Biochem. 66, 29–37. https://doi.org/10.1016/j.soilbio.2013.06.019

Peralta, A.L., Ludmer, S., Matthews, J.W., Kent, A.D., 2014. Bacterial community response to

changes in soil redox potential along a moisture gradient in restored wetlands. Ecol. Eng.

73, 246–253. https://doi.org/10.1016/j.ecoleng.2014.09.047

R Core Team, 2019. R: A language and environment for statistical computing. R Found. Stat.

Comput. Vienna, Austria.

Rabalais, N.N., Turner, R.E., Díaz, R.J., Justić, D., 2009. Global change and eutrophication of

coastal waters. ICES J. Mar. Sci. 66, 1528–1537. https://doi.org/10.1093/icesjms/fsp047

Rahman, M.M., Roberts, K.L., Warry, F., Grace, M.R., Cook, P.L.M., 2019. Factors controlling

dissimilatory nitrate reduction processes in constructed stormwater urban wetlands.

Biogeochemistry 142, 375–393. https://doi.org/10.1007/s10533-019-00541-0

Reisinger, A.J., Groffman, P.M., Rosi-Marshall, E.J., 2016. Nitrogen-cycling process rates

across urban ecosystems. FEMS Microbiol. Ecol. 92, fiw198.

https://doi.org/10.1093/femsec/fiw198

Rey-Sanchez, C., Bohrer, G., Slater, J., Li, Y.F., Grau-Andrés, R., Hao, Y., Rich, V.I., Davies,

G.M., 2019. The ratio of methanogens to methanotrophs and water-level dynamics drive

methane transfer velocity in a temperate kettle-hole peat bog. Biogeosciences 16, 3207–

3231. https://doi.org/10.5194/bg-16-3207-2019

Roberts, D., 2016. Package ' labdsv .'

Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open

source tool for metagenomics. PeerJ 4, e2584. https://doi.org/10.7717/peerj.2584

Schaller, J.L., Royer, T. V., David, M.B., Tank, J.L., 2004. Denitrification associated with plants

and sediments in an agricultural stream. J. North Am. Benthol. Soc. 23, 667–676.

https://doi.org/10.1899/0887-3593(2004)023<0667:DAWPAS>2.0.CO;2

Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski,

R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B., Thallinger, G.G., Van

Horn, D.J., Weber, C.F., 2009. Introducing mothur: Open-source, platform-independent,

community-supported software for describing and comparing microbial communities. Appl.

Environ. Microbiol. 75, 7537–7541. https://doi.org/10.1128/AEM.01541-09

Smith, M.S., Tiedje, J.M., 1979. Phases of denitrification following oxygen depletion in soil.

Soil Biol. Biochem. 11, 261–267. https://doi.org/10.1016/0038-0717(79)90071-3

Song, X., Wang, S., Wang, Y., Zhao, Z., Yan, D., 2016. Addition of Fe2+ increase nitrate

removal in vertical subsurface flow constructed wetlands. Ecol. Eng. 91, 487–494.

https://doi.org/10.1016/j.ecoleng.2016.03.013

Tiedje, J.M., 1988. Ecology of denitrification and dissimilatory nitrate reduction to ammonium.

Environ. Microbiol. Anaerobes 179–244.

Tiedje, J.M., Simkins, S., Groffman, P.M., 1989. Perspectives on measurement of denitrification

in the field including recommended protocols for acetylene based methods. Plant Soil 115,

261–284. https://doi.org/10.1007/BF02202594

Wall, L.G., Tank, J.L., Royer, T. V., Bernot, M.J., 2005. Spatial and temporal variability in

sediment denitrification within an agriculturally influenced reservoir. Biogeochemistry 76,

85–111. https://doi.org/10.1007/s10533-005-2199-6

Wen, X., Unger, V., Jurasinski, G., Koebsch, F., Horn, F., Rehder, G., Sachs, T., Zak, D.,

Lischeid, G., Knorr, K.H., Böttcher, M.E., Winkel, M., Bodelier, P.L.E., Liebner, S., 2018.

Predominance of methanogens over methanotrophs in rewetted fens characterized by high

methane emissions. Biogeosciences 15, 6519–6536. https://doi.org/10.5194/bg-15-6519-

2018

Zhang, Y., Liu, X., Fu, C., Li, X., Yan, B., Shi, T., 2019. Effect of Fe2+ addition on chemical

oxygen demand and nitrogen removal in horizontal subsurface flow constructed wetlands.

Chemosphere 220, 259–265. https://doi.org/10.1016/j.chemosphere.2018.12.144