# Multiple freeze-thaw cycles lead to a loss of consistency in poly(A)-enriched RNA sequencing

Benjamin P. Kellman[1,2,*], Hratch M. Baghdassarian[1,2,*], Tiziano Pramparo[3], Isaac Shamie[1,2],

Vahid Gazestani[1,3], Arjana Begzati[4], Shengzhong Li[1,6], Srinivasa Nalabolu[3], Sarah Murray[5],

Linda Lopez[3], Karen Pierce[3], Eric Courchesne[3], Nathan E. Lewis[1,6,7]

[1] Department of Pediatrics, University of California, San Diego

[2] Bioinformatics and Systems Biology Program, University of California San Diego

[3] Autism Center of Excellence, Department of Neuroscience, University of California San Diego

[4] Department of Medicine, University of California San Diego

[5] Department of Pathology, University of California San Diego

[6] Department of Bioengineering, University of California San Diego

[7] Novo Nordisk Foundation Center for Biosustainability, University of California San Diego

[*] These authors contributed equally

Correspondence to: Nathan E. Lewis, nlewisres@ucsd.edu

Keywords: RNA-Seq; quality control; freeze-thaw; sample preparation; differential expression

## **Abstract**

RNA-Seq is ubiquitous, but depending on the study, sub-optimal sample handling may be

required, resulting in repeated freeze-thaw cycles. However, little is known about how each cycle

impacts downstream analyses, due to a lack of study and known limitations in common RNA

quality metrics, e.g., RIN, at quantifying RNA degradation following repeated freeze-thaws.

Here we quantify the impact of repeated freeze-thaw on the reliability of downstream RNA-Seq

analysis. To do so, we developed a method to estimate the relative noise between technical

replicates independently of RIN. Using this approach we inferred the effect of both RIN and the

number of freeze-thaw cycles on sample noise. We find that RIN is unable to fully account for

the change in sample noise due to freeze-thaw cycles. Additionally, freeze-thaw is detrimental to

sample quality and differential expression (DE) reproducibility, approaching zero after three

cycles for poly(A)-enriched samples, wherein the inherent 3' bias in read coverage is more

exacerbated by freeze-thaw cycles, while ribosome-depleted samples are less affected by freeze-thaws. The use of poly(A)-enrichment for RNA sequencing is pervasive in library preparation of frozen tissue, and thus, it is important during experimental design and data analysis to consider the impact of repeated freeze-thaw cycles on reproducibility.

## Introduction

RNA sequencing (RNA-Seq) is a ubiquitous technology, used to answer a wide range of biological questions. Methods for aligning, quantifying, normalizing and analyzing expression data are available through popular packages such as Tophat, STAR, cufflinks, SVA, RUV, Combat, DESeq2, edgeR, Kallisto, Salmon, BWA-MEM, and many others[1–11]. Each method aims to accommodate and mitigate the unique challenges presented by RNA-Seq data. Some approaches attempt to account for characterized variability in RNA-Seq measurements due to factors such as sequencing depth, gene length, and transcripts' physical characteristics (e.g., GC content). Others account for "unwanted variance" due to technical, batch, or experimental variation. Yet, the influence of sample processing, such as tissue lysis and processing time[12–14], is not sufficiently characterized such that it can be explicitly controlled. It is important to adequately characterize noise introduced to RNA-Seq measurements by sample processing steps to optimize sample quality, account for transcript degradation, and improve the accuracy and reproducibility of sequencing.

Transcript degradation continues after sample acquisition and affects data quality. Sample storage conditions (e.g. temperature and the use of stabilizing reagents) affect sample quality via RNA degradation[14,15]. Yet, varying sources of degradation can impact RNA-Seq in different manners[16]. Degradation introduces variability in signal and can be impacted by sample handling.

Non-uniformity in degradation across genes and samples causes inaccurate normalization and transcript quantification[17]. Poly(A)-enrichment methods are commonly used to separate mRNA from other highly abundant RNA molecules (e.g., rRNA, tRNA, snoRNAs, etc.), but variable degradation directly impacts read counts by causing non-uniform transcript coverage[18]. Of particular interest, freeze-thaw can induce 20% degradation of spike-in standards per cycle, a factor that may be generalizable to mRNA transcripts[19]. Freeze-thaw cycles increase RNA degradation by disrupting lysosomes which store RNases, freeing the enzymes to promiscuously catalyze nuclease activity[20]. Furthermore, partially defrosted crystals create uneven cleaving pressure on mRNA strands[21,22]. Despite these observations, the extent to which freeze-thaw negatively impacts count and differential expression in RNA-Seq analyses has not been comprehensively characterized.

Standard sample quality control often relies on RNA integrity number (RIN), which quantifies the 28S to 18S rRNA ratio[23]. RIN-based quality control approaches rely on a heuristic threshold to assess sufficient quality[24,25]. RIN-based metrics have known confounders such as transcript level, and thus have been called into question as an appropriate quality metric[26]. For example, RIN failed to indicate a decrease in sample quality in lung cancer tissue samples that underwent five freeze-thaw cycles[27] and, in statistical analyses, failed to correct for the effects of degradation[28]. Despite this, many studies rely on RIN to correct for and assess sample quality confounders[16,29,30]. This is especially problematic in the case of transcript degradation because RIN scores are based on entire samples, while degradation effects can be transcript-specific[17,31,32]. Furthermore, existing studies on degradation are not simply generalizable to freeze-thaw, which has distinct and independent effects on sample quality and must be fully explored as such[16,33].

Here, we tested the susceptibility of poly(A)-enriched RNA-Seq results after multiple freeze-

thaw cycles. We assessed sample quality independently of RIN by simulating read count

variability to capture the noise between technical replicates. We found that each additional

freeze-thaw cycle increased the random counts between technical replicates by approximately

4%. Subsequently, differential expression reproducibility approached zero after three freeze-

thaw cycles. These effects are not captured by RIN. We find that these effects are reflected in

increasing 3' bias in read coverage when combining poly(A)-extraction with freeze-thaw, a

phenomena that appears to be generalizable to publically available datasets.

**Results**

**3' bias in read coverage of public datasets is associated to poly(A)-extraction**

**and freeze-thaw**

To examine the prevalence of bias from repeated freeze-thaw cycles, we search for the presence

of RNA-seq distortion in publically available datasets. Specifically, we analyze the gene-

coverage distribution in samples prepared with either poly(A)-extraction or ribosomal depletion.

Since freeze-thaw enhances transcript degradation and poly(A)-enriched samples select mRNA

by hybridization to the poly(A)-tail, we expect increased read coverage on the 3' end of

transcripts--3' bias--when these two factors are combined. To test this expectation, we compared

gene body coverage from the 5' to 3' end between poly(A)-enrichment and ribosomal depletion

prepared samples with and without freezing. Specifically, we examined the median coverage

percentile, the percentile-normalized nucleotide at which median cumulative coverage for a

given sample is achieved (**Fig. 1a**).

We analyzed the gene-coverage distribution in three studies: (1) RNA extraction in previously

frozen solid and liquid tissues, (2) RNA extracted immediately after lysis of cultured cells

without freezing, and (3) RNA extraction in previously frozen tissue from important public tissue

resources.

**Figure 1**: **3' Bias is Exacerbated in Frozen, Poly(A)-extracted Samples Across Multiple Studies:** *(a)*

*Demonstration for determining median coverage percentile (red vertical line). When coverage is*

*unbiased, reads (yellow) are distributed throughout the entire body of the transcript (green). In the*

*absence of read bias and observing coverage as a function of the nucleotide percentile, we see that*

*cumulative coverage along the transcript reaches 50% half-way through the gene body, at the 50th*

*percentile nucleotide. In contrast, given a 3' read bias, there is a shift in the distribution of reads and*

*cumulative coverage reaches 50% at, for example, the 60th percentile nucleotide. This results in a*

*rightward shift in median coverage percentile towards the 3' end of the transcript. In the middle row,*

*gene coverage (y-axis) at the $i^{th}$ nucleotide percentile from 5' to 3' (x-axis) is displayed for samples that*

*were extracted using either poly(A)-enrichment or ribosomal depletion. Gene body coverage distributions*

*were calculated for (b) for tissue samples that underwent an unspecified number of freeze-thaw cycles*

*and (c) cell-culture samples samples that underwent no freeze-thaw cycles. (d) Comparison of 5' to 3'*

*bias ratio (y-axis) of samples from the TCGA and UNC tissue repositories (x-axis) between extraction*

*methods (two-sample t-test).  Quantifying human RNA samples listed in GEO from 2008-2018, and*

*stratifying by those annotated as "frozen", we observe (d) the number of samples prepared with poly(A)-*

*extraction or ribosomal depletion (x-axis) (gray), (f) the proportion of samples extracted using either*

*method, and (g) the change in the number of samples over time.*

In the first study[34], a comparison of the performance of poly(A) and ribosomal depletion in liquid

and solid frozen tissue, we see a significant (one-sided Wilcoxon test, p = 0.01591) shift in the

median gene coverage percentile towards the 3' end in the poly(A)-extracted samples (**Fig. 1b**).

Using generalized linear regression, we found that the median coverage percentile of poly(A)-

extracted samples was 3.88% higher (Wald p = 0.011) than comparable ribosomal depletion

extracted samples. In the second study[35], cells were not frozen before extraction and there was a

small and insignificant difference in the 3' bias associated with library preparation (one-sided

Wilcoxon test, p = 0.13, **Fig. 1c**). While the first two studies extract RNA from tissue and cell-

culture respectively--tissue extraction is typically lower quality, they are internally controlled

and therefore comparable. Finally, the third study[36] examines the impact of RNA extraction in

frozen tissue from the UNC and TCGA tumor tissue repositories. We found a significant (two-

sample t-test, $p<1e16$) decrease in the 5'-to-3' coverage ratio in poly(A)-extracted samples

compared to ribosomal depletion (**Fig. 1d**), indicating an increase in 3' bias. 3' bias was

consistent across RNA extracted from tissue frozen at either repository.

Next, we explored how widespread the impact of these observations may be by quantifying the

prevalence of poly(A)-extraction from frozen tissue by examining metadata in the Gene

Expression Omnibus (GEO). With GEOmetadb [37], we queried all human RNA samples between

2008 and 2018 using either poly(A)-extraction or ribosomal depletion. There are tens to hundred

of thousands of samples annotated as "frozen" for both total and poly(A)-extraction methods

(**Fig. 1e**).  We note that this is an order of magnitude less than the total query, a value potentially

diminished by the complexity of the metadata. In samples annotated as "frozen", the frequency

of poly(A) extraction increases from less than 10% to over 25% **(Fig. 1f)** suggesting that the

problematic combination is prevalent and apparently preferred. Finally, stratifying this trend over

time, we see that poly(A)-extraction, as well as the relative proportion of poly(A)-extracted

frozen samples, is increasing in popularity relative to total RNA extraction, where usage has

remained fairly consistent (**Fig. 1g**).Taken together, these results indicate a potential, widespread

distortion in RNA-seq associated with a deleterious interaction between poly(A)-extraction and

freeze-thaw. To explore this potential more formally, the remainder of our analyses focus on a

specific experiment to address this question. Specifically we subjected whole-blood extracted

leukocyte samples--with technical replicates--from autistic or typically developing toddlers to a

varying number of freeze-thaw cycles, which we record along with other sample quality metrics

such as RIN.

## An Additional Freeze-Thaw Cycle Increases Random Read Counts 1.4-Fold

To address the scarcity in quantification of loss in sample quality due to freeze-thaw, we

compare changes in sample quality between technical replicates. We first note that neither RIN

nor TIN capture significant (one-sided Wilcoxon test) decreases in sample quality due to

increased freeze-thaw (**Fig. S1**). Given previous indications that these metrics may not

sufficiently address transcript degradation, we instead measure the introduction of noise—the

randomness in read counts between technical replicates—to samples by freeze-thaw. We

simulated this randomness to reflect the dissimilarity between technical replicates. Since noise

does not rely on RIN, we could compare freeze-thaw and RIN-effects independently.

**Figure 2**: **Higher Noise in Samples with More Freeze-Thaw Cycles.** *From left to right, noise—*

*the randomness in read counts between technical replicates—is estimated using Euclidean distance,*

*RMSE, Pearson correlation,and Spearman correlation. (a) Box plots of noise for samples that underwent*

*either one or two freeze-thaws. (b) A generalized linear model was used to determine the expected noise*

*at one freeze-thaw ($N_0$, pink) and the expected change in noise with each additional freeze-thaw ($\Delta N$,*

*green). All estimates are significant ($p \leq 0.05$).*

Median noise increased 1.4-fold from one to two freeze-thaw cycles  (one-sided Mann-Whitney

U test, $p \leq 0.007$) on average across all measures (**Fig. 2a**). Noise between technical replicates

when samples have only undergone one freeze-thaw was estimated to be 9.11-10.15% (Wald

test, $p \leq 5.77e-7$). The expected increase in noise per additional freeze-thaw cycle was estimated

to be 3.6-4.1 percentage points (Wald test, $p \leq 8.12e-3$) (**Fig. 2b**).

## RIN Does Not Predict Additional Noise After One Freeze-Thaw Cycle

To follow up on our observations that RIN does not sufficiently capture changes in sample

quality due to freeze-thaw, we asked whether RIN can reflect the differences in sample quality as

measured by noise.

**Figure 3**: **Discrepancy in the Relationship Between Noise and RIN due to additional Freeze-Thaw.** *Examining the relationship between noise, calculated by Euclidean distance, RMSE, pearson, and spearman correlation, for samples that underwent either one (blue) or two (red) freeze-thaw cycles. (a) Scatter plots comparing noise (y-axis) to RIN (x-axis). The solid lines show a linear regression fit and the shaded regions is the 95% confidence interval for this fit. (b) The expected change in noise due to a one point increase in RIN ($\Delta N$, y-axis) estimated by a generalized linear model. Significant estimates ($p \leq 0.05$) are marked by a circle and insignificant estimates are marked by a cross.*

When only considering samples that underwent one freeze-thaw, each unit increase in RIN decreases noise by 3.24-3.38 percentage points for all metrics (Wald test, $p \leq 6.3e-3$) (Fig. **3a-b**). Yet, when only accounting for samples that underwent two freeze-thaw cycles, noise does not significantly change as RIN increases. Taken together, these results indicate that while RIN can be a good measure of noise for samples that underwent one freeze-thaw, it does not capture the loss in sample quality induced by two freeze-thaw cycles.

## Differential Expression Similarity Increases 10.3% in High Quality Samples

Next, we investigated how the introduction of technical variation, noise, impacts downstream RNA-Seq analysis, specifically, differential expression (DE) analysis. As such, we assessed the reproducibility of DE results on combinations of samples with varying sample quality. We compared DE results between subsets of various sizes (4 - 14 samples). We measure reproducibility using similarity or discordance, based on correlation and dispersion, respectively. Higher similarity and lower discordance each represent higher reproducibility. We use these measures to assess differences that arise between subsets consisting of high quality (low freeze-thaw or high RIN) and low quality (high freeze-thaw or low RIN) samples.

We held two expectations regarding the effect of sample quality on DE reproducibility in the

context of similarity: 1) the reproducibility between subsets with high quality samples should be

higher than those with low quality samples, and 2) both subset size and sample quality should

interact to increase the reproducibility of DE analysis; the increase in stability is reflected by a

higher rate of increase in reproducibility with respect to subset size.

As expected, similarity increases with subset size. This is reflected in the upward shift in the

similarity distribution with increasing subset size (**Fig. S10**) and the estimated 0.02 (Wald test, p

= 2.2e-5)  increase in similarity per additional sample (**Fig. 4a)**; thus, expected similarity would

increase by 0.20 in a subset with 14 samples over a subset with 4 samples. Regression results for

each model predicting similarity are reported in **Supplementary Table 4**.

**Figure 4**: **Freeze-Thaw and RIN Both Demonstrate Higher Similarity with Increased**

**Quality.** *Top panels summarize generalized linear models used to quantify the change in similarity per*

*unit increase in (a) sample size, number of freeze-thaws and RIN combined, (b) only the number of*

*freeze-thaws, and (c) only RIN. $S_0$ represents the intercept estimate and sample size, freeze-thaw, and RIN*

*represent coefficient estimates. All estimates are significant. Bottom panels demonstrate fold-change in*

*median similarity of high quality subsets with respect to low quality subsets at each subset size. The*

*region shaded in blue (fold-change > 1) indicates instances where the median similarity for high quality*

*is larger than that of low quality. The region shaded in red (fold-change < 1) indicates instances where*

*the median similarity for low quality is larger than that of high quality. Average (d) freeze-thaw or (e)*

*RIN are used to place subset pairs into high or low quality sample bins. Significance (one-sided Mann-*

*Whitney U test) of comparisons in similarity distributions between high and low quality subset pairs are*

*displayed above each subset size.*

We tested our first expectation by placing subset pairs into high and low sample quality bins,

defined by either RIN or freeze-thaw, for each subset size and comparing their similarity values.

Regardless of sample quality, DE similarity increases with subset size. Yet, for nearly all subset

sizes, higher quality bins have significantly (one-sided Mann-Whitney U test, $p \leq 2.8e\text{-}17$)

higher similarity than low quality bins (**Fig. 4e-f**). Across subset sizes, we observed an average

1.13-fold and 1.06-fold increase in similarity from low to high quality samples for freeze-thaw

and RIN, respectively.

Similarity significantly (Wald test, $p \leq 9.2e\text{-}3$) decreases with the number of freeze-thaws and

increases with RIN when accounting for the effects of sample size (**Fig. 4a-c**), validating our

second expectation. Similarity decreases by 0.077 per additional freeze-thaw cycle (Wald test, p

= 8.77e-4). Given the estimated similarity of 0.23 for samples that have not undergone freeze-

thaw, this implies that DE reproducibility will approach zero after approximately three freeze-

thaw cycles (**Fig. 4b**). Even when accounting for subset size and the effects of RIN, the

estimated decrease in similarity from freeze-thaw is nearly the same--0.078 (Wald test, p =

8.77e-4); this further corroborated that RIN alone cannot capture the changes in sample quality

due to freeze-thaw. Taken together, these results indicate that higher sample quality increases DE

reproducibility as measured by similarity.

## Discordance Decreases ~5-fold In High Quality Samples

We further investigated the relationship between DE reproducibility and sample quality using an

effect size sensitive measure of discordance. Specifically, we explored how sample quality

affects the relationship between discordance and the mean-variance standardized effect at each

subset size. In this context, we expected 1) discordance at any given effect size to be lower in

high-quality subsets and 2) the rate of increase in discordance to be lower in high quality subsets

relative to low quality subsets.

Corresponding to the regression models used for this analysis, we label our expected discordance

when effect size is zero as $D_0$ and the change in discordance per unit increase in effect size as

$\Delta D$. We observed a significant (Wald test, $p \leq 9.45e\text{-}141$) decreasing trend in both $D_0$ and $\Delta D$

with increasing subset size (**Fig. S11, Supplementary Table 5)**.

As expected, independent of sample quality, $\Delta D$ demonstrates an overall decreasing trend with

respect to subset size for both RIN and freeze-thaw. Given freeze-thaw, at a subset size of 6,

there is a 1.1-fold decrease in the value of $\Delta D$ from low quality subsets to high quality subsets.

The disparity in $\Delta D$ between high and low sample quality ($\Delta m = \Delta D_{\text{Low Quality}} / \Delta D_{\text{High Quality}}$)

increases nearly monotonically through to the subset size of 14, at which point there is a 3.2-fold

decrease; the monotonicity is an indication of the stability of this relation between discordance

and sample quality. This causes notable differences in discordance values, even at low effect

sizes (**Fig. 5a**).

**Figure 5: Higher Sample Quality has Lower Discordance at Each Subset Size.** *GLM estimates*

*of discordance predicted from effect size and subset size at high and low quality subsets at each subset*

*size. High sample quality (blue) is compared to low sample quality (red). ΔD values represent the change*

*in discordance per unit increase in effect size. (a) The predicted discordance with respect to the mean-*

*variance standardized effect at a subset size of 14; sample quality is assessed by freeze-thaw. The*

*disparity (Δm) between the change in discordance per unit increase in effect size for high (ΔD$_H$) and low*

*(ΔD$_H$) quality subsets is also displayed. Summary of results for each subset size (x-axis) for sample*

*quality represented by either (b) freeze-thaw or (c) RIN. Significant estimates (Wald test, Benjamini-*

*Hochberg FDR correction, q ≤ 0.1) are marked by a circle and insignificant estimates are marked by a*

*cross. For freeze-thaw, Δm corresponding to panel A is also displayed.*

We estimate discordance with respect to effect size at each subset size and for subsets of either

high and low quality (**Fig. 5a**). Consistent with our expectations, $D_0$ and $\Delta D$ are consistently

lower for high quality subsets as compared to low quality subsets for both freeze-thaw and RIN

across all subset sizes (**Fig. 5b-c**). Nearly all estimates are significant after multiple test

correction (Wald test, Benjamini-Hochberg FDR correction, $q \le 0.07$), with the exception of

those for the smallest subset size for freeze-thaw.

Taken together, these results indicate that higher sample quality increases DE reproducibility as

measured by discordance.

## Additional freeze-thaw cycles show increased 3' bias in poly(A)-enriched but not ribosomal depletion samples

Finally, we asked whether repeated freeze-thaw cycles can induce a 3' bias, consistent with the

induction of random reads and the loss of DE reproducibility as well as our initial observation in

the public datasets.

Using the median coverage percentile, we found a shift in mRNA coverage towards the 3' end in

the poly(A)-enriched samples relative to ribosomal depletion (**Fig. S13a**). Specifically, the

median coverage percentile for poly(A)-enriched samples is significantly (one-sided Wilcoxon

test, $p < 2.2e-16$) larger than that of ribosome depletion (**Fig. S13b**). Samples prepared with

poly(A)-extraction have more 3' coverage bias compared to ribosomal depletion in both one

(one-sided Wilcoxon test, $p = 3.5e-15$ ) and two (one-sided Wilcoxon test, $p = 1.4e-4$) freeze-

thaw cycles (**Fig. S13d**). Altogether, this indicates an overall 3' bias of poly(A)-enriched

samples, even independently of freeze-thaw **(Fig. S13b**).

**Figure 6**: **Freeze-Thaw Cycles Exacerbate 3' Bias in poly(A)-enriched samples.** *(a) Gene coverage (y-*

*axis) at the $i^{th}$ nucleotide percentile (x-axis) for samples that underwent 1-5 freeze-thaws and were*

*extracted using either poly(A)-enrichment or ribosome depletion. Coverage is normalized to samples that*

*underwent one freeze-thaw. For each sample, coverage is averaged across all genes; samples are*

*aggregated using generalized additive model smoothing, with shaded regions representing 95%*

*confidence intervals. (b) Linear model fits comparing the change in median coverage percentile to the*

*number of freeze-thaw cycles for ribosome depletion (orange) or poly(A)-extraction (green).*

Crucially, this 3' bias is accentuated when samples are stratified by the number of freeze-thaw

cycles **(Fig. 6a).** We observe a significant increase (Wald test, p = 8.3e-4)  in median coverage

percentile due to the number of freeze-thaw cycles in poly(A)-enrichment. The increase was not

maintained when was isolated using  ribosome depletion (Wald test, p = 0.155) **(Fig. 6b,**

**Supplementary Table 7)**. For poly(A)-enriched samples, median coverage percentile increases

3.3 percentage points per root freeze-thaw cycle; the square-root of freeze-thaw was used to

stabilize variance. We further demonstrate a dependency of 3' bias on freeze-thaw cycles by

showing that median coverage percentile significantly increases with freeze-thaw in poly(A)-

enriched samples (Kruskal-Wallis test, p = 0.01). This 3' bias is particularly apparent after five

freeze-thaw cycles (one-sided Wilcoxon test, p = 2.4e-3). In contrast, we observe that median

coverage percentile does not significantly change across freeze-thaw cycle counts in ribosomal

depletion (Kruskal-Wallis test, p = 0.084) **(Fig. S13c)**.

Taken together, these analyses indicate that poly(A)-enrichment inherently introduces a 3' bias

in coverage as compared to ribosome depletion, and that this bias is exclusively exacerbated in

poly(A)-enriched samples due to freeze-thaw cycles. Thus, 3' bias may indicate the severity of

freeze-thaw induced signal degradation in poly(A)-extracted samples. If this 3' bias is the root

cause of freeze-thaw induced instability in absolute and differential RNA-seq quantification,

such instabilities may be subverted by substituting poly(A)-selection for ribosomal depletion

during library preparation; such instabilities may be an unnecessary and avoidable

inconvenience.

## Discussion

Despite the utility and ubiquity of RNA-Seq, many of the confounding elements associated with

the technology are still being characterized. In this work, we demonstrated how one such

confounder--sample freeze-thaw--impacts sample quality and downstream analyses. We

highlighted biases in publically available datasets, and observed an increased 3' bias when both

freeze-thaw and poly(A)-extraction are features of a sample.

To gain a comprehensive understanding of this effect, we first simulated technical replication to

measure the noise between technical replicates with different treatments. This allowed us to

examine the impact of freeze-thaw cycles and the ability of RIN to capture those impacts. Next,

we examined the impact of freeze-thaw cycles on the robustness and reproducibility of

differential expression analysis. We found freeze-thaw cycles were substantially detrimental to

the stability of gene expression analysis. By our estimates and at these subset sizes,

reproducibility of a differential expression signature approaches zero after three freeze-thaw

cycles (**Supplementary Table 4**). Finally, we demonstrated that poly(A)-enriched samples

demonstrate substantial 3' bias in read coverage with increased freeze-thaw cycles. Our results

have implications with regards to technical variation due to sample handling, the sensitivity of

differential gene expression analysis for frozen tissues and samples, and the utility of RIN.

Technical variation in RNA-Seq is substantial and can be attributed to a variety of factors,

including read coverage, mRNA sampling fraction, library preparation batch, GC content, and

sample handling[38,39]. As such, accounting for technical variation has been a major research area

of focus for the past decade[1,2,5,39,40]. Degradation in combination with poly(A)-enrichment is a

known source of variation in RNA-Seq. Yet, before technical variation can be accounted for, it

must be characterized. While studies have looked into the effect of degradation on RNA-Seq,

each mode of degradation impacts sample quality differently, and direct connections between

freeze-thaw and sample quality has mainly been assessed via RIN[16,41,42].

Our simulation of technical replicates helps delineate technical variation due to sample handling-

-specifically, freeze-thaw. Furthermore, the resulting noise provides an estimate for the number

of random read counts associated with a gene. For example, given an average 25 million reads

sequenced per sample, our approximate 4 percentage points difference in noise between one and

two freeze-thaw cycles gives an expected stochasticity in 1 million of those reads; approximating

the number of protein-coding genes in the human genome to be 20-25 thousand[43], we can expect

a difference of ~40-50  random counts per gene to exist between technical replicates due to a

freeze-thaw cycle (**Supplementary Methods, Fig. S15**). Thus, each freeze-thaw introduces a

non-negligible level of noise to the quantification of gene expression and differential expression

of such genes.

To check for the possibility that there is a signature which can help correct for freeze-thaw

distortion of RNA-Seq, we attempt to find a group of common DE genes across various DE

methods. We find no such signature (**Supplementary Results**). This is expected, given that a

major source of reduced sample quality due to freeze-thaw is mRNA degradation, which occurs

randomly for each transcript and sample. A possible path forward is to correct for sample

degradation. Several methods have been proposed for this. While some of these methods rely on

RIN or similar metrics (e.g. mRIN, TIN, etc.)[16,44], others have implemented statistical

frameworks which account for gene-specific biases. DegNorm, for example, accounts for the

gene-specific relative randomness in degradation in its correction approach[17]. Quality surrogate

variable analysis (qSVA) specifically improves differential expression by identifying transcript

features associated with RNA degradation for its correction[28]. Furthermore, there are recent

methods which only assay the 3' end of a transcript and therefore claim robustness in degraded

samples[45]. While these approaches could help account for noise introduced by RNA degradation

during repeated freeze-thaw cycles, they cannot necessarily remedy the associated loss of signal.

The effect of freeze-thaw and resultant degradation on RNA-Seq is particularly concerning when

considering differential gene expression analysis. It has been observed that RNA degradation can

induce the apparent differential expression in as many as 56% of genes[44]. To this end, we

quantified this loss of DE reproducibility by measuring similarity and discordance in the context

of sample quality. We found a decrease in reproducibility with both decreasing RIN and

increasing freeze-thaw. Interestingly, for most reproducibility assessments, we observed a

monotonic or near monotonic increase in disparity between low and high quality subsets with

respect to subset size. Similarity demonstrated a larger average magnitude of disparity for freeze-

thaw, whereas discordance demonstrated a larger average magnitude of disparity for RIN.

Based on our analysis, the utility of RIN in assessing quality when samples undergo freeze-thaw

is questionable. The non-uniformity in mRNA degradation[46–49] due to freeze-thaw sheds light on

these challenges, since RIN cannot quantify quality at the individual gene level[23]. This is

reflected in the fact that samples with RIN > 8 demonstrate degradation[32].  Furthermore, results

assessing the effect of freeze-thaw cycles on RIN are inconclusive. While some studies claim

RIN can be used to account for degradation effects in RNA-Seq[16], others suggest it does not

sufficiently capture the effects of degradation on sample quality[26,28]. When directly observing the

effect of freeze-thaw on RIN, studies have found a negligible effect[12]  or can only detect an

effect after numerous cycles[27,50].

As such, we re-examined the utility of RIN as a measure of sample quality in relation to our

noise estimation of random reads per sample[23]. We found that while noise increases with both

decreasing RIN and increasing freeze-thaw, RIN may be an insufficient indicator of quality for

samples that have undergone two or more freeze-thaws.  Given these results, RIN may not

always be a good metric to quantify the difference between technical replicates that have

undergone variable sample handling[17,26–28]. We validate noise by confirming that it does not

change with input RNA concentration, excepting outliers **(Fig. S12)**. Therefore, in the future,

noise could be a useful supplement to RIN when technical replicates are present.

The fact that our predicted decrease in similarity due to freeze-thaw does not change when

incorporating RIN into our model further indicates that RIN alone cannot capture the changes in

sample quality due to freeze-thaw. Despite this, RIN is a good indicator of sample quality, if not

specifically for freeze-thaw. This is reflected in the fact that RIN validates our expectations for

DE reproducibility analysis and the comparable range of noise, similarity, and discordance

values between freeze-thaw and RIN assessments.

Finally, to confirm our expectation that freeze-thaw decreases sample quality[18–22] and to further

characterize the underlying mechanism, we validated the presence of a 3' bias in coverage. This

builds on our and others' observations that a lower percentage of poly(A)-enriched transcripts

are covered[42]. We compared coverage to ribosome depleted RNA-Seq data, which does not use

3' hybridization to retain transcripts.  We find that poly(A)-enrichment does in fact introduce a

strong 3' bias in coverage as compared to ribosome depletion. This bias is further exacerbated

with additional freeze-thaw cycles in poly(A)-enriched but not ribosome depleted samples. This

implies that degradation due to freeze-thaw does not impact RNA-sequencing of ribosome

depleted samples to the extent that it does in poly(A)-enriched samples.  In light of  our

demonstrations that 3' bias is associated with a substantial increase in noise and a decrease in DE

reproducibility, these findings suggest that RNA-seq from samples that have both been poly(A)-

extracted and undergone freeze-thaw cycles likely has unknown, diminished stability. While not

all studies have technical replicates to estimate noise, the presence of exaggerated 3' bias when

poly(A)-extraction is combined with freeze-thaw can be a simple indicator of RNA-seq

distortion.

## Conclusion

Altogether, these results indicate that transcriptomics quality control steps cannot rely on RIN

alone for samples that have undergone poly(A)-enrichment and multiple freeze-thaws.

Furthermore, accounting for the effect of freeze-thaw on poly(A)-enriched RNA sequencing is

crucial. poly(A)-enrichment is prevalent for RNA-sequencing, and, in parallel, samples that

undergo multiple freeze-thaws are common in many protocols, especially rare tissues, e.g.,

postmortem neural tissue. Yet, there is no clear recommendation to avoid poly(A)-enrichment

following multiple freeze-thaws. Together, these results indicate that ribosomal depletion could

be a better alternative when freeze-thaw is necessary.

# Declarations

## Ethics Approval and Consent to Participate

In this study, we performed transcriptomics analyses of blood samples drawn from male toddlers

with the age range of 1–4 years. Research procedures were approved by the Institutional Review

Board of the University of California, San Diego. Parents of toddlers underwent informed

consent procedures with a psychologist or study coordinator at the time of their child's

enrollment. Additional details for the recruitment protocol are executed as described in Gazestani

et al.[51]

## Author contributions

B.P.K., N.E.L., T.P., S.M., and E.C. designed and planned the experiments. T.P., S.N., K.P.,

S.M. and L.L. collected the samples, managed diagnostics, conducted transcriptome assays and

managed the data.  B.P.K, H.M.B. and V.G. planned and conducted analyses. I.S. performed

RNA-seq QC. A.B. performed functional enrichment analyses. S.L. wrote the RNA-Seq

processing pipeline. B.P.K., H.M.B, and N.E.L wrote the manuscript. N.E.L. supervised the

project.

## Acknowledgements and Funding

440 This work was supported by NIMH R01-MH110558 (E.C., N.E.L., T.P., V.G., S.N., K.P.,L.L.),

441    NIDCD R01-DC016385 (E.C., T.P.), T32GM008806 (H.M.B.), R35 GM119850 (B.P.K., I.S.),

442    the Simons Foundation (E.C.), and generous funding from the Novo Nordisk Foundation through

443    Center for Biosustainability at the Technical University of Denmark (NNF10CC1016517 S.L.).

## Availability of Data and Materials

445    The datasets supporting the conclusions of this article are available in the NCBI Sequence Read

446    Archive repository (PRJNA627540,

447    https://www.ncbi.nlm.nih.gov/Traces/study/?acc=PRJNA627540&o=acc_s%3Aa). Associated

448    metadata is available in Supplementary Tables 1-2.

## Competing Interests

450    There are no competing interests in this study.

# Methods

## Sample Collection and Storage

Blood samples drawn from male toddlers with the age range of 1-4 years were usually taken at

the end of the clinical evaluation sessions. To monitor health status, the temperature of each

toddler was monitored using an ear digital thermometer immediately preceding the blood draw.

The blood draw was scheduled for a different day when the temperature was higher than 37 °C.

Moreover, blood draw was not taken if a toddler had some illness (for example, cold or flu), as

observed by us or stated by parents. We collected 4–6 ml blood into EDTA-coated tubes from

each toddler. Blood leukocytes were captured using LeukoLOCK filters (Ambion). After rinsing

the LeukoLOCK filters with PBS, the filters were flushed with RNAlater (Invitrogen) to stabilize

RNA within the intact leukocytes. After RNA stabilization, the LeukoLOCK filters were

immediately placed in a −20 °C freezer. Additional RNA standards were sourced from normal

human peripheral leukocytes pooled from 39 Asian individuals, ages 18 to 47 (Takara/ClonTech:

636592). The RNA standards underwent 1-5 simulated freeze-thaw cycles; 24hrs frozen at

−80 °C and 1hr defrosting on ice.

**RNA Extraction, Sequencing and Quantification**

For 47 samples (from 16 individuals), mRNA was extracted using polyA selection with the

TruSeq Stranded mRNA library preparation kit (Illumina). Ribosomal depletion was used to

prepare an additional 52 samples. Relevant metadata regarding polyA-enriched and ribosomal

depleted samples can be found in **Supplementary Table 1-2**. Ribosome depletion prepared

samples used the TruSeq Stranded Total RNA with RiboZero Gold library preparation kit

(Illumina). RNA Integrity Numbers (RIN) were measured using a NanoDrop ND-1000

(ThermoFisher). Poly-A selected samples were sequenced using 50-base pair single end

sequencing on a HiSeq4000 (Illumina) to a depth of 25M reads. The ribo-depletion prepared

libraries were sequenced using 100-base pair paired end sequencing on a HiSeq4000 (Illumina)

to a depth of 50M reads.

Fastq files for each sample underwent quality control using FastQC (v0.33). PolyA and adaptor-

trimming were conducted using Trimmomatic [52]. Reads were aligned to the gencode annotated

(v25) human reference genome (GRCh38) using STAR (v2.4.0) [7]. Alignments were processed to

sorted SAM files using SAMtools (v1.7) [53]. Finally, HTSeq using default settings (v0.6.1) was

used to quantify reads [53,54].

## Estimation of noise between technical replicates

To estimate the noise between technical replicates of the same individual blood samples, we simulate random loss and gain of reads **(Fig. S2)**. One technical replicate was chosen as the "reference" replicate, making the other technical replicate the "target" replicate. Specifically, we designated replicates that have undergone one freeze-thaw as the reference, and those that underwent two freeze-thaw cycles as the target. The dissimilarity between replicates is measured by one of four metrics (Euclidean distance, RMSE, Pearson correlation, and Spearman correlation). We iteratively add and remove random reads to the reference replicate until the dissimilarity between the simulated replicate and the reference replicate was equal to the dissimilarity between a target replicate and the reference replicate **(Fig. S3, Figure S2)**. We define the noise between the reference and target replicate as the fraction of reads added or removed per total reads in the reference replicate to achieve the aforementioned level of dissimilarity. We represent this as a percentage, e.g. 5% noise between a reference and target replicate can be interpreted as 5% randomness between their reads. For additional details on noise simulation, see Supplementary Methods.

## Measuring the Effect of Sample Quality on Noise

To measure the association between noise and sample quality metrics (number of freeze-thaw cycles, input RNA concentrations, and RNA integrity number), we used a generalized linear model (GLM). The significance of the model parameters is determined by the Wald test. All results are reported in **Supplementary Table 3**.

For each model, to mitigate the contribution of potential confounding variables, samples with input RNA concentrations in the top and bottom 5% ($|z| \geq 1.645$) were removed, decreasing the

504    total number of samples from 47 to 41. For noise prediction from concentration, samples with

505    more than one freeze-thaw were also excluded, decreasing the total number of samples to 35.

506    Noise prediction from RIN was run separately for samples that had undergone one freeze-thaw

507    and samples that had undergone two freeze-thaws.

## Differential Expression Analysis

We assess whether the observed sample qualities (freeze-thaw and RIN) have an impact on

differential expression (DE) reproducibility using a bootstrapping approach. DE was run on

random sample subsets of varying sizes **(Fig. S4)**. Before subsetting, we filtered our expression

matrix for genes with an average count $\leq$ 20 across all samples. This reduced the number of

genes from 10,028 to 4,520. The total number of samples considered was 46 when disregarding

samples that were industry standards, were not assigned to either an ASD or TD indication, or

did not have a recorded sample quality (RIN or freeze-thaw) value.

We generated subsets containing N = 4-14 samples. For each subset size N, we generated 2,000

unique subsets. Each subset had an equal number of TD or ASD samples. Additionally, only one

replicate from each blood sample could be included. These requirements limited our subset size

to a maximum of 14 samples.

DE between ASD and TD subjects was conducted using DESeq2[1]. **Fig. S10** summarizes DE

results for all subsets. To account for potential confounders, we used RUV[5]. Specifically, we use

a set of "in-silico empirical" negative control genes, including all but the top 5,000 differentially

expressed genes as described in section 2.4 of the documentation for RUVseq

(http://bioconductor.org/packages/release/bioc/vignettes/RUVSeq/inst/doc/RUVSeq.pdf). We

check that RUV produces consistent results with previous Autism leukocyte gene expression

signatures[51,55], see **Supplementary Results.**

**Similarity to Assess Differential Expression Reproducibility**

To assess DE reproducibility, we measure the similarity in log-fold-change (LFC) values

between DE runs. Similarity is calculated as the pairwise spearman correlation of LFC between

all subsets of the same size **(Fig. S6)**. Genes with a median base mean (the mean of counts of all

samples, normalizing for sequencing depth) or median LFC in the bottom $10^{th}$ percentile across

all subsets were excluded, filtering for low magnitude effects **(Fig. S5)**.

Average RIN and freeze-thaw were measured for all subset pairs. Resulting distributions for all

collected values from similarity analyses are displayed in **Fig. S7**.

Next, subsets of each size were split into two quantile bins for each quality metric separately.

High sample quality bins (low freeze-thaw or high RIN) were compared to low sample quality

bins.  High sample quality subsets were tested for higher similarity than low sample quality bins

using a one-sided Mann-Whitney U test.

Additionally, three generalized linear models (GLMs) were fit to quantify the contribution of

sample quality metrics to the change in similarity for DE results across subsets. We fit one model

to predict similarity from freeze-thaw and RIN, while also accounting for the improvement in

reproducibility due to increase in subset size (Similarity ~ Freeze-Thaw + RIN + Subset Size).

We also fit two models predicting similarity from freeze-thaw or RIN alone.

**Discordance to Assess Differential Expression Reproducibility**

We adapted a measure of concordance to measure discordance, or the lack of reproducibility,

between differential expression results[56]. Average RIN and freeze-thaw were calculated for each

subset **(Fig. S8-9)**. Subsets for each subset size were split into two quantile bins for either quality

metric (RIN and freeze-thaw). Genes with a median base mean across all subsets in the bottom

tenth percentile were excluded from the analysis **(Fig. S5)**.

We do not use the original concordance at the top (CAT) metric because we are not comparing

our results to a gold standard dataset. Instead, we use gene-wise LFC standard deviation across

subsets as a measure of discordance. Thus, the average LFC for each gene across DE runs is

analogous to the gold standard, and the dispersion from this average indicates a lack of

reproducibility. At each combination of subset size and quality bins, we calculate discordance

and compare it to the gene-wise median effect size **(Fig. S8)**. We measure effect size as the

mean-variance standardized effect[1]. This and two additional effect size metrics (Cohen's d and

absolute median LFC) we use are further described in **Fig. S9**. Results for all three effect size

metrics reflect similar trends and can be found in **Supplementary Tables 5-6.**

We used a GLM to predict discordance from effect size at each subset size. Additionally, in a

separate GLM, we account for the interaction between effect size and sample quality

(Discordance ~ Effect Size x Sample Quality) at each subset size. Here, sample quality is a

dummy variable, assuming a value of 0  for low quality and 1 for high quality.  We did not

include a term for subset size because regressions were fit within each subset size.

**Read Coverage Bias**

The distribution of read coverage over each gene body was measured using

*geneBody_coverage.py* from the *RSeQC* (v3.0.0) package [57]. We measure this coverage ranging

from the $0^{th}$ percentile (5' end) to the $100^{th}$ percentile (3' end) nucleotide. The i$^{th}$ percentile

nucleotide is calculated as $nucleotide_i / length_{gene}$. Coverage at the i$^{th}$ percentile nucleotide is

normalized across all genes within a sample.

For a given sample, the median coverage percentile is defined as the nucleotide percentile at

which median cumulative coverage is achieved; cumulative coverage is aggregated from the 5'

end to the 3' end. The larger the median coverage percentile value, the larger the 3' bias in

coverage.  We include 9 industry standards to our analysis--six of which had undergone five

freeze-thaw cycles and three of which had undergone one freeze-thaw cycle--to explore the

impact at higher freeze-thaw counts. We also include ribosomal depletion extracted samples as a

negative control.

We extended this read coverage bias to three additional datasets, all of which contained both

poly(A)-extracted and ribosomal depletion extracted samples. The first dataset (PRJNA427184)

contains samples for liquid and solid frozen tissue [34]. The second (PRJEB4197) contains

HEK293 samples that explicitly never underwent freeze-thaw cycles [35]. The third

(phs000676.v1.p1) contains frozen tissue samples from the UNC and TCG tumor tissue

repositories [36].  For the first two datasets, fastq files were aligned using STAR and gene body

coverage was calculated from alignments using RSeQC as previously described. We did not

directly analyze the raw files from TCGA or UNC, but instead reanalyzed the reported 5' to 3'

bias ratios. This ratio is similar to the inverse of the median coverage percentile: the smaller it is,

the larger the 3' bias in read coverage.

# References

1.    Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

2.    Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139

(2010).

3.    Trapnell, C. *et al.* Differential gene and transcript expression analysis of RNA-seq

experiments with TopHat and Cufflinks. *Nat. Protoc.* **7**, 562–578 (2012).

4.    Leek, J. T. & Storey, J. D. Capturing Heterogeneity in Gene Expression Studies by

Surrogate Variable Analysis. *PLoS Genet.* **3**, e161 (2007).

5.    Risso, D., Ngai, J., Speed, T. P. & Dudoit, S. Normalization of RNA-seq data using factor

analysis of control genes or samples. *Nat. Biotechnol.* **32**, 896–902 (2014).

6.    Johnson, W. E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray expression

data using empirical Bayes methods. *Biostatistics* **8**, 118–127 (2007).

7.    Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15 (2013).

8.    Conesa, A. *et al.* A survey of best practices for RNA-seq data analysis. *Genome Biol.* **17**, 1–

19 (2016).

9.    Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq

quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

10.    Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and

bias-aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

11.    Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

(2013).

12. Jun, E. *et al.* Method Optimization for Extracting High-Quality RNA From the Human

Pancreas Tissue. *Transl. Oncol.* **11**, 800–807 (2018).

13. Passow, C. N. *et al.* RNAlater and flash freezing storage methods nonrandomly influence

observed gene expression in RNAseq experiments. *bioRxiv* 379834 (2018)

doi:10.1101/379834.

14. Micke, P. *et al.* Biobanking of fresh frozen tissue: RNA is stable in nonfixed surgical

specimens. *Lab. Invest.* **86**, 202–211 (2006).

15. Ohmomo, H. *et al.* Reduction of Systematic Bias in Transcriptome Data from Human

Peripheral Blood Mononuclear Cells for Transportation and Biobanking. *PLoS One* **9**,

(2014).

16. Romero, I. G., Pai, A. A., Tung, J. & Gilad, Y. RNA-seq: impact of RNA degradation on

transcript quantification. *BMC Biol.* **12**, 42 (2014).

17. Xiong, B., Yang, Y., Fineis, F. R. & Wang, J. P. DegNorm: normalization of generalized

transcript degradation improves accuracy in RNA-seq analysis. *Genome Biol.* **20**, 75–75

(2019).

18. Li, S. *et al.* Multi-platform assessment of transcriptome profiling using RNA-seq in the

ABRF next-generation sequencing study. *Nat. Biotechnol.* **32**, 915–925 (2014).

19. Svensson, V. *et al.* Power analysis of single-cell RNA-sequencing experiments. *Nat.*

*Methods* **14**, 381–387 (2017).

20. Locksley E. McGann, Hongyou Yang, Michele Walterson. Manifestations of cell damage

after freezing and thawing. *Cryobiology* **25**, 178–185 (1988).

21. Röder, B., Frühwirth, K., Vogl, C., Wagner, M. & Rossmanith, P. Impact of long-term

storage on stability of standard DNA for nucleic acid-based methods. *J. Clin. Microbiol.* **48**,

4260–4262 (2010).

22.  Shao, W., Khin, S. & Kopp, W. C. Characterization of effect of repeated freeze and thaw

cycles on stability of genomic DNA using pulsed field gel electrophoresis. *Biopreserv.*

*Biobank.* **10**, 4–11 (2012).

23.  Schroeder, A. *et al.* The RIN: an RNA integrity number for assigning integrity values to

RNA measurements. *BMC Mol. Biol.* **7**, 3 (2006).

24.  Reiman, M., Laan, M., Rull, K. & Sõber, S. Effects of RNA integrity on transcript

quantification by total RNA sequencing of clinically collected human placental samples.

*FASEB J.* **31**, 3298–3308 (2017).

25.  Shen, Y. *et al.* Impact of RNA integrity and blood sample storage conditions on the gene

expression analysis. *Onco. Targets. Ther.* **11**, 3573 (2018).

26.  Sonntag, K.-C. *et al.* Limited predictability of postmortem human brain tissue quality by

RNA integrity numbers. *J. Neurochem.* **138**, 53 (2016).

27.  Yu, K. *et al.* Effect of multiple cycles of freeze–thawing on the RNA quality of lung cancer

tissues. *Cell and Tissue Banking* vol. 18 433–440 (2017).

28.  Jaffe, A. E. *et al.* qSVA framework for RNA quality correction in differential expression

analysis. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 7130 (2017).

29.  Bao, W.-G. *et al.* Biobanking of Fresh-frozen Human Colon Tissues: Impact of Tissue Ex-

vivo Ischemia Times and Storage Periods on RNA Quality. *Ann. Surg. Oncol.* **20**, 1737–

1744 (2012).

30.  Zeugner, S., Mayr, T., Zietz, C., Aust, D. E. & Baretton, G. B. RNA Quality in Fresh-

Frozen Gastrointestinal Tumor Specimens—Experiences from the Tumor and Healthy

Tissue Bank TU Dresden. in *Pre-Analytics of Pathological Specimens in Oncology* 85–93

(Springer, Cham, 2015).

31.   Li, J., Jiang, H. & Wong, W. H. Modeling non-uniformity in short-read rates in RNA-Seq

data. *Genome Biology* **11**, (2010).

32.   Hoen, P. A. C. *et al.* Reproducibility of high-throughput mRNA and small RNA sequencing

across laboratories. *Nat. Biotechnol.* **31**, 1015–1022 (2013).

33.   Thompson, K. L., Scott Pine, P., Rosenzweig, B. A., Turpaz, Y. & Retief, J.

Characterization of the effect of sample quality on high density oligonucleotide microarray

data using progressively degraded rat liver RNA. *BMC Biotechnol.* **7**, 57 (2007).

34.   Zhao, S., Zhang, Y., Gamini, R., Zhang, B. & von Schack, D. Evaluation of two main

RNA-seq approaches for gene quantification in clinical RNA sequencing: polyA+ selection

versus rRNA depletion. *Sci. Rep.* **8**, 1–12 (2018).

35.   Sultan, M. *et al.* Influence of RNA extraction methods and library selection schemes on

RNA-seq data. *BMC Genomics* **15**, 675 (2014).

36.   Zhao, W. *et al.* Comparison of RNA-Seq by poly (A) capture, ribosomal RNA depletion,

and DNA microarray for expression profiling. *BMC Genomics* **15**, (2014).

37.   Zhu, Y., Davis, S., Stephens, R., Meltzer, P. S. & Chen, Y. GEOmetadb: powerful

alternative search engine for the Gene Expression Omnibus. *Bioinformatics* **24**, 2798–2800

(2008).

38.   McIntyre, L. M. *et al.* RNA-seq: technical variability and sampling. *BMC Genomics* **12**, 1–

13 (2011).

39.   Hansen, K. D., Irizarry, R. A. & Wu, Z. Removing technical variability in RNA-seq data

using conditional quantile normalization. *Biostatistics* **13**, 204–216 (2012).

40.   Leek, J. T., Evan Johnson, W., Parker, H. S., Jaffe, A. E. & Storey, J. D. The sva package

680 for removing batch effects and other unwanted variation in high-throughput experiments.

681 *Bioinformatics* vol. 28 882–883 (2012).

682 41. Copois, V. *et al.* Impact of RNA degradation on gene expression profiles: assessment of

683 different methods to reliably determine RNA quality. *J. Biotechnol.* **127**, 549–559 (2007).

684 42. Adiconis, X. *et al.* Comparative analysis of RNA sequencing methods for degraded or low-

685 input samples. *Nat. Methods* **10**, 623–629 (2013).

686 43. Pertea, M. *et al.* Thousands of large-scale RNA sequencing experiments yield a

687 comprehensive new human gene list and reveal extensive transcriptional noise. *bioRxiv*

688 332825 (2018) doi:10.1101/332825.

689 44. Wang, L. *et al.* Measure transcript integrity using RNA-seq data. *BMC Bioinformatics* **17**,

690 58 (2016).

691 45. Foley, J. W. *et al.* Gene expression profiling of single cells from archival tissue with laser-

692 capture microdissection and Smart-3SEQ. *Genome Research* vol. 29 1816–1825 (2019).

693 46. Wang, Y. *et al.* Precision and functional specificity in mRNA decay. *Proceedings of the*

694 *National Academy of Sciences* vol. 99 5860–5865 (2002).

695 47. Yang, E. *et al.* Decay rates of human mRNAs: correlation with functional characteristics

696 and sequence attributes. *Genome Res.* **13**, 1863–1872 (2003).

697 48. Narsai, R. *et al.* Genome-wide analysis of mRNA decay rates and their determinants in

698 Arabidopsis thaliana. *Plant Cell* **19**, 3418–3436 (2007).

699 49. Feng, H., Zhang, X. & Zhang, C. mRIN for direct assessment of genome-wide and gene-

700 specific mRNA integrity from large-scale RNA-sequencing data. *Nat. Commun.* **6**, 7816

701 (2015).

702 50. Wang, Y. *et al.* The Impact of Different Preservation Conditions and Freezing-Thawing

Cycles on Quality of RNA, DNA, and Proteins in Cancer Tissue. *Biopreserv. Biobank.* **13**,

335–347 (2015).

51.  Gazestani, V. H. *et al.* A perturbed gene network containing PI3K/AKT, RAS/ERK,

WNT/β-catenin pathways in leukocytes is linked to ASD genetics and symptom severity.

*bioRxiv* 435917 (2019) doi:10.1101/435917.

52.  Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114 (2014).

53.  Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–

2079 (2009).

54.  Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-

throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).

55.  Pramparo, T. *et al.* Prediction of autism by translation and immune/inflammation

coexpressed genes in toddlers from pediatric community practices. *JAMA Psychiatry* **72**,

386–394 (2015).

56.  Collado-Torres, L. *et al.* Reproducible RNA-seq analysis using recount2. *Nat. Biotechnol.*

**35**, 319–321 (2017).

57.  Wang, L., Wang, S. & Li, W. RSeQC: quality control of RNA-seq experiments.

*Bioinformatics* **28**, 2184–2185 (2012).