Hyperactive end joining repair mediates resistance to DNA damaging therapy in p53-deficient cells

Rashmi J. Kumar[1,2], Hui Xiao Chao[3], Victoria R. Roberts[1], Aurora R. Sullivan[1], Sonam J. Shah[1], Dennis A. Simpson[1], Wanjuan Feng[1], Anne-Sophie Wozny[1], Sunil Kumar[1], Jeremy E. Purvis[3,4], Gaorav P. Gupta[1,2,5,6,#]

[1]Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

[2]Curriculum in Genetics and Molecular Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

[3]Curriculum in Bioinformatics and Computational Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

[3]Department of Computational Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

[4]Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

[5]Department of Radiation Oncology, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

[6]Department of Biochemistry and Biophysics, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

#Correspondence:
Gaorav P. Gupta, MD PhD
Assistant Professor
Department of Radiation Oncology
Department of Biochemistry and Biophysics
University of North Carolina at Chapel Hill
Chapel Hill, NC 27599
gaorav_gupta@med.unc.edu

Abstract

TP53 mutations in cancer are associated with poor patient outcomes and resistance to DNA damaging therapies[1–3]. However, the mechanisms underlying treatment resistance in p53-deficient cells remain poorly characterized. Here, we show that p53-deficient cells exhibit hyperactive repair of therapy-induced DNA double strand breaks (DSBs), which is suppressed by inhibition of DNA-dependent protein kinase (DNA-PK). Single-cell analyses of DSB repair kinetics and cell cycle state transitions reveal an essential role for DNA-PK in suppressing S phase DNA damage and mitotic catastrophe in p53-deficient cells. Yet, a subset of p53-deficient cells exhibit intrinsic resistance to therapeutic DSBs due to a repair pathway that is not sensitive to DNA-PK inhibition. We show that p53 deficiency induces overexpression of DNA Polymerase Theta (Pol θ), which mediates an alternative end-joining repair pathway that becomes hyperactivated by DNA-PK inhibition[4]. Combined inhibition of DNA-PK and Pol θ restores therapeutic DNA damage sensitivity in p53-deficient cells. Thus, our study identifies two targetable DSB end joining pathways that can be suppressed as a strategy to overcome resistance to DNA-damaging therapies in p53-deficient cancers.

Introduction

TP53 is the most commonly mutated tumor suppressor gene[5]. p53 mediates pleiotropic tumor suppressive effects through regulation of cell cycle arrest, apoptosis, and cellular metabolism in response to cellular stress[6,7]. Beyond its role as a tumor suppressor, loss of functional p53 is associated with poor prognostic outcomes across many different cancer types[1,8–10]. There is both clinical and preclinical evidence that p53-deficient cancers exhibit resistance to a variety of DNA damaging therapies[2,3,11–14].

The mechanisms for therapeutic resistance in p53-deficient cells remains poorly

characterized. Past work has suggested a role for loss of p53-mediated apoptosis[12,15].

However, the response of epithelial cancer cells to DNA damaging therapy is often

determined by the efficiency of inducing senescence or mitotic catastrophe, rather than

apoptosis[16,17]. p53 is also a transcription factor that responds to DNA double strand

breaks (DSBs) to determine cellular fate[7,18]. Recent insights have revealed the

importance of p53-signaling waves in regulation of cellular fate decisions of quiescence

versus cell cycle re-entry after DNA damage[19,20]. However, the mechanisms that

determine such cell fate decisions upon DNA damage induction in p53-mutant epithelial

cells have not been established, and may lead to novel strategies to restore treatment

sensitivity.

In this study, we investigate altered DNA repair mechanisms in p53 deficiency as

a major contributor to resistance to DNA damaging therapies. We find that p53-deficient

cells exhibit hyperactive repair and accelerated resolution of DNA damage foci. Utilizing

live-cell imaging, we show that this ability to resolve DNA damage rapidly is partially

dependent on DNA-PK, a critical serine/threonine kinase in the non-homologous end

joining (NHEJ) pathway[21]. Inhibition of DNA-PK using the small molecule inhibitor

NU7441 partially sensitizes p53-deficient cells to DSB inducing agents. We further show

that this effect is specifically due to propagation of S phase related damage leading to

mitotic catastrophe, highlighting a role for DNA-PK in S phase DNA damage repair that

was previously under appreciated. Furthermore, using chromosomal break repair assays

we show that in the context of inhibitor treatment, some p53-deficient cells utilize

alternative end-joining repair in a compensatory manner to escape cell death. Thus, our

work provides critical insight into a clinically-relevant mechanism for why p53-deficient

cells are resistant to DNA damaging therapies.

Results

p53-deficient cells exhibit radioresistance and accelerated resolution of DNA DSBs

We first established an isogenic cell system to investigate determinants of

treatment-induced cell fate in p53-deficient cells. In order to minimize potential

contributions of accessory mutations on phenotypes observed in cancer cell line models,

we used CRISPR/Cas9 to disrupt *TP53* in the p53-proficient immortalized epithelial cell

line model hTert-RPE1 ("RPE1"), which has also been a preferred model for investigating

p53-dependent cell fate[18,20,22]. Two independent CRISPR/Cas9-targeted *TP53*[-/-] RPE1

clones were selected for further study after confirming cells were deficient for p53 protein

and lacked p53-dependent transcriptional induction of p21 in response to ionizing

radiation (IR) (Supplementary Fig. 1a-c).

To assess whether p53 deficiency confers a proliferative advantage when treated

with ionizing radiation, we performed a mixed competition assay. We took mCherry

labelled RPE1 and mixed them with equal numbers of unlabeled *TP53*[-/-] RPE1 or p53-

proficient RPE1 (control) (Fig. 1a). We quantified the relative abundance of the unlabeled

cells after to exposure to IR (0 – 6Gy), normalized to untreated samples at each timepoint.

RPE1 labeled and unlabeled cells maintained stable representation across time and

treatment conditions (Supplementary Fig. 1d). Additionally, p53-deficient cells did not

demonstrate a proliferation advantage in the absence of RT. However, treatment with IR

at any dose level led to substantial positive selection for p53-deficient cells(Fig. 1b,c).

We also observed that p53 deficiency induced resistance to the radiomimetic clastogen,

Neocarzinostatin (NCS) by colony forming assay (Supplementary Fig. 1e-i). Thus p53

deficiency in this isogenic model is sufficient to induce radioresistance.

Unrepaired DSBs can suppress proliferation through the engagement of DNA

damage-induced cell cycle checkpoints. We examined kinetics of DSB repair by

performing immunofluorescence for 53BP1 and γH2AX after treatment of p53 WT and

$TP53^{-/-}$ cells with 5Gy IR (Fig. 1d,e and Supplementary Fig. 1j,k). We observed a

reduction in the number of 53BP1 damage foci in $TP53^{-/-}$ cells as early as 30 minutes

after treatment, that became even more pronounced by 4 hours post-treatment (Fig.

1d,e). Similar patterns of reduced foci formation were also apparent with γH2AX staining

at early timepoints (Supplementary Fig. 1j,k). Quantification of IR-induced DSBs by

neutral COMET assay revealed an equivalent DSB burden induced immediately after 5Gy

IR, irrespective of p53 status (Fig. 1f,g). However, by 4 hours post-treatment, tail DNA

percent was significantly reduced in the $TP53^{-/-}$ cells while remaining elevated in p53-

proficient RPE1 (Fig. 1f,g). Thus, p53 deficiency is sufficient to induce radioresistance

and accelerated DSB repair in an isogenic model.

Inhibition of DNA-PK restores DNA damage foci formation in p53-deficient cells

To directly assess the relationship between DSB repair kinetics, cell cycle status,

and cell fate at the single cell level, we established a live cell imaging platform (Fig. 2a).

RPE1 cells were dually labeled with PCNA-mCherry (to monitor cell cycle state

transitions) and 53BP1-mVenus (to monitor DSB foci kinetics) (Fig. 2b)[23,24]. These dual

labeled cells were treated with scrambled siRNA (si-Control) or siRNA targeting $TP53$

(si-$TP53$), the latter of which resulted in >90% knockdown of $TP53$ transcript and

elimination of p53-dependent $CDKN1A$ transcription in response to IR (Fig. 2c). 48 hours

after siRNA treatment, RPE1 cells were imaged for a total of 72 hours every 10 minutes, and 18 hours into imaging, the DSB inducing agent was added (Fig. 2a). To minimize time from radiation exposure to image capture and to induce equivalent DSBs in each population of cells, we utilized 100 ng/ml of Neocarzinostatin (NCS), a well-known radio-mimetic. NCS has been previously utilized in studies evaluating DNA DSB repair in conjunction with live-cell imaging and has been shown to induce peak DSBs within 10 minutes of drug addition[25,26]. This experimental design allowed us to determine the cell cycle status of each cell within the asynchronous cell population at the time of NCS exposure. After NCS treatment, single-cell analyses for DSB repair foci kinetics and cell cycle outcomes were performed. As anticipated from Fig. 1b,c, analysis of global proliferation by live-cell imaging revealed significantly greater proliferation of p53-deficient RPE1 cells relative to controls after NCS treatment (Supplementary Fig. 2a,b).

To analyze DSB repair kinetics in cells exposed to NCS, we tracked and quantified 53BP1 foci in single cells and plotted heatmaps of damage foci burden over time from cell birth to mitosis (Fig. 2d). Our results indicate that cells with functional p53 sustain high levels of damage foci in a prolonged manner after NCS exposure. In contrast, p53-deficient cells developed a lower peak burden of 53BP1 foci after NCS treatment, with accelerated resolution of damage foci to baseline levels (Fig. 2d,f). Given the rapidity with which 53BP1 foci were being resolved, we hypothesized that hyperactive NHEJ may be contributing. We thus performed the same experiment in the presence of an inhibitor of DNA-dependent Protein Kinase (DNA-PKi, NU7441 0.5μM), which targets the central kinase in the NHEJ pathway[27-29]. Strikingly, DNA-PKi qualitatively abolished the difference in 53BP1 kinetics after NCS treatment between p53-deficient and proficient

cells (Fig. 2e). To quantitatively assess the magnitude in damage burden, we calculated

peak maximum 53BP1 foci values for each cell represented in the heatmap (Fig. 2f).

Consistent with the heatmap representation, the median peak foci count after NCS

treatment was 40% lower in si-*TP53* treated cells relative to controls (Fig. 2f, p<0.0001).

Notably, DNA-PKi treatment resulted in a >2-fold increase in peak 53BP1 foci levels in

the p53-deficient cells, whereas there was no comparable effect in control cells (Fig. 2f).

These results indicate that DNA-PK activity is required for accelerated resolution of

clastogen-induced DNA damage foci in p53-deficient cells.

Following this analysis, we studied the effects of DNA-PKi in different phases of

the cell cycle during drug exposure. We used PCNA live-cell imaging to resolve cell cycle

phase transitions in cells tracked for 53BP1 foci kinetics. We performed area under the

curve (AUC) analyses in single cells to estimate total DNA damage burden during G1 and

S phase after NCS exposure (Fig. 2g). This analysis revealed that the diminished 53BP1

foci burden observed in p53-deficient cells was most pronounced during S phase relative

to control cells (Fig. 2g). DNA-PKi treatment significantly increased S phase 53BP1

burden in both si-Control and si-*TP53* treated RPE1 cells (Fig. 2g). While si-*TP53* treated

cells in G1 were also affected to a lesser degree, we were curious to examine if the effect

was in part due to loss of the p53-dependent G1/S checkpoint resulting in propagation of

unrepaired DNA damage into S phase. Indeed, we found that DNA-PKi induced a drastic

increase in 53BP1 foci as p53-deficient cells transitioned from G1 to S phase, which

subsequently diminished over time (Fig. 2h, p<0.00001 at t = start of S phase). Thus,

DNA-PK is required for hyperactive resolution of clastogen-induced DSB foci in p53-

deficient cells, and most prominently during S phase.

Checkpoint responses halt p53-proficient cells upon exposure to NCS while p53-deficient cells continue to cell cycle despite NCS exposure

To investigate the association between DNA damage and activation of cell cycle checkpoints, we quantified cell cycle phase durations for all treatment conditions (Fig. 3a,b). p53-proficient G1 cells exposed to NCS induced a significant prolongation of G1, indicative of G1/S checkpoint activation, with a substantial proportion of cells remaining arrested for the duration of imaging (Fig. 3c and Supplementary Fig. 3a-c). Similarly, cells exposed to NCS in S phase exhibited a G2-M checkpoint (Fig. 3d). p53-deficient cells exhibited no prolongation of G1 duration after NCS, consistent with the notion that G1/S checkpoint activation is p53-dependent (Fig. 3e)[30,31]. DNA-PK inhibition did not alter G1 duration in either p53-proficient or p53-deficient cells (Fig. 3c,e). In contrast, DNA-PKi increased the duration of G2-M checkpoints irrespective of p53 status (Fig. 3d,f). These observations suggest that increased levels of S phase DNA damage induced by DNA-PKi and NCS treatment (see Fig. 2g,h) result in activation of a G2/M checkpoint that is, at least partially, p53-independent. However, the duration of G2/M checkpoint activation differed by p53 status. While p53-proficient cells frequently remained arrested for the entire duration of imaging (open circles, Fig. 3c,d), p53-deficient cells experienced a more transient prolongation of G2 duration followed by progression into mitosis (Fig. 3e,f).

Inhibition of DNA-PK induces catastrophic mitoses in p53-deficient cells

We next used a heatmap representation to track the fate of individual cells from birth until mitosis (Fig. 4a,b, top panels). Red bars indicate a mitotic catastrophe or apoptosis event (Supplementary Fig. 4a,b). The median cell cycle time for both untreated p53-proficient and p53-deficient cells was approximately 22-24 hours. NCS treatment is

indicated as a dashed line at the 18 hour timepoint. Individual cells are ordered according

to cell cycle phase at the time of NCS treatment (G1 versus S) and eventual cell fate

(viable, G1 arrest, G2 arrest, or mitotic catastrophe/apoptosis). The majority (70%) of p53-

proficient (si-Control) G1 cells exposed to NCS activated a G1 checkpoint that was

maintained for the remainder of imaging (Fig. 4a). 26% of these cells underwent G2 arrest

or mitotic catastrophe, whereas only 3% retained their proliferative capacity (Fig. 4a).

Control cells exposed to NCS in S phase exhibited more diverse cell fates: 40% G2 arrest,

17% mitotic catastrophe, and 43% that retained proliferative capacity.   These

observations, made using single-cell tracking of asynchronous cell populations, are

consistent with observations of intrinsic radioresistance of S phase cells using cell

synchronization methods[32]. In contrast, the majority of p53-deficient (i.e., si-*TP53*

treated) cells in G1 or S at the time of NCS treatment remained viable without perceptible

engagement of any cell cycle checkpoints (Fig. 4b, 80% and 87%, respectively).

Consistent with prior 53BP1 analyses, S phase cells are most sensitized to DNA-PKi as

the addition of the inhibitor increased G2 arrest frequency in control cells (40% to 91%),

and increased mitotic catastrophe in p53-deficient cells (13% to 47%, Fig. 4a,b). In total,

the percentage of viable p53-deficient cells after NCS decreased from 87% to 47% when

treated in S phase with DNA-PK inhibition (p<0.0001, Fisher's exact test).

Despite the significant increase in mitotic catastrophe induced by combined

treatment with DNA-PKi and NCS, 47% of p53-deficient cells exhibit intrinsic resistance

to therapy with retained proliferative viability (Fig. 4b). We hypothesized that levels of

unrepaired DNA damage may be determinants of viable (*i.e.*, resistant) versus non-viable

(*i.e.*, sensitive) cell fates. To evaluate this hypothesis, we quantified integral DNA damage

burden in p53-deficient RPE1 with viable versus non-viable mitotic outcomes (Fig. 4c and

Supplementary Fig. 4c). The mean integral DNA damage burden was approximately 2-

fold higher in non-viable cells, relative to cells that viably completed mitosis (p<0.0001).

Further analysis revealed that integral DNA damage burden in S phase was most highly

associated with cell viability after drug treatment (Supplementary Fig. 4c). In addition, we

traced the average 53BP1 foci burden over time for these two cohorts (Fig. 4d). Our

results indicate that cells with non-viable mitotic outcomes have an increased peak value

of DNA damage after treatment with DNA-PKi and NCS, which remains elevated over

time (p<0.0001 at t=20 hrs, Fig. 4d). Conversely, these findings indicate that p53-

deficient cells that exhibit intrinsic therapeutic resistance may be utilizing compensatory

DSB repair pathways to counteract the effects of NCS and DNA-PKi prior to mitotic entry.

p53-deficient cells utilize alternative end-joining pathways in the absence of active

DNA-PK

Prior studies have demonstrated that cells with NHEJ deficiency exhibit a

compensatory increase in alternative end-joining repair mediated by DNA polymerase

theta (Pol θ, gene *POLQ*)[4,33,34]. Polymerase theta dependent end joining (TMEJ) of DNA

DSBs is characterized by deletions and templated insertions that are flanked by short

tracts of sequence identity, or microhomology (MH)[4]. We found that *POLQ* expression

was 10- to 20-fold higher in two independent *TP53*[-/-] RPE1 clones, relative to parental

TP53 wild-type cells (Fig. 5a). *POLQ* is also overexpressed in TCGA breast, lung,

bladder, colorectal, gastric, glioblastoma, pancreatic, prostate, melanoma, and uterine

cancers with *TP53* mutation, relative to their *TP53* wild-type counterparts (Fig. 5b).

To assess whether hyperactive TMEJ contributes to therapeutic resistance of

TP53[-/-] RPE1 cells to NCS and DNA-PKi, we sought to inhibit Pol θ. As pharmacological

inhibitors of Pol θ are not yet commercially available, we created a double knockout

POLQ[-/-]TP53[-/-] RPE1 line (Supplementary Fig. 5a). Bi-allelic frameshift mutations in

POLQ were confirmed by Sanger sequencing and functional deficiency was established

using an extrachromosomal TMEJ repair assay (Supplementary Fig. 5b-d)[4].

To directly assess whether TMEJ repair is increased after DNA-PKi treatment, we

analyzed chromosomal break repair patterns at a site-specific DSB in p53-deficient RPE1

cells. Cells were transfected with Cas9 ribonucleoprotein (RNP) complexes that target

the *LBR* locus, with or without DNA-PKi[35]. Genomic DNA was harvested 60 hours later

and analyzed for break repair patterns using next generation sequencing (NGS) (Fig. 5c).

Target amplification and TIDE analyses confirmed high rates of target site cleavage in all

samples transfected with a full complement of Cas9-RNP (Supplementary Fig. 5e,f)[36].

We applied a bioinformatic algorithm (ScarMapper, see methods) to characterize the

spectrum of repair products with at least 0.1% prevalence, classified according to the size

of left deletion (LD), right deletion (RD), insertion (Ins), and microhomology (MH)

(ScarMapper Methods). Indels <5bp were categorized as NHEJ, with the predominant

repair product being a +A 1bp insertion[35]. TMEJ was defined as repair products whose

frequency was diminished by at least 2-fold in *POLQ[-/-]* cells. All other repair products

were categorized as "Unclassified." DNA-PK inhibition in *TP53[-/-]* RPE1 cells results in a

substantial reduction in NHEJ repair, with a compensatory increase in TMEJ to nearly

45% of all DSB repair (Fig. 5d,e). In contrast, DNA-PK inhibition in *POLQ[-/-]TP53[-/-]* RPE1

cells did not result in a substantial increase TMEJ signature repair (Fig. 5f,g). However,

a higher proportion of Unclassified repair products were detected (Fig. 5f,g). A limitation

of NGS analysis of DSB break repair is that non-amplifiable target loci are not measured.

Thus, we used digital PCR to quantify the *LBR* locus detection rate, relative to a control

locus, upon inhibition of DNA-PK and/or Pol θ (Supplementary Fig. 5g,h). *LBR* locus

detection rates were most reduced upon inhibition of DNA-PK and Pol θ, consistent with

overall inhibition of DSB repair (Supplementary Fig. 5h). These observations confirm an

essential role for TMEJ in compensatory repair of chromosomal DSBs upon

pharmacologic inhibition of DNA-PK.

To determine the impact of *POLQ* inhibition on cellular viability, we performed

clonogenic survival assays in the parental *TP53$^{-/-}$* and *POLQ$^{-/-}$TP53$^{-/-}$* RPE1 lines treated

with NCS with or without DNA-PKi. Genetic deficiency in *POLQ* resulted in significantly

reduced viability after NCS treatment, particularly in combination with DNA-PKi (Fig. 5h).

TP53$^{-/-}$ RPE1 cells with inhibition of both TMEJ and NHEJ repair pathways had

comparable clonogenic survival to p53-proficient RPE1 cells (see Supplementary Fig.

1e). Collectively, these findings indicate that hyperactive end joining repair via NHEJ and

TMEJ mediate resistance to DNA damaging therapy induced by p53 deficiency (Fig. 5i).

Discussion

These results recognize enhanced DNA end joining repair capacity as a novel

component of therapeutic resistance induced by p53 deficiency, and that loss of functional

p53 alone is sufficient to increase hyperactive repair. Our findings indicate that DSB end

joining hyperactivity is particularly relevant for suppressing S phase DNA damage burden,

which we find is a key determinant of mitotic catastrophe (Fig. 5i). Although NHEJ Is

conventionally considered to be most critical for repair in G1, we observed a relatively

greater impact of DNA-PK inhibition on the fate of S phase cells after treatment with a radiomimetic. There are several potential explanations for this unanticipated observation. First, recent findings suggest that DNA-PK may be dispensable for synapsis formation during NHEJ[37]. Accordingly, repair of "simple" DSBs in G1 phase may have a reduced reliance on DNA-PK, whereas repair of more "complex" DSBs in S phase may require DNA-PK, possibly in partnership with the nuclease Artemis[38–40]. Second, it is possible that DNA-PK inhibition may be more impactful in S phase due to trapping of Ku proteins at DSBs, which inhibits the activation of homologous recombination pathways[41]. Third, DNA-PK may be particularly important in early S phase, when sister chromatids are not broadly present. Notably, we observed a prominent peak of unrepaired DSBs just as p53-deficient cells transitioned from G1 to S phase. Our observation that DSB end joining hyperactivity in p53-deficient cells is highly sensitive to DNA-PK inhibition warrants further mechanistic investigation. Recently, CYREN (cell cycle regulator of NHEJ) has been proposed to be a cell-cycle phase specific inhibitor of the Ku70/80 heterodimer that is critical for restricting NHEJ to G1[42]. It is therefore possible that p53-deficiency may transcriptionally reprogram cell cycle-inhibitors of NHEJ to enable hyperactive repair, though that is beyond the scope of this study.

Regulatory mechanisms that confer TMEJ hyperactivity in cancer are not well understood, although transcriptional overexpression of *POLQ* has also been observed in breast and ovarian cancers with *BRCA1*/*BRCA2* deficiency or mutations in other genes that confer Pol θ synthetic lethality[33,43]. Recent work investigating integrated pathway analysis of *TP53* deficiency noted *POLQ* to be frequently overexpressed in *TP53* pathway deficient cancers[1]. Our findings, in an isogenic p53-deficient cell line model, indicate that

this relationship may be causal. The mechanism for p53-dependent suppression of *POLQ*

expression remains to be elucidated, and may entail the regulation of non-coding RNAs[44].

The use of TMEJ can also be explained by the potential creation of more complex DSBs

upon NHEJ suppression that serve as poor substrates for homologous recombination

(HR). Indeed, the molecular mechanisms of NHEJ and TMEJ hyperactivity induced by

p53 deficiency warrant further investigation.

Radiotherapy and other forms of DNA damaging therapy are employed in the vast

majority of cancer patients[45]. Resistance to DNA damaging therapy may thus explain the

adverse clinical outcomes associated with *TP53* mutations in many different cancer

types[1]. Our study supports the investigation of DNA-PK inhibitors administered in

combination with DNA damaging therapy (including radiotherapy) in patients with p53-

deficient cancers. Additionally, as inhibitors of Pol θ are currently in development[46], our

study suggests that combined inhibition of both DNA-PK and Pol θ represents a promising

strategy to reverse the therapeutic DNA damage resistance in p53-deficient cancers.

Materials and Methods

Key Resources Table:

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
F(ab)2-Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 633 (1:10,000 for IF)	Thermo Fisher Scientific	Cat# A-21072, RRID:AB_2535733
Chicken anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 488, (1:10,000 for IF)	Thermo Fisher Scientific	Cat# A-21200, RRID:AB_2535786
Mouse Anti-beta-Actin Monoclonal Antibody, Unconjugated, Clone AC-15 (1:10,000 for WB)	Sigma-Aldrich	Cat# A1978, RRID:AB_476692
Rabbit Anti-53BP1 Polyclonal Antibody (1:500 for IF)	Bethyl	Cat# A300-272A, RRID:AB_185520
Mouse Anti-p53 (1C12) mAb Antibody (1:1000 for WB)	Cell Signaling Technology	Cat# 2524, RRID:AB_331743
Rabbit Anti-phosphorylated Histone H2AX (γ-H2AX) Polyclonal Antibody (1:500 for IF)	Trevigen	Cat# 4418-APC-100
Chemicals, Peptides, and Recombinant Proteins		
Trypsin EDTA	Gibco	25200-056
Polyethylenimine, Linear (MW 25,000)	Polysciences	23966
Bovine Serum Albumin	Fisher Scientific	BP9706-160
Corning® Cell-Tak™ and Tissue Adhesive	Corning	354240
RNAiMax	ThermoFisher	13778100
Critical Commercial Assays		
PlasmoTest	Invitrogen	REP-PT1

RNAeasy Plus Mini Kit	Qiagen	74136
Comet Assay Kit	Trevigen	4250-050-K
Q5® Hot Start High-Fidelity 2X Master Mix	NEB	M0494S
NEBuilder® HiFi DNA Assembly Master Mix	NEB	E2621L
TOPO® TA Cloning® Kit for Sequencing	Invitrogen	450030
T4 DNA Ligase	NEB	M0202S
EdU-Click 594	baseclick	BCK-Edu594
NEON Electroporation Kit	ThermoFisher	MPK1025
Cas9 Protein and TracrRNA for Alt-R Electroporation	IDT	Cas9 (s.p. high fidelity) #1081060 TracrRNA #1072532
Experimental Models: Cell Lines		
hTERT-RPE1-Tricolor Reporter (PCNA-mCherry, 53BP1-mVenus, H2B-mTurquoise)	Gift from Dr. Jeremy Purvis	(See Citations)
hTERT-RPE1	ATCC	ATCC® CRL-4000™
hTERT-RPE1-*TP53-/-*	This paper	
hTERT-RPE1-*TP53-/-POLQ-/-*	This paper	
Oligonucleotides (sgRNAs and Primers)		
sgLBR	GCCGATGGTGAAGTGGTAAG	Synthesized at: IDT
sgTP53_Exon2	TCGACGCTAGGATCTGACTG	IDT
sgTP53_Dwnstream_Intron	GAAACTGTGAGTGGATCCAT	IDT
sgPOLQ_1	ACTACTCTCAGCTTGA	IDT
sgPOLQ_2	TCAGGAGCATTGCAGCAGAG	IDT
LBR_Fwd	AAATGGCTGTCTTTCCCAGTAA	EtonBio
LBR_Rev	ACGCAGTGGCTAAATCATCC	EtonBio
TP53 RTqPCR Primer Fwd	GAGGTTGGCTCTGACTGTACC	EtonBio
TP53 RTqPCR Primer Rev	TCCGTCCCAGTAGATTACCAC	EtonBio
CDKN1A RTqPCR Primer Fwd	TCACTGTCTTGTACCCTTGTGCTT	EtonBio
CDKN1A RTqPCR Primer Rev	AGAAATCTGTCATGCTGGTCTGCC	EtonBio

ONTARGET plus Human TP53 Si-RNA SMARTPOOL	Horizon Discovery (previously Dharmacon)	L-003329-00-0010
ONTARGET plus NON-TARGETTING control siRNAs SMARTPOOL	Horizon Discovery (previously Dharmacon)	D-001810-10-05
ESR1 Genomic Locus Fwd Primer	ATCTGTACAGCATGAAGTGCAAGA	EtionBio
ESR1 Genomic Locus Rev Primer	CTAGTGGGCGCATGTAGGC	EtonBio
ESR1 Genomic Locus Probe	T+C+T +AT+G +A+CC TG (Locked nucleic acid probe with HEX)	IDT (LNA : Locked Nucleic Acid Probe)
LBR Locus Probe	TGAGATTGAATGTAGCCTTT CTGGCCCTAA (with FAM)	

LBR Nested Sequencing Primers

Purple -> Binds genomic DNA
Green -> Phasing portion of primer
chr1:225423928-225424162
Size: 235 base pairs
Forward Primer: 114 base pairs left of cut
Reverse Primer: 123 base pairs right of cut
Rcomp = reverse complimentary

TCAATTCAAGCTCTGTTCCATCTTTATACTTCACAGTGTAAAGCTGGGAGGTGCTGTCGTGGCTCAGAATTTCTACTTCATAATAAAGTGAACTCCCAGGCCATCGA**CCT**_CTTA_CCACTTCACCATCGGCAAATTTCCTACTTGGCATTTTCTATAATTAACCTGAATAGTTTTAAAGAAAAAAATTTGAGTCAATACATACACATTTATGTATTCGTCTTTTTCCACAGGCTGA

Primer Name	Orientation	Location	Sequence
LBR2.1 F0	Forward	chr1:225423928-225423949	CGACGCTCTTCCGATCTTCAATTCAAGCTCTGTTCCATC
LBR2.1 F1	Forward	chr1:225423927-225423949	CGACGCTCTTCCGATCTTTCAATTCAAGCTCTGTTCCATC
LBR2.1 F2	Forward	chr1:225423927-225423949	CGACGCTCTTCCGATCTCTTCAATTCAAGCTCTGTTCCATC
LBR2.1 F3	Forward	chr1:225423927-225423949	CGACGCTCTTCCGATCTACTTCAATTCAAGCTCTGTTCCATC

LBR2.1 F4	Forward	chr1:22542392 7-225423949	CGACGCTCTTCCGATCTGACTTCAATTCAAGCTCTGTTCCATC
LBR2.1 F5	Forward	chr1:22542392 7-225423949	CGACGCTCTTCCGATCTAGACTTCAATTCAAGCTCTGTTCCATC
LBR2.1 R0	Rcomp	chr1:22542416 2-225424143	CGTGTGCTCTTCCGATCTTCAGCCTGTGGAAAAAGACG
LBR2.1 R1	Rcomp	chr1:22542416 3-225424143	CGTGTGCTCTTCCGATCTATCAGCCTGTGGAAAAAGACG
LBR2.1 R2	Rcomp	chr1:22542416 4-225424143	CGTGTGCTCTTCCGATCTGATCAGCCTGTGGAAAAAGACG
LBR2.1 R3	Rcomp	chr1:22542416 5-225424143	CGTGTGCTCTTCCGATCTTGATCAGCCTGTGGAAAAAGACG
LBR2.1 R4	Rcomp	chr1:22542416 6-225424143	CGTGTGCTCTTCCGATCTCTGATCAGCCTGTGGAAAAAGACG
LBR2.1 R5	Rcomp	chr1:22542416 7-225424143	CGTGTGCTCTTCCGATCTAC

Software and Algorithms		
Python ≥v3.5	*G. van Rossum, Python tutorial, Technical Report CS-R9526, Centrum voor Wiskund e en Informatica* (CWI), *Amster dam*, May *1995*	https://www.python.org/
Flow Jo	*FlowJo™ Software (Mac) [proliferation assay analysis] Becton, Dickinson and Company; 2019.*	https://www.flowjo.com/
Graphpad Prism v8	N.A.	https://www.graphpad.com/
Fiji	*Schindelin, J.; Arganda-Carreras, I. & Frise, E. et al. (2012), "Fiji: an open-source platform for biological-image analysis", Nature methods 9(7): 676-682, PMID 22743772, doi:10.1038/nmeth.2019 (on Google Scholar).*	https://imagej.net/Fiji#Downloads
CellProfiler	*CellProfiler Program Citation:* McQuin C, Goodman A, Chernyshev V, Kamentsky L, Cimini BA, Karhohs KW, Doan M, Ding L, Rafelski SM, Thirstrup D, Wiegraebe W, Singh S, Becker T, Caicedo JC, Carpenter AE (2018). CellProfiler 3.0: Next-	www.cellprofiler.org

	generation image processing for biology. PloS Biol. 16(7):e2005970 / doi. PMID: 29969450 (Research article) *Analyst Software Citation:* Jones TR, Kang IH, Wheeler DB, Lindquist RA, Papallo A, Sabatini DM, Golland P, Carpenter AE (2008) CellProfiler Analyst: data exploration and analysis software for complex image-based screens. BMC Bioinformatics 9(1):482/doi: 10.1186/1471-2105-9-482. PMID: 19014601 PMCID: PMC2614436	
NIS Elements AR software		https://www.nikon.com/products/microscope-solutions/lineup/img_soft/nis-elements/
SnapGene software v4.3.4	GSL Biotech	https://www.snapgene.com
Open Comet v1.3.1	BM Gyori, G Venkatachalam, PS Thiagarajan, D Hsu and MV Clement. "OpenComet: An automated tool for comet assay image analysis", Redox Biology, 2:457-465, 2014.	http://www.cometbio.org
ScarMapper		https://github.com/pkMyt1/ScarMapper.git
Additional Image Analysis Scripts	Code EV1 Supplementary and Code Availability (MATLAB scripts)	PMID: 30886052 PMID: 29102360
Other		
Genes	www.Ensembl.org	Ensembl v91

Cell Culture

WT (p53$^{+/+}$), WT Fusion-Reporter (p53$^{+/+,}$PCNA-mCherry, 53BP1-mVenus), (p53$^{-/-}$, and

p53$^{-/-}$Polq$^{-/-}$cells are hTERT immortalized RPEs. Cells were maintained in Dulbecco's

modified Eagle's medium (DMEM), with 10% Fetal Bovine Serum (Hyclone FBS) and

2mM L-glutamine (ThermoFisher).All cells were maintained at 37 C in an atmosphere of

5% CO2. Cells were routinely tested for mycoplasma contamination using PlasmoTest

(Invivogen).

Establishment of Stable Cell Lines

For the TP53 and Polq mutant cell lines, we used the Alt-R-CRISPR-Cas9 system (IDT).

We performed Neon transfection (Invitrogen) and followed the manufacturer's protocol

with Alt-R HiFi Cas9 nuclease, crRNA and tracrRNA purchased from IDT. crRNA was

designed using MIT CRISPR (http://crispr.mit.edu) to target Exon 2 of the TP53 gene for

the p53 mutant cell line and the polymerase domain of the Polq gene for the Polq mutant

cell line. Forty-eight hours after transfection, cells were seeded for single clone selection.

For the p53 gene editing experiment, a homologous template with a stop codon and SCA-

I site was provided for selection of gene edited cells. Restriction Enzyme screening, PCR

screening, and Sanger sequencing confirmed gene targeting, post which we performed

functional tests.

Immunofluorescence

Cells were fixed with 3% Paraformaldehyde for 15 min at RT, followed by permeabilization

with 0.25% TritonX-100 in PBS. Cells were subsequently processed for immunostaining

experiments using the antibodies listed below. Nuclei were visualized by staining with

DAPI. The primary antibodies used were: γH2AX (1:500, Trevigen, 4418-APC-100), and

53BP1 (1:500 for immunofluorescence, Bethyl, A300-272A). The secondary antibodies

were: FITC Goat Anti-Mouse IgG (H + L) (1:500, Jackson ImmunoResearch, 115-095-

003) and FITC Goat Anti Rabbit IgG (H + L) (1:500, Jackson ImmunoResearch, 111-095-

144). Images were acquired using the GE IN CELL 2200 high through-put imaging system

at 40x magnification.

siRNA Treatment

WT Fusion-Reporter RPE cells were passaged twice after -80 thaw and plated on 12 well

plates at a density of 100,000 cells/ well for siRNA treatment. Twenty-four hours post

plating, cells were exposed to 10nM / well sip53 (SMART pool from Dharmacon), and

siControl (Non-targetting SMART pool from Dharmacon), in OPTIMEM with RNA-iMAX

(ThermoFisher) as a transfection reagent. As a no-treatment control, cells were exposed

to RNA-iMAX and OPTIMEM without siRNA. 48 hours post transfection, cells were

transferred onto 12 well Cell-Tak coated glass plates (Cellvis), at a concentration of

50,000 cells/well for imaging. Prior to imaging and at the end of imaging, samples were

taken for RT-qPCR analysis of p53 mRNA to confirm si-RNA knockdown.

Mixed Competition Assay - Flow Cytometry

mCherry labelled and unlabeled hTERT-RPE1 cell lines were plated on 96 well plates at

a 50:50 ratio, and irradiated 2 hrs post plating at 0, 2, 4, or 6 Gy, and left to grow. At

indicated timepoints cells were harvested by trypsinizing and quenching with PBS with

5% BSA. Cells were fixed with 2% PFA and subsequently transferred to V-bottom plates

(ThermoFisher, 249570). Cells were quantified by flow cytometry using the Intellicyt iQue

at a volume of 100 ul / sample, collecting all events per well. For each condition, 6

biological replicates.

Time-Lapse Imaging Microscopy

Cells stably expressing Proliferating Cell Nuclear Antigen (PCNA)-mCherry and Tumor

Suppressor p53 Binding Protein 1 (53BP1) – mVenus were treated with si-RNA for 48

hours prior to imaging. PCNA-mCherry and 53BP1-mVenus fusion reporter is a gift from

Dr. Jeremy Purvis and Hui Chao Xiao. Cells were plated on Cell-Tak (Corning) coated

glass-bottom 12-well plates (Cellvis) with Phenol-free DMEM (Invitrogen) supplemented

with 10% FBS, and L-glutamine. Twenty-four hours post plating, cells were image

captured every 10 min for 72 h in the mCherry and mVenus fluorescence channels. 18

hours into imaging, DNA PKi was added at a concentration of 0.5 uM / well, and/or NCS

at a concentration of 100ng/ mL/ well. We commenced imaging every 10 minutes in both

channels for another 48 hours. Fluorescence images were obtained using a Nikon Ti

Eclipse inverted microscope with a 40x objective and Nikon Perfect Focus (PFS) system

to maintain focus during acquisition period. Cells were maintained at constant

temperature (37 °C) and atmosphere (5% CO_2). Nikon, NIS Elements AR software was

utilized for image acquisition. Image analysis was performed on ImageJ – Fiji and Cell

Profiler.

Colony Forming Assays

Cells lines used in the assay are indicated in the figures. Cells were treated with NCS

and/ or DNA-PKi for twenty-four hours, after which we performed a media change. For

colony formation experiments with ionizing radiation, cells were plated for IR treatment

with or without DNA-PKi, and inhibitor treatment was washed off after twenty-four hours.

Cells were subsequently incubated for 10-12 days at 37 °C to allow colony formation.

Colonies were stained by Coomassie blue and counted.

DNA Repair Assay

Cell lines used in the assay are indicated in the figure. 5×10^5 cells were transfected with

sgLBR2 and TracrRNA complexed Cas9 protein at final concentrations of

sgRNA:tracrRNA duplex: 22 pmol and Cas9 : 18 pmol per reaction, with Neon transfection

kit (Invitrogen) using 2 1350 V, 30 ms pulses in a 10 µL chamber. 60 hours post

transfection, cells were harvested for genomic DNA extraction (Nucleospin). Part of the

gDNA was utilized for Sanger Sequencing and TIDE analysis post amplification of the

genomic LBR2 locus. Remaining gDNA was amplified using NGS nested sequencing

primers and sent for sequencing and/ or Digital PCR.

Digital PCR

Primers and 5' hydrolysis probes were designed to specifically detect the copies of *LBR*

locus. *ESR1* locus was used as genomic control. Each reaction assay contained 10 µLof

2x dPCR Supermix for Probes (No dUTP), 0.9 µmol/L of respective primers, 0.25 µmol/L

of respective probes, and 10 ng of DNA in a final volume of 20 µL. Droplets were

generated using automated droplet generator (Bio-Rad catalog #186-4101) following

manufacturer's protocol. PCR parameters for *LBR* locus were 10 sec at 95 °C, then 40

cycles of 94 °C for 30 sec, 60 °C for 30 sec, and 72°C for 2 min followed by 98°C for 10

min with a ramping of 2 °C/sec at all steps. The PCR cycling parameters for *ESR1*

genomic locus were 10 sec at 95 °C, then 40 cycles of 94 °C for 30 sec and 60 °C for 1

min followed by 98°C for 10 min with a ramping of 2 °C/sec at all steps. After PCR

amplification, droplet reader (Bio-Rad QX200™ Droplet Reader Catalog #1864003) was

used to measure the end-point fluorescence signal in droplets as per the manufacturer's

protocol. The recorded data was subsequently analyzed with QuantaSoft software

version 1.7.4.0917 (Bio-Rad). Each Taqman probe was evaluated for sensitivity and

specificity.

DNA Repair Assay

Cell lines used in the assay are indicated in the figure. 5×10^5 cells were transfected with

sgLBR2 and TracrRNA complexed Cas9 protein at final concentrations of

sgRNA:tracrRNA duplex: 22 pmol and Cas9 : 18 pmol per reaction, with Neon transfection

kit (Invitrogen) using 2 1350 V, 30 ms pulses in a 10 µL chamber. 60 hours post

transfection, cells were harvested for genomic DNA extraction (Nucleospin). Part of the

gDNA was utilized for Sanger Sequencing and TIDE analysis post amplification of the

genomic LBR2 locus. For analysis of INDELs, 100 ng of gDNA was amplified using

phased primers. These libraries were indexed with the Illumina unique dual combinatorial

indices. Following pooling, 2 x 150 cycle sequencing was done on an Illumina iSeq™.

INDELs were identified by comparing the target reference sequence to the resulting

sequence reads in the FASTQ files via a 10-nucleotide sliding window using the

ScarMapper program.

Acknowledgements

We thank Kasia Kedziora, Samuel Wolff, and Juan Carvajal-Garcia for data acquisition

and technical assistance. We are grateful to the Gupta and Purvis Lab members for

helpful discussions. UNC Core labs (Microscopy Services Laboratory, Hooker Imaging

Core, and Flow Cytometry Core Facility) used in this study are supported in part by P30

CA016086 Cancer Center Core Support Grant to the UNC Lineberger Comprehensive

Cancer Center. Funding support was provided by the NCI/NIH (CA222092), Dept of

Defense (W81XWH-18-1-0047), and the University Cancer Research Fund. G.P.G. holds

a Career Award for Medical Scientists from the Burroughs Wellcome Fund. R.J.K. is

supported by the Cancer Cell Biology T32 Training Program (2T32CA071341-21) and the

UNC Medical Scientist Training Program (MSTP).

Author contributions

R.J.K and G.P.G. designed and conceived experiments. G.P.G. supervised the study.

R.J.K and H.C.X performed the live-cell imaging experiments. H.C.X. and J.E.P. provided

critical reagents and image analysis guidance. R.J.K performed and implemented

computational analyses for image processing. R.J.K. performed all additional

experiments and data analyses with statistical review. V.R.R., A.R.S., S.J.S., W.F., A-

S.W., and S.K. provided technical assistance on imaging acquisition, colony forming

assays, and digital PCR. D.A.S. developed break site sequencing analysis platforms.

R.J.K. and G.P.G. wrote the manuscript, with contributions from all authors. All authors

read and accepted the manuscript.

Conflicts of interests

458   The authors declare no competing interests that are pertinent to this study.

Correspondence

461   Correspondence should be addressed to Gaorav P. Gupta: gaorav_gupta@med.unc.edu

References

463   1.   Donehower, L. A. *et al.* Integrated Analysis of TP53 Gene and Pathway

464   Alterations in The Cancer Genome Atlas. *Cell Rep.* **28**, 1370-1384.e5 (2019).

465   2.   Lowe, S. W. *et al.* p53 status and the efficacy of cancer therapy in vivo. *Science*

466   *(80-.).* **266**, 807–810 (1994).

3.   Lee, J. M. & Bernstein, A. p53 Mutations increase resistance to ionizing radiation. *Proc. Natl. Acad. Sci. U. S. A.* **90**, 5742–5746 (1993).

4.   Wyatt, D. W. *et al.* Essential Roles for Polymerase θ-Mediated End Joining in the Repair of Chromosome Breaks. *Mol. Cell* **63**, 662–673 (2016).

5.   Nigro, J., Baker, S., Preisinger, A., Nature, J. J.- & 1989, undefined. *Mutations in the p53 gene occur in diverse human tumour types. Springer* (1989).

6.   Hafner, A., Bulyk, M. L., Jambhekar, A. & Lahav, G. The multiple mechanisms that regulate p53 activity and cell fate. *Nature Reviews Molecular Cell Biology* **20**, 199–210 (2019).

7.   Kastenhuber, E. R. & Lowe, S. W. Leading Edge Putting p53 in Context. (2017). doi:10.1016/j.cell.2017.08.028

8.   Young, K. H. *et al.* Structural profiles of TP53 gene mutations predict clinical outcome in diffuse large B-cell lymphoma: An international collaborative study. *Blood* **112**, 3088–3098 (2008).

9.   Miller, L. D. *et al.* An expression signature for p53 status in human breast cancer predicts mutation status, transcriptional effects, and patient survival. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 13550–13555 (2005).

10.   Robles, A. I. & Harris, C. C. Clinical outcomes and correlates of TP53 mutations and cancer. *Cold Spring Harbor perspectives in biology* **2**, (2010).

11.   Freed-Pastor, W. A. & Prives, C. Mutant p53: one name, many proteins. *genesdev.cshlp.org* doi:10.1101/gad.190678.112

488   12.   Lee, M. K. *et al.* Cell-type, Dose, and Mutation-type Specificity Dictate Mutant p53

489   Functions In Vivo. *Cancer Cell* **22**, 751–764 (2012).

490   13.   Bykov, V. J. N. *et al.* Restoration of the tumor suppressor function to mutant p53

491   by a low-molecular-weight compound. *Nat. Med.* **8**, 282–288 (2002).

492   14.   Huang, S. *et al.* P53 modulates acquired resistance to EGFR inhibitors and

493   radiation. *Cancer Res.* **71**, 7071–7079 (2011).

494   15.   Aubrey, B. J., Kelly, G. L., Janic, A., Herold, M. J. & Strasser, A. How does p53

495   induce apoptosis and how does this relate to p53-mediated tumour suppression?

496   *Cell Death and Differentiation* **25**, 104–113 (2018).

497   16.   Nardella, C., Clohessy, J. G., Alimonti, A. & Pandolfi, P. P. Pro-senescence

498   therapy for cancer treatment. *Nature Reviews Cancer* **11**, 503–511 (2011).

499   17.   Vitale, I., Galluzzi, L., Castedo, M. & Kroemer, G. Mitotic catastrophe: a

500   mechanism for avoiding genomic instability. *Nat. Rev. Mol. Cell Biol.* **12**, 385–392

501   (2011).

502   18.   Williams, A. B. & Rn Schumacher, B. p53 in the DNA-Damage-Repair Process.

503   *perspectivesinmedicine.cshlp.org* doi:10.1101/cshperspect.a026070

504   19.   Reyes, J. *et al.* Fluctuations in p53 Signaling Allow Escape from Cell-Cycle Arrest.

505   *Mol. Cell* **71**, 581-591.e5 (2018).

506   20.   Zhang, X. P., Liu, F., Cheng, Z. & Wang, W. Cell fate decision mediated by p53

507   pulses. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 12245–12250 (2009).

508   21.   Chang, H. H. Y., Pannunzio, N. R., Adachi, N. & Lieber, M. R. Non-homologous

509   DNA end joining and alternative pathways to double-strand break repair. *Nature*

Reviews Molecular Cell Biology **18**, 495–506 (2017).

22.   Krenning, L., Feringa, F. M., Shaltiel, I. A., vandenBerg, J. & Medema, R. H.

Transient activation of p53 in G2 phase is sufficient to induce senescence. *Mol.*

Cell **55**, 59–72 (2014).

23.   Chao, H. X. *et al.* Evidence that the human cell cycle is a series of uncoupled,

memoryless phases. *Mol. Syst. Biol.* **15**, (2019).

24.   Chao, H. X. *et al.* Orchestration of DNA Damage Checkpoint Dynamics across the

Human Cell Cycle. *Cell Syst.* **5**, 445-459.e5 (2017).

25.   Chao, H. X. *et al.* Orchestration of DNA Damage Checkpoint Dynamics across the

Human Cell Cycle. *Cell Syst.* **0**, (2017).

26.   Chao, H. X. *et al.* Evidence that the cell cycle is a series of uncoupled,

memoryless phases. (2018). doi:10.1101/283614

27.   Blackford, A. N. & Jackson, S. P. Molecular Cell Review ATM, ATR, and DNA-PK:

The Trinity at the Heart of the DNA Damage Response. (2017).

doi:10.1016/j.molcel.2017.05.015

28.   Zhao, Y. *et al.* Preclinical evaluation of a potent novel DNA-dependent protein

kinase inhibitor NU7441. *Cancer Res.* **66**, 5354–5362 (2006).

29.   Chan, D. W. *et al.* Autophosphorylation of the DNA-dependent protein kinase

catalytic subunit is required for rejoining of DNA double-strand breaks. *Genes*

Dev. **16**, 2333–2338 (2002).

30.   Attardi, L. D., De Vries, A. & Jacks, T. Activation of the p53-dependent G1

checkpoint response in mouse embryo fibroblasts depends on the specific DNA

532   damage inducer. *Oncogene* **23**, 973–980 (2004).

533 31. Agarwal, M. L., Agarwal, A., Taylor, W. R. & Stark, G. R. p53 controls both the

534   G2/M and the G1 cell cycle checkpoints and mediates reversible growth arrest in

535   human fibroblasts. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 8493–8497 (1995).

536 32. Pawlik, T. M. & Keyomarsi, K. Role of cell cycle in mediating sensitivity to

537   radiotherapy. *Int. J. Radiat. Oncol.* **59**, 928–942 (2004).

538 33. Feng, W. *et al.* Genetic determinants of cellular addiction to DNA polymerase

539   theta. *Nat. Commun.* **10**, 1–13 (2019).

540 34. Wood, R. D. & Doublié, S. DNA polymerase θ (POLQ), double-strand break

541   repair, and cancer. *DNA Repair* **44**, 22–32 (2016).

542 35. Brinkman, E. K. *et al.* Kinetics and Fidelity of the Repair of Cas9-Induced Double-

543   Strand DNA Breaks Molecular Cell Article Kinetics and Fidelity of the Repair of

544   Cas9-Induced Double-Strand DNA Breaks. *Mol. Cell* **70**, 801–813 (2018).

545 36. Brinkman, E. K., Chen, T., Amendola, M. & Van Steensel, B. Easy quantitative

546   assessment of genome editing by sequence trace decomposition. *Nucleic Acids*

547   *Res.* (2014). doi:10.1093/nar/gku936

548 37. Zhao, B. *et al.* The essential elements for the noncovalent association of two DNA

549   ends during NHEJ synapsis. *Nat. Commun.* **10**, (2019).

550 38. Bétous, R. *et al.* DNA replication stress triggers Rapid DNA Replication Fork

551   Breakage by Artemis and XPF. *PLoS Genet.* **14**, (2018).

552 39. Ma, Y., Pannicke, U., Schwarz, K. & Lieber, M. R. Hairpin opening and overhang

553   processing by an Artemis/DNA-dependent protein kinase complex in

nonhomologous end joining and V(D)J recombination. *Cell* **108**, 781–794 (2002).

40.   Löbrich, M. & Jeggo, P. A Process of Resection-Dependent Nonhomologous End

Joining Involving the Goddess Artemis. *Trends in Biochemical Sciences* **42**, 690–

701 (2017).

41.   Zhou, Y. & Paull, T. T. DNA-dependent protein kinase regulates DNA end

resection in concert with Mre11-Rad50-Nbs1 (MRN) and Ataxia Telangiectasia-

mutated (ATM). *J. Biol. Chem.* **288**, 37112–37125 (2013).

42.   Arnoult, N. *et al.* Regulation of DNA repair pathway choice in S and G2 phases by

the NHEJ inhibitor CYREN. *Nature* **549**, 548–552 (2017).

43.   Ceccaldi, R. *et al.* Homologous recombination-deficient tumors are hyper-

dependent on POLQ-mediated repair. *Nature* **518**, 258–262 (2015).

44.   Huarte, M. *et al.* A large intergenic noncoding RNA induced by p53 mediates

global gene repression in the p53 response. *Cell* **142**, 409–419 (2010).

45.   Baskar, R., Dai, J., Wenlong, N., Yeo, R. & Yeoh, K. W. Biological response of

cancer cells to radiation treatment. *Frontiers in Molecular Biosciences* **1**, 24

(2014).

46.   Higgins, G. S. & Boulton, S. J. Beyond PARP-POLθ as an anticancer target.

Science **359**, 1217–1218 (2018).

Figure 1

Figure 1 | p53-deficient cells exhibit radioresistance and accelerated resolution of DNA DSBs.

a, Diagram of growth competition assay. mCherry-labelled RPE1 cells were mixed with unlabeled *TP53*[-/-] RPE1 (1:1), exposed to IR and grown for 6 days. **b,** Relative abundance of unlabeled *TP53* [-/-] Clone#1 measured by Intellicyte high-throughput cytometry $\pm$ SEM (n=6) is shown, normalized to the untreated (0Gy) cohort at each time point. **c,** Relative abundance of unlabeled *TP53* [-/-] Clone#2 $\pm$ SEM (n=6) is shown, normalized to the untreated (0Gy) cohort at each time point. **d,** Representative immunofluorescence images of 53BP1 foci in cells with indicated genotypes untreated (no IR) or treated with IR (5Gy) and collected at .5, 2, and 4 h after irradiation. **e,** Quantification of 53BP1 foci. Data shown are mean (n=50 cells per treatment condition) $\pm$ SEM (n=3), and are consistent across two independent biological replicates. *$p < 0.05$; **$p < 0.01$; by two-tailed Student's t-test. **f,** Representative Neutral COMET fluorescence staining for DNA tails in cells with indicated genotypes treated with or without 5Gy IR. For irradiated cells, 2 timepoints are shown: immediately after and 4 hours post IR. COMET tails and heads are denoted by OpenComet software analysis. **g,** Quantification of DNA DSBs via Neutral COMET assay reported as tail DNA percent at 0 and 4 hours post IR in RPE1 and two *TP53*[-/-] RPE1 cell lines. Data shown are mean (n= 50-150 cells per treatment condition) $\pm$ SEM, and are consistent across three independent biological replicates. *$p < 0.05$; **$p < 0.01$; ****$p < 0.0001$ by two-tailed Student's t-test.

Figure 2

Figure 2 | Inhibition of DNA-PK restores DNA damage foci formation in p53-deficient cells.

a, Live cell imaging procedure. Cells transfected with 10 nM si-control or si-*TP53* for 48 h prior to imaging. 18 h into imaging, cells are treated with NCS (100 nM), DNA-PKi (.5 uM) or both and imaged for 72 total hours. **b,** RPE1 cell expressing the PCNA-mCherry and 53BP1-mVenus reporters. Cell cycle phases delineated by PCNA foci and DNA DSBs are marked by 53BP1 foci. **c,** RT-qPCR for *TP53* mRNA levels (left) and *CDKN1A* mRNA levels (right) in si-control treated vs. si-*TP53* treated cells. To induce *CDKN1A* expression, cells irradiated at 5Gy and mRNA harvested 3 hrs post IR. **d,** Heatmap of 53BP1 foci tracings for single cells tracked from birth to mitosis or end of imaging. For si-control (n = 30 cells) and si-*TP53* treated RPE1 (n = 60 cells) treated with NCS 100 ng/ml. For visualization, cells are aligned to 10 frames prior to drug addition (black arrow). **e,** Heatmap of 53BP1 foci tracings for si-control (n = 25 cells) and si-*TP53* treated RPE1 cells (n = 55 cells) treated with 100 ng/ml NCS + 0.5 uM DNA-PKi. **f,** Peak 53BP1 foci counts for cells treated with 100 ng/ml NCS or NCS+0.5 uM DNA-PKi. Significance determined using two-tailed t-test. **g,** Area under the curve (AUC) analysis of 53BP1 burden showing integral DNA damage for cells treated with NCS vs. NCS and DNA-PKi. Cells are segregated into two groups: cells exposed to drug in G1 vs. S phase (n = 25-30 G1or S cells for si-*TP53* cohort, n = 10-15 G1 or S cells for si-control cohort). Significance determined by two-tailed t-test. ****$p < 0.0001$, ***$p < 0.001$, n.s. = non-significant. **h,** 53BP1 foci burden in G1 vs. S phase p53-deficient RPE1 upon exposure to NCS and DNA-PKi. Dashed line = S phase onset, blue line = mean 53BP1 foci burden for all cells in G1 with NCS and DNA-PKi addition, orange line = mean foci value for cells in G1 with NCS treatment alone, (n = 30 cells for each condition).

Figure 3

Figure 3 | Checkpoint responses halt p53-proficient cells upon exposure to NCS while p53-deficient cells continue to cell cycle despite NCS exposure.
a, Schematic depicting NCS treatment (50 ng/ml + 100 ng/ml for si-control and 100 ng/ml for si-*TP53* RPE1) and/or NCS + 0.5 uM DNA-PKi treatment, and phase of the cell cycle cells are exposed to drug (G1). **b,** Schematic of drug treatment for S phase cells. **c,** Distribution of cell cycle phase lengths, each colored dot is an individual cell with untreated cells (no NCS) shown in black, NCS treated cells shown in blue, and NCS+ 0.5uM DNA-PKi treated cells shown in red for si-control RPE1 in G1 phase. n = 20 untreated and n = 30 treated cells (for each treatment cohort). Statistical significance was determined by comparing untreated and treated groups at each phase. ****$p < 0.0001$, n.s. = non-significant. Open circles indicate arrested cells that did not enter the subsequent phase of cell cycle for remainder of imaging. **d,** Distribution of cell cycle phase lengths for si-control treated RPE1 in S phase, ***$p < 0.001$, ****$p < 0.0001$, n.s. = non-significant as evaluated by two-tailed t-test. **e,** Distribution of cell cycle phase lengths for si-*TP53* treated RPE1 in G1 phase, ****$p < 0.0001$, n.s. = non-significant as evaluated by two-tailed t-test. **f,** Distribution of cell cycle phase lengths for si-*TP53* treated RPE1 in S phase, ****$p < 0.0001$, n.s. = non-significant as evaluated by two-tailed t-test.

Figure 4

Figure 4 | Inhibition of DNA-PK induces catastrophic mitoses in p53-deficient cells.
a, Cell cycle outcome analyses for si-control treated RPE1, dashed white line indicates drug addition, each row is an individual cell (n = 60 cells for NCS and n=60 cells for NCS+DNA-PKi treatment). Colored bars indicate different phases of the cell cycle, legend shown with no treatment control for comparison. Cells with red bars at the end of mitosis indicate terminal cell cycle event (mitotic catastrophe or apoptosis). Event frequency is reported as a percentage on the right. Cells exposed in G1 vs. S cells are treated as separate cohorts. Fisher's exact test was performed between -/+ DNA-PKi cohorts using 2 outcome groups (viable, vs. non-viable (arrested cells + terminal outcomes). ****$p < 0.0001$, n.s. =non-significant **b,** Cell cycle outcome analyses for si-*TP53* treated RPE1, dashed line indicates drug addition, each row is an individual cell (n = 60 cells for NCS and n=60 cells for NCS+DNA-PKi treatment). **c,** AUC analysis of 53BP1 damage burden in viable vs. non-viable p53-deficient cells that were treated with NCS and DNA-PKi. Statistical significance was calculated using a Mann-Whitney test comparing ranks. ****$p < 0.0001$ **d,** Dynamics of 53BP1 foci burden p53-deficient RPE1 segregated by mitotic viability. The red line corresponds to mean 53BP1 foci burden for all p53-deficient cells treated with NCS and DNA-PKi that undergo catastrophic mitoses, black line indicates mean foci value for p53-deficient cells with NCS and DNA-PKi treatment that are viable post mitosis, (n = 20 viable cells and n = 33 non-viable cells).

Figure 5

Figure 5 | p53-deficient cells utilize alternative end-joining pathways in the absence of active DNA-PK

a, RT-qPCR for *POLQ* mRNA levels in 2 *TP53*[-/-]RPE1 clones compared to WT RPE1. Significance was determined using two-tailed t-test. ****$p < 0.0001$, **$p < 0.01$. **b,** *POLQ* gene expression depicted as log2 values of *TP53* wild-type vs. mutant cancers across a subset of TCGA tumor types. Tumor labels follow TCGA labeling format. BRCA: breast cancer, BLCA: B-cell lymphoma, UCEC: uterine cancer, PRAD: Prostate cancer, PAAD: pancreatic cancer, SKCM: melanoma, LUSC: lung squamous cell cancer, LUAD: lung adenocarcinoma, GBM: glioblastoma multiforme, STAD: stomach cancer, and COADREAD: colorectal cancer. ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *p<0.05, as calculated by one-way ANOVA. **c,** Schematic depicting chromosomal break repair assay. *TP53*[-/-] , and *POLQ*[-/-]*TP53*[-/-] RPE1 are segregated into 2 cohorts (+/- 3 uM DNA-PKi). Cells are electroporated using Cas9-RNP-sgRNA-*LBR* and evaluated by next generation sequencing for break repair products at target locus. **d,** Horizontal bar chart representation of individual break repair products at *LBR* locus in *TP53*[-/-] RPE1 by NGS. Position 0 denotes *LBR* locus cut site, with left and right positions denoting final INDEL size and orientation. Results are reported as average with SEM of n=3 independent biological replicates. **e,** Histogram of overall frequency of repair of NHEJ, TMEJ, and Unclassified products in *TP53*[-/-] RPE1 with or without DNA-PKi treatment. **f,** Horizontal bar chart representation of individual break repair products at *LBR* locus in *POLQ*[-/-]*TP53*[-/-] RPE1 by NGS. Position 0 denotes *LBR* locus cut site, with left and right positions denoting final INDEL size and orientation. Results are reported as average with SEM of n=3 independent biological replicates. **g,** Histogram of overall frequency of repair of NHEJ, TMEJ, and Unclassified products in *TP53*[-/-] RPE1 with or without DNA-PKi treatment. **h,** Colony forming efficiency assay evaluating *TP53*[-/-] and *POLQ*[-/-]*TP53*[-/-] RPE1 after treatment with NCS (at 25 ng/ml, 50 ng/ml, and 100 ng/ml) with or without .5 uM DNA-PKi, data shown are mean +/- SEM (n= 3). Statistical significance assessed with student's two-tail test. ***$p < 0.001$, **$p < 0.01$, * p<0.05. in comparison to the survival curve of *TP53*[-/-] + DNA-PKi. **i,** Graphical summary

a

Cas9-RNP #1
sgTP53_Exon2:
TCGACGCTAGGATCTGACTG

Cas9-RNP #2
sgTP53_Dwnstream_Intron:
GAAACTGTGAGTGGATCCAT

36 nt gap

TP53 Exon 2 → TP53 Intron

Chromosome 17p13.1, *TP53* Locus

b

p53
(53kDa)

B-actin
(42 kDa)

RPE C1 C2 C3 C4 C5

c

Relative *CDKN1A* mRNA Levles

Untreated, C1, C2, C3, C4, C5, Untreated + 5Gy, C1 + 5Gy, C2 + 5Gy, C3 + 5Gy, C4 + 5Gy, C5 + 5Gy

d

RPE1 : RPE1^mCherry

Relative Abundance of mCherry Negative Cells

- 0 Gy
- 2 Gy
- 4 Gy
- 6 Gy

Days

e

Cell Survival (%)

- RPE1
- *TP53*⁻ᐟ⁻ RPE1

NCS (ng/ml)

**

f

Cell Survival (%)

- RPE1
- RPE1 + DNA-PKi

NCS (ng/ml)

**

g

Cell Survival (%)

- *TP53*⁻ᐟ⁻ RPE1
- *TP53*⁻ᐟ⁻ RPE1 + DNA-PKi

NCS (ng/ml)

*   **

h

RPE1

No NCS   NCS 50 ng/ml   NCS 100 ng/ml

+ 0.5 uM DNA-PKi

i

TP53⁻ᐟ⁻ RPE1

No NCS   NCS 50 ng/ml   NCS 100 ng/ml

+ 0.5 uM DNA-PKi

j

NO IR   t.5 hr   t2 hr   t4 hr

RPE1

TP53⁻ᐟ⁻ C#1

TP53⁻ᐟ⁻ C#2

10 µm

k

yH2AX Foci /Nucleus

- TP53-/- RPE1 C#2
- TP53-/- RPE1 C#1
- RPE1

t.5 hr   t2 hr   t4 hr

Supplementary Figure 1

Supplementary Figure 1 | a, Schematic of CRISPR target locus in human *TP53* gene. Two sgRNAs were designed to target sites in the terminal region of exon 2 (which encodes the p53 transactivating domain) and a site in the downstream intron with a 36 nucleotide (nt) gap. sgRNAs were complexed with Cas9 in the RNP system and electroporated into RPE1 cells. **b,** Western Blot of 5 selected single-cell clones that were profiled for p53 protein. **c,** Functional assay evaluating p53-dependent *CDKN1A* transcriptional responses to treatment of 5Gy IR. RNA from cells exposed to IR were harvested 6 hrs post treatment. **d,** Relative abundance of unlabeled RPE1^{unlabelled} over RPE1^{mCherry} measured by Intellicyte high-throughput cytometry +/- SEM (n=6) is shown, normalized to the untreated (0Gy) cohort at each time point. **e,** Clonogenic survival assays performed in RPE1 vs *TP53$^{-/-}$* RPE1 cells exposed to NCS. **f,** Clonogenic survival assays of RPE1 treated NCS +/- 0.5 uM DNA-PKi. Reported values are mean of n = 3 replicates, and survival fraction was calculated by first calculating plating efficiency and normalizing it to the untreated samples. **g,** *TP53$^{-/-}$* RPE1 cells were treated with NCS +/- 0.5 uM DNA-PKi. **h,i,** Representative colony forming plates for e and f at NCS doses of 0, 50 and 100 ng/ mL +/- 0.5 uM DNA-PKi. Cell numbers for each conditions plated are the following: UT (500), NCS 50 ng/ ml (2000), NCS 100 ng/ ml (6000). **j,** Representative immunofluorescence images of yH2AX foci in cells with indicated genotypes untreated (no IR) or treated with IR (5Gy) and collected at .5, 2, and 4 h after irradiation. **k,** Quantification of yH2AX foci. Data shown are mean (n= 50 cells per treatment condition) +/- SEM (n=3), and are consistent across two independent biological replicates. *$p < 0.05$ by two-tailed Student's t-test.

Supplementary Figure 2

Supplementary Figure 2 | a, Quantification of cell proliferation from live-cell imaging experiments for si-Control treated RPE1. Cell counts were normalized to cell numbers at start of imaging. Here we show one representative imaging beacon for each treatment condition (untreated, NCS 100 ng/ ml at 18 hours, and NCS 100 ng/ml + 0.5 uM DNA-PKi at 18 hours). **b,** Cell proliferation counts for si-*TP53* treated RPE1 over live-cell imaging. **c,** Analysis of RPE1 with no exposure to NCS (untreated) but received si-Control or si-*TP53* 3 days prior to imaging. No significant differences are seen due to si-treatment alone. **d,** Quantification of baseline mitotic outcomes with no NCS exposure over the course of imaging. DNA-PKi treatment alone with no NCS showed no little to no additional effect on cells (Chi-squared analysis P>0.05 for each condition).

Please refer to the following file names for the Supplementary Figure 3 Video Files:
Each video is a split screen of the same cell depicted in 2 channels: PCNA (left) and 53BP1 (right)
3a: NormalMitosis.mp4
3b: TransientG2Delay.mp4
3c: G1Arrest.mp4

Supplementary Figure 3 | a, Normal Mitosis: RPE1 Cell cycle representative of normal mitosis, with NCS treatment only. For all cells in this figure both the PCNA and the 53BP1 channels are shown as two individual movies. **b, Transient G2 Delay:** RPE1 cell cycle representative of a transient cell cycle delay in G2 (length of G2 is significantly prolonged in comparison to untreated cells). This cell was treated with NCS and DNA-PKi. **c, G1 Arrest:** RPE1 cell cycle representative of a permanent G1 arrest. This is a p53 proficient cell treated with NCS and DNA-PKi.

a Apoptosis

b Mitotic Catastrophe

c

p53-Deficient RPE1 treated with NCS and DNA-PKi

Supplementary Figure 4

Supplementary Figure 4 | a, Time stamped image sequence of apoptotic cell (PCNA channel shown). Cells that experienced nuclear degradation during cell cycle prior to mitosis were categorized as "apoptotic cells." In this sequence a cell in G2 experiences cell death at 27 hours post birth, with indication of mitotic attempt, with nuclear envelope collapse or presence of any daughter cells. **b,** Time stamped image sequence of cell that experienced mitotic catastrophe (PCNA channel shown). Cell undergoes nuclear envelope collapse (24:10), and attempts mitosis, in subsequent images fragmentation of nucleus is clearly visible with no viable daughter cells present. Cell non-viability during mitosis was defined as mitotic catastrophe. **c,** Integral DNA damage burden for p53-deficient cells treated with NCS (100 ng/ml) and DNA-PKi (.5 uM) are calculated and segregated by viable (black) vs. non-viable outcomes (red). Legend indicates which phase of cell cycle the cells are in during drug exposure, followed by the phase for which the burden is calculated. Ex: G1 cells G1 burden = cells in G1 during drug exposure and total damage burden in G1. Area under the curve (AUC) analysis was performed by plotting 53BP1 foci counts over time for each cell and integrating burden over time. Statistical significance was determined using two-tailed Student's t-test.

Supplementary Figure 5

Supplementary Figure 5 | a, Schematic of CRISPR target locus in human *POLQ* gene. Two sgRNAs were designed to target sites in the polymerase domain, with an 87 nucleotide (nt) gap. sgRNAs were complexed with Cas9 in RNP system and electroporated into RPE1 cells with a *TP53*[-/-] background to create double knockout cell line. **b,c** POLQ specific substrates were introduced into the *TP53* [-/-] vs. *POLQ*[-/-]*TP53*[-/-] DKO cells to assess repair efficiency. Products were amplified and characterized by electrophoresis and end joining efficiency was normalized to RPE1 with *POLQ* expression. **d,** Sanger sequencing analysis of CRISPR edited locus in *POLQ*[-/-] *TP53*[-/-] RPE1 clones**.** The locus of interest was PCR-amplified and cloned into a TOPO vector for sequencing analyses. Each line of sequence shown was derived from a different TOPO clone and aligned to show differences. The *POLQ*[-/-]*TP53*[-/-] clone has 87bp deletion resulting in frameshift mutations. Red boxes indicated sgRNAs used for the CRIPSR. **e,** Schematic showing evaluation of NGS samples by TIDE analysis for efficiency of cleavage at *LBR* target site across cell lines. **f,** RPE1, *TP53*[-/-], and *POLQ*[-/-]*TP53*[-/-] RPE1 cutting efficiency, all three cell lines have comparable levels of cutting efficiency with sgLBR. **g,** Schematic depicting digital PCR method for assessing LBR detection rate. **h,** Results of digital PCR assay on *TP53* [-/-] vs. *POLQ*[-/-]*TP53*[-/-] cells with -/+RNP and subsequently -/+ DNA-PKi (3uM). Average of 3 independent biological replicates are shown with SEM. Statistical significance was calculated using Multiple-t tests. *p<0.05, **p<0.01.