Inferring the infant pain experience: a translational fMRI-based signature approach

Eugene P. Duff [a,b], Fiona Moultrie [a], Sezgi Goksan [c], Marianne van der Vaart [a], Alexandra Abos [d], Sean P. Fitzgibbon [b], Luke Baxter [a], Tor D. Wager [e], Rebeccah Slater [a,b]

[a]Department of Paediatrics, University of Oxford
[b]Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neurosciences, University of Oxford
[c]Institute of Education, University College London, London
[d]University of Barcelona, Barcelona, Spain;
[e] Psychological and Brain Sciences, Dartmouth College, & Psychology and Neurosciences, Univ. of Colorado Boulder Dept. of Psychology and Neuroscience

Corresponding Author:
Eugene Duff
Paediatric Neuroimaging Group
Department of Paediatrics,
John Radcliffe Hospital, OX39DU
University of Oxford

Abstract

An infant's lack of verbal communication makes it challenging to infer their sensory and emotional experiences. Improving our interpretation of their perceptions will enhance our understanding of human brain development and improve the clinical care of infants. Multivariate neuroimaging signatures have been validated and developed in adults, which capture aspects of the human pain experience. Here we translate these fMRI signatures to infants in order to draw inferences about their experience of nociceptive events. We find that the Neurologic Pain Signature (NPS), which reflects the nociceptive and intensity encoding aspects of pain experience, is consistently activated in a cohort of adults (n=10) and in two cohorts of infants (n=15 & n=22). However, evidence of higher-level cognitive modulation of the nociceptive input, characterised by the Stimulus Intensity Independent Pain Signature (SIIPS1), is

34 observed in adults but not observed in infants. This study expands the use of brain
35 signatures to infer and deconstruct the experience of pain in non-verbal populations
36 and provides a framework to better understand the early development of human
37 sensory and emotional experience.

Introduction

In the absence of verbal communication, the perceptions of infants are enigmatic.
Behavioural cues such as grimacing and vocalisation are relied upon by caregivers
and clinicians to make inferences regarding an infant's experience of pain [1–5], which
are broadly based on experiences of older children and adults capable of verbal
report. Nevertheless, behaviours have limited sensitivity and specificity and their
interpretation is subjective. Behavioural responses attributed to pain can be elicited
by non-noxious stimuli in premature infants[6] and infants can have high behavioural
pain scores in response to innocuous procedures such as the changing of a nappy[7,8].
While pain scoring systems such as the Premature Infant Pain Profile-Revised (PIPP-
R) have some construct validity [9], the extent to which these scores track pain
perception is unknown. Given that pain perception manifests in the brain,
neuroimaging provides a proxy measure of neural activity that can be compared
between infants and adults in order to make valuable inferences regarding the
experience of pain in this non-verbal population.

Pain has no specific nexus in the human brain. Instead, a disperse mode of neural
activity across sensory, limbic and other areas, with complex temporal and stochastic
characteristics is thought to underlie an individual's subjective perception [10]. The
adult experience of pain involves complex cognitive and emotional processes, which
are influenced by anticipation, attention, memory, salience, attribution of causality,
fear and anxiety [11,12]. However, many of these modulatory factors may not influence
pain perception in the youngest infants. As the complexity of brain circuitry develops
in terms of both structural and functional connectivity, a richer experience of pain
will likely evolve [1,4]. Functional magnetic resonance imaging (fMRI) has been
extensively used to characterise the nociceptive system in adults, and in infants
noxious stimulus-evoked neural activity has been reliably detected [13,14]. Regional
analysis of adult and infant responses has previously revealed broadly similar
patterns of noxious-evoked activity [13], with infants displaying more bilateral
activation, which may be due to exuberant cortico-cortical and interhemispheric

connectivity of the immature brain. Additional brain regions are also activated in
infants, such as the auditory cortices, hippocampus and caudate, and more limited
activity is reported in some brain regions such as the amygdala and orbitofrontal
cortex. However, inferring the perceptual consequences of functional activity or
inactivity within individual anatomical regions remains challenging.

Regionalised assessments of activity can be combined to make more robust
inferences from data. Multivariate spatial patterns of fMRI activity have been
identified that can predict specific aspects of a reported experience [15–17]. Recently, a
series of studies have characterised fMRI response signatures associated with pain
and negative affect [18–21]. The primary pain template was the Neurologic Pain
signature (NPS), which predicts reported pain intensity [19]. The NPS was trained on
responses to varying levels of heat pain, and has subsequently been shown to
predict variations in pain in over 40 independent published cohorts [15,18,19,22]
including noxious thermal [19,23] electrical [24] and mechanical [24] stimuli. This signature is
sensitive to nociceptive pain and does not correspond to activity evoked by non-
noxious control stimuli[19], vicarious pain[24], threat[24], negative affect[25] or social pain[19].
A related signature, the Stimulus Intensity Independent Pain Signature (SIIPS1), was
subsequently developed to explain additional variation in reported pain beyond the
NPS and stimulus intensity, relating to modulatory factors such as expectation,
motivation, context and perceived control [26,27].

Verbal report is the gold standard measure of pain in adults, but as infants cannot
use language to describe their pain, there is great value in identifying alternative
objective surrogate measures that can be used to quantify pain experience in
infants. Here, following the application of graded acute nociceptive stimuli to infants
and adults, we project functional neuroimaging signatures of pain onto infant
noxious-evoked brain activity to detect and quantify neural patterns of activity
underpinning pain experience. We determine whether infant fMRI responses express
several adult pain-related signatures and how their expression across stimulus
intensities compares to signature expression in adults. This work provides the proof
of concept for an approach to building more specific infant brain-derived biomarkers
of pain.

Results

fMRI brain-derived signatures predict reported pain in adults

We studied a single cohort of adults (Prisma scanner: n=10; age range 23-36 years) and two cohorts of healthy newborn infants within 10 days of birth (Verio scanner: n=15; Prisma scanner: n=22). Acute noxious experimental punctate stimuli of various intensities were applied to the foot of each participant (applied force = 64 and 128 mN (infants); 64, 128, 256 and 512 mN (adults); 10 trials per force, Fig 1A) and the evoked brain activity was recorded using fMRI. Both adult and infant response parameter maps were registered to the Montreal Neurological Institute (MNI) template space for the application of signatures (Fig 1B). The punctate stimuli were reported in adults to evoke a low to moderate pain sensation that adults most frequently described as pricking (n = 8 of 10) and sharp (n = 6 of 10), and verbal pain scores increased with stimulus intensity (Spearman r=0.89; p < 0.001; Fig 1C)[13].

For each adult participant, 6 brain signatures were applied to the fMRI response data and cosine similarity was used to measure the presence of the signature activity in response to the nociceptive input. The signatures included the Neurologic Pain Signature (NPS) and the Stimulus Intensity Independent Pain Signature (SIIPS1), and 4 control signatures, of which 3 correspond to aspects of negative affect (the vicarious pain (VPS) [24]; picture induced negative affect (PINES) [25]; and social rejection (SR) [21]) and one which measures global signal (GS). The NPS and SIIPS1 showed robust expression in the adults, whereas the 4 negative control signatures were not significantly expressed, confirming that the nociceptive responses specifically expressed pain-predictive signatures (Fig 1D). The NPS signature tracked the reported pain intensity scores (r = 0.61, p<0.001), while the SIIPS1 did not show a significant association with the reports (r=0.14, p=0.37).

A. Experimental design

B. Registration of infant and adult templates with NPS

Adults

Infants

C. Adult out-of-scanner pain ratings

D. Template pain responses across signatures in adults

Fig 1. A. Experimental design: stimuli were applied with a minimum inter-stimulus interval of 25s across a range of intensities in a randomised order. 4 stimulus intensities were used in adults and the lowest 2 stimulus intensities were used in infants. B. NPS signature overlaid on registered adult and infant templates, with both templates transformed into MNI template space[28]. C. Adult Cohort: the intensity of the verbal pain report (Visual Analogue Score (VAS)) significantly increased with increasing stimulus intensity (r=0.89; p < 0.001) D. Adult cohort: quantification of the presence of six signatures in response to the nociceptive input. Only the NPS and SIIPS1 show significant expression. * indicates that cosine similarity is significantly greater than zero.

Correspondence between adults and infant expression of the Neurologic Pain Signature

We considered the comparative expression of the NPS in the adult and infant participants. The NPS was consistently expressed in the adults (mean cosine similarity 0.01-0.063), and showed significant intensity encoding between 64 - 512 mN (p <.001, Fig. 2A), consistent with its correlation with verbal pain report. Similarly, in both infant data sets (Prisma and Verio scanner) there was strong evidence of the presence of the NPS at 128 mN (p=0.001 Prisma; p<0.05 Verio, Fig 2A). Mean cosine similarity levels in the infants (at both 64 mN and 128 mN) were

similar to adult values. With the reduced number of force levels, intensity coding (i.e.
a difference between 64 mN and 128 mN responses) was not significant. Overall,
these results suggest that the neural activity patterns that are indicative of intensity-
modulated pain in adults (evidenced by verbal report) are also present in infants.

Subregion analysis of NPS responses in adults and infants

The NPS signature extends across numerous brain regions and the strength of the
expression of the pain signature may vary across different spatial areas, particularly
in the immature brain[26,29]. At a gross anatomical level the NPS can be divided into a
"pro-pain" component, where the signature has positive weights (indicating a
positive relationship with pain report) in regions such as the insular cortex,
secondary somatosensory cortex, and anterior cingulate cortex (ACC), and an "anti-
pain" component, where the signature has negative weights (indicating a negative
relationship with pain report) in regions such as the inferior parietal lobule and
lateral occipital cortex[29]. These components can be further divided into spatially
contiguous subregions.

In the "pro-pain" NPS brain component both adults and infants showed strong
intensity-dependent activation (Fig 2B, red lines), with infants having greater average
expression compared to adults in response to the same intensity of stimulation.
"Pro-pain" subregions identified in adults also showed positive responses in infants
(Fig 2C, red bars), and activity generally increased with stimulus intensity (Fig 2D). In
adults the "anti-pain" NPS brain component responses showed weak cosine
similarity with the signature (Fig 2B adults, blue line). Weaker responses in these
areas may be a result of the brief transient nature of the punctate stimulus
compared to other experimental noxious stimuli. Infants showed similarly negligible
cosine similarity at 64mN intensity, and negative cosine similarity at 128mN.

When considering the "anti-pain" NPS on a regional basis, a divergence between
infant and adult responses becomes clear.   In adults, many constituent NPS "anti-
pain" brain regions, including the posterior cingulate cortex, the inferior parietal
lobule and the lateral occipital cortex, showed significant expression of the NPS in
response to the stimuli (Fig 2C, 2D adults – blue bars). However, these brain regions
did not show positive expression in the infant cohort (Fig 2C, 2D infants – blue bars).

 Instead an inverted response relative to the NPS template was observed,

 corresponding to increased activation in response to the stimulus.

Overall, these results indicate that "pro-pain" NPS brain regions, primarily targeted

by ascending nociceptive afferents[19,30,31], show similar activation patterns in adults

and infants, whereas, "anti-pain" regions, which are not directly targeted by

nociceptive afferents, show divergent responses between the adult and infant.

Fig 2 A. Neurologic Pain Signature (NPS) responses to the brief experimental noxious

stimuli for adults and the two infant cohorts, measured by cosine similarity. All three

cohorts showed positive cosine similarities with evidence of intensity encoding. B.

Plots of the NPS separated into "pro-pain" and "anti-pain" components. C. NPS

subregion analysis showing mean subregion responses following the application of

128 mN force (infants and adults) and 512 mN force (adults only). Adults and infants

show concordance in "pro-pain" NPS subregions (red); only the adults show

concordance in "anti-pain" NPS subregions (blue) D. Subregion analysis across

intensities for selected subregions. Abbreviations: ipsi – ipsilateral; cont –

contralateral; Ins – insula; dplns – dorsal posterior insula; S2 –secondary

somatosensory cortex; dACC – dorsal Anterior Cingulate Cortex; cerebellum (vermis);

PCC – posterior cingulate cortex; IPL – inferior parietal lobule; LOC – lateral occipital

cortex. * indicates cosine similarity is significantly greater than zero.

The Stimulus Intensity Independent Pain Signature (SIIPS1) is not present in

infants

We next assessed the expression of the SIIPS1 in infants. The SIIPS1 is designed to

predict variability in subjective pain report that is not predicted by stimulus intensity

or the NPS, and is known to be driven by factors such as expectation, motivation,

context and perceived control. [26] The SIIPS1 was reliably expressed in adults (cosine

similarity: 0.023-0.062). As expected, it did not show significant intensity encoding

(Fig. 3A, adults). In contrast, noxious-evoked brain activity in infants did not

positively express the SIIPS1 signature (Fig. 3A, infants). The infant cohort imaged

with the Verio scanner showed a negative concordance at 128 mN.

SIIPS1 can also be divided into "pro-pain" and "anti-pain" components. [26] "Pro-pain"

components active in the adults include SIIPS1 specific patterns within the insula and

central operculum, and have been reported to exhibit intensity encoding[26]. "Anti-

pain" subregions include areas of the somatomotor cortex, hippocampus and cuneus

and have been reported to not show intensity encoding.

In the "pro-pain" brain component both adults and infants showed a similar pattern

of SIIPS1 related activation (Fig 3B, red lines). Adults (p=0.006) and infants from the

Prisma cohort (p=0.04) showed significant intensity encoding (Verio cohort p=0.20).

A subregion decomposition of the SIIPS1 "pro-pain" component found those

subregions reflecting the signature in adults showed similar responses in infants (Fig

3C, red bars).

While the "anti-pain" SIIPS1 component was consistently present in adult responses

a positive expression was not seen in infants (Fig. 3B). A significant negative

contribution of SIIPS1 was observed when a force of 128 mN was applied in both

cohorts of infants. This divergence in the adult and infant responses to the

nociceptive input was observed across all subregions that responded in adults, and

follows a similar pattern to that observed with the NPS (Fig. 3C and Fig 3D).

Fig 3. A. SIIPS1 signatures responses across stimulus intensity levels for the adult cohort and two infant cohorts. Adults show consistent positive concordance with the SIIPS1, while infants do not. B. Plots of SIIPS1 separated into reliable "pro-pain" and "anti-pain" subcomponents. C. SIIPS1 subregion analysis showing mean subregion responses following the application of 128 mN force (infants and adults) and 512 mN force (adults only). Consistent with the NPS, both adults and infants show concordance in "pro-pain" NPS subregions (red), but only adults show concordance in "anti-pain" SIIPS subregions (blue) D. Subregion analysis across intensities for selected subregions. Adults show some evidence of intensity encoding in "pro-pain" subregions. ipsi – ipsilateral; cont – contralateral; Thal – thalamus; midIns – middle insula; dpIns - dorsal posterior insula; aINS – anterior insula; ; COp – central operculum; Cb – cerebellum; SN - substantia nigra; SMC – somatomotor cortex; Precen – precentral gyrus; Cun – Cuneus; STG – superior temporal gyrus ; HC – hippocampus; LG – lingual gyrus; SPL - superior parietal lobule; MOG - middle occipital gyrus. * indicates cosine similarity is significantly greater than zero.

Discussion

Human language provides an exquisite means by which subtle sensory and affective states can be communicated. However, in non-verbal populations, such as newborn infants, inference about sensory and emotional experiences are guided by behavioural and physiological observations. For the quantification and treatment of infant pain, this approach is relied upon in clinical practice but these measures lack specificity and the inferences that can be drawn are subjective. How behaviours relate to a conscious experience of pain is ultimately unknown. Neuroimaging provides more direct insight into the neural processes underlying infant experience[32,33] and has the potential to improve the inferences we make about pain in nonverbal populations. Multivariate fMRI signatures aim to robustly link features of brain activity to specific experiences [15,19], and a key potential application of these signatures is to make valuable inferences in challenging patient populations with limited communication.

In infants, noxious-evoked behaviour and brain activity have been studied and correlated[34–36]. However, determining a quantitative association between behaviour, or brain activity, and perception has remained elusive. Here we translate whole-brain fMRI signatures that predict aspects of pain perception in adults to the newborn infant brain, in order to advance our understanding of how infants experience noxious events in early life. Two pain-related brain signatures (NPS and SIIPS1) that represent different aspects of the adult pain experience were applied to two independent cohorts of infants. The NPS is a multivariate template that tracks reported pain intensity associated with a variety of stimuli[19,23,24] and is robust to potential confounds such as non-physical pain and other salient aversive and sensory events [19,24,25]. The SIIPS1 signature aims to quantify further cerebral contributions to reported pain experience, beyond features of brain activity that directly relate to nociceptive input and intensity encoding [26]. Unlike the NPS, it is found to explain the effects of psychological factors on reported pain, such as expectation and control. These signatures identify combinations of regional activity rather than activity in single brain subregions, which allows for more accurate and specific prediction of pain reports [17,26].

Although caution is advised when translating and interpreting adult signatures in the immature brain[37], there is considerable evidence to support the validity of this approach. Core structural features of the mature adult brain are developed in term-aged infants[38,39]. We see functional responses[40–43], connectivity patterns, and resting

state activity [44,45] exhibit similar features to adult brain activity. Furthermore, strong

neural activity overlap between adult and infant sensory-evoked and noxious-evoked

brain activity patterns have been reported across a variety of brain imaging

modalities [13,36,46].

Our results provide quantitative evidence that noxious stimulation in infants evokes

changes in brain activity that would be attributed to the sensory discriminative

aspects of pain experience in an adult. We found the NPS to be expressed similarly

across infants and adults. This result aligns with a recent EEG study comparing pain

responses of adults and infants[36]. Here, infants displayed noxious evoked potentials

and gamma oscillations, understood in adults to be markers of primary nociceptive

processing and subjective pain[47]. In contrast to the NPS results, we also show that

endogenous cerebral contributions to pain reports that are observed in adults but

indirectly relate to the nociceptive input, are not significantly represented in infants.

In adults this activity, which is characterised by the SIIPS1, is reported to reflect

variation in expectation, motivation, context and perceived control [26]. The

observation that nociceptive input in newborn infants does not evoke patterns of

cerebral activity that relate to these complex psychological constructs may reflect a

lack of contextual awareness. As infants mature they become more engaged with

their external environment but it is not known at what developmental stage they are

capable of interpretation, judgement and modulation of their subjective experience

of pain.

A lack of concordance of infant brain activity with the SIIPS1 was primarily driven by

divergent responses in "anti-pain" brain regions, which occurs when a brain region

that is normally de-activated with increasing pain intensity is instead activated, for

example in fibromyalgia patients [29]. This divergence of responses in "anti-pain"

regions, including the cuneus, hippocampus, and areas of the precentral gyrus, was

also seen in the NPS signature, and was the most striking axis of differentiation

between infants and adults. A number of sources could potentially contribute to

this observation. It is possible that immaturities in the hemodynamic response in

infants produce altered BOLD responses in these areas. However, while differences

in BOLD mechanisms have been observed in infant rats[48], we used a hemodynamic

response function derived from somatosensory responses in newborn human

infants, where differences compared to adults appear more subtle[49]. Alternatively,

the differences in "anti-pain" responses may be underpinned by differences in the

noxious-evoked neural activity. Slower maturation of association and attentional

networks is well established in infants[50,51], and has been observed in resting state

fMRI[52]. Some differences between infant and adult EEG responses to nociceptive

stimuli have been reported, including delays in event-related potentials (ERPs), and a

late ERP component in infants corresponding to an increase in delta band energy[36].

The positive BOLD response association with the nociceptive stimuli in the "anti-

pain" regions of infants could be due to exuberant connections in the infant brain[50],

coupled with a potentially underdeveloped GABAergic inhibitory system[53]. However,

it is speculative to consider the exact mechanisms that underpin differences in the

evoked haemodynamic activity in the infant and adult brain[54].

The translational application of validated adult brain signatures to the immature

brain can be further expanded and refined. Applying these methods to a cohort of

young children who can reliably report their experiences would reveal the extent to

which these signatures are maintained once they appear during development.

Developmentally varying neural signatures could be characterised, with alternate

trajectories for infants born prematurely. Potential confounding factors could also be

further addressed. These signatures can be optimised for infants by including testing

and validation in contexts designed to replicate the reduced knowledge of the

experimental context of infants [55]. Despite its advantages, it is important to

appreciate that a single modality, such as fMRI, cannot capture the full complexity of

the processes that underlie a conscious experience. Therefore, we strongly advocate

a multimodal measurement approach to provide the best proxy of infant pain[56,57].

Infant behavioural pain scores have been extensively studied, and brain-derived EEG

signatures have been linked to infant nociception [34,58–63]. Linking these approaches

to validated fMRI signatures will enhance our confidence in these methods.

Given that infants are non-verbal, finding the best proxy to quantify their pain

experience is critical if we are to minimise pain exposure in clinical care and mitigate

the potential long-term and short-term impact of pain in early life[1,4,63]. Improving the

validity of inferences we make about infant perceptions will also advance our

understanding of human cognitive development. This study expands the use of brain

signatures to infer and deconstruct the experience of pain in non-verbal populations,

and provides a framework to better understand the early development of human

sensory and emotional experience.

Methods

This work extends previous work that used a subset of the data analysed here, which identified qualitative correspondences between adult and infant fMRI pain responses [64].

Experimental Data

All data were acquired at the FMRIB Centre, University of Oxford. Data were acquired in three studies. Informed written consent for the studies was provided by adult participants or by the infant's parent before the study commenced. The infants were studied under ethical approval given by the National Research Ethics Service. The adult study was approved by the University of Oxford Central University Research Ethics Committee. Both studies were carried out in accordance with the Declaration of Helsinki and Good Clinical Practice guidelines.

The adult study used a 3T Siemens Verio scanner with a 32-channel adult head coil to measure pain responses in 10 healthy adults (mean age = 28.3 years; range: 23–36; number of females: 5). BOLD images were acquired using a T2*-weighted EPI acquisition (sequence parameters: TR/TE = 3280/40ms; flip angle = 90°; FOV = 192×192 mm; imaging matrix 64×64; resolution 3×3×3 mm; slices = 50, collected in descending order; average total volumes = 142).

The second study, *Infants-Verio,* used the same protocol to study 15 healthy infants scanned within 7 days of birth (GA 38 - 43 weeks; number of females: 6) [13]. The infants were wrapped, fed and fitted with appropriate ear protection before being placed in the scanner. These infant data were acquired with the same sequence as the adults except using 33 slices, and an infant optimised TR/TE of 2500/40ms [65].

The final experiment studied 22 healthy term infants within 10 days of birth (GA 35-42 weeks; number of females: 12) using a Siemens Prisma 3T scanner with an adult 32 channel receive coil (Infants-Prisma). Infants were fed, wrapped and fitted with appropriate ear protection before being placed in the scanner. BOLD images were acquired using a T2* BOLD-weighted, GRE acquisition with EPI readout, 70° flip angle, TE= 50 ms[65], TR= 1,300 ms, mean TA= 6 mins (approx.), multiband 4 [66,67], 90 x 90 in-plane matrix size, 56 slices, 2 mm isotropic voxels.

Participants were scanned while receiving trains of 10 very brief non-tissue damaging punctate stimuli of a specific intensity, applied to the left foot by a trained

experimenter with a minimum inter-stimulus interval of 25s (Fig 1A). Participants

had scans with 64mN and 128mN force, and adults additionally had scans with

256mN and 512mN stimuli. Some subjects also had 32mN scans, which were not

studied here. The order of scans was randomised across subjects. The noxious

stimuli were time-locked to the fMRI recording using Neurobehavioural Systems

(Presentation, www.neurobs.com) software.

After the scanning session, application of the stimuli was repeated in the adults and

each participant was asked to lie down on a patient bed and verbally state a pain

score following application of each stimulus. Once all stimuli had been presented,

the participants were asked to describe the type of pain they experienced by

completing the McGill Pain Questionnaire[68].

Data Preprocessing

All infant FMRI data were processed using the DHCP functional and structural

pipelines[45,69,70]. Boundary-based-registration (BBR) to register functional data to

high-resolution structural data. Dynamic motion-by-susceptibility distortions were

corrected using FSL's EDDY[71,72], and ICA-based denoising was used to remove

additional artefactual signals [73,74]. 3mm spatial smoothing and 100s high-pass

temporal filtering was applied.

Time series modelling

The pain responses were modelled using a general linear model (GLM) in FEAT with a

regressor defined for responses at each stimulus intensity. The experimental design

(Fig 1A) was convolved with term-infant-specific optimal basis functions generated

by Arichi and colleagues [49]. Additional regressors modelling periods of excessive

motion-related signal as identified by DVARS [75] were also included. Parameter maps

were generated for each subject at each stimulus intensity and transformed into

MNI space.

The adult data were processed using FSL 6.0 [76]. BOLD data were high pass filtered

with a cut-off of 100s, and 5mm spatial smoothing was applied. FIX clean-up was

employed and motion parameters were regressed from the data [74].

Registration

Parameter estimate maps for individual subject and intensity levels were

transformed into adult MNI space via a series of transformations generated using

418	ANTs's SyN (Advanced Normalization Tools Symmetric image Normalization method)
419	[77]. Multi-modal non-linear registration was first used to register structural images T2
420	to the age-matched T2 template [78]. This was combined with the appropriate atlas
421	week-to-week nonlinear transforms to transform to 44 week template, and then to
422	MNI standard space with the NIHPD2 lifespan atlases (ages birth to 21 years) used as
423	intermediate transform targets [79,80]. For adults, FNIRT nonlinear registration was
424	used to transform data to MNI space [81].

Signature Analysis

426	Parameter map data derived from responses to specific intensities were analysed for
427	their correspondences to nociceptive signatures using a modified pipeline from the
428	CANlab toolbox [19,27,82](canlab.github.io), using additional MATLAB and Python code.
429	The CANLAB toolbox provides a pipeline for processing, QC and visualization of
430	parameter maps for the signature-based analysis of group fMRI data. This includes
431	the extraction, assessment, regression and of white matter and CSF components, the
432	calculation of signature similarity measures, estimation of contrasts and SVM
433	classifier maps, and NPS subregion analysis. Concordances between brain responses
434	and signatures are based on cosine-similarity scores – equivalent to correlation
435	without centring – between the response parameter maps and the predefined
436	spatial signatures. A cosine similarity of 1 represents perfect alignment of the data
437	with the signature and a cosine similarity of 0 indicates that the signature and the
438	data are orthogonal. To assess whether the signature responses tracked stimulus
439	intensity, we used a linear modelling across stimulus levels. We used the adult study
440	to validate the signature responses for our specific experimental protocol, to identify
441	responsive subregions for the infant analysis, and for direct comparison of adult and
442	infant responses.

443	The primary signature employed was the Neurologic Pain Signature (NPS) [19]. This
444	signature was developed using a regularised regression procedure that identified a
445	pattern of brain regions whose activity tracks pain ratings across trials of painful
446	thermal stimuli. The pattern has been validated across a variety of datasets, and has
447	been shown to exhibit specificity to pain relative to non-painful warm stimuli,
448	romantic rejection, and other emotions of negative affect[15,20]. Presence of the NPS in
449	infant responses to nociceptive stimuli, and any evidence of intensity encoding,
450	constitutes evidence for the experience of pain in infants. We also performed
451	analyses of subregions of the NPS and SIIPS1. The subregions were obtained using
452	the Canlab toolbox, and previously used in assessments of NPS [19] and SIIPS1 [26]. We

excluded regions that showed negligible or negative signature responses to our
nociceptive stimulus in the adult dataset (average cosine similarity <0.01).

We also assess the Stimulus Intensity Independent Pain signature (SIIPS1) [26], a
signature associated with factors affecting reported pain in adults independent of
stimulus intensity and the NPS. The SIIPS1 signature was trained using data from
which the NPS had been removed via regression. We performed the same procedure
here. If pain is processed in a similar fashion in adults and infants, we would expect
infant brain responses to painful stimuli to be concordant with the NPS. On the other
hand, correlation with the SIIPS1 signature might be expected to vary depending on
the level of development of the neurobiological systems captured (e.g.,
ventromedial prefrontal cortex and nucleus accumbens figure prominently) and
individual infants' capacity to engage these systems, which are associated with both
descending modulation[83,84] and self-regulation[27]. Therefore, in addition to the mean
response across subjects, we assessed the variability of responses across individuals,
measuring responses of individual trials to identify confidence intervals. We adapted
CANLAB code to include regression of the NPS prior to SIIPS1, and analysis of SIIPS1
subregions and single trial variability.

In addition, to determine the specificity of signature responses, we looked for the
presence of a number of other validated negative affect signatures, namely, picture
induced negative affect (PINES) [25]; signature of vicarious pain (VPS) [24], and social
rejection (SR) [21], and a marker of global signal (GS).

Bibliography

1. Anand, K. J. & Hickey, P. R. Pain and its effects in the human neonate and fetus.

The New England Journal of Medicine **317**, 1321–1329 (1987).

2. Grunau, R. V. & Craig, K. D. Pain expression in neonates: facial action and cry.

Pain **28**, 395–410 (1987).

3. Moultrie, F., Slater, R. & Hartley, C. Improving the treatment of infant pain. *Curr*

Opin Support Palliat Care **11**, 112–117 (2017).

4. Roofthooft, D. W. E., Simons, S. H. P., Anand, K. J. S., Tibboel, D. & van Dijk, M.

Eight years later, are we still hurting newborn infants? *Neonatology* **105**, 218–

226 (2014).

5. Stevens, B., Johnston, C., Petryshen, P. & Taddio, A. Premature {Infant} {Pain}

{Profile}: development and initial validation. *The Clinical Journal of Pain* **12**, 13–

22 (1996).

6. Green, G. *et al.* Behavioural discrimination of noxious stimuli in infants is

dependent on brain maturation. *Pain* **160**, 493–500 (2019).

7. Comaru, T. & Miura, E. Postural support improves distress and pain during diaper

change in preterm infants. *J Perinatol* **29**, 504–507 (2009).

8. Mörelius, E., Hellström-Westas, L., Carlén, C., Norman, E. & Nelson, N. Is a nappy

change stressful to neonates? *Early Hum. Dev.* **82**, 669–676 (2006).

9. Gibbins, S. *et al.* Validation of the Premature Infant Pain Profile-Revised (PIPP-R).

Early Human Development **90**, 189–193 (2014).

10. Baliki, M. N. & Apkarian, A. V. Nociception, pain, negative moods and behavior

selection. *Neuron* **87**, 474–491 (2015).

11. Wiech, K. Deconstructing the sensation of pain: The influence of cognitive

processes on pain perception. *Science* **354**, 584–587 (2016).

12. Legrain, V., Iannetti, G. D., Plaghki, L. & Mouraux, A. The pain matrix reloadedA

salience detection system for the body. *Progress in Neurobiology* **93**, 111–124

(2011).

13. Goksan, S. *et al.* {fMRI} reveals neural activity overlap between adult and infant

pain. *eLife* **4**, e06356 (2015).

14. Williams, G. *et al.* Functional magnetic resonance imaging can be used to explore

tactile and nociceptive processing in the infant brain. *Acta Paediatrica* **104**, 158–

166 (2015).

15. Kragel, P. A., Koban, L., Barrett, L. F. & Wager, T. D. Representation, Pattern

Information, and Brain Signatures: From Neurons to Neuroimaging. *Neuron* **99**,

257–273 (2018).

16. Mano, H. & Seymour, B. Pain: a distributed brain information network? *PLoS*

biology **13**, e1002037 (2015).

17. Norman, K. A., Polyn, S. M., Detre, G. J. & Haxby, J. V. Beyond mind-reading:

multi-voxel pattern analysis of fMRI data. *Trends in Cognitive Sciences* **10**, 424–

430 (2006).

18. Duff, E. P. *et al.* Learning to identify CNS drug action and efficacy using

multistudy fMRI data. *Science translational medicine* **7**, 274ra16 (2015).

19. Wager, T. D. *et al.* An fMRI-based neurologic signature of physical pain. *The New*

England journal of medicine **368**, 1388–97 (2013).

20. Woo, C.-W., Chang, L. J., Lindquist, M. A. & Wager, T. D. Building better

biomarkers: brain models in translational neuroimaging. *Nat. Neurosci.* **20**, 365–

377 (2017).

21. Woo, C.-W. *et al.* Separate neural representations for physical pain and social

rejection. *Nat Commun* **5**, 5380 (2014).

22. Zunhammer, M., Bingel, U., Wager, T. D. & Placebo Imaging Consortium. Placebo

Effects on the Neurologic Pain Signature: A Meta-analysis of Individual

Participant Functional Magnetic Resonance Imaging Data. *JAMA Neurol* **75**,

1321–1330 (2018).

23. Bräscher, A.-K., Becker, S., Hoeppli, M.-E. & Schweinhardt, P. Different Brain

Circuitries Mediating Controllable and Uncontrollable Pain. *J. Neurosci.* **36**, 5013–

5025 (2016).

24. Krishnan, A. *et al.* Somatic and vicarious pain are represented by dissociable

multivariate brain patterns. *eLife* **5**, e15166 (2016).

25. Chang, L. J., Gianaros, P. J., Manuck, S. B., Krishnan, A. & Wager, T. D. A Sensitive

and Specific Neural Signature for Picture-Induced Negative Affect. *PLOS Biology*

13, e1002180 (2015).

26. Woo, C.-W. *et al.* Quantifying cerebral contributions to pain beyond nociception.

Nat Commun **8**, 14211 (2017).

27. Woo, C.-W., Roy, M., Buhle, J. T. & Wager, T. D. Distinct brain systems mediate

the effects of nociceptive input and self-regulation on pain. *PLoS Biol.* **13**,

e1002036 (2015).

28. Evans, A. C. *et al.* 3D statistical neuroanatomical models from 305 MRI volumes.

in *1993 IEEE Conference Record Nuclear Science Symposium and Medical Imaging*

Conference 1813–1817 vol.3 (1993). doi:10.1109/NSSMIC.1993.373602.

29. López-Solà, M. *et al.* Towards a neurophysiological signature for fibromyalgia.

Pain **158**, 34–47 (2017).

30. Dum, R. P., Levinthal, D. J. & Strick, P. L. The Spinothalamic System Targets Motor

and Sensory Areas in the Cerebral Cortex of Monkeys. *J. Neurosci.* **29**, 14223–

14235 (2009).

31. Apkarian, A. V., Bushnell, M. C., Treede, R.-D. & Zubieta, J.-K. Human brain

mechanisms of pain perception and regulation in health and disease. *European*

Journal of Pain (London, England) **9**, 463–484 (2005).

32. Ellis, C. T. & Turk-Browne, N. B. Infant fMRI: A Model System for Cognitive

Neuroscience. *Trends in Cognitive Sciences* **22**, 375–387 (2018).

33. Cusack, R., Ball, G., Smyser, C. D. & Dehaene-Lambertz, G. A neural window on

the emergence of cognition. *Ann N Y Acad Sci* **1369**, 7–23 (2016).

34. Slater, R. *et al.* Evoked potentials generated by noxious stimulation in the human

infant brain. *European Journal of Pain (London, England)* **14**, 321–326 (2010).

35. Slater, R. *et al.* Cortical Pain Responses in Human Infants. *J. Neurosci.* **26**, 3662–

3666 (2006).

36. Fabrizi, L. *et al.* Encoding of mechanical nociception differs in the adult and infant

brain. *Scientific Reports* **6**, 28642 (2016).

37. Cusack, R., McCuaig, O. & Linke, A. C. Methodological challenges in the

comparison of infant fMRI across age groups. *Developmental Cognitive*

Neuroscience **33**, 194–205 (2018).

38. Kuklisova-Murgasova, M. *et al.* A dynamic 4D probabilistic atlas of the developing

brain. *NeuroImage* **54**, 2750–2763 (2011).

39. Zubiaurre-Elorza, L. *et al.* Auditory structural connectivity in preterm and healthy

term infants during the first postnatal year. *Developmental Psychobiology* **60**,

256–264 (2018).

40. Arichi, T. *et al.* Somatosensory cortical activation identified by functional MRI in

preterm and term infants. *NeuroImage* **49**, 2063–2071 (2010).

41. Perani, D. *et al.* Functional specializations for music processing in the human

newborn brain. *PNAS* **107**, 4758–4763 (2010).

42. Deen, B. *et al.* Organization of high-level visual cortex in human infants. *Nature*

Communications **8**, 13995 (2017).

43. Wild, C. J. *et al.* Adult-like processing of naturalistic sounds in auditory cortex by

3- and 9-month old infants. *NeuroImage* **157**, 623–634 (2017).

44. Doria, V. *et al.* Emergence of resting state networks in the preterm human brain.

Proceedings of the National Academy of Sciences **107**, 20015–20020 (2010).

45. Fitzgibbon, S. P. *et al.* The developing Human Connectome Project (dHCP)

automated resting-state functional processing framework for newborn infants.

bioRxiv 766030 (2019) doi:10.1101/766030.

46. Seghier, M. L., Lazeyras, F. & Huppi, P. S. Functional MRI of the newborn.

Seminars in Fetal and Neonatal Medicine **11**, 479–488 (2006).

47. Zhang, Z. G., Hu, L., Hung, Y. S., Mouraux, A. & Iannetti, G. D. Gamma-Band

Oscillations in the Primary Somatosensory Cortex—A Direct and Obligatory

Correlate of Subjective Pain Intensity. *J. Neurosci.* **32**, 7429–7438 (2012).

48. Kozberg, M. G., Chen, B. R., DeLeo, S. E., Bouchard, M. B. & Hillman, E. M. C.

Resolving the transition from negative to positive blood oxygen level-dependent

responses in the developing brain. *PNAS* **110**, 4380–4385 (2013).

49. Arichi, T. *et al.* Development of {BOLD} signal hemodynamic responses in the

human brain. *Neuroimage* **63**, 663–673 (2012).

50. Tau, G. Z. & Peterson, B. S. Normal Development of Brain Circuits.

Neuropsychopharmacol **35**, 147–168 (2010).

51. Kostovic, I. & Rakic, P. Developmental history of the transient subplate zone in

the visual and somatosensory cortex of the macaque monkey and human brain.

Journal of Comparative Neurology **297**, 441–470 (1990).

52. Fransson, P. *et al.* Early development of spatial patterns of power-law frequency

scaling in {FMRI} resting-state and {EEG} data in the newborn brain. *Cerebral*

Cortex (New York, N.Y.: 1991) **23**, 638–646 (2013).

53. Ben-Ari, Y. The GABA excitatory/inhibitory developmental sequence: A personal

journey. *Neuroscience* **279**, 187–219 (2014).

54. Colonnese, M. & Khazipov, R. Spontaneous activity in developing sensory

circuits: Implications for resting state fMRI. *NeuroImage* **62**, 2212–2221 (2012).

55. Hu, L. & Iannetti, G. D. Painful Issues in Pain Prediction. *Trends in Neurosciences*

39, 212–220 (2016).

56. Worley, A., Fabrizi, L., Boyd, S. & Slater, R. Multi-modal pain measurements in

infants. *J Neurosci Methods* **205**, 252–257 (2012).

57. Vaart, M. van der *et al.* Multimodal pain assessment improves discrimination

between noxious and non-noxious stimuli in infants. *Paediatric and Neonatal*

Pain **1**, 21–30 (2019).

58. Hartley, C. *et al.* The relationship between nociceptive brain activity, spinal reflex

withdrawal and behaviour in newborn infants. *Scientific Reports* **5**, 12519 (2015).

59. Hartley, C. *et al.* Nociceptive brain activity as a measure of analgesic efficacy in

infants. *Science Translational Medicine* **9**, (2017).

60. Fabrizi, L. *et al.* A {Shift} in {Sensory} {Processing} that {Enables} the {Developing}

{Human} {Brain} to {Discriminate} {Touch} from {Pain}. *Current Biology* **21**, 1552–

1558 (2011).

61. Ahola Kohut, S. & Pillai Riddell, R. Does the Neonatal Facial Coding System

Differentiate Between Infants Experiencing Pain-Related and Non-Pain-Related

Distress? *The Journal of Pain* **10**, 214–220 (2009).

62. DiLorenzo, M. G., Pillai Riddell, R., Flora, D. B. & Craig, K. D. Infant Clinical Pain

Assessment: Core Behavioral Cues. *The Journal of Pain* **19**, 1024–1032 (2018).

63. Slater, R., Cantarella, A., Franck, L., Meek, J. & Fitzgerald, M. How well do clinical

pain assessment tools reflect pain in infants? *PLoS medicine* **5**, e129 (2008).

64. Goksan, S. *et al.* The influence of the descending pain modulatory system on

infant pain-related brain activity. *eLife* **7**, e37125 (2018).

65. Goksan, S. *et al.* Optimal echo time for functional MRI of the infant brain

identified in response to noxious stimulation. *Magnetic Resonance in Medicine*

78, (2017).

66. Moeller, S. *et al.* Multiband multislice {GE}-{EPI} at 7 tesla, with 16-fold

acceleration using partial parallel imaging with application to high spatial and

temporal whole-brain {fMRI}. *Magnetic Resonance in Medicine* **63**, 1144–1153

(2010).

67. Xu, J. *et al.* Evaluation of slice accelerations using multiband echo planar imaging

at 3 {T}. *NeuroImage* **83**, 991–1001 (2013).

68. Melzack, R. & Torgerson, W. S. On the language of pain. *Anesthesiology* **34**, 50–

59 (1971).

69. Baxter, L. *et al.* Optimising neonatal fMRI data analysis: Design and validation of

an extended dHCP preprocessing pipeline to characterise noxious-evoked brain

activity in infants. *Neuroimage* **186**, 286–300 (2018).

70. Makropoulos, A. *et al.* The developing human connectome project: A minimal

processing pipeline for neonatal cortical surface reconstruction. *Neuroimage*

173, 88–112 (2018).

71. Andersson, J. L. R. *et al.* Towards a comprehensive framework for movement and

distortion correction of diffusion {MR} images: {Within} volume movement.

NeuroImage **152**, 450–466 (2017).

72. Andersson, J. L. R. & Sotiropoulos, S. N. An integrated approach to correction for

off-resonance effects and subject movement in diffusion {MR} imaging.

NeuroImage **125**, 1063–1078 (2016).

73. Griffanti, L. *et al.* {ICA}-based artefact removal and accelerated {fMRI} acquisition

for improved resting state network imaging. *NeuroImage* **95**, 232–247 (2014).

74. Salimi-Khorshidi, G., Smith, S. M., Keltner, J. R., Wager, T. D. & Nichols, T. E.

Meta-analysis of neuroimaging data: A comparison of image-based and

coordinate-based pooling of studies. *NeuroImage* **45**, 810–823 (2009).

75. Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L. & Petersen, S. E.

Spurious but systematic correlations in functional connectivity MRI networks

arise from subject motion. *NeuroImage* **59**, 2142–2154 (2012).

76. Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W. & Smith, S. M.

FSL. *NeuroImage* **62**, 782–90 (2012).

77. Avants, B. B., Epstein, C. L., Grossman, M. & Gee, J. C. Symmetric diffeomorphic

image registration with cross-correlation: evaluating automated labeling of

elderly and neurodegenerative brain. *Medical Image Analysis* **12**, 26–41 (2008).

78. Schuh, A. *et al.* A deformable model for the reconstruction of the neonatal

cortex. in *2017 IEEE 14th International Symposium on Biomedical Imaging (ISBI*

2017) 800–803 (2017). doi:10.1109/ISBI.2017.7950639.

79. Evans, A. C. The NIH MRI study of normal brain development. *NeuroImage* **30**,

184–202 (2006).

670 80. Fonov, V., Evans, A., Mckinstry, R., Almli, C. R. & Collins, L. Unbiased nonlinear

671   average age-appropriate brain templates from birth to adulthood. *Neuroimage*

672   **47**, (2009).

673 81. Andersson, J., Jenkinson, M. & Smith, S. M. Non-linear registration, aka spatial

674   normalisation. *FMRIB technical report TR07JA2* (2007).

675 82. Wager, T. D. *et al.* Pain in the ACC? *Proc Natl Acad Sci U S A* **113**, E2474–E2475

676   (2016).

677 83. Eippert, F. *et al.* Activation of the Opioidergic Descending Pain Control System

678   Underlies Placebo Analgesia. *Neuron* **63**, 533–543 (2009).

679 84. Leknes, S. & Tracey, I. A common neurobiology for pain and pleasure. *Nat. Rev.*

680   *Neurosci.* **9**, 314–320 (2008).

694