Integrated analysis of cervical squamous cell carcinoma
cohorts from three continents reveals conserved subtypes of
prognostic significance.

Ankur Chakravarthy[1,*], Ian Reddin[2,*], Stephen Henderson[3], Cindy Dong[4], Nerissa
Kirkwood[4], Maxmilan Jeyakumar[4], Daniela Rothschild Rodriguez[4], Natalia Gonzalez
Martinez[4], Jacqueline McDermott[5], Xiaoping Su[6], Nagayasau Egawa[7], Christina S
Fjeldbo[8], Vilde Eide Skingen[8], Mari Kyllesø Halle[10], Camilla Krakstad[10], Afschin
Soleiman[11], Susanne Sprung[12], Peter Ellis[4], Mark Wass[4], Martin Michaelis[4], Heidi
Lyng[8, 9], Heidi Fiegl[13], Helga Salvesen[10], Gareth Thomas[2], John Doorbar[7], Kerry
Chester[5, #], Andrew Feber[14, 15, #], Tim R Fenton[2, 4, #].

[1] Princess Margaret Cancer Centre, University Health Network. Toronto, Ontario,
Canada

[2] School of Cancer Sciences, Cancer Research UK Centre, Faculty of Medicine,
University of Southampton, Southampton, UK

[3] UCL Cancer Institute, Bill Lyons Informatics Centre, University College London,
London, UK

[4] School of Biosciences, Division of Natural Sciences, University of Kent,
Canterbury, UK

[5] UCL Cancer Institute, University College London, London, UK

[6] MD Anderson Cancer Center, Houston, Texas, USA

[7] Department of Pathology, University of Cambridge, Cambridge, UK

[8] Department of Radiation Biology, Oslo University Hospital, Oslo, Norway

[9] Department of Physics, University of Oslo, Oslo, Norway

[10] Department of Obstetrics and Gynaecology, Haukeland University Hospital,

Bergen, Norway; Centre for Cancer Biomarkers, Department of Clinical Science,

University of Bergen, Norway

[11] INNPATH, Institute of Pathology, Tirol Kliniken Innsbruck, Innsbruck, Austria.

[12] Institute of Pathology, Medical University of Innsbruck, Innsbruck, Austria.

[13] Department of Obstetrics and Gynaecology, Medical University of Innsbruck,

Innsbruck, Austria.

[14] Centre for Molecular Pathology, Royal Marsden Hospital Trust, London, UK

[15] Division of Surgery and Interventional Science, University College London,

London, UK

*Equal contribution

[#]Correspondence   to: k.chester@ucl.ac.uk;   a.feber@ucl.ac.uk   or

t.fenton@soton.ac.uk

Abstract

Human papillomavirus (HPV)-associated cervical cancer represents one of the

leading causes of cancer death worldwide. Although low-middle income countries

are disproportionately affected, our knowledge of the disease predominantly

originates from populations in high-income countries. Using the largest multi-omic

analysis of cervical squamous cell carcinoma (CSCC) to date, totalling 643 tumours

and representing patient populations from the USA, Europe and Sub-Saharan Africa,

we identify two CSCC subtypes (C1 and C2) with differing prognosis. C1 tumours

are largely HPV16-driven, display increased cytotoxic T-lymphocyte infiltration and

frequently harbour *PIK3CA* and *EP300* mutations. C2 tumours are associated with

shorter overall survival, are frequently driven by HPVs from the HPV18-containing

alpha-7 clade, harbour alterations in the Hippo signalling pathway and increased

expression of immune checkpoint genes, *B7-H3* (also known as *CD276*) and *NT5E*

(also known as *CD73*) and *PD-L2* (also known as *PDCD1LG2*). In conclusion, we

identify two novel, therapy-relevant CSCC subtypes that share the same defining

characteristics across three geographically diverse cohorts.

--

Despite screening and the introduction of prophylactic human papillomavirus (HPV)

vaccination in developed countries, cervical cancer continues to be one of the

leading worldwide causes of cancer-related deaths in women[1]. Prognosis for

patients with metastatic disease remains poor, thus new treatments and effective

molecular markers for patient stratification are urgently required. Cervical cancer is

caused by at least 14 high-risk human papillomaviruses (hrHPVs), with HPV16 and

HPV18 together accounting for over 70% of cases worldwide, with some variation by

region[1,2]. Cervical squamous cell carcinoma (CSCC) is the most common

histological subtype of cervical cancer, accounting for approximately 60-70% of

cases, again with some variation seen across different populations[2]. Adeno- and

adenosquamous histology are both associated with poor prognosis[3–6], while the

relationship, if any, between HPV type and cervical cancer prognosis remains

unclear[7]. HPV type is also associated with histology, with HPV16 more commonly

found in CSCC, while adenocarcinomas are more likely to harbour HPV18[2]. Previous

landmark studies described the genomic landscape of cervical cancer in different

populations[8–11] and in some cases identified subtypes based on gene expression,

DNA methylation and/or proteomic profiles[8,9]. The Cancer Genome Atlas (TCGA)

network identified clusters based on RNA, micro-RNA, protein/phospho-protein, DNA

copy number alterations and DNA methylation patterns and combined data from

multiple platforms to define integrated iClusters[8]. In their analysis, only clustering

based on the expression levels and/or phosphorylation state of 192 proteins as

measured by reverse-phase protein array (RPPA) was associated with outcome,

with significantly shorter overall survival (OS) observed for a cluster of cervical

cancers exhibiting increased expression of Yes-associated protein (YAP) and

features associated with epithelial-to-mesenchymal transition (EMT) and a reactive

tumour stroma. Since TCGA's RPPA analysis was restricted to 155 tumours

including SCCs, adeno- and adenosquamous carcinomas, we set out to test the

hypothesis that with data from more samples, we could identify a set of

transcriptional and epigenetic features associated with prognosis within CSCC and

to establish whether it is also present in independent patient cohorts representing

different geographical locations and ethnicities. To identify molecular subtypes and

prognostic correlates, we identified a set of 643 CSCCs (all HPV-positive), for which

clinico-pathological data and genome-wide DNA methylation profiles were either

publicly available or generated in this study, and for which in most cases, matched

gene expression and somatic mutation data were also available (Table 1).

Table 1: Summary of clinicopathological characteristics for five cervical cancer

cohorts.

		Discovery Cohort		Validation Cohorts			
		TCGA	Bergen	Innsbruck	Oslo	Uganda	Total
Cohort Numbers							
		236	37	28	248	94	643
Stage							
	I	122	33	16	23	15	209
	II	54	3	6	173	45	281
	III	39	0	4	61	31	135
	IV	14	1	2	13	2	32
	NA	7	0	0	0	1	8
Age							
	Median (Range)	47 (20-88)	42 (28-64)	49 (29-91)	54 (22-82)	45 (26-82)	
HPV Type							
	16	136	22	14	168	39	379
	18	26	2	5	30	14	77
	45	19	4	0	9	14	46
	Other	55	9	9	41	27	141
HPV Clade							
	Alpha 7	57	6	6	40	32	141
	Alpha 9	172	29	18	183	54	456
	Other	7	2	4	25	8	46
Treatment							
	Surgery alone	NA	18	5	0	NA	23
	Surgery and radiotherapy	NA	14	17	0	NA	31
	Surgery, radiotherapy and chemotherapy	NA	5	6	0	NA	11
	Radiotherapy alone	NA	0	0	47	NA	47
	Radiotherapy and chemotherapy	NA	0	0	201	NA	201
Overall Survival							
	Median (Range)	1.9 (0-17.5)	8 (1.8-13.2)	9.8 (0.1-23.2)	4.2 (0.3-12.7)	1.1 (0-2.4)	
Survival Status							
	Alive	181	32	16	187	42	458
	Dead	55	5	12	61	52	185
Cluster Assignment							
	C1	175	32	24	198	69	498
	C2	61	5	4	50	25	145
	% C2	25.8	13.5	14.3	20.2	26.6	22.6
HIV Status							
	Positive	NA	NA	NA	NA	59	59
	Negative	NA	NA	NA	NA	35	35
Available Data							
	RNA-seq	236	37	NA	NA	94	367
	Methylation	236	37	28	248	94	643
	Mutation	236	37	NA	NA	94	367
	RPPA	137	NA	NA	NA	NA	137
	Microarray (Illumina HumanWG-6 v3)	NA	NA	NA	137	NA	137
	Microarray (Illumina HumanHT-12 v4)	NA	NA	NA	109	NA	109
	Copy number (methylation derived)	236	37	28	248	NA	549

Identification of two gene expression-based clusters in cervical squamous cell carcinoma

Molecular and clinical differences between cervical adeno/adenosquamous and CSCCs are well documented[12–14] and gene expression differences were apparent in multi-dimensional TSNE analysis based on the top 10% most variable genes of three previously published cervical cancer cohorts[8,9,11] with available RNA-seq data (Supplementary Fig. S1a-d). To examine molecular and clinical heterogeneity specifically within SCC we focused all subsequent analysis on a collection of confirmed HPV-positive CSCCs from the USA, Europe and Uganda, as shown in Table 1.

236 cervical SCCs profiled by TCGA were defined as our discovery cohort (Table 1, Supplementary Table S1) and consensus clustering was performed using the top 10% most variable genes (n=1377 genes, Supplementary Table S2). Consensus cluster membership heatmaps, delta area plot, consensus cumulative distribution function (CDF) and proportion of ambiguous clusters (PAC) indicated the optimal number of clusters was two (Fig. 1a, Supplementary Fig. 2), the larger of which (n=175) was designated C1 while the smaller cluster (n=61) was designated C2 (Supplementary Table S1). Modelling transcriptomic differences between these two clusters identified 938 differentially expressed genes (DEGs, FDR=0.01, FC > 2) (Fig. 1b, Supplementary Table S3). Tumours in C1 predominantly harbour HPV types from the HPV16-containing alpha-9 clade (150/175) while 38 of 61 C2 tumours contained HPV types from the HPV18-containing alpha-7 clade. C2 tumours were

13.3 times more likely to harbour alpha 7 HPVs than C1 tumours (p = 1.8 x 10^{-14},

Fishers Exact Test) (Fig. 1b).

Univariate analysis of 5-year overall survival (OS) revealed worse outcomes for

patients with C2 tumours (HR = 2.54, p = 0.001; Fig. 1c) and in Cox regression

including age, tumour stage and HPV type as covariates along with cluster

membership, only membership of the C2 cluster (HR = 2.44, p = 0.017 95% CI 1.18,

5.05) and a tumour stage of IV (HR versus stage I = 4.74, p<0.001, 95% CI 2.1,

10.7) were independent predictors of five-year OS (Table 2). The relationship

between cluster and OS is also clear when restricting the analysis to HPV16-

containing tumours in each cluster in both univariate analysis (HR = 3.39, p = 0.004;

Fig. 1d) and multivariate analysis, including age and tumour stage as covariates (HR

= 3.89, p = 0.003, 95% CI 1.57, 9.67; Supplementary Table S4).

Identification of C1 and C2 CSCCs and association with prognosis in

independent SCC cohorts

To further investigate the association between C1/C2 cluster membership and OS,

we assembled a combined validation cohort consisting of 313 CSCC patients treated

at three centres in Europe (Bergen (n = 37), Oslo (n = 248) and Innsbruck (n = 28)),

for which detailed clinical information were available and for which genome-wide

DNA methylation profiles from Illumina Infinium 450k arrays (the same platform used

by TCGA) were either available or generated in this study (Table 1). Since RNA-seq

data were not available for all European samples, cluster membership was assigned

using a support vector machine (SVM) classification model based on 129 CpG sites

(methylation variable positions, MVPs) at which methylation differed significantly

between tumours in C1 vs C2 clusters in the discovery cohort (Fig. 2a, b; mean

delta-Beta > 0.25, FDR < 0.01, Supplementary Table S5), 18 of which were located

within 12 genes differentially expressed between the clusters (Supplementary Fig.

S3). MVP and DEG signatures were also used to assign cluster membership to 94

CSCCs from a Ugandan cohort originally profiled by the Cancer Genome

Characterization Initiative (CGCI)[9], for which both DNA methylation and RNA-seq

data were available. C2 tumours from all cohorts clustered together using TSNE

analysis based on the MVP signature (Fig. 2c) and high concordance between DEG

and MVP-based cluster allocation was observed in all cohorts for which both gene

expression (RNA-seq for Uganda and Bergen or Illumina bead chip arrays for Oslo)

and DNA methylation data were available (Supplementary Fig. S4a, b). Single-

sample gene set enrichment analysis (ssGSEA) confirmed differential expression of

the signature genes in tumours classified as C1 or C2 using DNA methylation data

(Supplementary Fig. 4c). 59 of 313 (18.8%) tumours in the combined European

cohort (Fig. 2b, Supplementary Table S7) and 25 of 94 (26.6%) tumours in the

Ugandan cohort were classified as C2 (Supplementary Fig. S5a, Supplementary

Table S6). As in the discovery cohort, most C1 tumours from the European and

Ugandan cohorts harboured alpha-9 HPV types (260/325) while C2 tumours were

3.9 times more likely to harbour alpha-7 HPVs than C1 tumours ($p = 1.07 \times 10^{-6}$,

Fishers Exact Test) (Fig. 2b, Supplementary Fig. S5a). Interestingly 80% (20/25) of

Ugandan C2 patients were human immunodeficiency virus (HIV) positive, while only

56% (39/69) of C1 patients were HIV positive (Supplementary Fig. S5a).

Univariate analysis indicated lower 5-year OS in C2 tumours from the European

cohort (Fig. 2d) and Cox regression controlling for FIGO stage, age, HPV type and

treatment (surgery alone, surgery with radio-chemotherapy, surgery with

radiotherapy alone, radio-chemotherapy and radiotherapy alone) again identified C2

status but not HPV type to be an independent predictor of 5-year OS (HR = 2.54, p

=0.003, 95% CI 1.4, 4.7) along with tumour stage and inclusion of chemotherapy in

the treatment regimen (Table 2). As in the discovery cohort, a significant prognostic

difference was identified between the C1 and C2 subgroups when considering only

the HPV16-positive tumours (n = 204) in both univariate (Supplementary Fig. S5b)

and multivariate analyses (HR = 2.64, p = 0.02, 95% CI = 1.16, 6; Supplementary

Table S4). Interestingly the prognostic difference was even greater among 78

patients in the European cohort that did not receive chemotherapy (Supplementary

Fig. S5c; multivariate HR = 4.4, p = 0.005, 95% CI = 1.58, 12.3). At 94 patients, the

Ugandan cohort was underpowered for comparing survival between C1 and C2

tumours and survival rates in the Ugandan cohort were much lower than in the other

cohorts (Supplementary Fig. S5d), thus we did not attempt a combined survival

analysis including these patients. Taken together, the C1/C2 clusters identified in the

TCGA cohort (USA) are apparent in cohorts of CSCC patients from Europe and

Uganda and tumours can be accurately assigned to cluster using either gene

expression or DNA methylation profiles. C1/C2 cluster is an independent predictor of

5-year OS in both the TCGA (n = 236) and European (n = 313) cohorts and remains

so when only HPV16+ tumours are considered. There is no difference in the

breakdown of C1 and C2 tumours by stage (Supplementary Table S7).

Table 2 – Five-year survival analysis for all cohorts

	Univariate			Multivariate		
	Hazard Ratio	p Value	95% CI	Hazard Ratio	p Value	95% CI
TCGA						
	2.54	0.002	1.42, 4.56	2.44	0.02	1.18, 5.05
Bergen						
	5.28	0.07	0.87, 31.9	98.1	<0.001	8.41, 1145
Innsbruck						
	0	1	0, Inf	0	1	0, Inf
Oslo						
	1.74	0.07	0.96, 3.14	2.36	0.012	1.21, 4.62
Europe Combined						
	1.68	0.07	0.97, 2.90	2.54	0.003	1.40, 4.67
Uganda						
	NA	NA	NA	NA	NA	NA

Relationships between C1/C2 and clusters previously identified by TCGA

Of the 178 tumour samples that made up the core set in the TCGA's landmark study

into cervical cancer genomics/epigenomics[8], 140 CSCCs were present in our

discovery cohort of 236 (Supplementary Table S8). This enabled comparisons

between our gene expression-based cluster allocations and the subtypes defined by

TCGA (Fig. 3). TCGA analysis included integrated clustering using multiomics data

(three iClusters, two of which ('keratin-high' and 'keratin-low' were composed entirely

of CSCCs) and clustering based on transcriptomic data (three mRNA clusters).

There is considerable overlap between our C1 cluster and TCGA's mRNA C2 cluster

(84/106) and keratin-high iCluster (80/106), and between our C2 cluster and TCGA's

mRNA C3 cluster (19/34) and keratin-low iCluster (27/34). Neither the mRNA C3 nor

the keratin-low iCluster were associated with poor prognosis in TCGA's analysis and

given the increased expression of a subset of keratin genes (including *KRT7*, *KRT8*

and *KRT18*) in C2 tumours (Fig. 3), we decided against adopting the keratin-high /

keratin-low nomenclature for our clusters. We also examined the relationship

between our subtypes and three clusters defined by TCGA based on reverse phase

protein array (RPPA) data. Notably, 57% of C2 TCGA tumours with RPPA data

available belong to the EMT cluster compared with only 25% of C1 tumours (Fig. 3)

and, consistent with the proteomic classification, C2 tumours display higher EMT

mRNA expression scores, as defined by TCGA[8] than C1 tumours (Supplementary

Fig. S6). Although there is greater concordance between C2 and the TCGA EMT

cluster compared to C1, it is clearly distinct from the EMT cluster.

Genomic analyses of prognostic clusters

To investigate whether C1 and C2 tumours differ at the genomic level in addition to

the transcriptomic and epigenomic differences observed above, whole-exome data

was obtained for SCCs from three cohorts, TCGA[8], Bergen[11] and Uganda[9]. This

amounted to 367 samples, 29 of which were classed as hypermutated by standards

set by TCGA[8] (>600 mutations). The median tumour mutation burden (TMB) was

2.04/Mb for all tumour, 2.11/Mb for C1 tumours and 1.82/Mb for C2 tumours

(1.92/Mb, 1.94/Mb and 1.72/Mb respectively after removal of hypermutated

samples). We detected four mutation signatures for the combined cohorts

(Supplementary Fig. S7): as expected based on previous studies[8,9,11], COSMIC

signatures 2 and 13 (characterised by C>T transitions or C>G transversions

respectively at TpC sites attributed to cytosine deamination by APOBEC enzymes);

age-related COSMIC signature 1 (characterised by C>T transitions attributed to

spontaneous deamination of 5' methylated cytosine) and COSMIC signature 5, for

which the underlying mutational process is unknown[15]

(https://cancer.sanger.ac.uk/signatures/). The proportion of mutations attributable to

each signature did not vary between clusters (Fig. 4).

Having excluded the hypermutated samples, we next performed dNdScv analysis[16]

on each cohort, followed by p-value combination using sample size weighted

Fisher's method followed by FDR correction[17] to permit identification of significantly

mutated genes (SMGs) across the entire dataset. This combined approach, followed

by analysis of individual samples by cluster identified 34 SMGs (Fig. 4,

Supplementary Table S9), 21 of which (highlighted by [†]) have not previously been

identified as SMGs in cervical cancer[8,9,11]. Of the 34 SMGs, 21 were significantly

mutated in only C1 samples, two genes in only C2 samples, three genes in both C1

and C2 individual analysis, and eight genes were only significantly mutated when

both C1 and C2 clusters were analysed together (Fig. 4, Supplementary Table S9).

The frequency of mutations in SMGs that had been previously observed was

comparable between combined cohort and each respective SMG study

(Supplementary Table S10). Among the 21 genes that have not previously been

identified as significantly mutated in cervical cancer, six are SMGs in other SCCs,

including head and neck (*NOTCH1, JUB (*also known as *AJUBA), MLL2 (*also known

as *KMT2D), RB1, PIK3R1*)[18], oesophageal (*MLL2, NOTCH1, RB1*)[19] and lung SCC

(*NOTCH1, RB1, MLL2, CREBBP (*also known as *KAT3A*))[20]. Conversely, several

genes previously identified as SMGs in cervical cancer, including *TP53*, *ARID1A* and

TGFBR2 are significantly mutated in adenocarcinoma but not in CSCC[8,11].

Comparing somatic mutation rates in SMGs between clusters using binomial

regression identified *PIK3CA* (FDR = 0.001) and *EP300* (FDR = 0.046) mutations as

disproportionally more common in C1 tumours and *STK11* (FDR = 0.005) and *NF2*

(FDR = 0.045) as enriched in C2 tumours (Fig. 4). *STK11* is also under-expressed in

C2 tumours compared with C1 tumours (Supplementary Table S3).

C2 tumours display Hippo pathway alterations and increased YAP1 activity

Two SMGs from our analysis (*LATS1* and *NF2*) are core members of the HIPPO

signalling pathway, while SMGs *FAT1, JUB* and *STK11* are known regulators of

HIPPO signalling[21–23]. Mutations in *LATS1*, *FAT1, JUB, STK11* or *NF2* (the latter two

of which are significantly mutated specifically in C2 tumours, Fig. 4) result in aberrant

activation of the downstream transcription factor, yes1 associated transcriptional

regulator (YAP1)[24–28], the expression of which is also elevated at the mRNA level in

C2 tumours (Table S3).

We generated segmented copy number data for all tumours (combining TCGA and

European validation cohort samples for which the necessary data were available for

maximum statistical power), which identified 211 focal candidate copy number

alterations (CNAs) at FDR < 0.1. Following binomial regression, we identified five

discrete CNAs that differed in frequency between C1 and C2 clusters (Fig. 5a; FDR

< 0.1, log2 (Odds Ratio) > 1). All five were more prevalent in C2 tumours and

included 11q11 and 1q21.2 deletions and 6p22.1, 11q22.1 and 11q22.2 gains.

11q22.2 contains matrix metalloproteinase genes (MMPs) which are well known to

be involved in metastasis[29], but notably 11q22.1 contains the *YAP1* gene.

Furthermore, analysis of Reverse Phase Protein Assay (RPPA) data from TCGA

revealed significantly higher YAP1 protein expression in C2 tumours (Fig. 5b). We

confirmed that of the 137 TCGA cases for which RPPA data were available, cases

with *YAP1* amplification (8/37 C2 tumours and 6/100 C1 tumours) also showed

increased YAP1 mRNA and protein expression (Supplementary Fig. S8). In total 10

genes from a 22 gene signature that predicts HIPPO pathway activity in cancer[30] are

differentially expressed between C1 and C2 tumours (Supplementary Table S3).

Differences in the tumour immune microenvironment between C1 and C2

tumours.

The nature of the tumour immune microenvironment, particularly the abundance of

tumour infiltrating lymphocytes (TILs) is a strong prognostic factor in cervical

cancer[31–33]. We used DNA methylation data to compare the cellular composition of

TCGA tumours[34], observing differences in the proportions of multiple cell types

between the subgroups (Fig. 6a); most notably decreased CD8+ (cytotoxic T

lymphocyte (CTL)), and a marked elevation of neutrophil and CD56+ natural killer

(NK)-cells in C2 tumours. Repeating this method with the validation cohorts

produced results that were remarkably similar (Fig. 6b). Differences in the

proportions of cell types between C1 and C2 in the validation cohort mirrored those

in the TCGA cohort, decreased CTL, and elevated neutrophil, NK-cell and

endothelial cell levels were observed in C2 tumours. Importantly, this was not driven

by any single validation cohort, as individual cohorts displayed consistent patterns of

differences in the proportion of cell types between C1 and C2 tumours, especially

with regards to CTLs, neutrophils and NK-cells (Supplementary Fig. S9a-d). C2

tumours also exhibit markedly higher neutrophil:CTL ratios (Supplementary Fig. S9e,

f) and neutrophil:lymphocyte (CTL, B-cell and Treg) ratios (NLR, Supplementary Fig.

S10); established adverse prognostic factors in cervical cancer[35–37]. At 0.7, the NLR

in C1 tumours across all cohorts was less than half that observed in C2 tumours

(1.85).

Validation of MethylCIBERSORT cell estimates was performed for a subset of

samples from the Innsbruck cohort using CD8 (CTLs) and myeloperoxidase (MPO,

neutrophils) immunohistochemistry (IHC)-based scores from a pathologist blinded to

cluster designation (Supplementary Fig. S11a-c) and for CTLs in the Oslo cohort

samples using comparison of MethylCIBERSORT estimates to CD8 IHC-based

digital pathology scores (Supplementary Fig. S11d).

Also of potential significance regarding the tumour immune microenvironment, is the

presence of two immune checkpoint genes, *CD276* (also known as *B7-H3*) and

NT5E (also known as *CD73*) in the set of 938 signature DEGs that separate the

clusters (Table S3). Both *B7-H3* and *NT5E*, along with a third immune checkpoint

gene (*PD-L2*) are expressed at higher levels in C2 tumours (Supplementary Fig.

S12) and hypomethylation of two CpGs in the *NT5E* promoter is evident in C2

tumours (Supplementary Table S5). All three suppress T-cell activity[38–40] and *B7-H3*

expression has been linked to poor prognosis in cervical cancer[41,42].

Evidence for differences in stromal fibroblast phenotype between C1 and C2

tumours

Gene set enrichment analysis using Metascape[43] suggested increased EMT (Supplementary Table S9) in C2 tumours, with 52 of 200 genes in in the EMT Hallmark gene set upregulated. As noted above there is also greater overlap between the C2 cluster and an EMT cluster defined by TCGA and based on RPPA data (Figure 3). Single-cell RNA sequencing and xenografting studies strongly suggest that rather than arising from the tumour cells (few of which have undergone EMT at any given time[44–46]), mesenchymal gene signatures in bulk tumour expression data instead derive from stromal cells including fibroblasts, which can adopt various phenotypes and play an important role in shaping the tumour immune microenvironment[47,48]. In addition to YAP1, which has been linked to the formation of cancer-associated fibroblasts (CAFs)[49] (as well as EMT[50–52] and angiogenesis[53]), C2 tumours display increased expression of the CAF marker genes *FAP* and *SERPINE1* (also known as *PAI-1*)[54]; the latter evidenced at both mRNA and protein levels (Supplementary Table S3, Fig. 5b). Overall fibroblast content as estimated by MethylCIBERSORT is similar between C1 and C2 tumours (Fig. 6a, b) but given recent findings regarding the extent and prognostic significance of CAF heterogeneity in the tumour microenvironment[55–59], we hypothesized that CAF phenotype rather than overall abundance, may differ between C1 and C2 tumours. To examine this, hierarchical clustering was performed based on the expression of eight gene sets (68 genes) curated by Qian et al[56], representing CAF-related biological processes and which are differentially expressed across six CAF phenotypes recently identified in a pan-cancer analysis[59]. C2 tumours cluster together, displaying increased expression of proinflammatory genes associated with an inflammatory (pan-iCAF2) CAF phenotype, C1 tumours appear more

heterogenous with respect to expression of the signature genes used to define CAF

phenotypes; there is upregulation of assorted myofibroblastic (myoCAF) genes in a

subgroup C1 tumours, including various collagens, ECM genes and TGFb-

associated genes, as well as 'contractile' genes such as smooth muscle actin

(*ACTA2*, Fig. 6c). While *ACTA2* is commonly used to identify myoCAF, it is also

expressed by pericytes and smooth muscle cells, which share the contractile

phenotype (and express for example, *MYH11*)[47,59,60]. Consistent with this, C2

tumours are 4.8x (p = 1.78 x 10^{-9}, Fisher's Exact Test) more likely to be classified as

'CAF-high' than C1 tumours using a four-gene CAF index defined by Ko et al[48].

Indeed, three of the four CAF index genes (*TGFBI*, *TGFB2* and *FN1*) appear in the

938 DEG signature that separates C2 from C1 tumours (Supplementary Table S3).

Discussion

In this study we hypothesized that by drawing upon several cervical cancer cohorts

for which 'omics data, clinical information and HPV typing were either available or for

which we were able to profile samples ourselves, we would be able to gain further

insight into CSCC – the most common histological cervical cancer subtype.

Clustering of CSCCs according to the 10% most variable genes identified two

clusters (C1 and C2) that bear resemblance to the keratin-high and keratin-low

iClusters originally defined by TCGA[8]. Cluster membership is an independent

predictor of 5-year OS and CSCCs can be accurately assigned to cluster using either

a 938 gene expression signature or a 129 MVP DNA methylation signature,

providing a means by which to gain prognostic information for cervical cancer

patients. While HPV16 and the alpha-9 clade to which it belongs have been

associated with longer PFS and OS in several studies[61–66], the relationship between

HPV genotype and cervical cancer prognosis remains unclear, as highlighted by a

recent meta-analysis[7]. In our multivariate analyses, membership of the C2 cluster

but not HPV type was an independent predictor of poor prognosis in both the

discovery and validation cohorts and remained so when only HPV16-positive

tumours in either cohort were considered. Possibly, the reason that HPV16 and other

alpha-9 HPV types have been associated with more favourable outcomes in certain

studies is that these viruses are more likely to cause C1-type tumours.

Adeno- and adenosquamous carcinomas, which are thought to arise from the

columnar epithelium of the endocervix, have been linked to poor prognosis in

cervical cancer [3–6] and to avoid differences due to histology, we focused our study

entirely on CSCC. Interestingly, of the 14 keratin genes that are differentially

expressed between C1 and C2 tumours, three (*KRT7*, *KRT8* and *KRT18*) that are

upregulated in C2 were classified as marker genes for columnar-like tumours with a

possible endocervical origin in a recent study that used single cell RNA-sequencing

and lineage tracing experiments to explore cell-of-origin for CSCC and

adenocarcinoma[67]. In contrast, C1 tumours display increased expression of *KRT5*, a

marker of the squamous-like subtype with a proposed ectocervical origin identified

by Chumduri et al[67] (Fig. 3). Other signature genes (*TP63, CERS3, CSTA, CLCA2,*

DSC3 and *DSG3*) upregulated in C1 tumours are also markers of the squamous-like

subtype, while further columnar-like marker genes (*MUC5B* and *RGL3*) are

upregulated in C2 tumours (Supplementary Table S3). Squamous-like tumours are

significantly enriched in the C1 sub-group, a C1 tumour is 4.9x more likely to be

squamous-like than columnar-like or unclassified (Fisher Exact Test, p = 0.0003).

426 This suggests that C2 tumours, although SCCs, harbour features associated with

427 adenocarcinoma; possibly even hinting at a different cell-of-origin for C1 versus C2

428 tumours. The greater frequency with which alpha 7 HPV types are found in C2 SCCs

429 is another feature shared with adenocarcinoma.

431 Our analysis suggests differences in the tumour immune microenvironment between

432 C1 and C2 CSCCs, that are highly reproducible across cohorts from the USA,

433 Europe and Uganda and that might explain the differential prognosis associated with

434 these clusters. In addition to the high neutrophil:lymphocyte ratio, the increased

435 expression of cytokines including IL-6, TGF-β and G-CSF and of the chemokines

436 CXCL1-3 in C2 tumours suggests pro-tumourigenic (N2) polarisation of these

437 neutrophils[68–73], which is typical of tumours with a high NLR[74]. The observation that

438 CSCCs occurring in HIV$^+$ patients from the Ugandan/CGCI cohort are much more

439 likely to be of the C2 subtype than those in HIV$^-$ patients hints at a possible

440 relationship between the immune competence of the patient and the likelihood of

441 developing a C2 tumour. This requires further investigation but is consistent with

442 greater evidence of existing anti-tumour immune responses in C1 tumours.

444 Finally, it is interesting to note that three targetable immune checkpoint proteins (B7-

445 H3, NT5E and PD-L2) are expressed at higher levels in C2 tumours. In addition to its

446 immune suppressive effects, B7-H3 has been linked to key processes that are

447 upregulated in these tumours including EMT and angiogenesis, through the

448 activation of NF-κB signalling and the downregulation of E-cadherin expression[75,76].

449 Interestingly, the expression of B7-H3 and NT5E on CAFs has been linked to poor

450 prognosis in gastric and colorectal cancer, respectively[40,77]. Also of relevance given

our observation of differing CAF phenotype between clusters is the report that a CAF

subtype (CAF-S1) identified in breast cancer that displays high levels of B7-H3 and

NT5E expression is seen in tumours with low levels of CTL infiltration[78]. PD1/PD-L1

immune checkpoint blockade (pembrolizumab) was recently FDA-approved for first-

line treatment of metastatic cervical cancer in combination with chemotherapy in

patients whose tumours express PD-L1[79,80], while CTLA4 blockade (Ipilimumab) has

also shown promising activity, both as a single agent[81,82] and in combination with

PD1 blockade (Nivolumab)[83]. Efficacy of PD1 blockade in cervical cancer has been

linked to the presence of a CD8+FoxP3+CD25+ T-cell subset[84] and an important

limitation of our study is the inability to differentiate between CD8+ T-cell

phenotypes. Nonetheless, identification of alternative, targetable immune checkpoint

molecules in C2 tumours provides a potential therapeutic strategy for a subset of

cervical cancers that respond poorly to chemoradiotherapy and that, given their low

overall levels of T-cell infiltrates, are maybe less likely to respond to PD1 blockade

than C1 tumours.

In conclusion, we show that CSCCs can be categorised in two novel tumour types,

C1 and C2, among which C1 tumours have a more favourable outcome. Although

HPV16 is more likely to cause C1 tumours and HPV18 C2 tumours, HPV type is not

an independent predictor of prognosis, suggesting it is the tumour type rather than

the causative HPV type that is critical for the disease outcome. Notably, the key

molecular and cellular characteristics of C1 and C2 tumours are consistent among

cohorts from the US, Europe, and Sub-Saharan Africa. This suggests that the

findings and underlying principle: that CSCC can develop along two trajectories

associated with differing clinical behaviour that can be identified using defined gene

expression or DNA methylation signatures, are of broad relevance.

Methods

Patient samples

All patients gave written, informed consent before inclusion. Samples from Bergen

were collected in a population-based setting from patients treated at the Department

of Obstetrics and Gynaecology, Haukeland University Hospital, Bergen, Norway,

from May 2001 to May 2011. The study has been approved by the regional ethical

committee (REK 2009/2315, 2014/1907 and 2018/591). For more details on sample

collection see[11,79]. Samples from Innsbruck were collected and processed at the

Department of Obstetrics and Gynaecology of the Medical University of Innsbruck.

The study was reviewed and approved by the Ethics committee of the Medical

University of Innsbruck (reference number: AN2016-0051 360/4.3; 374/5.4: 'Biobank

study: Validation of a DNA-methylation based signature in cervical cancer') and

conducted in accordance with the Declaration of Helsinki. Samples from Oslo (n =

268) were collected from patients participating in a previously published prospective

clinical study[80] approved by the Regional Committee for Medical Research Ethics in

southern Norway (REK no. S-01129). Limited quantities of patient tumour samples

and extracted DNA may remain and the distribution of these materials is subject to

ethical approval at the institutions from which they were collected. Note that the

cases in the Oslo cohort were not treated with surgery. The samples used for

molecular analysis were diagnostic biopsies from the primary tumour. In all other

cases, specimens were from resections of the primary tumour. Those interested in

working with these samples should contact the authors to discuss their requirements.

Dataset assembly

DNA methylation (Illumina Infinium 450k array) and RNAseq data were obtained for

CESC from the TCGA data portal. TCGA mutation data were obtained from the MC3

project on SAGE Synapse (syn7214402). RNAseq data for the Uganda cohort was

obtained from the TCGA data portal and DNA methylation (Illumina Infinium EPIC

array) and mutation data from National Cancer Institute's Genome Data Commons

Publication Page at https://gdc.cancer.gov/about-data/publications/CGCI-HTMCP-

CC-2020. DNA methylation (Illumina Infinium 450k array) and gene expression

(Illumina HumanHT-12 V4.0 expression beadchip) data from the Oslo cohort were

obtained from the Gene Expression Omnibus (GSE68339). RNAseq data were

obtained for the Bergen cohort from dbGaP (phs000600/DS-CA-MDS 'Genomic

Sequencing of Cervical Cancers') under the authorisation of project #14589

"Investigating the mechanisms by which viruses and carcinogens contribute to

cancer development" and were converted to fastq files using SRA-dump from the

SRA Toolkit (http://ncbi.github.io/sra-tools/). Kallisto[81] was then used to quantify

expression of GENCODE GrCh37 transcripts, repbase repeats and transcripts from

20 different high-risk HPV types with bias correction. Where IDAT files for 450k data

were available, they were parsed using *minfi*[82] and were subjected to Functional

Normalisation[83], followed by BMIQ-correction[84] for probe type distribution (which

was performed for all methylation data). For TCGA samples, viral type allocation was

performed using VirusSeq[85].

Only squamous cell carcinomas were considered in this study to avoid confounding

from histology. Multidimensional visualisation of the molecular differences in

histology was performed using Rtsne R package with parameters available in

Supplementary Table S12, and the top 10% most variable genes using mean

absolute deviation after pre filtering of low count genes (n = 1,385). Final cohort

numbers and summaries are shown in Table 1.

Generation of 450k methylation profiles

100ng DNA was bisulphite converted using the EZ DNA Methylation kit (Zymo

Research) as per manufacturer's instructions. Bisulphite converted DNA hybridised

to the Infinium 450K Human Methylation array and processed in accordance with the

manufacturer's recommendations.

HPV typing

HPV16 or 18 was detected in 208 samples from the Oslo cohort by PCR, using the

primers listed in86. The PCR products were detected by polyacrylamide gene

electrophoresis or the Agilent DNA 1000 kit (Agilent Technologies Inc, Germany).

Samples from the Innsbruck cohort and the remaining non-HPV16/18 samples from

the Oslo cohort (n=40) were HPV-typed by DDL Diagnostic Laboratory (Netherlands)

using the SPF10 assay, in which a PCR-based detection of over 50 HPV types is

followed by a genotyping assay (LIPA$_{25}$) that identifies 25 HPV types (HPV 6, 11, 16,

18, 31, 33, 34, 35, 39, 40, 42, 43, 44, 45, 51, 52, 53, 54, 56, 58, 59, 66, 68/73, 70

and 74). If more than one HPV type was identified in a sample (e.g. HPV16 and

HPV18), that sample was designated "Other" as HPV type in the study. HPV type

data for the remaining samples were published previously[8,9,11].

Prognostic analyses and tumour clustering

Unsupervised consensus clustering was performed on TCGA SCC samples using r

package ConsensusClusterPlus. After prefiltering of genes to remove those with low

read counts (75% samples read count < 1), only the top 10% most variable genes

using mean absolute deviation were considered for clustering (n = 1,385). 80% of

tumours were sampled over 1000 iterations using all genes. PAM clustering

algorithm was used and clustering distance was measured using Pearson's

correlation. An optimum number of clusters (K) of 2 was obtained by using the

proportion of ambiguously clustered pairs (PAC) using thresholds of 0.1 and 0.9 to

define the intermediate sub-interval. PAC was used as it accurately infers K[87].

Limma-voom on RNAseq data and limma on BMIQ and Functionally-normalised

450k and EPIC data were used to identify differentially expressed genes (DEGs,

FDR = 0.01, FC > 2) and methylation variable positions (MVPs, FDR = 0.01, mean

delta-Beta > 0.25) between the 2 clusters, C1 and C2. The 116 MVPs

(Supplementary Table S13) common to the 450k and EPIC arrays were used to

allocate clusters for the Ugandan cohort. The mean delta-Beta threshold for MVPs

was determined as it delivered the highest concordance between DEG and MVP

signature cluster allocation in the Bergen cohort (89.5%) and high concordance in

the Ugandan cohort (91.5%). The caret R package and limma were used to develop

an SVM using 5 iterations of 5-fold Cross-Validation using DEGs and MVPs to

allocate RNAseq samples in Ugandan and Bergen cohorts, 450k samples in Bergen,

Innsbruck and Oslo cohorts and EPIC samples in Ugandan cohort to these

subgroups. Multidimensional visualisation using R package Rtsne was performed on

the TCGA and European cohorts with available DNA methylation data combined

using the 129 MVPs and parameters as shown in Supplementary Table S12.

Samples from our validation cohort, comprise of cases from three European centres

(Bergen and Oslo in Norway and Innsbruck, Austria) and one African centre

(Uganda) were binned into these categories, and were used for subsequent

statistical analyses to identify genomic and microenvironmental correlates. Survival

analyses of epigenetic allocations were carried out using Cox Proportional Hazards

regression with age, tumour stage, HPV type, and with surgery, radiotherapy and

chemotherapy (given/not given) as covariates. R packages used were survival and

survminer. For all clinical analyses, stages were collapsed into Stages I, II, III and IV.

RNAseq data for Bergen and Ugandan samples, Illumina HumanWG-6 v3

microarray data for 137 of the Oslo samples and Illumina HumanHT-12 v4

microarray data for 109 of the Oslo samples were used to explore cluster allocation

concordance accuracy between DEG and MVP signature cluster allocation. ROC

curve and ssGSEA analysis were performed using R (scripts available at request).

Previous study comparison

140 TCGA samples from the core set analysis (TCGA, 2017) were present in our

TCGA SCC cohort. Previous cluster analysis by TCGA (2017) and Chumduri *et al.*

(2021) was compared with our C1 and C2 cluster allocation.

Pathway analyses

Pathway and gene sets were analysed with Metascape[43]. Settings used were

minimum gene set overlap of 10, p value cutoff of 0.01 and minimum enrichment of

1.5. All functional set, pathway, structural complex and miscellaneous gene sets

were included in the analysis. Only hits with an FDR of less than 0.05 were included

in final results.

Mutational analyses

For TCGA data, mutation calls were obtained from SAGE synapse as called by the

MC3 project. Mutations for the Bergen cohort were obtained from[11]. Ugandan

mutation calls were obtained from National Cancer Institute's Genome Data

Commons Publication Page at https://gdc.cancer.gov/about-data/publications/CGCI-

HTMCP-CC-2020. VCFs obtained for the Ugandan cohort samples were converted

to maf files using R package vcf2maf, filtered for whole-exome mutations only, and

combined. Significantly mutated genes (SMGs) were identified using dNdScv[16]

individually for the three cohorts. Hypermutated samples (>600 mutations[8]) were

excluded from this analysis. A weighted approach was used to combine p values for

each gene for the three cohorts. R package metapro[17] function wFisher was used to

perform this task. Genes were considered SMGs if after FDR correction of combined

p values, q < 0.1. Analysis was repeated for only C1 and C2 samples individually.

Two genes were removed from our list. *MUC4* was removed due to the large size of

the gene and *GOLGA6L18* was removed as this gene and it's aliases were not

recognised by R package maftools88.

R package maftools was used to produce an oncoplot for SMGs, calculating tumour

mutational burden for individual samples, SMG mutation frequency and mutational

signatures for the combined cohorts. Binomial GLMs were used to estimate

associations between C1 and C2 clusters and SMG mutation frequencies.

The estimated exposures of each sample to the identified mutational signatures were

calculated using R package mutsignatures89 and converted to proportion of

signature exposure per sample.

Copy number analysis

450k total intensities (Methylated and Unmethylated values) were used to generate

copy number profiles with normal blood samples from Renius et al90 as the germline

reference. Functional normalisation83 was used to regress out technical variation

across the reference and tumour datasets before merging and quantile normalisation

was used to normalise combined intensities followed by Circular Binary

Segmentation as previously described91. Median density peak correction was

performed to ensure centering before further analysis. GISTIC2.092 was then used

to identify regions of significant copy number change at both arm and gene levels.

Candidate copy number changes were evaluated for association with cluster using

binomial GLMs. The parameters chosen were a noise threshold of 0.1 with arm-level

peel off and a confidence level of 0.95 was used to nominate genes targeted by copy

number changes. Binomial regression was finally used to estimate rates of

differential alteration.

Reverse Phase Protein Assay analysis

Reverse Phase Protein Assay (RPPA) data for the core TCGA CESC samples were

obtained from the NCI GDC Legacy Archive. Differentially expressed proteins

between C1 and C2 clusters were determined using R package limma (FDR = 0.05,

FC > 1.3).

Tumour microenvironment analyses

MethylCIBERSORT[34] was used to estimate tumour purity and abundances of nine

other microenvironmental cellular fractions using TCGA and validation cohort

methylation beta values. Fraction numbers were then normalised by cellular

abundance and differences between clusters C1 and C2 were estimated using

Wilcoxon's rank sum test with Benjamini Hochberg correction for multiple testing.

This analysis was performed separately on TCGA cohort and combined validation

cohort, as well as on each individual cohort.

Cancer associated fibroblast associated gene set lists were obtained from Qian et

al[56]. TCGA, Bergen and Ugandan cohort sample RNAseq data was combined and

visualised for these gene set genes using R package NMF93.

CAF Index calculation

For cohorts that RNAseq data was available (TCGA, Bergen and Uganda), a CAF

index was calculated as described in Ko et al[48]. The median CAF index value was

used as a threshold to allocate high or low CAF in tumour samples.

Immunohistochemistry

Immunohistochemical staining of samples from the Innsbruck cohort was conducted

by HSL-Advanced Diagnostics (London, UK) using the Leica Bond III platform with

Leica Bond Polymer Refine detection as per manufacturer's recommendations.

Sections from a series of 17 tumour samples from the validation cohort were stained

for CD8 (mouse monoclonal 4B11, Leica Biosystems PA0183, used as supplied for

15 minutes at room temperature. HIER was performed on-board using Leica ER2

solution (high pH) for 20 minutes), CD68 (mouse monoclonal PGM1, Agilent

M087601-2, used at a dilution of 1/50 for 15mins at room temperature. HIER was

performed on-board using Leica ER1 solution (low pH) for 20 minutes) or MPO

(rabbit polyclonal, Agilent A039829-2, used at a dilution of 1/4000 for 15 minutes at

room temperature without epitope retrieval. Scoring was performed blinded to cluster

membership by a histopathologist (JM) as follows: 0 = no positive cells / field (200X

magnification); 1 = 1 − 10 positive cells; 2 = 11 − 100 positive cells; 3 = 101 − 200

positive cells; 4 = 201 = 300 positive cells; 5 = over 300 positive cells.

For the Oslo cohort, manual CD8 staining was conducted using the Dako

EnVision^{TM} Flex+ System (K8012, Dako). Deparaffinization and unmasking of epitopes

were performed using PT-Link (Dako) and EnVision^{TM} Flex target retrieval solution at a

high pH. The sections were incubated with CD8 mouse monoclonal antibody (clone

4B11, 1:150, 0.2 µg IgG$_{2b}$/ml) from Novocastra (Leica Microsystems, Newcastle Upon

Tyne, UK) for 45 minutes. All CD8 series included positive controls. Negative controls

included substitution of the monoclonal antibody with mouse myeloma protein of the

same subclass and concentration as the monoclonal antibody. All controls gave

satisfactory results. CD8 pathology scores were given to each sample (blinded to

cluster membership) for connective tissue only, tumour only and both as follows: 0 = no

positive: 1 = <10% CD8 positive cells; 2 = 10-25% CD8 positive cells; 3 = 25-50% CD8

positive cells; 4 = >50% CD8 positive cells. For digital quantification scanned images

of all sections at a high resolution of 0.46 um/pixel (20x), which was reduced to 0.92

um/pixel for analysis, were used. Digital score was calculated by quantifying the area

fraction of stained CD8 cells in relation to the entire section in the digital assessment.

703
704

Data availability

Illumina Infinium 450k array DNA methylation data generated in-house from Bergen

and Innsbruck validation cohort samples   have been deposited in the Gene

Expression Omnibus   (accession number GSEXXXXX (to be deposited upon

publication)). For detailed information on all other datasets see 'Dataset Assembly'.

Code availability

All packages used have been published, are freely available and are referenced in

the methods. R markdowns used to run the analyses specific to this study are

available from the authors on request.

Acknowledgements

AC was supported by postgraduate research scholarships from UCL and received

additional research support from a Debbie Fund grant to KC and TRF. TRF was

supported by Rosetrees Trust (M229-CD1), Cancer Research UK (A25825), the

Biotechnology and Biosciences Research Council (Grant Ref: BB/V010271/1), the

Royal Society (IEC\R2\202256) and the Global Challenges Doctoral Centre at the

University of Kent. DNA methylation data were generated through funding provided

by the Debbie Fund and the results shown here are in part based upon data generated by the TCGA Research Network: https://www.cancer.gov/tcga and the Cancer Genome Characterization Initiative: https://ocg.cancer.gov/programs/cgci. AF was supported by grants from the MRC (MR/M025411/1), PCUK(MA-TR15-009), BBSRC (BB/R009295/1), TUF, Orchid and the UCLH BRC. The authors dedicate this manuscript to the late Dr Helga Salvesen, a wonderful collaborator and colleague who played a key role in the project.

References

1. de Martel, C., Plummer, M., Vignat, J. & Franceschi, S. Worldwide burden of cancer attributable to HPV by site, country and HPV type. *International journal of cancer* **141**, 664–670 (2017).

2. Li, N., Franceschi, S., Howell-Jones, R., Snijders, P. J. F. & Clifford, G. M. Human papillomavirus type distribution in 30,848 invasive cervical cancers worldwide: Variation by geographical region, histological type and year of publication. *International journal of cancer* **128**, 927–935 (2011).

3. Jung, E. J. *et al.* Cervical adenocarcinoma has a poorer prognosis and a higher propensity for distant recurrence than squamous cell carcinoma. *International Journal of Gynecological Cancer* **27**, 1228–1236 (2017).

4. Huang, Y. T. *et al.* Clinical behaviors and outcomes for adenocarcinoma or adenosquamous carcinoma of cervix treated by radical hysterectomy and adjuvant radiotherapy or chemoradiotherapy. *International Journal of Radiation Oncology Biology Physics* **84**, 420–427 (2012).

5. Zhou, J. *et al.* Comparison of clinical outcomes of squamous cell carcinoma, adenocarcinoma, and adenosquamous carcinoma of the uterine cervix after definitive radiotherapy: a population-based analysis. *Journal of cancer research and clinical oncology* **143**, 115–122 (2017).

6. Galic, V. *et al.* Prognostic significance of adenocarcinoma histology in women with cervical cancer. *Gynecologic Oncology* **125**, 287–291 (2012).

7. Chen, X. *et al.* Better or Worse? The Independent Prognostic Role of HPV-16 or HPV-18 Positivity in Patients With Cervical Cancer: A Meta-Analysis and Systematic Review. *Frontiers in oncology* vol. 10 1733 (2020).

8. Burk, R. D. *et al.* Integrated genomic and molecular characterization of cervical cancer. *Nature* **543**, 378–384 (2017).

9. Gagliardi, A. *et al.* Analysis of Ugandan cervical carcinomas identifies human papillomavirus clade–specific epigenome and transcriptome landscapes. *Nature genetics* **52**, 800–810 (2020).

10. Huang, J. *et al.* Comprehensive genomic variation profiling of cervical intraepithelial
neoplasia and cervical cancer identifies potential targets for cervical cancer early
warning. *Journal of medical genetics* **56**, 186–194 (2019).
11. Ojesina, A. I. *et al.* Landscape of genomic alterations in cervical carcinomas. *Nature*
506, 371–375 (2014).
12. Chen, J. L. *et al.* Differential clinical characteristics, treatment response and prognosis
of locally advanced adenocarcinoma/adenosquamous carcinoma and squamous cell
carcinoma of cervix treated with definitive radiotherapy. *Acta Obstetricia et*
Gynecologica Scandinavica **93**, 661–668 (2014).
13. Williams, N. L., Werner, T. L., Jarboe, E. A. & Gaffney, D. K. Adenocarcinoma of the
cervix: should we treat it differently? *Current oncology reports* **17**, 16–17 (2015).
14. Hu, K., Wang, W., Liu, X., Meng, Q. & Zhang, F. Comparison of treatment outcomes
between squamous cell carcinoma and adenocarcinoma of cervix after definitive
radiotherapy or concurrent chemoradiotherapy. *Radiation oncology (London, England)*
13, 245–249 (2018).
15. Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature*
500, 415–421 (2013).
16. Martincorena, I. *et al.* Universal Patterns of Selection in Cancer and Somatic Tissues.
Cell **171**, 1029-1041.e21 (2017).
17. Yoon, S., Baik, B., Park, T. & Nam, D. Powerful p-value combination methods to
detect incomplete association. *Scientific Reports* **11**, 6980 (2021).
18. Lawrence, M. S. *et al.* Comprehensive genomic characterization of head and neck
squamous cell carcinomas. *Nature* **517**, 576–582 (2015).
19. Lin, D.-C. *et al.* Genomic and molecular characterization of esophageal squamous cell
carcinoma. *Nature genetics* **46**, 467–473 (2014).
20. Hammerman, P. S. *et al.* Comprehensive genomic characterization of squamous cell
lung cancers. *Nature* **489**, 519–525 (2012).
21. Rauskolb, C., Sun, S., Sun, G., Pan, Y. & Irvine, K. D. Cytoskeletal tension inhibits
Hippo signaling through an Ajuba-Warts complex. *Cell* **158**, 143–156 (2014).
22. Nguyen, T. H., Ralbovska, A. & Kugler, J.-M. RhoBTB Proteins Regulate the Hippo
Pathway by Antagonizing Ubiquitination of LKB1. *G3 (Bethesda, Md.)* **10**, 1319–
1325 (2020).
23. Mohseni, M. *et al.* A genetic screen identifies an LKB1-MARK signalling axis
controlling the Hippo-YAP pathway. *Nature cell biology* **16**, 108–117 (2014).
24. Martin, D. *et al.* Assembly and activation of the Hippo signalome by FAT1 tumor
suppressor. *Nature communications* **9**, 2372 (2018).
25. Sourbier, C. *et al.* Targeting loss of the Hippo signaling pathway in NF2-deficient
papillary kidney cancers. *Oncotarget* **9**, 10723–10733 (2018).
26. Petrilli, A. M. & Fernández-Valle, C. Role of Merlin/NF2 inactivation in tumor
biology. *Oncogene* **35**, 537–548 (2016).
27. White, S. M. *et al.* YAP/TAZ Inhibition Induces Metabolic and Signaling Rewiring
Resulting in Targetable Vulnerabilities in NF2-Deficient Tumor Cells. *Developmental*
cell **49**, 425-443.e9 (2019).
28. Yang, H. *et al.* NF2 and Canonical Hippo-YAP Pathway Define Distinct Tumor
Subsets Characterized by Different Immune Deficiency and Treatment Implications in
Human Pleural Mesothelioma. *Cancers* **13**, 1561. doi: 10.3390/cancers13071561
(2021).
29. Gonzalez-Avila, G. *et al.* Matrix metalloproteinases participation in the metastatic
process and their diagnostic and therapeutic applications in cancer. *Critical reviews in*
oncology/hematology **137**, 57–83 (2019).

30.   Wang, Y. *et al.* Comprehensive Molecular Characterization of the Hippo Signaling
Pathway in Cancer. *Cell reports* **25**, 1304-1317.e5 (2018).

31.   Gooden, M. J., de Bock, G. H., Leffers, N., Daemen, T. & Nijman, H. W. The
prognostic influence of tumour-infiltrating lymphocytes in cancer: a systematic review
with meta-analysis. *British journal of cancer* **105**, 93–103 (2011).

32.   Jordanova, E. S. *et al.* Human leukocyte antigen class I, MHC class I chain-related
molecule A, and CD8+/regulatory T-cell ratio: which variable determines survival of
cervical cancer patients? *Clinical cancer research : an official journal of the*
American Association for Cancer Research **14**, 2028–2035 (2008).

33.   Nedergaard, B. S., Ladekarl, M., Thomsen, H. F., Nyengaard, J. R. & Nielsen, K. Low
density of CD3+, CD4+ and CD8+ cells is associated with increased risk of relapse in
squamous cell cervical cancer. *British journal of cancer* **97**, 1135–1138 (2007).

34.   Chakravarthy, A. *et al.* Pan-cancer deconvolution of tumour composition using DNA
methylation. *Nature communications* **9**, 3220–3221 (2018).

35.   Mizunuma, M. *et al.* The pretreatment neutrophil-to-lymphocyte ratio predicts
therapeutic response to radiation therapy and concurrent chemoradiation therapy in
uterine cervical cancer. *International journal of clinical oncology* **20**, 989–996 (2015).

36.   Lee, Y. Y. *et al.* Pretreatment neutrophil:lymphocyte ratio as a prognostic factor in
cervical carcinoma. *Anticancer Research* **32**, 1555–1561 (2012).

37.   Huang, Q. T. *et al.* Prognostic significance of neutrophil-to-lymphocyte ratio in
cervical cancer: A systematic review and meta-analysis of observational studies.
Oncotarget **8**, 16755–16764 (2017).

38.   Prasad, D. V. R. *et al.* Murine B7-H3 Is a Negative Regulator of T Cells. *The Journal*
of Immunology **173**, 2500 (2004).

39.   Zang, X. *et al.* B7x: a widely expressed B7 family member that inhibits T cell
activation. *Proceedings of the National Academy of Sciences of the United States of*
America **100**, 10388–10392 (2003).

40.   Yu, M. *et al.* CD73 on cancer-associated fibroblasts enhanced by the A2B-mediated
feedforward circuit enforces an immune checkpoint. *Nature Communications* **11**, 515
(2020).

41.   Han, S. *et al.* Roles of B7-H3 in Cervical Cancer and Its Prognostic Value. *Journal of*
Cancer **9**, 2612–2624 (2018).

42.   Huang, C. *et al.* B7-H3, B7-H4, Foxp3 and IL-2 expression in cervical cancer:
Associations with patient outcome and clinical significance. *Oncology reports* **35**,
2183–2190 (2016).

43.   Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. *Nature communications* **10**, 1523–1526 (2019).

44.   Ruscetti, M., Quach, B., Dadashian, E. L., Mulholland, D. J. & Wu, H. Tracking and
Functional Characterization of Epithelial-Mesenchymal Transition and Mesenchymal
Tumor Cells during Prostate Cancer Metastasis. *Cancer research* **75**, 2749–2759
(2015).

45.   Fischer, K. R. *et al.* Epithelial-to-mesenchymal transition is not required for lung
metastasis but contributes to chemoresistance. *Nature* **527**, 472–476 (2015).

46.   Zheng, X. *et al.* Epithelial-to-mesenchymal transition is dispensable for metastasis but
induces chemoresistance in pancreatic cancer. *Nature* **527**, 525–530 (2015).

47.   Puram, S. v *et al.* Single-Cell Transcriptomic Analysis of Primary and Metastatic
Tumor Ecosystems in Head and Neck Cancer. *Cell* **171**, 1611-1624.e24 (2017).

48.   Ko, Y.-C. *et al.* Index of Cancer-Associated Fibroblasts Is Superior to the Epithelial-
Mesenchymal Transition Score in Prognosis Prediction. *Cancers* **12**, 1718 (2020).

49.   Shen, T. *et al.* YAP1 plays a key role of the conversion of normal fibroblasts into
861   cancer-associated fibroblasts that contribute to prostate cancer progression. *Journal of*
862   *Experimental & Clinical Cancer Research* **39**, 36 (2020).

863   50.   Zanconato, F. *et al.* Genome-wide association between YAP/TAZ/TEAD and AP-1 at
864   enhancers drives oncogenic growth. *Nature cell biology* **17**, 1218–1227 (2015).

865   51.   Shao, D. D. *et al.* KRAS and YAP1 converge to regulate EMT and tumor survival.
866   *Cell* **158**, 171–184 (2014).

867   52.   Schlegelmilch, K. *et al.* Yap1 acts downstream of α-catenin to control epidermal
868   proliferation. *Cell* **144**, 782–795 (2011).

869   53.   Kim, J. *et al.* YAP/TAZ regulates sprouting angiogenesis and vascular barrier
870   maturation. *The Journal of clinical investigation* **127**, 3441–3461 (2017).

871   54.   Sakamoto, H. *et al.* PAI-1 derived from cancer-associated fibroblasts in esophageal
872   squamous cell carcinoma promotes the invasion of cancer cells and the migration of
873   macrophages. *Laboratory Investigation* **101**, 353–368 (2021).

874   55.   Neuzillet, C. *et al.* Inter- and intra-tumoural heterogeneity in cancer-associated
875   fibroblasts of human pancreatic ductal adenocarcinoma. *The Journal of pathology* **248**,
876   51–65 (2019).

877   56.   Qian, J. *et al.* A pan-cancer blueprint of the heterogeneous tumor microenvironment
878   revealed by single-cell profiling. *Cell research* **30**, 745–762 (2020).

879   57.   Mhaidly, R. & Mechta-Grigoriou, F. Fibroblast heterogeneity in tumor micro-
880   environment: Role in immunosuppression and new therapies. *Seminars in Immunology*
881   **48**, 101417 (2020).

882   58.   Hutton, C. *et al.* Single-cell analysis defines a pancreatic fibroblast lineage that
883   supports anti-tumor immunity. *Cancer cell* **39**, 1227-1244.e20 (2021).

884   59.   Galbo, P. M., Zang, X. & Zheng, D. Molecular Features of Cancer-associated
885   Fibroblast Subtypes and their Implication on Cancer Pathogenesis, Prognosis, and
886   Immunotherapy Resistance. *Clinical Cancer Research* **27**, 2636 (2021).

887   60.   Chen, Z. *et al.* Single-cell RNA sequencing highlights the role of inflammatory cancer-
888   associated fibroblasts in bladder urothelial carcinoma. *Nature Communications* **11**,
889   5077 (2020).

890   61.   Rader, J. S. *et al.* Genetic variations in human papillomavirus and cervical cancer
891   outcomes. *International journal of cancer* **144**, 2206–2214 (2019).

892   62.   Wright, J. D. *et al.* Human papillomavirus type and tobacco use as predictors of
893   survival in early stage cervical carcinoma. *Gynecologic oncology* **98**, 84–91 (2005).

894   63.   Yang, S. H., Kong, S. K., Lee, S. H., Lim, S. Y. & Park, C. Y. Human papillomavirus
895   18 as a poor prognostic factor in stage I-IIA cervical cancer following primary surgical
896   treatment. *Obstetrics & gynecology science* **57**, 492–500 (2014).

897   64.   Burger, R. A. *et al.* Human papillomavirus type 18: association with poor prognosis in
898   early stage cervical cancer. *Journal of the National Cancer Institute* **88**, 1361–1368
899   (1996).

900   65.   Schwartz, S. M. *et al.* Human papillomavirus and prognosis of invasive cervical
901   cancer: a population-based study. *Journal of clinical oncology : official journal of the*
902   *American Society of Clinical Oncology* **19**, 1906–1915 (2001).

903   66.   Hang, D. *et al.* Independent prognostic role of human papillomavirus genotype in
904   cervical cancer. *BMC Infectious Diseases* **17**, 391 (2017).

905   67.   Chumduri, C. *et al.* Opposing Wnt signals regulate cervical squamocolumnar
906   homeostasis and emergence of metaplasia. *Nature cell biology* **23**, 184–197 (2021).

907   68.   Fridlender, Z. G. *et al.* Polarization of tumor-associated neutrophil phenotype by TGF-
908   beta: "N1" versus "N2" TAN. *Cancer cell* **16**, 183–194 (2009).

69. Zhu, Q. *et al.* The IL-6–STAT3 axis mediates a reciprocal crosstalk between cancer-
derived mesenchymal stem cells and neutrophils to synergistically prompt gastric
cancer progression. *Cell Death & Disease* **5**, e1295–e1295 (2014).

70. Ohms, M., Möller, S. & Laskay, T. An Attempt to Polarize Human Neutrophils
Toward N1 and N2 Phenotypes in vitro. *Frontiers in immunology* **11**, 532 (2020).

71. SenGupta, S. *et al.* Triple-Negative Breast Cancer Cells Recruit Neutrophils by
Secreting TGF-β and CXCR2 Ligands. *Frontiers in immunology* **12**, 659996 (2021).

72. Casbon, A. J. *et al.* Invasive breast cancer reprograms early myeloid differentiation in
the bone marrow to generate immunosuppressive neutrophils. *Proceedings of the*
National Academy of Sciences of the United States of America **112**, E566-75 (2015).

73. Shaul, M. E. *et al.* Tumor-associated neutrophils display a distinct N1 profile
following TGFβ modulation: A transcriptomics analysis of pro- vs. antitumor TANs.
Oncoimmunology **5**, e1232221 (2016).

74. Kim, Y., Lee, D., Lee, J., Lee, S. & Lawler, S. Role of tumor-associated neutrophils in
regulation of tumor growth in lung cancer development: A mathematical model. *PloS*
one **14**, e0211041 (2019).

75. Xie, C. *et al.* Soluble B7-H3 promotes the invasion and metastasis of pancreatic
carcinoma cells through the TLR4/NF-κB pathway. *Scientific reports* **6**, 27528 (2016).

76. MacGregor, H. L. *et al.* High expression of B7-H3 on stromal cells defines tumor and
stromal compartments in epithelial ovarian cancer and is associated with limited
immune activation. *Journal for ImmunoTherapy of Cancer* **7**, 357 (2019).

77. Zhan, S. *et al.* Overexpression of B7-H3 in α-SMA-Positive Fibroblasts Is Associated
With Cancer Progression and Survival in Gastric Adenocarcinomas. *Frontiers in*
Oncology **9**, 1466 (2020).

78. Costa, A. *et al.* Fibroblast Heterogeneity and Immunosuppressive Environment in
Human Breast Cancer. *Cancer cell* **33**, 463-479.e10 (2018).

79. Chung, H. C. *et al.* Efficacy and Safety of Pembrolizumab in Previously Treated
Advanced Cervical Cancer: Results From the Phase II KEYNOTE-158 Study. *Journal*
of Clinical Oncology **37**, 1470–1478 (2019).

80. Colombo, N. *et al.* Pembrolizumab for Persistent, Recurrent, or Metastatic Cervical
Cancer. *New England Journal of Medicine* (2021) doi:10.1056/nejmoa2112435.

81. Lheureux, S. *et al.* Association of Ipilimumab With Safety and Antitumor Activity in
Women With Metastatic or Recurrent Human Papillomavirus-Related Cervical
Carcinoma. *JAMA oncology* **4**, e173776–e173776 (2018).

82. Mayadev, J. S. *et al.* Sequential Ipilimumab After Chemoradiotherapy in Curative-
Intent Treatment of Patients With Node-Positive Cervical Cancer. *JAMA oncology* **6**,
92–99 (2020).

83. Naumann, R. W. *et al.* Efficacy and safety of nivolumab (Nivo) + ipilimumab (Ipi) in
patients (pts) with recurrent/metastatic (R/M) cervical cancer: Results from
CheckMate 358. in *Annals of Oncology* v898–v899 (2019).
doi:10.1093/annonc/mdz394.

84. Heeren, A. M. *et al.* Efficacy of PD-1 blockade in cervical cancer is related to a
CD8+FoxP3+CD25+ T-cell subset with operational effector functions despite high
immune checkpoint levels. *Journal for ImmunoTherapy of Cancer* **7**, 43 (2019).

85. Halle, M. K. *et al.* Clinicopathologic and molecular markers in cervical carcinoma: a
prospective cohort study. *American journal of obstetrics and gynecology* **217**, 432.e1-
432.e17 (2017).

86. Lando, M. *et al.* Identification of eight candidate target genes of the recurrent 3p12-
p14 loss in cervical cancer by integrative genomic profiling. *The Journal of pathology*
230, 59–69 (2013).

87. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-
seq quantification. *Nature Biotechnology* **34**, 525–527 (2016).
88. Aryee, M. J. *et al.* Minfi: a flexible and comprehensive Bioconductor package for the
analysis of Infinium DNA methylation microarrays. *Bioinformatics (Oxford, England)*
30, 1363–1369 (2014).
89. Fortin, J.-P. *et al.* Functional normalization of 450k methylation array data improves
replication in large cancer studies. *Genome biology* **15**, 503 (2014).
90. Teschendorff, A. E. *et al.* A beta-mixture quantile normalization method for correcting
probe design bias in Illumina Infinium 450 k DNA methylation data. *Bioinformatics*
(Oxford, England) **29**, 189–196 (2013).
91. Chen, Y. *et al.* VirusSeq: software to identify viruses and their integration sites using
next-generation sequencing of human cancer tissue. *Bioinformatics (Oxford, England)*
29, 266–267 (2013).
92. Lyng, H. *et al.* Intratumor chromosomal heterogeneity in advanced carcinomas of the
uterine cervix. *International Journal of Cancer* **111**, 358–366 (2004).
93. enbabaoğlu, Y., Michailidis, G. & Li, J. Z. Critical limitations of consensus
clustering in class discovery. *Scientific reports* **4**, 6207 (2014).
94. Mayakonda, A., Lin, D.-C., Assenov, Y., Plass, C. & Koeffler, H. P. Maftools:
efficient and comprehensive analysis of somatic variants in cancer. *Genome research*
28, 1747–1756 (2018).
95. Fantini, D., Vidimar, V., Yu, Y., Condello, S. & Meeks, J. J. MutSignatures: an R
package for extraction and analysis of cancer mutational signatures. *Scientific Reports*
10, 18217 (2020).
96. Reinius, L. E. *et al.* Differential DNA methylation in purified human blood cells:
implications for cell lineage and studies on disease susceptibility. *PloS one* **7**, e41361–
e41361 (2012).
97. Feber, A. *et al.* Using high-density DNA methylation arrays to profile copy number
alterations. *Genome biology* **15**, R30–R30 (2014).
98. Mermel, C. H. *et al.* GISTIC2.0 facilitates sensitive and confident localization of the
targets of focal somatic copy-number alteration in human cancers. *Genome Biology* **12**,
R41 (2011).
99. Gaujoux, R. & Seoighe, C. A flexible R package for nonnegative matrix factorization.
BMC Bioinformatics **11**, 367 (2010).

Figure Legends

Figure 1 Consensus clustering produces two prognostic clusters in TCGA SCC cohort. a) Consensus clustering of 236 TCGA HPV+ SCC patients. **b)** There were 938 differentially expressed genes between the two clusters. **c)** 5 year survival between the 2 SCC subgroups. **d)** 5 year survival between the 2 SCC subgroups considering only HPV16+ tumours. Statistics from univariate Cox regression.

1001

Figure 2 Cluster allocation of validation cohorts using methylation signature.
a) A signature of DNA methylation (dB > 0.25, FDR < 0.01) separates C1 and C2 SCC subgroups in the TCGA cohort. **b)** The methylation patterns are reproduced in a validation dataset from three European centres (n = 313). **c)** C2 tumours from TCGA and European validation cohorts cluster together based on the 129 MVP signature. **d)** 5 year survival curve for combined European validation cohorts. Statistics from univariate Cox regression.

Figure 3 Comparison of SCC subgroups with previous studies. Cluster analysis had previously been performed on 140 TCGA SCC tumours in two studies – one determined clusters based on cell of origin markers (Chumduri *et al,* 2021, red), one determined clusters based on integrated omics data (TCGA Network, 2017, orange). The heatmap at the bottom of plot represents expression levels of cytokeratin genes present in our C2 gene signature.

Figure 4 Genomic summary of significantly mutated genes (SMGs) in SCC cohorts. Main plot shows mutation type and frequencies for 34 SMGs identified using dNdSCV on TCGA, Bergen and Ugandan cohorts (367 total patients). Grey bars at top of plot represent TMB per sample. Grey bars to left of plot represent significance of SMG, larger bar is more significant. Barchart to the right shows proportion of a genes mutations in by cluster (blue = C1, red = C2). Black box around bar represents a significant difference in mutation frequency between the clusters (p<0.05) while a gold box means no significant difference between the

clusters. The plot at the bottom of figure represents the mutational signatures that

contribute towards each individuals tumour mutational burden.

[Gene name key – blue – unique to C1 analysis, red = unique to C2 analysis, black =

both in C1 and C2 individual analyses, black* = only significant when combining both

clusters for analysis, † = novel SMG in cervical cancer, ‡ = not significant in

combined cluster analysis but significant in C1 only analysis].

1031

Figure 5 Copy number and protein level differences between SCC subgroups.

a) Volcano plot showing differences in GISTIC copy number peak frequencies

between C1 and C2 tumours, with –log10(FDR) on the y axis and the odds ratio on

the x axis. **b)** Volcano plot showing differentially abundant proteins and phospho-

proteins (FDR < 0.05, FC > 1.3, represented by yellow dots) between C1 and C2

TCGA tumours, as measured by Reverse Phase Protein Array.

1038

1039 **Figure 6 Differences in the tumour microenvironment between cervical cancer**

1040 **subgroups.** Plot showing median abundances (x-axis) and median differences (%,

1041 y-axis) for different cell types estimated using MethylCIBERSORT, with significant

1042 differences in orange, for **a)** TCGA discovery cohort and **b)** combined validation

1043 cohorts. **c)** C2 tumours cluster together using CAF geneset genes.

1044

1045 **Supplementary Figure Legends**

1046

1047 **Supplementary Figure S1 – tSNE clustering by histology in cervical cancer**

1048 **cohorts.** Unsupervised tSNE analysis using top 10% most variable genes for

1049 cervical cancer cohorts **a)** TCGA (1385 most variable genes), **b)** Ugandan (1371)

1050   and **c)** Bergen (1430). Concordance of most variable genes was high amongst the 3

1051   cohorts **(d)**.

1052

1053   **Supplementary Figure S2 Consensus clustering using ConsensusClusterPlus.**

1054   **a)** Consensus CDF plot. PAC score = CDF at 0.9 consensus index − CDF at 0.1

1055   consensus index for each curve. **b)** Delta area plot used in decision of optimum

1056   number of clusters.

1057

Supplementary Figure S3 Genes that are both differentially expressed and

differentially methylated between C1 and C2 subgroups. Datapoints represent

methylated variable positions (in either the 3'UTR, body of gene, intergenic region or

gene promoter) in genes that are also differentially expressed between C1 and C2

subgroups. Datapoints in the top left quadrant are MVPs that are hypomethylated in

genes that are also upregulated in C2 tumours. Those in the bottom right quadrant

are hypermethylated in genes that are downregulated in C2 tumours.

1065

Supplementary Figure S4 Concordance between gene expression and DNA

methylation-derived cluster membership. a) The percentage of samples that are

designated the same cluster allocation by gene expression signature and

methylation signatures based on varying delta Beta thresholds. **b)** ROC curves

showing the accuracy with which C1 or C2 cluster membership can be predicted

using DNA methylation differences (MVPs) in samples from the validation cohorts for

which either RNA-seq (Bergen, n=37, and Uganda, n=94, HPV+ SCC cases),

Illumina HumanHT-12 V4.0 expression beadchip array (Oslo SCC cases, n=109) or

Illumina HumanWG-6 v3.0 expression beadchip array (Oslo SCC cases, n=139)

gene expression data were available. **c)** Single sample gene set enrichment analysis

(ssGSEA) for validation cohorts used in panel B. The y-axis represents the ssGSEA

score for each sample, compared with the genes from the C2 gene expression

signature. P-values from Wilcoxon rank-sum test.

1079

1080 **Supplementary Figure S5 Validation SCC cohorts. a)** Ugandan validation cohort

1081 clustering based on 116 MVP signature. Kaplan-meier curves for **b)** HPV16+

1082 European validation cohort SCC patients; **c)** European validation cohort SCC

1083 patients without chemotherapy treatment and **d)** 5 year survival for the 5 individual

1084 cohorts in this study.

1085

1086 **Supplementary Figure S6 Elevation of epithelial mesenchymal transition (EMT)**

1087 **score is evident in C2 tumours. a)** EMT score derived by TCGA for 140 HPV+

1088 squamous TCGA cervical cancer tumours in our study. EMT score is higher in C2

1089 tumours.

1090

1091 **Supplementary Figure S7 Mutational signatures of combined HPV+ squamous**

1092 **cervical cancer cohorts.** COSMIC mutational signatures identified in combined

1093 HPV+ squamous cervical cancer cohort including genomic data from TCGA, Bergen

1094 and Ugandan cohorts.

1095

1096

1097 **Supplementary Figure S8 Increased levels of YAP in tumours with YAP1**

1098 **amplification.** YAP1 expression **(a)**, and YAP protein levels **(b)** unphosphorylated,

1099 **c)** phosphorylated) are higher in tumours that contain YAP1 amplifications.

Supplementary Figure S9 Differences in immune microenvironment between SCC subgroups in individual cohorts. Median abundances (x-axis) and median differences (%, y-axis) for different cell types estimated using MethylCIBERSORT, with significant differences in orange for cohorts from **a)** Bergen, **b)** Innsbruck, **c)** Oslo and **d)** Uganda. C2 tumours display increased neutrophil:CTL ratios as estimated using MethylCIBERSORT for **e)** TCGA discovery cohort and **f)** combined validation cohorts.

Supplementary Figure S10 Immune cell ratios by cluster using MethylCIBERSORT estimates. a) Neutrophil:CD19 estimate ratios for combined cohorts. **b)** Neutrophil:Treg estimate ratios for combined cohorts.

Supplementary Figure S11 Comparison of MethylCIBERSORT estimates and immunohistochemistry(IHC)-based scoring. Correlations between MethylCIBERSORT estimates and IHC-based scoring for **a)** CD8+ T-cells, **b)** neutrophils (MPO+), **c)** CD8+ T-cell:neutrophil ratio in 14 SCCs from the Innsbruck validation cohort and **d)** CD8+ T-cells for 229 SCCs from the Oslo validation cohort. Trendlines are derived from linear modelling, shaded areas represent 95% CI of trendlines.

Supplementary Figure S12 Upregulation of immune checkpoint genes in C2 SCCs. Upregulation of **a)** *B7-H3* (*CD276*), **b)** *NT5E* (*CD73*) and **c)** *PD-L2* (*PDCD1LG2*) was observed in poor prognosis C2 tumours. Analysis performed with RNA-seq data from TCGA, Bergen and Ugandan cohorts.

1125

1126 **Tables**

1127

1128 Table 1: Summary of clinicopathological characteristics for five cervical cancer

1129 cohorts.

1130

1131 Table 2 – Five-year survival analysis for all cohorts

1132

1133 **Supplementary Tables**

1134

1135 Table S1 - Clinical and pathaologic characteristics of TCGA squamous cervical

1136 cancer cohort samples

1137

1138 Table S2 - Top 10% most variable genes in TCGA squamous cervical cancer cohort

1139

1140 Table S3 - 938 Differentially expressed genes between TCGA squamous cervical

1141 cancer clusters C1 and C2

1142

1143 Table S4 - 5 year survival uni- and multivariate analysis for HPV16+ patients in

1144 squamous cervical cancer cohorts

1145

1146 Table S5 - 129 MVP signature probes (European validation cohorts)

1147

1148 Table S6 - Combined validation cohort cluster allocation

1149

1150   Table S7 - Breakdown of tumour stage in C1 and C2 cluster by percentage

1151

1152   Table S8 - Clusters and EMT scores for TCGA squamous cervical cancer samples

1153

1154   Table S9 - Significantly mutated genes using dNdSCV analysis and combining

1155   cohorts

1156

1157   Table S10 - Mutation frequency in SMGs observed in previous studies

1158

1159   Table 11 - Gene set enrichment analysis of C2 gene expression signature genes

1160   using Metascape

1161

1162   Table S12 - Paramaters for TSNE multidimensional visualisation analyses

1163

1164   Table S13 - 116 MVP signature probes (Ugandan validation cohort)

1165

1166

1167

1168

1169

1170

1171

Figure 1 Consensus clustering produces two prognostic clusters in TCGA SCC cohort. a) Consensus clustering of 236 TCGA HPV+ SCC patients. **b)** There were 938 differentially expressed genes between the two clusters. **c)** 5 year survival between the 2 SCC subgroups. **d)** 5 year survival between the 2 SCC subgroups considering only HPV16+ tumours. Statistics from univariate Cox regression.

Figure 2 Cluster allocation of validation cohorts using methylation signature. a) A signature of DNA methylation (dB > 0.25, FDR < 0.01) separates C1 and C2 SCC subgroups in the TCGA cohort. **b)** The methylation patterns are reproduced in a validation dataset from three European centres (n = 313). **c)** C2 tumours from TCGA and European validation cohorts cluster together based on the 129 MVP signature. **d)** 5 year survival curve for combined European validation cohorts. Statistics from univariate Cox regression.

Figure 3 Comparison of SCC subgroups with previous studies. Cluster analysis had previously been performed on 140 TCGA SCC tumours in two studies – one determined clusters based on cell of origin markers (Chumduri *et al,* 2021, red), one determined clusters based on integrated omics data (TCGA Network, 2017, orange). The heatmap at the bottom of plot represents expression levels of cytokeratin genes present in our C2 gene signature.

Figure 4 Genomic summary of significantly mutated genes (SMGs) in SCC cohorts. Main plot shows mutation type and frequencies for 34 SMGs identified using dNdSCV on TCGA, Bergen and Ugandan cohorts (367 total patients). Grey bars at top of plot represent TMB per sample. Grey bars to left of plot represent significance of SMG, larger bar is more significant. Barchart to the right shows proportion of a genes mutations in by cluster (blue = C1, red = C2). Black box around bar represents a significant difference in mutation frequency between the clusters (p<0.05) while a gold box means no significant difference between the clusters. The plot at the bottom of figure represents the mutational signatures that contribute towards each individuals tumour mutational burden.

[Gene name key – blue – unique to C1 analysis, red = unique to C2 analysis, black = both in C1 and C2 individual analyses, black* = only significant when combining both clusters for analysis, † = novel SMG in cervical cancer, ‡ = not significant in combined cluster analysis but significant in C1 only analysis]

Figure 5 Copy number and protein level differences between SCC subgroups. a) Volcano plot showing differences in GISTIC copy number peak frequencies between C1 and C2 tumours, with –log10(FDR) on the y axis and the odds ratio on the x axis. **b)** Volcano plot showing differentially abundant proteins and phospho-proteins (FDR < 0.05, FC > 1.3, represented by yellow dots) between C1 and C2 TCGA tumours, as measured by Reverse Phase Protein Array.

Figure 6 Differences in the tumour microenvironment between cervical cancer subgroups. Plot showing median abundances (x-axis) and median differences (%, y-axis) for different cell types estimated using MethylCIBERSORT, with significant differences in orange, for **a)** TCGA discovery cohort and **b)** combined validation cohorts. **c)** C2 tumours cluster together using CAF geneset genes.