Pro-metastatic gene expression, immune evasion and an altered HPV spectrum characterize an aggressive subtype of cervical cancer.

Ankur Chakravarthy[1,*], Ian Reddin[2,*], Stephen Henderson[3], Cindy Dong[2], Nerissa Kirkwood[2], Maxmilan Jeyakumar[2], Jacqueline McDermott[4], Xiaoping Su[5], Nagayasau Egawa[6], Christina S Fjeldbo[7], Mari Kyllesø Halle[8], Camilla Krakstad[8], Afschin Soleiman[9], Susanne Sprung[10], Heidi Lyng[7], Heidi Fiegl[11], Helga Salvesen[8], John Doorbar[6], Kerry Chester[4, #], Andrew Feber[12, #], Tim R Fenton[2,#].

[1] Princess Margaret Cancer Centre, University Health Network. Toronto, Ontario, Canada

[2] School of Biosciences, University of Kent, Canterbury, Kent, UK

[3] UCL Cancer Institute, Bill Lyons Informatics Centre, University College London, London, UK

[4] UCL Cancer Institute, University College London, London, UK

[5] MD Anderson Cancer Center, Houston, Texas, USA

[6] Department of Pathology, University of Cambridge, Cambridge, UK

[7] Department of Radiation Biology, Oslo University Hospital, Oslo, Norway

[8] Department of Obstetrics and Gynaecology, Haukeland University Hospital, Bergen, Norway; Centre for Cancer Biomarkers, Department of Clinical Science, University of Bergen, Norway

[9] INNPATH, Institute of Pathology, Tirol Kliniken Innsbruck, Innsbruck, Austria.

[10] Institute of Pathology, Medical University of Innsbruck, Innsbruck, Austria.

[11] Department of Obstetrics and Gynaecology, Medical University of Innsbruck, Innsbruck, Austria.

[12] Division of Surgery and Interventional Science, University College London, London, UK

*Equal contribution

[#]Correspondence to: k.chester@ucl.ac.uk; a.feber@ucl.ac.uk or t.fenton@kent.ac.uk

Abstract

Cervical cancer is caused by carcinogenic human papillomavirus infection and represents one of the

leading causes of cancer death worldwide. Effective means of tumour classification are required for

better disease understanding. We performed an integrated multi-omic analysis of 655 cervical

cancers, using epigenomic and transcriptomic signatures to discover two distinct cervical cancer

subtypes we named "typical" and "atypical". Typical tumours were largely HPV16-driven and

frequently displayed an 'immune-hot' tumour microenvironment. Atypical tumours were associated

with poor prognosis; they were more likely to be driven by HPVs from the HPV18-containing α7

clade, displayed distinct genomic aberrations, greater evidence of past immunoediting and a

microenvironment associated with immune-evasion and failure of anti-PD1 checkpoint inhibition.

The finding that atypical tumours encounter stronger anti-tumour immune responses during

development may explain the lower frequency at which α7 HPV infected-lesions progress from pre-

invasive disease. However those escaping this selection pressure evolve into aggressive tumours

(independent of HPV-type) in which more intensive adjuvant treatment may be warranted.

--

Despite screening and the introduction of prophylactic human papillomavirus (HPV) vaccination in

developed countries, cervical cancer continues to be one of the leading worldwide causes of cancer-

related deaths in women. Prognosis for patients with metastatic disease remains poor, thus new

treatments and effective molecular markers for patient stratification are urgently required. Cervical

cancer is caused by at least 14 high-risk human papillomaviruses (hrHPVs), with HPV16 and HPV18

together accounting for over 70% of cases worldwide[1]. As the two dominant HPV types in cervical

cancer, much work has focused on HPV16 and HPV18. Although HPV18 displays a more potent

ability to transform keratinocytes than HPV16 *in vitro*[2], persistent HPV16 infections are associated

with a higher risk of progression to CIN3 or invasive carcinoma, suggesting additional factors play an

important role in modulating HPV-driven carcinogenesis *in vivo*[3]. We have previously shown that the

majority of HPV16+ cervical tumours profiled by The Cancer Genome Atlas (TCGA) project fall into a

good prognosis group based on similarity at the DNA methylation level to HPV16+ HNSCC and penile

cancer[4], and in their integrated molecular analysis of cervical cancer TCGA reported differential

micro-RNA expression and transcription factor activation between tumours harbouring different

HPV types[5].   Here we made use of molecular data from 655 cervical cancers to extend these

observations, leading to the finding that cervical cancers can be broadly classified into two groups:

one containing the majority of HPV16+ cases shares gene expression, DNA methylation and genomic

features with HPV16-driven malignancies at other anatomic sites and another group, in which HPV

types from the HPV18-containing α7 clade are more likely to be found, in which the HPV16-

associated transcriptional programme is less evident. Tumours in this latter group also display

increased YAP1 expression, *STK11* mutations and lower numbers of tumour infiltrating lymphocytes,

both of which have potential implications for therapeutic approaches.

To address this question we used data from 281 cervical tumours profiled by TCGA[5] as our discovery

cohort (**Table 1, Table S1**) and 374 cervical cancers from three European centres, all with detailed

clinical annotation and long term follow-up as our validation cohort[6,7] (**Table 1, Table S2**).

Representing to our knowledge, the largest study of its kind in cervical cancer, we defined two

cervical cancer subtypes; an "atypical" aggressive subtype defined by a lymphocyte-depleted

microenvironment and evidence for epithelial-mesenchymal transition, and a "typical" subgroup

comprised almost entirely of tumours harbouring α9 HPVs and associated with longer overall

survival. The molecular, cellular and clinical differences identified between typical and atypical

tumours also reveal new potential therapeutic options for the treatment of cervical cancers.

Results

		TCGA Training Cohort	European Validation Cohort
Histology			
	Adenocarcinoma	44	32
	Squamous cell Carcinoma	237	335
	Adenosquamous	0	7
Stage			
	I	153	98
	II	61	194
	III	43	65
	IV	17	17
	NA	7	0
Age	Median (Range)	46 (20-88)	51 (22-91)
HPV Sub Type			
	16	166	235
	18	38	53
	45	22	14
	Other	55	45
	Negative	0	27
Survival Status			
	Alive	226	275
	Dead	55	99
Cluster Assignment			
	Typical	231	301
	Atypical	50	73

8   **Table 1:** Summary of clinicopathological characteristics for the two cervical cancer cohorts.

Generation of HPV type-specific gene expression and DNA methylation profiles in cervical cancer

To look for associations between HPV type and overall survival while avoiding confounding from

histology where clear molecular differences to squamous cell carcinomas (SCCs) was observed

(Figure S1A) and advanced stage, we initially reduced our discovery cohort to 139 stage I and II SCCs,

which we confirmed using VirusSeq[8] to be transcript-positive for at least one of the three most

common HPV types (HPV16, HPV18 and HPV45), and for which the covariates age and tumour stage

were available. Multivariate Cox regression identified a significantly worse prognosis in HPV45-

driven tumours relative to HPV16 (HR = 5.040, p < 1e-3), while HPV18+ tumours exhibited an

intermediate prognosis (**Figure 1A**).

2 **Figure 1. Clinical and molecular variation among TCGA cervical cancers driven by different HPV types. A)**

3 HPV45+ early stage Cervical Squamous Cancers display markedly worse prognosis compared to HPV16+

4 cancers with HPV18+ tumours showing intermediate survival. HR and p-value from Cox regression controlling

5 for stage. **B)** Comparisons between HPV45+ and HPV16+ tumours identify large scale variation in

transcriptional and **C)** epigenetic (genome-wide DNA methylation) profiles. **D)** Twelve genes from a pan-tissue

signature for HPV-driven tumorigenesis show significant variation between HPV16+ and HPV45+ tumours. **E)**

tSNE plot showing separation of HPV16+ from HPV18+ tumours based on the differential gene expression

signature. **F)** tSNE plot showing separation of HPV16+ from HPV18+ tumours based on the MVP (differential

methylation) signature.

Modelling transcriptomic and epigenomic (DNA methylation) differences between HPV16 and

HPV45-associated early-stage tumours identified 713 DEGs (Differentially Expressed Genes,

FDR=0.01, FC > 2; **Figure 1B**, **Table S3**) and 689 MVPs (Methylation Variable Positions, delta-Beta 0.1,

FDR < 0.01), (**Figure 1C, Table S4**). Multi-dimensional analysis using tSNE illustrates the molecular

differences between the two HPV types using these transcriptomic and epigenomic signatures

(Figure 1D and 1E). 10 DEGs previously shown to be aberrantly expressed in HPV-associated cancers

from different anatomic sites[9] displayed greater dysregulation in HPV16+ than in HPV45+ cervical

tumours, while two (*PLOD2* and *KRT18*) were more strongly upregulated in HPV45+ tumours (**Figure**

1D). Several DEGs were also differentially methylated between HPV16+ and HPV45+ tumours (**Figure**

S1B). These findings indicate molecular differences between cervical cancers driven by different

hrHPV types which may manifest in clinical differences.

18
Development and validation of a prognostic classification.

Interestingly, when using either the 713 DEG or 689 MVP signatures, a minority (10% gene

expression-based, 11% DNA methylation-based) early-stage HPV16+ SCCs clustered with the

HPV45+ tumours in the discovery cohort (**Figures 1B and 1C**). We subsequently used consensus

clustering based on the 713 DEG signature, which identified two robust clusters, which we termed

"typical" (as most tumours are classified in this subgroup and the gene expression profile resembles

that seen in HPV16+ cancers from other anatomic sites) and "atypical" cervical cancer subtypes. 6 of

104 HPV16+ tumours co-clustered with the majority (13 of 17) of HPV45+ tumours in the atypical

subgroup, which also contained 28% (5/18) of the HPV18+ tumours (**Figure 2A, Table S1**). To assess

if these typical and atypical subgroups also exist at the epigenetic level, we developed a DNA

methylation Support Vector Machine (SVM) classification model, using TCGA DNA methylation data

reduced to 178 CpG sites at which methylation differed significantly between tumours in typical

versus atypical clusters (**Figure 2B,** mean delta-Beta > 0.3, FDR < 0.01, **Table S5**). Using this signature

we allocated cluster membership to a further 374 cervical cancers from our validation cohort (**Figure**

2C and Table S2). Adenocarcinomas (12 of 32) and adenosquamous carcinomas (5 of 7) were more

likely to be classified as atypical than SCCs (56/335, p = 4.526e-09, Fisher's Exact Test) and again, the

majority (209 of 234) HPV16+ tumours were designated as typical (**Figure 2C**). We could accurately

predict the DEG-based cluster assignment using the MVP signature for all validation cohort tumours

for which gene expression data were available (RNA-seq for Bergen samples (n = 65) and Illumina

HumanHT-12 V4.0 expression beadchip arrays for Oslo samples (n = 268), **Figure S2A**), confirming

that both MVP and DEG signatures classify the same tumours as typical or atypical. Single-sample

gene set enrichment analysis (ssGSEA)[10] of the same tumours confirmed differential expression of

the signature genes in tumours classified as typical or atypical using DNA methylation (**Figure S2B**).

Having derived our typical and atypical clusters directly from the HPV45 vs HPV16 expression

signature and shown that they were consistent whether assigned from gene expression or from DNA

methylation data, for clarity we henceforth refer to all comparisons as atypical and typical.

Integrating DNA and RNA-based HPV typing where available, we confirmed that co-infection with

HPV45 or other HPV types was not responsible for the assignment of HPV16 transcript-positive

samples to the atypical group. In both the validation and discovery cohorts, HPV types from the α7

clade (HPV18, 45, 59, 68, 70) were strongly enriched in atypical tumours (atypical tumours were 5.6X

more likely to harbour α7 HPVs than typical tumours, p < 2.2e-16 Fisher's Exact Test).

Figure 2. Derivation of two type-associated prognostic subgroups in cervical cancer and validation across independent cohorts. A) HPV45-like transcriptional profiles are also shared by small numbers of HPV16+ and HPV18+ tumours, coalescing into HPV45-like ('Atypical') and HPV16-like ('Typical') clusters. **B)** A signature of DNA methylation (dB > 0.3, FDR < 0.01) separates these groups based on consensus clustering (see methods for details). **C)** The methylation patterns are reproduced in a validation dataset from three European centres (n = 374). **D)** Survival curves and statistics from multivariate Cox regression of overall survival in TCGA cervical

cancer cohort stratified by cluster. **E)** Survival curves and statistics from multivariate Cox regression of overall

survival in the European validation cohort stratified by cluster.

Multivariate analysis of survival data from the 274 TCGA tumours for which tumour stage was

available identified a significant prognostic difference between the typical and atypical subgroups

(**Figure 2D;** HR = 2.24, p = 0.01). This difference became even greater when restricting to stage I/II

tumours (**Figure S3A**; n = 139, HR = 4.88, p = 0.0006), and was retained even upon removal of the

HPV45+ tumours (**Figure S3B**; n = 122, HR = 4.91, p = 0.03).   Cox regression stratifying by histology

and controlling for FIGO stage and treatment (surgery alone, surgery with radio-chemotherapy and

surgery with chemotherapy alone) identified typical/atypical status to be an independent predictor

of overall survival in the validation cohort (n = 374, HR = 1.65 , p = 0.043**, Figure 2E)**. Again, the

survival difference between the typical and atypical groups was greater when stage IV tumours were

excluded (n = 357, HR = 1.73, p = 0.04) and was most pronounced in stage II tumours (n = 194, HR =

2.59, p = 0.02).

Evidence for epithelial to mesenchymal transition (EMT) in atypical tumours.

Gene set enrichment using Ingenuity Pathway Analysis 'Diseases and Functions Ontology' (**Figure S4**,

Table S6) which identified cellular movement as the most activated pathway in atypical tumours,

with 85 of 121 genes in the set expressed consistent with increased metastatic potential. Prominent

pro-metastatic genes in this pathway included the transcription factor *SNAI1* (a master regulator of

EMT that accompanies invasion through the basement membrane and dissemination from the

primary tumour[11] , fibronectin 1, which is known to trigger EMT-associated transcriptional

cascades[12], *RHOF*, a prominent player in invasion through pseudofilopodia formation[13] and *VEGFC,*

involved in prometastatic lymphangiogenesis[14]. Multiple other gene sets also pertaining to cell

movement were strongly enriched and associated with high activation z-scores in atypical tumours

(**Figure S4, Table S6**).   Moreover, these tumours expressed high levels of Transforming Growth

Factor (*TGFB1* and *TGFB2*, **Table S3**), which have been implicated as key inducers of EMT that are

potentially amenable to therapeutic targeting (reviewed in[15]). Given these findings, we examined the

relationship between our subtypes and the poor-prognosis cervical cancer EMT cluster defined by

TCGA based on reverse phase protein array (RPPA) data[5]. 62% of atypical TCGA tumours with RPPA

data available belong to the EMT cluster compared with only 20% of typical tumours. Consistent

with the proteomic classification, atypical tumours displayed higher EMT gene expression scores, as

defined by TCGA[5], than typical tumours (**Figure S5).** Upstream regulator analysis identified *EZH2* and

SMARCA4, both chromatin modifiers, as the leading differentially-activated regulators in atypical

tumours (**Figure S6, Table S7**). Notably *EZH2* expression has previously been linked to poor prognosis

in cervical cancer[16]. Other key activated regulators include β-catenin and HIF1α, both of which have

been linked to chemo or radio- resistance and poor prognosis in cervical cancer[17-19]. These analyses

link the atypical expression signature to several independently discovered poor prognostic factors.

Genomic analyses of prognostic clusters

To search for genomic differences between typical and atypical cancers, we analysed mutation

(WEX) and copy number data from samples with matched methylation data[20].   We first generated

segmented copy number data for all tumours (combining the TCGA and validation cohort samples

for which the necessary data were available for maximum statistical power), which identified 387

focal candidate copy number alterations at FDR < 0.1. Following binomial regression, we identified

12 discrete copy

Figure 3. Genomic differences between cervical cancer subgroups. A) Volcano plot showing differences in

GISTIC copy number peak frequencies between typical and atypical tumours, with -log10(FDR) on the y axis

and odds ratio on the x axis. **B)** Volcano plot showing differentially abundant proteins and phospho-proteins

(FDR < 0.05, FC > 1.3, represented by yellow dots) between typical and atypical TCGA tumours, as measured by

Reverse Phase Protein Array. **C)** Bar chart showing mutation frequencies for candidate driver mutations in

typical and atypical cancers. Y axis indicates percentage of tumours mutated within tumour subtype, and the

outline colour indicates statistical significance of differences in mutation frequencies. **D)** Histograms show

overall mutational burdens are greater in HPV16-like cancers. Odds ratios and p values are from a negative

binomial GLM. **E)** Neoantigen burdens are elevated in typical tumours cervical cancers (estimates and p-values

from negative binomial regression). **F)** Atypical tumours display greater evidence of past immunoediting as

measured by depletion of predicted neoantigens versus total mutations.

number alterations between typical and atypical clusters (**Table S8**, **Figure 3A;** FDR < 0.1, log2 (Odds

Ratio) > 1). These included 1p31.3 loss and 11q22.1 gain which were more prevalent in atypical

tumours and multiple 3q gains and 1p13.3 gain, and 11q22.1 loss, which were disproportionately

common in typical cancers (**Figure 3A**). Notably, the 11q22.1 gain seen in atypical tumours are

centred on the Yes-Associated Protein 1 (YAP1): a key transcription factor downstream of the HIPPO

signalling pathway. Analysis of Reverse Phase Protein Assay (RPPA) data from TCGA also revealed

significantly higher YAP1 protein expression in the atypical tumours (**Fig 3B**). We confirmed that

those same cases with *YAP1* amplification (7/28 atypical tumours and 12/141 typical tumours) also

showed increased YAP1 mRNA and protein expression.

We next compared the somatic mutation rates in a defined set of candidate driver genes, using a

binomial regression. This identified *PIK3CA, FBXW7* and *PTEN* mutations as disproportionately more

common in typical cancers and loss-of-function *STK11* mutations as more frequent in atypical

tumours (FDR < 0.25, **Figure 3C),** *STK11* (LKB1) is also under-expressed in atypical tumours compared

with typical tumours (**Table S1**). We observed a higher overall mutation burden in typical tumours

(OR= 0.48, p = 1.4e-5, **Figure 3D**), leading us to investigate whether there is also a difference in

neoantigen load between the subgroups. Fitting a negative binomial GLM to neoantigen data from

TCGA (31 atypical, 157 typical for which neoantigen estimates were available from the Cancer

Immunome Atlas [21]) revealed markedly more predicted neoantigens in typical tumours, at both the

gene and individual MHC class 1-binding peptide level (OR = 1.72, p = 0.01 and OR = 1.99, p = 0.001

respectively, **Figure 3E**). Interestingly, the ratio of expected versus observed neoantigens per tumour

(neoantigen depletion, see methods) is greater in atypical tumours (p = 0.02 (Wilcoxon's Rank Sum

test), **Fig 3F**), suggesting more extensive immunoediting during their development and leading us to

compare the tumour immune microenvironment between subgroups.

Immunological analyses of prognostic clusters implicate microenvironmental

differences and highlight potential therapeutic interventions.

The nature of the tumour immune microenvironment, particularly the abundance of tumour

infiltrating lymphocytes (TILs) is a strong prognostic factor in HPV-associated cancers[9,22-26]. Pathway

analysis revealed activation of granulocyte and aggranulocyte adhesion along with diapedesis in

atypical tumours, suggesting increased neutrophil infiltration. Other pathways activated in atypical

tumours include inflammatory processes such as Acute Phase Response, TREM1 signalling,

complement activation and AHR (Aryl Hydrocarbon Receptor) signalling which are often associated

with macrophages (**Figure S7, Table S9**), suggesting a strong microenvironmental component may

mediate type-associated pathology. This is supported by the presence of multiple cytokines and

chemokines in the atypical-associated transcriptional signature, including *IL11, IL18, IL1B, IL24, IL6,*

IL8, CCL2, CXCL2, CXCL3, CXCL5, TNF (TNF-α) and *TNFAIP6* (**Table S3**). We next used DNA methylation

data to compare the cellular composition of tumours[27], observing differences in the proportions of

multiple cell types between the subgroups (**Figure 4A**); most notably decreased CD8+ (cytotoxic T

lymphocytes (CTL)), a marked elevation of neutrophil and natural killer (NK)-cells and lower tumour

purity in atypical cancers (the latter confirmed using genomic estimates[28] (ABSOLUTE); **Figure S8**, p =

0.02 (Wilcoxon's Rank Sum test). Integrating these cell type estimates with our previously-published

pan-cancer immune hot/cold classification[27], we also found significant enrichment for immune-hot

cancers in the typical subgroup (**Figure 4B;** p = 1.9e-10), consistent with increased CTL infiltration,

CTL:Treg ratios (**Figure S9**) and higher MHC class I neoantigen loads in these tumours.

Figure 4. Differences in the immune microenvironment between cervical cancer subgroups. A) Plot showing

median abundances (x-axis) and median differences (%, y-axis) for different cell types estimated using

MethylCIBERSORT, with significant differences in orange. **B)** Applying a pan-cancer classifier to DNA

methylation data from the cervical cancer samples shows typical tumours are significantly more likely to be

immune-hot than atypical tumours. **C)** Atypical tumours display increased neutrophil:CTL ratios as estimated

using MethylCIBERSORT. **D)** Representative images showing Immunohistochemistry for MPO (neutrophils) and

CD8 in cervical tumour sections. **E)** Correlations between MethylCIBERSORT estimates and

immunohistochemistry-based scoring for neutrophils (MPO+). **F)** Correlations between MethylCIBERSORT

estimates and immunohistochemistry-based scoring for CD8+ T-cells. **G)** Correlations between

MethylCIBERSORT estimates and immunohistochemistry-based scoring for CD8+ T-cell:neutrophil ratios in 17

cervical tumours from the validation cohort. Trendlines in parts E-G are derived from linear modelling, shaded

areas represent 95% CI of trendlines. **H)** Typical tumours show higher fractions of proinflammatory (M1)

compared to immunosuppressive (M2/M0) macrophages (y axis = ratios, p value from Wilcoxon's Rank Sum

Test). **I)** ssGSEA scores (y-axis) showing enrichment for a TGFβ-associated extracellular matrix gene expression

signature in atypical tumours.

The estimation of increased neutrophil abundance in atypical tumours (**Figure 4A)** is supported by

the upregulation of genes associated with granulocyte diapedesis inferred from pathway analysis

(**Table S4**) and with increased ssGSEA scores for a neutrophil gene set derived from publicly-

available gene expression data (see methods; p<9.2e-6 (Wilcoxon's Rank Sum test) **Figure S10**).

Atypical tumours also exhibit a markedly higher neutrophil:CTL ratio (**Figure 4C**); an established

adverse prognostic factor in cervical cancer[29,30]. Enumerating CTLs and neutrophils by IHC for CD8

and MPO respectively in 17 tumours from our validation cohort (representative images shown in

Figure 4D, see methods for details) revealed strong correlations with MethylCIBERSORT estimates

(**Figure 4E-G**). Given the upregulation of multiple CXCR2-ligands (CXCL1, CXCL2, CXCL3, CXCL3) as

well as G-CSF, PTGS1, PTGS2, IL-1B, CCL2, IL6 (**Table S3**) and activation of TREM1 signalling (**Figure**

S7, Table S9) [31-36] in atypical tumours, all of which are associated with Monocytic-Myeloid Derived

Suppressor Cells or tolerogenic macrophages, we used CIBERSORT[37] to derive estimates of

macrophage subpopulations for samples with expression data in our dataset. Atypical tumours

displayed significantly lower pro-effector (M1) macrophage proportions relative to suppressive (M0

and M2-like) macrophages (p = 0.016 (Wilcoxon's Rank Sum test), **Figure 4H**). Finally, we observed

enrichment for a TGFβ-associated extracellular matrix gene expression signature (C-ECM) in atypical

tumours that we have previously linked to immune evasion and failure of PD1 blockade in

melanoma and bladder cancer[27]. Although total fibroblast content was similar between typical and

atypical tumours (**Figure 4A**) enrichment for the C-ECM signature indicates increased myofibroblast

trans-differentiation, which could explain the lack of CTL infiltration and the poor prognosis seen in

atypical tumours.

Discussion

Of the studies indicating an association between HPV type and prognosis in cervical cancer, both

HPV18 and the α7 clade to which it belongs have been linked to worse prognosis in early-stage

tumours[38-40]. Our analysis of TCGA cervical cancer data agrees with these studies; among early-stage

SCCs, HPV18 was associated with worse prognosis than HPV16 and for tumours harbouring HPV45

(also an α7 type), prognosis was worse still. Although the numbers are small, this observation could

help to explain why in a further study, positivity for HPV16 or 18 was associated with favourable

prognosis compared with other HPV types or no HPV detection[41].

Examining the molecular alterations and tumour microenvironment in a large set of tumours

provided insight into the complex interplay of HPV type, histology and changes to the host genome

that drive cervical carcinogenesis. Firstly, when clustering based on gene expression differences

between HPV16+ and HPV45+ tumours, a small minority of HPV16+ tumours co-clustered with the

majority of HPV45+ tumours, while HPV18+ tumours were found in both clusters. These clusters

were robust; they could be recapitulated at the DNA methylation level and in a larger independent

cohort. In both cohorts, we observed a clear difference in prognosis between the clusters, even after

inclusion of tumours displaying poor prognostic features, including locally advanced tumours,

adenocarcinomas and adenosquamous carcinomas[42]. Likewise, the survival differences were not

driven by particularly poor outcomes among HPV45+ tumours, as they remained following exclusion

of these samples (**Figure S3**) and there were very few HPV45+ samples in the validation cohort

(**Table S4**). Consistent with these observations, membership of the atypical subgroup was an

independent prognostic factor in the validation cohort.

Our aggressive, atypical subgroup is enriched for genes linked to EMT and displays significant

overlap with the EMT cluster defined for the subset of TCGA samples with RPPA data available; in

particular increased expression of YAP1, which in most cases appears to be driven by *YAP1* gene

amplification. Our finding that a subgroup of HPV16+ tumours co-cluster with tumours driven almost

exclusively by α7 HPV types suggests that cervical cancers driven by different HPV types evolve along

different trajectories but that HPV16+ tumours can occasionally develop via the atypical route more

commonly associated with the α7 types. HPV18+ tumours can likewise evolve via either route but

like the other α7 types, HPV18+ tumours are frequently atypical. We postulate that the host

immune response is the key driver here: a cervical cancer developing in the presence of a stronger

immune response will become an atypical tumour, hence the greater neoantigen depletion and

immune-evasive/suppressive features displayed in the tumours at time of resection, including M2

macrophage polarization, increased neutrophil abundance, reduced CTL:Treg ratios and increased

ECM deposition. By extension, this implies that α7 HPV types elicit more effective

immunosurveillance than HPV16 and the other α9 types, however when HPV16-transformed cells do

encounter an effective immune response, the resulting tumours develop down the atypical

trajectory. This model could also explain why HPV16 infections are at much higher risk to progress to

CIN3 than other HPV types[3]. Greater immunogenicity of α7 HPVs might also explain why they are so

rarely, if ever, associated with cancers developing at sites such as the oropharynx, where HPV16

dominates but other α9 types are occasionally seen[43].

Two genomic features of atypical tumours are particularly likely to be selected for in cells under

immune surveillance, *STK11* and *YAP1*. *STK11* loss-of-function drives immunosuppressive TGF-beta

signalling[44] and activates glycolysis, in turn suppressing T-cell homing and activity[45][46,47] . Consistent

with this, *STK11* mutations have recently been implicated in mediating resistance to PD1/PD-L1

blockade [48]. *YAP1*, is frequently amplified in atypical tumours, has been shown to drive cervical

cancer development in mice even in the absence of HPV, and strongly synergizes with HPV16 E6 and

E7 to promote tumorigenesis [49,50]. It is also associated with poor prognosis, reduced lymphocyte

activation and resistance to PD1/PD-L1 blockade in HNSCC [51]. Downstream of these genomic

alterations, TGF-beta is known to modulate several features of the microenvironment observed in

atypical cancers: it suppresses NK cell activation[52]; drives M2-polarisation in macrophages[53], induces

neutrophil switching to an immunosuppressive phenotype[54] and drives immunotherapy resistance

through T-cell exclusion by CAFs[55] in addition to its established role in driving metastasis. Our

analysis suggests that typical cervical cancers, with their higher mutation burdens and greater T-

lymphocyte infiltration will be good candidates for immunotherapy. What therapeutic strategies

might be efficacious in the atypical tumours, which display worse prognosis, even when resected at

early stage? Although TGF-beta inhibition for cancer treatment has thus far been limited by toxicity,

inhibiting NOX4, an NADPH oxidase required for fibroblast differentiation into ECM-depositing

myofibroblasts has recently been shown to permit CTL infiltration and to potentiate immunotherapy

in a mouse model of HPV-associated cancer [56]. GKT137831, the NOX4 inhibitor used in this study is

already approved for use in fibrosis, so could readily be trialled in patients. In those atypical cervical

cancers harbouring loss-of-function *STK11* mutations, treatment with the mitochondrial inhibitor

Phenformin is a possible therapeutic option[57].

In summary, by assembling the largest multi-omics cervical cancer dataset to date, we have gained

novel insights into the development and progression of this disease. It has also allowed us to

develop a prognostic classification that captures variation in HPV type, host genomic alterations and

the tumour microenvironment and offers the potential for stratification of cervical cancer patients

for improved clinical management. Genome-wide DNA methylation profiling is already being used in

the clinic for diagnosis and stratification of brain tumours[58], thus a similar strategy could readily be

tested for cervical cancer.

Methods

Patient samples

All patients gave written, informed consent before inclusion. Samples from Bergen were collected in

a population based setting from patients treated at the Department of Obstetrics and Gynaecology,

Haukeland University Hospital, Bergen, Norway, from May 2001 to May 2011. The study has been

approved by the regional ethical committee (REK 2009/2315, 2014/1907 and 2018/591). For more

details on sample collection see[7,42]. Samples from Innsbruck were collected and processed at the

Department of Obstetrics and Gynaecology of the Medical University of Innsbruck. The study was

reviewed and approved by the Ethics committee of the Medical University of Innsbruck (reference

number: AN2016-0051 360/4.3; 374/5.4: 'Biobank study: Validation of a DNA-methylation based

signature in cervical cancer') and conducted in accordance with the Declaration of Helsinki. Samples

from Oslo (n = 268) were collected from patients participating in a previously published prospective

clinical study[59] approved by the Regional Committee for Medical Research Ethics in southern Norway

(REK no. S-01129). Limited quantities of patient tumour samples and extracted DNA may remain and

the distribution of these materials is subject to ethical approval at the institutions from which they

were collected. Those interested in working with these samples should contact the authors to

discuss their requirements.

Dataset assembly

DNA methylation (Illumina Infinium 450k array) and RNAseq data were obtained for CESC from the

TCGA data portal. TCGA mutation data were obtained from the MC3 project on SAGE Synapse

(syn7214402). DNA methylation (Illumina Infinium 450k array) and gene expression (Illumina

HumanHT-12 V4.0 expression beadchip) data from the Oslo cohort were obtained from the Gene

Expression Omnibus (GSE68339). RNAseq data were obtained for the Bergen cohort from dbGaP

(phs000600/DS-CA-MDS 'Genomic Sequencing of Cervical Cancers') under the authorisation of

project #14589 "Investigating the mechanisms by which viruses and carcinogens contribute to

cancer development" and were converted to fastq files using SRA-dump from the SRA Toolkit

(http://ncbi.github.io/sra-tools/). Kallisto[60] was then used to quantify expression of GENCODE

GrCh37 transcripts, repbase repeats and transcripts from 20 different high-risk HPV types with bias

correction. Where IDAT files for 450k data were available, they were parsed using *minfi*[61] and were

subjected to Functional Normalisation[62], followed by BMIQ-correction[63] for probe type distribution

(which was done for all methylation data). For TCGA samples, viral type allocation was performed

using VirusSeq[8].

Generation of 450k methylation profiles.

100ng DNA was bisulphite converted using the EZ DNA Methylation kit (Zymo Research) as per

manufacturer's instructions. Bisulphite converted DNA hybridised to the Infinium 450K Human

Methylation array, and processed in accordance with the manufacturer's recommendations.

23
24   *HPV typing.*
25

HPV16 or 18 was detected in 230 samples from the Oslo cohort by PCR, using the primers listed in[64].

The PCR products were detected by polyacrylamide gene electrophoresis or the Agilent DNA 1000 kit

(Agilent Technologies Inc, Germany). Samples from the Innsbruck cohort and the remaining non-

HPV16/18 samples from the Oslo cohort (n=38) were HPV-typed by DDL Diagnostic Laboratory

(Netherlands) using the SPF10 assay, in which a PCR-based detection of over 50 HPV types is

followed by a genotyping assay (LIPA$_{25}$) that identifies 25 HPV types (HPV 6, 11, 16, 18, 31, 33, 34,

35, 39, 40, 42, 43, 44, 45, 51, 52, 53, 54, 56, 58, 59, 66, 68/73, 70 and 74). HPV type data for the

remaining samples were published previously[5,7]

Prognostic analyses and tumour clustering

Multidimensional visualisation of clusters was performed using Rtsne R package. For tSNE analysis of

DEGs between only HPV16 and HPV45, 15 principal components were considered, at a perplexity of

11 over 1000 iterations. For tSNE analysis of MVPs (using BMIQ normalised data) between only

HPV16 and HPV45, 11 principal components were considered, at a perplexity of 11 over 1000

iterations. Unsupervised tSNE analysis was performed on all TCGA samples considering 41 principal

components, with a perplexity of 17 over 1000 iterations. Associations between HPV type and

survival were tested for early stage (Stage I and II) CESC in the TCGA cohort containing either HPV16,

HPV18 or HPV45. Limma-voom and limma on BMIQ and Functionally-normalised 450k data were

used to identify differentially expressed genes and methylation variable positions between HPV45

and HPV16 associated tumours. Expression profiles were clustered to yield HPV45-like (atypical) and

HPV16-like (typical) cancers using the clusterCons package[65]. The caret R package and limma were

used to develop an SVM using 5 iterations of 5-fold Cross-Validation to allocate 450k samples to

these subgroups.

Samples from our validation cohort, comprise of cases from three European centres (Bergen and

Oslo in Norway and Innsbruck, Austria) were binned into these categories, along with TCGA samples

not in the original training set, and taken together were used for subsequent statistical analyses to

identify genomic and microenvironmental correlates. Survival analyses of epigenetic allocations

were carried out using Cox Proportional Hazards regression with stratification by histology, and with

surgery, radiotherapy and chemotherapy (given/not given) as covariates. For all clinical analyses,

stages were collapsed into Stages I, II, III and IV.

Pathway analyses

Pathways were analysed with Ingenuity Pathway Analysis. Settings used were experimentally

validated interactions in human models. Z-score cutoffs of 2 and FDR cutoffs of 0.05 were used to

identify significant hits from Canonical Pathways, Upstream Regulator and Functions ontology

analyses for plotting.

Copy number analysis

450k total intensities (Methylated and Unmethylated values) were used to generate copy number

profiles with normal blood samples from Renius et al[66] as the germline reference. Functional

normalisation [62] was used to regress out technical variation across the reference and tumour

datasets before merging and quantile normalisation was used to normalise combined intensities

followed by Circular Binary Segmentation as previously described[20]. Median density peak correction

was performed to ensure centering before further analysis. GISTIC2.0[67] was then used to identify

regions of significant copy number change at both arm and gene levels. Candidate copy number

changes were evaluated for association with cluster using binomial GLMs. The parameters chosen

were a noise threshold of 0.1 with arm-level peel off and a confidence level of 0.95 was used to

nominate genes targeted by copy number changes. Binomial regression was finally used to estimate

rates of differential alteration.

Mutational analyses

For TCGA data, mutation calls were obtained from SAGE synapse as called by the MC3 project.

Mutations for the Bergen cohort were obtained from[7]. Binomial GLMs were then used to estimate

associations between the aggressiveness clusters and mutation frequencies.

Immunological Analyses

MHC-class I neoantigens were retrieved using The Cancer Immunome Atlas[21] for TCGA samples.

Immunoediting estimates were computed based on silent mutation rates as previously described in

Rooney et al[68] . Comparisons with immune infiltration in HPV+ Head and Neck Tumours were carried

out using a previously published, manually-curated signature of immune-checkpoints, infiltration

markers and effector molecules[9]. MethylCIBERSORT[27] was used to estimate tumour purity and

abundances of seven other microenvironmental cellular fractions. Monocyte polarisation was

computed using CIBERSORT on the basis of the original LM22 matrix (1000 permutations, data

supplied in counts per million) provided with the software. We normalised the estimates to total

monocyte fractions and estimated the fraction of proinflammatory (M1 and dendritic cells) relative

to all monocytes to yield a proinflammatory monocyte fraction, which was tested for associations

with prognostic cluster using Wilcoxon's Rank Sum Test.

To generate a neutrophil gene expression signature for assessing neutrophil content by ssGSEA,

RNA-seq data were downloaded from the European Nucleotide Archive for the following datasets -

PRJEB11844[69], GSE60424[70], and E-MTAB-2319[71] in order to derive an RNAseq dataset of immune cell

types. *Kallisto*[60] was used to quantify gene expression with a reference transcriptome consisting of

Gencode Grch37 assembly of protein coding and lincRNA transcripts. Data were then modelled using

limma trend and overexpressed markers (3FC, FDR < 0.05) were selected for each cell subset from

one versus all comparisons.

The GSVA R package was then used to compute ssGSEA scores for T-cell subsets of interest with

absolute ranking, score normalisation and RNA-seq flags set to true. Enrichment scores were then

normalised by cellular abundance and differences were estimated using Wilcoxon's rank sum test

with Benjamini Hochberg correction for multiple testing.

Immunohistochemistry

All immunohistochemical staining was conducted by HSL-Advanced Diagnostics (London, UK) using

the Leica Bond III platform with Leica Bond Polymer Refine detection as per manufacturer's

recommendations. Sections from a series of 17 tumour samples from the validation cohort were

stained for CD8 (mouse monoclonal 4B11, Leica Biosystems PA0183, used as supplied for 15 minutes

at room temperature. HIER was performed on-board using Leica ER2 solution (high pH) for 20

minutes), CD68 (mouse monoclonal PGM1, Agilent M087601-2, used at a dilution of 1/50 for 15mins

at room temperature. HIER was performed on-board using Leica ER1 solution (low pH) for 20

minutes) or MPO (rabbit polyclonal, Agilent A039829-2, used at a dilution of 1/4000 for 15 minutes

at room temperature without epitope retrieval. Scoring was performed blinded to cluster

membership by a histopathologist (JM) as follows: 0 = no positive cells / field (200X magnification); 1

= 1 − 10 positive cells; 2 = 11 − 100 positive cells; 3 = 101 − 200 positive cells; 4 = 201 = 300 positive

cells; 5 = over 300 positive cells.

Data availability

Illumina Infinium 450k array DNA methylation data generated in-house from Bergen and Innsbruck

validation cohort samples (used to generate Figure 2C and for all subsequent analyses based on

DNA methylation-derived clustering of validation cohort samples and DNA methylation-based copy

number estimates)   have been deposited in the Gene Expression Omnibus (accession number

GSEXXXXX (*to be deposited upon publication*)). RNA-seq data from the Bergen samples are available

under controlled access from dbGAP (accession number phs000600/DS-CA-MDS 'Genomic

Sequencing of Cervical Cancers') subject to authorisation from NCBI. DNA methylation and gene

expression data from the Oslo samples are available from the Gene Expression Omnibus (accession

number GSE68339) and all TCGA data used are available through the TCGA Data Portal (DNA

methylation and RNA-seq) or the SAGE Synapse (DNA mutation data, accession number

syn7214402).

Code availability

All packages used have been published, are freely available and are referenced in the methods. R markdowns used to run the analyses specific to this study are available from the authors on request.

Acknowledgements

AC was supported by postgraduate research scholarships from UCL and received additional research support from a Debbie Fund grant to KC. Research in TRF's lab is supported by Rosetrees Trust (M229-CD1) and Cancer Research UK (A25825). DNA methylation data were generated through funding provided by the Debbie Fund to KC and analyses were carried out in part using data generated by The Cancer Genome Atlas. AF was supported by grants from the MRC (MR/M025411/1), PCUK(MA-TR15-009), BBSRC (BB/R009295/1), TUF, Orchid and the UCLH BRC. The authors dedicate this manuscript to the late Dr Helga Salvesen, a wonderful collaborator and colleague who played a key role in the project.

Disclosures

United Kingdom Patent Application No.1703406.7 was filed based on some of the results in this paper (TRF, AC, KC, AF).

Contributions

TRF, AC, AF and KC conceived the project; AC and SH carried out computational analyses; AF, CD, MJ, NK, NE, JM performed experiments; HL, HF, HS, XS, CSF, MKH, JD, CK, AS, SS provided samples and data. AC and TRF and AF wrote the paper and TRF supervised the study.

References

1.   Li, N., Franceschi, S., Howell-Jones, R., Snijders, P.J.F. & Clifford, G.M. Human papillomavirus type distribution in 30,848 invasive cervical cancers worldwide: Variation by geographical region, histological type and year of publication. *International Journal of Cancer* **128**, 927-935 (2011).

2.   Villa, L.L. & Schlegel, R. Differences in transformation activity between HPV-18 and HPV-16 map to the viral LCR-E6-E7 region. *Virology* **181**, 274-377 (1991).

3.   Sand, F.L., Munk, C., Frederiksen, K., Junge, J., Iftner, T., Dehlendorff, C. & Kjaer, S.K. Risk of CIN3 or worse with persistence of 13 individual oncogenic HPV types. *Int J Cancer* 144, 1975-82 (2019).

4.   Feber, A. *et al.* Epigenetics markers of metastasis and HPV-induced tumorigenesis in penile cancer. *Clin Cancer Res* **21**, 1196-206 (2015).

5.   Cancer Genome Atlas Research, N. *et al.* Integrated genomic and molecular characterization of cervical cancer. *Nature* **543**, 378-384 (2017).

6.   Lando, M. *et al.* Interplay between promoter methylation and chromosomal loss in gene silencing at 3p11-p14 in cervical cancer. *Epigenetics* **10**, 970-80 (2015).

7.   Ojesina, A.I. *et al.* Landscape of genomic alterations in cervical carcinomas. *Nature* **506**, 371-5 (2014).

8.   Chen, Y. *et al.* VirusSeq: software to identify viruses and their integration sites using next-generation sequencing of human cancer tissue. *Bioinformatics* **29**, 266-7 (2013).

9.   Chakravarthy, A. *et al.* Human Papillomavirus Drives Tumor Development Throughout the Head and Neck: Improved Prognosis Is Associated With an Immune Response Largely Restricted to the Oropharynx. *J Clin Oncol* **34**, 4132-4141 (2016).

10.   Barbie, D.A. *et al.* Systematic RNA interference reveals that oncogenic KRAS-driven cancers require TBK1. *Nature* **462**, 108-12 (2009).

11.   Kaufhold, S. & Bonavida, B. Central role of Snail1 in the regulation of EMT and resistance in cancer: a target for therapeutic intervention. *J Exp Clin Cancer Res* **33**, 62 (2014).

12.   Park, J. & Schwarzbauer, J.E. Mammary epithelial cell interactions with fibronectin stimulate epithelial-mesenchymal transition. *Oncogene* **33**, 1649-57 (2014).

13.   Reymond, N., d'Agua, B.B. & Ridley, A.J. Crossing the endothelial barrier during metastasis. *Nat Rev Cancer* **13**, 858-70 (2013).

14.   Skobe, M. *et al.* Induction of tumor lymphangiogenesis by VEGF-C promotes breast cancer metastasis. *Nat Med* **7**, 192-8 (2001).

15.   Steeg, P.S. Targeting metastasis. *Nat Rev Cancer* **16**, 201-18 (2016).

16.   Liu, Y. *et al.* Increased EZH2 expression is associated with proliferation and progression of cervical cancer and indicates a poor prognosis. *Int J Gynecol Pathol* **33**, 218-24 (2014).

17.   Halle, C. *et al.* Hypoxia-induced gene expression in chemoradioresistant cervical cancer revealed by dynamic contrast-enhanced MRI. *Cancer Res* **72**, 5285-95 (2012).

18.   Kim, B.W. *et al.* Prognostic assessment of hypoxia and metabolic markers in cervical cancer using automated digital image analysis of immunohistochemistry. *J Transl Med* **11**, 185 (2013).

19.   Zhang, Y., Liu, B., Zhao, Q., Hou, T. & Huang, X. Nuclear localizaiton of beta-catenin is associated with poor survival and chemo-/radioresistance in human cervical squamous cell cancer. *Int J Clin Exp Pathol* **7**, 3908-17 (2014).

20.   Feber, A. *et al.* Using high-density DNA methylation arrays to profile copy number alterations. *Genome Biol* **15**, R30 (2014).

21.   Charoentong, P. *et al.* Pan-cancer Immunogenomic Analyses Reveal Genotype-Immunophenotype Relationships and Predictors of Response to Checkpoint Blockade. *Cell Rep* **18**, 248-262 (2017).

22.   Gilbert, D.C. *et al.* Tumour-infiltrating lymphocyte scores effectively stratify outcomes over and above p16 post chemo-radiotherapy in anal cancer. *Br J Cancer* (2016).

23. Gooden, M.J.M., de Bock, G.H., Leffers, N., Daemen, T. & Nijman, H.W. The prognostic
influence of tumour-infiltrating lymphocytes in cancer: a systematic review with meta-analysis.
British Journal Of Cancer **105**, 93 (2011).
24. Jordanova, E.S. *et al.* Human leukocyte antigen class I, MHC class I chain-related molecule A,
and CD8+/regulatory T-cell ratio: which variable determines survival of cervical cancer patients? *Clin*
Cancer Res **14**, 2028-35 (2008).
25. Nedergaard, B.S., Ladekarl, M., Thomsen, H.F., Nyengaard, J.R. & Nielsen, K. Low density of
CD3+, CD4+ and CD8+ cells is associated with increased risk of relapse in squamous cell cervical
cancer. *British Journal Of Cancer* **97**, 1135 (2007).
26. Ward, M.J. *et al.* Tumour-infiltrating lymphocytes predict for outcome in HPV-positive
oropharyngeal cancer. *Br J Cancer* **110**, 489-500 (2014).
27. Chakravarthy, A. *et al.* Pan-cancer deconvolution of tumour composition using DNA
methylation. *Nat Commun* **9**, 3220 (2018).
28. Carter, S.L. *et al.* Absolute quantification of somatic DNA alterations in human cancer. *Nat*
Biotechnol **30**, 413-21 (2012).
29. Mizunuma, M. *et al.* The pretreatment neutrophil-to-lymphocyte ratio predicts therapeutic
response to radiation therapy and concurrent chemoradiation therapy in uterine cervical cancer. *Int*
J Clin Oncol **20**, 989-96 (2015).
30. Lee, Y.Y. *et al.* Pretreatment neutrophil:lymphocyte ratio as a prognostic factor in cervical
carcinoma. *Anticancer Res* **32**, 1555-61 (2012).
31. Condamine, T. & Gabrilovich, D.I. Molecular mechanisms regulating myeloid-derived
suppressor cell differentiation and function. *Trends Immunol* **32**, 19-25 (2011).
32. Zelenay, S. *et al.* Cyclooxygenase-Dependent Tumor Growth through Evasion of Immunity.
Cell **162**, 1257-70 (2015).
33. Elkabets, M. *et al.* IL-1beta regulates a novel myeloid-derived suppressor cell subset that
impairs NK cell development and function. *Eur J Immunol* **40**, 3347-57 (2010).
34. Chun, E. *et al.* CCL2 Promotes Colorectal Carcinogenesis by Enhancing Polymorphonuclear
Myeloid-Derived Suppressor Cell Population and Function. *Cell Rep* **12**, 244-57 (2015).
35. Hartwig, T. *et al.* The TRAIL-Induced Cancer Secretome Promotes a Tumor-Supportive
Immune Microenvironment via CCR2. *Mol Cell* **65**, 730-742.e5 (2017).
36. Xu, M. *et al.* Interactions between interleukin-6 and myeloid-derived suppressor cells drive
the chemoresistant phenotype of hepatocellular cancer. *Exp Cell Res* **351**, 142-149 (2017).
37. Newman, A.M., Liu, C.L. & Green, M.R. Robust enumeration of cell subsets from tissue
expression profiles. **12**, 453-7 (2015).
38. Lai, C.H. *et al.* Role of human papillomavirus genotype in prognosis of early-stage cervical
cancer undergoing primary surgery. *J Clin Oncol* **25**, 3628-34 (2007).
39. Burger, R.A. *et al.* Human papillomavirus type 18: association with poor prognosis in early
stage cervical cancer. *J Natl Cancer Inst* **88**, 1361-8 (1996).
40. Wang, C.C. *et al.* Clinical effect of human papillomavirus genotypes in patients with cervical
cancer undergoing primary radiotherapy. *Int J Radiat Oncol Biol Phys* **78**, 1111-20 (2010).
41. Cuschieri, K. *et al.* Influence of HPV type on prognosis in patients diagnosed with invasive
cervical cancer. *Int J Cancer* **135**, 2721-6 (2014).
42. Halle, M.K. *et al.* Clinicopathologic and molecular markers in cervical carcinoma: a
prospective cohort study. *Am J Obstet Gynecol* **217**, 432 e1-432 e17 (2017).
43. Gillison, M.L., Chaturvedi, A.K., Anderson, W.F. & Fakhry, C. Epidemiology of Human
Papillomavirus-Positive Head and Neck Squamous Cell Carcinoma. *J Clin Oncol* **33**, 3235-42 (2015).
44. Moren, A., Raja, E., Heldin, C.H. & Moustakas, A. Negative regulation of TGFbeta signaling by
the kinase LKB1 and the scaffolding protein LIP1. *J Biol Chem* **286**, 341-53 (2011).
45. Faubert, B. *et al.* Loss of the tumor suppressor LKB1 promotes metabolic reprogramming of
cancer cells via HIF-1alpha. *Proc Natl Acad Sci U S A* **111**, 2554-9 (2014).

46.   Ottensmeier, C.H. *et al.* Upregulated glucose metabolism correlates inversely with CD8+ T
cell infiltration and survival in squamous cell carcinoma. *Cancer Res* (2016).
47.   Ngwa, V.M., Edwards, D.N., Philip, M. & Chen, J. Microenvironmental Metabolism Regulates
Antitumor Immunity. *Cancer Res* **79**, 4003-4008 (2019).
48.   Skoulidis, F. *et al.* STK11/LKB1 Mutations and PD-1 Inhibitor Resistance in KRAS-Mutant Lung
Adenocarcinoma. *Cancer Discov* **8**, 822-835 (2018).
49.   Eun, Y.G. *et al.* Clinical significance of YAP1 activation in head and neck squamous cell
carcinoma. *Oncotarget* **8**, 111130-111143 (2017).
50.   He, C. *et al.* A Human Papillomavirus-Independent Cervical Cancer Animal Model Reveals
Unconventional Mechanisms of Cervical Carcinogenesis. *Cell Rep* **26**, 2636-2650 e5 (2019).
51.   Ayers, M. *et al.* IFN-gamma-related mRNA profile predicts clinical response to PD-1 blockade.
J Clin Invest **127**, 2930-2940 (2017).
52.   Viel, S. *et al.* TGF-beta inhibits the activation and functions of NK cells by repressing the
mTOR pathway. *Sci Signal* **9**, ra19 (2016).
53.   Zhang, F. *et al.* TGF-beta induces M2-like macrophage polarization via SNAIL-mediated
suppression of a pro-inflammatory phenotype. *Oncotarget* **7**, 52294-52306 (2016).
54.   Fridlender, Z.G. *et al.* Polarization of tumor-associated neutrophil phenotype by TGF-beta:
"N1" versus "N2" TAN. *Cancer Cell* **16**, 183-94 (2009).
55.   Mariathasan, S. *et al.* TGFbeta attenuates tumour response to PD-L1 blockade by
contributing to exclusion of T cells. *Nature* **554**, 544-548 (2018).
56.   Ford, K. *et al.* NOX4 inhibition potentiates immunotherapy by overcoming cancer-associated
fibroblast-mediated CD8 T-cell exclusion from tumours. *Cancer Res* (2020).
57.   Shackelford, D.B. *et al.* LKB1 inactivation dictates therapeutic response of non-small cell lung
cancer to the metabolism drug phenformin. *Cancer Cell* **23**, 143-58 (2013).
58.   Capper, D. *et al.* DNA methylation-based classification of central nervous system tumours.
Nature **555**, 469-474 (2018).
59.   Lando, M. *et al.* Identification of eight candidate target genes of the recurrent 3p12-p14 loss
in cervical cancer by integrative genomic profiling. *J Pathol* **230**, 59-69 (2013).
60.   Bray, N.L., Pimentel, H. & Melsted, P. Near-optimal probabilistic RNA-seq quantification. *Nat*
Biotechnol **34**, 525-7 (2016).
61.   Aryee, M.J. *et al.* Minfi: a flexible and comprehensive Bioconductor package for the analysis
of Infinium DNA methylation microarrays. *Bioinformatics* **30**, 1363-9 (2014).
62.   Fortin, J.P. *et al.* Functional normalization of 450k methylation array data improves
replication in large cancer studies. *Genome Biol* **15**, 503 (2014).
63.   Teschendorff, A.E. *et al.* A beta-mixture quantile normalization method for correcting probe
design bias in Illumina Infinium 450 k DNA methylation data. *Bioinformatics* **29**, 189-96 (2013).
64.   Lyng, H. *et al.* Intratumor chromosomal heterogeneity in advanced carcinomas of the uterine
cervix. *Int J Cancer* **111**, 358-66 (2004).
65.   Simpson, T.I., Armstrong, J.D. & Jarman, A.P. Merged consensus clustering to assess and
improve class discovery with microarray data. *BMC Bioinformatics* **11**, 590 (2010).
66.   Reinius, L.E. *et al.* Differential DNA methylation in purified human blood cells: implications
for cell lineage and studies on disease susceptibility. *PLoS One* **7**, e41361 (2012).
67.   Mermel, C.H. *et al.* GISTIC2.0 facilitates sensitive and confident localization of the targets of
focal somatic copy-number alteration in human cancers. *Genome Biol* **12**, R41 (2011).
68.   Rooney, M.S., Shukla, S.A., Wu, C.J., Getz, G. & Hacohen, N. Molecular and genetic
properties of tumors associated with local immune cytolytic activity. *Cell* **160**, 48-61 (2015).
69.   De Simone, M. *et al.* Transcriptional Landscape of Human Tissue Lymphocytes Unveils
Uniqueness of Tumor-Infiltrating T Regulatory Cells. *Immunity* **45**, 1135-1147 (2016).
70.   Linsley, P.S., Speake, C., Whalen, E. & Chaussabel, D. Copy number loss of the interferon
gene cluster in melanomas is linked to reduced T cell infiltrate and poor patient prognosis. *PLoS One*
9, e109760 (2014).

71.   Bonnal, R.J. *et al*. De novo transcriptome profiling of highly purified human lymphocytes
primary cells. *Sci Data* **2**, 150051 (2015).