Expansion of frozen hybrids in guppy ectoparasite, *Gyrodactylus turnbulli*

Authors: Mateusz Konczal[1], Karolina J. Przesmycka[1], Ryan S. Mohammed[2,3], Christoph Hahn[4],

Jo Cable[3], Jacek Radwan[1]

[1] Evolutionary Biology Group, Faculty of Biology, Adam Mickiewicz University, 61-614

Poznań, Poland.

[2] The University of the West Indies Zoology Museum, Department of Life Sciences, Faculty of

Science and Technology, UWI, St. Augustine, Trinidad and Tobago, WI.

[3] School of Biosciences, Cardiff University, Cardiff, CF10 3AX, UK.

[4] Institute of Biology, University of Graz, Austria

* Corresponding author

E-mail: mateusz.konczal@amu.edu.pl (MK)

Keywords: hybridization, clonal reproduction, heterosis, invasion, *Gyrodactylus*

Abstract

Hybridization is one of the major factors contributing to the emergence of highly successful parasites.

Hybrid vigor can play an important role in this process, but its contribution to the evolutionary

success of parasites has not been demonstrated. Here, we identify an invasion of a "frozen hybrid"

genotype in natural populations of *Gyrodactylus turnbulli*, ectoparasitic flatworms with facultative

sexual reproduction and significant impact on cultured fish. We re-sequenced genomes of these

parasites infecting guppies from six Trinidad and Tobago populations, and found surprisingly high

discrepancy in genome-wide nucleotide diversity between islands. The elevated heterozygosity on

Tobago is maintained by predominantly clonal reproduction of hybrids formed from two diverged

genomes. Hybridization has been followed by spread of the hybrids across the island, implying a

selective advantage compared to native genotypes. Our results highlight the potential consequences

of heterosis in the epidemiology of parasite hybrids and its impact on population genomic diversity.

Introduction

Hybridization can be a potent evolutionary force, creating opportunities for adaptive evolution by

increasing genetic variation on which natural selection can act (Arnold, 1997; Oziolor et al., 2019;

Rieseberg et al., 2003; Taylor and Larson, 2019). The ecological and evolutionary consequences are

potentially large, and not just for the hybrid organism itself: such hybridization may, for example,

exert strong influence on the evolution of pathogenic organisms ('parasites' henceforth) (King et al.,

2015). This role is likely to increase in importance because human activities increase opportunities

for hybridization by inducing environmental changes and incidental translocations (Brooks and

Hoberg, 2007; Cable et al., 2017; Patz et al., 2000). Understanding the role of hybridization in

parasite evolution is therefore essential if we are to predict hot spots of disease risk and emergence

(Laine et al., 2019).

Parasite strains of hybrid origin can gain an advantage because recombinants produced after

hybridization may show enhanced phenotypic characteristics compared to the parental generation,

such as higher infectivity (Grigg et al., 2001), expanded host range (Detwiler and Criscione, 2010),

increased transmission potential (Volf et al., 2007) or increased tolerance to harsh environmental

conditions (Laine et al., 2019). The probability of hybridization increases when species co-infect the

same host, and such systems may be considered hotspots for studying its phenotypic effects.

However, hybridization is often associated with asexual reproduction, precluding benefits resulting

from recombination between divergent genomes. Asexual reproduction is thought to be enforced by

Bateson-Dobzhansky-Muller incompatibilities accumulated in parental lineages (Neaves and Baumann,

2011). The genomes of parental species, divergent enough to disrupt meiosis in hybrids, yet not

diverged enough to seriously affect hybrid viability, could drive asexual reproduction of hybrids

(Janko et al., 2018; Moritz et al., 1989). In fact, many asexually reproducing organisms have

originated from hybridization events (Neaves and Baumann, 2011).

Conversely, asexual reproduction may be selectively beneficial if clonal reproduction "freezes"

heterozygosity that confers short-term evolutionary advantage (Neiman et al., 2014). Hybrids often

show greater vigor, productivity, disease resistance, or overall increased Darwinian fitness

(Kingsbury, 2009). Two alternative hypotheses may explain the existence of hybrid vigor (heterosis)

(Lippman and Zamir, 2007). The overdominance hypothesis, attributes heterosis to superior fitness of

heterozygous genotypes over homozygous wild types. The dominance hypothesis assumes that

recessive deleterious mutations are to some extent masked in a heterozygous stage. Recent

theoretical work indicates that dominance-induced heterosis might play an important role in

evolution, especially if deleterious alleles are not removed effectively from populations as is

expected when populations are small (Gilbert et al., 2020; Kim et al., 2018; Zhao and Charlesworth,

2016). This mechanism may be particularly relevant to parasite evolution. Many parasite populations

are small or experience frequent bottlenecks, therefore they are prone to accumulation of

deleterious mutations (Lynch et al., 1993), but also to elevated homozygosity due to inbreeding. In

such case, one could expect higher fitness of hybrid parasites. Clonal reproduction can then preserve

heterosis in so called 'frozen hybrids' and facilitate expansion of successful parasites. However, a

case of successful frozen hybrid parasites has not, to our knowledge, been demonstrated.

Here, we report genomic analysis of the guppy parasite, *Gyrodactylus turnbulli*, which allowed us to

infer the expansion of a predominantly clonal hybrid strain in a species otherwise reported to

regularly reproduce sexually (Schelkle et al., 2012). Interactions between gyrodactylids and guppies

have been the subject of intensive eco-evolutionary research focusing on coevolution between host

and parasites (e.g. Houde & Torio, 1992; Jacquin et al., 2016; Phillips et al., 2018; Reynolds, Arapi, &

Cable, 2018; van Oosterhout, Harris, & Cable, 2003), while the effect of different reproductive

strategies (Cable and Harris, 2002) on population structure and dynamics of infections in the parasite

wild populations are poorly understand (Xavier et al., 2015). Our earlier work on another

monogenean guppy parasite, *G. bullatarudis*, identified the potential of recombination between

divergent strains in the emergence of invasive parasites (Konczal et al., in press). In contrast, the

results reported here indicate the importance of heterosis in rapid clonal invasions of *G. turnbulli* and

highlight that hybridization can lead to the emergence of highly successful parasites, despite

suppressing recombination.

Results

A total of 30 *G. turnbulli* samples from three Trinidadian (Aripo, Caura and Lopinot Rivers) and three Tobagonian populations (Dog River, Health Center and Spring Site; Figure 1A, details in Supplementary Table 1) collected in two consecutive years were sequenced to an average coverage of 38.7 per sample. Sequencing reads were then mapped to the reference genome and used for SNP calling. We identified 776,881 single nucleotide- and 161,255 indel-polymorphisms. Principal Component Analyses of SNP genotypes clearly separated Trinidad and Tobago populations, while differentiation between local rivers was present only in some cases (Figure 1B). In particular, samples from Health Center and Dog River (Tobago) showed virtually no differentiation and, in contrast to the Trinidadian populations, do not form monophyletic groups according to the populations (Figure 1C). The Trinidadian samples are also more similar to the reference genome, obtained from a strain on commercial guppies, *Gt3,* than samples collected from Tobago (Figure 1C). To remove the potential effect of linkage on population differentiation, we pruned linked SNPs, but the general pattern remained the same (Supplementary Figure 1). Furthermore, a similar pattern was observed with ADMIXTURE analyses: the most likely number of genetic groups is two, with pronounced distinction between the Trinidad and Tobago samples (Supplementary Figure 2). We calculated F_{ST} between Trinidad and Tobago populations in 25 kb non-overlapping windows and found that 90% of the windows show F_{ST} values between 0.25 and 0.55 (mean F_{ST} = 0.42; Supplementary Figure 3). Pairwise F_{ST} values were higher among Trinidadian populations (Supplementary Figures 4) than among populations from Tobago (Supplementary Figures 5). To resolve whether moderate F_{ST} between islands resulted from moderate diversification or from elevated level of genetic variation within groups, we calculated nucleotide diversity across Trinidad and Tobago samples separately. Nucleotide diversity on Tobago was 2.4 times greater than that for Trinidad (0.0028 and 0.0012 respectively, Figure 1D). The difference does not result from high differentiation among Tobagonian populations: nucleotide diversity was also higher within Tobagonian populations, compared with those from Trinidad (Supplementary Figure 6).

Increased nucleotide diversity on Tobago was mainly driven by high heterozygosity of most individuals (Figures 1D/E). Similarly, inbreeding coefficient, F_{IS}, was negative for most of the samples from Tobago, suggesting over-representation of heterozygous positions (Supplementary Figure 7). In contrast, most of the Trinidadian samples showed positive values of F_{IS}.

High heterozygosity was observed in all samples from two populations on Tobago (Health Center and Dog River), whereas the third population (Spring Site) showed a mixture of patterns. One Spring Site sample (G334) had low level of heterozygosity, reminiscent of the general pattern observed in Trinidadian samples. Three other samples (G328, G367 and G368) showed bimodal distribution, with

genomic regions presenting either high or low level of heterozygosity. Finally, two Spring Site samples (G331 and G370) contained only highly heterozygous regions, similar to the pattern observed in Health Center and Dog River (Figure 1E).

Figure 1 Patterns of genetic variation in *Gyrodactylus turnbulli* populations from Trinidad and Tobago. A: Map of Trinidad and Tobago with sampling sites. **B:** Principal Component Analyses of *G. turnbulli* samples. PCA based on 673,104 SNPs which passed filtering criteria. SNPs were identified within 30 isolates from three Trinidadian and three Tobagonian populations. **C:** Neighbor Joining tree (Saitou and Nei 1987) constructed using distance matrix, with one minus identity-by-state units, calculated based on 673,104 SNPs. Populations are represented by different colours. The reference sequence (obtained from a strain on commercial guppies) is shown in black. **D:** Distribution of nucleotide diversity (π) calculated across the genome in 25 kb non-overlapping windows. **E:** Distribution of heterozygosity calculated in 25 kb non-overlapping windows for each sample collected from Trinidad (white background) and Tobago (shaded background) populations.

The majority of polymorphic Tobagonian SNPs (325,473 out of 498,111) were heterozygous in Health
Center, Dog River,and two (G331 and G370) Spring Site samples, and 88% of these SNPs contained an
allele from the reference genome and its alternative found in the Tobagonian G334 sample (Figure
2). Genome-wide pattern of divergence between re-sequenced samples and the reference genome
demonstrated that G334 sample is the most divergent, while all samples from Trinidad are more
closely related to the reference genome (Figure 1C, Supplementary Figure 8). Based on these
patterns we infer that highly heterozygous Tobagonian samples are hybrids between two divergent
haplotypes: one similar to the reference sequence and the second similar to the G334 sequence. The
same pattern with shared heterozygous sites and high, genome-wide heterozygosity was observed in
the Health Center and Dog River samples from 2017 and 2018, indicating that the hybrid population
is stable, and implying an absence of sexual reproduction (otherwise recombination would have fixed
one of the two haplotypes in at least some genomic regions). In contrast, divergent haplotypes have
clearly recombined in the Spring Site population. Three samples (G328, G367 and G368) show a
mosaic pattern of the two haplotypes fixed in different genomic locations, with other parts of the
genome still remaining heterozygous (Figure 2, Supplementary Figure 9).

Figure 2. Graphical representation of genotypes in *Gyrodactylus turnbulli* samples. Genotypes
identified as reference homozygotes are shown in green, alternative homozygotes in red,
heterozygous in yellow. Samples collected in 2017 and 2018 are labeled in grey and black
respectively, and those collected from different local populations are separated by horizontal grey
lines. A subset of polymorphic sites (10,000 SNPs) from the longest scaffolds is shown.

Signatures of sexual reproduction are also observed at least in some of the samples from Trinidad.
For example, SNPs localized in the first scaffold show admixture of another haplotype in sample G314
coming from Lopinot, and another two samples in Lopinot are heterozygous in this region (Figure 2).
Another sample from the same population (G308) has negative F_{IS} (Supplementary Figure 7) and
slightly elevated genome-wide heterozygosity (Figure 1D), although the effect is weaker compared to
samples from Tobago. Nevertheless, heterozygous sites are on average rare in Trinidadian samples.

Clustering analyses (Figure 1B, 1C) and F_{ST} results (Supplementary Figure 5) suggest low divergence
between samples identified as hybrids. We explored it further by calculating divergence in the
haploid mitochondrial genomes. One Trinidadian sample (G126) was used for mitochondrial genome
assembly, and then reads from all other samples were mapped to that genome for haplotype
reconstruction. The resulting mt genome spans 14,159 bp. We found very little divergence between
samples identified as hybrids, compared to divergence amongst other samples (Figure 3,
Supplementary Figure 10). Putative hybrid samples from Health Center, Dog River and Spring Site
populations were found more closely related to Trinidadian samples than to remaining SPS samples,
confirming introgression inferred from genomic data. Within hybrids (Health Center, Dog River two
hybrid samples from Spring Site) the mean divergence between populations was 0.02%. In contrast,
three other samples collected from Spring Site had strongly divergent mitochondria - the nucleotide
divergence between the two Spring Site groups was 1.58%. To independently verify low divergence
between Tobagonian hybrids from Tobago and samples from Trinidad we sequenced a 262 bp
fragment of COII from 81 samples collected by us on Trinidad and Tobago. The results confirmed
little divergence between all Tobagonian samples except for some derived from the Spring Site
population (Supplementary Figure 11).

Figure 3. Phylogenetic relationship between *Gyrodactylus turnbulli* samples, reconstructed based on mitochondrial genome sequences (14,159 bp). The trees were derived via a Bayesian phylogenetic method, using 10,000,000 generations of MCMC samples, with HKY substitution model and strict clock model, assuming 13.1% divergence per million years. The estimated time before present (in years) for tree nodes is shown at the bottom of the tree.

We compared genome wide nucleotide diversity of Tobagonian samples collected in 2017 and 2018 and did not find evidence for a substantial difference between time points (π_{2017} = 0.0018 vs π_{2018} = 0.0017). We also plotted relationships between heterozygosity calculated in non-overlapping windows for each pair of samples and did not find any indication for reduction in heterozygosity in any sample (Supplementary Figure 12). Little variation between years in both nucleotide diversity and heterozygosity demonstrate that the populations from Tobago reproduce mostly clonally.

The mostly clonal reproduction in Tobago hampers identification of fast evolving regions in the genome, because of strong linkage between sites. However, there was no evidence that the Trinidadian populations experienced clonal expansion, and therefore rapidly evolving genes can be potentially identified. For this reason, we tested for correlations between nucleotide diversity

calculated in windows separately for Tobagonian hybrid populations and for all Trinidadian samples.
We used nucleotide diversity calculated jointly for Health Center and Dog River samples as a proxy
for the long-term evolution of genomic regions (divergence between two haplotypes forming
hybrids). Regions in which the nucleotide diversity calculated jointly for all samples from Trinidad
deviates from predictions based on this long-term evolutionary rate are therefore candidates for
recent rapid evolutionary change between years or populations. The divergence calculated in 25 kb
windows were highly correlated between Tobagonian hybrid populations and Trinidadian
populations, while few windows demonstrated elevated (according to overall trend) diversity in
Trinidad (Figure 4A). One scaffold (scf7180000445747; 19.4 kb long) showed extensive divergence
between Trinidadian samples collected in 2017 and 2018 (Figure 4B). It contains 7 protein coding
genes, including putative methylotransferase NSUN5, Endonuclease/exonuclease/phosphatase
family domain-containing protein 1, synaptogyrin and prohibitin.

Figure 4 Genetic variation in genomic region with elevated nucleotide diversity in populations from
Trinidad. A: Correlation between nucleotide diversity calculated jointly for Health Center and Dog

River populations and nucleotide diversity calculated jointly for all samples from Trinidad. Black line

represents linear regression (y = 0.00 + 0.38*x). Scaffold scf7180000443747 is shown in red. **B:**

Graphical representation of genotypes in polymorphic sites of *Gyrodactylus turnbulli* localized in

scaffold scf7180000443747 (707 SNPs). Genotypes identified as reference homozygotes are shown in

green, alternative homozygotes in red, heterozygous in yellow. Samples collected in 2017 and 2018

are shown in grey and black respectively, and those collected from different local populations are

separated by horizontal grey lines.

Discussion

We examined whole genomes of 30 *G. turnbulli* parasites, sampled from two Caribbean islands in

two consecutive years, providing insight into the evolution of their genomic diversity. On Tobago,

parasite populations tend to have much higher nucleotide diversity than those on Trinidad. This

higher diversity was caused by the presence of highly heterozygous individuals in all studied

Tobagonian populations. These individuals shared most of the heterozygous loci, with a repeatable

pattern of having one reference-like allele and the other allele present in a homozygous state only in

one individual from Tobago (G334 sample). This pattern led us to conclude that these highly

heterozygous individuals are hybrids of two divergent lineages, one of which is close to a strain

isolated from ornamental guppies (Trinidad-like, used for the reference genome assembly), and the

second which is close to the Tobagonian G334 sample, suggesting that these two lineages originated

from different islands. In the G334 population of origin (Tobago, Spring Site) two considerably

diverged mitochondrial genomes (diverged 145,000 years ago based on 13.1% divergence per Mya,

(Kuusela et al., 2007), 95% HPD: 133,900 − 155,700) were observed. Most likely they represent

mitochondrial genomes originating from two parental populations that formed Tobagonian hybrids.

The mitochondrial divergence between these two lineages was much larger than between

Tobagonian hybrids and Trinidadian populations (7,600 years, 95% HPD: 5,200 − 9,800 years),

demonstrating that hybrids inherited mitochondria from Trinidad-like ancestors. Initial hybridization

must have been followed by predominantly clonal reproduction, as otherwise recombination would

produce both homozygous and heterozygous genotypes at least in some of the genomic regions. This

was not the case: heterozygosity was elevated genome-wide (Figure 1D) and heterozygous

genotypes were the same for all samples (Figure 2). Genetic divergence between putative hybrids as

inferred based on whole mitochondrial genomes was low. The mitochondrial divergence corresponds

to 2,700 (95% HPD: 1,500 − 4,000) years of divergence between these samples (Figure 3), indicating

that clonally reproducing hybrids spread relatively recently across Tobago. The divergence times are

based on the indirectly derived substitution rate in a single mitochondrial locus of *Gyrodactylus*

salaris (13.1% per Mya; Kuusela et al., 2007), and other indirect estimates suggest lower (5.1% per

Mya; Hahn et al., 2015) or higher divergence rate (up to 20.3% per Mya; Meinilä et al., 2004) in

Gyrodactylus species, and therefore divergence dating should be taken with caution. Nevertheless,

little nucleotide divergence between mitochondrial genomes indicates a recent origin of the hybrids,

and their widespread occurrence across Tobago provides the evidence for their evolutionary success.

Hybrid genomes that were most widespread on Tobago did not show any signs of recombination,

and apparently remained stable. This suggests that the selective advantage of the hybrids might have

arose due to heterosis. Thus, our results suggest that hybrid vigor can have dramatic consequences

for the evolution of natural parasite populations.

Hybridization of parasites can lead to the emergence of highly invasive, virulent strains and therefore

is an emerging public health and conservation concern (King et al., 2015). Human activity, migrations

and climate change shift the geographic distribution of many species (Brooks and Hoberg, 2007;

Cable et al., 2017; Lafferty, 2009), which increases the probability of contact between distinct

populations, potentially leading to hybridization followed by adaptive evolution. Parasites are often

transported with their hosts, and even if hosts do not contribute to genetic pools of local

populations, hitchhiked parasites can do so. This could be the case for *G. turnbulli*, a common guppy

parasite*. Gyrodactylus* species have no free-living stage and are transmitted primarily through direct

physical contact between hosts. Our approximate dating of divergence of mtDNA between hybrids

from Trinidad and Tobago is consistent with the end of the last glacial maximum, when a land bridge

that probably existed between the islands during glaciation-related sea level decline was disrupted

again (Alexander et al., 2006; Lambeck, 2004). This might have facilitated migration of guppies

between the islands, and possibly other fish species capable of spreading *G. turnbulli* (Cable et al.,

2013; King and Cable, 2007). Alternatively, human activity could increase the potential for

hybridization between divergent gyrodactylid strains. Guppies are commonly kept in aquaria, and *G.*

turnbulli are regularly reported in aquarium populations (Maceda-Veiga and Cable, 2019) from where

they can potentially be released to nature. The potential to increase virulence of *G. turnbulli* by

combining different genomes was suggested in an earlier report of laboratory experiments, in which

infection with two inbred stains led to higher parasite load compared to single strain infections

(Schelkle et al., 2012). However, the genomic divergence between these strains was unknown and, as

discussed by the authors, this result could be due to competition between clones leading to

evolution of increased virulence (Schelkle et al., 2012).

Our recent study on the natural populations of another monogenean parasite infecting guppies

(*Gyrodactylus bullatarudis*) suggested that recent hybridization has also played a predominant role in

shaping genetic variation, but in that case in populations from Trinidad (Konczal et al., in press).
Hybridization between two diverged linages and its subsequent recombination resulted in a mosaic
genome composition of *G. bullatarudis*. Around half of the genome originates from one or another
lineage. High divergence between recombined lineages and relatively little divergence between local
populations led us to conclude that recent hybridization was followed by rapid expansion of a
recombinant *G. bulllatarudis* strain. Similar patterns of successful admixed hybrid genotypes have
been reported for many other parasites, including schistosomes (Kincaid-Smith et al., 2019; Platt et
al., 2019), *Trypanosoma* (see Tihon et al., 2017) and *Leishmania* (see Rogers et al., 2014) species,
highlighting the role of recombination between divergent genomes in parasite evolution. Our
analysis of *G. turnbulli* genomes indicates, that unlike its congener *G. bullatarudis,* natural expansion
occurred without prior recombination, highlighting that invasive hybrid parasites can arise without
recombination. So far, there has been little evidence for clonal expansion of hybrid parasites. An
interesting suggestion comes from *Gyrodactylus salaris* – a significant pathogen of Atlantic salmon.
The Baltic-salmon specific *G. salaris* could have originated from a single hybridization event between
two clades, followed by the clonal reproduction and frozen heterozygosity, however this suggestion
is based on analyses of single nuclear marker (Kuusela et al., 2007). Indirect observations in other
systems also suggest that non-recombined hybrids might have significant evolutionary advantage.
Hybridization between schistosomes - parasites of great medical and veterinary importance - results
in higher fecundity, faster maturation time, and wider intermediate host range, although this
evidence is based only on laboratory experiments (Wright and Ross, 1980). Another potential
example of non-recombined hybrid advantage occurs in a common fungal parasite of amphibians,
Batrachochytrium dendrobatidis, in which asexually reproducing hybrids have higher virulence than
both parental strains in some host species (Greenspan et al., 2018). However, no clear evidence for
expansion of these asexually reproducing hybrids has yet emerged.

Heterosis arising from hybridization between species or strains is known for traits other than
pathogen invasiveness, and is a widespread phenomenon in plant and animal species (Lippman and
Zamir, 2007). For example, spontaneous heterosis has been demonstrated in European water frog
hybrids, increasing survival and growth rate in F1 hybrids. These frog hybrids reproduce asexually,
freezing heterozygosity, which may contribute to their ecological success (Hotz et al., 1999). Another
example, freshwater snails, has been used to demonstrate that hybrid vigor can be more important
than an increase in genetic variance in the evolution of invasiveness (Facon et al., 2005). Our results
highlight that heterosis could influence epidemiological dynamics. In such case, heterosis may then
interplay   with   other   factors   influencing   parasite   distribution,   including   for   example

metapopulational structure (Tadiri et al., 2018), host density (Johnson et al., 2011) and host

immunological gene repertoire (Phillips et al., 2018).

Parasites may be particularly prone to accumulation of deleterious mutations in their genomes

(Criscione and Blouin, 2005). In general, the rate of accumulation depends primarily on the effective

population size. In small populations, random genetic drift can overpower selection making it easier

for deleterious mutations, particularly those of small effects, to become fixed (Kimura et al., 1963).

This can cause mutational melt-down leading to population extinction (Lynch and Gabriel, 1990).

Several features of macroparasite life cycles can act in concert to reduce effective population sizes

and increase the risk of such melt-down (Criscione and Blouin, 2005). In particular, frequent

bottlenecks associated with colonization/recolonization of new hosts can significantly reduce the

efficiency of selection in removing mildly deleterious mutations from populations. Additionally,

observations of *G. turnbulli* in the wild suggest that census population sizes are often small and

fluctuating over time (Stephenson et al., 2015). If this leads to homozygosity for a number of

recessive or partially recessive mutations, hybridization between divergent strains carrying those

mutations at different genomic locations would result in hybrid vigor.

Alternatively, the advantage of hybridization can be due to alleles carried by heterozygotes being

adapted to different, and a wider range of, environments. In the case of parasites, divergent alleles

may each allow infection of different host species or genomes, allowing heterozygotes to infect a

wider range of hosts. For example, laboratory experiments on tapeworms in sticklebacks showed

that two genetic lines were only able to infect a single respective host species, whereas their hybrids

could infect both species (Henrich et al., 2013). In the case of *G. turnbulli,* however, hybridization

occurred among two strains infecting the same host species.

Clonal expansion of hybrids is likely facilitated by facultative sexual reproduction of *Gyrodactylus*

species. The first generation is always produced clonally, whereas the following can be produced

sexually or asexually (Cable and Harris, 2002). In such systems, switching between facultative and

obligate clonal reproduction does not appear to be evolutionarily constrained and can easily evolve

when favored by selection (Neiman et al., 2014). However, though it has been proposed that

hybridization is the main mechanism responsible for evolution of asexual reproduction, its causality

and underlying mechanisms are poorly understood (Janko et al., 2018; Neiman et al., 2014; Schön et

al., 2009; Xu et al., 2013). Mortiz et al. (1989) suggested that reproducing asexually results from the

accumulation of incompatibilities between hybridizing species that disrupts cellular regulation of

sexual reproduction. In fact, the majority of species which recently evolved asexual reproduction are

of hybrid origin (Neaves and Baumann, 2011). In the case of *G. turnbulli,* however, recombination is

not completely suppressed, because it was observed in one of the hybrid populations on Tobago

(Spring Site). The fact that recombinant individuals are rare on Tobago might result either from rarity

of recombination events, or from effective removal of recombinants from the population by

selection. The disadvantage of recombinants may stem from restoring homozygotes for recessive

deleterious mutations. Discrimination between these two possibilities ideally requires direct

estimates of recombination rate, possibly based on changes in genotype frequencies over time in

both hybrid and non-hybrid populations (Becheler et al., 2017).

Unlike the Tobagonian hybrids, the patterns observed in the Trinidadian *G. turnbulli* populations

suggest that recombination might be more common there. On Trinidad, only a single individual

showed slightly elevated heterozygosity, possibly as a result of sexual reproduction between two

moderately diverged strains (G308, Fig 1D, Supplementary Figure 7), and indeed another individual

from the same population showed a pattern consistent with being a subsequent generation

recombinant (G312). All other Lopinot and Aripo individuals had low genome-wide heterozygosity

and positive inbreeding coefficients (Supplementary Figure 7). This suggests relatively common

sexual reproduction because long term clonal reproduction would have led to accumulation of

different mutations on homologous chromosomes, resulting in the excess of heterozygosity, i.e.

negative inbreeding coefficients (Meselson effect; Birky, 1996; Stoeckel and Masson, 2014).

Recombination on Trinidad could, at least to some degree, have enabled independent evolution of

different genomic regions, allowing identification of genomic regions with elevated genomic

variation between the two sampling years, possibly driven by selection. We found that one genomic

region showed an extreme pattern of genetic diversity, with several interesting genes previously

associated with modulating organismal lifespan, stress resistance (Methylotrasferase NSUN5;

Schosserer et al., 2015) and host-parasite interactions (Prohibitin; Jain et al., 2010). Future work

including more individuals and experimental validation is needed to confirm involvement of these

genes in coevolutionary dynamics.

Concluding, our population genomic dataset of *G. turnbulli* allowed us to infer that hybridization

followed by clonal reproduction contributed to evolutionary success of these parasites on Tobago. It

is likely that the evolutionary success stems from heterotic effects, which are frozen in hybrids.

Benefits of such freezing may only be temporary, as the lack of recombination may hinder the

pathogen in keeping up in the evolutionary arms-race with the host's immune system. That

recombination seemed to be maintained in Trinidadian populations suggesting that it may be

favoured in the long run in *G. turnbulli,* as well. Hybrids on Tobago appear to have persisted for a

while, so it would be interesting to see if after initial success, recombination would again be favored,

as might have been the case with *G. bullatarudis* (see Konczal et al., in press). Our results thus
highlight that both heterosis and increased evolutionary potential associated with recombination
between divergent genome, can play important role in the evolution of invasive parasites, possibly at
different stages following hybridization.

Materials and Methods

Field collections

During two expeditions (2017 and 2018), guppies (*Poecilia reticulata*) were sampled from six
locations on Trinidad and Tobago (Trinidad: Aripo River [ARI], Caura River [CAU], Lopinot River [LOP];
Tobago: Dog River [DR], Health Center [HC], Spring Site [SPS]). Fish were transported to our field
station in Charlotteville, northeast Tobago, where each population was kept in separate aquarium.
Over four weeks, guppies were subsequently screened for ectoparasites under a dissecting
microscope. If gyrodactylids were observed, a single worm was allowed to move to a naïve,
anesthetized fish, with transfer closely monitored under the microscope to ensure movement of just
a single worm. Experimentally infected fish were then screened every 2-3 days for the presence of
gyrodactylids. Approximately 8-13 days post-infection, hosts were euthanized with an overdose of
MS-222, the number of worms was counted, and samples were preserved in 97% molecular grade
ethanol and transported to Poland. Samples with at least 10 worms were used for DNA extraction
and species identification as described below. Parasites collected in 2017 and identified as *G.*
bullatarudis were reported in a separate study (Konczal et al., in press), while this work focused
solely on *G. turnbulli*.

DNA extraction, libraries preparation and sequencing

DNA was extracted from each pool of individuals derived from a single worm (sample) with a MagJET
Genomic DNA kits (Thermo Scientific). DNA concentration was measured using Qubit High Sensitivity
reagents. *Gyrodactylus* were identified to species using Sanger sequencing of a 262 bp fragment of
the mitochondrial protein coding COII gene (Xavier et al., 2015). Sequences were aligned together
with records downloaded from the NCBI Genbank, and a Neighbor Joining tree was reconstructed
with MEGA X (Kumar et al., 2018) to determine species. Thirty samples were selected for library
preparation (Nextera DNA Flex Library Prep Kit, Illumina) based on DNA quantity and population of
origin. Subsets of samples were sequenced in 2018 with HiSeq2500 (2 x 100 bp). All other samples

were sequenced in 2019 on the NovaSeq6000 machine in 2 x 150 bp mode (see Supplementary Table

1 for more details).

SNP calling and population genomic analyses

Quality of raw reads was assessed with Fastqc (Andrews and Babraham Bioinformatics, 2010). Reads

were then filtered with trimmomatic (version 0.36; Bolger, Lohse, & Usadel, 2014) and mapped to

the reference genome (Hahn et al., in prep.) with bwa mem (version 0.7.10-r789). Mapping results

was visualized with Qualimap (García-Alcalde et al., 2012) and duplicated reads were marked with

picardtools (version 2.18.5-6). SNPs were then called with samtools mpileup (version 1.6.0, options -

R -C50 -t DP, ADF, ADR). Based on inspection of the distributions of diagnostic statistics, we removed

SNPs with DP > 1200 or DP < 400 (for all samples together), with QUAL < 30, with MQ < 30 or in 5 bp

distance from identified indels.

We then used SNPs to calculate PCA with plink (version 1.90, Purcell et al., 2007) using only SNPs

with minor allele frequency greater than 0.05. The same dataset with additional sample representing

reference genotype (all genotypes encoded as reference homozygous) was used to calculate distance

matrix (in one minus identity-by-state units) with plink and to construct Neighbor Joining tree with

ape package in R. In addition, we ran PCA and Admixture analyses with SNPs, that were previously

pruned for Linkage Disequilibrium. We first pruned SNPs with –indep-pairwise 100 10 0.1 option in

plink, and then used the same program to calculate PCA. Admixture (version 1.3.0; Alexander et al.,

2009) analysis was performed to determine the number of ancestral populations (K) from one to ten.

Vcftools was used to calculate nucleotide diversity (π) and genetic divergence between populations

(F_{ST}) in 25 kb non-overlapping windows, as well as inbreeding coefficient (F_{IS}, option –het) per sample

(version v0.1.12b; Danecek et al., 2011). To calculate F_{IS} per sample we considered only SNPs

polymorphic within population of origin. Custom script calculating fraction of sites being

heterozygous in given genomic window was used to estimate genome wide heterozygosity per

sample. For each sample we calculated also divergence form the reference genome in 25 kb non-

overlapping windows, by counting number of non-reference variants (alternative homozygous sites

were counted as one and reference/alternative allele heterozygous sites were counted as half) and

divide it by a window length. Windows shorter than 12.5 kb (ends of scaffolds and scaffolds shorter

than that length) were excluded from all analyses.

Mitochondrial genome assembly and mtDNA analyses

To reconstruct phylogenetic relationships between samples we reconstructed the entire
mitochondrial haplotype for sample G126 using MITObim (*de novo* approach of Hahn, Bachmann, &
Chevreux, 2013) and COII fragment downloaded from NCBI (accession KP168411.1) as an initial bait.
The resulting sequence spanned 14,159 bp. We then mapped reads from all sequenced samples to
that sequence using bwa mem with default parameters. Samples were inspected manually using IGV.
Duplicates were marked with picardtools and SNPs were called using mpileup and bcftools assuming
haploid samples. SNPs with quality lower than phred 30 were discarded from analyses. The
nucleotide sequence for each sample was reconstructed with bcftools consensus. Sequences were
then merged and used for phylogenetic analyses.

The evolutionary relationship between samples was inferred using ML with a GTR model,
implemented in MEGA X Software (Kumar et al., 2018). The same software and model were used to
calculate mean evolutionary distances between groups. The BEAST2 (v2.6.1; Bouckaert et al., 2014)
was then used to perform Bayesian phylogenetic analysis of mitochondrial genomes and to estimate
time to most recent common ancestor (TMRCA). TMRCAs of nodes of interest were estimated based
on 10,000,000 generations, logging every 1,000 steps, with the first 20% generations discarded as a
burn-in. We used a Yule Model as prior, HKY substitution model, and strict clock model. Divergence
time estimates were based on 13.1% divergence per million years estimated for *G. salaris* (see
Kuusela et al., 2007).

Data accessibility

The raw sequences will be available as FASTQ files in the GeneBank (BioProject accession no.
PRJNA603444).

Acknowledgements

We thank K. Phillips for his help in improving this manuscript, N. Cook, A. Szubert-Kruszyńska, A.
Sadowska-Konczal and staff of the Environmental Research Institute Charlotteville (ERIC), Tobago, for
support in the field; P. Turpin of Man O'War Bay Cottages for renting us the field station; Tobago
House of Assembly (THA) for granting permission to conduct field surveys; K. Dudek and J. Raubic for
help in molecular laboratory. The research was funded by Polish National Science Center Fuga Grant
UMO-509-2016/20/S/NZ8/00208. Computations were performed at the Poznan Supercomputing and
Networking Center.

References

Alexander DH, Novembre J, Lange K. 2009. Fast model-based estimation of ancestry in unrelated
individuals. *Genome Res* **19**:1655–1664. doi:10.1101/gr.094052.109

Alexander HJ, Taylor JS, Sze-Tsun Wu S, Breden F. 2006. PARALLEL EVOLUTION AND VICARIANCE IN
THE GUPPY (*POECILIA RETICULATA*) OVER MULTIPLE SPATIAL AND TEMPORAL SCALES. *Evolution*
(N Y) **60**:2352–2369. doi:10.1554/06-288.1

Andrews S, Babraham Bioinformatics. 2010. FastQC: A quality control tool for high throughput
sequence data. *Manual*. doi:citeulike-article-id:11583827

Arnold ML. 1997. Natural hybridization and evolution. Oxford University Press on Demand.

Becheler R, Masson JP, Arnaud-Haond S, Halkett F, Mariette S, Guillemin ML, Valero M, Destombe C,
Stoeckel S. 2017. ClonEstiMate, a Bayesian method for quantifying rates of clonality of
populations genotyped at two-time steps. *Mol Ecol Resour* **17**:e251–e267. doi:10.1111/1755-
0998.12698

Birky CW. 1996. Heterozygosity, heteromorphy, and phylogenetic trees in asexual eukaryotes.
Genetics **144**:427–437. doi:10.1007/978-90-481-2770-2_23

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina sequence data.
Bioinformatics **30**:2114–2120. doi:10.1093/bioinformatics/btu170

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu CH, Xie D, Suchard MA, Rambaut A, Drummond AJ.
2014. BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. *PLoS Comput Biol* **10**.
doi:10.1371/journal.pcbi.1003537

Brooks DR, Hoberg EP. 2007. How will global climate change affect parasite-host assemblages?
Trends Parasitol **23**:571–574. doi:10.1016/j.pt.2007.08.016

Cable J, Archard GA, Mohammed RS, Mcmullan M, Stephenson JF, Hansen H, van Oosterhout C.
2013. Can parasites use predators to spread between primary hosts? *Parasitology* **140**:1138–
1143. doi:10.1017/S003118201300067X

Cable J, Barber I, Boag B, Ellison AR, Morgan ER, Murray K, Pascoe EL, Sait SM, Wilson AJ, Booth M.
2017. Global change, parasite transmission and disease control: Lessons from ecology. *Philos*
Trans R Soc B Biol Sci **372**:20160088. doi:10.1098/rstb.2016.0088

Cable J, Harris PD. 2002. Gyrodactylid developmental biology: Historical review, current status and
future trends. *Int J Parasitol* **32**:255–280. doi:10.1016/S0020-7519(01)00330-7

Criscione CD, Blouin MS. 2005. Effective sizes of macroparasite populations: A conceptual model.
Trends Parasitol **21**:212–217. doi:10.1016/j.pt.2005.03.002

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT,
Sherry ST, McVean G, Durbin R. 2011. The variant call format and VCFtools. *Bioinformatics*
27:2156–2158. doi:10.1093/bioinformatics/btr330

Detwiler JT, Criscione CD. 2010. An infectious topic in reticulate evolution: Introgression and
hybridization in animal parasites. *Genes (Basel)* **1**:102–123. doi:10.3390/genes1010102

Facon B, Jarne P, Pointier JP, David P. 2005. Hybridization and invasiveness in the freshwater snail
Melanoides tuberculata: Hybrid vigour is more important than increase in genetic variance. *J*
Evol Biol **18**:534–535. doi:10.1111/j.1420-9101.2005.00887.x

García-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, Götz S, Tarazona S, Dopazo J, Meyer TF,
Conesa A. 2012. Qualimap: Evaluating next-generation sequencing alignment data.
Bioinformatics **28**:2678–2679. doi:10.1093/bioinformatics/bts503

Gilbert KJ, Pouyet F, Excoffier L, Peischl S. 2020. Transition from background selection to associative
overdominance promotes diversity in regions of low recombination. *Curr Biol* **30**:101–107.
doi:10.1101/748004

Greenspan SE, Lambertini C, Carvalho T, James TY, Toledo LF, Haddad CFB, Becker CG. 2018. Hybrids
of amphibian chytrid show high virulence in native hosts. *Sci Rep* **8**:9600. doi:10.1038/s41598-
018-27828-w

Grigg ME, Bonnefoy S, Hehl AB, Suzuki Y, Boothroyd JC. 2001. Success and virulence in Toxoplasma as
the result of sexual recombination between two distinct ancestries. *Science (80-)* **294**:161–165.
doi:10.1126/science.1061888

Hahn C, Bachmann L, Chevreux B. 2013. Reconstructing mitochondrial genomes directly from
genomic next-generation sequencing reads - A baiting and iterative mapping approach. *Nucleic
Acids Res* **41**:e129. doi:10.1093/nar/gkt371

Hahn C, Weiss SJ, Stojanovski S, Bachmann L. 2015. Co-speciation of the ectoparasite *Gyrodactylus
teuchis* (*Monogenea, Platyhelminthes*) and its salmonid hosts. *PLoS One* **10**.
doi:10.1371/journal.pone.0127340

Henrich T, Benesh DP, Kalbe M. 2013. Hybridization between two cestode species and its
consequences for intermediate host range. *Parasites and Vectors* **6**:33. doi:10.1186/1756-3305-
6-33

Hotz H, Semlitsch RD, Gutmann E, Guex GD, Beerli P. 1999. Spontaneous heterosis in larval life-
history traits of hemiclonal frog hybrids. *Proc Natl Acad Sci U S A* **96**:2171–2176.
doi:10.1073/pnas.96.5.2171

Houde AE, Torio AJ. 1992. Effect of parasitic infection on male color pattern and female choice in
guppies. *Behav Ecol* **3**:346–351. doi:10.1093/beheco/3.4.346

Jacquin L, Reader SM, Boniface A, Mateluna J, Patalas I, Pérez-Jvostov F, Hendry AP. 2016. Parallel
and nonparallel behavioural evolution in response to parasitism and predation in Trinidadian
guppies. *J Evol Biol* **29**:1406–1422. doi:10.1111/jeb.12880

Jain R, Ghoshal A, Mandal C, Shaha C. 2010. *Leishmania* cell surface prohibitin: Role in host-parasite
interaction. *Cell Microbiol* **12**:432–452. doi:10.1111/j.1462-5822.2009.01406.x

Janko K, Pačes J, Wilkinson-Herbots H, Costa RJ, Roslein J, Drozd P, Iakovenko N, Rídl J, Hroudová M,
Kočí J, Reifová R, Šlechtová V, Choleva L. 2018. Hybrid asexuality as a primary postzygotic
barrier between nascent species: On the interconnection between asexuality, hybridization and
speciation. *Mol Ecol* **27**:248–263. doi:10.1111/mec.14377

Johnson MB, Lafferty KD, van Oosterhout C, Cable J. 2011. Parasite transmission in social interacting
hosts: Monogenean epidemics in guppies. *PLoS One* **6**. doi:10.1371/journal.pone.0022634

Kim BY, Huber CD, Lohmueller KE. 2018. Deleterious variation shapes the genomic landscape of
introgression. *PLoS Genet* **14**:1–30. doi:10.1371/journal.pgen.1007741

Kimura M, Maruyama T, Crow JF. 1963. The mutation load in small populations. *Genetics*.

Kincaid-Smith J, Tracey A, Augusto R de C, Bulla I, Holroyd N, Rognon A, Rey O, Chaparro C, Oleaga A,
Mas-Coma S, Allienne J-F, Grunau C, Berriman M, Boissier J, Toulza E. 2019. Morphological and
genomic characterisation of the hybrid schistosome infecting humans in Europe reveals a

complex admixture between *Schistosoma haematobium* and *Schistosoma bovis* parasites.
bioRxiv. doi:10.1101/387969

King KC, Stelkens RB, Webster JP, Smith DF, Brockhurst MA. 2015. Hybridization in Parasites:
Consequences for Adaptive Evolution, Pathogenesis, and Public Health in a Changing World.
PLOS Pathog **11**. doi:10.1371/journal.ppat.1005098

King TA, Cable J. 2007. Experimental infections of the monogenean *Gyrodactylus turnbulli* indicate
that it is not a strict specialist. *Int J Parasitol* **37**:663–672. doi:10.1016/j.ijpara.2006.11.015

Kingsbury N. 2009. Hybrid: the history and science of plant breeding. University of Chicago Press.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular evolutionary genetics analysis
across computing platforms. *Mol Biol Evol* **35**:1547–1549. doi:10.1093/molbev/msy096

Kuusela J, Ziętara MS, Lumme J. 2007. Hybrid origin of Baltic salmon-specific parasite *Gyrodactylus*
salaris: A model for speciation by host switch for hemiclonal organisms. *Mol Ecol* **16**:5234–
5245. doi:10.1111/j.1365-294X.2007.03562.x

Lafferty KD. 2009. The ecology of climate change and infectious diseases. *Ecology* **90**:888–900.
doi:10.1890/08-0079.1

Laine A-L, Barrès B, Numminen E, Siren JP. 2019. Variable opportunities for outcrossing result in
hotspots of novel genetic variation in a pathogen metapopulation. *Elife* **8**:e47091.
doi:10.7554/elife.47091

Lambeck K. 2004. Sea-level change through the last glacial cycle: geophysical, glaciological and
palaeogeographic consequences. *Comptes Rendus Geosci* **336**:677–689.
doi:10.1016/j.crte.2003.12.017

Lippman ZB, Zamir D. 2007. Heterosis: revisiting the magic. *Trends Genet* **23**:60–66.
doi:10.1016/j.tig.2006.12.006

Lynch M, Butcher D, Bürger R, Gabriel W. 1993. The mutational meltdown in asexual populations. *J*
Hered **84**:339–344. doi:10.1093/oxfordjournals.jhered.a111354

Lynch M, Gabriel W. 1990. Mutation Load and the Survival of Small Populations. *Evolution (N Y)*
44:1725–1737. doi:10.2307/2409502

Maceda-Veiga A, Cable J. 2019. Diseased fish in the freshwater trade: From retailers to private
aquarists. *Dis Aquat Organ* **132**:157–162. doi:10.3354/dao03310

Meinilä M, Kuusela J, Ziętara MS, Lumme J. 2004. Initial steps of speciation by geographic isolation
and host switch in salmonid pathogen *Gyrodactylus salaris* (*Monogenea: Gyrodactylidae*). *Int J*
Parasitol **34**:515–526. doi:10.1016/j.ijpara.2003.12.002

Moritz C, Brown WM, Densmore LD, Wright JW, Vyas D, Donnellan S, Adams M, Baverstock P. 1989.
Genetic diversity and the dynamics of hybrid parthenogenesis in *Cnemidophorus* (*Teiidae*) and
Heteronotia (*Gekkonidae*). *Evol Ecol unisexual Vertebr* **466**:87–112.

Neaves WB, Baumann P. 2011. Unisexual reproduction among vertebrates. *Trends Genet* **27**:81–88.
doi:10.1016/j.tig.2010.12.002

Neiman M, Sharbel TF, Schwander T. 2014. Genetic causes of transitions from sexual reproduction to
asexuality in plants and animals. *J Evol Biol* **27**:1346–1359. doi:10.1111/jeb.12357

Oziolor EM, Reid NM, Yair S, Lee KM, VerPloeg SG, Bruns PC, Shaw JR, Whitehead A, Matson CW.
2019. Adaptive introgression enables evolutionary rescue from extreme environmental
pollution. *Science (80-)* **364**:455–457. doi:10.1126/science.aav4155

Patz JA, Graczyk TK, Geller N, Vittor AY. 2000. Effects of environmental change on emerging parasitic
diseases. *Int J Parasitol* **30**:1395–1405. doi:10.1016/S0020-7519(00)00141-7

Phillips KP, Cable J, Mohammed RS, Herdegen-Radwan M, Raubic J, Przesmycka KJ, van Oosterhout C,
Radwan J. 2018. Immunogenetic novelty confers a selective advantage in host–pathogen
coevolution. *Proc Natl Acad Sci U S A* **115**:1552–1557. doi:10.1073/pnas.1708597115

Platt RN, McDew-White M, Le Clec'H W, Chevalier FD, Allan F, Emery AM, Garba A, Hamidou AA, Ame
SM, Webster JP, Rollinson D, Webster BL, Anderson TJC, Shapiro B. 2019. Ancient Hybridization
and Adaptive Introgression of an Invadolysin Gene in Schistosome Parasites. *Mol Biol Evol*
36:2127–2142. doi:10.1093/molbev/msz154

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker
PIW, Daly MJ, Sham PC. 2007. PLINK: a tool set for whole-genome association and population-
based linkage analyses. *Am J Hum Genet* **81**:559–575. doi:10.1086/519795

Reynolds M, Arapi EA, Cable J. 2018. Parasite-mediated host behavioural modifications: *Gyrodactylus*
turnbulli infected Trinidadian guppies increase contact rates with uninfected conspecifics.
Parasitology **145**:920–926. doi:10.1017/S0031182017001950

Rieseberg LH, Raymond O, Rosenthal DM, Lai Z, Livingstone K, Nakazato T, Durphy JL, Schwarzbach
AE, Donovan LA, Lexer C. 2003. Major ecological transitions in wild sunflowers facilitated by
hybridization. *Science (80-)* **301**:1211–1216. doi:10.1126/science.1086949

Rogers MB, Downing T, Smith BA, Imamura H, Sanders M, Svobodova M, Volf P, Berriman M, Cotton
JA, Smith DF. 2014. Genomic Confirmation of Hybridisation and Recent Inbreeding in a Vector-
Isolated *Leishmania* Population. *PLoS Genet* **10**. doi:10.1371/journal.pgen.1004092

Schelkle B, Faria PJ, Johnson MB, van Oosterhout C, Cable J. 2012. Mixed infections and hybridisation
in monogenean parasites. *PLoS One* **7**. doi:10.1371/journal.pone.0039506

Schön I, van Dijk P, Martens K. 2009. Lost sex: The evolutionary biology of parthenogenesis, Lost Sex:
The Evolutionary Biology of Parthenogenesis. Springer. doi:10.1007/978-90-481-2770-2

Schosserer M, Minois N, Angerer TB, Amring M, Dellago H, Harreither E, Calle-Perez A, Pircher A,
Gerstl MP, Pfeifenberger S, Brandl C, Sonntagbauer M, Kriegner A, Linder A, Weinhäusel A,
Mohr T, Steiger M, Mattanovich D, Rinnerthaler M, Karl T, Sharma S, Entian KD, Kos M,
Breitenbach M, Wilson IBH, Polacek N, Grillari-Voglauer R, Breitenbach-Koller L, Grillari J. 2015.
Methylation of ribosomal RNA by NSUN5 is a conserved mechanism modulating organismal
lifespan. *Nat Commun* **6**:1–17. doi:10.1038/ncomms7158

Stephenson JF, van Oosterhout C, Mohammed RS, Cable J. 2015. Parasites of Trinidadian guppies:
Evidence for sex- and age-specific trait-mediated indirect effects of predators. *Ecology* **96**:489–
498. doi:10.1890/14-0495.1

Stoeckel S, Masson JP. 2014. The exact distributions of FIS under partial asexuality in small finite
populations with mutation. *PLoS One* **9**. doi:10.1371/journal.pone.0085228

Tadiri CP, Scott ME, Fussmann GF. 2018. Microparasite dispersal in metapopulations: A boon or bane
to the host population? *Proc R Soc B Biol Sci* **285**:20181519. doi:10.1098/rspb.2018.1519

Taylor SA, Larson EL. 2019. Insights from genomes into the evolutionary importance and prevalence
of hybridization in nature. *Nat Ecol Evol* **3**:170–177. doi:10.1038/s41559-018-0777-y

Tihon E, Imamura H, Dujardin JC, van Den Abbeele J, van den Broeck F. 2017. Discovery and genomic
analyses of hybridization between divergent lineages of Trypanosoma congolense, causative
agent of Animal African Trypanosomiasis. *Mol Ecol* **26**:6524–6538. doi:10.1111/mec.14271

van Oosterhout C, Harris PD, Cable J. 2003. Marked variation in parasite resistance between two wild
populations of the Trinidadian guppy, *Poecilia reticulata* (*Pisces: Poeciliidae*). *Biol J Linn Soc*
79:645–651. doi:10.1046/j.1095-8312.2003.00203.x

Volf P, Benkova I, Myskova J, Sadlova J, Campino L, Ravel C. 2007. Increased transmission potential of
Leishmania major/Leishmania infantum hybrids. *Int J Parasitol* **37**:589–593.
doi:10.1016/j.ijpara.2007.02.002

Wright CA, Ross C. 1980. Hybrids between *Schistosoma haematobium* and *S. mattheei* and their
identification by isoelectric focusing of enzymes. *Trans R Soc Trop Med Hyg* **74**:326–332.
doi:10.1016/0035-9203(80)90091-7

Xavier R, Faria PJ, Paladini G, van Oosterhout C, Johnson M, Cable J. 2015. Evidence for cryptic
speciation in directly transmitted gyrodactylid parasites of trinidadian guppies. *PLoS One* **10**.
doi:10.1371/journal.pone.0117096

Xu S, Innes DJ, Lynch M, Cristescu ME. 2013. The role of hybridization in the origin and spread of
asexuality in *Daphnia*. *Mol Ecol* **22**:4549–4561. doi:10.1111/mec.12407

Zhao L, Charlesworth B. 2016. Resolving the conflict between associative overdominance and
background selection. *Genetics* **203**:1315–1334. doi:10.1534/genetics.116.188912

659