Expansion of frozen hybrids in the guppy ectoparasite, *Gyrodactylus turnbulli*

Mateusz Konczal[1], Karolina J. Przesmycka[1], Ryan S. Mohammed[2,3], Christoph Hahn[4], Jo

Cable[3], Jacek Radwan[1]

[1] Evolutionary Biology Group, Faculty of Biology, Adam Mickiewicz University, 61-614

Poznań, Poland.

[2] The University of the West Indies Zoology Museum, Department of Life Sciences, Faculty of

Science and Technology, UWI, St. Augustine, Trinidad and Tobago, WI.

[3] School of Biosciences, Cardiff University, Cardiff, CF10 3AX, UK.

[4] Institute of Biology, University of Graz, Austria

* Corresponding author

E-mail: mateusz.konczal@amu.edu.pl

Abstract

Hybridization is one of the major factors contributing to the emergence of highly successful parasites.

Hybrid vigor can play an important role in this process, but subsequent rounds of recombination in

the hybrid population may dilute its effects. Increased fitness of hybrids can, however, be frozen by

asexual reproduction. Here, we identify invasion of a "frozen hybrid" genotype in natural populations

of *Gyrodactylus turnbulli*, a facultatively sexual ectoparasitic flatworm that causes significant damage

to its fish host. We re-sequenced genomes of these parasites infecting guppies from six Trinidad and

Tobago populations, and found surprisingly high discrepancy in genome-wide nucleotide diversity

between islands. The elevated heterozygosity on Tobago is maintained by predominantly clonal

reproduction of hybrids formed from two diverged genomes. Hybridization has been followed by

spread of the hybrids across the island, implying a selective advantage compared to native

genotypes. Our results thus highlight that a single outcrossing event may be independently sufficient

to cause pathogen expansion.

Keywords: hybridization, clonal reproduction, heterosis, invasion, *Gyrodactylus*

Introduction

Hybridization refers to a successful mating between individuals from two populations, which are

distinguishable on the basis of one or more heritable characters and have divergent genomes

(Arnold, 2004). It can be a potent evolutionary force, creating opportunities for adaptive evolution by

increasing genetic variation on which natural selection can act (Arnold, 1997, 2004; Oziolor et al.,

2019; Rieseberg et al., 2003; Taylor & Larson, 2019). The ecological and evolutionary consequences

are potentially large and may include significant effects on the evolutionary trajectories of

pathogenic organisms ('parasites' henceforth) (King, Stelkens, Webster, Smith, & Brockhurst, 2015).

This role is likely to increase in importance because human activities increase opportunities for

hybridization by inducing environmental changes and incidental translocations (Brooks & Hoberg,

2007; Cable et al., 2017; Patz, Graczyk, Geller, & Vittor, 2000). Understanding the role of

hybridization in parasite evolution is therefore essential if we are to predict hot spots of disease risk

and emergence (Laine, Barrès, Numminen, & Siren, 2019).

Parasite strains of hybrid origin can gain an advantage because recombinants – individuals with

shuffled genomes - produced after hybridization may show enhanced phenotypic characteristics

compared to the parental generation, such as higher infectivity (Grigg, Bonnefoy, Hehl, Suzuki, &

Boothroyd, 2001), expanded host range (Detwiler & Criscione, 2010), increased transmission

potential (Volf et al., 2007) or increased tolerance to harsh environmental conditions (Laine et al.,

2019). Hybridization, however, is often associated with asexual reproduction, which may in the long

term preclude the initial benefits resulting from recombination between divergent genomes, since

asexual lineages tend to be less adaptable to environmental change compared to sexually

reproducing lineages (Neaves & Baumann, 2011). Asexual reproduction is thought to be enforced by

Bateson-Dobzhansky-Muller incompatibilities accumulated in parental lineages (Janko et al., 2018;

Neaves & Baumann, 2011). Asexual lineages frequently originate from hybridization events, for

example in cases when the genomes of parental species are divergent enough to disrupt meiosis in

hybrids, yet not diverged enough to seriously affect hybrid viability (Janko et al., 2018; Lamelza et al.,

2019; Moritz et al., 1989). Alternatively, the hybrid origin of asexual animals can arise from

hampered pairings of homologous chromosomes due to decreasing sequence homology (De Storme

& Mason, 2014) or from asynchronous expression of genes brought together by hybridization

(Carman, 1997).

Conversely, asexual reproduction may be selectively beneficial if clonal reproduction "freezes"

heterozygosity that confers short-term evolutionary advantage (Neiman, Sharbel, & Schwander,

2014). Hybrids often show greater vigor, productivity, disease resistance, or overall increased

Darwinian fitness (Kingsbury, 2009). Two alternative hypotheses may explain the existence of hybrid

vigor (heterosis) (Lippman & Zamir, 2007). The overdominance hypothesis, attributes heterosis to

superior fitness of heterozygous genotypes over homozygous wild types. The dominance hypothesis

assumes that recessive deleterious mutations are to some extent masked in a heterozygous stage.

Recent theoretical work indicates that dominance-induced heterosis might play an important role in

evolution, especially if deleterious alleles are not effectively removed from populations as it is

expected when populations are small (Gilbert, Pouyet, Excoffier, & Peischl, 2020; Kim, Huber, &

Lohmueller, 2018; Zhao & Charlesworth, 2016). This mechanism may be particularly relevant to

parasite evolution. Many parasite populations are small or experience frequent bottlenecks,

therefore are prone to the accumulation of deleterious mutations (M. Lynch, Butcher, Bürger, &

Gabriel, 1993), but also to elevated homozygosity due to inbreeding. In such a case, one could expect

higher fitness of hybrid parasites. Successful parasite strains originating from a hybridization event

have been documented in several species, including *Batrachochytrium dendrobatidis* (Farrer et al.,

2011; Greenspan et al., 2018), *Toxoplasma gondii* and *Sarcocystis neurona* (Grigg et al., 2001;

Wendte et al., 2010). However, because in all these cases some degree of recombination occurred,

the relative role of heterozygosity versus recombination between highly divergent genomes remains

unresolved.

Here, we report frozen genome-wide heterozygosity in wild *Gyrodactylus turnbulli*, a common guppy

parasite and well-known model species in epidemiological and eco-evolutionary research (e.g. Houde

& Torio, 1992; Jacquin et al., 2016; Phillips et al., 2018; Reynolds, Arapi, & Cable, 2018; van

Oosterhout, Harris, & Cable, 2003). Previous laboratory work demonstrated that *G. turnbulli* from

mixed strain infections often hybridize and have higher fitness (more parasites per host, longer

infection, but not increased host mortality), suggesting that in fact hybrid individuals might have

increased fitness, possibly due to heterosis or recombination (Schelkle, Faria, Johnson, van

Oosterhout, & Cable, 2012). However, the effect of hybridization on population structure and

dynamics of infections in wild populations has not been investigated to date (Xavier et al., 2015). We

analyze thirty whole genomes of *G. turnbulli* originated from six natural populations from Trinidad

and Tobago. On Tobago we identified a widespread, recently formed "frozen hybrid" genotype,

suggesting the potential importance of heterosis in clonal invasions of *G. turnbulli* and highlight that

a single hybridization event can lead to the emergence of highly successful parasites, despite

suppressing recombination.

Materials and Methods

Field collections

During two expeditions (2017 and 2018), guppies (*Poecilia reticulata*) were sampled from six

locations on Trinidad and Tobago (Trinidad: Aripo River [ARI], Caura River [CAU], Lopinot River [LOP];

Tobago: Dog River [DR], Health Center [HC], Spring Site [SPS]). Fish were transported to our field

station in Charlotteville, northeast Tobago, where each population was kept in a separate aquarium.

Over four weeks, guppies were subsequently screened for ectoparasites under a dissecting

microscope. If gyrodactylids were observed, a single worm was allowed to move to a naïve,

anaesthetized fish, obtained from *Gyrodactylus*-free mesocosms established by Phillips et al. (2018).

The transfer was closely monitored under the microscope to ensure movement of just a single worm.

Fish experimentally infected in this way were kept in separate tanks in order to establish strains from

which we could obtain enough DNA required to prepare libraries for sequencing. Fish were screened

every 2-3 days to verify the presence of gyrodactylids, and hosts were euthanized with an overdose

of MS-222 8-13 days post-infection. The number of worms was counted, samples were preserved in

97% molecular grade ethanol and transported to Poland. Life expectancy for *G. turnbulli* was

estimated to 4.2 days (Scott, 1982). The first offspring is born approx. 1 day after the birth of the

parent, and subsequent offspring are born at 2-2.5 days intervals (Scott, 1982). We thus expect that

most individuals develop from the first or second births. The first-born always develops clonally,

while the second-born daughter originates from parthenogenesis (Bakke, Cable, & Harris, 2007;

Cable & Harris, 2002). Even though it is unclear which mechanism restores the diploid component

during parthenogenesis (Cable & Harris, 2002), it has been suggested that heterozygosity is

maintained during that process, as reported for *G. salaris.* Third- and subsequent born daughters

may result from outcrossing, selfing, or parthenogenesis. Given our experimental design we expect

that the majority of gyrodactylid individuals on each experimentally infected fish developed from

first or second births and should have preserved founder's genomes. Samples with at least 10 worms

were used for DNA extraction and for species identification as described below. Parasites collected in

2017 and identified as *G. bullatarudis* were reported in a separate study (Konczal et al., 2020), while

this work focused solely on *G. turnbulli.*

DNA extraction, libraries preparation and sequencing

DNA was extracted from pools of individuals ('strain' henceforth) derived from a single worm

collected from a single experimentally infected fish using MagJET Genomic DNA kits (Thermo

Scientific). DNA concentration was measured using Qubit High Sensitivity reagents. *Gyrodactylus*

were identified to species using Sanger sequencing of a 262 bp fragment of the mitochondrial protein

coding COII gene (Xavier et al., 2015). Sequences were aligned together with records downloaded

from the NCBI Genbank, and a Neighbor Joining tree was reconstructed with MEGA X (Kumar,

Stecher, Li, Knyaz, & Tamura, 2018) to determine species. Thirty strains were selected for library

preparation (Nextera DNA Flex Library Prep Kit, Illumina) based on DNA quantity and population of

origin. Subsets of strains were sequenced in 2018 with HiSeq2500 (2 x 100 bp). All other strains were

sequenced in 2019 on the NovaSeq6000 platform in 2 x 150 bp mode (see Supplementary Table 1 for

more details).

SNP calling and population genomic analyses

The quality of raw reads was assessed with Fastqc (Andrews & Babraham Bioinformatics, 2010).

Reads were then filtered with trimmomatic (version 0.36; Bolger, Lohse, & Usadel, 2014) and

mapped using bwa mem (version 0.7.10-r789) to the reference nuclear genome (Hahn et al., in

prep.), which was assembled from individuals from the *Gt3* laboratory strain kept at Cardiff

University. Mapping results was visualized with Qualimap (García-Alcalde et al., 2012) and duplicated

reads were marked with picardtools (version 2.18.5-6). SNPs were then called with samtools mpileup

(version 1.6.0, options -R -C50 -t DP, ADF, ADR). Based on inspection of the distributions of diagnostic

statistics, we removed SNPs with DP > 1200 or DP < 400 (for all samples together), with QUAL < 30,

with MQ < 30 or in 5 bp distance from identified indels.

We then used SNPs to calculate PCA with plink (version 1.90, Purcell et al., 2007) using only SNPs

with minor allele frequency greater than 0.05. The same dataset with additional sample representing

reference genotype (all genotypes encoded as reference homozygous) was used to calculate distance

matrix (in one minus identity-by-state units) with plink and to construct a Neighbor Joining tree using

the ape package in R. In addition, we ran PCA analyses with SNPs, that were previously pruned for

Linkage Disequilibrium. We first pruned SNPs with –indep-pairwise 100 10 0.1 option in plink, and

then used the same program to calculate PCA (D. H. Alexander, Novembre, & Lange, 2009).

Vcftools (version v0.1.12b; Danecek et al., 2011) was used to calculate nucleotide diversity (π) and

genetic divergence between populations (F_{ST}) in 25 kb non-overlapping windows, as well as

inbreeding coefficient (F_{IS}, option –het) per strain. To calculate F_{IS} per strain we considered only SNPs

polymorphic within population of origin. The fraction of heterozygous sites per genomic window was

calculated using a custom script and our estimates of genome wide heterozygosity per strain were

inferred as the average of these fractions across all genomic windows. For each strain we also

calculated divergence form the reference genome in 25 kb non-overlapping windows, by counting

the number of non-reference variants (alternative homozygous sites were counted as one and

reference/alternative allele heterozygous sites were counted as half) and dividing it by a window

length. Windows shorter than 12.5 kb (ends of scaffolds and scaffolds shorter than that length) were

excluded from all analyses.

Mitochondrial genome assembly and mtDNA analyses

To reconstruct phylogenetic relationships between strains we reconstructed the entire mitochondrial

haplotype for strain G126 using MITObim (*de novo* approach of Hahn, Bachmann, & Chevreux, 2013)

with COII fragment downloaded from NCBI (accession KP168411.1) as an initial bait. We then

mapped reads from all sequenced strains to that sequence using bwa mem with default parameters.

Results were inspected manually using IGV. Duplicates were marked with picardtools and SNPs were

called using mpileup and bcftools assuming haploid samples. SNPs with quality lower than phred 30

were discarded from analyses. The nucleotide sequence for each strain was reconstructed with

bcftools consensus. Sequences were then aligned and used for phylogenetic analyses.

The evolutionary relationship between strains was inferred using ML with a GTR model, implemented

in MEGA X Software (Kumar et al., 2018). The same software and model were used to calculate mean

evolutionary distances between groups. BEAST2 (v2.6.1; Bouckaert et al., 2014) was then used to

perform Bayesian phylogenetic analysis of mitochondrial genomes and to estimate time to most

recent common ancestor (TMRCA). TMRCAs of nodes of interest were estimated based on

10,000,000 generations, logging every 1,000 steps, with the first 20% generations discarded as a

burn-in. We used a Yule Model as prior, HKY substitution model, and strict clock model. Divergence

time estimates were based on 13.1% divergence per million years estimated for *G. salaris* (see

Kuusela et al., 2007).

Results

A total of thirty *G. turnbulli* strains from three Trinidadian (Aripo, Caura and Lopinot Rivers) and three

Tobagonian populations (Dog River, Health Center and Spring Site; Figure 1A, details in

Supplementary Table 1) collected in two consecutive years were sequenced to an average coverage

of 38.7 per strain. Sequencing reads were then mapped to the reference nuclear genome and used

for SNP calling. We identified 776,881 single nucleotide- and 161,255 indel-polymorphisms. Principal

Component Analyses of SNP genotypes clearly separated Trinidad from Tobago, while differentiation

between local populations was present only in some cases (Figure 1B). In particular, strains from

Health Center and Dog River (Tobago) showed virtually no differentiation and, in contrast to the

Trinidadian populations, do not form monophyletic groups according to the populations of origin

(Figure 1C). The Trinidadian strains were more similar to the reference genome, obtained from a

strain on commercial guppies, *Gt3,* than strains collected from Tobago (Figure 1C). To remove the

potential effect of linkage on population differentiation, we pruned linked SNPs, but the general

pattern remained the same (Supplementary Figure 1). We calculated F_{ST} between Trinidad and

Tobago populations in 25 kb non-overlapping windows and found that 90% of the windows show F_{ST}

values between 0.25 and 0.55 (mean F_{ST} = 0.42; Supplementary Figure 2). Pairwise F_{ST} values were

higher among Trinidadian populations (Supplementary Figures 3) than among populations from

Tobago (Supplementary Figures 4). To resolve whether moderate F_{ST} between islands resulted from

moderate diversification between or from elevated levels of genetic variation within populations, we

calculated nucleotide diversity across Trinidad and Tobago strains separately. Nucleotide diversity on

Tobago was 2.4 times greater than that on Trinidad (0.0028 and 0.0012 respectively, Figure 1D). The

difference does not result from high differentiation among Tobagonian populations: nucleotide

diversity was also higher within Tobagonian populations, compared with those from Trinidad

(Supplementary Figure 5).

Increased nucleotide diversity on Tobago was mainly driven by high heterozygosity of most

individuals (Figures 1D-E). Similarly, inbreeding coefficient, F_{IS}, was negative for most of the strains

from Tobago, suggesting over-representation of heterozygous positions (Supplementary Figure 6). In

contrast, most of the Trinidadian strains showed positive values of F_{IS}.

High heterozygosity was observed in all strains from the Tobago populations Health Center and Dog

River. The third Tobago population (Spring Site) showed a mixture of patterns. One Spring Site strain

(G334) had low level of heterozygosity, reminiscent of the general pattern observed in Trinidadian

strains. Three other strains (G328, G367 and G368) showed bimodal distribution, with genomic

regions presenting either high or low level of heterozygosity. Finally, two Spring Site strains (G331

and G370) contained only highly heterozygous regions, similar to the pattern observed in Health

Center and Dog River (Figure 1E).

The majority of positions found polymorphic among Tobagonian strains (325,473 out of 498,111)

were heterozygous in all Health Center and Dog River strains, as well as and two (G331 and G370)

Spring Site strains. A total of 88% of these SNPs combined one allele from the reference genome and

the alternative allele found in the Tobagonian G334 strain (Figure 2). Genome-wide pattern of

divergence between re-sequenced strains and the reference genome demonstrated that strain G334

was the most divergent. Strains from Trinidad were consistently more closely related to the

reference genome (Figure 1C, Supplementary Figure 7). Based on these patterns we infer that highly

heterozygous Tobagonian strains consist of hybrids between two divergent haplotypes: one similar

to the reference sequence and the second similar to the G334 sequence. The same pattern with

shared heterozygous sites and high, genome-wide heterozygosity was observed in the Health Center

and Dog River strains from 2017 and 2018, indicating that the hybrid population is stable. This

implies an absence of recombination, as recombination would have fixed one of the two haplotypes

in at least some genomic regions. Auxiliary analysis comparing heterozygosity across all contigs to

that of a simulated, maximally heterozygous hybrid genome, created *in silico* from the reference

sequence and the G334 strain, confirmed that no such homozygous regions are observed in these

populations (Supplementary Figure 8). Furthermore, we compared genome wide nucleotide diversity

of Tobagonian strains collected in 2017 and 2018 and did not find evidence for a substantial

difference between time points (π_{2017} = 0.0018 vs π_{2018} = 0.0017). We also plotted relationships

between heterozygosity calculated in non-overlapping windows for each pair of strains and did not

find any indication for a reduction in heterozygosity in any strain (Supplementary Figure 9),

suggesting no ongoing recombination.

In contrast, divergent haplotypes have clearly recombined in the Spring Site population.

Three strains (G328, G367 and G368) show a mosaic pattern of the two haplotypes fixed in different

genomic locations, with other parts of the genome still remaining heterozygous (Figure 2,

Supplementary Figures 8 and 10). Signatures of sexual reproduction are also observed at least in

some of the strains from Trinidad. For example, SNPs localized in the scaffold scf7180000443747

(first sequence in the reference genome) show admixture of another haplotype in strain G314

coming from Lopinot, and another two strains in Lopinot are heterozygous in this region (Figure 2).

Another strain from the same population (G308) has negative F_{IS} (Supplementary Figure 6) and

slightly elevated genome-wide heterozygosity (Figure 1D), although the effect is weaker compared to

strains from Tobago. Nevertheless, heterozygous sites are on average rare in Trinidadian strains.

Clustering analyses (Figures 1B, 1C) and F_{ST} results (Supplementary Figure 4) based on nuclear SNPs

suggest low divergence between samples identified as hybrids. We explored it further by calculating

divergence in the haploid mitochondrial genomes. One Trinidadian strain (G126) was used for

mitochondrial genome assembly, and then reads from all other strains were mapped to that genome

for haplotype reconstruction. The resulting mt genome spans 14,159 bp. We found very little

divergence between strains identified as hybrids, compared to divergence amongst other samples

(Figure 3, Supplementary Figure 11). Putative hybrid strains from Health Center, Dog River and Spring

Site populations were found more closely related to Trinidadian strains than to remaining SPS strains,

confirming introgression inferred from genomic data. Within hybrids (Health Center, Dog River two

hybrid strains from Spring Site) the mean divergence between populations was 0.02%. In contrast,

the three other strains collected from Spring Site had strongly divergent mitochondria - the

nucleotide divergence between the two Spring Site groups was 1.58%. To independently verify low

divergence between Tobagonian hybrids and strains from Trinidad we sequenced a 262 bp fragment

of COII from 81 samples collected by us on Trinidad and Tobago. The results confirmed little

divergence between all Tobagonian samples except for some derived from the Spring Site population

(Supplementary Figure 12).

Lack of recombination in Tobago hampers identification of fast evolving regions in the genome,

because of strong linkage between sites. However, there was no evidence that the Trinidadian

populations experienced clonal expansion, and therefore rapidly evolving genes can be potentially

identified. For this reason, we tested for correlations between nucleotide diversity calculated in

windows separately for Tobagonian hybrid populations and for all Trinidadian strains. We used

nucleotide diversity calculated jointly for Health Center and Dog River strains as a proxy for the long-

term evolution of genomic regions (divergence between two haplotypes forming hybrids). Genomic

regions in which the nucleotide diversity calculated jointly for all strains from Trinidad deviates from

predictions based on this long-term evolutionary rate are therefore candidates for recent rapid

evolutionary change between years or populations. The divergence calculated in 25 kb windows

were highly correlated between Tobagonian hybrid populations and Trinidadian populations, while

few windows demonstrated elevated (according to overall trend) diversity in Trinidad (Figure 4A).

One scaffold (scf7180000445747; 19.4 kb long) showed extensive divergence between Trinidadian

strains collected in 2017 and 2018 (Figure 4B). It contains 7 protein coding genes, including putative

methylotransferase NSUN5, Endonuclease/exonuclease/phosphatase family domain-containing

protein 1, synaptogyrin and prohibitin.

Discussion

We examined whole genomes of thirty *G. turnbulli* strains, sampled from two Caribbean islands in

two consecutive years, providing insight into the evolution of their genomic diversity. On Tobago,

parasite populations tend to have much higher nucleotide diversity than those on Trinidad. This

higher diversity was caused by the presence of highly heterozygous strains in all studied Tobagonian

populations (50-100% of individuals per population). These strains shared most of the heterozygous

loci, with a repeatable pattern of having one reference-like allele and the other allele present in a

homozygous state only in one strain from Tobago (G334). This pattern led us to conclude that these

highly heterozygous strains comprise hybrids of two divergent lineages, one of which is close to a

strain isolated from ornamental guppies (Trinidad-like, used for the reference genome assembly),

and the second which is close to the Tobagonian G334 strain. In the Spring Site population (Tobago),

we observed two distinct mitochondrial lineages (diverged 145,000 years ago based on 13.1%

divergence per Mya, (Kuusela et al., 2007), 95% HPD: 133,900 – 155,700). Most likely they represent

mitochondrial genomes originating from two parental populations that formed Tobagonian hybrids.

The mitochondrial divergence between these two lineages was much larger than between

Tobagonian hybrids and Trinidadian populations (7,600 years, 95% HPD: 5,200 – 9,800 years),

demonstrating that hybrids inherited mitochondria from Trinidad-like ancestors. Initial hybridization

must have been followed by reproduction without recombination, otherwise recombination would

have produced both homozygous and heterozygous genotypes at least in some of the genomic

regions (as observed in two strains at Spring Site). In contrast, in most Tobagonian strains

heterozygosity was elevated genome-wide (Figure 1D), heterozygous genotypes were very similar for

all samples (Figure 2), suggesting that these genomes represent "frozen hybrids". Indeed, our

analyses imply that no recombination occurred since the hybrids were formed. If recombination

happened before expansion of the hybrid genotype, some regions with loss of heterozygosity would

be found and shared for all HC and DR strains. We did not find any evidence for loss of heterozygosity

in HC nor DR (Supplementary Figure 8). In contrast to HC and DR, patterns suggesting recombination

were obvious in some strains from the Spring Site population. Strains G328, G367 and G368 showed

non-identical patterns of heterozygosity across the nuclear genome, consistent with recent

independent outcrossing/selfing events followed by recombination. Two of these three Spring Site

strains (G328, G368) have a mitochondrial genome significantly divergent from other hybrids,

suggesting that subsequent rounds of outcrossing occurred between hybrid and non-hybrid

individuals, the latter carrying the divergent mtDNA. On Trinidad, a single individual showed slightly

elevated heterozygosity, possibly as a result of sexual reproduction between two moderately

diverged strains (G308, Figure 1D, Supplementary Figure 6), and indeed another individual from the

same population showed a pattern consistent with being a subsequent generation recombinant

(G312). Thus, in contrast to a set of pure hybrid populations we found on Tobago (Dog River and

Health Centre), recombination is not completely absent in other populations both on Tobago (Spring

Site) and on Trinidad. This evidence is consistent with the data from laboratory cross infections with

G. turnbulli, which demonstrated that 4-11% of individuals in mixed infections are hybrids with

ancestors from different strains used in the experiment (Schelkle et al., 2012).

Not only sexual reproduction, but also most forms of parthenogenesis involve meiotic

recombination, which is expected to lead to loss of heterozygosity spanning large genomic regions.

However, automictic parthenogenesis (automixis) with central fusion can freeze heterozygosity in

spite of recombination.(Engelstädter, 2017; Jaron, Bast, Ranallo-Benavidez, Robinson-Rechavi, &

Schwander, 2018) Under automixis with central fusion the probability of recombination increases

with distance from the centromere, and particularly regions distal to the centromere are affected if

recombination does occur. While, we are not able identify centromeres/telomeres in the reference

genome of *G. turnbulli,* we did not find any evidence for large genomic regions with loss of

heterozygosity (Supplementary Figures 11 and 12). Thus, either Tobagonian hybrids (i) reproduce

only mitotically, (ii) recombination is strongly suppressed during automictic parthenogenesis, or (iii)

recombinants are removed by selection leading to clonal-like evolution. Our data suggest that the

inferred, clonally-like reproducing hybrids spread relatively recently across Tobago, as indicated by

high similarity of their mitochondrial genomes, corresponding to 2,700 (95% HPD: 1,500 − 4,000)

years of divergence (Figure 3). The divergence time is based on the indirectly derived substitution

rate in a single mitochondrial locus of *Gyrodactylus salaris* (13.1% per Mya; Kuusela et al., 2007).

Other indirect estimates suggest lower (5.1% per Mya; Hahn et al., 2015) or higher substitution rates

(up to 20.3% per Mya; Meinilä et al., 2004) in *Gyrodactylus* species, and therefore divergence dating

should be taken with caution. Nevertheless, low nucleotide divergence between mitochondrial

genomes indicates a recent origin of the hybrids, and their widespread occurrence across Tobago

provides evidence for their evolutionary success. Hybrid genomes that were most widespread on

Tobago did not show any signs of recombination, and apparently remained stable, as would be

expected under predominantly clonal evolution.

What evolutionary forces are responsible for the apparent success of clonal hybrids on Tobago?

Tibayrenc and Ayala (2012, 2017) proposed that a predominance of clonal evolution is a common

adaptation to parasitism in micropathogens. Their hypothesis considers the total set of reproductive

strategies used by pathogens to escape 'recombination load'. In contrast, the 'epidemic clonality

model' (Smith, Smith, O'Rourke, & Spratt, 1993) advocates for the occurrence of occasional bouts of

clonally reproduced strains in otherwise recombining species. Ephemeral clones are frequently

replaced by other genotype combinations, maintaining a dominant role of sexual reproduction in

evolution. Complex patterns of outcrossing events in *G. turnbulli* populations seem to be consistent

with an 'epidemic clonality model' of evolution.

In the case of *G. turnbulli* a bout of clonality is associated with hybridization between two, relatively

divergent strains, confirming that hybridization of parasites can lead to the emergence of highly

successful, and potentially virulent strains. The potential to increase the virulence of *G. turnbulli* by

combining different genomes was suggested in an earlier report of laboratory experiments, in which

infection with two inbred stains led to higher parasite load compared to single strain infections

(Schelkle et al., 2012). However, the genomic divergence between these strains was unknown and, as

discussed by the authors, this result could be due to competition between clones leading to the

evolution of increased virulence (Schelkle et al., 2012).

Human activity, migrations and climate change shift the geographic distribution of many species

(Brooks & Hoberg, 2007; Cable et al., 2017; Lafferty, 2009), which increases the probability of

secondary contact between former distinct populations, potentially leading to hybridization followed

by adaptive evolution. Parasites are often transported with their hosts, and even if hosts do not

contribute to genetic pools of local populations, hitchhiking parasites can do so. This could be the

case for *G. turnbulli.* Guppies are commonly kept in aquaria, and *G. turnbulli* are regularly reported in

aquarium populations (Maceda-Veiga & Cable, 2019) from where they can potentially be released to

nature. Alternatively, hybrids might have appeared naturally. Our approximate dating of divergence

of mtDNA between hybrids from Trinidad and Tobago is consistent with the end of the last glacial

maximum, when a land bridge that probably existed between the islands during glaciation-related

sea level decline was disrupted again (H. J. Alexander, Taylor, Sze-Tsun Wu, & Breden, 2006;

Lambeck, 2004). This might have facilitated migration of guppies between the islands, and possibly

other fish species capable of spreading *G. turnbulli* (Cable et al., 2013; T. A. King & Cable, 2007).

Our recent study on the natural populations of another monogenean parasite infecting guppies

(*Gyrodactylus bullatarudis*) suggested that recent hybridization has also played a predominant role in

shaping genetic variation, but in that case in populations from Trinidad (Konczal et al., 2020).

Hybridization between two diverged lineages and its subsequent recombination resulted in a mosaic

genome composition of *G. bullatarudis*. Around half of the genome originates from one or another

lineage. High divergence between recombined lineages and relatively little divergence between local

populations led us to conclude that recent hybridization was followed by rapid expansion of a

recombinant *G. bulllatarudis* strain. Similar patterns of successful admixed hybrid genotypes have

been reported for many other parasites, including schistosomes (Kincaid-Smith et al., 2019; Platt et

al., 2019), *Trypanosoma* (see Tihon et al., 2017) and *Leishmania* (see Rogers et al., 2014) species, as

well as Dutch elm disease pathogens (Brasier, 2001; Hessenauer et al., 2020) and other fungi

pathogens (Mixão & Gabaldón, 2018) highlighting the role of recombination between divergent

genomes in parasite evolution. Our analysis of *G. turnbulli* genomes indicates, that unlike its

congener *G. bullatarudis,* natural expansion occurred without prior recombination. This scenario is

similar to suggestions for *Gyrodactylus salaris* – a significant pathogen of Atlantic salmon. The Baltic-

salmon specific *G. salaris* was proposed to have originated from a single hybridization event between

two clades, followed by the clonal reproduction of 'frozen hybrids', however this suggestion was

based on analyses of a single anonymous nuclear marker (Kuusela et al., 2007). Therefore, not only

clonal expansion of hybrids, but also linkage to lethal variants for both alleles (balanced lethal

system), or epistatic interactions could potentially explain the pattern observed in *G. salaris.* Our

genome-wide data suggest that clonal expansion of 'frozen hybrids' can indeed take place amongst

gyrodactylids.

Similar observations in other pathogens also suggest that 'frozen hybrids' might have a significant

evolutionary advantage. An interesting example comes from a common fungal parasite of

amphibians, *Batrachochytrium dendrobatidis,* in which asexually reproducing hybrids have higher

virulence than both parental strains in some host species (Farrer et al., 2011; Greenspan et al., 2018).

That hybrid is highly heterozygous and widespread. Most likely contact between previously

genetically isolated allopatric populations of the fungus facilitated hybridization, resulting in

generation, spread and invasion of the hypervirulent strain (Farrer et al., 2011). However, the strain

is not completely clonal and diversification can proceed by either mitotic or sexual recombination.

Similarly, in *Toxoplasma gondii* and *Sarcocystis neurona* outcrossing can precede a disease outbreak

(Grigg et al., 2001; Wendte et al., 2010), but subsequent selfing events were responsible for epidemic

expansions, and, as pointed out by the authors, it is not clear whether an out-cross is independently

sufficient to cause an epidemic (Wendte et al., 2010). Our data are therefore unique, in

demonstrating parasite expansion despite complete genome-wide suppression of recombination,

implying that 'frozen hybrids' might gain advantage from masking recessive, deleterious mutations or

from widening ecological niche.

Parasites may be particularly prone to accumulation of deleterious mutations in their genomes

(Criscione & Blouin, 2005). In general, the rate of accumulation depends primarily on the effective

population size. In small populations, random genetic drift can overpower selection making it easier

for deleterious mutations, particularly those with small effects, to become fixed (Kimura, Maruyama,

& Crow, 1963). This can cause mutational melt-down leading to population extinction (Michael Lynch

& Gabriel, 1990). Several features of macroparasite life cycles can act in concert to reduce effective

population sizes and increase the risk of such melt-down (Criscione & Blouin, 2005). In particular,

frequent bottlenecks associated with colonization/recolonization of new hosts can significantly

reduce the efficiency of selection in removing mildly deleterious mutations from populations.

Additionally, observations of *G. turnbulli* in the wild suggest that census population sizes are often

small and fluctuating over time (Stephenson, van Oosterhout, Mohammed, & Cable, 2015), which is

consistent with the low heterozygosity and generally positive inbreeding coefficient in the non-

hybrid populations that we investigated (Figure 1E, Supplementary Figure 6). If this leads to

homozygosity for a number of recessive or partially recessive mutations, hybridization between

divergent strains carrying those mutations at different genomic locations would result in hybrid vigor.

Alternatively, the advantage of hybridization can be due to alleles carried by heterozygotes being

adapted to different, and a wider range of, environments. In the case of parasites, divergent alleles

may each allow infection of different host species or genomes, allowing heterozygotes to infect a

wider range of hosts. For example, laboratory experiments on tapeworms in sticklebacks showed

that two genetic lines were only able to infect a single respective host species, whereas their hybrids

could infect both species (Henrich, Benesh, & Kalbe, 2013). In the case of *G. turnbulli,* however,

hybridization occurred among two strains infecting the same host species.

Clonal expansion of hybrids is likely facilitated by facultative sexual reproduction of *Gyrodactylus*

species. In such systems, switching between facultative and obligate clonal reproduction does not

appear to be evolutionarily constrained and can easily evolve when favoured by selection (Neiman et

al., 2014). However, though it has been proposed that hybridization is the main mechanism

responsible for evolution of asexual reproduction, its causality and underlying mechanisms are

poorly understood (Janko et al., 2018; Neiman et al., 2014; Schön, van Dijk, & Martens, 2009; Xu,

Innes, Lynch, & Cristescu, 2013). Mortiz et al. (1989) suggested that reproducing asexually results

from the accumulation of incompatibilities between hybridizing species that disrupts cellular

regulation of sexual reproduction. In fact, the majority of species which recently evolved asexual

reproduction are of hybrid origin, although it concerns primarily vertebrates and it is not clear

whether it applies also to other taxa (Neaves & Baumann, 2011). In the case of *G. turnbulli,* however,

recombination is not completely suppressed, because it was observed in one of the hybrid

populations on Tobago (Spring Site), where it might have occurred between hybrid and non-hybrid

individuals. The fact that recombinant individuals are rare on Tobago might result either from rarity

of recombination events, or from effective removal of recombinants from the population by

selection. The disadvantage of recombinants may stem from restoring homozygotes for recessive

deleterious mutations. Discrimination between these two possibilities ideally requires direct

estimates of recombination rate, possibly based on changes in genotype frequencies over time in

both hybrid and non-hybrid populations (Becheler et al., 2017).

Recombination can, at least to some degree, enable independent evolution of different genomic

regions, allowing identification of genomic regions with elevated genomic variation between the two

sampling years, possibly driven by selection. We therefore searched for such regions in Trinidad

populations and found that one genomic region showed an extreme pattern of genetic diversity, with

several interesting genes previously associated with modulating lifespan, stress resistance

(Methylotrasferase NSUN5; Schosserer et al., 2015) and host-parasite interactions (Prohibitin; Jain et

al., 2010). Future work including more individuals and experimental validation is needed to confirm

involvement of these genes in coevolutionary dynamics.

Concluding, our population genomic dataset of *G. turnbulli* allowed us to infer that hybridization

followed by clonal reproduction contributed to evolutionary success of these parasites on Tobago. It

is likely that the evolutionary success stems from heterotic effects, which are frozen in hybrids.

Benefits of such freezing may only be temporary, as the lack of recombination may hinder the

pathogen in keeping up in the evolutionary arms-race with the host's immune system. That

recombination seemed to be maintained in Trinidadian populations suggests that it may be favoured

in the long run as well. It would be interesting to see if after initial success, recombination would

again be favoured, as might have been the case with *G. bullatarudis* (see Konczal et al. 2020). Our

results highlight that a single outcrossing event is independently sufficient to cause pathogen

expansion, thus emphasizing that such processes can result in emergence of invasive pathogens,

some of which may be of public health and conservation concern.

Acknowledgements

We thank K. Phillips for his help in improving this manuscript, N. Cook, A. Szubert-Kruszyńska, A.

Sadowska-Konczal and staff of the Environmental Research Institute Charlotteville (ERIC), Tobago, for

support in the field; P. Turpin of Man O'War Bay Cottages for renting us the field station; Tobago

House of Assembly (THA) for granting permission to conduct field surveys; K. Dudek and J. Raubic for

help in molecular laboratory. All animal work was approved by Cardiff Universities Animal Ethics

Committee and followed UK Home Office Licence (PPL 303424). This work was supported by Polish

National Science Center Fuga Grant UMO-509-2016/20/S/NZ8/00208. Computations were performed

at the Poznan Supercomputing and Networking Center.

References

Alexander, H. J., Taylor, J. S., Sze-Tsun Wu, S., & Breden, F. (2006). Parallel evolution and vicaraince in
the guppy (*Poecilia reticulata*) over multiple spatial and temporal scales. *Evolution*, *60*(11),
2352–2369. doi: 10.1554/06-288.1

Andrews, S., & Babraham Bioinformatics. (2010). FastQC: A quality control tool for high throughput
sequence data. *Manual*. doi: citeulike-article-id:11583827

Arnold, M. L. (1997). *Natural hybridization and evolution*. Oxford University Press.

Arnold, M. L. (2004). Natural hybridization and the evolution of domesticated, pest and disease
organisms. *Molecular Ecology*. doi: 10.1111/j.1365-294X.2004.02145.x

Bakke, T. A., Cable, J., & Harris, P. D. (2007). The Biology of Gyrodactylid Monogeneans: The "Russian-
Doll Killers." *Advances in Parasitology*, *64*, 161–376. doi: 10.1016/S0065-308X(06)64003-7

Becheler, R., Masson, J. P., Arnaud-Haond, S., Halkett, F., Mariette, S., Guillemin, M. L., … Stoeckel, S.
(2017). ClonEstiMate, a Bayesian method for quantifying rates of clonality of populations
genotyped at two-time steps. *Molecular Ecology Resources*, *17*(6), e251–e267. doi:
10.1111/1755-0998.12698

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence
data. *Bioinformatics*, *30*(15), 2114–2120. doi: 10.1093/bioinformatics/btu170

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C. H., Xie, D., … Drummond, A. J. (2014). BEAST
2: A Software Platform for Bayesian Evolutionary Analysis. *PLoS Computational Biology*, *10*(4).
doi: 10.1371/journal.pcbi.1003537

Brasier, C. M. (2001). Rapid Evolution of Introduced Plant Pathogens via Interspecific Hybridization.
BioScience. doi: 10.1641/0006-3568(2001)051[0123:reoipp]2.0.co;2

Brooks, D. R., & Hoberg, E. P. (2007). How will global climate change affect parasite-host
assemblages? *Trends in Parasitology*, *23*(12), 571–574. doi: 10.1016/j.pt.2007.08.016

Cable, J., Archard, G. A., Mohammed, R. S., Mcmullan, M., Stephenson, J. F., Hansen, H., & van
Oosterhout, C. (2013). Can parasites use predators to spread between primary hosts?
Parasitology, *140*(9), 1138–1143. doi: 10.1017/S003118201300067X

Cable, J., Barber, I., Boag, B., Ellison, A. R., Morgan, E. R., Murray, K., … Booth, M. (2017). Global
change, parasite transmission and disease control: Lessons from ecology. *Philosophical
Transactions of the Royal Society B: Biological Sciences*, *372*(1719), 20160088. doi:
10.1098/rstb.2016.0088

Cable, J., & Harris, P. D. (2002). Gyrodactylid developmental biology: Historical review, current status
and future trends. *International Journal for Parasitology*, *32*(3), 255–280. doi: 10.1016/S0020-
7519(01)00330-7

Carman, J. G. (1997). Asynchronous expression of duplicate genes in angiosperms may cause
apomixis, bispory, tetraspory, and polyembryony. *Biological Journal of the Linnean Society*. doi:
10.1111/j.1095-8312.1997.tb01778.x

Criscione, C. D., & Blouin, M. S. (2005). Effective sizes of macroparasite populations: A conceptual
model. *Trends in Parasitology*, *21*(5), 212–217. doi: 10.1016/j.pt.2005.03.002

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin, R. (2011). The
variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158. doi:
10.1093/bioinformatics/btr330

De Storme, N., & Mason, A. (2014). Plant speciation through chromosome instability and ploidy

change: Cellular mechanisms, molecular factors and evolutionary relevance. *Current Plant*
Biology. doi: 10.1016/j.cpb.2014.09.002

Detwiler, J. T., & Criscione, C. D. (2010). An infectious topic in reticulate evolution: Introgression and
hybridization in animal parasites. *Genes*, *1*(1), 102–123. doi: 10.3390/genes1010102

Engelstädter, J. (2017). Asexual but not clonal: Evolutionary processes in automictic populations.
Genetics. doi: 10.1534/genetics.116.196873

Farrer, R. A., Weinert, L. A., Bielby, J., Garner, T. W. J., Balloux, F., Clare, F., … Fisher, M. C. (2011).
Multiple emergences of genetically diverse amphibian-infecting chytrids include a globalized
hypervirulent recombinant lineage. *Proceedings of the National Academy of Sciences*. doi:
10.1073/pnas.1111915108

García-Alcalde, F., Okonechnikov, K., Carbonell, J., Cruz, L. M., Götz, S., Tarazona, S., … Conesa, A.
(2012). Qualimap: Evaluating next-generation sequencing alignment data. *Bioinformatics*,
28(20), 2678–2679. doi: 10.1093/bioinformatics/bts503

Gilbert, K. J., Pouyet, F., Excoffier, L., & Peischl, S. (2020). Transition from background selection to
associative overdominance promotes diversity in regions of low recombination. *Current Biology*,
30(1), 101–107. doi: 10.1101/748004

Greenspan, S. E., Lambertini, C., Carvalho, T., James, T. Y., Toledo, L. F., Haddad, C. F. B., & Becker, C.
G. (2018). Hybrids of amphibian chytrid show high virulence in native hosts. *Scientific Reports*,
8(1), 9600. doi: 10.1038/s41598-018-27828-w

Grigg, M. E., Bonnefoy, S., Hehl, A. B., Suzuki, Y., & Boothroyd, J. C. (2001). Success and virulence in
Toxoplasma as the result of sexual recombination between two distinct ancestries. *Science*,
294(5540), 161–165. doi: 10.1126/science.1061888

Hahn, C., Bachmann, L., & Chevreux, B. (2013). Reconstructing mitochondrial genomes directly from
genomic next-generation sequencing reads - A baiting and iterative mapping approach. *Nucleic*
Acids Research, *41*(13), e129. doi: 10.1093/nar/gkt371

Hahn, C., Weiss, S. J., Stojanovski, S., & Bachmann, L. (2015). Co-speciation of the ectoparasite
Gyrodactylus teuchis (*Monogenea, Platyhelminthes*) and its salmonid hosts. *PLoS ONE*, *10*(6).
doi: 10.1371/journal.pone.0127340

Harris, P. D., Cable, J., & Tinsley, R. C. (1997). Detection of sperm within *Gyrodactylus*
(Platyhelminthes, Monogenea) tissues using fluorescence and transmission electron
microscopy. *Parasitology Research*, *83*(6), 533–538. doi: 10.1007/s004360050294

Henrich, T., Benesh, D. P., & Kalbe, M. (2013). Hybridization between two cestode species and its
consequences for intermediate host range. *Parasites and Vectors*, *6*(1), 33. doi: 10.1186/1756-
3305-6-33

Hessenauer, P., Fijarczyk, A., Martin, H., Prunier, J., Charron, G., Chapuis, J., … Landry, C. R. (2020).
Hybridization and introgression drive genome evolution of Dutch elm disease pathogens.
Nature Ecology and Evolution. doi: 10.1038/s41559-020-1133-6

Houde, A. E., & Torio, A. J. (1992). Effect of parasitic infection on male color pattern and female
choice in guppies. *Behavioral Ecology*, *3*(4), 346–351. doi: 10.1093/beheco/3.4.346

Jacquin, L., Reader, S. M., Boniface, A., Mateluna, J., Patalas, I., Pérez-Jvostov, F., & Hendry, A. P.
(2016). Parallel and nonparallel behavioural evolution in response to parasitism and predation
in Trinidadian guppies. *Journal of Evolutionary Biology*, *29*(7), 1406–1422. doi:
10.1111/jeb.12880

Jain, R., Ghoshal, A., Mandal, C., & Shaha, C. (2010). *Leishmania* cell surface prohibitin: Role in host-
parasite interaction. *Cellular Microbiology*, *12*(4), 432–452. doi: 10.1111/j.1462-
5822.2009.01406.x

Janko, K., Pačes, J., Wilkinson-Herbots, H., Costa, R. J., Roslein, J., Drozd, P., … Choleva, L. (2018).
Hybrid asexuality as a primary postzygotic barrier between nascent species: On the
interconnection between asexuality, hybridization and speciation. *Molecular Ecology*, *27*(1),
248–263. doi: 10.1111/mec.14377

Jaron, K. S., Bast, J., Ranallo-Benavidez, T. R., Robinson-Rechavi, M., & Schwander, T. (2018). Genomic
features of asexual animals. *BioRxiv*, 497495.

Kim, B. Y., Huber, C. D., & Lohmueller, K. E. (2018). Deleterious variation shapes the genomic
landscape of introgression. *PLoS Genetics*, *14*(10), 1–30. doi: 10.1371/journal.pgen.1007741

Kimura, M., Maruyama, T., & Crow, J. F. (1963). The mutation load in small populations. *Genetics*.

Kincaid-Smith, J., Tracey, A., Augusto, R. de C., Bulla, I., Holroyd, N., Rognon, A., … Toulza, E. (2019).
Morphological and genomic characterisation of the hybrid schistosome infecting humans in
Europe reveals a complex admixture between *Schistosoma haematobium* and *Schistosoma*
bovis parasites. *BioRxiv*. doi: 10.1101/387969

King, K. C., Stelkens, R. B., Webster, J. P., Smith, D. F., & Brockhurst, M. A. (2015). Hybridization in
Parasites: Consequences for Adaptive Evolution, Pathogenesis, and Public Health in a Changing
World. *PLOS Pathogens*, *11*(9). doi: 10.1371/journal.ppat.1005098

King, T. A., & Cable, J. (2007). Experimental infections of the monogenean *Gyrodactylus turnbulli*
indicate that it is not a strict specialist. *International Journal for Parasitology*, *37*(6), 663–672.
doi: 10.1016/j.ijpara.2006.11.015

Kingsbury, N. (2009). *Hybrid: the history and science of plant breeding*. University of Chicago Press.

Konczal, M., Przesmycka, K. J., Mohammed, R. S., Phillips, K. P., Camara, F., Chmielewski, S., …
Radwan, J. (2020). Gene duplications, divergence and recombination shape adaptive evolution
of the fish ectoparasite, *Gyrodactylus bullatarudis*. *Molecular Ecology*. doi: 10.1111/mec.15421

Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular evolutionary
genetics analysis across computing platforms. *Molecular Biology and Evolution*, *35*(6), 1547–
1549. doi: 10.1093/molbev/msy096

Kuusela, J., Ziętara, M. S., & Lumme, J. (2007). Hybrid origin of Baltic salmon-specific parasite
Gyrodactylus salaris: A model for speciation by host switch for hemiclonal organisms. *Molecular*
Ecology, *16*(24), 5234–5245. doi: 10.1111/j.1365-294X.2007.03562.x

Lafferty, K. D. (2009). The ecology of climate change and infectious diseases. *Ecology*, *90*(4), 888–900.
doi: 10.1890/08-0079.1

Laine, A.-L., Barrès, B., Numminen, E., & Siren, J. P. (2019). Variable opportunities for outcrossing
result in hotspots of novel genetic variation in a pathogen metapopulation. *ELife*, *8*, e47091.
doi: 10.7554/elife.47091

Lambeck, K. (2004). Sea-level change through the last glacial cycle: geophysical, glaciological and
palaeogeographic consequences. *Comptes Rendus Geoscience*, *336*(7–8), 677–689. doi:
10.1016/j.crte.2003.12.017

Lamelza, P., Young, J. M., Noble, L. M., Caro, L., Isakharov, A., Palanisamy, M., … Ailion, M. (2019).
Hybridization promotes asexual reproduction in *Caenorhabditis* nematodes. *PLoS Genetics*. doi:
10.1371/journal.pgen.1008520

Lippman, Z. B., & Zamir, D. (2007). Heterosis: revisiting the magic. *Trends in Genetics*, *23*(2), 60–66.
doi: 10.1016/j.tig.2006.12.006

Lynch, M., Butcher, D., Bürger, R., & Gabriel, W. (1993). The mutational meltdown in asexual
populations. *Journal of Heredity*, *84*(5), 339–344. doi: 10.1093/oxfordjournals.jhered.a111354

Lynch, Michael, & Gabriel, W. (1990). Mutation Load and the Survival of Small Populations. *Evolution*,
44(7), 1725–1737. doi: 10.2307/2409502

Maceda-Veiga, A., & Cable, J. (2019). Diseased fish in the freshwater trade: From retailers to private
aquarists. *Diseases of Aquatic Organisms*, *132*(2), 157–162. doi: 10.3354/dao03310

Meinilä, M., Kuusela, J., Ziȩtara, M. S., & Lumme, J. (2004). Initial steps of speciation by geographic
isolation and host switch in salmonid pathogen *Gyrodactylus salaris* (Monogenea:
Gyrodactylidae). *International Journal for Parasitology*, *34*(4), 515–526. doi:
10.1016/j.ijpara.2003.12.002

Mixão, V., & Gabaldón, T. (2018). Hybridization and emergence of virulence in opportunistic human
yeast pathogens. *Yeast*. doi: 10.1002/yea.3242

Moritz, C., Brown, W. M., Densmore, L. D., Wright, J. W., Vyas, D., Donnellan, S., … Baverstock, P.
(1989). Genetic diversity and the dynamics of hybrid parthenogenesis in *Cnemidophorus*
(Teiidae) and *Heteronotia* (Gekkonidae). *Evolution and Ecology of Unisexual Vertebrates*, *466*,
87–112.

Neaves, W. B., & Baumann, P. (2011). Unisexual reproduction among vertebrates. *Trends in Genetics*,
27(3), 81–88. doi: 10.1016/j.tig.2010.12.002

Neiman, M., Sharbel, T. F., & Schwander, T. (2014). Genetic causes of transitions from sexual
reproduction to asexuality in plants and animals. *Journal of Evolutionary Biology*, *27*(7), 1346–
1359. doi: 10.1111/jeb.12357

Oziolor, E. M., Reid, N. M., Yair, S., Lee, K. M., VerPloeg, S. G., Bruns, P. C., … Matson, C. W. (2019).
Adaptive introgression enables evolutionary rescue from extreme environmental pollution.
Science, *364*(6439), 455–457. doi: 10.1126/science.aav4155

Patz, J. A., Graczyk, T. K., Geller, N., & Vittor, A. Y. (2000). Effects of environmental change on
emerging parasitic diseases. *International Journal for Parasitology*, *30*(12–13), 1395–1405. doi:
10.1016/S0020-7519(00)00141-7

Phillips, K. P., Cable, J., Mohammed, R. S., Herdegen-Radwan, M., Raubic, J., Przesmycka, K. J., …
Radwan, J. (2018). Immunogenetic novelty confers a selective advantage in host–pathogen
coevolution. *Proceedings of the National Academy of Sciences of the United States of America*,
115(7), 1552–1557. doi: 10.1073/pnas.1708597115

Platt, R. N., McDew-White, M., Le Clec'H, W., Chevalier, F. D., Allan, F., Emery, A. M., … Shapiro, B.
(2019). Ancient Hybridization and Adaptive Introgression of an Invadolysin Gene in *Schistosome*
Parasites. *Molecular Biology and Evolution*, *36*(10), 2127–2142. doi: 10.1093/molbev/msz154

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., … Sham, P. C. (2007).
PLINK: a tool set for whole-genome association and population-based linkage analyses.
American Journal of Human Genetics, *81*(3), 559–575. doi: 10.1086/519795

Reynolds, M., Arapi, E. A., & Cable, J. (2018). Parasite-mediated host behavioural modifications:
Gyrodactylus turnbulli infected Trinidadian guppies increase contact rates with uninfected
conspecifics. *Parasitology*, *145*(7), 920–926. doi: 10.1017/S0031182017001950

Rieseberg, L. H., Raymond, O., Rosenthal, D. M., Lai, Z., Livingstone, K., Nakazato, T., … Lexer, C.

(2003). Major ecological transitions in wild sunflowers facilitated by hybridization. *Science*,
301(5637), 1211–1216. doi: 10.1126/science.1086949

Rogers, M. B., Downing, T., Smith, B. A., Imamura, H., Sanders, M., Svobodova, M., … Smith, D. F.
(2014). Genomic Confirmation of Hybridisation and Recent Inbreeding in a Vector-Isolated
Leishmania Population. *PLoS Genetics*, *10*(1). doi: 10.1371/journal.pgen.1004092

Schelkle, B., Faria, P. J., Johnson, M. B., van Oosterhout, C., & Cable, J. (2012). Mixed infections and
hybridisation in monogenean parasites. *PLoS ONE*, *7*(7). doi: 10.1371/journal.pone.0039506

Schön, I., van Dijk, P., & Martens, K. (2009). Lost sex: The evolutionary biology of parthenogenesis. In
Lost Sex: The Evolutionary Biology of Parthenogenesis. doi: 10.1007/978-90-481-2770-2

Schosserer, M., Minois, N., Angerer, T. B., Amring, M., Dellago, H., Harreither, E., … Grillari, J. (2015).
Methylation of ribosomal RNA by NSUN5 is a conserved mechanism modulating organismal
lifespan. *Nature Communications*, *6*(1), 1–17. doi: 10.1038/ncomms7158

Scott, M. E. (1982). Reproductive Potential of Gyrodactylus Bullatarudis (Monogenea) on Guppies
(*Poecilia reticulata*). *Parasitology*. doi: 10.1017/S0031182000055207

Smith, J. M., Smith, N. H., O'Rourke, M., & Spratt, B. G. (1993). How clonal are bacteria? *Proceedings*
of the National Academy of Sciences of the United States of America. doi:
10.1073/pnas.90.10.4384

Stephenson, J. F., van Oosterhout, C., Mohammed, R. S., & Cable, J. (2015). Parasites of Trinidadian
guppies: Evidence for sex- and age-specific trait-mediated indirect effects of predators. *Ecology*,
96(2), 489–498. doi: 10.1890/14-0495.1

Taylor, S. A., & Larson, E. L. (2019). Insights from genomes into the evolutionary importance and
prevalence of hybridization in nature. *Nature Ecology and Evolution*, *3*(2), 170–177. doi:
10.1038/s41559-018-0777-y

Tihon, E., Imamura, H., Dujardin, J. C., van Den Abbeele, J., & van den Broeck, F. (2017). Discovery
and genomic analyses of hybridization between divergent lineages of *Trypanosoma congolense*,
causative agent of Animal African Trypanosomiasis. *Molecular Ecology*, *26*(23), 6524–6538. doi:
10.1111/mec.14271

van Oosterhout, C., Harris, P. D., & Cable, J. (2003). Marked variation in parasite resistance between
two wild populations of the Trinidadian guppy, *Poecilia reticulata* (Pisces: Poeciliidae). *Biological*
Journal of the Linnean Society, *79*(4), 645–651. doi: 10.1046/j.1095-8312.2003.00203.x

Volf, P., Benkova, I., Myskova, J., Sadlova, J., Campino, L., & Ravel, C. (2007). Increased transmission
potential of *Leishmania major/Leishmania infantum* hybrids. *International Journal for*
Parasitology, *37*(6), 589–593. doi: 10.1016/j.ijpara.2007.02.002

Wendte, J. M., Miller, M. A., Lambourn, D. M., Magargal, S. L., Jessup, D. A., & Grigg, M. E. (2010).
Self-mating in the definitive host potentiates clonal outbreaks of the apicomplexan parasites
Sarcocystis neurona and Toxoplasma gondii. *PLoS Genetics*. doi: 10.1371/journal.pgen.1001261

Xavier, R., Faria, P. J., Paladini, G., van Oosterhout, C., Johnson, M., & Cable, J. (2015). Evidence for
cryptic speciation in directly transmitted gyrodactylid parasites of trinidadian guppies. *PLoS*
ONE, *10*(1). doi: 10.1371/journal.pone.0117096

Xu, S., Innes, D. J., Lynch, M., & Cristescu, M. E. (2013). The role of hybridization in the origin and
spread of asexuality in *Daphnia. Molecular Ecology*, *22*(17), 4549–4561. doi:
10.1111/mec.12407

Zhao, L., & Charlesworth, B. (2016). Resolving the conflict between associative overdominance and

background selection. *Genetics*, *203*(3), 1315–1334. doi: 10.1534/genetics.116.188912

Data accessibility

The raw sequences are available as FASTQ files in the GeneBank (BioProject accession no.

PRJNA603444).

Author Contributions

M.K. and J.R. designed research; M.K., K.J.P. and R.S.M. collected samples; C.H. provided reference

genome; M.K. analysed data with suggestions from J.R., J.C. and C.H.; MK drafted manuscript, J.R.

and J.C. contributed to manuscript writing. All authors corrected manuscript and approved the final

version.

Figure captions

Figure 1 Patterns of genetic variation in *Gyrodactylus turnbulli* populations from Trinidad and
Tobago. A: Map of Trinidad and Tobago with sampling sites. **B:** Principal Component Analyses of *G.*
turnbulli strains. PCA based on 673,104 SNPs which passed filtering criteria. SNPs were identified
within 30 isolates from three Trinidadian and three Tobagonian populations. **C:** Neighbor Joining tree
(Saitou and Nei 1987) constructed using distance matrix, with one minus identity-by-state units,
calculated based on 673,104 SNPs. Populations are represented by different colours. The reference
sequence (obtained from a strain on commercial guppies) is shown in black. **D:** Distribution of
nucleotide diversity (π) calculated across the genome in 25 kb non-overlapping windows. **E:**
Distribution of heterozygosity calculated in 25 kb non-overlapping windows for each strain collected
from Trinidad (white background) and Tobago (shaded background) populations.

Figure 2. Graphical representation of genotypes in *Gyrodactylus turnbulli* strains. Genotypes
identified as reference homozygotes are shown in green, alternative homozygotes in red,
heterozygous in yellow. Strains collected in 2017 and 2018 are labeled in grey and black respectively,
and those collected from different local populations are separated by horizontal grey lines. A subset
of polymorphic sites (10,000 SNPs) from the longest scaffolds is shown.

Figure 3. Phylogenetic relationship between *Gyrodactylus turnbulli* strains, reconstructed based on
mitochondrial genome sequences (14,159 bp). The trees were derived via a Bayesian phylogenetic
method, using 10,000,000 generations of MCMC samples, with HKY substitution model and strict

clock model, assuming 13.1% divergence per million years. The estimated time before present (in
years) for tree nodes is shown at the bottom of the tree.

Figure 4 Genetic variation in genomic region with elevated nucleotide diversity in populations from
Trinidad. A: Correlation between nucleotide diversity calculated jointly for Health Center and Dog
River populations and nucleotide diversity calculated jointly for all strains from Trinidad. Black line
represents linear regression (y = 0.00 + 0.38*x). Scaffold scf7180000443747 is shown in red. **B:**
Graphical representation of genotypes in polymorphic sites of *Gyrodactylus turnbulli* localized in
scaffold scf7180000443747 (707 SNPs). Genotypes identified as reference homozygotes are shown in
green, alternative homozygotes in red, heterozygous in yellow. Strains collected in 2017 and 2018
are shown in grey and black respectively, and those collected from different local populations are
separated by horizontal grey lines.

SNP Number

A

π_{TRI}

scf7180000443747

$\pi_{HC\&DR}$

B

LOP
G314
G312
G308
G117
G116
G113

ARI
G347
G277
G273
G136
G85
G65

CAU
G126
G35

SPS
G370
G368
G367
G334
G331
G328

DR
G285
G240
G236
G234
G190
G23

HC
G299
G291
G218
G10

SNPs in scf7180000443747