# Rapid stimulus-driven modulation of slow ocular position drifts

Tatiana Malevich[1, 2, 3]**, Antimo Buonocore[1, 2]**, Ziad M. Hafed[1, 2]*

[1] Werner Reichardt Centre for Integrative Neuroscience, Tuebingen University, Tuebingen, Germany

[2] Hertie Institute for Clinical Brain Research, Tuebingen University, Tuebingen, Germany

[3] Graduate School of Neural and Behavioural Sciences, International Max-Planck Research School, Tuebingen University, Tuebingen, Germany

* Correspondence to: ziad.m.hafed@cin.uni-tuebingen.de

** Contributed equally

**Corresponding author information:**

Ziad M. Hafed
Werner Reichardt Centre for Integrative Neuroscience
Tübingen University
Otfried-Müller Str. 25
Tübingen, Germany
72076
ziad.m.hafed@cin.uni-tuebingen.de

**Abstract**

The eyes are never still during maintained gaze fixation. When microsaccades are not occurring, ocular position exhibits continuous slow changes, often referred to as drifts. Unlike microsaccades, drifts remain to be viewed as largely random eye movements. Here we found that ocular position drifts can, instead, be very systematically stimulus-driven, and with very short latencies. We used highly precise eye tracking in three well trained macaque monkeys and found that even fleeting (~8 ms duration) stimulus presentations can robustly trigger transient and stimulus-specific modulations of ocular position drifts, and with only approximately 60 ms latency. Such drift responses are binocular, and they are most effectively elicited with large stimuli of low spatial frequency. Intriguingly, the drift responses exhibit some image pattern selectivity, and they are not explained by convergence responses, pupil constrictions, head movements, or starting eye positions. Ocular position drifts have very rapid access to exogenous visual information.

**Introduction**

Eye position changes subtly even when perfect gaze fixation is attempted (Barlow,

1952; Nachmias, 1959, 1961; Ratliff & Riggs, 1950; Steinman, Haddad, Skavenski, &

Wyman, 1973). Such "fixational" eye movement comes in two primary flavors:

microsaccades, which resemble large saccades (Hafed, 2011; Rolfs, 2009; Zuber,

Stark, & Cook, 1965) and rapidly shift gaze position by a minute amount; and ocular

position drifts, which are even smaller and slower eye movements (Barlow, 1952;

Ditchburn & Ginsborg, 1953; Martins, Kowler, & Palmer, 1985; Nachmias, 1959,

1961; Poletti & Rucci, 2015; Ratliff & Riggs, 1950; Skinner, Buonocore, & Hafed,

2019; St Cyr & Fender, 1969).

The mechanisms for generating and influencing microsaccades have received much

attention (Hafed, 2011; Hafed, Chen, & Tian, 2015; Krauzlis, Goffart, & Hafed, 2017).

Despite contrary ideas in the past century, a now accepted property of

microsaccades is that, like larger saccades, they are not random but are very

systematically (Engbert, 2006; Engbert & Kliegl, 2003; Hafed & Clark, 2002; Hafed &

Ignashchenkova, 2013; Hafed, Lovejoy, & Krauzlis, 2011; Ko, Poletti, & Rucci, 2010;

Rolfs, Kliegl, & Engbert, 2008; Thaler, Schutz, Goodale, & Gegenfurtner, 2013; Tian,

Yoshida, & Hafed, 2016) and rapidly (Buonocore, Chen, et al., 2017; Hafed &

Ignashchenkova, 2013; Rolfs et al., 2008; Tian et al., 2016; Tian, Yoshida, & Hafed,

2018) influenced by peripheral as well as foveal visual stimuli, among other factors

(Hafed et al., 2015; Willeke et al., 2019). In stark contrast, the brain mechanisms for

controlling ocular position drifts are unknown; these movements continue to be

thought of as random, often being modeled as random walk processes (Burak,

Rokni, Meister, & Sompolinsky, 2010; Engbert & Kliegl, 2004; Engbert, Mergenthaler,

Sinn, & Pikovsky, 2011; Herrmann, Metzler, & Engbert, 2017; Kuang, Poletti, Victor,

& Rucci, 2012). This is despite the fact that slow ocular position drifts exhibit

behavioral characteristics (e.g. Ditchburn & Ginsborg, 1953; Nachmias, 1961)

supporting some level of central nervous system control over them. For example,

these eye movements aid in stabilizing eye position near a certain location (e.g.

Epelboim & Kowler, 1993); they can compensate for subtle head movements through

mechanisms likely related to the vestibular ocular reflex (e.g. Ditchburn & Ginsborg,

1953; Poletti, Aytekin, & Rucci, 2015; Schor & Westall, 1984; Skavenski, Hansen,

Steinman, & Winterson, 1979; Steinman & Collewijn, 1980); and they also appear to

be associated with internal knowledge of their occurrence for regulating the

perception (or lack thereof) of the image motions that they cause on the retina (e.g.

Poletti, Listorti, & Rucci, 2010).

All of the above hallmarks of brain control over drifts, however, still do not rule out the

possibility that such control may be somewhat "loose". For example, the movements

could still be random walks but nonetheless have statistics sufficient to modulate

retinal image information for the purposes of aiding spatial vision (Kuang et al., 2012;

Rucci & Victor, 2015). This leaves the question of how precisely these eye

movements may potentially be controlled still unanswered. Here, we used precise

eye tracking in three well trained rhesus macaque monkeys to demonstrate that

ocular position drifts can exhibit highly systematic stimulus-driven modulations in

both speed and direction. These modulations have a very short latency, and they are

stimulus-tuned and binocular. These modulations are also independent of head

movements, convergence responses, and starting eye positions. Moreover, their

amplitudes are sufficient to move images across individual cone photoreceptors in

the fovea or multiple rod photoreceptors in some peripheral zones.

Our results, coupled with evidence that drift statistics adapt to a variety of behavioral

task constraints (Chen & Hafed, 2013; Cherici, Kuang, Poletti, & Rucci, 2012;

Epelboim & Kowler, 1993; Intoy & Rucci, 2020; Poletti et al., 2015; Poletti et al.,

2010; Shelchkova, Tang, & Poletti, 2019; Skinner et al., 2019; Tian et al., 2018),

strongly motivate deeper research into the neurophysiological mechanisms

controlling incessant ocular position drifts in between saccades.

## Results

*Short-latency, stimulus-driven ocular position drift response*

We analyzed microsaccade-free fixation after a visual transient in monkeys highly

trained to fixate on tiny targets. In a first experiment, each monkey fixated a white

spot over a gray background spanning approximately +/- 15 deg horizontally and +/-

11 deg vertically. At a random time, the display changed: one entire half (right or left)

became black (split view stimulus); and a half-circle of 0.74-deg radius around the

fixated position remained gray, in order to maintain view of the stable fixation spot

(Methods). We explicitly used such a patterned stimulus configuration (as opposed to

just a flash as in some subsequent experiments described below) in order to explore

any potential stimulus selectivity of ocular position drifts, as we elaborate on shortly.

In trials without microsaccades (-100 to 200 ms from stimulus onset), a highly

repeatable short-latency drift response occurred. This drift response is illustrated in

Fig. 1A, B for monkey A, in which we tracked the left eye using the precise scleral

search coil technique (Fuchs & Robinson, 1966; Judge, Richmond, & Chu, 1980); the

response measured in the two other monkeys, in which we measured right eye

position, is shown in Fig. 1 – figure supplement 1. Regardless of the tracked eye, and

regardless of the steady-state stereotypical drift direction exhibited individually by a

given monkey, there was always a robust upward position drift after stimulus onset.

Importantly, it was clearly visible at the individual trial level (Fig. 1C, Fig. 1 – figure

supplement 1C, I). In one monkey (N), this upward drift response was sometimes

initiated first by a much more short-lived and downward shift of eye position (of very

small magnitude) before the flip to the upward drift (Fig. 1 – figure supplement 1G-L).

We statistically assessed the onset and duration of the drift response in each monkey

by computing 95% confidence intervals (Methods) around the average eye velocity

traces. When the 95% confidence intervals between the stimulus and control velocity

traces did not overlap for at least 20 consecutive milliseconds, we deemed the

velocity response to be significant. In monkey A, the upward velocity pulse started at

~80 ms and lasted for ~60 ms (Fig. 1B). These values were ~60 ms and ~90 ms,

respectively, for monkey M (Fig. 1 – figure supplement 1B) and ~70 ms and ~100

ms, respectively, for monkey N (Fig. 1 – figure supplement 1H). Note that Fig. 1A and

Fig. 1 – figure supplement 1A, G also show significance intervals for the eye position

traces for completeness, although these provide a more conservative estimate of

response onset time (Methods).

**Figure 1 Short-latency, stimulus-driven ocular position drift response. (A)** Vertical eye position
(plus 95% confidence intervals) after split view stimulus onset. Compared to the no-stimulus condition
(gray), the eye drifted upwards with short latency after stimulus onset. This happened for both right and
left darkness stimuli (the two panels). We vertically aligned all starting eye positions at time 0 before
averaging trials, to highlight the systematic change in eye position shortly after stimulus onset. **(B)** Same
data as in **A** but for vertical eye velocity. A clear velocity pulse (much slower than that of
microsaccades/saccades) is visible. **(C)** Individual trials demonstrating the drift response. The vertical
position of each curve was jittered across the shown trials to facilitate visibility. **(D)** When we included
progressively longer intervals of saccade-free fixation, we confirmed that the drift response (e.g. **B**) is
transient. **(E, F)** Relative to starting eye position at stimulus onset, the drift response was predominantly
vertical, but the horizontal component of eye position was also stimulus-dependent. Error bars in **F**
denote 95% confidence intervals. The trajectories in **E** are two-dimensional plots of the average curves
in **A**, **F**. All pink lines on x-axes indicate intervals in which the 95% confidence intervals of the two
compared curves did not overlap. Also see figure supplements 1 and 2.

In all three monkeys, the drift response consisted of a predominantly upward velocity

pulse reaching a peak speed of ~33-45 min arc/s, or ~0.55-0.75 deg/s (Fig. 1B, D,

Fig. 1 – figure supplement 1B, D, H, J) and a resultant displacement of ~2-3 min arc,

or 0.034-0.05 deg (Fig. 1A, Fig. 1 – figure supplement 1A, G). Such speeds and

displacements are well within the detectability ranges of the visual system. For

example, foveal cone photoreceptor separation in the rhesus macaque retina is ~1

min arc (Rolls & Cowey, 1970), and rod photoreceptor density can be even higher in

the rod peak region (Wikler, Williams, & Rakic, 1990). Thus, the stimulus-driven drift

response clearly modulates input images to the visual system.

The drift response was also transient: when we considered longer periods of

saccade-free fixation, we did not notice increasing durations of drift response.

Instead, the eye velocity "pulse" subsided, and eye position reverted back to its

steady-state drift direction that was prominent, and stereotyped, in each monkey (Fig.

1D, Fig. 1 – figure supplement 1D, J).

The drift response also showed horizontal direction dependence on the stimulus

properties, and this is why our use of the split view stimulus in the first experiment

was relevant. In all three monkeys, the horizontal component of the eye movement

during the drift pulse was systematically biased by the side of darkness (Fig. 1E, F,

Fig. 1 – figure supplement 1E, F, K, L): the upward drift tilted rightward when the

darkness was on the right of gaze and leftward when it was on the left, and these

results were statistically significant (pink intervals on all relevant x-axes of plots;

Methods). The horizontal component of eye position therefore reacted to the spatial

position of the luminance transient. This was our first evidence that the drift response

can be stimulus-specific, a property that we revisit below.

*The drift response is neither an eye tracking artifact nor a convergence reflex*

We considered the possibility that the stimulus-driven drift response was part of a

near reflex triad (convergence, accommodation, and pupil constriction) (Myers &

Stark, 1990; Nachmias, 1961). For example, the luminance transient may have been

perceived by the monkey as a change in depth plane. However, this is unlikely to

explain our observations: the horizontal bias as a function of stimulus appearance

(e.g. Fig. 1E, F) occurred regardless of whether we tracked the right (monkeys M and

N) or left (monkey A) eye. For example, if the drift was part of a convergence

movement, it would not be biased rightward when tracked in the right eye (monkey

M). We also explicitly tested for convergence in a later experiment, described below,

in which we measured eye movements binocularly.

Our results also cannot be explained by variations in pupil diameter, which can cause

artifactual eye drift measurements in video-based eye trackers (Choe, Blake, & Lee,

2016; Drewes, Zhu, Hu, & Hu, 2014; Hooge, Hessels, & Nystrom, 2019; Kimmel,

Mammo, & Newsome, 2012; Wildenmann & Schaeffel, 2013; Wyatt, 2010); in all of

our analyses, we measured eye position using scleral search coils. Moreover,

pupillary responses to stimulus transients exhibit significantly slower dynamics

(Clarke, Zhang, & Gamlin, 2003; Pong & Fuchs, 2000). To confirm this, we

performed a control experiment in which we explicitly measured pupil diameter in one

monkey (A) using a video-based eye tracker. Using a stimulus manipulation in which

the upward drift response was clearly visible in the (left) eye (tracked with a scleral

search coil), simultaneous measurement of pupil diameter in the other eye (tracked

with a video-based eye tracker; Methods) revealed that the pupillary constriction

caused by stimulus onset occurred later than the ocular position drift response (Fig. 1

– figure supplement 2). It was futile to look for the drift response in the right eye,

confirming that video-based eye trackers are not capable of resolving such small

detail (Choe et al., 2016; Kimmel et al., 2012; Wyatt, 2010).

*The drift response is a binocular eye movement*

To directly investigate the convergence question, we next implanted one monkey (M)

with a second scleral search coil, this time in the left eye. We could therefore

measure binocular movements simultaneously. The monkey performed a new task in

which we avoided a prolonged period of darkness in half of the display. Instead, the

monkey steadily fixated with the original gray background. Then, for only one display

frame (~8 ms), we flipped the luminance of the gray screen to black before it returned

to gray again (Methods). We still observed a similar drift response (Fig. 2A, C).

Importantly, the drift response occurred in both eyes simultaneously (Fig. 2A, C), and

the horizontal component of eye position did not show an appreciable convergence

response of similar magnitude (Fig. 2B, D). We confirmed this statistically; in a time

interval capturing the drift response (50-140 ms), there was no statistically significant

difference between right and left peak vertical or peak horizontal eye velocity

(p=0.249, Z=1.15 for vertical and p=0.8321, Z=0.21 for horizontal; Wilcoxon rank sum

test). Therefore, the stimulus-driven drift response (Figs. 1, 2) is a binocular eye

movement phenomenon that is independent of a convergence response.

**Figure 2 The drift response is a binocular eye movement. (A)** Vertical eye position from monkey M during simultaneous measurements of the right and left eyes. The upward drift response (for a brief ~8 ms black flash covering the whole display) occurred binocularly. Note that this response superseded the monkey's stereotypical baseline slow upward drift, which was much slower (each monkey had a stereotypical baseline drift direction; Fig. 1, Fig. 1 – figure supplement 1). **(B)** Horizontal eye position showed no evidence of convergence. **(C, D)** Same data with vertical and horizontal eye velocity, respectively. The drift response (**C**) was not associated with a convergence response (**D**). The lack of horizontal pink lines on the x-axes of **C**, **D** confirm that both eyes moved similarly. **(E)** In monkey A, we confirmed that the same paradigm (brief black flash) was still sufficient (bluish curve) to cause an upward drift response. When we switched the flash to white (reddish curve), the drift response still occurred. **(F)** Monkey M also showed a drift response for a white flash. Error bars in all panels denote 95% confidence intervals, and pink lines on the x-axes show periods in which the 95% confidence intervals of the two compared curves did not overlap.

Naturally, we also tested a second monkey (A) on this new black fixation flash

paradigm (albeit monocularly), and we still observed the drift response (Fig. 2E,

bluish curve). Therefore, the drift response is not a peculiarity of the split view

stimulus, and it still occurred in two monkeys under a completely different protocol.

*The drift response is not the same as a steady-state up-shift in gaze in darkness*

We next asked whether our observations were related to a previously reported

steady-state upward eye position deviation in darkness (Barash, Melikyan, Sivakov,

& Tauber, 1998; Goffart, Quinet, Chavane, & Masson, 2006; Snodderly, 1987). After

all, our split view stimulus introduced substantial darkness (in an already dark

laboratory), and the fixation flash paradigm (Fig. 2A-D) also introduced darkness,

although for just a fleeting moment. To test this, we ran the same two monkeys (A

and M) on a third condition in which the entire display went from gray to white instead

of black (again, for only ~8 ms; Methods). For both white and black flashes, a similar

stimulus-driven transient ocular position drift response occurred (Fig. 2E, F).

Statistically, peak eye velocity was significantly higher with black than white flashes

in both monkeys (p=4.2x10$^{-14}$, Z=7.55 for monkey A and p=4.3x10$^{-5}$, Z=4.09 for

monkey M; Wilcoxon rank sum test), suggesting a secondary impact of stimulus

polarity on the oculomotor system (Malevich, Buonocore, & Hafed, 2020).

*The drift response is not associated with a reflexive head movement*

Our results so far involved measuring only eye position; our monkeys' heads were

robustly fixed with a metal holder (Methods). Nonetheless, it could be argued that the

drift response is potentially a stabilizing reflex due to stimulus-driven head

movements (e.g. due to, say, startle by the monkeys after sudden flashes). Indeed,

microscopic head movements (Corneil & Munoz, 2014) may be associated with small

compensatory eye movements (Ditchburn & Ginsborg, 1953; Poletti et al., 2015;

Schor & Westall, 1984; Skavenski et al., 1979; Steinman & Collewijn, 1980).

Alternatively, if the head were to move in a stimulus-driven manner, then the drift

response could be an artifactual measurement aberration due to movement of the

head within the fixed magnetic fields of the scleral search coil system used to track

eye movements; such potential head movement would, then, alter the measured eye

position, but only due to physical displacement of the eye (by the head movement) in

the magnetic fields and not because of a genuine eyeball rotation.

While it would be surprising to get a startle-induced head movement with such short

latencies for our stimuli (some of which were very subtle and brief; Fig. 2), we

nonetheless performed additional control experiments in which we simultaneously

measured eye and head position. We placed a "head coil" in three different

positions/orientations on the monkey's head, across three different sets of

experiments, and we digitized the head coil output (into two channels) like we

digitized the eye coil measurements into two channels (Methods). We used three

different head coil positions on the head (in three sets of experiments) because

head-fixation apparati and recording chambers implanted on the head resulted in

non-standard head coil orientations relative to the magnetic fields around the head

used to induce coil currents; we therefore wanted to ensure that there was no

particular axis of head movement that we would potentially miss from using only one

potential unfortunate head coil orientation in one given experiment. We also used the

same black fixation flash paradigm of Fig. 2, and we performed these additional

control experiments in monkey M.

For all three sets of new experiments, we replicated the drift response observation in

the same monkey (Fig. 2 – figure supplement 1A). Simultaneous measurements of

head position (Fig. 2 – figure supplement 1B, C) did not reveal any concomitant head

movements. To assess this statistically, we used a velocity estimate for both eye and

head position measurements in each experiment (similar to our earlier analyses

above). We specifically defined a pre-stimulus baseline interval and a post-stimulus

"drift" interval (indicated in pink in Fig. 2 – figure supplement 1). Within each interval,

we averaged the velocity measure, resulting in a paired measurement between

intervals across trials. For eye movements, there was a significant difference

between vertical eye velocity during the post-stimulus drift interval and the pre-

stimulus baseline interval (p=5.55x10$^{-19}$ [Z=-8.9], p=3.27x10$^{-18}$ [Z=-8.7], p=1.05x10$^{-13}$

[Z=-7.44], respectively, for the three head coil positions in Fig. 2 – figure supplement

1A; Wilcoxon rank sum test). This directly replicates our earlier drift response

observations in multiple monkeys above. For head movements, none of the head coil

positions that we tested resulted in substantial movements after stimulus onset

(p=0.1394 to 0.8911 [Z=-0.91 to 1.48] across all positions and channels in Fig. 2 –

figure supplement 1B, C; Wilcoxon rank sum test). These results mean that the head

did not move at the same time as the ocular position drift response, ruling out both a

compensatory eye movement explanation as well as an artifactual change in sensed

position due to displacement of the eye's 3-dimensional position in the search coil

system's magnetic fields.

*The drift response is feature-tuned*

A significant swath (i.e. low spatial frequency) of luminance transience seemed

necessary for the drift response. We did not observe the short-latency response as

strongly with small localized targets briefly flashed to the right or left of fixation (Fig. 2

– figure supplement 2). Our earlier experiments with small localized stimuli (Tian et

al., 2018) also only revealed a later position modulation than the drift response being

characterized in the current study (which could have been caused by stimulus-

modulated microsaccades).

Given that visual responses in oculomotor areas like the superior colliculus (SC)

preferentially represent low spatial frequencies (Chen, Sonnenberg, Weller, Witschel,

& Hafed, 2018), we therefore wondered whether the ocular position drift response

could exhibit tuning to spatial frequency. We replicated the same full-screen flash

paradigm in one monkey (M) but now with different spatial frequencies (0.55-6.8

cycles/deg; cpd). We saw clear preference for low spatial frequencies in the drift

response, both in response latency and magnitude (Fig. 3A). To confirm this

statistically, we searched for peak eye velocity in the interval 80-140 ms after

stimulus onset (Fig. 3). We then characterized peak eye velocity and peak eye

velocity latency as a function of spatial frequency using a Kruskal-Wallis test. There

was an effect of spatial frequency on both peak velocity (p=$2.2 \times 10^{-16}$, $\chi^2(4)$=79.49)

and peak velocity latency (p=$3.1 \times 10^{-17}$, $\chi^2(4)$=83.56). Post-hoc tests revealed that the

drift response was earliest (adjusted p<$9.3 \times 10^{-8}$ against all other spatial frequencies)

and strongest (adjusted p<0.0021 against the third to fifth spatial frequencies) for the

lowest spatial frequency. With our split view stimulus and display geometry (tested in

all three monkeys), the effective spatial frequency was even lower than 0.55 cpd, and

the drift response appeared to be the strongest (Fig. 1, Fig. 1 – figure supplement 1).

Therefore, the ocular position drift response not only occurs early enough to reflect

potential effects of early sensory responses in the oculomotor system (e.g. SC visual

bursts; Chen et al., 2018), but it is also sensitive to stimulus specifics. In fact, when

we ran yet another experiment with grating orientation, rather than spatial frequency,

the drift response of the same monkey was again altered, and it displayed a form of

orientation tuning (Fig. 3B; p=0.01, $\chi^2(4)$=11.0 for peak velocity and p=0.0004,

$\chi^2(4)$=18.5 for peak velocity latency; Kruskal-Wallis test). Thus, ocular position drifts

are sensitive, with short latency, to stimulus transients and their image features.

**Figure 3 Stimulus tuning of the ocular position drift response. (A)** We replaced the full-screen flash of Fig. 2 with a vertical grating of different spatial frequencies (Methods). The upward velocity pulse depended on spatial frequency; it occurred earlier, and had stronger amplitude, for low spatial frequencies. This is reminiscent of visual neural activity in the superior colliculus, an important oculomotor control circuit (Chen et al., 2018). **(B)** Similarly, when we fixed spatial frequency and altered grating orientation, the drift response was also modulated. Ocular position drifts are therefore sensitive to stimulus manipulations. Error bars denote 95% confidence intervals.

*The drift response happens at a time of complete saccade inhibition and is*

*independent of starting eye position*

In all of our analyses above, we isolated the properties of the drift response by

analyzing saccade-free epochs. However, in reality, saccades could still occur even

in our fixation paradigms. We therefore now considered the trials in which fixational

microsaccades did occur. We found a remarkably complementary nature between

the timing of microsaccades and the timing of the ocular position drift response.

Specifically, it was long known that saccade and microsaccade likelihood decreases

dramatically shortly after visual transients (Buonocore, Chen, et al., 2017; Buonocore

& McIntosh, 2008; Edelman & Xu, 2009; Engbert & Kliegl, 2003; Hafed &

Ignashchenkova, 2013; Hafed et al., 2011; Reingold & Stampe, 2002). We replicated

this finding in all of our three monkeys, and we also compared the timing of the drift

response to the timing of such saccadic inhibition (Fig. 4A-F; split view stimulus). The

drift response occurred when it was least likely to observe microsaccades. Even

though the mechanisms of (micro)saccadic inhibition are still debated, we recently

hypothesized that sensory transients in the oculomotor system (e.g. SC) play a

particularly important role (Buonocore, Chen, et al., 2017). The stimulus dependence

of the ocular drift response (e.g. Fig. 3) and its timing relative to saccadic inhibition

(and indeed relative to the timing of early sensory responses in the oculomotor

system; Chen et al., 2018) suggest that the ocular drift response is mechanistically

linked to the circuits mediating saccadic inhibition; and particularly to the short-

latency visual responses in these circuits. That is, stimulus onset triggers both

saccadic inhibition and drift response. Future neurophysiological experiments will

have to directly test this hypothesis.

Moreover, we found that the drift response could still occur with nearby

microsaccades before or after it. Specifically, for a given experiment, we picked trials

in which microsaccades occurred within a given time range (e.g. up to 0 ms relative

to stimulus onset). We then plotted the average eye velocity in the remainder of

these trials (i.e. the portion of eye velocity that was non-saccadic). For different

microsaccade times relative to stimulus onset, we observed similar amplitudes of the

ocular drift response (Fig. 4G-I). This also happened in our other tasks (e.g. Fig. 4 –

figure supplement 1). Interestingly, the amplitude of the drift response seemed similar

regardless of the timing of nearby microsaccades (Fig. 4G-I). This is an interesting

contrast to our earlier findings that the amplitudes of the faster (and later) initial

ocular following responses after full-screen motion pulses are strongly modulated

when microsaccades happen near the time of stimulus (i.e. full-screen motion)

onsets (Chen & Hafed, 2013).

**Figure 4 The drift response starts when saccades are inhibited. (A-C)** Our monkeys exhibited
known saccadic inhibition (Buonocore, Chen, et al., 2017; Buonocore & McIntosh, 2008; Edelman & Xu,
2009; Hafed & Ignashchenkova, 2013; Hafed et al., 2011; Reingold & Stampe, 2002) after stimulus
onset (split view condition shown; figure supplement 1 shows the fixation flash condition). Error bars
denote 95% confidence intervals. **(D-F)** In the same sessions, saccade-free fixation epochs (no
saccades from -100 to 350 ms) showed the ocular position drift response (error bars denote 95%
confidence intervals). The timing of this drift response was linked to the timing of saccadic inhibition
(individual dots show individual trial rasters of microsaccade onset times from **A-C**). The drift response
started when microsaccades were inhibited. **(G-I)** When we combined **A-C** and **D-F**, we found that even
when microsaccades occurred near the time of drift response (i.e. before or after with shorter saccade-
free epochs than in **D-F**), the drift response could still occur (in **G**, two curves are missing because they
had <10 trials due to the prolonged saccadic inhibition profile in **A**). The different colors show traces in

which there were microsaccades within a given time range relative to stimulus onset. For example,
"Saccades up to -50 ms" means that the curve shows average smooth eye velocity from -50 ms
onwards, since earlier times had saccadic velocities in them. Similarly, the curve for "Saccades after
175 ms" would mean that average eye velocity is shown up to 175 ms, since later times relative to
stimulus onset had saccadic velocities in them. Error bars in **D**-**F** denote 95% confidence intervals. Also
see figure supplement 1 and Fig. 2 – figure supplements 3 and 4.

Finally, we checked whether the drift response could still occur with different starting

eye positions. We divided trials based on eye position being above or below median

vertical eye position across trials (Methods), and the same upward drift response was

present (Fig. 2 – figure supplement 3 shows these results from the black fixation

flash paradigm in monkeys A and M). And, the drift response was too small to even

elicit corrective downward microsaccades after it occurred: there was no clear bias

for downward microsaccades after the drift response (Fig. 2 – figure supplement 4).

**Discussion**

Our results demonstrate that slow fixational drift eye movements are systematically

influenced by visual transients. These results motivate much deeper

neurophysiological investigation of drift eye movement control, especially given that

these eye movements are incessant and modulate the spatio-temporal statistics of

input images to the visual system (Kuang et al., 2012; Rucci, Iovin, Poletti, & Santini,

2007; Rucci & Victor, 2015). Classic brainstem neurophysiological studies of the

oculomotor system (e.g. reviewed in: Takahashi & Shinoda, 2018) are too coarse to

provide the mechanistic insights necessary for explaining our observations. They

need to be revisited with the perspective of understanding millisecond-by-millisecond

tiny deviations of eye position during fixational drift.

Visual representations also need to be considered. Given the temporal transience of

the drift response and its dependence on spatial frequency, it is likely that visual

responses in oculomotor circuitry play a role. One possibility is that low spatial

frequency stimuli increase the spatial uncertainty of intended gaze position. It is

conceivable that such increased uncertainty, coupled with overrepresentation of the

upper visual field in the SC (Hafed & Chen, 2016), as well as earlier visual bursts for

upper visual field locations, results in transient upward drift responses. That is,

increased spatial uncertainty increases the size of the active population of neurons,

and if more neurons represent the upper visual field (Hafed & Chen, 2016), then the

transient response would be biased in the upward direction. Indeed, the SC

influences eye position (Hafed, Goffart, & Krauzlis, 2008); what remains is to

understand whether instantaneous SC activity landscapes, in cooperation with the

deeper brainstem, can explain instantaneous variations in fixational eye position.

This would suggest fairly high level central nervous system control of these eye

movements.

Given the short latency of the drift response, an even more intriguing possibility than

SC involvement is the contribution of downstream oculomotor control structures. The

question of whether primate pre-motor and motor brainstem nuclei possess early,

and feature-tuned, visual responses remains completely open. In the cat, visual

responses in the nucleus raphe interpositus (rip) have been reported (Evinger,

Kaneko, & Fuchs, 1982), and there are hints of visual responses in monkey rip

(Busettini & Mays, 2003; Everling, Pare, Dorris, & Munoz, 1998; Missal & Keller,

2002), although no visual tuning was explicitly investigated. Given both the

preponderance and richness of SC visual responses (e.g. Chen & Hafed, 2018;

Chen, Hoffmann, Distler, & Hafed, 2019; Chen et al., 2018), and given the fact that

the SC projects to brainstem oculomotor control circuits, we hypothesize the

presence of feature-tuned visual responses in such oculomotor control circuits. Such

feature-tuned responses could be particularly useful to interpret the intriguing

orientation tuning effects that we observed in Fig. 3B. Our ongoing

neurophysiological experiments do support this view (Buonocore, Baumann, &

Hafed, 2020), and they allow us to predict that lower brainstem visual responses

might be additionally critical for the ocular position drift characteristics that we

observed in the current study.

Naturally, one might also wonder whether the drift response is specific to monkeys.

We think not. Indeed, in an earlier human study in which natural images (rich in low

spatial frequency content) were presented during fixation, drift speeds were notably

the highest in the very first analyzed time bin after image onset (Poletti & Rucci,

2010). While the time bins in that study were too coarse to explore the short latency

responses that we saw, these human results are, at least, consistent with our

observation that image onsets can rapidly modulate ocular drifts. Other oculomotor

phenomena are also very well preserved between monkeys and humans: some

notable examples from our own prior work include ocular following responses (Chen

& Hafed, 2013) as well as saccade-related implications of SC visual field

asymmetries (Grujic, Brehm, Gloge, Zhuo, & Hafed, 2018; Hafed & Chen, 2016;

Hafed & Goffart, 2020) and SC foveal magnification (Chen et al., 2019; Hafed &

Goffart, 2020; Willeke et al., 2019).

Functionally, we believe that the timing of the drift response in relation to the timing of

saccadic inhibition (Fig. 4) might provide potential hints about the functional roles of

the drift response for visual coding and perception. Saccadic inhibition is thought to

reflect a resetting of ongoing saccadic rhythms to enable reacting (often with a

subsequent eye movement) to the external stimulus arriving asynchronously to such

rhythms (Buonocore, Chen, et al., 2017; Hafed & Ignashchenkova, 2013). Increasing

behavioral work is also suggesting that the brief period of such saccadic inhibition

additionally provides an important time window during which perceptual and cognitive

processing of the incoming external stimulus could proceed (Bompas, Campbell, &

Sumner, 2020; Buonocore, Purokayastha, & McIntosh, 2017; Salinas & Stanford,

2018). As part of this brief processing, under some situations like in our image

patterns that caused the drift response, it might be important to modulate the visual

statistics of the input images with drift eye movements. Therefore, the drift response

(which introduces a coherent motion of certain speed) could serve this function, and

this would suggest a short time-scale version of the longer term modulations of how

drifts are believed to modulate image statistics (e.g. to enhance edges and equalize

spectral content of images; Kuang et al., 2012). Coupled with evidence that different

subclasses of visual neurons (e.g. in primary visual cortex) will be differentially

activated by different speeds and amplitudes of fixational eye movements

(Snodderly, 2016), it is conceivable that the drift response that we observed could

functionally serve to preferentially activate the most suitable classes of visual

neurons in cortex for processing the kinds of images that we presented in our

experiments (and that were also presented in: Poletti & Rucci, 2010). Indeed, during

smooth pursuit initiation (i.e. with very small eye velocities, albeit still faster than the

ocular position drifts in our current study), when we used the same flash paradigms

as in the current study, we found that the stimulus onsets consistently reduced

smooth eye velocity rather than enhanced it (Buonocore, Skinner, & Hafed, 2019).

This suggests that, during fixation, flexibility to enhance smooth eye velocity is

maintained, potentially to aid in visual processing.

In all, our results directly complement an increasing body of research on slow ocular

position drifts during fixation, both on how they might be controlled as well as how

they might be used for supporting vision. Early behavioral work focused on control,

and even eventually invoked the term "slow control" to describe these eye

movements (as opposed to "drifts"). This already suggested a kind of central nervous

system control over these eye movements (Epelboim & Kowler, 1993; Nachmias,

1961). Similarly, vestibular ocular and other reflexes due to subtle head movements

could result in small eye movements on the scale of ocular position drifts (Ditchburn

& Ginsborg, 1953; Poletti et al., 2015; Schor & Westall, 1984; Skavenski et al., 1979;

Steinman & Collewijn, 1980), and ocular tracking of very slow and small-amplitude

motion trajectories is possible in both humans and monkeys (Cunitz, 1970; Martins et

al., 1985; Skinner et al., 2019). Our results add to these observations the possibility

of very rapid access to early sensory information by the oculomotor system, and they

will hopefully motivate studies and models of how ocular position drifts can be

generated on a millisecond-by-millisecond basis. This complements other recent

work investigating the visual and perceptual consequences of drifts. For example, in

(Poletti et al., 2010), it was suggested that the brain has some internal knowledge of

these movements, again consistent with central control. And, in theoretical accounts,

drifts have been suggested to modulate image statistics for aiding in visual image

processing (Kuang et al., 2012; Rucci & Victor, 2015). Interestingly, such accounts

can still be functional if drifts were completely random movements, as long as these

random movements possessed statistics that could functionally whiten the spectral

content of natural images. Our results complement these latest theoretical accounts

by demonstrating robust drift modulations on much shorter time scales, and which

are also anything but random.

**Acknowledgements**

We were funded by the Deutsche Forschungsgemeinschaft (DFG) through the

Research Unit: FOR 1847 (project: HA6749/2-1). We were also funded by the

Werner Reichardt Centre for Integrative Neuroscience (CIN; DFG EXC307) and the

Hertie Institute for Clinical Brain Research. TM and ZMH were additionally supported

by a CIN intramural grant (Mini_GK 2017-04). We thank Tong Zhang and Matthias

Baumann for helping us to perform the control experiments measuring both eye and

head position simultaneously, and we also thank Joachim Bellet for help in collecting

monkey N data.

**Author Contributions**

TM, AB, ZMH collected and analyzed the data. TM, AB, ZMH wrote and edited the

manuscript.

**Declaration of Interests**

The authors declare no competing interests.

**References**

Barash, S., Melikyan, A., Sivakov, A., & Tauber, M. (1998). Shift of visual fixation dependent on background illumination. *J Neurophysiol, 79*(5), 2766-2781. doi:10.1152/jn.1998.79.5.2766

Barlow, H. B. (1952). Eye movements during fixation. *J Physiol, 116*(3), 290-306.

Bellet, M. E., Bellet, J., Nienborg, H., Hafed, Z. M., & Berens, P. (2019). Human-level saccade detection performance using deep neural networks. *J Neurophysiol, 121*(2), 646-661. doi:10.1152/jn.00601.2018

Bompas, A., Campbell, A. E., & Sumner, P. (2020). Cognitive control and automatic interference in mind and brain: A unified model of saccadic inhibition and countermanding. *Psychol Rev, 127*(4), 524-561. doi:10.1037/rev0000181

Buonocore, A., Baumann, M. P., & Hafed, Z. M. (2020). Visual pattern analysis by motor neurons (Abstract). *Computational and Systems Neuroscience (Cosyne) 2020 Conference*, 147.

Buonocore, A., Chen, C. Y., Tian, X., Idrees, S., Munch, T. A., & Hafed, Z. M. (2017). Alteration of the microsaccadic velocity-amplitude main sequence relationship after visual transients: implications for models of saccade control. *J Neurophysiol, 117*(5), 1894-1910. doi:10.1152/jn.00811.2016

Buonocore, A., & McIntosh, R. D. (2008). Saccadic inhibition underlies the remote distractor effect. *Exp Brain Res, 191*(1), 117-122. doi:10.1007/s00221-008-1558-7

Buonocore, A., Purokayastha, S., & McIntosh, R. D. (2017). Saccade Reorienting Is Facilitated by Pausing the Oculomotor Program. *J Cogn Neurosci, 29*(12), 2068-2080. doi:10.1162/jocn_a_01179

Buonocore, A., Skinner, J., & Hafed, Z. M. (2019). Eye Position Error Influence over "Open-Loop" Smooth Pursuit Initiation. *J Neurosci, 39*(14), 2709-2721. doi:10.1523/JNEUROSCI.2178-18.2019

Burak, Y., Rokni, U., Meister, M., & Sompolinsky, H. (2010). Bayesian model of dynamic image stabilization in the visual system. *Proc Natl Acad Sci U S A, 107*(45), 19525-19530. doi:10.1073/pnas.1006076107

Busettini, C., & Mays, L. E. (2003). Pontine omnipause activity during conjugate and disconjugate eye movements in macaques. *J Neurophysiol, 90*(6), 3838-3853. doi:10.1152/jn.00858.2002

Chen, C. Y., & Hafed, Z. M. (2013). Postmicrosaccadic enhancement of slow eye movements. *The Journal of neuroscience : the official journal of the Society for Neuroscience, 33*(12), 5375-5386. doi:10.1523/JNEUROSCI.3703-12.2013

Chen, C. Y., & Hafed, Z. M. (2018). Orientation and Contrast Tuning Properties and Temporal Flicker Fusion Characteristics of Primate Superior Colliculus Neurons. *Front Neural Circuits, 12*, 58. doi:10.3389/fncir.2018.00058

Chen, C. Y., Hoffmann, K. P., Distler, C., & Hafed, Z. M. (2019). The Foveal Visual Representation of the Primate Superior Colliculus. *Curr Biol, 29*(13), 2109-2119 e2107. doi:10.1016/j.cub.2019.05.040

Chen, C. Y., Sonnenberg, L., Weller, S., Witschel, T., & Hafed, Z. M. (2018). Spatial frequency sensitivity in macaque midbrain. *Nature communications, 9*(1), 2852. doi:10.1038/s41467-018-05302-5

Cherici, C., Kuang, X., Poletti, M., & Rucci, M. (2012). Precision of sustained fixation in trained and untrained observers. *Journal of vision, 12*(6). doi:10.1167/12.6.31

Choe, K. W., Blake, R., & Lee, S. H. (2016). Pupil size dynamics during fixation impact the accuracy and precision of video-based gaze estimation. *Vision Res, 118*, 48-59. doi:10.1016/j.visres.2014.12.018

Clarke, R. J., Zhang, H., & Gamlin, P. D. (2003). Characteristics of the pupillary light reflex in the alert rhesus monkey. *J Neurophysiol, 89*(6), 3179-3189. doi:10.1152/jn.01131.2002

Corneil, B. D., & Munoz, D. P. (2014). Overt responses during covert orienting. *Neuron, 82*(6), 1230-1243. doi:10.1016/j.neuron.2014.05.040

Cunitz, R. J. (1970). *Relationship Between Slow Drift and Smooth Pursuit Eye Movements (Ph.D. Dissertation).* (Doctor of Philosophy), University of Maryland,

Ditchburn, R. W., & Ginsborg, B. L. (1953). Involuntary eye movements during fixation. *J Physiol, 119*(1), 1-17.

Drewes, J., Zhu, W., Hu, Y., & Hu, X. (2014). Smaller is better: drift in gaze measurements due to pupil dynamics. *PLoS One, 9*(10), e111197. doi:10.1371/journal.pone.0111197

Edelman, J. A., & Xu, K. Z. (2009). Inhibition of voluntary saccadic eye movement commands by abrupt visual onsets. *J Neurophysiol, 101*(3), 1222-1234. doi:10.1152/jn.90708.2008

Engbert, R. (2006). Microsaccades: A microcosm for research on oculomotor control, attention, and visual perception. *Prog Brain Res, 154*, 177-192. doi:S0079-6123(06)54009-9 [pii]
10.1016/S0079-6123(06)54009-9

Engbert, R., & Kliegl, R. (2003). Microsaccades uncover the orientation of covert attention. *Vision Res, 43*(9), 1035-1045. doi:S0042698903000841 [pii]

Engbert, R., & Kliegl, R. (2004). Microsaccades keep the eyes' balance during fixation. *Psychol Sci, 15*(6), 431-436. doi:10.1111/j.0956-7976.2004.00697.x
PSCI697 [pii]

641 Engbert, R., Mergenthaler, K., Sinn, P., & Pikovsky, A. (2011). An integrated model
642   of fixational eye movements and microsaccades. *Proceedings of the National*
643   *Academy of Sciences of the United States of America, 108*(39), E765-770.
644   doi:10.1073/pnas.1102730108

645 Epelboim, J., & Kowler, E. (1993). Slow control with eccentric targets: evidence
646   against a position-corrective model. *Vision Res, 33*(3), 361-380.
647   doi:10.1016/0042-6989(93)90092-b

648 Everling, S., Pare, M., Dorris, M. C., & Munoz, D. P. (1998). Comparison of the
649   discharge characteristics of brain stem omnipause neurons and superior
650   colliculus fixation neurons in monkey: implications for control of fixation and
651   saccade behavior. *J Neurophysiol, 79*(2), 511-528.

652 Evinger, C., Kaneko, C. R., & Fuchs, A. F. (1982). Activity of omnipause neurons in
653   alert cats during saccadic eye movements and visual stimuli. *J Neurophysiol,*
654   *47*(5), 827-844. doi:10.1152/jn.1982.47.5.827

655 Fuchs, A. F., & Robinson, D. A. (1966). A method for measuring horizontal and
656   vertical eye movement chronically in the monkey. *Journal of applied*
657   *physiology, 21*(3), 1068-1070.

658 Goffart, L., Quinet, J., Chavane, F., & Masson, G. S. (2006). Influence of background
659   illumination on fixation and visually guided saccades in the rhesus monkey.
660   *Vision Res, 46*(1-2), 149-162. doi:10.1016/j.visres.2005.07.026

661 Grujic, N., Brehm, N., Gloge, C., Zhuo, W., & Hafed, Z. M. (2018). Peri-saccadic
662   perceptual mislocalization is different for upward saccades. *J Neurophysiol*.
663   doi:10.1152/jn.00350.2018

664 Hafed, Z. M. (2011). Mechanisms for generating and compensating for the smallest
665   possible saccades. *Eur J Neurosci, 33*(11), 2101-2113. doi:10.1111/j.1460-
666   9568.2011.07694.x

667 Hafed, Z. M., Chen, C.-Y., & Tian, X. (2015). Vision, perception, and attention
668   through the lens of microsaccades: mechanisms and implications. *Front Syst*
669   *Neurosci, 9*, 167. doi:10.3389/fnsys.2015.00167

670 Hafed, Z. M., & Chen, C. Y. (2016). Sharper, Stronger, Faster Upper Visual Field
671   Representation in Primate Superior Colliculus. *Curr Biol, 26*(13), 1647-1658.
672   doi:10.1016/j.cub.2016.04.059

673 Hafed, Z. M., & Clark, J. J. (2002). Microsaccades as an overt measure of covert
674   attention shifts. *Vision Res, 42*(22), 2533-2545. doi:S0042698902002638 [pii]

675 Hafed, Z. M., & Goffart, L. (2020). Gaze direction as equilibrium: more evidence from
676   spatial and temporal aspects of small-saccade triggering in the rhesus
677   macaque monkey. *J Neurophysiol, 123*(1), 308-322.
678   doi:10.1152/jn.00588.2019

679 Hafed, Z. M., Goffart, L., & Krauzlis, R. J. (2008). Superior colliculus inactivation
680   causes stable offsets in eye position during tracking. *J Neurosci, 28*(32), 8124-
681   8137. doi:28/32/8124 [pii]

682    10.1523/JNEUROSCI.1317-08.2008

683    Hafed, Z. M., & Ignashchenkova, A. (2013). On the dissociation between
684           microsaccade rate and direction after peripheral cues: microsaccadic inhibition
685           revisited. *J Neurosci, 33*(41), 16220-16235. doi:10.1523/JNEUROSCI.2240-
686           13.2013

687    Hafed, Z. M., Lovejoy, L. P., & Krauzlis, R. J. (2011). Modulation of microsaccades in
688           monkey during a covert visual attention task. *Journal of Neuroscience, 31*(43),
689           15219-15230. doi:10.1523/JNEUROSCI.3106-11.2011

690    Herrmann, C. J. J., Metzler, R., & Engbert, R. (2017). A self-avoiding walk with neural
691           delays as a model of fixational eye movements. *Sci Rep, 7*(1), 12958.
692           doi:10.1038/s41598-017-13489-8

693    Hooge, I. T. C., Hessels, R. S., & Nystrom, M. (2019). Do pupil-based binocular video
694           eye trackers reliably measure vergence? *Vision Res, 156*, 1-9.
695           doi:10.1016/j.visres.2019.01.004

696    Intoy, J., & Rucci, M. (2020). Finely tuned eye movements enhance visual acuity.
697           *Nature communications, 11*(1), 795. doi:10.1038/s41467-020-14616-2

698    Judge, S. J., Richmond, B. J., & Chu, F. C. (1980). Implantation of magnetic search
699           coils for measurement of eye position: an improved method. *Vision Res, 20*(6),
700           535-538.

701    Kimmel, D. L., Mammo, D., & Newsome, W. T. (2012). Tracking the eye non-
702           invasively: simultaneous comparison of the scleral search coil and optical
703           tracking techniques in the macaque monkey. *Frontiers in behavioral*
704           *neuroscience, 6*, 49. doi:10.3389/fnbeh.2012.00049

705    Ko, H. K., Poletti, M., & Rucci, M. (2010). Microsaccades precisely relocate gaze in a
706           high visual acuity task. *Nat Neurosci, 13*(12), 1549-1553. doi:nn.2663 [pii]
707    10.1038/nn.2663

708    Krauzlis, R. J., Goffart, L., & Hafed, Z. M. (2017). Neuronal control of fixation and
709           fixational eye movements. *Philos Trans R Soc Lond B Biol Sci, 372*(1718).
710           doi:10.1098/rstb.2016.0205

711    Krauzlis, R. J., & Miles, F. A. (1996). Release of fixation for pursuit and saccades in
712           humans: evidence for shared inputs acting on different neural substrates. *J*
713           *Neurophysiol, 76*(5), 2822-2833.

714    Kuang, X., Poletti, M., Victor, J. D., & Rucci, M. (2012). Temporal encoding of spatial
715           information during active visual fixation. *Current biology : CB, 22*(6), 510-514.
716           doi:10.1016/j.cub.2012.01.050

717    Malevich, T., Buonocore, A., & Hafed, Z. M. (2020). Dependence of the stimulus-
718           driven microsaccade rate signature on visual stimulus polarity. *bioRxiv*.
719           doi:10.1101/2020.05.23.112417

720    Martins, A. J., Kowler, E., & Palmer, C. (1985). Smooth pursuit of small-amplitude
721           sinusoidal motion. *J Opt Soc Am A, 2*(2), 234-242.

Missal, M., & Keller, E. L. (2002). Common inhibitory mechanism for saccades and
smooth-pursuit eye movements. *J Neurophysiol, 88*(4), 1880-1892.

Myers, G. A., & Stark, L. (1990). Topology of the near response triad. *Ophthalmic*
*Physiol Opt, 10*(2), 175-181. doi:10.1111/j.1475-1313.1990.tb00972.x

Nachmias, J. (1959). Two-dimensional motion of the retinal image during monocular
fixation. *J Opt Soc Am, 49*, 901-908.

Nachmias, J. (1961). Determiners of the drift of the eye during monocular fixation. *J*
*Opt Soc Am, 51*, 761-766.

Poletti, M., Aytekin, M., & Rucci, M. (2015). Head-Eye Coordination at a Microscopic
Scale. *Curr Biol, 25*(24), 3253-3259. doi:10.1016/j.cub.2015.11.004

Poletti, M., Listorti, C., & Rucci, M. (2010). Stability of the visual world during eye
drift. *The Journal of neuroscience : the official journal of the Society for*
*Neuroscience, 30*(33), 11143-11150. doi:10.1523/JNEUROSCI.1925-10.2010

Poletti, M., & Rucci, M. (2010). Eye movements under various conditions of image
fading. *J Vis, 10*(3), 6 1-18. doi:10.1167/10.3.6
/10/3/6/ [pii]

Poletti, M., & Rucci, M. (2015). A compact field guide to the study of microsaccades:
Challenges and functions. *Vision Res*. doi:10.1016/j.visres.2015.01.018

Pong, M., & Fuchs, A. F. (2000). Characteristics of the pupillary light reflex in the
macaque monkey: metrics. *J Neurophysiol, 84*(2), 953-963.
doi:10.1152/jn.2000.84.2.953

Ratliff, F., & Riggs, L. A. (1950). Involuntary motions of the eye during monocular
fixation. *J Exp Psychol, 40*(6), 687-701.

Reingold, E. M., & Stampe, D. M. (2002). Saccadic inhibition in voluntary and
reflexive saccades. *J Cogn Neurosci, 14*(3), 371-388.
doi:10.1162/089892902317361903

Rolfs, M. (2009). Microsaccades: small steps on a long way. *Vision Res, 49*(20),
2415-2441. doi:S0042-6989(09)00369-1 [pii]
10.1016/j.visres.2009.08.010

Rolfs, M., Kliegl, R., & Engbert, R. (2008). Toward a model of microsaccade
generation: the case of microsaccadic inhibition. *J Vis, 8*(11), 5 1-23.
doi:10.1167/8.11.5
/8/11/5/ [pii]

Rolls, E. T., & Cowey, A. (1970). Topography of the retina and striate cortex and its
relationship to visual acuity in rhesus monkeys and squirrel monkeys. *Exp*
*Brain Res, 10*(3), 298-310.

Rucci, M., Iovin, R., Poletti, M., & Santini, F. (2007). Miniature eye movements
enhance fine spatial detail. *Nature, 447*(7146), 851-854. doi:nature05866 [pii]
10.1038/nature05866

Rucci, M., & Victor, J. D. (2015). The unsteady eye: an information-processing stage,
not a bug. *Trends Neurosci*. doi:10.1016/j.tins.2015.01.005

Salinas, E., & Stanford, T. R. (2018). Saccadic inhibition interrupts ongoing
oculomotor activity to enable the rapid deployment of alternate movement
plans. *Sci Rep, 8*(1), 14163. doi:10.1038/s41598-018-32224-5

Schor, C. M., & Westall, C. (1984). Visual and vestibular sources of fixation instability
in amblyopia. *Invest Ophthalmol Vis Sci, 25*(6), 729-738.

Shelchkova, N., Tang, C., & Poletti, M. (2019). Task-driven visual exploration at the
foveal scale. *Proc Natl Acad Sci U S A, 116*(12), 5811-5818.
doi:10.1073/pnas.1812222116

Skavenski, A. A., Hansen, R. M., Steinman, R. M., & Winterson, B. J. (1979). Quality
of retinal image stabilization during small natural and artificial body rotations in
man. *Vision Res, 19*(6), 675-683. doi:0042-6989(79)90243-8 [pii]

Skinner, J., Buonocore, A., & Hafed, Z. M. (2019). Transfer function of the rhesus
macaque oculomotor system for small-amplitude slow motion trajectories. *J
Neurophysiol, 121*(2), 513-529. doi:10.1152/jn.00437.2018

Snodderly, D. M. (1987). Effects of light and dark environments on macaque and
human fixational eye movements. *Vision Res, 27*(3), 401-415. doi:0042-
6989(87)90089-7 [pii]

Snodderly, D. M. (2016). A physiological perspective on fixational eye movements.
*Vision Res, 118*, 31-47. doi:10.1016/j.visres.2014.12.006

St Cyr, G. J., & Fender, D. H. (1969). The interplay of drifts and flicks in binocular
fixation. *Vision Res, 9*(2), 245-265. doi:10.1016/0042-6989(69)90004-2

Steinman, R. M., & Collewijn, H. (1980). Binocular retinal image motion during active
head rotation. *Vision Res, 20*(5), 415-429. doi:10.1016/0042-6989(80)90032-2

Steinman, R. M., Haddad, G. M., Skavenski, A. A., & Wyman, D. (1973). Miniature
eye movement. *Science, 181*(102), 810-819.

Takahashi, M., & Shinoda, Y. (2018). Brain Stem Neural Circuits of Horizontal and
Vertical Saccade Systems and their Frame of Reference. *Neuroscience, 392*,
281-328. doi:10.1016/j.neuroscience.2018.08.027

Thaler, L., Schutz, A. C., Goodale, M. A., & Gegenfurtner, K. R. (2013). What is the
best fixation target? The effect of target shape on stability of fixational eye
movements. *Vision Res, 76*, 31-42. doi:10.1016/j.visres.2012.10.012

Tian, X., Yoshida, M., & Hafed, Z. M. (2016). A Microsaccadic Account of Attentional
Capture and Inhibition of Return in Posner Cueing. *Front Syst Neurosci, 10*,
23. doi:10.3389/fnsys.2016.00023

Tian, X., Yoshida, M., & Hafed, Z. M. (2018). Dynamics of fixational eye position and
microsaccades during spatial cueing: the case of express microsaccades. *J
Neurophysiol, 119*(5), 1962-1980. doi:10.1152/jn.00752.2017

Wikler, K. C., Williams, R. W., & Rakic, P. (1990). Photoreceptor mosaic: number and
distribution of rods and cones in the rhesus monkey retina. *J Comp Neurol,*
*297*(4), 499-508. doi:10.1002/cne.902970404

Wildenmann, U., & Schaeffel, F. (2013). Variations of pupil centration and their
effects on video eye tracking. *Ophthalmic Physiol Opt, 33*(6), 634-641.
doi:10.1111/opo.12086

Willeke, K. F., Tian, X., Buonocore, A., Bellet, J., Ramirez-Cardenas, A., & Hafed, Z.
M. (2019). Memory-guided microsaccades. *Nature communications, 10*(1),
3710. doi:10.1038/s41467-019-11711-x

Wyatt, H. J. (2010). The human pupil and the use of video-based eyetrackers. *Vision*
*research, 50*(19), 1982-1988. doi:10.1016/j.visres.2010.07.008

Zuber, B. L., Stark, L., & Cook, G. (1965). Microsaccades and the velocity-amplitude
relationship for saccadic eye movements. *Science, 150*(702), 1459-1460.
813

**Methods**

*Ethics approvals*

We tracked eye movements in three male rhesus macaque monkeys trained on behavioral eye movement tasks under head-stabilized conditions. The experiments were part of a larger neurophysiological investigation in the laboratory. All procedures and behavioral paradigms were approved (CIN3/13 and CIN4/19G) by ethics committees at the Regierungspräsidium Tübingen, and they complied with European Union directives on animal research.

*Animal preparation and laboratory setup*

Monkey N was 12 years old at the time of the experiments; monkeys A and M were 6-9 years old. Each monkey was initially implanted with a titanium head holder under aseptic surgical conditions, as described earlier (Chen & Hafed, 2013). In a subsequent procedure, each monkey was additionally implanted with a scleral search coil in one eye (left for monkey A and right for monkeys N and M), in order to allow precise eye tracking using the magnetic induction technique (Fuchs & Robinson, 1966; Judge et al., 1980). For some experiments in monkey M (e.g. Fig. 2), we also tracked eye movements binocularly. In this case, we performed an additional procedure to implant a search coil in this monkey's other eye. Eye coil implant procedures in our laboratory were described in detail earlier (Chen & Hafed, 2013).

The laboratory setup consisted of a cathode ray tube (CRT) display placed in front of the monkey in an otherwise dim room. The display subtended approximately +/- 15 deg horizontally and +/- 11 deg vertically when the monkey fixated at its center, and it had a 120 Hz refresh rate. The monkey's head was surrounded by a cube generating

electromagnetic fields for eye tracking (Remmel Labs). When we performed

pupillometry (e.g. Fig. 1 – figure supplement 2), we placed a video-based eye

tracking camera (EyeLink 1000, desktop mount; SR-Research, Ontario, Canada) in

front of the monkey under the CRT display. The eye tracking software output eye

positions and pupil diameters (of the right eye) for synchronization with our existing

real-time experiment control system, described in (Chen & Hafed, 2013; Tian et al.,

2016).

All experiments involved initially fixating a white fixation spot over a gray background.

The luminance of the gray background was 29.7 cd/m$^2$. The luminance of the white

fixation spot was 86 cd/m$^2$. When we presented black stimuli, the luminance was <

0.02 cd/m$^2$. For all other stimuli (e.g. spatial frequencies), the details of the stimuli are

provided below. Display resolution (~34-36 pixels/deg) was always higher than the

Nyquist limit associated with the highest spatial frequency stimulus that we tested,

and the display was linearized and calibrated before the experiments.

The monkeys were also all highly trained on fixation tasks that exercised their

abilities for accurate and precise fixation of a small target. For example, monkey N

previously contributed to gaze-contingent experiments in which forced foveal motor

errors of only a few minutes of arc could be corrected for by the monkey's eye

position (Tian et al., 2016, 2018), and monkeys M and A successfully used smooth

tracking of very slow motion trajectories, demonstrating that they could control their

smooth eye movement system at speeds similar to or even lower than the speeds of

ocular drift eye movements during fixation (Skinner et al., 2019).

*Behavioral tasks*

Experiment 1 was called the split view stimulus paradigm. The monkey fixated a small white spot (approximately 5 x 5 min arc) presented over a gray background (Chen & Hafed, 2013). After an initial fixation interval (approximately 500-1400 ms), a stimulus onset occurred, which had the following properties. The stimulus onset was marked by turning one entire half of the display (right or left, randomly picked across trials) to black until the end of the trial (approximately 500-2000 ms after stimulus onset). The vertical edge associated with such a split view stimulus was gaze-contingently (Chen & Hafed, 2013; Tian et al., 2016, 2018) updated, such that the edge between the light and dark regions of the retinal image was experimentally stabilized in real time relative to the fovea. This gaze-contingent image manipulation was relevant for other aspects of the task that are beyond the scope of the current analyses, especially because the current study focuses on time periods that are too short-lived to be affected by whether we used retinal image stabilization of the split view stimulus or not. To maintain visibility of the fixation spot despite the half-display of darkness that went right down its middle, we additionally enforced a kind of foveal sparing of the split view stimulus. Specifically, for a circle of radius 0.74 deg centered on instantaneous gaze position, we maintained background luminance even for the half of the circle that resided in the darkened half of the display (i.e. the circle was not split into a dark and bright half like the rest of the split view display). The fixation spot was always visible and always stable in the display (i.e. did not move gaze-contingently). We also interleaved control trials in this experiment that had identical timing; however, in this case, there was never a stimulus onset on the display (i.e. the monkey simply fixated steadily until trial end with no visual transients at all). We analyzed 5193, 2156, and 3860 trials in this experiment from monkeys M, A, and N,

respectively. Typically, we collected 150-330 trials per session and repeated the

experiment across multiple sessions.

Experiment 2 was called the black fixation flash experiment. Each monkey fixated a

white spot over a gray background like in Experiment 1. After an initial fixation

interval (approximately 550-1800 ms after initiating fixation), the entire display turned

black for one single display frame (~8 ms duration given our 120 Hz refresh rate)

before returning to its original state. Therefore, for one display frame, the fixation spot

was rendered momentarily invisible. However, such a short duration of fixation spot

occlusion is not expected to affect the eye movements that we analyzed (e.g. the drift

response that we observed in this experiment was similar to that in Experiment 1 with

a continuously visible fixation spot; see Results). In other control conditions in this

experiment, we interleaved trials in which either no flash occurred at all, or in which

the single frame flash was now a flash of a localized square of 1 x 1 deg dimensions

centered at 2.1 deg horizontally either to the right or left of the fixation spot. The

flashed square was also black, and the fixation spot was still visible in this case

(because the flash did not occlude it). This experiment was similar to one we used

recently under smooth pursuit conditions (Buonocore et al., 2019). We analyzed

1671 and 1837 trials from monkeys M and A, respectively, in this experiment.

In our third experiment, we replicated Experiment 2 but with white rather than black

flashes (the white fixation flash paradigm). We analyzed 5191 and 3112 trials from

monkeys M and A, respectively, in this experiment.

Finally, Experiments 4 and 5 tested spatial frequency and orientation tuning,

respectively, of the ocular position drift response in monkey M. We repeated the

fixation flash paradigms described above. However, instead of full-screen black or

white flashes, we presented a full-screen grating of a given stimulus property for

approximately 260-700 ms. Across trials, we varied the spatial frequency

(Experiment 4) or the orientation (Experiment 5) of the presented grating. There were

no control or localized flash trials in these new experiments. Instead, we exploited the

available numbers of trials to collect by varying the stimulus properties being tested

across trials. For spatial frequency tuning, we tested 5 different spatial frequencies

across trials (0.55, 1.1, 2.2, 4.2, and 6.8 cycles/deg; cpd). For orientation tuning, we

tested 4 different orientations (vertical, 45 deg clockwise from vertical, horizontal, and

45 deg counterclockwise from vertical). Grating contrast was always maximal

(100%). Also, across trials, we picked from two different grating phases (one in which

the grating was white at the center of the display, near the fixation spot, and one in

which the grating was black at the center of the display; i.e. a phase difference of $\pi$

from the other condition). We analyzed 9967 trials from the spatial frequency

manipulation and 9856 trials from the orientation manipulation.

To test for the relationship between the ocular position drift response and pupillary

constrictions associated with stimulus onset, we measured pupil diameter in monkey

A (Experiment 6). We ran the monkey on a variant of the fixation flash tasks that we

knew would evoke a drift response (i.e. a full-screen flash of a visual pattern

consisting of a high-contrast vertical grating). We confirmed such a drift response by

measuring eye position using a scleral search coil. Simultaneously, we analyzed

pupil diameter using a video-based eye tracker, and we analyzed variations in pupil

diameter as a function of time from stimulus onset. This allowed us to relate the

pupillary constriction response after stimulus onset to the ocular position drift

response that we report in this study. Note that the video-based eye tracker was not

suitable to detect the drift response in the same eye for which we measured the

pupil. That is why we tracked the other eye with a scleral search coil. That is, we

exploited our observation that the drift response is a binocular phenomenon (Fig. 2).

We could have measured the pupil diameter in the coil eye, but our initial intent was

to measure the drift response binocularly (with a coil in one eye and a video-based

eye tracker for the other). However, it quickly became clear that the video-based eye

tracker was simply not suitable for properly measuring the drift response, consistent

with prior observations in the literature (Choe et al., 2016; Drewes et al., 2014;

Hooge et al., 2019; Kimmel et al., 2012; Wyatt, 2010). We analyzed 6203 trials from

this comparison.

To investigate whether the drift response was associated with a simultaneous head

movement, we also repeated Experiment 2 (with only the full-screen flash condition)

on monkey M while simultaneously recording eye and head position. Eye position

recording (left eye) was identical to the procedures described above. For head

position measurements, we placed a search coil (identical to the coils that we

implanted in the monkeys' eyes) at one of three possible positions on the head (e.g.

attached to a recording chamber implanted on the head or attached to the titanium

head holder screwed on the monkey's skull, and so on). The head coil had lower

resistance (40 ohms) than the eye coils (typically around 90 ohms), resulting in an

even more sensitive measure of potential head movements. We used three positions

of head coils, across three different sets of experiments, to make sure that we could

catch all possible axes of head motion that could occur in our experiments.

Specifically, the magnetic fields that were generated around the monkey's head were

optimized for measuring horizontal and vertical eye position (i.e. with a nominal

orientation of the eye coil within the magnetic fields). However, with head coil

placement, positions of recording chambers on the head or the titanium head holder

screwed on the skull resulted in non-standard orientations of the head coil relative to

the magnetic fields. This meant that some axes of potential head movements could

be missed with only one inopportunely chosen head coil position. As a result, we

varied head coil position on the head, across three different experiments, in order to

be absolutely sure that our drift response could not be explained by a reflexive head

movement after stimulus onset. We digitized the head coil output using the same

amplification and acquisition system as our eye coil outputs.

To obtain an approximate calibration of head coil measurements, we brought the

monkey chair to the laboratory without a monkey, and we then placed the head coil

at one position relative to the head holder (which is always at a fixed position relative

to both the chair and the monkey's head). We then measured calibration output by

physically moving the head coil by 3 mm in different directions from its standard

position. At each position, we recorded 5000 ms of data, which we then averaged.

This gave the approximate calibration values in Fig. 2 – figure supplement 1.

*Data analyses*

We detected saccades and microsaccades using established methods in our

laboratory (Bellet, Bellet, Nienborg, Hafed, & Berens, 2019; Chen & Hafed, 2013).

We also manually inspected all trials to ensure correct detection, and to also remove

blinks.

To analyze our drift responses, we considered only trials in which there were no

saccadic or blink events within an interval starting from -100 ms relative to stimulus

onset and ending at +200 ms relative to stimulus onset. In additional control

analyses, we extended this so-called saccade-free interval forward in time in steps of

50 ms until +350 ms after stimulus onset. That is, in the longest analysis interval that

we considered, there were no saccadic or blink events between -100 ms and 350 ms

from stimulus onset.

1002

To absolutely ensure that we excluded any possibility of saccadic eye speeds (for

even the smallest microsaccades) that might bias our measurements of average eye

position across trials, we added one extra step after saccade detection, just as a

sanity check: we inspected all accepted trials based on saccade detection and the

above interval, and we ensured that eye velocity profiles did not have any clear

outliers above the levels expected from ocular position drifts and measurement noise

for a given session and monkey. This ensured that even rare electronic artifacts

(lasting for only 1 ms or so) in scleral search coil measurements were not corrupting

our ocular position drift interpretations.

1012

We summarized the drift response properties by averaging eye position or velocity

traces for a given stimulus type. When averaging eye positions, we aligned traces

horizontally on stimulus onset time and vertically on the eye position that existed at

this time. This way, we could observe the drift responses regardless of the absolute

eye position that existed at trial onset. In other analyses (e.g. Fig. 2 – figure

supplement 3), we explicitly considered the effect of absolute eye position.

Specifically, in each session, we estimated the trial-to-trial variance of eye position

(due to fixational eye movements) at the time of stimulus onset (we averaged the

final 50 ms of eye position data right before stimulus onset in each trial, and the

across-trial distribution provided an estimate of fixation variability). Then, we

considered only trials above the median vertical eye position across the population or

only trials below the median vertical eye position. We then pooled the above-median

and below-median populations across sessions to increase the numbers of

observations (and, therefore, the statistical confidence of our results). For

presentation in Fig. 2 – figure supplement 3 of these median split data, we still

aligned the eye positions at the time of stimulus onset as we did for the other figures

(see above). This allowed us to directly demonstrate that the drift response was the

same in timing and amplitude whether the starting eye position was below or above

median eye position.

1032

For analyzing microsaccades, we estimated microsaccade rate as a function of time

from stimulus onset using similar methods to our earlier work (Buonocore, Chen, et

al., 2017; Hafed et al., 2011). Briefly, we used a running window of width 80 ms,

during which we estimated rate as the average number of movements observed per

time window (and normalized to a unit of movements per second) (Hafed et al.,

2011). We then stepped the window forward in time, in steps of 5 ms, starting from -

300 ms and ending at +600 ms relative to stimulus onset.

1040

We also checked whether the drift response happened when a nearby microsaccade

still occurred (e.g. Fig. 4G-I). To do this, we binned trials based on the timing of

microsaccades. For example, we looked for trials in which there was a microsaccade

(or more) occurring up to a time of, say, +50 ms after stimulus onset. Then, we

plotted saccade-free eye velocity after this time to look for the drift response. We

used a similar procedure for other times earlier than +50 ms. For microsaccades after

the drift response, we did the same. For example, we checked if there was a

microsaccade (or more) starting from a time of, say, +225 ms after stimulus onset

(i.e. after the ocular drift response). In this case, the drift response occurred earlier,

so we plotted saccade-free eye velocity up to the time of, say, +225 ms.

1051

To further relate microsaccades to the ocular drift response, we investigated whether

the first microsaccade after the drift response was biased downwards in direction

(e.g. to compensate for the predominantly upward drift response). We used the

following procedure. In all of our saccade-free trials in the standard interval of -100

ms to +200 ms relative to stimulus onset, we searched for the first microsaccade to

occur after this interval. We then plotted the angular distribution of this first

microsaccade and related it to the control data without any stimulus onsets, but with

the same analysis workflow of saccade-free fixation between -100 ms and +200 ms

(Fig. 2 – figure supplement 4).

1061

For pupil diameter measurements, our data acquisition system output analog

voltages proportional to pupil diameter values calculated by the video-based eye

tracker. We stored digitized measurements of these voltages in synchrony with our

stimulus events, and that is why we reported these measurements using arbitrary

units in Fig. 1 – figure supplement 2. We used the average pupil diameter in the final

100 ms before stimulus onset in a given trial as the baseline measurement. We

subtracted this baseline measurement from all pupil measurements within the trial.

Then, we averaged across trials. This allowed us to display the relative change in

pupil diameter after stimulus onset (again, in arbitrary units). We ensured that we

analyzed saccade-free epochs of pupil diameter, like we analyzed the ocular position

drift responses in all other experiments. We measured saccades in both the video-

based eye tracker signal as well as the other eye, which was tracked with a scleral

search coil. This way, our analyses of pupil diameter dynamics after stimulus onset

were comparable to the analyses of saccade-free ocular position drift responses that

we report in this study. For the linear fits in Fig. 1 – figure supplement 2, we used a

piece-wise linear model to estimate the onset of the smooth ocular drift response or

the onset of the pupil constriction. This approach is the same as that used for smooth

pursuit eye movements earlier (Krauzlis & Miles, 1996).

1080

Because we were dealing with very subtle eye position effects even in no-stimulus

controls (e.g. Fig. 2A), we first confirmed that potential electronic drifts for our scleral

search coil system cannot explain our results, and we did this before performing any

of the analyses above. Specifically, sometimes, scleral search coil systems might

exhibit a subtle drift in eye position calibration during the course of a given session

(i.e. across tens of minutes or even hours). To confirm that such a potential technical

artifact was too slow (if it existed at all) to account for even the baseline no-stimulus

drift pattern of a given monkey in our data (e.g. Fig. 2A), we compared such baseline

drift speed (which we deemed physiological) to any potential electronic drifts in eye

tracker calibration across trials within a session. To do this, we measured average

eye position in the final 30 ms before stimulus onset (including sham stimulus onset

in control conditions). We then plotted this measurement as a function of trial

number. Given the trial durations, this allowed us to estimate an "electronic drift

speed" measure. When it existed, this electronic drift speed was at least an order of

magnitude slower than baseline no-stimulus drift speeds of our monkeys. We are,

therefore, confident that our results are physiological drifts of eye position, and not

measurement artifacts.

1098

Finally, for the simultaneous eye and head position measurements, we analyzed eye

position and velocity as described above, including our measures to ensure saccade-

free epochs for analyzing the drift response. For head position, we obtained the

output of each channel of the head coil, and we also obtained a differential of it using

the same smoothing differentiation filter that we used to obtain eye velocity. We

visualized the position measurements in Fig. 2 – figure supplement 1 to confirm that

the drift response was not accompanied by a head response (the head data in the

figure are very similar in variance to the measurements obtained during head coil

calibration with no monkey at all; i.e. at the level of electronic noise of the system).

Head movement data were analyzed from the very same trials that we included for

the eye movement analyses.

1110

*Statistics*

We used descriptive statistics throughout the study. All figures show and define error

bars. We also did not show any average curves having fewer than 10 repetitions,

because such few repetitions were deemed unreliable; this is the reason for the

missing two curves in Fig. 4G in relation to the similar panels of Fig. 4H, I (saccades

were simply too rare due to the saccadic inhibition shown in Fig. 4A). Typically, we

had many more numbers of repetitions than 10, as shown in the insets of the figures.

1118

To estimate temporal parameters such as the onset time of an ocular drift response,

we compared measurements on trials with a stimulus onset to corresponding control

trials without any stimulus onset. For each of the stimulus or control populations of

repetitions, we estimated the 95% confidence intervals surrounding the mean

measurement. We then marked all time points in which such 95% confidence

intervals of the two compared curves did not overlap. This was done using pink

horizontal lines on all relevant x-axes in all figures. Due to noise (particularly in

differentiated eye velocity measurements), there could be individual milliseconds or

pairs of milliseconds in which there was no overlap between the 95% confidence

intervals. We still displayed these time points in all figures. However, we consider a

significant difference as one in which the 95% confidence intervals do not overlap for

at least 20 consecutive milliseconds.

1131

We used the above procedures on either eye position or eye velocity independently,

and we include results in all figures for both eye position and eye velocity curves.

Because eye velocity computations necessarily involve some smoothing operation to

regularize the noise associated with numerical differentiation, they introduce some

temporal blurring of signals (both backwards and forwards in time). On the other

hand, lack of overlap in the 95% confidence intervals of position traces would be too

conservative, because it would estimate a later onset of the drift response (velocity is

more sensitive than position for detecting onset of smooth eye movements). We

therefore elected to show in the figures all statistical comparisons independently for

both eye position and eye velocity traces, for the sake of completeness. In this

regard, the position results can be considered to be conservative estimates of drift

response onset time, whereas the velocity results can be considered to be slightly

more liberal estimates. Numbers (e.g. p-values) reported in the text were based on

eye velocity statistics, because eye velocity also allowed estimating the duration of

the drift response (the end of which being when the eye velocity curve returned back

to baseline with overlapping 95% confidence intervals).

1148

In some cases, we additionally statistically compared drift responses across

conditions (e.g. for right versus left eye). To do this, we defined a measurement

interval based on the timing of the drift response that we observed (50-140 ms in

black or white fixation flash paradigms and 80-140 ms in the spatial frequency and

orientation paradigms). During this interval, we searched for peak eye velocity and

characterized its magnitude and latency. When comparing two measurements (e.g.

right versus left eye peak velocity latency), we used non-parametric Wilcoxon rank

sum tests. When comparing multiple conditions (e.g. 5 different spatial frequencies),

we used non-parametric Kruskal-Wallis tests. We used non-parametric tests because

our measurements were not normally distributed. Finally, when we reported "adjusted

p-values" in Results, we were reporting corrections for multiple comparisons using

Bonferroni methods.

1161

For confirming a lack of head movement around the time of the drift response (Fig. 2

– figure supplement 1), we obtained a paired measurement of eye or head velocity

from each trial of a given experiment. The first measurement of the pair was average

eye or head velocity in the interval (-100-0 ms from stimulus onset), and it served as

a baseline velocity. The second measurement of the pair was average eye or head

velocity in the interval (50-150 ms from stimulus onset), chosen to coincide with the

time of the expected drift response. For a given head coil position, we then

performed a Wilcoxon rank sum test on vertical eye velocity across both intervals. If

there was a drift response, then the vertical eye velocity in the second interval was

expected to be statistically significantly different from the vertical eye velocity in the

second interval (Fig. 2 – figure supplement 1A). We then performed the Wilcoxon

rank sum test on each channel of head velocity (Fig. 2 – figure supplement 1B, C). If

the drift response was accompanied by a head movement, then there should have

been a statistically significant change in head velocity (in either channel or both) after

stimulus onset when compared to before stimulus onset. Note that we could have

also performed the statistics directly on the raw position measurements, and we

would have reached similar conclusions (Fig. 2 – figure supplement 1). However,

since the monkey had a slight baseline drift even without any stimulus onset,

statistics on eye position before and after stimulus onset would have been significant

(for eye position) even if there was no drift response at all. We therefore used

velocity because it effectively detrends the eye position signal. We also analyzed

each channel of head coil output individually because head coil orientation was not

always perfectly orthogonal to the inducing magnetic field in the laboratory (see

above).

1186

1187

**Supplemental Figures**

 **Figure 1 – figure supplement 1 Short-latency, stimulus-driven ocular position drift response in**
 **two additional monkeys besides the monkey shown in Fig. 1.** This figure is formatted identically to
 Fig. 1. It shows very similar results from two additional monkeys, M and N. **(A-F)** Monkey M results. **(G-**
 **L)** Monkey N results. Error bars are defined in all relevant panels. The pink horizontal lines on various
 x-axes denote intervals in which the 95% confidence intervals around the two compared curves did not
 overlap.

1200

**Figure 1 – figure supplement 2 The ocular position drift response occurred earlier than pupil diameter constriction occurring after stimulus onset.** We measured pupil diameter in monkey A when a stimulus transient (high-contrast, full-screen vertical grating) was presented. The blue curve shows the ocular position drift response, replicating the results of Fig. 1, Fig. 1 – figure supplement 1 in new experiments. This response was measured with a scleral search coil in the left eye. The gray curve shows pupil diameter from the same set of trials, measured with a video-based eye tracker aimed at the right eye (the ocular position drift response was unresolvable in the right eye with the video-based eye tracker; Methods). To highlight the change in pupil diameter after stimulus onset within each trial, we averaged pupil diameter in the final 100 ms before stimulus onset, and we then subtracted this average from all pupil measurements within the trial. This aligned the pupil measurements across all trials in the y-axis dimension, isolating the relative changes in pupil diameter occurring after stimulus onset (Methods). This is a similar approach to our procedures for ocular position drift responses (e.g. Fig. 1 and the blue curve in the current figure). As can be seen, the inflection point of pupil diameter (indicated by the gray dashed vertical line at 145 ms), demonstrating stimulus-driven pupillary constriction, came significantly later than the onset of the ocular position drift response (indicated by the light blue dashed vertical line at 98 ms). Each inflection point was obtained by fitting a piece-wise linear model to each curve: a linear fit in an early interval before the onset of the ocular position drift response or pupil constriction, and a second linear fit in a later interval after the onset of the ocular position drift response or pupil constriction. This is similar to methods used earlier for estimating smooth pursuit onset latencies (Krauzlis & Miles, 1996). The two linear fits per curve are shown in red. Error bars denote 95% confidence intervals across trials.

1224

**Figure 2 – figure supplement 1 The drift response is independent of head movements. (A)** Each row shows vertical eye position in a format identical to that shown in earlier figures (e.g. Figs. 1, 2). The same drift response was replicated in these three new sets of experiments of the black flash condition. **(B, C)** Simultaneous measurements of head position for the same sets of trials shown in **A**. In each row, we placed a head coil at a different position on the monkey's head to explore different axes of potential head movements (Methods). Each coil provided two measurements (**B**, **C**) of induced electrical current from two orthogonal magnetic fields used to measure position (these were the same two fields used to induce electrical current in the eye coil to allow measuring horizontal and vertical eye position). There was no head movement concomitant with the drift response in any of the channels. The pink intervals illustrate the measurement intervals that we used for statistical assessment of whether there was a response after stimulus onset relative to before (see text); the calibration values (in mm) were obtained using an approximate calibration of head position measurement with one of the head coil positions (Methods). Error bars denote 95% confidence intervals.

1238

1239

**Figure 2 – figure supplement 2 Small localized stimulus onsets had minimal effects on ocular position drifts when compared to larger flashes. (A, B)** Horizontal (top row in each panel) and vertical (bottom row in each panel) eye position in each of two monkeys (A and M) under different conditions. The left column in each panel shows data when small localized flashes were presented in the black fixation flash paradigm (Methods; similar results were obtained with the white fixation flash paradigm). The right column shows the full-screen flash results of Fig. 2 for comparison. All other conventions are like in Fig. 2. As can be seen, the upward drift response with full-screen flashes was much bigger than any responses to small localized stimuli. Consistent with this, there was no single contiguous interval longer than 8 ms in duration for which horizontal or vertical eye position differed between whether a localized flash was presented to the right or to the left of fixation (pink intervals on the x-axis of the bottom left panel of **B**, denoting when the 95% confidence intervals did not overlap with each other between the two compared curves). **(C, D)** Eye velocity measurements for the same data as in **A**, **B**. There were no time intervals during which the right and left flash eye velocity curves differed from each other (that is, there were no pink intervals on x-axes denoting a lack of overlap between 95% confidence intervals). We also confirmed that in the interval of 50-140 ms after stimulus onset, peak vertical eye velocity was not different between rightward and leftward localized flashes (p=0.64, Z=0.47 for monkey A and p=0.92, Z=-0.10 for monkey M; Wilcoxon rank sum test). A similar conclusion was also reached for peak horizontal eye velocity (p=0.5, Z=-0.68 for monkey A and p=0.52, Z=-0.65 for monkey M). Error bars in all panels denote 95% confidence intervals.

1262

**Figure 2 – figure supplement 3 Independence of the ocular position drift response from starting eye position.** For each trial of the black fixation flash paradigm (full-screen flashes), we measured average eye position in the final 50 ms before stimulus onset. Across trials within a given session, this resulted in a population of starting eye positions. We split the trials within a session based on the median value of starting vertical eye position. We then measured the ocular position drift response for all sessions when only taking the "above" median group of the sessions (purple) and again when taking only the "below" median group of all sessions (bluish). **(A, B)** Position measurements in two monkeys. We vertically aligned traces (based on starting position at stimulus onset) in order to highlight the relative change in eye position associated with the ocular position drift response (similar to how we analyzed data in all other figures; e.g. Fig. 1). **(C, D)** Eye velocity measurements. In both eye position and eye velocity, the ocular drift response measurements were statistically very similar; there were no time intervals in which the 95% confidence intervals between the "above" and "below" median groups did not overlap with each other. We also confirmed that in the interval of 50-140 ms after stimulus onset, peak vertical eye velocity was not different between the two groups (p=0.63, Z=0.48 for monkey A and p=0.46, Z=0.74 for monkey M; Wilcoxon rank sum test). All other conventions are similar to our other analyses of the ocular position drift response (e.g. Fig. 1). Similar results were obtained with the split view stimulus and white fixation flash paradigms (e.g. p=0.32 [Z=-1.01], 0.62 [Z=-0.5] for monkeys A and M, respectively, in the white fixation flash paradigm, and p>0.15 [Z=-1.46, -0.82], p>0.2 [Z=-0.55, 1.27], and p>0.85 [Z=0.01, 0.19] for monkeys A, M, and N, respectively, in the split view stimulus paradigm; Z-values reported for right and left darkness for each monkey in the split view stimulus paradigm).

1284

**Figure 2 – figure supplement 4 The drift response was not systematically related to the direction of the first microsaccade to occur after it.** We searched for the first microsaccade to occur after our standard designated saccade-free fixation interval of -100 ms to 200 ms relative to full-screen flash onset. Across trials, we then plotted the direction distribution of these first microsaccades. Each row shows the stimulus onset condition (left column) and the corresponding control condition without any full-screen stimulus flash (right column). **(A, B)** Black fixation flash paradigm. **(C, D)** White fixation flash paradigm. Similar results were obtained with the split view stimulus paradigm. In all cases, each monkey showed a stereotypical distribution of microsaccade directions in the control condition without any stimulus onset (e.g. predominantly leftward in the monkey A control data; right column). This distribution was largely unaltered after the upward ocular position drift response, despite some idiosyncratic variations in the ranges of microsaccade directions that occurred. This is expected: given how small the ocular position drift response is (e.g. Fig. 1), its amplitude is much smaller than the amplitude of microsaccades, and it is therefore unlikely to trigger a corrective downward microsaccade.

1301

**Figure 4 – figure supplement 1 The ocular position drift response occurred during saccadic inhibition, and it also occurred even with nearby saccades.** This figure is formatted identically to Fig. 4. It shows similar results, but this time from the black fixation flash paradigm using full-screen flashes (the white fixation flash paradigm produced similar observations). **(A, B)** Microsaccade rate after full-screen flash onset in each monkey. The gray curve shows baseline microsaccade rate with no stimulus transient. **(C, D)** Ocular position drift response in each monkey during saccade-free fixation. To illustrate the timing of the drift response to saccadic inhibition, the dots indicate trial-by-trial rasters of microsaccade onset times during fixation. The drift response occurred when microsaccades were inhibited. **(E, F)** In each monkey, we inspected the ocular drift response when microsaccades still occurred near stimulus onset (different colors denote different microsaccade times as per the legends). The drift response still occurred, as in Fig. 4 with another paradigm and the inclusion of a third monkey. Error bars are defined in the respective panels. All other conventions are similar to Fig. 4.

1314