1   **Low-frequency neural tracking of speech amplitude envelope reflects the convolution of**

2   **evoked responses to acoustic edges, not oscillatory entrainment**

Katsuaki Kojima*[1,2,3], Yulia Oganian*[2], Chang Cai[3], Anne Findlay[3], Edward Chang[2]ⵥ, Srikantan Nagarajan[3]ⵥ

1. Department of Pediatrics, University of California, San Francisco, Box 0734, 550 16th Street, 5th Floor, San Francisco, CA 94158, USA

2. Department of Neurological Surgery, University of California, San Francisco, 675 Nelson Rising Lane, San Francisco, CA 94158, USA

3   3. Department of Radiology, University of California, San Francisco, 513 Parnassus Avenue, S362,

4   San Francisco, CA 94143-0628

*, ⵥ authors contributed equally

Corresponding authors:

Edward Chang: Edward.chang@ucsf.edu

Srikantan Nagarajan: Srikantan.Nagarajan@ucsf.edu

Abstract

The amplitude envelope of speech is crucial for accurate comprehension, and several studies have

shown that the phase of neural activity in the theta-delta bands (1 - 10 Hz) tracks the phase of the speech

amplitude envelope during listening, a process referred to as envelope tracking. However, the

mechanisms underlying envelope tracking have been heavily debated. A dominant model posits that

envelope tracking reflects continuous entrainment of endogenous low-frequency oscillations to the speech

envelope. However, it has proven challenging to distinguish this from the alternative model that envelope

tracking reflects a convolution of evoked responses to acoustic landmarks within the envelope. To address

this, we recorded magnetoencephalography while participants listened to natural and slowed speech. First,

we found that peaks in the rate of envelope change, acoustic edge landmarks, induced a strong evoked

response. We then contrasted central diverging predictions of evoked response and oscillatory models

regarding the spectral and temporal extent of neural phase locking in the theta-delta range at different

speech rates. Our analyses revealed transient theta phase locking at both speech rates and delta phase

locking in slow speech. This pattern was predicted by the evoked response model but not by oscillatory

entrainment. Furthermore, the encoding of acoustic edge magnitudes was invariant to contextual speech

rate, demonstrating speech rate normalization of acoustic edge representations. Taken together, our results

show that an evoked response model provides a better account of neural phase locking to the speech

envelope than oscillatory entrainment.

Introduction

Speech comprehension is essential to human communication. A major computational step in neural processing of speech is tracking of its amplitude envelope, the overall intensity of speech across spectral bands. The speech envelope is dominated by fluctuations in the range of ~ 1 – 10 Hz, which cue the syllabic structure of speech, and the removal of which from the speech signal severely impairs intelligibility (Drullman et al., 1994a, 1994b). A multitude of studies have shown a consistent relationship between the phase of band-limited low-frequency neural activity measured in M/EEG over auditory cortical areas and the phase of the amplitude envelope of speech, a phenomenon widely known as envelope tracking (Ahissar et al., 2001; Luo & Poeppel, 2007). Subsequent studies found that the strength of envelope tracking is correlated with speech intelligibility, suggesting that it could constitute an essential stage in speech comprehension (Abrams et al., 2008; Peelle et al., 2013). However, the neural computations underlying speech envelope tracking are controversially debated (Gwilliams, 2019; Obleser & Kayser, 2019; Zoefel, ten Oever, et al., 2018).

A dominant theory of speech envelope tracking is "oscillatory entrainment," which posits that envelope tracking reflects the neural entrainment (i.e., phase alignment) of endogenous neural oscillations in the delta-theta range (1-10 Hz) to fluctuations in the amplitude envelope of speech, with the primary entrained frequency possibly shifting dynamically to reflect the syllabic rate of the speech signal (Ding et al., 2015; Zoefel, 2018; Kösem et al., 2018). Functionally, oscillatory entrainment is thought to benefit speech processing due to its resonance properties (Haegens & Zion Golumbic, 2018; Helfrich et al., 2019), implementing automatic prediction of the timing of upcoming speech (e.g., the next syllable (Hovsepyan et al., 2020)). According to this theory, in natural speech, phase resetting of entrained oscillations is driven by discrete acoustic landmark events in the speech signal.

An alternative theory suggests that representations of the speech envelope reflect the convolution of evoked responses to discrete acoustic landmark events in the speech signal, as has been found previously in the visual domain (Breska & Deouell, 2017; Capilla et al., 2011). These acoustic landmarks could either be amplitude peaks, marking vowel midpoints (Ghitza, 2013), or acoustic edges (i.e., peaks in the rate of amplitude envelope change) that mark the perceived onset of vowels (Mermelstein, 1975). Recent intracranial evidence of acoustic edge encoding in speech cortical areas suggested the latter (Oganian & Chang, 2019), also supported by findings from electroencephalography (Brodbeck et al., 2018). However, to date, due to the rapid dynamics of natural speech, it has not been tested directly whether landmark events in the envelope drive an evoked response or entrain an oscillator (Ruhnau et al., 2020; Zoefel et al., 2019).

Here, we approach this problem by comparing neural responses to speech at a regular speech rate and equally intelligible speech slowed to 1/3 of its original rate, using magnetoencephalography (MEG)

recordings (Figure 1A). Slowing increased the time interval between consecutive landmark events in the speech envelope, as well as the time interval between different envelope landmarks and other (e.g., phonetic) content in the speech signal, allowing us to examine the effects of this shift on neural phase-locking patterns. Prior to examining the predictions of evoked response and oscillatory accounts of envelope tracking, we confirmed that our data exhibited phase locking between speech envelope and neural responses at the frequency of the speech rate in each condition, as reported previously (Peelle & Davis, 2012). Furthermore, we identified the phase-resetting envelope landmark, confirming it to be acoustic edges (Oganian & Chang, 2019).

Regarding the main model comparisons, an increase in neural phase locking following phase resets by acoustic landmarks is expected under both evoked response and oscillatory models. However, we hypothesized that the spectral and temporal extent of this phase locking, and its dependency on speech rate, might differ. To derive qualitative predictions, we examined the responses predicted by simple simulation models of an evoked response mechanism and two different oscillatory mechanisms, as previously proposed in the literature: one with a fixed entrainment frequency in the delta range and one with a flexible adjustment of the entrained frequency to the stimulus speech rate (Figure 1B). We then focused our analyses on patterns of phase locking that diverged between predictions of the three models.

Examining cortical dynamics in response to slow speech also allowed us to address a second question, namely the neural mechanisms of speech rate normalization. Speech rate normalization allows listeners to adjust perceptual processes, in particular envelope tracking, to differences in speech rate. It has previously been proposed that speech rate normalization is implemented through shifts in the frequency of the phase-locked oscillator to match the speech rate (Ahissar et al., 2001; Bosker & Kösem, 2017; Kösem et al., 2018; Nourski et al., 2009; Pefkou et al., 2017). Moreover, previous work with music stimuli suggested that evoked responses to edges might be reduced in slowed speech due to overall softer amplitude slopes (Doelling et al., 2019). In contrast, by comparing MEG responses to regular and slowed speech in this study, we demonstrate that the frequency of phase locking to edges does not shift with slowing but that the strength of phase resets at acoustic edges is readjusted to compensate for the overall softer amplitude slopes at slow speech rates.

Figure 1. Task design and envelope tracking in neural data A. The acoustic waveform of an example utterance ('Tarantino

says…"), with syllable boundaries, amplitude envelope, and first temporal derivative of the envelope superimposed on it. The sa[...]

utterance is shown at a regular rate (left) and slowed (right) speech rate. Arrows mark candidate temporal landmark that might in[...]

phase locking (Black: local peaks in the envelope, peakEnv; Purple: acoustic edges, defined as local peaks in the first temporal

derivative (rate of change) of the envelope, peakRate). **B.** Frequency of peakRate/peakEnv events. Dashed vertical lines mark the

average frequency of peakRate events in slow (blue, 1.9 Hz) and regular speech (green, 5.7 Hz). **C.** Single-subject (black) and gr[...]

average (red) comprehension performance. **D.** Sensor selection was based on M100 response to utterance onsets. Top: Group-

averaged evoked response across all 20 sensors included in the analysis. Error bars are ± 1 SEM across subjects. Bottom: Topogr[...]

map of a group-averaged M100 response with selected sensors marked in red. **E.** Group-averaged evoked response aligned to

peakRate and peakEnv events. Dotted lines mark clusters with $p < 0.05$ with a cluster-based permutation test against 0. Error bars

± 1 SEM across subjects. **F**. Cerebro-acoustic phase coherence (CAC) between MEG responses and speech envelope (upper pane[...]

and the difference between slow and regular speech (ΔCAC, lower panel). Data were filtered in semi-logarithmically spaced band[...]

between 0.2 and 10 Hz for this analysis. Dashed vertical lines mark the average frequency of peakRate events in each condition, [...]

shown in B. * $p < 0.001$ with 2-way repeated-measures ANOVA. Error bars are ± 1 SEM across subjects.

17   **Results**

Speech Envelope Tracking for regular and slow speech as seen in MEG

We recorded MEG while participants ($n = 12$) listened to ten minutes of continuous speech, once at the

original rate (regular speech condition) and once slowed to 1/3 of the original speech rate (Slow speech

condition, Figure 1A). Stimulus materials were selected from the Boston University Radio Speech Corpus,

BURSC (Ostendorf, Price, & Shattuck-Hufnagel, 1995); see Tables S1 and S2 for full stimulus transcript

and list of comprehension questions.

In a first step, we characterized the temporal dynamics of acoustic landmark events in our speech
stimulus (Figure S1), with a focus on peaks in the rate of envelope change (peakRate, purple in Figure 1A)
and peaks in the envelope (peakEnv, black in Figure 1A). At the regular speech rate, the average interval
between landmarks (similar for peakRate and peakEnv) was 175ms, corresponding to an average
landmark frequency in the theta band at 5.7 Hz (SD = 2.9 Hz, Figure 1B), in line with the typical power
spectrum of the amplitude envelope in natural speech (Ding et al., 2017). When slowed to 1/3 of the
original speech rate, the average frequency of landmarks was 1.9 Hz (SD = 1 Hz, similar for peakRate
and peakEnv), shifting the peak in the envelope power spectrum to the delta band. Slowing did not impair
participants' comprehension behavior, as indicated by responses to 3-4 multiple choice comprehension
questions after each story (chance: 50 %, regular speech: mean = 83 %, SD = 13 %; slow speech: mean =
90%, SD = 9.5 %; $t(11) = -1.85$, $p = 0.09$; Figure 1C). To focus our analyses on sensors that capture
auditory sensory processing, we selected ten sensors with the largest M100 response to speech onsets
after silence periods from each hemisphere (Figure 1D) for all further analyses. The M100 response
showed the typical dipole pattern in each hemisphere (Chait et al., 2004).
15
Acoustic edges drive MEG evoked responses
We first asked which landmark in the speech envelope drives evoked responses and phase locking to
the envelope in regular speech. First, we tested whether averaging the time-domain neural data (band-pass
filtered 1-40 Hz) with alignment to peakRate and peakEnv landmark events led to evoked responses and
examined their characteristics. While peakEnv closely follows on peakRate in regular speech, the interval
between them varied. Thus, aligning to the incorrect landmark should lead to (1) a reduced magnitude of
the averaged evoked neural signal due to smearing and (2) shifts in response onset times away from the
acoustic event. We found that alignment to either peakRate or peakEnv led to a robust transient evoked
response (Figure 1E). Crucially, the evoked response was of larger magnitude with alignment to peakRate
than to peakEnv (peak magnitude: $t(11) = 5.9$, $p < 0.001$). Moreover, this response started after peakRate
events, but before peakEnv events (response latency relative to the event for peakEnv: -12.5 ms; peakRate:
+50 ms, determined as the first significant time point in a cluster-based permutation test against 0).
Together, these analyses indicated that peakRate events, not peakEnv events, drove the evoked response.
This analysis of regular speech confirms that acoustic edges, and not envelope peaks, drive evoked
responses to the speech envelope in MEG, in line with previous results (Brodbeck et al., 2018; Doelling et
al., 2014; Oganian & Chang, 2019).
32
Cerebro-acoustic phase coherence between speech envelope and MEG

To confirm that cortical speech envelope tracking was present in our data (Peelle & Davis, 2012), we

calculated the cerebro-acoustic phase coherence (CAC) between neural responses and the speech

envelope in frequency bands below 10 Hz. CAC is typically increased at the frequency corresponding to

the speech rate (Pefkou et al., 2017), which in our data corresponds to the frequency of peakRate and

peakEnv in each rate condition (regular: 5.7 Hz, slow: 1.9 Hz). Indeed, speech rate had opposite effects

on CAC in these two frequency bands (2-way repeated-measures ANOVA, interaction $F(1, 11) = 31.20$,

$p < 0.001$, $\eta^2 = 0.30$, Figure 1F). At 5.7 Hz, CAC was higher for regular speech ($t(11) = 5.6$, $p < 0.001$,

$\eta^2 = 0.42$), while at 1.9 Hz it was higher for slow speech ($t(11) = 3.4$, $p = 0.006$, $\eta^2 = 0.29$). Moreover,

CAC was overall higher at lower frequencies ($F(1, 11) = 16.44$, $p < 0.001$, $\eta^2 = 0.39$), reflecting the 1/f

shape of MEG frequency spectrum. No other frequency band showed a significant effect of speech rate on

CAC (all Bonferroni-corrected $p > 0.05$). Overall, this result replicates previous findings of cortical

speech envelope tracking in frequency bands corresponding to the speech rate of the stimulus. However,

as this measure is calculated across the entire stimulus time course, it cannot capture local dynamics in

neural phase, driven by phase resets to acoustic edges.

Figure 2. Inter-event phase coherence (IEPC) to peakRate events in models and data. **A.** Schematic
illustrations of the predicted neural response to the utterance in Figure 1A using three different models. Top: Evoked
response to peakRate events; Middle: Fixed-frequency oscillatory model; Bottom: Flexible-frequency oscillatory
model. Colored bars illustrate time windows around peakRate events in which models predict different IEPC
magnitudes, in the absence of a repeated phase reset, as in D. **B.** IEPC patterns predicted by each model show
distinct spectral patterns for slowed speech. Dashed lines indicate the frequency of peakRate events in each
condition. **C**. Average IEPC for each model 0 – 250 ms after peakRate events. **D.** Illustration of predicted IEPC
following peakRate events that are at least 500 ms apart from neighboring events (~40 % of events in slow speech).
Note the shorter temporal extent of IEPC increase in evoked response than in oscillatory models. **E.** Neural data
IEPC with alignment to peakRate events in regular and slow speech. Outlined area: $p < 0.01$ with a cluster-based
permutation test. **F.** Mean IEPC 0 – 250 ms after peakRate events. **G**. Correlation between IEPC patterns predicted
by the three models (shown in B) and observed in MEG data (shown in E). Evoked response model predicts neural

IEPC better than oscillatory models, ** $p < 0.001$. All average responses are averaged across subjects and sensors;

error bars indicate ± 1 SEM.

Model Predictions for Delta-theta phase locking

We built three simple toy models of neural envelope tracking: an evoked response model, a fixed-

frequency oscillatory model, and a flexible-frequency oscillator model. The oscillator models were

designed to implement two central properties associated with oscillatory models in the speech literature:

phase alignment of an oscillatory response to external stimulation and reverberation of this response

following the phase reset for multiple (at least 2) oscillatory cycles. To compare the predictions of these

models with experimental data, we quantified inter-event phase coherence (IEPC) with an alignment to

peakRate. In contrast to prior studies of CAC, which quantified phase consistency across time, IEPC is

calculated across single event occurrences (i.e., single trials) for each time point. IEPC thus enables

tracking of the temporal dynamics of phase locking (Gross et al., 2013). To calculated IEPC as predicted

by each model for our regular and slow speech stimuli, we derived predicted response time series from

each of the three models (Figure 2A). To generate predictions of the evoked response model, we

convolved the peakRate event time series with a stereotypical auditory evoked response shape. For the

oscillator models, neural responses were modeled as a continuous sinusoidal oscillator, with a phase reset

at each peakRate event. In the fixed-frequency oscillator model, the oscillator's frequency was fixed at

5.7 Hz for both regular and slow speech conditions. For the flexible-frequency oscillator model, oscillator

frequencies were set to 5.7 Hz for regular speech and 1.9 Hz for slowed speech.

We then calculated model IEPC in the theta-delta range for each of the models, for both conditions

(Figure 2B). All three models predicted a transient increase in theta IEPC following peakRate events in

regular speech. However, their predictions for the slow speech condition diverged significantly (Figure

2C). The evoked response model predicted a two spectral peak in IEPC, around 6 Hz and around 2 Hz,

reflective of the shape of the evoked response and its frequency of occurrence, respectively. In contrast,

both oscillatory models predicted single spectral peaks in IEPC following peakRate events, centered

around the oscillatory frequency of each model. Notably, although the spectral range of peakRate IEPC

in the fixed-frequency oscillator model was wider in regular than in slow speech and spanned the entire

delta-theta range, its temporal extent and spectral dynamics clearly differed from the patterns predicted by

the evoked response model. To further distinguish between these two models, we analyzed the temporal

extent of neural phase-locking predicted by each model for peakRate events with a large temporal

distance to neighboring peakRate events (> 500 ms, $n = 456$ events, roughly 40% of peakRate events in

slow speech). In the absence of a new phase reset, the oscillatory model predicted that IEPC would

remain increased for multiple oscillatory cycles (Figure 2D). In contrast, in the evoked response model,

the increase in IEPC was temporally limited to the duration of the evoked response. Note that the precise temporal extent of the IEPC increase depends on the decay parameters of the oscillator model. However, a hallmark prediction of oscillatory models is that phase locking will continue after phase reset beyond a single oscillatory cycle. Thus, we focused our analyses on the first two cycles after peakRate events, which was sufficient to distinguish between models. Therefore, it was unnecessary to include a decay parameter in our models (see Methods for additional details).

Taken together, the qualitative model comparison identified two divergent model predictions for IEPC patterns in slow speech. Next, we performed these identical analyses on our neural data and compared the patterns in the data against the models' predictions.

Delta-Theta phase-locking to acoustic edges is best described by the evoked response model

First, we examined whether peakRate events induced theta phase reset in regular speech. Consistent with the time-domain analysis and predictions of all three models, we observed an increase in IEPC in the theta band (4-8 Hz) following peakRate events (Figure 2E left, two-dimensional cluster-based permutation against pre-event baseline $p < 0.001$).

Next, we tested how slowing would affect the frequency range of phase locking to peakRate events. If neural phase locking reflects evoked responses in slowed speech, phase locking should still be observed in the theta range with a second peak in the delta range. In contrast, if neural phase locking reflects an oscillatory phase reset, we should observe a single spectral peak in IEPC, with the frequency corresponding to the oscillator's frequency. In line with the former, we found two spectral peaks in IEPC to peakRate events in slow speech: a peak in theta, as also seen in regular speech (two-dimensional cluster-based permutation against pre-event baseline $p < 0.001$) and a second peak in the delta band, around the frequency of peakRate events in slow speech (Figures 2E right, two-dimensional cluster-based permutation against pre-event baseline $p < 0.001$). This pattern was predicted by the evoked response model but not by either oscillatory model.

Figure 3. Theta phase locking is limited to the duration of the evoked response after peakRate

events. A. IEPC across selected peakRate events that were at least 500 ms apart from preceding and

subsequent events. Black dots mark the extent of one oscillatory cycle in each frequency band. **B**. Group

distribution IEPC in 3 - 4 cycles around peakRate events and baseline in each frequency band. * $p < 0.05$.

Boxplots reflect median, 25^{th} and 75^{th} percentile, and range of values. Outliers are plotted individually.

Theta phase locking is limited to a single cycle after peakRate events.

A second prediction, differentiating between an evoked response and a fixed-frequency oscillator

model, was the temporal extent of phase alignment after a phase reset. In our data, this should be

expressed as an increase in IEPC, extending beyond a single oscillatory cycle after peakRate events

(Haegens & Zion Golumbic, 2018; Helfrich et al., 2019; Meyer et al., 2019). In contrast, if phase locking

results from evoked responses to peakRate events, the increase in IEPC should be limited to the duration

of the evoked response (approximately 250 ms, see Figure 1E). To test this, we again focused on those

peakRate events in the slowed speech that were at least 500 ms apart from preceding and subsequent

peakRate events ($n = 456$ events, 40 % of all events). For these events, IEPC increases in the theta band

(4 - 8 Hz, average cycle length 125 to 250 ms) would not be affected by preceding/subsequent peakRate

events for at least two oscillatory cycles, allowing us to test for continuous entrainment (Figures 3A, B).

Notably, such cycles were distributed throughout the slowed speech stimulus and not limited to sentence

or phrase ends. We extracted IEPC for each frequency band, for a single theoretical cycle width preceding

and up to three-cycle widths after each peakRate event (defined based on the center frequency of each

frequency band used in the analysis). Additionally, using a shuffling distribution, we calculated an IEPC

baseline. We first tested whether IEPC in the cycle immediately following peakRate events was elevated

as compared to surrounding cycles. In a second step, we tested whether IEPC in the pre- and post-event

cycles differed significantly from baseline.

For bands with two post-event cycles (4 bands in the range of 4 – 6Hz, see Figure 3), IEPC differed between cycles (repeated-measures ANOVA: main effect of cycle, $F (2, 22) = 14.82$, $p < 0.001$, $\eta^2 = 0.26$), and this modulation did not differ between bands (interaction: $p = 0.6$, n.s., main effect of band: Greenhouse-Geyser $p > 0.05$). Planned comparisons showed that IEPC was increased in the first post-event cycle relative to the pre-event cycle ($p < 0.001$) as well as relative to the 2^{nd} post-event cycle ($p < 0.001$). In contrast, we observed no significant difference in IEPC between the pre-event and the second post-event cycles ($p = 0.3$, n.s.), and IEPC in both cycles did not differ from the permutation-based baseline (all Bonferroni-corrected p's > 0.05). Similarly, for bands with three post-event cycles (three bands in the range of 6-8 Hz), IEPC differed between cycles ($F (3, 33) = 6.58$, $p < 0.001$, $\eta^2 = 0.15$), being higher in the cycle immediately following peakRate events relative to the other cycles (planned contrast of 1^{st} post-event cycle to all other cycles: $p = 0.002$). Here as well, we saw no significant IEPC difference between frequency bands ($p = 0.15$, n.s.), and the effect of the cycle did not differ significantly between bands ($p = 0.054$). Moreover, we found that IEPC did not differ between the permutation-based baseline and cycles that were not immediately following peakRate events (baseline vs. 1^{st} post-event cycle: Bonferroni-corrected $p = 0.006$; baseline vs. all other cycles Bonferroni-corrected p's > 0.05, n.s.). Notably, this pattern was evident in single subjects (see Figure S3 for single-subject data). Overall, the IEPC increase in the theta band extended for the duration of the evoked response, but not beyond. This analysis thus illustrates the transient nature of theta phase locking to peakRate events, which is more in line with an evoked response mechanism of speech envelope tracking than with an oscillatory entrainment model.

Collectively, these findings disagree with an oscillatory entrainment account, which postulates an oscillatory phase reset after an event, followed by continuous oscillatory reverberation. A more parsimonious account of our results is that the low-frequency phase locking to the speech envelope in MEG is driven by evoked responses to peaks in the envelope rate of change (peakRate). Furthermore, our analysis shows that IEPC to peakRate events reflects the superposition of two different sources: (1) local responses to individual peakRate events and (2) the rate of occurrence of responses to peakRate events. Furthermore, our analyses demonstrate that the shift in IEPC frequency bands with changes in speech rate may be the product of a time-frequency decomposition of a series of evoked responses, rather than a shift in the frequency of an entrained oscillator. In a final analysis, we thus tested whether IEPC to peakRate is normalized to account for changes in speech envelope dynamics induced by changes in speech rate.

Speech rate normalization of peakRate IEPC

The perceptual ability to adapt to variation in the speech signal resulting from changes in the speech rate, i.e., the number of syllables produced per second, is referred to as speech rate normalization.

Changes in speech rate result in acoustic changes in the speech signal, including slower amplitude increases at acoustic edges, that is, lower peakRate magnitudes (Figure 4A-C). We had previously found that responses to peakRate monotonically scale with peakRate magnitude, being larger for faster changes in the speech amplitude (Oganian & Chang, 2019). Efficient envelope tracking across speech rates would thus require remapping of neural responses to peakRate magnitude to account for this overall reduction. Here, we assessed the effect of speech rate on the magnitude of theta IEPC to peakRate events. In the slowed speech, stimuli peakRate magnitudes were 1/3 of those in regular speech (Figures 4A-C). If no normalization occurs, IEPC magnitudes in slow speech should reflect absolute peakRate values, resulting in an overall reduction in IEPC (Figure 4F, dark dots). In contrast, if theta IEPC to peakRate is invariant to speech rate, it should reflect peakRate values relative to the contextual speech rate, resulting in similar IEPC magnitudes in both speech rate conditions (Figure 4F, light dots).

An evaluation of IEPC after peakRate events, split by peakRate magnitude quantiles, showed comparable theta IEPC in both speech rate conditions (Figure 4D-E), such that average theta IEPC was more robust for larger peakRate magnitudes across both rate conditions (the main effect of peakRate quantile: $b = 0.01$, SD $= 0.001$, $t = 1.4$, $\chi^2 = 55.0$, $p = 10^{-13}$). Crucially, they did not differ between regular and slow speech (Interaction effect: $b = 0.003$, SD $= 0.005$, $t = 0.6$, n.s., Figure 4G), as expected in the case of speech rate normalization (Figure 4F, dark dots). Thus, the magnitude of phase reset induced by peakRate depended on its magnitude relative to the local speech rate context, allowing for the flexible encoding of peakRate information at different speech rates.

Figure 4. Normalization of peakRate IEPC for contextual speech rate. A. Histogram of peakRate magnitudes in regular speech, with quantile boundaries marked in red. **B.** Same as A for slow speech **C.** Quantile-Quantile plot of peakRate magnitudes in regular and slowed speech stimulus. peakRate values in slowed speech stimulus are 1/3 of peakRate values in regular speech stimulus. **D.** IEPC in 1st, 3rd, 5th peakRate magnitude quantile. Horizontal lines mark the theta frequency range (4 – 8 Hz). **E.** Same as D for slow speech. **F.** Predicted quantile-quantile plots of theta IEPC in regular and slowed speech with (dark) or without (light) normalization. **G.** Quantile-quantile plot of theta-band IEPC (mean, error bars mark ± 1 SEM across subjects) in regular and slow speech. Theta IEPC quantile-quantile values are close to the diagonal, indicating similar magnitudes of theta IEPC in regular and slowed speech conditions.

Discussion

We evaluated local temporal dynamics in MEG neural responses to continuous speech under the predictions of evoked response and oscillatory entrainment models. Confirming previous results, we found that acoustic edges (peakRate events), but not peaks in the envelope (peakEnv events), drove evoked responses and phase locking in sensors over auditory cortical areas (Brodbeck et al., 2018;

Hertrich et al., 2012; Oganian & Chang, 2019). Only an evoked response model, but not oscillatory entrainment models captured the spectral and temporal extent of phase locking to acoustic edges: a transient local component in the theta range, reflective of the evoked response, and - additionally in slow speech - a separate global component, which captured the frequency of evoked responses in the stimulus. An analysis of portions of the speech signal with temporally sparse acoustic events showed that phase locking to acoustic edges was transient and limited to the duration of the evoked response, providing no evidence for sustained oscillatory phase locking in the theta band, a hallmark marker of entrainment models (Helfrich et al., 2019; Peelle & Davis, 2012). Moreover, we found that the magnitude of the evoked phase reset to acoustic edges reflected the amplitude slope at the acoustic edge after normalization for speech rate, offering novel neural evidence for speech rate normalization. Overall, our findings confer with the predictions of the evoked response model and suggest that neural phase locking induced by evoked responses to acoustic edges is the primary source of speech envelope tracking in the theta-delta band in MEG.

Although phase locking between low-frequency MEG components and the amplitude envelope of speech is well established (Ahissar et al., 2001; Luo & Poeppel, 2007), considerable dispute surrounds the question of which acoustic events in the speech signal drive this phase locking (Aubanel et al., 2016; Doelling et al., 2014; Ghitza, 2011; Peelle & Davis, 2012). In natural speech, which does not show robust temporal rhythmicity, such phase locking must rely on the repeated realignment of neural phase to the speech signal (Haegens & Zion Golumbic, 2018). Here, we show that acoustic edges drive such phase resets, in line with recent intracranial results (Oganian & Chang, 2019). This supports the central role of acoustic edges in speech comprehension, as they mark the onsets of stressed vowel nuclei and closely correspond to perceptual centers (P-centers) of syllables (Harsin, 1993; Scott, 1998), and establishes them as the basis for the cortical representation of the speech envelope.

Neural phase resetting may be due to either the superposition of evoked responses or the entrainment of endogenous oscillatory activity. To distinguish between them, we derived the spectral and temporal extent of phase locking to acoustic edges using simulations of evoked responses and oscillatory models. We found two local spectral peaks in phase reset to acoustic edges in slow speech – in the theta (4 - 8 Hz) and in the delta frequency range (1 - 4 Hz). Crucially, only an evoked response model predicted this pattern, whereas oscillatory models predicted phase locking centered around a single spectral location. As theta phase locking to acoustic edges was prominent both in regular and in slow speech, to further distinguish whether it reflected evoked responses or oscillatory entrainment, we tested a second hallmark prediction attributed to oscillatory entrainment: the persistence of an oscillation after the phase-resetting stimulus (Haegens & Zion Golumbic, 2018; Helfrich et al., 2019). To this end, we focused on speech portions where the intervals between consecutive acoustic edges were long enough to detect reverberation

in the theta band (4 - 8 Hz). We found that phase locking increased for the duration of the evoked response, but diminished right after to the baseline level, showing no evidence of reverberation and supporting the interpretation of theta phase locking as a transient evoked response rather than a theta oscillator. Moreover, this analysis showed no phase alignment in the cycle preceding peakRate events, providing no evidence for the anticipatory phase locking to the speech envelope in the theta band.

Taken together, these analyses show that a linear convolution of evoked responses to discrete acoustic edge events in the speech signal is sufficient to account for the pattern of neural phase locking observed in response to continuous speech. In particular, a spectral analysis of an evoked response model shows that it is capable of generating a low-frequency component of neural phase locking, corresponding to the frequency of acoustic edge events in slow speech. It should be noted that our results do not refute the possibility that an additional oscillatory component contributes to phase locking at the speech rate frequency in each condition. However, our results show that such a component is not necessary to produce phase-locking patterns as observed for natural speech in MEG, as these can be explained more parsimoniously by an evoked response model. Our finding that speech envelope tracking reflects the convolution and superposition of repeated evoked responses has major implications for theories of speech perception. For instance, instead of oscillatory resonance, predictive processing of speech could rely on non-oscillatory temporal prediction mechanisms guided by statistical learning (Breska & Deouell, 2017; Friston et al., 2020; Sohoglu & Davis, 2016).

Previous research has suggested distinct functional roles of delta and theta band entrainment in linguistic processing, with theta implicated in envelope tracking and delta in the processing of phrasal boundaries (Donhauser & Baillet, 2020; Meyer et al., 2017). In the present study, however, both theta and delta effects reflect responses to the acoustic amplitude envelope of speech, as the effect in the delta band arose with the slowing of the speech stimulus.

Speech rate normalization is a central behavioral (Reinisch, 2016; Wade & Holt, 2005) and neural phenomenon in speech perception. Shifting of the entrained oscillatory frequency to match the overall speech rate of the input was previously proposed as its neural mechanism (Alexandrou et al., 2018b; Kösem et al., 2018). Here, however, we found that the shift of neural phase locking to lower frequencies with speech slowing might be an epiphenomenon of spectral analysis of a neural signal containing a series of evoked responses. Instead, we found that the magnitude of phase locking to acoustic edges was normalized to reflect the distribution of peakRate magnitudes at each speech rate. Namely, while phase locking to acoustic edges was more robust for sharper edges (Biermann & Heil, 2000; Oganian & Chang, 2019), it was comparable in regular and slowed speech, despite acoustic edges being flatter in slow speech. This result suggests that the cortical representations of acoustic edges reflect the magnitude of an edge relative to the contextual speech rate. Such shifting of the dynamic range for the representation of

acoustic edge magnitude constitutes a flexible mechanism that maximizes the dynamic range for the representation of speech temporal dynamics (Diehl, Souther, & Convis, 1980; Hirataa & Lambacher, 2004). Further studies are necessary to elucidate the neural mechanisms underlying this process at the circuit level. One possibility is that the local distribution of peakRate values is continuously learned via dynamic adjustment of synaptic weights.

Our approach represents a methodological departure from previous investigations of speech envelope tracking. Namely, previous studies focused on cerebro-acoustic coherence (CAC), which reflects the consistency of phase differences between the neural signal and the acoustic stimulus across time (Peelle, Gross, & Davis, 2013). CAC is primarily sensitive to regularities across time, such as the rate of phase resets. In contrast, we used inter-event phase coherence (IEPC), which focuses on assessing temporally local similarities in neural phase across repeated occurrences of the same acoustic event (see (Gross et al., 2013) for IEPC to speech onsets). As IEPC is sensitive to local phase resets as well as to their rate of occurrence, this approach revealed that both components are reflected in IEPC to acoustic edges. In regular speech, both components overlapped within the theta frequency range. In contrast, slowing of the speech signal separated them in an evoked response component in the theta band and a slower event frequency component.

Speech rate manipulations are frequently used to study neural tracking of speech, in particular, speech envelope tracking (Ahissar et al., 2001; Ghitza & Greenberg, 2009; Nourski et al., 2009; Pefkou et al., 2017). Most previous studies used compressed speech, in which speech rate is increased, to study temporal boundaries on envelope tracking and intelligibility. In contrast, here, we used slowed speech as we aimed to spread distinct acoustic envelope features out in time. It is essential to reconsider previous findings in light of the evoked response framework. For example, while envelope tracking and intelligibility deteriorate for speech rates higher than 8 Hz, insertion of brief silence periods in compressed speech, which returns the effective speech rate to below 8 Hz, improves intelligibility (Ghitza & Greenberg, 2009). While this result is typically interpreted as evidence for oscillatory envelope tracking in the theta range, within an evoked response framework, it might be reflective of the minimal refractory period of neural populations that encode acoustic edges in speech.

Recent studies have demonstrated that neural responses to speech in M/EEG reflect not only the amplitude envelope but also phonetic and semantic information (Broderick et al., 2019; Di Liberto et al., 2015; O'Sullivan et al., 2015). These studies focused on impulse responses (evoked responses) to different speech features and their modulation by higher-order information content. We extended upon these studies by pinpointing the envelope feature that drives neural responses to the envelope and showed that neural phase locking to this feature, acoustic edges, can be fully driven by an impulse response. Furthermore, we found responses to speech onsets and amplitude modulations in an ongoing speech on

the same sensors over temporal cortical areas. This stands in contrast to intracranial results, which showed that distinct areas of the superior temporal gyrus represent speech onsets and the content of ongoing speech (Hamilton et al., 2018; Oganian & Chang, 2019). However, this is not surprising as we focused on an analysis of MEG data from sensors over the temporal cortex, reflecting the summation of neural sources across auditory cortical areas, such that the shape of the evoked responses to acoustic edges is likely to also be influenced by other temporally confounded events in the speech envelope (such as vowel onsets).

Natural speech does not have robust temporal rhythmicity (Alexandrou et al., 2018a). Our focus on envelope tracking for natural speech indicates that, in this case, neural signatures of envelope tracking are well explained by an evoked response model without the need for an oscillatory component. These results seemingly contradict recent findings of predictive entrainment to music (Doelling et al., 2019). However, our study employed natural speech with considerable variability in inter-edge intervals, unlike in rhythmic musical stimuli. Critically, recent neuropsychological work dissociated neural mechanisms for prediction based on rhythmic streams from predictions in non-rhythmic streams (Breska & Ivry, 2018). This adds an important caveat to the current debate, suggesting that the results of Doelling et al. may perhaps not extend to natural speech with inherent temporal variability and reduced rhythmicity. The present study thus calls to reevaluate the role of oscillatory entrainment in natural speech comprehension. However, it does not preclude the possibility that the introduction of additional rhythmicity to speech, e.g., in poetry or song, recruits additional neural processes associated with the processing of rhythms. Such additional processes might support speech comprehension and could underlie some of the recent findings obtained with a rhythmic speech stimulus (Ding et al., 2015; ten Oever & Sack, 2015; Zoefel et al., 2019). Therefore, caution needs to be exercised when extending findings obtained with rhythmic stimuli (e.g., (Ding et al., 2015; Doelling et al., 2019; Zoefel, Archer-Boyd, et al., 2018)) to natural speech.

In summary, our results show that an evoked response model accounts for the main neural signatures of speech envelope tracking in MEG. The neural representation of acoustic edges serves as a cue to syllabic nucleus onsets and informs about speech rate via the inter-event interval. Moreover, the speech rate normalization of these responses renders this mechanism flexibly adaptable to changes in speech rate. To conclude, evoked responses to acoustic edges track the syllabic rate in speech and provide a flexible framework for temporal analysis and prediction during speech perception.

Methods

Participants

Twelve healthy, right-handed volunteers (six females; age range 22 - 44 years, median 25 years)

participated in the study. All participants were native speakers of English. All participants provided

informed written consent and received monetary compensation for their participation. The study was

approved by the University of California, San Francisco Committee on Human Research. All participants

gave informed written consent before experimental testing.

Speech stimulus

Task stimuli contained a selection of two 3-minute new excerpts from the Boston University Radio

Speech Corpus (BURSC, Table S1) (Ostendorf et al., 1995), produced by a male and a female voice. The

stimuli were split into 62 utterances of 2.2 – 14.9 s, with silent periods of 500 – 1100 ms between

utterances. Slowed speech stimuli consisted of the same utterances slowed to 1/3 of the original speech

rate. Speech stimuli were slowed using the PSOLA (Pitch Synchronous Overlap and Add) algorithm, as

implemented in the software Praat (Boersma & Weenik, 2019), which slows down the temporal structure

of the speech signal while keeping its spectral structure constant (Moulines & Charpentier, 1990).

Stimulus presentation

All stimuli were presented binaurally at a comfortable ambient loudness (~ 70 dB) through MEG

compatible headphones using custom-written MATLAB R2012b scripts (Mathworks,

https://www.mathworks.com). Speech stimuli were sampled at 16,000 Hz. Participants were asked to

listen to the stimuli attentively and to keep their eyes closed during stimulus presentation.

Regular and slow speech were presented interleaved in 4 blocks, such that half the stimulus was

heard first in the regular condition and half in the slowed condition. Comprehension was assessed with 3 -

4 unique comprehension questions after each block (Table S2). Comprehension was compared between

regular and slow speech using a two-sided paired t-test.

Data analysis

All analyses were conducted in MATLAB R2019a (Mathworks, https://www.mathworks.com) using

standard toolboxes and custom-written scripts. The FieldTrip toolbox (Oostenveld et al., 2011) was used

for visualization of topography.

Acoustic feature extraction

We extracted the broad amplitude envelope of speech stimuli by low-pass filtering to 10 Hz and

down-sampling to 100 Hz. We then calculated the derivative of the resulting loudness contours as a

measure of the rate of change in the amplitude envelope. Finally, we extracted the sparse time series of

local peaks in the amplitude envelope (peakEnv) and its derivative (peakRate). All features are depicted in Figure 1A, for an example stimulus excerpt. Overall the stimulus set contained 2,106 peakRate and peakEnv events per condition.

Speech preprocessing

To assess cerebro-acoustic phase coherence between the speech envelope and MEG responses, the speech envelope was processed using the same procedure that was applied to the MEG responses: Down-sampling to 400 Hz and band-passed between 0.2 and 10 Hz using the second-order Butterworth filter. We then used the Hilbert transform to extract the phase in the delta-theta frequency range using eighteen band-pass filters (Gaussian filters, logarithmically increasing center frequencies between 0.2 and 10 Hz with semi-logarithmically increasing bandwidth).

Neural data acquisition and preprocessing

MEG recordings were obtained with a 275-axial gradiometers whole-head MEG system (CTF, Coquitlam, British Columbia, Canada) at a sampling rate of 1,200 Hz. Three fiducial coils were placed on the nasion and left and right pre-auricular points to triangulate the position of the head relative to the MEG sensor array. The position of the patient's head in the device relative to the MEG sensors was determined using indicator coils before and after each recording interval to verify an adequate sampling of the entire field. The fiducial markers were later co-registered onto a structural magnetic resonance imaging scan to generate head shape (Teichmann et al., 2013).

Offline data preprocessing included (in this order) artifact rejection with dual signal subspace projection (DSSP) and down-sampling to 400 Hz. DSSP is a MEG interference rejection algorithm based on spatial and temporal subspace definition (Sekihara et al., 2016). Its performance has been recently validated using clinical data (Cai et al., 2019). To assess the broadband response, data were then band-pass filtered between 1 and 40 Hz using a second-order Butterworth filter. To evaluate the power and phase of each frequency component, we extracted the phase and analytic amplitude from eighteen bands between 0.2 and 10 Hz, using band-pass filters (Gaussian filters, logarithmically increasing center frequencies with semi-logarithmically increasing bandwidths) and the Hilbert transform.

Data segmentation

The evoked response, band-passed power, and phase were segmented into symmetric epochs of 1,000 ms duration around onsets of utterances, peakEnv events, and peakRate events, respectively. Segments containing sensor data above 1.5 pT and visually identified artifacts (including muscle, eye blink, and motion) were flagged as bad events and removed from further processing (0.2 % of segments).

1   For the evoked response and narrow-band power analysis, baseline correction (- 200 ms to 0 ms
2   relative to event onset for the evoked response, - 400 ms to - 200 ms for narrow-band power) was applied
3   before averaging across events. No baseline correction was applied to the instantaneous phase.
4
5   **Sensor selection**
6   We selected sensors based on the magnitude of the group averaged M100 waveform, 60 - 100 ms
7   after the onset of utterance. To focus all further analyses on responses originating in temporal auditory
8   areas, we selected ten sensors with maximal M100 responses from each hemisphere.
9
10   **Cerebro-acoustic phase coherence (CAC)**
11   Phase locking between the speech envelope and MEG response was calculated across the entire
12   duration of every utterance within each frequency band, using the Cerebro-acoustic phase coherence
13   (CAC):

$$CAC(\varphi) = \frac{1}{N} \left| \sum_{t=1}^{T} \exp\left(i * (ph(\varphi, t) - phs(\varphi, t))\right) \right|$$

14
where φ is the center frequency of a frequency band, T is the number of time samples in an utterance,
ph is the phase of the neural signal, and *phs* is the phase of the speech envelope in band φ at time *t*. To
equate the number of time points entering the analysis for slow and regular speech, slow speech
utterances were split into three equal parts before CAC calculation, and resultant CAC values were
averaged. CAC was averaged across all twenty sensors included in the analysis.
A priori, we hypothesized that CAC would differ between conditions in the frequency bands
corresponding to the average frequency of peakRate events in each rate condition (regular: 5.7 Hz; slow:
1.9 Hz, Figure 1F). We tested this hypothesis using a 2-way repeated-measures ANOVA with factors
frequency band (high/low), factor speech rate (slow/regular).
To test for further differences in each frequency band, we assessed the effect of speech rate onto
CAC using a repeated-measures *t*-test between speech rate (slow/regular) across frequency bands.
Significance in this analysis was Bonferroni-corrected for multiple comparisons across bands.
27
28   **Model simulations**
29   We created simple simulations of neural response time series in response to our speech stimuli for an
30   evoked response and two oscillatory models (see details below). After creating the predicted response
31   time series for each model, we added noise with 1/f spectral content and applied additional filtering using
32   the same procedures as for the MEG data. Noise and neural response were normalized to the same range,

such that the model data had an SNR of 1. Changes to the SNR and the slope of the 1/f noise had no effect on the qualitative pattern of IEPC results.

Evoked response model for IEPC simulation

We used the group-averaged N100 response estimated from our data for this model (similar to recent approaches in the field, Doelling et al. 2019). While we found no effect of speech slowing on the shape of the neural response to peakRate events in our previous intracranial work, we assume that neural responses recorded with MEG will be additionally shaped by other speech features that occur in temporal proximity to peakRate events (e.g., vowel onsets). As the relative latencies of peakRate events and such other components of the speech signal increase with slowing, we slowed this stereotypical evoked response to 75% of its original duration in the slow speech condition. Note that this value was determined a-priori to create a slight widening uninformed by the neural data. We chose this approach as fairer for the overall model comparison, as it is agnostic to the evoked responses in the actual data, just as the oscillator models used stereotypical oscillators without additional fitting to the neural data. To create the predicted response time series for our continuous speech stimulus, we linearly convolved the model response with the time series of peakRate events.

Oscillatory entrainment model for IEPC simulation

To simulate neural phase-locking under the assumption of endogenous oscillatory entrainment, we focused on two alternative models. For the fixed-frequency oscillator model, we assumed a single theta-band oscillator with its frequency centered at the speech rate in our regular speech condition, which coincided with the center of the theta band (6 Hz, with theta at 4-8Hz). For the flexible-frequency oscillator model, we assumed that the oscillator frequency would shift to reflect the speech rate in each condition; thus, its frequency was set at 5.7 Hz for regular speech and at 1.9 Hz for slow speech. For ease of illustration, we adopted a very simple model with a full phase-reset at each peakRate event and no decay of oscillator magnitude after phase-reset. Change to these parameters would vary the magnitude of oscillatory entrainment; however, it would not change its spectral distribution, which was in the focus of the present study.

Inter-event phase coherence (IEPC)

From the phase time series for each frequency band, epochs with 1,000 ms duration (- 500 ms to + 500 ms relative to event onset) were extracted for peakRate events. Inter-event phase coherence (IEPC) (Lachaux, Rodriguez, Martinerie, & Varela, 1999) was computed for each condition, for segmentations around peakRate events. IEPC was calculated using the following formula:

$$IEPC(\varphi, t) = \frac{1}{N} \left| \sum_{k=1}^{N} \exp\left(i * ph_k(\varphi, t)\right) \right|$$

where N is the number of events, ph is the phase of the neural signal in trial k, for the frequency band

φ and timepoint t. IEPC were averaged across subjects and selected sensors. The resulting time x

frequency IEPC was compared with the pre-event baseline using 2-D cluster-based permutation t-tests

(Maris & Oostenveld, 2007) with 3,000 permutations, a peak threshold of $p < 0.05$, and a cluster

threshold of $p < 0.01$. Baseline IEPC was calculated as the average IEPC between - 400 ms to - 100 ms

relative to event onset in each frequency band.

Theta-band IEPC for peakRate events with large inter-event distance

To assess the temporal extent of theta-band IEPC between peakRate events, we analyzed IEPC

following peakRate events with a temporal interval of more than 500 ms to adjacent peakRate events. For

each frequency band between 4 and 8 Hz, we computed the average IEPC in time windows corresponding

to a single oscillatory cycle at the band center frequency, for a single cycle before and up to 3 cycles post-

event. Post-event, the maximal number of cycles within the 500 ms window was included for each band.

Bands were split into two sets, according to the number of post-event cycles in the analysis (2 cycles: 4

Hz, 4.5 Hz, 5 Hz, 5.5 Hz; 3 cycles: 6.1 Hz, 6.7 Hz, 7.4 Hz). The IEPC baseline was calculated using

circular permutation of peakRate time series across sentences. In each frequency band, baseline IEPC was

averaged across the time window of a single oscillatory cycle. For each frequency range, IEPC in

neighboring oscillatory cycles was compared using 2-way repeated-measures ANOVAs with factors

frequency band x cycle relative to the event. Then, IEPC in the pre- and end post-event cycles was

compared with baseline using 2-way repeated-measures ANOVAs with factors frequency band x cycle

time window relative to event time. Pairwise comparisons between cycles were Holm-corrected for

multiple comparisons. Pairwise comparisons to the permutation-based baseline were Bonferroni-corrected

for multiple comparisons across frequency bands. Note that IEPC in the cycle before peakRate events and

the last cycle following the peakRate event are not necessarily identical: increased IEPC in the cycle

before peakRate event would indicate predictive processing based on the preceding peakRate event

(Doelling et al., 2019), whereas increased IEPC in the last extracted cycle would be an indicator of

continuous entrainment, even if it does not capture the timing of the next event. Those two would be

identical for an isochronous stimulus sequence but diverge otherwise, for example, in natural speech.

Effect of peakRate magnitude on IEPC

In each rate condition, peakRate events were split into five quantiles, and IEPC was separately

calculated within each quantile. Then, we extracted the average IEPC in the theta band (4 – 8 Hz) across

all the time points for one cycle of the given frequency band after the event. IEPC in each quantile was

compared using 2-way ANOVA with factors quantile and speech rate (regular speech, slow speech).

Data and code availability

All custom-written analysis code will be publicly available upon publication on github

(https://github.com/ChangLabUcsf/MEG-SlowSpeech). Data will be made available upon request from

the corresponding authors.

Author contributions: Y.O and E.F.C conceived the study, Y.O and K.K designed the experiments

and analyzed the data. K.K., C.C., and A.F collected and preprocessed the data. K.K. and Y.O. wrote the

manuscript, K.K., Y.O, E.F.C, and S.N revised the manuscript.

References

Abrams, D. A., Nicol, T., Zecker, S., & Kraus, N. (2008). Right-Hemisphere Auditory Cortex Is Dominant for Coding Syllable Patterns in Speech. *J Neurosci*, *28*(15), 3958–3965. https://doi.org/10.1523/JNEUROSCI.0187-08.2008

Ahissar, E., Nagarajan, S. S., Ahissar, M., Protopapas, A., Mahncke, H., & Merzenich, M. M. (2001). Speech comprehension is correlated with temporal response patterns recorded from auditory cortex. *Proceedings of the National Academy of Sciences of the United States of America*, *98*(23), 13367–13372. https://doi.org/10.1073/pnas.201400998

Alexandrou, A. M., Saarinen, T., Kujala, J., & Salmelin, R. (2018a). Cortical entrainment: What we can learn from studying naturalistic speech perception. *Language, Cognition and Neuroscience*, *0*(0), 1–13. https://doi.org/10.1080/23273798.2018.1518534

Alexandrou, A. M., Saarinen, T., Kujala, J., & Salmelin, R. (2018b). Cortical Tracking of Global and Local Variations of Speech Rhythm during Connected Natural Speech Perception. *Journal of Cognitive Neuroscience*, *26*(3), 1–16. https://doi.org/10.1162/jocn_a_01295

Aubanel, V., Davis, C., & Kim, J. (2016). Exploring the Role of Brain Oscillations in Speech Perception in Noise: Intelligibility of Isochronously Retimed Speech. *Frontiers in Human Neuroscience*, *10*. https://doi.org/10.3389/fnhum.2016.00430

Biermann, S., & Heil, P. (2000). Parallels Between Timing of Onset Responses of Single Neurons in Cat and of Evoked Magnetic Fields in Human Auditory Cortex. *Journal of Neurophysiology*, *84*(5), 2426–2439.

Boersma, P., & Weenik, D. (2019). Praat: Doing phonetics by computer [Computer program]. Version 6.1.08, retrieved 5 December 2019 from http://www.praat.org/. *Amsterdam: University of Amsterdam.*

Bosker, H. R., & Kösem, A. (2017). An Entrained Rhythm's Frequency, Not Phase, Influences Temporal Sampling of Speech. *Interspeech 2017*, 2416–2420. https://doi.org/10.21437/Interspeech.2017-73

Breska, A., & Deouell, L. Y. (2017). Neural mechanisms of rhythm-based temporal prediction: Delta phase-locking reflects temporal predictability but not rhythmic entrainment. *PLOS Biology*, *15*(2), e2001665. https://doi.org/10.1371/journal.pbio.2001665

Breska, A., & Ivry, R. B. (2018). Double dissociation of single-interval and rhythmic temporal prediction in cerebellar degeneration and Parkinson's disease. *Proceedings of the National Academy of Sciences*, *115*(48), 12283–12288. https://doi.org/10.1073/pnas.1810596115

Brodbeck, C., Hong, L. E., & Simon, J. Z. (2018). Rapid Transformation from Auditory to Linguistic Representations of Continuous Speech. *Current Biology*, *28*(24), 3976-3983.e5. https://doi.org/10.1016/j.cub.2018.10.042

Broderick, M. P., Anderson, A. J., & Lalor, E. C. (2019). Semantic Context Enhances the Early Auditory Encoding of Natural Speech. *The Journal of Neuroscience*, 0584–19. https://doi.org/10.1523/JNEUROSCI.0584-19.2019

Cai, C., Xu, J., Velmurugan, J., Knowlton, R., Sekihara, K., Nagarajan, S. S., & Kirsch, H. (2019). Evaluation of a dual signal subspace projection algorithm in magnetoencephalographic recordings from patients with intractable epilepsy and vagus nerve stimulators. *Neuroimage*, *188*, 161–170. https://doi.org/10.1016/j.neuroimage.2018.11.025

Capilla, A., Pazo-Alvarez, P., Darriba, A., Campo, P., & Gross, J. (2011). Steady-State Visual Evoked Potentials Can Be Explained by Temporal Superposition of Transient Event-Related Responses. *PLoS ONE*, *6*(1), e14543. https://doi.org/10.1371/journal.pone.0014543

Chait, M., Simon, J. Z., & Poeppel, D. (2004). Auditory M50 and M100 responses to broadband noise: Functional implications: *NeuroReport*, *15*(16), 2455–2458. https://doi.org/10.1097/00001756-200411150-00004

Di Liberto, G. M., O'Sullivan, J. A., & Lalor, E. C. (2015). Low-frequency cortical entrainment to speech reflects phoneme-level processing. *Current Biology*, *25*(19), 2457–2465. https://doi.org/10.1016/j.cub.2015.08.030

Ding, N., Melloni, L., Zhang, H., Tian, X., & Poeppel, D. (2015). Cortical tracking of hierarchical linguistic structures in connected speech. *Nature Neuroscience*, *19*(1), 158–164. https://doi.org/10.1038/nn.4186

Doelling, K. B., Arnal, L. H., Ghitza, O., & Poeppel, D. (2014). Acoustic landmarks drive delta-theta oscillations to enable speech comprehension by facilitating perceptual parsing. *NeuroImage*, *85*(0 2), 761–768. https://doi.org/10.1016/j.neuroimage.2013.06.035

Doelling, K. B., Assaneo, M. F., Bevilacqua, D., Pesaran, B., & Poeppel, D. (2019). An oscillator model better predicts cortical entrainment to music. *Proceedings of the National Academy of Sciences*, *116*(20), 201816414. https://doi.org/10.1073/pnas.1816414116

Donhauser, P. W., & Baillet, S. (2020). Two Distinct Neural Timescales for Predictive Speech Processing. *Neuron*, *105*(2), 385-393.e9. https://doi.org/10.1016/j.neuron.2019.10.019

Drullman, R., Festen, J. M., & Plomp, R. (1994a). Effect of temporal envelope smearing on speech reception. *The Journal of the Acoustical Society of America*, *95*(2), 1053–1064. https://doi.org/10.1121/1.408467

Drullman, R., Festen, J. M., & Plomp, R. (1994b). Effect of reducing slow temporal modulations on speech reception. *The Journal of the Acoustical Society of America*, *95*(5), 2670–2680. https://doi.org/10.1121/1.409836

Friston, K. J., Sajid, N., Quiroga-Martinez, D. R., Parr, T., Price, C. J., & Holmes, E. (2020). Active listening. *Hearing Research*, 107998. https://doi.org/10.1016/j.heares.2020.107998

Ghitza, O. (2011). Linking speech perception and neurophysiology: Speech decoding guided by cascaded oscillators locked to the input rhythm. *Frontiers in Psychology*, *2*(JUN), 1–13. https://doi.org/10.3389/fpsyg.2011.00130

Ghitza, O. (2013). The theta-syllable: A unit of speech information defined by cortical function. *Frontiers in Psychology*, *4*(MAR), 138. https://doi.org/10.3389/fpsyg.2013.00138

Ghitza, O., & Greenberg, S. (2009). On the Possible Role of Brain Rhythms in Speech Perception: Intelligibility of Time-Compressed Speech with Periodic and Aperiodic Insertions of Silence. *Phonetica*, *66*(1–2), 113–126. https://doi.org/10.1159/000208934

Gross, J., Hoogenboom, N., Thut, G., Schyns, P., Panzeri, S., Belin, P., & Garrod, S. (2013). Speech Rhythms and Multiplexed Oscillatory Sensory Coding in the Human Brain. *PLoS Biology*, *11*(12), e1001752. https://doi.org/10.1371/journal.pbio.1001752

Gwilliams, L. (2019). Hierarchical oscillators in speech comprehension: A commentary on Meyer Sun and Martin 2019. *Language, Cognition and Neuroscience*. https://doi.org/10.1080/23273798.2020.1740749

Haegens, S., & Zion Golumbic, E. (2018). Rhythmic facilitation of sensory processing: A critical review. *Neuroscience & Biobehavioral Reviews*, *86*, 150–165. https://doi.org/10.1016/j.neubiorev.2017.12.002

Hamilton, L. S., Edwards, E., & Chang, E. F. (2018). A Spatial Map of Onset and Sustained Responses to Speech in the Human Superior Temporal Gyrus. *Current Biology*, *28*(12), 1860-1871.e4. https://doi.org/10.1016/j.cub.2018.04.033

Harsin. (1993). Acoustics of perceptual centers in speech. *The Journal of the Acoustical Society of America*, *94*(3), 1864–1864. https://doi.org/10.1121/1.407646

Helfrich, R. F., Breska, A., & Knight, R. T. (2019). Neural entrainment and network resonance in support of top-down guided attention. *Current Opinion in Psychology*, *29*, 82–89. https://doi.org/10.1016/j.copsyc.2018.12.016

Hertrich, I., Dietrich, S., Trouvain, J., Moos, A., & Ackermann, H. (2012). Magnetic brain activity phase-locked to the envelope, the syllable onsets, and the fundamental frequency of a perceived speech signal. *Psychophysiology*, *49*(3), 322–334. https://doi.org/10.1111/j.1469-8986.2011.01314.x

Hovsepyan, S., Olasagasti, I., & Giraud, A.-L. (2020). Combining predictive coding and neural oscillations enables online syllable recognition in natural speech. *Nature Communications*, *11*(1). https://doi.org/10.1038/s41467-020-16956-5

Kösem, A., Bosker, H. R., Takashima, A., Meyer, A., Jensen, O., & Hagoort, P. (2018). Neural Entrainment Determines the Words We Hear. *Current Biology*, *28*(18), 2867-2875.e3. https://doi.org/10.1016/j.cub.2018.07.023

Luo, H., & Poeppel, D. (2007). Phase Patterns of Neuronal Responses Reliably Discriminate Speech in Human Auditory Cortex. *Neuron*, *54*(6), 1001–1010. https://doi.org/10.1016/j.neuron.2007.06.004

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Mermelstein, P. (1975). Automatic segmentation of speech into syllabic units. *The Journal of the Acoustical Society of America*, *58*(4), 880–883. https://doi.org/10.1121/1.380738

Meyer, L., Henry, M. J., Gaston, P., Schmuck, N., & Friederici, A. D. (2017). Linguistic Bias Modulates Interpretation of Speech via Neural Delta-Band Oscillations. *Cerebral Cortex*, *27*(9), 4293–4302. https://doi.org/10.1093/cercor/bhw228

Meyer, L., Sun, Y., & Martin, A. E. (2019). Synchronous, but not entrained: Exogenous and endogenous cortical rhythms of speech and language processing. *Language, Cognition and Neuroscience*, 1–11. https://doi.org/10.1080/23273798.2019.1693050

Moulines, E., & Charpentier, F. (1990). Pitch-synchronous waveform processing techniques for text-to-speech synthesis using diphones. *Speech Communication*, *9*(5–6), 453–467. https://doi.org/10.1016/0167-6393(90)90021-Z

Nourski, K. V., Reale, R. A., Oya, H., Kawasaki, H., Kovach, C. K., Chen, H., Howard, M. A., & Brugge, J. F. (2009). Temporal Envelope of Time-Compressed Speech Represented in the Human Auditory Cortex. *Journal of Neuroscience*, *29*(49), 15564–15574. https://doi.org/10.1523/JNEUROSCI.3065-09.2009

Obleser, J., & Kayser, C. (2019). Neural Entrainment and Attentional Selection in the Listening Brain. *Trends Cogn Sci*, *23*(11), 913–926. https://doi.org/10.1016/j.tics.2019.08.004

Oganian, Y., & Chang, E. F. (2019). A speech envelope landmark for syllable encoding in human superior temporal gyrus. *Science Advances*. https://doi.org/10.1101/388280

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput Intell Neurosci*, *2011*, 156869. https://doi.org/10.1155/2011/156869

Ostendorf, M., Price, P. J., & Shattuck-Hufnagel, S. (1995). The Boston University radio news corpus. *Linguistic Data Consortium*, 1–19.

O'Sullivan, J. A., Power, A. J., Mesgarani, N., Rajaram, S., Foxe, J. J., Shinn-Cunningham, B. G., Slaney, M., Shamma, S. A., & Lalor, E. C. (2015). Attentional Selection in a Cocktail Party Environment Can Be Decoded from Single-Trial EEG. *Cerebral Cortex*, *25*(7), 1697–1706. https://doi.org/10.1093/cercor/bht355

Peelle, J. E., & Davis, M. H. (2012). Neural oscillations carry speech rhythm through to comprehension. *Frontiers in Psychology*, *3*(SEP), 1–17. https://doi.org/10.3389/fpsyg.2012.00320

Peelle, J. E., Gross, J., & Davis, M. H. (2013). Phase-locked responses to speech in human auditory cortex are enhanced during comprehension. *Cerebral Cortex*, *23*(6), 1378–1387. https://doi.org/10.1093/cercor/bhs118

Pefkou, M., Arnal, L. H., Fontolan, L., & Giraud, A.-L. (2017). θ-Band and β-Band Neural Activity Reflects Independent Syllable Tracking and Comprehension of Time-Compressed Speech. *The Journal of Neuroscience*, *37*(33), 7930–7938. https://doi.org/10.1523/JNEUROSCI.2882-16.2017

Reinisch, E. (2016). Speaker-specific processing and local context information: The case of speaking rate. *Applied Psycholinguistics*, *37*(6), 1397–1415. https://doi.org/10.1017/S0142716415000612

Ruhnau, P., Rufener, K. S., Heinze, H.-J., & Zaehle, T. (2020). Pulsed transcranial electric brain stimulation enhances speech comprehension. *Brain Stimulation*. https://doi.org/10.1016/j.brs.2020.07.011

Scott, S. K. (1998). The point of P-centres. *Psychological Research Psychologische Forschung*, *61*(1), 4–11. https://doi.org/10.1007/PL00008162

Sekihara, K., Kawabata, Y., Ushio, S., Sumiya, S., Kawabata, S., Adachi, Y., & Nagarajan, S. S. (2016). Dual signal subspace projection (DSSP): A novel algorithm for removing large interference in biomagnetic measurements. *J Neural Eng*, *13*(3), 036007. https://doi.org/10.1088/1741-2560/13/3/036007

Sohoglu, E., & Davis, M. H. (2016). Perceptual learning of degraded speech by minimizing prediction error. *Proceedings of the National Academy of Sciences*, *113*(12), E1747–E1756. https://doi.org/10.1073/pnas.1523266113

Teichmann, M., Kas, A., Boutet, C., Ferrieux, S., Nogues, M., Samri, D., Rogan, C., Dormont, D., Dubois, B., & Migliaccio, R. (2013). Deciphering logopenic primary progressive aphasia: A clinical, imaging and biomarker investigation. *Brain*, *136*(Pt 11), 3474–3488. https://doi.org/10.1093/brain/awt266

ten Oever, S., & Sack, A. T. (2015). Oscillatory phase shapes syllable perception. *Proceedings of the National Academy of Sciences*, *112*(52), 15833–15837. https://doi.org/10.1073/pnas.1517519112

Wade, T., & Holt, L. L. (2005). Perceptual effects of preceding nonspeech rate on temporal properties of speech categories. *Perception & Psychophysics*, *67*(6), 939–950. https://doi.org/10.3758/BF03193621

Zoefel, B. (2018). Speech Entrainment: Rhythmic Predictions Carried by Neural Oscillations. *Current Biology*, *28*(18), R1102–R1104. https://doi.org/10.1016/j.cub.2018.07.048

Zoefel, B., Allard, I., Anil, M., & Davis, M. H. (2019). Perception of Rhythmic Speech Is
Modulated by Focal Bilateral Transcranial Alternating Current Stimulation. *Journal of*
Cognitive Neuroscience, *32*(2), 226–240. https://doi.org/10.1162/jocn_a_01490
Zoefel, B., Archer-Boyd, A., & Davis, M. H. (2018). Phase Entrainment of Brain Oscillations
Causally Modulates Neural Responses to Intelligible Speech. *Current Biology*, *28*(3),
401-408.e5. https://doi.org/10.1016/j.cub.2017.11.071
Zoefel, B., ten Oever, S., & Sack, A. T. (2018). The Involvement of Endogenous Neural
Oscillations in the Processing of Rhythmic Input: More Than a Regular Repetition of
Evoked Neural Responses. *Frontiers in Neuroscience*, *12*.
https://doi.org/10.3389/fnins.2018.00095
11

Figure S1. Frequency of peakEnv events. Frequency of peakEnv events in slow speech (blue) and

regular speech (green). Dashed vertical lines mark average event frequency in slow (blue, 1.8 Hz) and

regular speech (green, 5.5 Hz).

Evoked low-frequency power following peakRate events

Evoked increase in power is a marker of evoked neural responses and is used to distinguish between

evoked responses and oscillatory activity. In addition to calculating the ERP to peakRate events, we thus

also tested whether band-passed power would increase after peakRate events. However, we found no

significant effects of peakRate on evoked power in theta or delta bands ($p > 0.05$, cluster-based

permutation test, data not shown). Our hypothesis that this was due to higher susceptibility of power

measures to noise was confirmed in a simulation of the evoked response model (see below).

We hypothesized that this lack of increase in power in theta or delta bands following peakRate

events might reflect the high susceptibility of power increases to noise. To assess the effect of noise on

power and phase measures, we tested the evoked response model at noise levels of 1 to 10 relative to

response magnitude. We evaluated the effect of noise onto power and IEPC in the theta band ($4 - 8$ Hz) in

the window of a single cycle for a given frequency band after event onset. The effects of noise on power

and IEPC were compared using two-sided paired t-tests at each noise level ($n = 20$ simulated responses),

with Bonferroni correction for the number of comparisons. As predicted, we found continuously large

effect sizes for IEPC even at high levels of noise, whereas the effect size for power deteriorated rapidly

with the addition of noise.

Figure S2. Effect of noise level on IEPC (black) and power (red) after peakRate events in theta

band (4 – 8 Hz) for regular speech. $* p < 0.01$

Figure S3. The duration of MEG phase locking is consistent across subjects. Single-subject data for Figure 3

Average IEPC during baseline (red), pre-event (black), 1st post-event (pink), 2nd post-event (blue), and 3^{rd} post-event

cycle (light blue) in each frequency band. IEPC increased from pre-event to 1st post-event and decreased from 1st post-

event to 2nd post-event across subjects.

Figure S4. Normalization of peakRate evoked response for contextual speech rate.

A. Evoked response for peakRate events in 1^{st}, 3^{rd}, 5^{th} peakRate magnitude quantile. Group-averaged

ERPs, error bars are ± 1 SEM. **B**. Same as A for slow speech. **C.** Quantile-quantile plot of peak evoked

response (mean (SEM)) in regular and slow speech. Peak evoked response quantile-quantile values are

close to the diagonal, indicating the similar distribution of evoked response in regular and slow speech

conditions.

2 **Table S1. Speech stimulus transcription**

Stimulus ID	Transcript
f1ajrlp1	Wanted: Chief Justice of the Massachusetts Supreme Court. In April, the S.J.C.'s current leader Edward Hennessy reaches the mandatory retirement age of seventy, and a successor is expected to be named in March. It may be the most important appointment Governor Michael Dukakis makes during the remainder of his administration and one of the toughest. As WBUR's Margo Melnicove reports, Hennessy will be a hard act to follow.
f1ajrlp2	In nineteen seventy-six, Democratic Governor Michael Dukakis fulfilled a campaign promise to de-politicize judicial appointments. He named Republican Edward Hennessy to head the State Supreme Judicial Court. For Hennessy, it was another step along a distinguished career that began as a trial lawyer and led to an appointment as associate Supreme Court Justice in nineteen seventy- one. That year Thomas Maffy, now president of the Massachusetts Bar Association, was Hennessy's law clerk.
f1ajrlp3	The author of more than eight hundred State Supreme Court opinions, Hennessy is widely respected for his legal scholarship and his administrative abilities. Admirers give Hennessy much of the credit for sweeping court reform that began a decade ago, and for last year's legislative approval of thirty-five new judgeships and three hundred million dollars to restore crumbling court houses. Despite the state's massive budget deficit, Hennessy recently urged colleagues in the bar association not to retreat from these hard won gains.
f1ajrlp4	Hennessy is the S.J.C.'s thirty-second chief justice. Holding the court system on the course he has set and plotting it's future agenda won't be an easy job for his successor.
f1ajrlp5	Attorney Haskel Kassler chairs the Judicial Nominating Council, eighteen attorneys and laypeople charged with screening applicants for vacancies on the bench. Usually the J.N.C. refers three nominees to the Governor. His top choice is rated by bar associations and grilled by the Governor's executive council. Kassler says, unlike the Federal Supreme Court, there's no litmus test on particular issues that Massachusetts high court nominees must pass.
f1ajrlp6	All but one of the Chief Justices since eighteen ninety- nine, when Oliver Wendell Holmes was appointed, came from the ranks of S.J.C. associate justices. If he sticks with

	tradition, Dukakis is likely to elevate one of his appointees to chief. That means Paul Leocos or Ruth Abrams, the only woman on the court. Another possible choice is Herbert Wilkins, a Governor Sargent appointee, and next to Hennessy, the court's most senior member. The other three associate justices were put on the bench by Governor Edward King. And many lawyers say that despite Dukakis' promise to keep the judiciary above the political fray, it's unlikely Dukakis will choose a King appointee to run the state's highest court. For WBUR, I'm Margo Melnicove.
m2b rrlp1	Massachusetts may now have the toughest drunken driving law in the nation, thanks to the Safe Roads Act that became law this week. The legislation came complete with a message from the governor to those seeking a little too much comfort and joy this holiday season and beyond.
m2b rrlp2	Under the new law the consequences of driving drunk will be swift and severe. Under the old, already tough law, first time offenders used to lose their license for thirty days. Well now they could lose it for as many as ninety. The prison sentence for a second time offender has now doubled. The law is also tougher for vehicular homicide, and it cracks down on minors who drink and drive. State trooper Joseph Holly says this tough new law will mean fewer drunken drivers.
m2b rrlp3	Well that's what supporters of the Safe Roads Act are hoping anyway, but will it work? Massachusetts got tough on drunken drivers years ago. What's going to make the difference this time around? Ralph Henksen is chief of the social and behavioral science section at the Boston University school of public health. He says research into what deters the drunken driver shows that often those who drink and drive are risk takers. Even when they're sober they're more likely to run red lights and shun the seat belt. If they think they can drink and drive and get away with it, they'll try.
m2b rrlp4	And in fact the state is not planning on putting more police on the road. State officials say there will be the same levels of enforcement. B.U.'s professor Henksen believes the key to the new law's success is its enforcement.
m2b rrlp5	The state Safe Roads Act is only a few days old, but it's already facing a challenge. The man who directed the successful effort last election to get the mandatory seat belt law stripped from the law books is now vowing to try to repeal part of this law. Lynn activist Chip Ford has yet to give a name to his committee, but he's already set its goal: to repeal what is the most controversial clause in the Safe Roads Act, the per se clause, which allows someone to be punished for drinking and driving before they've even been tried in court.

	Under the per se part of this new law, if you're arrested, take a breath test, and your blood content level registers point one oh or higher, you can automatically lose your license for ninety days. And that's just at your arraignment. Then you still got to face DWI charges at a trial. State officials say about a quarter of all those arrested for drunken driving are repeat offenders. They say the per se clause will get those people off the road sooner. Chip Ford says it may indeed get them off the road, but it will do it at the expense of their civil rights.
m2b rrlp6	Ford and other opponents of the per se clause say it relies exclusively on the breath test, and they say breathalizers are simply unreliable. John Tarantino is a Providence attorney who specializes in drunken driving cases.
m2b rrlp7	Tarantino says because of people's physiological differences, the breathalizer test will always be inaccurate on about fourteen percent of the population. Those people will register as having more alcohol in their blood than they actually do. But state officials aren't convinced that's a problem. While they don't say breathalizers are one hundred percent accurate, they do say they are extremely reliable, and if they err at all it's on the conservative side, the side favorable to the driver. As for those who say that taking away someone's license based on the breathalizer results is tantamount to trial by machine, state officials say that simply isn't so. Losing one's license under the per se clause is not a criminal offense. You won't get a criminal record for it. It is a civil infraction and one that they believe will be a key deterrent to the drunken driver. If the per se clause is tested in court, the state says it will pass constitutional muster. And if opponents gather enough signatures to make it a ballot question in nineteen eighty-eight, state officials predict voters will keep it intact.

2   **Table S2. Comprehension questions**

Stimulus ID	Question	Answer
f1ajrlp1	What is the mandatory retirement age for chief justice?	**a: 70** b: 30
f1ajrlp2	What did Michael Dukakis promise in his campaign?	a: To legalize medical Marijuana, **b: To de-politicize judicial appointments.**
f1ajrlp3	How many laws were passed under Hennessy?	**a: more than 800**

			b: more than 8000
f1ajrlp4	How many chief justices were there before Hennessy?		**a: 31** b: 10
f1ajrlp5	How many nominees for the court are passed to the governor?		a: 5 **b: 3**
f1ajrlp6	How many chief Justices so far were not from the ranks of SJC associate justices?		a: 5 **b: 1**
m2brrlp 1	What is the name of the new law in Massachusetts?		**a: Safe Roads Act**, b: Seatbelt act
m2brrlp 2	For how long do first-time offenders lose their license under the new law?		a: 10 days, **b: 90 days**
m2brrlp 3	Research shows that those that drink, and drive are		**a: risk-takers**; b: not aware of the risks
m2brrlp 4	Are there more police on the road after the new law?		a: Yes, **b: No**
m2brrlp 5	How many of those arrested for drunk driving are repeat offenders?		**a: a quarter** b: More than half
m2brrlp 6	Is the breathalyzer test reliable?		a: yes, **b: no**
m2brrlp 7	Does one get a criminal record for losing their driver's license?		a: yes, **b: no**