First-in-class matrix anti-assembly peptide prevents staphylococcal biofilm *in vitro* and *in vivo*

Rafael Gomes Von Borowski[1,2], Sophie Chat[1], Rafael Schneider[1,2], Sylvie Nonin-Lecomte[3], Serge Bouaziz[3], Emmanuel Giudice[1], Aline Rigon Zimmer[2], Simone Cristina Baggio Gnoatto[2], Alexandre José Macedo[2,4,*], and Reynald Gillet[1,*]

[1] Univ. Rennes, CNRS, Institut de Génétique et de Développement de Rennes (IGDR), UMR6290, Rennes, France; [2]Programa de Pós-Graduação em Ciências Farmacêuticas, Faculdade de Farmácia, Universidade Federal do Rio Grande do Sul, 90610-000 Porto Alegre, Brazil; [3]Université de Paris, CNRS, CiTCoM (Cibles Thérapeutiques et Conception de Médicaments) UMR 8038, Faculté de Pharmacie, 75006 Paris, France; [4]Centro de Biotecnologia da Universidade Federal do Rio Grande do Sul, 91501-970 Porto Alegre, Brazil

[*]To whom correspondence may be addressed. Email reynald.gillet@univ-rennes1.fr or alexandre.jose.macedo@gmail.com.

Abstract

Staphylococci are pathogenic biofilm-forming bacteria, source of multidrug-resistance and/or – tolerance causing a broad spectrum of infections. These bacteria are enclosed in a matrix that allows them to colonize medical devices such as catheters and tissue, and which protects against antibiotics and immune systems. Advances in antibiofilm strategies for targeting this matrix are therefore extremely relevant. Plants are constantly attacked by a wide range of pathogens, and have protective factors such as peptides to defend themselves. These peptides are common

components in *Capsicum* peppers (CP). Here, we describe the development of CP bioinspired peptide "capsicumicine". We demonstrate that capsicumicine strongly prevents methicillin-resistant *S. epidermidis* biofilm *via* a new extracellular "matrix anti-assembly" mechanism of action. Catheters pre-coated with capsicumicine decreased *S. aureus* colonization leading to the attenuation of infection, decreasing mice systemic infection. Capsicumicine is the first-in-class non-antibiotic, carbohydrate-binding peptide.

Keywords: antibiofilm, biofilm, matrix, peptides, anti-assembly, resistance, tolerance.

Introduction

Antimicrobial failure is a worldwide challenge, currently addressed by a WHO global action plan (*1*). A lack of new antibiotics and the inappropriate use of older treatments mean that multidrug-resistant strains are increasing (*2*). This process is favored by biofilm development, and microorganisms enclosed in biofilm matrix have antibiotic resistance that is up to 1000 times higher than planktonic ones (*3*). This makes the matrix itself an important target for biofilm control. Biofilms are organized microbial clusters made of a self-assembled matrix that usually attaches to a surface, whether abiotic (medical devices such as catheters, teeth, etc.) or biotic (host tissues, mucus, chronic wounds, etc.) (*4, 5*). Since the bacteria are embedded into this matrix, they are harder to treat because of their increased tolerance and resistance to antibiotics, disinfectants, and host defenses (*6, 7*). Other advantages over planktonic forms include physiological and biochemical changes, beneficial quorum sensing, higher (up to 100 times) mutation rates, and persister cell development (*8-10*).

Staphylococci are the most frequent sources of nosocomial infections, particularly *S. epidermidis* and *S. aureus*. While *S. aureus* expresses many virulence factors such as toxins and proteases, for *S. epidermidis* the formation of biofilm is the most important mechanism in infection development (*11*) . *Staphylococcus epidermidis* is the most frequent coagulase-negative staphylococcal (CoNS)

infection-causing disease (*12*), surviving on various surfaces for months (*13*). It is present in 30% of

health care-involving bloodstream infections, and is significantly associated with medical device

infections, including 15-40% of prosthetic valve endocarditis (*14*) and 30-43% of prosthetic

orthopedic device infections (*15*). More than 150 million intravascular catheters are used per year in

the USA, and there are about 250,000 catheter-related infections (*16, 17*). These bacteria are

developing antibiotic multi-resistances such as elevated glycopeptide minimal inhibitory

concentrations (*18*), and 73-88% of isolates display resistance to oxacillin, fluoroquinolones,

macrolides, clindamycin, and trimethoprim/sulfamethoxazole (*19-21*).

In this circumstance, the extracellular matrix is a complex physicochemical barrier representing one

of the biggest challenges in microbial treatments (*4*). Therefore, the development of antivirulence

strategies such as antibiofilm agents is crucial for the current antibiotic crisis, and peptides are an

increasingly arsenal for controlling pathogenic biofilms (*22-24*).

Here, we describe the discovery of the antibiofilm peptide capsicumicine, inspired by natural

peptides from the seeds of the red pepper *Capsicum baccatum.* We elucidated its mechanism of

action (MoA), the matrix anti-assembly (MAA). Capsicumicine is the first-in-class non-antibiotic

peptide displaying this extracellular MoA. Notably, we report an *in vivo* anti-infective proof of

concept towards to the use of capsicumicine for complementary treatment of infectious diseases.

Results

Capsicumicine prevents biofilm formation without antibiotic activity. We synthesized three

peptides inspired by a natural antibiofilm fraction previously identified from *Capsicum baccatum* var.

pendulum pepper seeds (*25*): P1 (RVQSEEGEDQISQRE), P2 (RAEAFQTAQALPGLCRI), and P3

(RSCQQQIQQAQQLSSCQQYLKQ). To find the most active one, we exposed these compounds to

strong biofilm-forming *S. epidermidis* RP62A (ATCC 35984). After 24 h, crystal violet was used to

quantify the remaining biofilm (Fig. 1*A*). The P3, named "capsicumicine," was the most active with

particularly strong antibiofilm activity. Biofilm decreases were observed at all tested concentrations,

but especially at 10 µM. There, biofilm was reduced by over 91%, independently of cell growth

inhibition (Fig. 1*B*). To examine the effects of capsicumicine on growth, we checked *S. epidermidis*

colony-forming unit (CFU) counts after peptide exposure. As expected, the CFUs were unchanged by

capsicumicine, so the peptide's biofilm inhibition is not due to bactericidal activity (Fig. 1*B*). To verify

the interactions between capsicumicine and established matrices, we exposed a pre-existing *S.*

epidermidis biofilm to a single concentration (100 µM) of the peptide. At that concentration,

capsicumicine accounts for 15% of the disruption of pre-existing biofilm (Fig. S1).

Fig. 1. Antibiofilm activity of bioinspired peptides and cytotoxicity. *(A)* Antibiofilm activity of peptides
P1, P2, and P3 (capsicumicine) at 1, 10, and 100 µM. Quantification of *Staphylococcus epidermidis*
(ATCC 35984) biofilm and growth were done at an optical density; bacteria without peptide exposure
(Control) and the antibiotic control, 16 µg/mL rifampicin (Rif.). Student's *t*-test: *, $p \leq 0.05$; **, $p \leq$
0.01. *(B)* Antibiofilm activity and growth of capsicumicine at 10 µM. Yellow represents 100% of
formed biofilm and red represents the remaining biofilm after exposure to capsicumicine. Growth

was verified using colony-forming units (CFUs). *(C)* Gene expression (mean log fold changes ±
standard errors of the means) of the encoding genes involved in *S. epidermidis* biofilm formation as
compared to the planktonic (grey) and biofilm controls (black), with the *ssrA* gene used as a
reference. The group exposed to capsicumicine is shown in white. CWA, cell wall-anchored proteins.
D) Capsicumicine cytotoxicity evaluation in representative human cell lines shown via automated
image-based cellular content analysis. Cell counts are presented as residual cell percentages (%)
compared to the average of the DMSO control, with water control also shown (grey bars). The black
bars on the left show cytotoxic controls (roscovitine, doxycycline, and taxol), while the white bars are
cells exposed to 10 µM of capsicumicine.
97

To study the peptide's possible mechanisms of action, we selected several genes involved in different

stages of biofilm development (*atlE*, *aap*, *agrC*, *icaA*, *leuA*, *saeR*, *saeS,* and *sarA*); primers are listed in

Table S1. Fold changes were analyzed by quantitative real-time PCR (qRT-PCR). Since exposed

bacteria remain planktonic, we compared their relative gene expressions to planktonic control cells.

For all tested genes, capsicumicine-exposed cells show the same fold changes as the control (Fig. 1*C*).

Capsicumicine is not cytotoxic in mammalian cells. To ensure that capsicumicine is safe before

propose *in vivo* trials, we verified its cytotoxicity in seven different representative human cell lines.

We used automated image-based cellular content analysis and found that capsicumicine-treated

cells perform exactly the same as untreated controls, displaying no cytotoxicity (Fig. 1*D*).

Independently of cell interactions, capsicumicine impairs biofilm attachment, aggregation, and

accumulation. To explore its activity during the first stages of biofilm development, we analyzed

biofilm cultures on polystyrene coupons with and without capsicumicine after 1, 4, and 24 h.

Scanning electron microscopy (SEM) analysis shows that bacterial attachment decreases after 1 h of

capsicumicine exposure, with biofilm accumulation and cell aggregation profiles strongly reduced

after 4 and 24 h (Fig. 2*A*). This demonstrates that capsicumicine prevents *S. epidermidis* coupon

adhesion, and notably this activity still occurs after 24 h incubation. In order to localize the peptide,

we used capsicumicine conjugated to fluorescein isothiocyanate (capsicumicine-FITC) and confocal

fluorescence microscopy (CFM). Analysis of the CFM images showed that capsicumicine-FITC stays

associated with extracellular matrix components, not entering into bacterial cells or the walls or

membranes (Fig. 2*B*).

Fig. 2. Biological characterization of capsicumicine by microscopic approaches. (*A)* SEM images of
polystyrene coupons after 1, 4, or 24h of culture with *Staphylococcus epidermidis* (ATCC 35984). *Top:*
peptide-less biofilm control; *bottom:* cultures exposed to 10 µM capsicumicine. Magnification x500,
with insets at x5,000; scale bars, 10 µm. (*B*) CFM images: *(left) S. epidermidis* after exposure to 10 µM
capsicumicine-FITC, with the matrix in green (fluorescence), and the bacterial cells in black (no
fluorescence); (*right*) control, *S. epidermidis* after exposition to pseudonajide-FITC, an antibacterial
peptide, this time showing the matrix in gray (no fluorescence) and bacterial cells in green
(fluorescence). White arrows indicate matrix content. Scale bars, 5 µm. *(C)* CFM images: *(left) S.*
epidermidis after exposure to 10 µM capsicumicine-FITC; *(center)* after exposure to concanavalin A
conjugates; and (*right*) their colocalization. Scale bars, 5 µm. (*D-G*) These images explore the
organizational state of *S. epidermidis* biofilm after 24 h in the presence (*right*) or absence (control,
left) of capsicumicine. (*D*) Macroscopic examination by pictures from the bottom of 24-well plates.
The "sterile control" shows no bacteria or biofilm formation; the "biofilm control" shows
homogenous adhered layers of bacteria; and "capsicumicine" shows non-adhered bacteria but
agglutinates; scale bar, 5 mm. (*E*) TLM images show the biofilm control with overlapping attached
bacterial clusters, while the capsicumicine-exposed culture shows agglutinated non-adhered cells;
scale bar, 20 µm. (*F*) SEM images show the control with dense globular-like matrix features, while the
capsicumicine-exposed culture shows fibrillary branch-like oligomer structures; scale bar, 200 nm.
(*G*) TEM images of the biofilm control show denser assembled structures, while the capsicumicine-
exposed displays thin fibrillary oligomer structures; scale bar, 200 nm. Arrows indicate the matrices.

Capsicumicine disturbs *S. epidermidis* matrix assembly. Since capsicumicine's antibiofilm activity

was not associated with direct bacterial interactions or gene expression modulation, we used various

microscopic approaches to investigate the interactions between the peptide and the extracellular

matter. In the biofilm control, macroscopic observation shows a homogenous whitish adhered layer

covering the walls and bottoms of the well (Fig. 2D, middle), but when capsicumicine is present, we

see whitish flocculent non-adhered heterogeneous agglutinates (Fig. 2*D*, right). Transmitted light

microscopy (TLM) images match with these macroscopic observations (Fig. 2*E*). Crucially, SEM and

TEM technique yield ultra-structural descriptions that support these results, with the control biofilm

matrix showing denser assembled globular-like structures, while the capsicumicine-exposed matrix is

less dense and has thin fibrillary branch-like structures (Fig. 2*F* and *G*). However, cellular

morphologies are not different from the controls. Therefore, different imaging techniques prove that

matrix assembly changes when capsicumicine is present.

Capsicumicine interacts with matrix polysaccharides. To explore the peptide's affinity with these

major matrix components, we exposed *S. epidermidis* cultures to capsicumicine (-FITC; green) and

saccharides staining, then analyzed them all by CFM. We used concanavalin A and calcofluor to

selectively target matrix saccharides (blue). Amounts of capsicumicine-FITC appear exclusively on the

matrix (Fig. 2*B* left; S2*B*) and its colocalization indicates the interaction between both marked

elements (Fig. 2*C*, right; S2*A*). Visualization was done by individual (Fig. 2*C*, left and center) and

colocalization fluorescence (Fig. 2*C*, right). The peptide control was pseudonajide-FITC, an

antibacterial peptide; it showed green fluorescence in the cells but not in the matrix (Fig. 2*B* right;

S2*C*). To explore whether capsicumicine features are compatible with carbohydrate-binding modules

(CBMs), we performed a BLAST and amino acid alignments between capsicumicine and chitin-,

chitosan-, and polysaccharide intercellular adhesin (PIA), PIA-binding proteins (Table S2). UniProt

tools (*26*) and CAZy information crossing (*27*) showed that capsicumicine does in fact present CBM

homology with all tested proteins (Fig. S3).

Capsicumicine shifts Staphylococcal synthetic matrix. To confirm these interactions in the absence

of bacteria metabolic or regulatory influences, we adapted a model of artificial staphylococcal biofilm

assembly to test it. Briefly, we monitored the real-time molecular self-assembly (RTMSA) reaction by

measuring the optical density at 600 nm (OD_{600}) as a function of time with or without capsicumicine.

OD increases when capsicumicine is present, which shows that the molecular self-assembly reaction

is quicker overall (Fig. 3*A*). The profiles of assembled matrices are visually different, with larger

agglutinates in the presence of capsicumicine, although for both controls are similar (Fig. 3*A*).

Remarkably, these profiles are comparable to those previously observed in the presence of bacteria

(Fig. 2*D*).

Fig. 3. Molecular interactions between capsicumicine and target-saccharides. *(A)* RTMSA curves of
artificial staphylococcal biofilm assembly. Optical densities (OD_{600}) were recorded as a function of the
time in the presence of capsicumicine (blue dots) or PA1 as peptide control (orange dots) and the
synthetic matrix without peptides (black dots); each reaction tub is shown under the graph. *(B-D)*
NMR titration of capsicumicine by chitosan at 600MHz, 10°C. *(B)* From the bottom to the top:
increasing concentrations of soluble chitosan [< 0.5 eq. and 8% of volume variation] were gradually
added to capsicumicine water solution. Specific line broadenings and frequency shifts are shown in
the NMR spectrum for some amide protons upon titration; white bullets for terminal R1 and Q22 and

black bullets for S2, C3 and Q4. *(C)* Assembled chitosan gradually added to soluble capsicumicine [0.5
eq.]; an overall line broadening was observed. *(D)* Assembled chitosan suddenly added to soluble
capsicumicine [0.5 eq.]; the orange spectrum was recorded after few minutes of stirring. *(E)*
Superposition of capsicumicine 1D NMR and TOCSY spectra immediately after sample preparation
(blue) and chitosan [10 eq.] addition (red) and; capsicumicine structure predicted by Phyre2 server;
according to NMR results the aggregation nucleating regions are shown as orange triangles and
favored interacting region with sugar moieties is shown in grey.

To evidence the interactions between capsicumicine and target-saccharides we performed NMR

titration experiments using chitosan as a mimic of the matrix poly-N-acetyl glucosamine (*PNAG*). The

evolutions of the capsicumicine NMR spectra upon chitosan addition were monitored. Reference 1D

proton NMR spectra were recorded at all tested conditions. Chitosan was gradually added from a

concentrated solution and spectra changes were monitored. This solution remained clear during the

7 days of recording. The addition of soluble chitosan drives noticeable changes of the resonance

frequency of some of the amide protons mainly in the N-terminal part of the peptide < S2, C3, Q4

and R1 (not shown) > (Fig. 3*B*). Conversely, to < Y19, L20 and K21 NH and the Y19 > aromatic proton

resonances, between 6.8 and 7.2 are not sensitive to the presence of chitosan (Fig. 3*E*). The

concentration variation of chitosan was estimated to less than 0.5eq at the end of the titration. This

small quantity of chitosan drives spectral modifications in the N-terminal part of capsicumicine and

strongly supports the interaction between both partners. The important broadening observed after 1

day shows that the peptide structure keeps evolving over time and this is not due to cysteines'

oxidation (Fig 4*D*, *E*). These experiments were repeated adding gelled chitosan instead of soluble

chitosan, mimicking an assembled matrix. It does not drive any significant spectral changes, except

for a general line broadening which probably arises from an increased viscosity of the solution (Fig.

3*C*). The suddenly addition of gelled chitosan pellets in the NMR solution, does not modify the

spectrum either (Fig 3*D*).

Structural requirements toward to molecular interactions with target-saccharides. To better

understand the mechanisms involved in capsicumicine bioactivity we step toward the knowledge of

its conformation in solution and molecular interactions with matrix representative saccharides. To

monitor the conformation of free capsicumicine in response to time and temperature we first used

Circular Dichroism (CD) and Nuclear Magnetic Resonance (NMR) spectroscopies. Spectral

deconvolution using the CDSSTR algorithm discloses a helical structure (about 43-44%) with a non-

negligible proportion of β strands (31-33%) and unfolded (about 20%) (Table S3). The structure

stability was assessed over 5 days recording the CD spectra at 5°C (Fig. 4A). The main structure

observed over the first 24h is helical. After 5 days, deconvolutions show a slight decrease of the

helical content. Furthermore, on the fifth day, CD spectra were also recorded at 15 and 30°C (Fig.

4*B*). The helical content decreases to 18-23% while the β strand and turn mean proportions rose

around 37% and 23%, respectively. The proportion of unfolded structures also increased. After 2

days at room temperature, a new CD spectrum was recorded at 30°C. The helix proportion became

less than 15 % in favor of β strand and unfolded conformations (respectively 38-41% and 31-35%).

During these days, the peptide solution remained clear, showing no macroscopic signs of

aggregation.

The 1D and 2D NMR spectra of a 0.3mM freshly prepared solution of capsicumicine were recorded at

10°C, pH 5.0. All amide resonances were unambiguously assigned using TOCSY and NOESY spectra,

except for Q9 and Q11 residues (Fig. S4). The spectral dispersion and the spreading of the amide

proton resonances disclose that the peptide is folded with one conformation in these conditions. In

accordance with CD spectra, a second set of NH resonances appears as a function of time (Fig. 4*C, D*),

revealing conformational changes in the slow exchange regime on the NMR time scale. After 5 days,

the most important chemical shift perturbations observed on the TOCSY spectrum are clustered

around the two cysteines apart from the two terminal residues (boxed cross-peaks, Fig. 4*E*).

Interestingly, I7 located one helix turn apart C3 is among the most sensitive residues to the

conformational perturbation. This is due to a disulfide bridge (DB), since no reducing agent was

added. Consequently, large proton chemical shift perturbations (CSP) would be expected for every -

NH amino acid and this was not detected (Fig. 4*D*, *E*). The prevalence of the second conformation

reaches more than 50% after 7 days (Fig. 4C, top). The solution remained clear overall the spectra

recording time.

Fig. 4 Capsicumicine conformational changes. CD spectras *(A)* Evolution at 5°C over 5 days; *(B)*
Evolution *vs* time and temperature. *(C)* NMR spectra recorded at pH 5.0, 0.3mM and 10°C
immediately after extemporaneous preparation (bottom); a second conformation appears over the
time corresponding to the second set of amide resonances (stars). *(D)* Superposition of capsicumicine
TOCSY spectra immediately after sample preparation (red) and at day 5 (brown). The most important
chemical shit perturbations between the original and the new conformations are boxed and
clustered around the two cysteines. *(E)* Proton chemical shift perturbations (CSP) computed from the
TOSCY spectra are displayed in the box.

Capsicumicine attenuates the dynamics of *S. aureus* (Xen36) infection on central venous catheter (CVC). We performed a translational proof-of-concept evaluating long-term bacterial biofilm and related infection in mice implanted with capsicumicine pre-coated CVC. These CVCs were previously coated using an immobilization polymer ("hydrogel") encompassing capsicumicine. This coating confers amorphous and biocompatible surfaces to CVC (Fig. 5*A*). They were first validated *in vitro*, decreasing ≥51 % of *S. aureus* colonization (Fig. 5*B*). Then, bacterial development was evaluated *in vivo* by bioluminescence imaging and bacterial load of harvested CVC (Fig. 5*C-F*). Two days (D2) after *S. aureus* systemic infection, 40% of the control group (CVC-Hydrogel) presented high bioluminescence signal (red zones) related to ROI against none in the treated group (CVC-Hydrogel + capsicumicine), (Fig. 5*C*). Four days (D4) after infection, 75% of the control presented high ROI red zones against 20% in the treated group (Fig. 5*C*). In addition, at D4, 1 animal was found dead in the control group. Therefore, bioluminescence quantifications show that the treated group decreased 56% (D2) and 54% (D4) of the total flux compared to the control (Fig. 5*D*). This trend was also observed on the CFU load at D4 (Fig. 5*E*). After image acquisitions (D4), 2 animals / group were euthanized due to ethical criteria. At the end of the experiment, 7 days after infection (D7), the treated group showed a decrease of 86% of bacterial load compared to control (Fig. 5*E*). Macroscopic observation revealed that one animal from the control presented several organs with necrosis (liver, spleen, intestine, kidneys and bladder). Finally, the treated group increased survival rate of 50% at D7 compared to control (Fig. 5*E*).

Fig. 5. Assessment of capsicumicine pre-coated central venous catheter (CVC) in mice (SKH1) infected
by *Staphylococcus aureus* Xen36. *(A-B) In vitro* CVC development; *(A)* From the top to the bottom: a
picture of a up to half capsicumicine pre-coated CVC (whitish), scale bar 1 cm; SEM image of pre-
coated CVC, scale bar 0.5 mm and its magnification, scale bar 1 μm and; a compilation of images of
fluorescence microscopy of pre-coated CVC (capsicumicine-FITC in green); *(B)* Bacterial load on CVC.
(C-F) In vivo CVC assessment; *(C)* IVIS images (radiance-thresholded and smoothed with a
bidimensional Gaussian filter) of ventral position for each mouse at each time point; the control
group (CVC-Hydrogel) is shown in the top and the treated (CVC-Hydrogel + capsicumicine) in the
bottom; white circles delimitate the CVC localization; the luminescence scale is presented in
radiance. *(D)* Bioluminescence quantification (ROI mean signals ± SD) is shown as photons total flux
[p/s] for each group at each time point. *(E)* Bacterial load (CFU/mL) on harvested CVC for each mouse
at each time point (day 4 and 7 after infection) and; *(F)* The Kaplan-Meier survival curve (% of
survival) at each group, during 7 days after infection. All bioluminescence data and analysis were
performed using IVIS Spectrum of Perkin Elmer. All experiments were conducted in accordance with
ethical committee and the French authorities (n = 5 / group). Bioluminescence images are presented
with a defined pseudo-color scale to visualize the intensity of signals emitted.

Discussion

Bioinspired peptides are increasingly being explored as alternative biofilm controls, and become

important allies in the fight against bacterial tolerance and resistance (*28, 29*). As shown here,

capsicumicine, a peptide derived from *Capsicum baccatum* red pepper seeds, possesses strong

antibiofilm activity *in vitro* and *in vivo*. Capsicumicine prevents the establishment and maintenance

of biofilm architecture through a new mechanism of action the "matrix anti-assembly" (MAA). MAA

differs from matrix *dis*assembly (*30*) as instead of de-structuring pre-established matrices it acts on

the initial phase of assembly preventing its correct structuration. In fact, established biofilms are

harder to treat than initial biofilms, because they have more complex structures (*31*), and increased

structural complexity means that more energy is required for disassembly (*32*).

Bacterial surface proteins can passively interact with surfaces such as medical devices, generating an

initial and reversible adhesion after electrostatic and hydrophobic interactions, Van der Waals forces

and others (*33*). Bacteria will then require extracellular matrix production in order to remain

attached after these weak interactions (*34*). During this process, physicochemical interactions drive

molecular and colloidal matrix self-assembly, establishing a chain of dense architecture that results in

stable adhesion (Fig. 6*A*) (*35*). In contrast, capsicumicine interacts with matrix saccharides and

modifies the self-assembly chain, resulting in a less dense and nonfunctional matrix, and impaired

biofilm formation (Fig. 6*B*).

Fig. 6. The matrix anti-assembly (MAA), antibiofilm mechanism of action (MoA) identified in capsicumicine. *(A)* Untreated biofilm development on abiotic surfaces: planktonic cells interact passively with the surface and start the extracellular matrix production. Initial adhesion is established, after which matrix self-assembly occurs, leading to irreversible biofilm structuring and adhesion. *(B)* Biofilm development in the presence of capsicumicine shows the MAA MoA: planktonic cells are still able to interact passively with the surface. At that point, capsicumicine acts, attracting extracellular saccharides and shifting the matrix self-assembly; *(C)* Capsicumicine (red) tends to organize itself in beta leaves favoring the interactions with matrix target-saccharides (yellow). These interactions occur between the non-catalytic carbohydrate binding modules of the peptide and the saccharides. This peptide supra organization can compete with matrix assembly, preventing/destroying intra matrix h-bonds over a large range of residues. Consequently, the MAA decreases adhesion and aggregation, forcing bacteria to remain planktonic. The arrows indicate reversible (white background) and irreversible stages (black background) during biofilm formation.

Capsicumicine's amino acid sequence contains several residue characteristics for recognizing putative chitin-binding domains, including polar and hydrophobic residue (45% of Q) and cysteines (*36-39*). The chitin-binding domain is a well-conserved amino acid stretch that binds specifically to N-acetyl glucosamine, a homologous structure of PIA (*27, 40, 41*). Additionally, non-catalytic carbohydrate binding modules (CBMs) are contiguous amino acid sequences with a discreet fold displaying carbohydrate-binding activity. In this context, capsicumicine displays homologies with all tested CBM-proteins, notably with icaA protein that is a PIA synthetase from the same *S. epidermidis* strain studied here (Fig. S3 and Table S2).

According to the CAZY web site, there are 86 CBMs defined families up to now (http://www.cazy.org/Carbohydrate-Binding-Modules.html). CBMs exhibit different folds, with secondary structures ranging from mainly β-sheet based organization to a mixture of α-helices and β-sheets and usually display a high content of loops and unfolded parts. Likewise, capsicumicine secondary fold is a mixture of α-helix, β-sheet and unfolded regions, with conformational time and temperature interconversion in favor of β-sheet and unfolded segments. The β-sheet sub-structure observed in the CD concerns the Nter part of the peptide according to the chemical shifts of the corresponding amino acids (between 8.6 and 9.0). The TOCSY spectra analyses disclose that the conformational changes observed through the time are originated around the two cysteines, possibly involved in intermolecular disulfide bonds. This disulfide bond is possibly intermolecular since the

formation of an intramolecular disulfide bond is only possible by bending the structure to get both

cysteines close to each other. The size of different conformers and potential polymers are limited as

shown by the NMR peaks of free peptides which are not broadened at day 7 (Fig. 4*C*). As a result,

these conformational changes may correspond to structural requirements toward to interactions

with staphylococcal matrix.

Antibiofilm, matrix anti-assembly (MAA) mechanism of action

This proposed mechanism is based on a set of intermolecular *(i)* and cooperative *(ii)* forces triggered

by capsicumicine which then disrupt matrix assembly:

(i) Intermolecular forces

The staphylococcal matrix is composed mainly of polysaccharides, but also contains proteins (AtlE,

Aap, Empb), teichoic acids, and extracellular DNA (*42*). These saccharides are PIA/PNAG,

homoglycans of beta-1,6-linked 2-amino-2-deoxy-D-glucopyranosyl residues. The matrix contains

positively-charged amino groups (PNAG), as well as negative charges from O-succinylation, which

confers electrical charge lability on the matrix.

Supported by NMR observations, capsicumicine interacts with free chitosan, a PNAG mimetic, and

then switches to a higher molecular weight organization (Fig 3*B*). The line broadening observed with

the peptide-chitosan mixture after one day evidence the aggregation of chitosan particles with

capsicumicine over the time as schematized on Fig 6. Likewise, CSPs analysis reveals that chitosan is

able to bind preferentially capsicumicine N-terminus (Fig 4*E*) before intermolecular disulfide bridges

formation and the relative conformational transition. This is supported by the fact that chitosan is

sub-stoichiometric at the end of the NMR titration and because the capsicumicine spectrum of the

mixture is different from the spectra of the free peptide (Fig 4*C*). These interactions may proceed by

both hydrogen-bonding between the hydrophilic Q's lateral chains and hydrophobic/stacking

interactions involving I and L residues. According to NMR, the N-terminal part of the peptide is

probably the favored interacting region with the carbohydrate moieties. Thus, these data support

the MAA model in which the peptide supra organization would compete with matrix self-assembly

preventing intra h-bonds over a large range of residues. However, the NMR results obtained with

assembled chitosan, mimetizing a preformed biofilm, demonstrate that capsicumicine does not

interact with pre-assembled matrix (Fig 3*C*,*D*) such as demonstrated in the eradication test.

Additionally, the amino acids in capsicumicine are mostly neutral, counting 13 Q and 3 S although,

due to Q's amide group and S's hydroxyl group, they can generate electronegative (dipole-dipole)

zones. This high electronegativity suggests that capsicumicine may interact with the positive free

charges of the PIA/PNAG. Thus, these polar non-ionic forces perform moderate interactions with the

polysaccharides. Since strong interactions such as ionic forces may trigger unwanted effects including

matrix repulsion or sequestration (*43, 44*), the moderate interactions of capsicumicine seems to be

ideal.

(ii) Cooperative polymerization forces

In living systems, biomolecules perform their functions in the presence of various macromolecules of

different shapes and sizes, and these interactions can include cooperative polymerization forces such

as depletion forces (DFs) and subsequent molecular crowding (MC) (*45, 46*). These forces are

noncovalent and non-specific physicochemical interactions leading to bridging, aggregation, and

rheological variations (*47, 48*), as is observed in the presence of capsicumicine (Fig. 2*D-G* and 3*A*),

and also may contribute to MAA. In a suspension containing different molecules, DFs are the

pressures exerted by small particles, which in turn cause attractive forces between the

macromolecules. DFs are only expressed in crowded environments like biofilms, driving the assembly

and final shape of these structures (*49*). Taking together NMR and CD experimental results as well as

the TANGO and SALSA analysis (Fig. S5*B*), we can picture that capsicumicine can self-organize to form

extended structures both by disulfide bridges and β-sheet mediated fibrils formation. We propose

that capsicumicine-associated DFs may coagulate in polymer solutions forming fibers and parallel

bundles (*50*), explaining the observed branch-like profile (Fig. 2*D-G*) (*51*). This higher organization

should facilitate the interaction with the saccharides of the matrix preventing their pattern assembly

and resulting in flocculation (*52*). In a same way, MC in macromolecule solutions is characterized by

a decrease in accessible volume due to high macromolecule concentrations, as well as attractive and

repulsive forces between them (*45, 53*). Some molecular crowders modulate refolding kinetics and

decrease competing aggregation and segregation (*54*). In this way, capsicumicine enhances the

aggregation kinetics (Fig. 3*A*), acting as an agent of MC, decreasing entropic forces and leading to

segregation (*55*).

Thus, capsicumicine' structures are chemically enable to interact with staphylococcal matrices

saccharides. These combined intermolecular and cooperative forces disturb matrix self-assembly at

both molecular and colloidal levels. Consequently, matrix functionality shifts and antibiofilm activity

occurs. In other hand, planktonic microorganisms are more available for the innate immune system

to recognize and clear them (*56*). Furthermore, non-antibiotic activity appears less susceptible to the

development of bacteria resistance because these microorganisms are under less evolutionary

pressure then when exposed to conventional antibiotics (*57, 58*).

Finally, we demonstrated that capsicumicine strongly prevents biofilm formation for the most

frequent bacteria related to nosocomial infections, *S. epidermidis* and *S. aureus*. We first confirmed

that capsicumicine was not cytotoxic thus being safe for *in vivo* tests. Then, we trialed the peptide in

a translational pre-clinical model that mimics medical device-related infection. As a result, pre-

coated CVCs enhanced mice survival decreasing *S. aureus* colonization and consequently attenuated

the infection.

Although there are no antibiofilm drugs available here we relate the discovery of the first-in-class

carbohydrate-binding peptide as a promising candidate for complementary drug/treatment of

infectious diseases. In particular, we elucidated its antibiofilm mechanism of action, matrix anti-

assembly (MAA) and validated a proof of concept towards to an *in vivo* application.

Materials and Methods

Peptides. All peptides were synthesized by Biomatik and ProteoGenix at purity grades over 95% in salts suitable for cell culture. For the assays, the peptides were all solubilized in ultra-pure sterile water.

Bacterial strains and growth conditions. *Staphylococcus epidermidis* ATCC 35984 was grown overnight on blood agar (Thermo Scientific Oxoid PB5039A) at 37 °C. Oxoid LB agar was used for the colony-forming unit (CFU) assay. The other assays were done using a bacterial suspension of 3×10^8 CFU/mL in tryptone soya broth (TSB, Oxoid) or 0.9% NaCl.

Biofilm formation. At least three technical and biological replicates were done for each assay of 1, 10, or 100 μM peptide concentrations. *Biofilm formation inhibition:* A protocol adapted from Trentin et al (*59*) was used, with crystal violet in 96-well BD Falcon polyvinyl chloride (PVC) microtiter plates. The cell-bound stains were solubilized with absolute ethanol (Sigma-Aldrich), and absorbance was measured at 570 nm using a BIO-TEK PowerWave XS plate reader. The biofilm formation control represents 100% of biofilm formation. *Biofilm eradication:* Biofilm was pre-formed as described above for 24 h at 37 °C without treatment. Afterwards, the wells were washed to remove planktonic cells, peptide solutions and controls were added, and all were incubated for 24 h. Biofilm eradication was verified by evaluating the remaining content by crystal violet.

Bacterial growth assays. *Microtiter plates:* Bacterial growth was evaluated by comparing OD_{600} values at the start and end of incubation in 96-well PVC microtiter plates. *Colony-forming units:* After incubation at 37 °C for 24 h, CFU/mL was calculated to determine the peptide solution's bactericidal effects. The untreated growth control was considered to be 100% planktonic cells. At least three technical and biological replicates were performed for all assays.

Quantitative reverse transcription PCR (qRT-PCR): After culturing for 24 h, RNAs were isolated from planktonic controls, biofilm controls, and from total cells exposed to 10 µM capsicumicine. An Invitrogen TRIzol Max bacterial RNA isolation kit and an Ambion TURBO DNase treatment were used as per manufacturer instructions. Total RNA concentrations and purities were assessed using a Biochrom SimpliNano spectrophotometer, and PCR reactions was done to ensure the complete absence of DNA. Each qRT-PCR reaction was then subjected to previously established quantities of cDNA (10 ng) and primers (0.2 µM). Reactional volumes were calculated using SYBR Select Master Mix (Applied Biosystems), as per the manufacturer's instructions. Primers (see Table S1) were designed using the Primer3 program, then produced by Eurofins Genomics. Applied Biosystems StepOnePlus equipment and software were used. Relative transcript levels were determined by the $2^{-\Delta\Delta ct}$ method (*60*).

Cytotoxicity assays: Cytotoxicity assays were performed on the ImPACcell robotic platform (BIOSIT, Université de Rennes 1). Multiparameter high-content screening (HCS) and high-content analysis (HCA) were done on 7 different mammalian lines: HuH7, CaCo-2, MDA, HCT116, PC3, NCI-H727, and MCF7. The number of normal cells is presented as residual cell percentage compared to the DMSO control average.

Microscopic analysis. *S. epidermidis* ATCC 35984 biofilm was cultured as described above.

Scanning electron microscopy (SEM): Sterile 10x4mm polystyrene coupons were inserted into bacterial cultures in the presence or absence of capsicumicine for 1, 4, and 24 h. The coupons were then washed with sterile 0.9% NaCl and fixed with 2.5% glutaraldehyde, 2% paraformaldehyde, and 0.1 M cacodylate buffer (pH 7.2). Afterwards, they were washed with 0.1 M cacodylate buffer and 0.2 M sucrose, then dehydrated with increasing concentrations of ethanol. A Leica EM CPD300 was used for critical point drying of the dehydrated samples. These were then sputtered with palladium in a Leica EM ACE200, and analyzed with a JEOL JSM-7100F microscope with EDS and EBSD at 10 kV.

Transmission electron microscopy (TEM): All well content was carefully detached at 1, 4, and 24 h, centrifuged at 10,000 g for 15 min at 4 °C, then washed with sterile 0.9% NaCl. Fixation was performed at 4 °C with sodium 0.1 M cacodylate, 2% paraformaldehyde, 2.5% glutaraldehyde, and 75 mM lysine. Samples were washed with 0.1 M sodium cacodylate and 0.2 M sucrose and contrasted with 1% osmium tetroxide and 1.5% potassium ferrocyanide. Dehydration was done with a gradual solution of ethanol and infiltration of increasing concentrations of LR White resin (Delta Microscopies, France). LR White resin inclusion and polymerization were then performed over 24 h at 60 °C in the absence of O_2. Thin 80 nm sections were collected onto carbon grids, and visualized at 200 kV with a FEI Tecnai Sphera microscope equipped with a Gatan 4k x 4k CCD UltraScan camera.

Confocal fluorescence microscopy (CFM): Capsicumicine-fluorescein isothiocyanate (capsicumicine-ITC, 10 µM) was used to detect the capsicumicine peptide, whose antibiofilm activity was previously verified. After incubation for 1, 4, or 24 h, the well contents were carefully detached, centrifuged at 11,000g for 2 min at 4 °C, then washed with sterile 0.9% NaCl. The suspension was visualized directly or after adding 0.1µg/µL concanavalin A conjugates (Alexa Fluor® 633, Invitrogen) or 2 mg/mL Calcofluor White dye (Fluorescent Brightener 28, Sigma-Aldrich). To find bacterial cells permeated by the peptide control, we used pseudonajide FITC-labelled antimicrobial peptide (Ref). Images were acquired via resonant scanner with a Leica SP8 DMI 6000 CS confocal microscope with hybrid detector, and ImageJ software was used for image analysis.

Real-time molecular self-assembly (RTMSA) assay: After checking the starting point (pH 7.2) of the assembly reaction for the synthetic staphylococcal matrix (*35*), we recorded the OD_{600} as a function of the time every 30 sec until 30 min. Molecular self-assembly reactions were calculated to a final volume of 4 mL, with 0.3% chitosan (medium molecular weight, 75-85% deacetylation), 0.15% bovine serum albumin, and 0.015% lambda DNA (all from Sigma) in TSB. The concentration (µM) of tested peptides was calculated for a final volume of 4 mL. Before getting the assembly reaction pH starting points, a calibration record was done using the same reactional tube containing all reagents (auto

zero). Acetic acid and NaOH were used to adjust pH, and the reaction temperature was about 30 °C.

As a negative control a similarly sized peptide was used, PA-1 (*61*).

NMR: NMR spectra were recorded in 3mm tubes on a Bruker Avance III 600MHz spectrometer

equipped with a TXI (1H,13C,15N) probe and a Z-gradient unit. Spectra were processed with Topspin

4.0.8 (Bruker Biospin) and CcpNmr Analysis (*62*). TOCSY and NOESY experiments were respectively

recorded at 10°C and pH5.0 with a 70ms and 300ms mixing time. Capsicumicine concentration in

water was set at 0.3mM and the pH adjusted either 5.0 or to 3.5 with few microliters of deuterated

HCl 0.1M and/or NaOH 0.1M solutions. Chitosan powder (purchased from Sigma) was dissolved in

pure water and the pH adjusted to either 5.0 or 3.5. Reference 1D proton NMR spectra were

recorded at 10°C and either at pH3.5 and 5.0. Chitosan was stepwise added from a concentrated

solution at pH 5.0 or 3.5 and spectra changes were monitored. We first titrated capsicumicine

solution with chitosan solution at pH 5.0. The concentration variation of chitosan was estimated

using the peaks intensities, resulting in 3.6 and 3.8 ppm (not shown) in the beginning to less than

0.5eq at the end of the titration. These experiments were repeated with a starting capsicumicine

solution at pH 3.5 and the gradually addition of gelled chitosan, at pH 3.5. The suddenly addition of

gelled chitosan pellets in the NMR solution, at pH3.5.

Circular Dichroism (CD): The peptide was dissolved in 18M water at a concentration of 50µM, at pH

5.0**.** The ultraviolet CD spectra were recorded at 5, 15 and 30°C, in a 0.1cm path length quartz cell on

a Jobin Yvon CD6 spectrometer equipped with a temperature controller unit, over a 180-260 nm

range, with a 2nm bandwidth, a step size of 1 nm and an integration time of 2s per point. The

samples were conserved at 5°C between each recording. Spectra were averaged over 5 records.

Water CD contributions were subtracted from CD spectra before processing. Spectra were processed

using Kaleidagraph (Synergy Software). Molar circular dichroism ( ) per residue and molar ellipticity

per residue (, MER) were computed from the difference of the delta absorbance recorded by the

spectrometer. Raw delta epsilon per residue spectra were analyzed using the CDSSTR program (*63*)

and different reference protein data sets provided by the DICHROWEB facility (*64*). MER data curves

were smoothed for presentation, using the interpolate and weighted data (5%) routines provide by

Kaleidagraph.

TANGO (*65*) and SALSA (*66*) predictions: Tango predictions were run at 298K. Predictions were

identical for pH set either to 5.0, 7.0 or 9.0. SALSA predictions were run with a window size dynamic

of 4-20 residues, a cutoff of 1.2 and a minimal hot spot length of 5.

CVC coatings: We adapted an approach to immobilize peptides based on poly(ethylene

glycol)diacrylate (PEGDA) hydrogel (*67*). Briefly, polyurethane (PU) tubes (Instech) of 2Fr or

25G equivalent were used as coating framework. Hydrogel base was prepared using

pentaerythritol tetrakis(3-mercaptopropionate) - PTMP (4.1 mmol), PEGDA (10 mmol), PEG-

600 (20mM), 2,2-Dimethoxy-2-phenylacetophenone - DMPA (0.1 wt%), THF soluble PU (10

wt%) and methanol (qs). Peptides were solubilized in DMSO and added to hydrogel base

under vortex agitation. PU tubes were first internally coated by suction using a needle-

syringe and then externally by immersion. The reaction and polymerization conditions such

as room temperature and oxygen tolerance were convenient. After polymerization,

successive washes with methanol under agitation were used to eliminate undesirable

monomers. All reagents were purchased from Sigma Aldrich.

CVC infection assay: This study was conducted by Voxcan s.a.r.l. The ethical agreement is part of the

project n° APAFiS# 10756-2017072522272676 v4, approved by Voxcan ethical committee (CEAA-129)

and the French authorities (ministry of national education, the higher education and research). This

study used SKH1 mice, females, immunocompetent and specific pathogen free (SPF) provided by

Charles River Laboratories. Animals were acclimated at least 2 days, housed collectively in disposable

standard cages in ventilated racks A3, at +21 ± 3°C, 30-70% of humidity, 12 hours of dark and light

cycles, with filtered water and autoclaved standard food provided *ad libitum*. Catheters were blind

implanted (n = 5 / group) in mice jugular vein following by an intravenous (IV) inoculation of *S.*

aureus Xen36, 5 days days later, which was in turn to colonize the device from the blood circulation.

The bacterial development was evaluated and compared between the different catheters by *in vivo*

bioluminescence imaging (IVIS Spectrum, acquisition and analysis with Living Image 4.5.5 version)

performed 2 and 4 days after mice infection. In addition, 7 days after inoculation or at the time of

mice euthanasia for ethical reason, catheters were harvested and bacterial load were evaluated by

CFU counting (SCAN500). All along the study, mice clinical state was evaluated using a scoring grid

and body weight measurements 3 times a week. At each timepoint, a bioluminescence acquisition

was also performed on the background (BKG) mouse to measure the flux level corresponding to the

auto-bioluminescence. Images were radiance-thresholded with respect to the background radiance

level and smoothed with a bidimensional Gaussian filter (3x3).

Acknowledgments:

General: We thank Daniel Thomas for his support, and Juliana Berland for insightful comments on

the manuscript. Thanks to the Microscopy Rennes Imaging Center (MRic) and CMEBA facilities.

Funding: This study was funded by the CAPES-COFECUB program, whose institutional partners are

the Brazilian Ministry of Education's CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível

Superior) agency and the French Ministère de l'Europe et des Affaires Étrangères (MEAE) and

Ministère de l'Enseignement supérieur, de la Recherche et de l'Innovation (MESRI). Support was also

received from the Transfer Acceleration Company SATT Ouest-Valorisation. **Author contributions:**

R.G.V.B, S.C.B.G, A.J.M, S.N-L. and R.G. designed research; R.G.V.B, S.C., R.S., S.N-L. and S.B.

performed research; R.G.V.B, A.R.Z., E.G., S.C.B.G, A.J.M, S.N-L., S.B and R.G. analyzed data; and

R.G.V.B, A.J.M, S.N-L. and R.G. wrote the paper. **Competing interests:** R.G.V.B, S.C.B.G, A.R.Z, A.J.M

and R.G. are co-authors of a patent register of capsicumicine. Application number WO 2020/169709

A1 at European Patent Office. Specific aspect of manuscript covered in patent application: its amino

acid sequence and bioactivity.

Data and materials availability: All data needed to evaluate the conclusions in the paper are present

in the paper and/or the Supplementary Materials. Additional data related to this paper may be

requested from the authors.

References:

1. WHO, W. H. Organization. (2015).
2. WHO, W. H. Organization. (2018).
3. A. Penesyan, M. Gillings, I. T. Paulsen, Antibiotic discovery: combatting bacterial resistance in
cells and in biofilm communities. *Molecules* **20**, 5286-5298 (2015).
4. H.-C. Flemming, J. Wingender, The biofilm matrix. *Nature Reviews Microbiology* **8**, 623-633
(2010).
5. A. Taglialegna *et al.*, Staphylococcal Bap Proteins Build Amyloid Scaffold Biofilm Matrices in
Response to Environmental Signals. *PLoS Pathog* **12**, e1005711 (2016).
6. N. Hoiby, T. Bjarnsholt, M. Givskov, S. Molin, O. Ciofu, Antibiotic resistance of bacterial
biofilms. *International Journal of Antimicrobial Agents* **35**, 322-332 (2010).
7. C. Beloin, S. Renard, J. M. Ghigo, D. Lebeaux, Novel approaches to combat bacterial biofilms.
Curr Opin Pharmacol **18C**, 61-68 (2014).
8. D. Davies, Understanding biofilm resistance to antibacterial agents. *Nat Rev Drug Discov* **2**,
114-122 (2003).
9. A. Brauner, O. Fridman, O. Gefen, N. Q. Balaban, Distinguishing between resistance,
tolerance and persistence to antibiotic treatment. *Nat Rev Microbiol* **14**, 320-330 (2016).
10. V. Defraine, M. Fauvart, J. Michiels, Fighting bacterial persistence: Current and emerging
anti-persister strategies and therapeutics. *Drug Resist Updat* **38**, 12-26 (2018).
11. M. Otto, Physical stress and bacterial colonization. *FEMS Microbiol Rev* **38**, 1250-1270 (2014).
12. I. Uçkay *et al.*, Meticillin resistance in orthopaedic coagulase-negative staphylococcal
infections. *J Hosp Infect* **79**, 248-253 (2011).
13. M. Otto, Staphylococcal biofilms. *Bacterial Biofilms* **322**, 207-228 (2008).
14. Y. Nishizaki *et al.*, Japanese features of native valve endocarditis caused by coagulase-
negative staphylococci: case reports and a literature review. *Intern Med* **52**, 567-572 (2013).
15. C. L. Abad, A. Haleem, Prosthetic Joint Infections: an Update. *Curr Infect Dis Rep* **20**, 15
(2018).
16. M. E. Rupp, Clinical characteristics of infections in humans due to Staphylococcus
epidermidis. *Methods Mol Biol* **1106**, 1-16 (2014).
17. CDCP, Center for Disease Control and Prevention. Vital signs: central line-associated blood
stream infections -United States, 2001, 2008, and 2009. *MMWR Morb Mortal Wkly Rep* **60**,
243-248 (2011).
18. J. Y. H. Lee *et al.*, Global spread of three multidrug-resistant lineages of Staphylococcus
epidermidis. *Nat Microbiol*, (2018).

19. J. M. Streit, R. N. Jones, H. S. Sader, T. R. Fritsche, Assessment of pathogen occurrences and
resistance profiles among infected patients in the intensive care unit: report from the
SENTRY Antimicrobial Surveillance Program (North America, 2001). *Int J Antimicrob Agents*
24, 111-118 (2004).
20. R. K. Flamm *et al.*, Linezolid Surveillance Results for the United States (LEADER Surveillance
Program 2014). *Antimicrob Agents Chemother* **60**, 2273-2280 (2016).
21. R. Hope *et al.*, Non-susceptibility trends among staphylococci from bacteraemias in the UK
and Ireland, 2001-06. *J Antimicrob Chemother* **62 Suppl 2**, ii65-74 (2008).
22. N. Strempel, J. Strehmel, J. Overhage, Potential Application of Antimicrobial Peptides in the
Treatment of Bacterial Biofilm Infections. *Current Pharmaceutical Design* **21**, 67-84 (2015).
23. C. de la Fuente-Nunez *et al.*, D-Enantiomeric Peptides that Eradicate Wild-Type and
Multidrug-Resistant Biofilms and Protect against Lethal Pseudomonas aeruginosa Infections
(vol 22, pg 196, 2015). *Chemistry & Biology* **22**, 1280-1282 (2015).
24. O. M. El-Halfawy *et al.*, Discovery of an antivirulence compound that reverses β-lactam
resistance in MRSA. *Nat Chem Biol* **16**, 143-149 (2020).
25. R. G. V. Borowski *et al.*, Red pepper peptide coatings control Staphylococcus epidermidis
adhesion and biofilm formation. *Int J Pharm* **574**, 118872 (2020).
26. The UniProt Consortium, UniProt: the universal protein knowledgebase. *Nucleic Acids Res* **45**,
D158-D169 (2017).
27. V. Lombard, H. Golaconda Ramulu, E. Drula, P. M. Coutinho, B. Henrissat, The carbohydrate-
active enzymes database (CAZy) in 2013. *Nucleic Acids Res* **42**, D490-495 (2014).
28. C. Feuillie *et al.*, Molecular interactions and inhibition of the staphylococcal biofilm-forming
protein SdrC. *Proc Natl Acad Sci U S A* **114**, 3738-3743 (2017).
29. S. Schmitt *et al.*, Analysis of modular bioengineered antimicrobial lanthipeptides at nanoliter
scale. *Nat Chem Biol* **15**, 437-443 (2019).
30. R. Roy, M. Tiwari, G. Donelli, V. Tiwari, Strategies for combating bacterial biofilms: A focus on
anti-biofilm agents and their mechanisms of action. *Virulence* **9**, 522-554 (2018).
31. S. Jabbouri, I. Sadovskaya, Characteristics of the biofilm matrix and its role as a possible
target for the detection and eradication of Staphylococcus epidermidis associated with
medical implant infections. *FEMS Immunol Med Microbiol* **59**, 280-291 (2010).
32. D. Fleming, K. P. Rumbaugh, Approaches to Dispersing Medical Biofilms. *Microorganisms* **5**,
(2017).
33. C. R. Armbruster, M. R. Parsek, New insight into the early stages of biofilm formation. *Proc
Natl Acad Sci U S A* **115**, 4317-4319 (2018).
34. M. Otto, Staphylococcal Infections: Mechanisms of Biofilm Maturation and Detachment as
Critical Determinants of Pathogenicity. *Annual Review of Medicine, Vol 64* **64**, 175-188
(2013).
35. E. J. Stewart, M. Ganesan, J. G. Younger, M. J. Solomon, Artificial biofilms establish the role of
matrix interactions in staphylococcal biofilm assembly and disassembly. *Sci Rep* **5**, 13081
(2015).
36. H. Hemmi, J. Ishibashi, T. Tomie, M. Yamakawa, Structural basis for new pattern of conserved
amino acid residues related to chitin-binding in the antifungal peptide from the coconut
rhinoceros beetle Oryctes rhinoceros. *J Biol Chem* **278**, 22820-22827 (2003).
37. T. Suetake *et al.*, Chitin-binding proteins in invertebrates and plants comprise a common
chitin-binding structural motif. *J Biol Chem* **275**, 17929-17932 (2000).
38. H. T. Wright, G. Sandrasegaram, C. S. Wright, Evolution of a family of N-acetylglucosamine
binding proteins containing the disulfide-rich domain of wheat germ agglutinin. *J Mol Evol*
33, 283-294 (1991).
39. J. Yin, S. Yang, K. Li, W. Guo, Y. Cao, Identification and molecular characterization of a chitin-
binding protein from the beet webworm, Loxostege sticticalis L. *Int J Mol Sci* **15**, 19147-
19161 (2014).

40. S. El-Gebali *et al.*, The Pfam protein families database in 2019. *Nucleic Acids Res* **47**, D427-
D432 (2019).
41. W. Suginta, P. Sirimontree, N. Sritho, T. Ohnuma, T. Fukamizo, The chitin-binding domain of a
GH-18 chitinase from Vibrio harveyi is crucial for chitin-chitinase interactions. *Int J Biol*
Macromol **93**, 1111-1117 (2016).
42. A. E. Paharik, A. R. Horswill, The Staphylococcal Biofilm: Adhesins, Regulation, and Host
Response. *Microbiol Spectr* **4**, (2016).
43. G. Batoni, G. Maisetta, S. Esin, Antimicrobial peptides and their interaction with biofilms of
medically relevant bacteria. *Biochimica Et Biophysica Acta-Biomembranes* **1858**, 1044-1060
(2016).
44. A. Sharma, P. Gupta, R. Kumar, A. Bhardwaj, dPABBs: A Novel in silico Approach for
Predicting and Designing Anti-biofilm Peptides. *Sci Rep* **6**, 21839 (2016).
45. S. F. Banani, H. O. Lee, A. A. Hyman, M. K. Rosen, Biomolecular condensates: organizers of
cellular biochemistry. *Nat Rev Mol Cell Biol* **18**, 285-298 (2017).
46. C. Rest, R. Kandanelli, G. Fernández, Strategies to create hierarchical self-assembled
structures via cooperative non-covalent interactions. *Chem Soc Rev* **44**, 2543-2572 (2015).
47. A. Kudlay, M. S. Cheung, D. Thirumalai, Influence of the shape of crowding particles on the
structural transitions in a polymer. *J Phys Chem B* **116**, 8513-8522 (2012).
48. C. Even *et al.*, Recent advances in studying single bacteria and biofilm mechanics. *Adv Colloid*
Interface Sci **247**, 573-588 (2017).
49. D. Marenduzzo, K. Finan, P. R. Cook, The depletion attraction: an underappreciated force
driving cellular organization. *J Cell Biol* **175**, 681-686 (2006).
50. H. X. Zhou, Effect of mixed macromolecular crowding agents on protein folding. *Proteins* **72**,
1109-1113 (2008).
51. A. Sakaguchi, K. Higashiguchi, K. Matsuda, Bundle formation of supramolecular fibers of
amphiphilic diarylethene by depletion force. *Chem Commun (Camb)* **54**, 4298-4301 (2018).
52. R. Fantoni, A. Santos, Depletion force in the infinite-dilution limit in a solvent of nonadditive
hard spheres. *J Chem Phys* **140**, 244513 (2014).
53. W. M. Aumiller, B. W. Davis, C. D. Keating, Phase separation as a possible means of nuclear
compartmentalization. *Int Rev Cell Mol Biol* **307**, 109-149 (2014).
54. S. Biswas, J. Kundu, S. K. Mukherjee, P. K. Chowdhury, Mixed Macromolecular Crowding: A
Protein and Solvent Perspective. *ACS Omega* **3**, 4316-4330 (2018).
55. J. M. Polson, D. R. Kerry, Segregation of polymers under cylindrical confinement: effects of
polymer topology and crowding. *Soft Matter* **14**, 6360-6373 (2018).
56. C. de la Fuente-Núñez, F. Reffuveille, L. Fernández, R. E. Hancock, Bacterial biofilm
development as a multicellular adaptation: antibiotic resistance and new therapeutic
strategies. *Curr Opin Microbiol* **16**, 580-589 (2013).
57. A. M. Krachler, K. Orth, Targeting the bacteria-host interface: strategies in anti-adhesion
therapy. *Virulence* **4**, 284-294 (2013).
58. L. Travier, O. Rendueles, L. Ferrieres, J.-M. Herry, J.-M. Ghigo, Escherichia coli Resistance to
Nonbiocidal Antibiofilm Polysaccharides Is Rare and Mediated by Multiple Mutations Leading
to Surface Physicochemical Modifications. *Antimicrobial Agents and Chemotherapy* **57**, 3960-
3968 (2013).
59. D. S. Trentin *et al.*, Natural Green Coating Inhibits Adhesion of Clinically Important Bacteria.
Scientific Reports **5**, (2015).
60. K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408 (2001).
61. P. Liu *et al.*, Targeting Inhibition of SmpB by Peptide Aptamer Attenuates the Virulence to
Protect Zebrafish against. *Front Microbiol* **8**, 1766 (2017).
62. S. P. Skinner *et al.*, CcpNmr AnalysisAssign: a flexible platform for integrated NMR analysis. *J*
Biomol NMR **66**, 111-124 (2016).

63.   N. Sreerama, R. W. Woody, Estimation of protein secondary structure from circular dichroism
699   spectra: comparison of CONTIN, SELCON, and CDSSTR methods with an expanded reference
700   set. *Anal Biochem* **287**, 252-260 (2000).
701   64.   L. Whitmore, B. A. Wallace, DICHROWEB, an online server for protein secondary structure
702   analyses from circular dichroism spectroscopic data. *Nucleic Acids Res* **32**, W668-673 (2004).
703   65.   A. M. Fernandez-Escamilla, F. Rousseau, J. Schymkowitz, L. Serrano, Prediction of sequence-
704   dependent and mutational effects on the aggregation of peptides and proteins. *Nat*
705   *Biotechnol* **22**, 1302-1306 (2004).
706   66.   S. Zibaee, O. S. Makin, M. Goedert, L. C. Serpell, A simple algorithm locates beta-strands in
707   the amyloid fibril core of alpha-synuclein, Abeta, and tau using the amino acid sequence
708   alone. *Protein Sci* **16**, 906-918 (2007).
709   67.   R. T. C. Cleophas *et al.*, Convenient Preparation of Bactericidal Hydrogels by Covalent
710   Attachment of Stabilized Antimicrobial Peptides Using Thiol–ene Click Chemistry. *ACS Macro*
711   *Letters* **3**, 477-480 (2014).
712   68.   L. A. Kelley, S. Mezulis, C. M. Yates, M. N. Wass, M. J. Sternberg, The Phyre2 web portal for
713   protein modeling, prediction and analysis. *Nat Protoc* **10**, 845-858 (2015).

715   **Supplementary Materials**

716   **Table S1.** Primers used in this study. These were previously designed using the Primer3 program and
717   standardized by our team.

Gene	Forward primer 5'-3'	Reverse primer 5'-3'	Amplicon
atlE	TACCAGGGTTTGCAGGATTC	GGCGCTAAATTCATTGGAAA	85 pb
aap	AGGCCGTACCAACAGTGAAT	ATGGGCAAACGTAGACAAGG	100 pb
agrC	TCATCAATATCGCATTCATCG	CCTAAACCGCGATTATCACC	136 pb
icaA	TTATCAATGCCGCAGTTGTC	CCGTTGGATATTGCCTCTGT	104 pb
leuA	GATGATCTCGGAATGGCAGT	TGAGGCATTTCCTGCTCTTT	108 pb
saeR	GCTAACACTGTCAATGTCCACA	AGGCCCCACACAGTTGTAAT	92 pb
saeS	GGCGTCAATTTGTTGTGCTA	AGGGCATAGGTATCGTTCCA	140 pb
sarA	TTTGCTTCTGTGATACGGTTGT	CGTAATGAACACGATGAAAGAACT	107 pb
gyrB	ATCAACATCGGCATCAGTCA	GCATTTGGTACGGGTATTGG	87 pb
rrsA	AAGCAACGCGAAGAACCTTA	ATGCACCACCTGTCACTCTG	95 pb

720 **Table S2.** Carbohydrate-binding domain proteins that are capsicumicine homologs. These proteins
721 are available in the UniProtKB database and were used for BLAST and amino acid alignments to
722 perform similarity analysis.

Entry	Name	Accession ID
ICAA_STAEQ	Intercellular adhesion protein icaA	Q5HKQ0.1
ICAD_STAEQ	Poly-beta-1,6-N-acetyl-D-glucos. synthase prot. icaD	Q5HKP9
A7Z8H9_BACVZ	Chitosanase	A7Z8H9
A0A0N0MLT5_9ACTN	Chitosanase	A0A0N0MLT5
CBP2_MOROL	Chitin-binding protein 2	C0HKC5
A0A1R0GTZ5_9FUNG	Chitin synthase 8	A0A1R0GTZ5
A0A194V113_9PEZI	Chitin biosynthesis protein CHS5	A0A194V113

723
724
725 **Table S3.** Results of the CD spectra deconvolution using CDSSTR algorithm.

Day	Temp (°C)	Helix (%)	Bêta (%)	Turn (%)	Unfolded (%)
0	5	42-46	29-34	5-8	16-21
1	5	42-45	30-32	6-8	20-21
5	5	38-42	24-31	8-12	17-23
5	15	22-26	36-45	13-14	20-26
5	30	16-24	33-40	18-20	20-25
7	30	6-17	38-41	10-23	31-35

729 **Fig. S1.** Biofilm eradication test. Shown are *Staphylococcus epidermidis* (ATCC 35984) biofilm
730 quantifications at OD_{570} for the bacterial biofilm without peptide exposure ("Growth Control"), after
731 exposure to the rifampicin antibiotic control, and after 24 h treatment with 100 μM capsicumicine.
732

Fig. S2. Confocal fluorescence microscopy (CFM) of *Staphylococcus epidermidis* (ATCC 35984).
Calcofluor was used to highlight matrix polysaccharides (blue), and FITC used for the peptides
(green). Visualization was done by fluorescence (left) and transmitted light (right) microscopy. (*A*)
Cultures exposed to capsicumicine and calcofluor. (*B*) Cultures exposed to capsicumicine-FITC. (*C*)
Peptide negative control cultures exposed to an antimicrobial peptide-FITC (Pseudonajide). The
matrix is highlighted (circles). Scale bars, 5 µm.

Fig. S3. Amino acid alignment of capsicumicine and its carbohydrate-binding domain protein
homologs. (*a*) The icaA (protein accession number Q5HKQ0.1) fragment from 1 to 60. (*b*)
Capsicumicine fragment from 1 to 21. (*c*) Chitosanase (A7Z8H9) fragment from 1 to 60 and from 61
to 120. (*d*) Capsicumicine fragment from 1 to 15 and from 16 to 22. (*e*) Chitin synthase 8
(A0A1R0GTZ5) fragment from 1801 to 1860 and from 1921 to 1967. (*f*) Capsicumicine fragment from
1 to 14 and from 16 to 22. Equal (* and grey highlighting), similar (.), and highly similar (:) amino
acids are indicated, as well as amino acid polar characteristics (purple). Support for this analysis is
available at UniProt.
750

Fig. S4. Superposition of the TOCSY (red), NOESY (green) and 1D (blue) spectra of capsicumicine recorded at pH 5.0 and 10°C. All amide NH resonances were assigned using nNH-(n-1)Hα NOESY cross-peaks, except for Q9 and Q11 residues due to poor resolution. For clarity, only NH-Hα and NH-Hβ labels were reported. The specta display 22 spin systems as expected for a single conformation. According to the spectral dispersion, capsicumicine is mainly folded.

Results of aggregation predictions

Position	Residue	TANGO	SALSA
1	R	0.0000	1.2020
2	S	0.0000	3.7070
3	C	9.0330	6.2770
4	Q	30.2200	6.2770
5	Q	85.3950	6.2770
6	Q	94.9340	6.2770
7	I	96.5820	6.2770
8	Q	99.2880	2.4720
9	Q	99.7080	0.0000
10	A	99.8980	0.0000
11	Q	99.7820	1.2010
12	Q	99.0890	4.9220
13	L	89.8320	10.0960
14	S	36.0550	13.9070
15	S	3.3470	16.5480
16	C	0.3860	20.5120
17	Q	0.0130	19.1140
18	Q	0.0060	16.6110
19	Y	0.0000	16.6110
20	L	0.0000	10.1030
21	K	0.0000	1.2080
22	Q	0.0000	0.0000

Fig. S5. (**A**) Predicted capsicumicine structure using the Phyre[2] prediction server. (**B**) Aggregation structure predictions using TANGO (*68*) and SALSA algorithms (*66*).