Article Type: Letter

Water-Body Type is More Important than Ecozone in Size-Based

Characterization of Freshwater Dissolved Organic Matter across Canada

Pieter J. K. Aukes[1,2]*, Sherry L. Schiff[1], Jason J. Venkiteswaran[2], Richard J. Elgood[1], John Spoelstra[1,3]

1) Department of Earth and Environmental Sciences, University of Waterloo, 200 University Avenue West,

Waterloo, Ontario, N2L 3G1, Canada

2) Department of Geography and Environmental Studies, Wilfrid Laurier University, 75 University Avenue West,

Waterloo, Ontario, N2L 3C5, Canada

3) Environment and Climate Change Canada, Canada Centre For Inland Waters, 867 Lakeshore Road, Burlington,

Ontario, L7S 1A1, Canada

*corresponding author: paukes@wlu.ca

Author Contribution Statement:

PJKA and SLS conceived and co-led the entire manuscript and contributed equally. PJKA and SLS

designed the research. Field work was conducted by PJKA, SLS, RJE, and JS. PJKA and SLS analyzed and

co-led the writing, as JJV, RJE, and JS contributed to writing of the manuscript. Statistical analyses was

conducted by PJKA and JJV.

SCIENTIFIC SIGNIFICANCE STATEMENT

Aquatic health and drinking water quality are affected by the differing mixtures of molecules that comprise dissolved organic matter (DOM), yet quantifying differences in DOM mixtures has been mostly focussed on light-absorbing techniques or confined to specific water-body types (i.e. only lakes). This makes it difficult to fully understand how DOM composition, and its reactivity, differs across the aquatic continuum, let alone across climatic gradients. This study uses molecular size-based groupings of DOM to quantify differences in DOM composition from both ground and surface waters across a range of Canadian ecozones, finding that DOM size distribution is quite similar within a water-body type regardless of surrounding vegetation and climate.

DATA AVAILABILITY STATEMENT

All data and metadata is available in the Wilfrid Laurier University Dataverse data repository and can be accessed at
https://dataverse.scholarsportal.info/dataset.xhtml?persistentId=doi:10.5683/SP2/ZRGXQ5

ABSTRACT

Dissolved Organic Matter (DOM) represents a mixture of organic molecules that vary due to different source materials and degree of processing. Characterizing how DOM composition evolves along the aquatic continuum can be difficult. Using a size-exclusion chromatography technique (LC-OCD), we assessed the variability in DOM composition from both surface and groundwaters across a number of Canadian ecozones (mean annual temperature spanning -10 to +6 C). A wide range in DOM concentration was found from 0.2 to 120 mg C/L. Proportions of different size-based groupings across ecozones were variable, yet similarities between specific water-body types, regardless of location, suggest commonality in the processes dictating the evolution of DOM composition. A principal-component analysis identified 70% of the variation in LC-OCD derived DOM compositions could be explained by the water-body type. We find that water-body type has a greater influence on DOM composition than differences in climate or surrounding vegetation.

Keywords:
Dissolved organic matter, size-exclusion chromatography, DOM composition, Canadian ecozones, water quality

Highlights
- Size-exclusion chromatography (using LC-OCD) is a fast and effective tool to quantify differences in DOM composition across different environments
- Proportions of biopolymers and low molecular weight fractions can distinguish different characteristics of surface and groundwater DOM
- Similar water-body types have comparable DOM size compositions across ecozones that have mean annual air temperatures ranging from -10 to $6\,^{\circ}C$

1 INTRODUCTION

Dissolved organic matter (DOM) is a ubiquitous component of terrestrial and aquatic

ecosystems. DOM influences light penetration within lakes (Schindler et al. 1996) and provides an

energy source for microbial metabolism (Biddanda and Cotner 2002). Comprised of thousands of

molecules with differing structures and properties, DOM concentration and composition can vary

greatly among environments due to different physical, chemical, and biological processes. Future

drinking water treatment options may be significantly impacted as climate change is predicted to alter

the quantity and quality of DOM in surface waters (Ritson et al. 2014). Harmful disinfection by-products

(DBP), formed by reactions between DOM and chlorine in drinking water supplies is more strongly

correlated with DOM composition than overall DOM concentration (Awad et al. 2016). Increased

terrestrial DOM contributions observed among northern surface waters are thought to result in the

'brownification' of these systems, affecting lake characteristics and food webs (Creed et al. 2018;

Wauthy et al. 2018). Hence, the evolution of DOM composition across the aquatic continuum has

important implications for downstream ecosystems, and must be monitored or observed to better

understand its impact on water treatment effectiveness and downstream ecosystems.

DOM concentration and composition can change along the aquatic continuum (i.e. transport

pathway from headwaters to the ocean) as it encounters different sources and degradation processes.

Watershed characteristics, such as type of terrestrial land cover or number of upstream lakes, can

dictate the concentration and composition of DOM that is transported through the aquatic continuum

(Mueller et al. 2012; Jaffé et al. 2012). The persistence of certain DOM components within freshwaters

has been linked to its initial composition (Kellerman et al. 2015). Determination of the variability in

DOM composition across the aquatic continuum can indicate potential avenues of future DOM change,

allowing us to better anticipate water treatment requirements and costs. However, due to the inherent

complexity and heterogeneity of DOM, a number of different techniques have been used to quantify

compositional differences.

Many of the techniques for analyzing DOM composition measure either bulk characteristics or

only a subset of all DOM molecules by capturing select components. Generally, the need for enhanced

information on DOM composition and molecular moieties results in increased cost and complexity to

implement (McCallister et al. 2018). Fortunately, comprehensive chemical characterization of DOM is

not always required to make useful predictions on how DOM will behave in the environment. For

instance, DOM composition has been assessed via molecular ratio assays (Hunt et al. 2000), light

absorption (Weishaar et al. 2003), fluorescence (Jaffé et al. 2008), resin fractionation (Kent et al. 2014),

and mass spectrometry (Kellerman et al. 2014; Hutchins et al. 2017). Bulk optical indices, although

common due to the relative ease of analysis, only respond to compounds that absorb or fluoresce in a

specific range of wavelengths and can include non-organic components in the matrix (Weishaar et al.

2003) or miss non-absorbing DOM components (Her et al. 2002). The ideal DOM characterization

method, or combination of methods, would provide sufficient detail on the key aspects of DOM

composition that control its fate and function, yet be analytically simple enough to be cost effective and

practical for environmental applications.

Size exclusion chromatography (SEC) has been used to characterize freshwater DOM in lakes

(Kent et al. 2014), rivers or streams (He et al. 2016), and groundwaters (Szabo and Tuhkanen 2010).

Liquid Chromatography – Organic Carbon Detection (LC-OCD) is a simple, fast (<120 minutes per

sample), and quantitative SEC method that separates DOM based upon hydrodynamic radii (Huber et al.

2011), characterizing both absorbing and non-absorbing DOM components. Studies using LC-OCD have

focussed on wastewater and water treatment applications (Ciputra et al. 2010), with some data from

agricultural and forested catchments in Germany (Heinz et al. 2015), and rivers in South Korea (He et al.

2016) and Spain (Catalán et al. 2017). LC-OCD-based DOM groupings provide a relatively easy measure

of DOM size-based composition across a large suite of environments.

Light absorbing and fluorescing components of DOM vary in the environment (Jaffé et al. 2008,

2012) but few studies examine how non-light absorbing DOM components differ across areas with

different climate and vegetation. Here we determine the effectiveness of using size-based groupings of

DOM to quantify differences in DOM composition and, to our knowledge, present one of few studies

that pairs DOM characterization of both groundwater and surface water across a gradient of climate

and vegetation regimes. The objectives of this study are: 1) to determine whether differences in DOM

composition, from various water-body types and ecozones (an ecosystem unit that groups similar areas

of biodiversity (Marshall et al. 1999)), can be identified using different LC-OCD fractions, and 2) to

quantify the degree of similarity in DOM composition among water-body types in different ecozones.

2 METHODS

2.1 Site Descriptions

Samples were collected in the summer months (June to August) across various Canadian

ecozones (Marshall et al. 1999) that span a mean annual temperature from -9.8 to 6.5°C, precipitation

from 313 to 940 mm, and overlie continuous to no permafrost (Figure 1; Suppl. Table 1). Surface water

samples were collected from the Boreal Shield (IISD Experimental Lakes Area (ELA)), Mixedwood

Plains (agriculturally-impacted Grand River (GR)), Taiga Shield (Yellowknife (YW) and Wekweètì

(WK)), and Southern Arctic ecozones (Daring Lake (DL); Suppl. Info Table 1). Groundwater, defined

here as water collected beneath the ground surface, was sampled from various depths from the Boreal

Shield (Turkey Lakes Watershed (TLW)) Mixedwood Plains (Nottawasaga River Watershed (NRW)),

Atlantic Maritime (Black Brook Watershed (BBW)), Taiga Shield (YK, WK), and Southern Arctic (DL).

Both NRW and BBW are areas of extensive agriculture. Shallow groundwater samples were collected

from the Northwest Territories (deepest extent of active-layer in July or August, ~0.1-0.5 m.b.s; meters

below surface) and TLW (0-7 m.b.s), while samples from NRW and BBW were collected at depths

ranging between 1 and 30 m.b.s.

2.2 DOM Characterization

All samples were filtered to 0.45µm (Whatman GD/X) into acid-washed and pre-rinsed glass

vials. Samples were immediately stored cool (<4°C) and dark until analyses (within two weeks). DOM

composition was quantitatively assessed using LC-OCD (DOC-Labor, Karlsruhe, Germany; Huber et al.

2011) at the University of Waterloo. A description of analytical equipment and DOM fraction

characterization has been described elsewhere (Huber et al. 2011) and a detailed explanation can be

found in the Supplementary Information. Briefly, the sample is injected through a size-exclusion

column (Toyopearl HW-50S, Tosoh Bioscience) that continuously elutes DOM based on hydrodynamic

radii into five hydrophilic fractions (from largest to smallest): biopolymers (BP; polysaccharides or

proteins), humic substances fraction (HSF; humic and fulvic acids), building blocks (BB; lower weight

humic substances), low molecular weight neutrals (LMWN; aldehydes, small organic materials), and

LMW-acids (LMWA; saturated mono-protic acids). Part of the sample bypasses the column for

measurement of total dissolved organic carbon concentration, herein referred to as DOM concentration

in mg C/L. As the hydrophilic component comprised 90±9% across all DOM samples, fraction

percentages were normalized to the sum of the eluted components (total hydrophilic fraction, herein

referred to as DOM_{hyphl}). The LC-OCD is both calibrated and run with potassium hydrogen phthalate,

IHSS-HA, and IHSS-FA standards to check DOM concentrations, minimize differences in machine drift,

and compare elution times over the period of analysis. Duplicates run at six concentrations yield a

precision of ±0.09 mg C/L or better. Samples also pass through a UV-Detector (Smartline UV Detector

200, Germany) for determination of the specific ultra-violet absorbance at 254 nm (SUVA), calculated by

normalizing the absorbance at this wavelength to the DOM concentration.

2.3 Statistical Analyses

Multiple sampling events from the same site have been averaged into one value per site (Suppl.

Fig S1; (Aukes et al. 2020)). Principal component analyses (PCA) was performed using *prcomp* in R (R

Core Team 2019). Differences in DOM composition across water-body types was assessed via a non-

parametric multivariate analysis of variance PERMANOVA using the *adonis2* function in the *vegan*

package (Oksanen et al. 2019). Further pairwise comparisons were calculated using a permutational

MANOVA via the *EcolUtils* package (Salazar 2020).

3 RESULTS

3.1 DOM Quantity

Highest DOM concentrations were found in groundwater samples in organic-rich peats from

Yellowknife (mean: 97 mg C/L, 1σ: ±40 mg C/L) and ELA (48±22 mg C /L; Figure 2a, Table 1). Lowest

concentrations were found in organic-poor groundwater sites (BBW: 0.6±0.4 mg C/L; NRW: 2.1±0.5 mg

C /L). The heavily agriculturally-impacted river (GR: 6.1±1.0 mg C /L) contained similar DOM

concentrations to pristine Canadian Shield subarctic rivers (YW: 6.6±3.1 mg C /L). Groundwater DOM

concentration exhibited a greater difference between ecozones than pond, lake, stream, or river DOM,

and were lower in southern areas with mineral strata compared to northern areas with organic-rich

deposits. In general, a larger range in DOM concentration was found across groundwater than other

water-body types.

3.2 DOM UV-Absorbance

The UV-absorbing capability of DOM can be compared across samples using SUVA values.

Lowest SUVA values were measured in agriculturally-impacted groundwater DOM (BBW, NRW) and

highest in northern groundwater (DL) and boreal stream DOM (ELA; Figure 2b; Table 1). Unlike

samples from ELA, northern ecozone groundwaters (DL, WK, and YW) had higher SUVA values than

other water-body types. Highest SUVA values were encountered in three Boreal Shield lakes at depth

(~5-10 m below surface) and may represent samples with iron interference (Weishaar et al. 2003) and

are therefore not included in subsequent discussion. Overall, differences in groundwater SUVA values

exist across ecozones but SUVA values are more similar among other water-body types.

3.3 LC-OCD Characterization

Different water-body types within an ecozone contained different proportions of BP.

Biopolymers ranged from 1-45% of DOM_{hyphl} but generally contributed less than 15% to DOM_{hyphl}

(Figure 3a; Table 1). Streams and lakes contained higher BP proportions than groundwaters.

Agricultural groundwater sites contained the lowest BP proportions (Figure 3a). High BP proportions

were found in Boreal and Arctic lakes, while moderate BP proportions were found in rivers and were

similar to other studies using LC-OCD (He et al. 2016; Catalán et al. 2017) (Figure 3a). Similar BP

fraction averages are found across a water-body type compared to different water-body types within an

ecozone, and high BP proportions are indicative of lake or stream DOM.

The HSF comprised the largest proportion, representing up to 85% of DOM_{hyphl} in some

environments (Figure 3b). Overall, the HSF proportion ranged from 15-85% with the highest proportions

found in Taiga Shield groundwater and Boreal streams (Figure 3). Taiga Shield and Southern Arctic lakes and ponds contained higher HSF proportions than Boreal lakes (Table 1). Taiga Shield rivers had slightly lower proportions than the Mixedwood Plains agriculturally-impacted river. Boreal and agriculturally-impacted (NRW and BBW) groundwaters had the lowest HSF proportions, as well as Boreal wetland groundwater (Table 1; Figure 3).

Smaller molecular weight humics, defined as building blocks (BB), ranged from 5-37% of DOM_{hyphl}. High BB proportions were found from southern agricultural groundwater samples, while lowest proportions were observed among a Boreal shield stream and groundwater samples in organic rich peats in YW and ELA (<10%; Figure 3c). Across ecozones, ponds, lakes, streams, and rivers had comparable proportions of BB between 10-30% of DOM_{hyphl}.

Proportions of LMWN ranged from 3-76% and were much lower than HSF except among groundwater samples (Figure 3d). DOM from Boreal wetland groundwater sites contained higher LMWN proportions than any other environment (Table 1). Groundwater environments with the lowest HSF proportion had higher LMWN proportions, specifically the agriculturally-impacted sites. Lowest proportions were found from Boreal streams and Arctic groundwater and pond samples. LMWA generally comprised a minor component of total DOM (Figure 3e).

The variability across different sites in LC-OCD defined DOM composition was assessed using PCA. Initially, two principal axes accounted for 60% of the variability within the dataset (Suppl. Fig S2). However, high LMWN proportions from Boreal wetland groundwaters did not allow for a good resolution of other components. For this reason, only these wetland groundwater samples were omitted (n=28) and the PCA was recalculated based on the remaining samples (n=122; Figure 4; Suppl. Table 2). Two principal axes now account for 71% of the variability and illustrates a similarity in DOM composition when comparing water-body types across ecozones. The first principal component (PC) axis represents degraded humics or smaller DOM components, whereas the second PC represents differences in the BP fraction (Figure 4). Distinct compositions are observed comparing lake and groundwater DOM, while rivers and streams contain intermediary compositions between these. Broad

variation in groundwater DOM composition resulted from HSF and LMWN or BB proportions, while

higher BP proportions were common for ponds, lakes, streams and rivers. Lake DOM composition was

also highly variable and mainly characterized by LMWN, BB, and BP. Groundwater DOM from Arctic

ecozones (organic soils) grouped separately from other groundwater samples (mineral substrate), but

were similar in composition to Boreal streams. Lake DOM can be identified by higher BB and LMWN,

lower HSF, and occasionally high BP.

4 DISCUSSION

4.1 Can LC-OCD Analyses Differentiate DOM Composition?

This study provided a comprehensive dataset to quantify DOM heterogeneity across a range of

environmental conditions that spans the Southern Arctic, with long-cold winters and short-cool

summers, to the Mixedwood Plains with warm-wet climates and productive soils. Differences in both

SUVA and size-based properties of DOM fractions are apparent (Figure 2, 3) and result from differences

in organic matter sources, catchment characteristics, residence times, and processing history (Curtis

and Schindler 1997; Jaffé et al. 2008; Mueller et al. 2012). We find that size-based groupings can identify

differences in DOM composition not observed measuring SUVA alone. For instance, ELA and TLW

wetland groundwater samples contained similar SUVA values (Figure 2b) but different LC-OCD

compositions: a greater proportion of degraded humics at TLW than ELA (Figure 3c; Table 1**)**. The use

of various characterization techniques would provide different information on DOM and may be the

best way to holistically quantify DOM. Differences in size-based groupings of DOM provide additional

information suited to identifying differences in DOM composition that is not provided by UV-based

techniques alone.

Although our study design limits the ability to conclusively determine whether DOM

composition differed significantly with ecozone, PCA and PERMANOVA analyses suggests a similarity

in DOM composition within a water-body type (Table 2). Thus, although climate and vegetation differ,

DOM composition within a water-body type is comparable across ecozones to be significantly different

when compared to other water-body types.

4.2 Similarity in Groundwater DOM

Groupings of similar water-body types in the PCA indicates a commonality of DOM

composition across spatial scales. The general conception, especially in groundwater environments, is

that processing of DOM results in a loss of heterogeneity. For instance, physiochemical and biological

processes were observed to conform South Carolina soil DOM composition (Shen et al. 2015). The

grouping of groundwater DOM in the PCA indicates various ecozones contain similar mixtures of LC-

OCD defined components. Groundwater DOM is differentiated from DOM in other water-body types

by the proportion of BP, but can be even further separated by HSF and LMWN to compare organic-rich

contributions (ELA and TLW) to agriculturally-impacted sites within mineral strata (BBW and NRW;

Figure 3a; Figure 4). Biodegradation and accumulation of DOM in porewaters (Chin et al. 1998) may be

responsible for wetland groundwater DOM from Boreal shield sites being easily identified by high

proportions of LMWN (Suppl Fig 2). Hence, size-based groupings of groundwater DOM indicates that,

while composition can differ among groundwater environments, groundwater contains much different

DOM than other water-body types.

4.3 Trends in DOM Composition across Water-Body Types

Photolysis and *in-situ* production are two processes that occur among surface waters that alter

the composition of DOM. Exposure of DOM to sunlight and photochemical transformations of humics

results in smaller molecules and increased proportions of BB (Tercero Espinoza et al. 2009; He et al.

2016). This is observed in the dataset by a shift from PCA groupings with high BP (indicative of lakes,

ponds, and streams) towards higher BB and LMWN suggestive of exposure to sunlight (Figure 4). High

BP in lakes, rivers, and most streams indicate high molecular weight polysaccharides or proteins that

form during *in-situ* production or microbial processing of DOM (Ciputra et al. 2010; Huber et al. 2011;

Catalán et al. 2017).

272   Rivers and streams can act as conduits of DOM transport and internal producers and processors

273 of DOM (Creed et al. 2015), forming a DOM composition intermediary between lake and groundwater

274 end-members. Further, stream DOM composition can identify DOM source as some samples plot closely

275 with either groundwater or lake DOM (Figure 4). Thus, size-based DOM groupings can be used to

276 evaluate the importance of autochthonous versus groundwater sources of DOM in rivers and streams.

277   Along a hypothetical flow path, terrestrial groundwater DOM is mobilized into surrounding

278 ponds and lakes, and eventually exported from the watershed via rivers or streams. Changes to LC-

279 OCD components, concurrent with decreases to overall DOM concentration, are found along this

280 hypothetical flow with the loss of high-HSF in groundwaters as LMW and BP proportions increase

281 (Figure 3, 4). These changes are in agreement with watershed-scale observations of DOM composition

282 change along an aquatic continuum, shifting from relatively high-molecular weight groundwater

283 components to smaller, more degraded, and more aliphatic components (Kellerman et al. 2014; Hutchins

284 et al. 2017). The similarity in DOM composition within water-body types across different ecoregions

285 (Figure 4) indicates the processes responsible for altering DOM composition may be similar regardless

286 of surrounding vegetation or climate.

287 *4.4 Implications of Different Size-Based Groupings of DOM*

288   Differences in DOM composition are associated with water-body type, hence shifts or changes

289 to the hydrological regime, such as recent increases to terrestrial-derived DOM, can alter DOM

290 composition and its effect upon the surrounding environment (Creed et al. 2018). Size-exclusion DOM

291 analysis provides a quantitative measure of DOM composition variability across aquatic environments,

292 and may help identify lakes experiencing stronger terrestrial influences, such as higher HSF and lower

293 BP. In particular, a shift towards heavier fall rains in the Northwest Territories, Canada (Spence et al.

294 2011), can enhance terrestrial DOM contributions to surrounding surface waters, which may result in

295 increased pre-treatment costs (Ritson et al. 2014) and higher DBP concentrations (Awad et al. 2016).

296 Implementation of this size-based grouping scheme provides a quantitative tool to measure DOM

297 composition while acknowledging the inherent heterogeneity when attempting to characterize the

complex mixtures characteristic of naturally occurring DOM. The ability to characterize and compare

specific DOM compositions is important to better predict, plan, and adapt water treatment methods for

these climate-induced changes.

ACKNOWLEDGEMENTS

Thanks to the Environmental Geochemistry Laboratory at the University of Waterloo for assistance

with sample collection and processing. Staff of the Fredericton Research and Development Centre,

Agriculture and Agri-Food Canada, collected samples from the Black Brook Watershed, New

Brunswick. This work benefited from the assistance of Monica Tudorancea with operation of the LC-

OCD instrument. We would also like to thank two anonymous reviewers for their suggestions and

comments. Funding was provided by a Natural Science and Engineering Research Council (NSERC) of

Canada grant to S. L. Schiff, and an Ontario Graduate Scholarship, University of Waterloo Scholarship,

and International Association of GeoChemistry Research Award (IAGC) to P. J. K. Aukes. Additional

financial support was provided to J. Spoelstra by Environment and Climate Change Canada and the

Ontario Ministry of Agriculture, Food, and Rural Affairs (OMAFRA). The authors declare no conflicts of

interest related to the study. All data can be found in the Supplementary Information, while code used

to generate figure and statistical analysis used in this manuscript is available at

github.com/paukes/canada-ecozone-DOM.

REFERENCES

Aukes, P. J. K., S. L. Schiff, J. J. Venkiteswaran, R. J. Elgood, and J. Spoelstra. 2020. Data used in Water

Bodies are More Important than Ecozone in Size-Based Characterization of Freshwater Dissolved

Organic Matter across Canada. Wilfrid Laurier Univ. Dataverse. doi:10.5683/SP2/ZRGXQ5

Awad, J., J. van Leeuwen, C. W. K. Chow, M. Drikas, R. J. Smernik, D. J. Chittleborough, and E. Bestland.

2016. Characterization of dissolved organic matter for prediction of trihalomethane formation

potential in surface and sub-surface waters. J. Hazard. Mater. **308**: 430–439.

doi:10.1016/j.jhazmat.2016.01.030

Biddanda, B. A., and J. B. Cotner. 2002. Love handles in aquatic ecosystems: The role of dissolved

organic carbon drawdown, resuspended sediments, and terrigenous inputs in the carbon balance

of Lake Michigan. Ecosystems **5**: 431–445. doi:10.1007/s10021-002-0163-z

Catalán, N., J. P. Casas-Ruiz, D. von Schiller, L. Proia, B. Obrador, E. Zwirnmann, and R. Marcé. 2017.

Biodegradation kinetics of dissolved organic matter chromatographic fractions in an intermittent

river. J. Geophys. Res. Biogeosciences **122**: 131–144. doi:10.1002/2016JG003512

Chin, Y., S. J. Traina, C. R. Swank, and D. Backhus. 1998. Abundance and properties of dissolved organic

matter in pore waters of a freshwater wetland. Limnol. Oceanogr. **43**: 1287–1296.

Ciputra, S., A. Antony, R. Phillips, D. Richardson, and G. Leslie. 2010. Comparison of treatment options

for removal of recalcitrant dissolved organic matter from paper mill effluent. Chemosphere **81**:

86–91. doi:10.1016/j.chemosphere.2010.06.060

Creed, I. F., A. K. Bergström, C. G. Trick, and others. 2018. Global change-driven effects on dissolved

organic matter composition: Implications for food webs of northern lakes. Glob. Chang. Biol. **24**:

3692–3714. doi:10.1111/gcb.14129

Creed, I. F., D. M. Mcknight, B. A. Pellerin, and others. 2015. The river as a chemostat : fresh

perspectives on dissolved organic matter flowing down the river continuum. Can. J. Fish. Aquat.

Sci. **14**: 1–14.

Curtis, P. J., and D. W. Schindler. 1997. Hydrologic control of dissolved organic matter in low-order

Precambrian Shield lakes. Biogeochemistry **36**: 125–138. doi:10.1023/A:1005787913638

He, W., I. Choi, J.-J. Lee, and J. Hur. 2016. Coupling effects of abiotic and biotic factors on molecular

composition of dissolved organic matter in a freshwater wetland. Sci. Total Environ. **544**: 525–534.

doi:10.1016/j.scitotenv.2015.12.008

Heinz, M., D. Graeber, D. Zak, E. Zwirnmann, J. Gelbrecht, and M. T. Pusch. 2015. Comparison of

Organic Matter Composition in Agricultural versus Forest Affected Headwaters with Special

Emphasis on Organic Nitrogen. Environ. Sci. Technol. **49**: 2081–2090. doi:10.1021/es505146h

Her, N., G. L. Amy, and D. Foss. 2002. Variations of molecular weight estimation by HP-size exclusion

chromatography with UVA versus online DOC detection. Environ. Sci. Technol. **36**: 3393–3399.

Huber, S. A., A. Balz, M. Abert, and W. Pronk. 2011. Characterisation of aquatic humic and non-humic

matter with size-exclusion chromatography – organic carbon detection – organic nitrogen

detection (LC-OCD-OND). Water Res. **45**: 879–885. doi:10.1016/j.watres.2010.09.023

Hunt, A. P., J. D. Parry, and J. Hamilton-Taylor. 2000. Further evidence of elemental composition as an

indicator of the bioavailability of humic substances to bacteria. Limnol. Oceanogr. **45**: 237–241.

Hutchins, R. H. S., P. J. K. Aukes, S. L. Schiff, T. Dittmar, Y. T. Prairie, and P. A. del Giorgio. 2017. The

Optical, Chemical, and Molecular Dissolved Organic Matter Succession Along a Boreal Soil-

Stream-River Continuum. J. Geophys. Res. Biogeosciences **122**: 2892–2908.

doi:10.1002/2017JG004094

Jaffé, R., D. McKnight, N. Maie, R. M. Cory, W. H. McDowell, and J. L. Campbell. 2008. Spatial and

temporal variations in DOM composition in ecosystems: The importance of long-term monitoring

of optical properties. J. Geophys. Res. **113**: 1–15. doi:10.1029/2008JG000683

Jaffé, R., Y. Yamashita, N. Maie, and others. 2012. Dissolved Organic Matter in Headwater Streams:

Compositional Variability across Climatic Regions of North America. Geochim. Cosmochim. Acta

94: 95–108. doi:10.1016/j.gca.2012.06.031

Kellerman, A. M., T. Dittmar, D. N. Kothawala, and L. J. Tranvik. 2014. Chemodiversity of dissolved

organic matter in lakes driven by climate and hydrology. Nat. Commun. **5**: 1–8.

doi:10.1038/ncomms4804

Kellerman, A. M., D. N. Kothawala, T. Dittmar, and L. J. Tranvik. 2015. Persistence of dissolved organic

matter in lakes related to its molecular characteristics. Nat. Geosci. **8**. doi:10.1038/ngeo2440

Kent, F. C., K. R. Montreuil, A. K. Stoddart, V. A. Reed, and G. A. Gagnon. 2014. Combined use of resin

fractionation and high performance size exclusion chromatography for characterization of natural

organic matter. J. Environ. Sci. Heal. **4529**: 1615–1622. doi:10.1080/10934529.2014.950926

Marshall, I. B., P. H. Schut, and M. Ballard. 1999. A National Ecological Framework for Canada:

Attribute Data. Ecosystem Stratification Working Group, Agriculture and Agri-Food Canada &

Environment Canada.

McCallister, S. L., N. F. Ishikawa, and D. N. Kothawala. 2018. Biogeochemical tools for characterizing

organic carbon in inland aquatic ecosystems. Limnol. Oceanogr. Lett. **3**: 444–457.

doi:10.1002/lol2.10097

Mueller, K. K., C. Fortin, and P. G. C. Campbell. 2012. Spatial Variation in the Optical Properties of

Dissolved Organic Matter (DOM) in Lakes on the Canadian Precambrian Shield and Links to

Watershed Characteristics. Aquat. Geochemistry **18**: 21–44. doi:10.1007/s10498-011-9147-y

Oksanen, J., F. Guillaume Blanchet, M. Friendly, and others. 2019. vegan: Community Ecology Package.

R Core Team. 2019. R: A language and environment for statistical computing.

Ritson, J. P., N. J. D. Graham, M. R. Templeton, J. M. Clark, R. Gough, and C. Freeman. 2014. The impact

of climate change on the treatability of dissolved organic matter (DOM) in upland water supplies:

A UK perspective. Sci. Total Environ. **473–474**: 714–730. doi:10.1016/j.scitotenv.2013.12.095

Salazar, G. 2020. EcolUtils: Utilities for commnity ecology analysis.

Schindler, D. W., P. J. Curtis, B. R. Parker, and M. P. Stainton. 1996. Consequences of climate warming

and lake acidification for UV-B penetration in North American boreal lakes. Nature **379**: 705–708.

Shen, Y., F. H. Chapelle, E. W. Strom, and R. Benner. 2015. Origins and bioavailability of dissolved

organic matter in groundwater. Biogeochemistry **122**: 61–78. doi:10.1007/s10533-014-0029-4

Spence, C., S. V. Kokelj, and E. Ehsanzadeh. 2011. Precipitation trends contribute to streamflow regime

shifts in northern Canada. *Cold Region Hydrology in a Changing Climate*. 3–8.

Szabo, H. M., and T. Tuhkanen. 2010. The application of HPLC-SEC for the simultaneous

characterization of NOM and nitrate in well waters. Chemosphere **80**: 779–86.

doi:10.1016/j.chemosphere.2010.05.007

Tercero Espinoza, L. A., E. ter Haseborg, M. Weber, and F. H. Frimmel. 2009. Investigation of the

photocatalytic degradation of brown water natural organic matter by size exclusion

chromatography. Appl. Catal. B Environ. **87**: 56–62. doi:10.1016/j.apcatb.2008.08.013

Wauthy, M., M. Rautio, K. S. Christoffersen, L. Forsström, I. Laurion, H. L. Mariash, S. Peura, and W. F.

Vincent. 2018. Increasing dominance of terrigenous organic matter in circumpolar freshwaters due

to permafrost thaw. Limnol. Oceanogr. Lett. doi:10.1002/lol2.10063

Weishaar, J. L., G. R. Aiken, B. A. Bergamaschi, M. S. Fram, R. Fujii, and K. Mopper. 2003. Evaluation of

specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of

dissolved organic carbon. Environ. Sci. Technol. **37**: 4702–4708. doi:10.1021/es030360x

TABLES

Table 1: Dissolved organic matter concentration, specific UV-absorbance at 254 nm (SUVA), and LC-OCD characterization for different sites and water-body types across a range of Canadian ecozones. The LC-OCD percentages are relative to the total hydrophilic fraction of the samples and include biopolymers (BP), humic substances fraction (HSF), building blocks (BB), low molecular weight neutrals (LMWN), and LMWN acids (LMWA).

Ecozone	Location	Water body			DOM (mg C/L)		SUVA (L / (mg m))		BP (%)		HSF (%)		BB (%)		LMWN (%)		LMWA (%)	
		Type	n		Mean	σ (±)	Mean	σ (±)	Mean	σ (±)	Mean	σ (±)	Mean	σ (±)	Mean	σ (±)	Mean	σ (±)
Southern Arctic	Daring Lake	Groundwater	4		21.0	9.35	5.3	0.5	5.39	3.95	73.1	4.52	14.8	2.75	6.39	2.37	0.30	0.35
		Pond	1		4.00		2.2		33.5		39.8		15.3		11.3		0.00	
		Lake	8		5.64	2.95	3.4	1.2	14.6	14.4	58.5	15.8	15.6	3.41	10.8	1.93	0.54	0.42
		Stream	10		7.96	3.44	3.8	0.8	13.3	9.50	57.4	10.4	17.4	3.30	11.0	1.15	0.95	0.48
Taiga Shield	Wekweètì	Groundwater	1		10.2		4.7		2.09		70.1		14.8		6.91		6.18	
		Lake	6		11.0	5.05	3.4	1.1	7.24	4.07	59.9	10.3	20.7	4.75	10.3	2.90	1.91	2.36
		Stream	1		6.24		2.6		12.3		57.2		19.6		9.94		0.91	
Taiga Shield	Yellowknife	Groundwater	17		94.1	41.4	4.4	0.8	5.79	3.90	75.8	5.93	10.3	2.91	7.15	1.61	1.02	1.75
		Pond	2		31.9	2.97	3.6	0.4	5.08	1.89	70.5	1.70	14.8	0.53	8.56	0.65	1.04	0.31
		Lake	1		27.8		1.5		21.9		52.0		15.8		9.84		0.60	
		Stream	1		16.0		2.0		19.5		55.4		15.8		8.67		0.64	
		River	3		6.49	1.25	2.0	0.2	10.7	3.10	59.8	2.36	17.0	1.27	10.7	0.56	1.76	2.08
Boreal Shield	IISD-Experimental Lakes Area	Wetland Groundwater	17		47.8	22.3	2.9	0.9	7.21	2.76	37.2	13.6	7.82	2.65	46.9	15.0	0.81	1.60
		Lake	21		7.53	2.80	3.2	1.7	14.8	6.21	49.2	6.95	23.0	3.74	12.2	1.82	0.79	0.57
		Wetland Complex	2		12.2	3.00	4.3	0.2	11.8	0.71	58.7	4.41	18.1	2.85	10.7	0.54	0.83	0.30
		Stream	11		20.4	6.77	5.3	0.6	2.50	2.66	80.7	2.85	9.40	3.60	7.27	1.62	0.17	0.27
Boreal Shield	Turkey Lakes Watershed	Wetland Groundwater	9		1.83	0.69	3.2	1.4	3.20	4.22	43.5	17.7	21.5	5.05	28.0	18.8	6.49	15.8
		Groundwater	7		4.20	5.56	3.4	1.0	0.51	0.40	58.3	12.2	23.5	7.20	17.3	11.5	0.68	1.13
Mixedwood Plains	Nottawasaga River Watershed	Groundwater	6		2.07	0.53	3.6	0.1	0.42	0.26	54.9	8.02	28.8	5.30	15.9	3.24	0.03	0.08
	Grand River	River	7		6.07	0.73	2.6	0.6	9.54	0.77	65.9	1.78	14.6	0.93	9.60	0.79	0.28	0.45
Atlantic Maritime	Black Brook	Groundwater	15		0.67	0.47	2.3	0.9	1.32	1.39	53.6	10.2	27.3	7.87	17.7	6.66	0.12	0.47

413   Table 2: Non-parametric PERMANOVA on Euclidean distances for dissolved organic matter (DOM) compositions collected across various

414   water-body types, with ecozone nested within the function. Included are the degrees of freedom (d.f), sum of squares (SS), R-squared value (R^2),

415   F-statistic (F), and p-value (P). Further post hoc comparisons were conducted using Dunn's Test. All code used to generate these results can be

416   found on github.com/paukes/canada-ecozone-DOM.

Parameter	d.f.	SS	R^2	F	P
Water-Body Type	5	0.7859	0.19644	5.7695	0.001
Residual	118	3.2146	0.80356		
Total	123	4.0005	1		

Comparison*	Corrected p-value
Lake vs River	0.05
Lake vs Stream	<0.01
Lake vs Groundwater	<0.01

Pair-wise a posteriori tests (only significant pairs included)

418 **FIGURES**

Figure 1: Ecozones (Marshall et al. 1999) and DOM collection sites: Daring Lake (DL; Southern Arctic ecozone),

Wekweeti (WK; Taiga Shield ecozone), Yellowknife (YW; Taiga Shield ecozone), International Institute for

Sustainable Development - Experimental Lakes Area (IISD-ELA; Boreal Shield ecozone), Turkey Lakes Watershed

(TLW; Boreal Shield ecozone), Nottawasaga River Watershed (NRW; Mixedwood Plains ecozone), Grand River

Watershed (GR; Mixedwood Plains ecozone), and Black Brook Watershed (BBW; Atlantic Maritime ecozone).

Figure 2: Comparison of dissolved organic matter concentration (DOM; logarithmic y-scale, a) and specific ultra-violet absorbance at 254 nm (SUVA; b) across different water-body types. For each water-body type, different ecozones are grouped together and denoted by different shapes, while separate locations within each ecozone are differentiated by colour. Boxplots for each ecozone in a water-body type indicate the mean value and extend to the first and third quartile, while whiskers represent 1.5x inter-quartile range. Included are literature values that use LC-OCD in South Korea (He et al. 2016) and Spain (Catalán et al. 2017), and are denoted by symbols in grey. Literature values from South Korea include a maximum and minimum value found within the study (open circle). Random scatter is introduced in the x-axis for ease of seeing all data points.

Figure 3: Differences in the composition of DOM based on proportions of LC-OCD groups across water-body types:

biopolymers (BP; a), humic substances fraction (HSF; b), building blocks (BB; c), low molecular weight neutrals (LMWN; d), and

low molecular weight acids (LMWA; e). For each water-body type, different ecozones are grouped together and denoted by

different shapes, while separate locations within each ecozone are differentiated by colour. Boxplots for each ecozone in a

water-body type represent the mean value and extend to the first and third quartile, while whiskers represent 1.5x inter-

 quartile range. Included are literature values that use LC-OCD in South Korea (He et al. 2016) and Spain (Catalán et al. 2017),

441 and are denoted by symbols in grey. South Korea include a maximum and minimum value found within the study (open circle).

442 Random scatter is introduced in the x-axis for ease of seeing all data points.

444
445 *Figure 4*: Principal component analyses (PCA) of LC-OCD data (minus wetland groundwater samples) with first and second

446 principal component axes plotted. PCA vectors are included for the proportion of DOM for each LC-OCD group: biopolymers

447 (BP), humic substances fraction (HSF), building blocks (BB), low molecular weight neutrals (LMWN), and low molecular weight

448 acids (LMWA).

SUPPLEMENTARY INFORMATION

LC-OCD Methodology

Dissolved organic matter characterization using liquid chromatography – organic carbon detection (LC-OCD) was performed in the Department of Civil and Environmental Engineering, University of Waterloo. A detailed description of the instrument, setup, and analysis can be found in Huber et al. (2011). First, samples are diluted to a DOM concentration between 1 – 5 mg C/L to obtain an optimal chromatogram to analyse. Dilution factors were used to correct concentrations for comparison. The sample is added to a mobile phase (phosphate buffer exposed to UV-irradiation) and passed through an in-line 0.45μm filter and into either a by-pass or the size-exclusion column (Toyopearl HW-50S (Tosoh, Japan)). The by-pass allows for overall determination of the dissolved organic carbon concentration, as well as the overall UV-absorbance at 254 nm. The novel organic carbon detector (OCD) is comprised of a thin film UV-reactor (Gräntzel thin-film reactor, DOC-Labor, Karlsruhe, Germany). As the acidified sample enters the OCD , it is thinly spread over a UV-lamp, allowing irradiation to oxidize organic carbon into CO_2. The CO_2 is then measured with a highly-sensitive infrared detector, where the integration of the peak area provides the total DOC concentration. Sample that is not passed through the by-pass enters the size-exclusion column (SEC) where it is divided into different fractions based on nominal molecular weight and is continuously analyzed by the UV detector and OCD, producing OCD and UV chromatograms.

LC-OCD Molecular Groupings

Chromatograms are analyzed using customized software (ChromCALC, DOC-LABOR, Karlsruhe, Germany) which calculates the concentration of each 'size grouping' by integrating areas based on specific elution times. First, the overall DOC concentration is determined from the bypass peak. Second, the hydrophilic portion (sample that elutes from the SEC) is subdivided into five categories based upon elution time: biopolymers (BP), humic substances fraction (HSF; which includes both humic and fulvic acids), building blocks (BB), and low-molecular-weight neutrals (LMW-N) and acids (LMW-A).

Grouping names and boundaries of each fraction are arbitrarily defined based on elution times and

characteristics as determined by Huber et al. (2011). Briefly, the groupings are (from largest to smallest):

• Biopolymers: >10 kDa; polysaccharides, proteins, or amino sugars

• Humic substances fraction: 100 – 1200 Da; a heterogeneous mixture of large, complex molecules

that elute at similar times as the IHSS humic standards

• Building blocks: degraded humic substances with humic-like characteristics, but of lower

molecular weight

• LMW acid and LMW neutrals: contain monoprotic acids, amino sugars, ketones, and aldehydes

Finally, the software determines a 'hydrophobic' parameter quantified as the residual between the sum

of all measured fractions and the overall concentration of DOC (determined from the by-pass). As the

difference between the overall DOM concentration and the sum of the five eluted fractions. As the

hydrophilic component comprised 90±9% across all DOM samples, group percentages were normalized

to the sum of the eluted components.

SUPPLEMENTARY TABLES

Table S1: Description of sampling sites. The number of samples represents the number of distinct sites sampled in that water-body

for that ecozone. Mean annual air temperatures and total annual precipitation taken from Marshall et al. (1999).

SURFACE WATERS

Ecozone	Location	Name	n	Description
Boreal Shield	49° 39' 40"N, 93° 43' 48"W Ontario, Canada	*IISD-Experimental Lakes Area, ON (ELA)*	**34**	Boreal forest, underlain by Precambrian bedrock with discontinuous surficial layer of sandy-gravel till. Sampled from 2010 to 2012. Ecozone mean annual temperature: 0.7 C; Ecozone total annual precipitation: 825 mm.
		Lake	21	
		Stream	11	
		Wetland Complex	2	
Mixedwood Plains	43° 30' 41"N, 80° 29' 43"W Ontario, Canada	*Grand River (GR)*	7	Surrounding land predominately agricultural (~80%) Receives wastewater treatment plant effluent from approximately ~900 000 urban residents. Sampled from 6 consecutive locations along a 90km stretch every two months from 2011 to 2012. Ecozone mean annual temperature: 6.5 C; Ecozone total annual precipitation: 940 mm.
Taiga Shield	62° 27' 14"N, 114° 22' 18"W Northwest	*Yellowknife (YW)*	7	Samples from the Taiga Shield underlain by discontinuous permafrost. Surface waters are surrounded by bedrock and peat plateau around Yellowknife. Sampled in July or October between 2013 and 2017.
		Lake	1	
		Pond	2	

Ecozone	Location	Name	n	Description
	Territories, Canada	River	*3*	Ecozone mean annual temperature: -4.8 C; Ecozone total annual precipitation: 565 mm.
		Stream	*1*	
Taiga Shield	64° 11' 24"N, 114° 11' 10"W Northwest Territories, Canada	*Wekweètì (WK)*	**7**	Situated in the Taiga Shield, below treeline, continuous permafrost. Samples taken in October of 2015 and 2016. Ecozone mean annual temperature: -4.8 C; Ecozone total annual precipitation: 565 mm.
		Lake	*6*	
		Stream	*1*	
Southern Arctic	64° 31' 29"N, 111° 40' 24"W Northwest Territories, Canada	*Daring Lake (DL)*	**19**	Found in the Southern Arctic above treeline, continuous permafrost. Ecozone mean annual temperature: -9.8 C; Ecozone total annual precipitation: 313 mm.
		Lake	*8*	
		Pond	*1*	
		Stream	*10*	

GROUNDWATERS

Ecozone	Location	Name	n	Description
Boreal Shield	47° 2' 54"N, 84° 24' 25"W Ontario, Canada	*Turkey Lakes Watershed*	**16**	Relatively un-impacted watershed in the Great Lakes-St. Lawrence forest region. Area consists of Precambrian bedrock and surficial glacial deposits of glaciofluvial outwash. Samples collected from depths ranging between 0.90 - 6.89m below surface. Ecozone mean annual temperature: 0.7 C; Ecozone total annual precipitation: 825 mm.
		Upper Reaches	*7*	
		Wetland	*9*	
Boreal Shield	49° 39' 40"N, 93° 43' 48"W Ontario, Canada	*IISD-Experimental Lakes Area, ON (ELA)*	**17**	Piezometers constructed in transect along a wetland, ranging from 0.70 - 3.85m below surface. Ecozone mean annual temperature: 0.7 C; Ecozone total annual precipitation: 825 mm.
		Wetland		
Mixedwood Plains	44° 7' 26"N, 79° 49' 12"W Ontario, Canada	*Nottawasaga River Watershed (NRW)*	**6**	Surficial deposits of glaciolacustrine deposits in an agriculturally-impacted aquifer. Samples collected from single multi-level piezometer within an unconfined surficial sand aquifer at depths of 4.35m, 5.13m, 6.68m, 9.90m, and 11.3m below surface. Ecozone mean annual temperature: 6.5 C; Ecozone total annual precipitation: 940 mm.
Atlantic Maritime	47° 6' 11"N, 67° 45' 40"W New Brunswick, Canada	*Black Brook Watershed (BBW)*	**15**	Site is an agriculturally-impacted aquifer, sampled during the summer of 2012. Surficial deposits of till and small deposits of glacial outwash. Samples taken from twelve domestic wells and three multi-level piezometers (6.1 - 30m below surface). Ecozone mean annual temperature: 4.6 C; Ecozone total annual precipitation: 1185 mm.
Taiga Shield	62° 27' 14"N, 114° 22' 18"W Northwest Territories, Canada	*Yellowknife (YW)*	**17**	Samples from the Taiga Shield underlain by discontinuous permafrost. Surface waters are surrounded by bedrock and peat plateau around Yellowknife. Sampled in July or October between 2013 and 2017. Ecozone mean annual temperature: -4.8 C; Ecozone total annual precipitation: 565 mm.
Taiga Shield	64° 11' 24"N, 114° 11' 10"W Northwest Territories, Canada	*Wekweètì (WK)*	**1**	Situated in the Taiga Shield, below treeline, continuous permafrost. Samples taken in October of 2015 and 2016. Ecozone mean annual temperature: -4.8 C; Ecozone total annual precipitation: 565 mm.

Southern Arctic	64° 31' 29"N, 111° 40' 24"W Northwest Territories, Canada	Daring Lake (DL)	4	Found in the Southern Arctic above treeline, continuous permafrost. Ecozone mean annual temperature: -9.8 C; Ecozone total annual precipitation: 313 mm.

492 *Table S2: Summary statistics for the principal component analysis used in Figure 4. Included are the LC-OCD variables and their*

493 *correlation with each principal component (PC) axis, variance explained by each PC, and eigenvalues for each PC. All code used to*

494 *generate these results can be found on github.com/paukes/canada-ecozone-DOM.*

	PC1	**PC2**	**PC3**	**PC4**	**PC5**
Correlation					
BP	0.287836	0.892915	-0.34591	0.012806	0.005893
HSF	-0.95566	-0.29157	0.011116	0.038546	0.009544
BB	0.824618	-0.28994	0.205789	-0.43995	0.005492
LMWN	0.794954	-0.33186	0.013731	0.507655	0.003726
LMWA	-0.07828	0.418298	0.89968	0.097332	0.000835
Variance					
Proportion	0.4628	0.2503	0.1943	0.09248	0.00003
Cumulative	0.4628	0.7131	0.9075	0.99997	1
Eigenvalue	2.314212	1.251482	0.971741	0.462394	0.000171

SUPPLEMENTARY FIGURES

Figure S1: Comparison of LC-OCD fractions for each water-body type between all sampling events (left graph; including

numerous sampling events at the same site) and averaged-sampling events per site (right panel; one averaged number per

unique site) for different locations (denoted by colour). A boxplot of the data is overlain by the actual data points for each site

(with random scatter on the x-axis to improve visibility of all points) and the total number of points is included as a black

number above each boxplot. Each boxplot represents the mean with 25th and 75th percentiles, with whiskers that extend up to

1.5x the inter-quartile range.

Figure S2: Original principal component analysis (PCA) that included wetland groundwater samples (pink cross symbol) with

507   high DOM and high LMWN proportions.