Temporal Analysis of T-Cell Receptor-Imposed Forces via Quantitative Single Molecule FRET Measurements

Janett Göhring[1,2*], Florian Kellner[1*], Lukas Schrangl[2*], René Platzer[1], Enrico Klotzsch[2,3,4], Hannes Stockinger[1], Johannes B. Huppa[1,‡], Gerhard J. Schütz[2,‡]

[1] Institute for Hygiene and Applied Immunology, Center for Pathophysiology, Infectiology and Immunology, Medical University of Vienna, Vienna, Austria

[2] Institute of Applied Physics, TU Wien, Vienna, Austria

[3] Laboratory of Applied Mechanobiology, Department for Health Sciences and Technology, ETH Zürich, 8093 Zürich, Switzerland

[4] Institute of Biology, Experimental Biophysics/ Mechanobiology, Humboldt Universität zu Berlin, 10115 Berlin, Germany

* Authors contributed equally

‡ please send correspondence to: johannes.huppa@meduniwien.ac.at (Johannes B. Huppa) or schuetz@iap.tuwien.ac.at (Gerhard J. Schütz)

ABSTRACT

Mechanical forces acting on ligand-engaged T-cell receptors (TCRs) have been implicated in T-cell antigen recognition. To measure these molecular forces in real time, we devised a FRET-based force sensor that is equipped with a TCR-reactive single chain antibody fragment and tethered to planar supported lipid bilayers (SLBs) which serve as surrogate antigen presenting cell. When confronting T-cells with sensor-functionalized gel-phase SLBs, we measured 4.6 pN force-peaks for activated T-cells and 5.5 pN force-peaks for non-activated scanning T-cells, which vary in the range of seconds with force loads of 1.5 pN per second. When using more physiological fluid-phase SLBs, TCR-imposed forces were found to have a ~3-fold reduction for scanning T-cells, yet no longer detectable for activated T-cells. Hence, if they are essential for ligand discrimination and TCR-proximal signaling during initial antigen scanning, TCR-exerted pulling forces are already effective below 2pN. Once activated, T-cells appear to create a synaptic environment that is devoid of noticeable TCR-exerted mechanical forces.

INTRODUCTION

A remarkable feature of the vertebrates' immune system is its inherent ability to distinguish harmful from harmless based on the primary antigen structure. T-cells embody this trait of adaptive immunity through their unique detection of antigens, which are displayed as peptide fragments in the context of the products of the major histocompatibility complex (MHC) on the surface of antigen-presenting cells (APCs). The formation of a productive immunological synapse, the area of contact between a T-cell and an APC (1), is driven by T-cell antigen receptors (TCRs) on the T-cell binding to peptide/MHC complexes (pMHC) on the APC. Importantly, T-cells specifically detect the presence of even a single antigenic pMHC molecule among millions of structurally similar yet non-stimulatory pMHCs (2). This is despite the fact that nominal pMHC-TCR interactions are of rather moderate affinity, at least when measured *in vitro*. Despite copious amounts interest, the molecular, biophysical and cellular mechanisms underlying this phenomenal sensitivity and specificity are not well understood.

Tensile forces have been implicated in the discrimination of antigenic peptides from non-activating pMHCs, especially if T-cells gauge TCR-proximal signaling based on TCR-pMHC bond lifetimes as is proposed by the kinetic proofreading model (3-6). Over the course of the last decade such forces have garnered significant attention and have even become regarded an instrumental trigger in T-cell antigen detection (7). With the use of a single-molecule Förster Resonance Energy Transfer (FRET)-based live cell imaging assay, we have observed synaptic unbinding between TCR and pMHC with significantly increased off-rates compared to TCR-pMHC binding in solution (8), possibly as the result of cellular motility. Additionally, cell mechanical (9, 10) and *in vitro* (11) studies have revealed altered TCR-pMHC unbinding under force. When confronted with beads that have been functionalized with TCR-ligands, T-cells undergo calcium signaling upon mechanical bead retraction (7, 12). Activation thresholds were found markedly reduced when pMHC-coated beads were moved tangentially to the T-cell surface (13). In line with this, T-cells reportedly push and pull against model APCs in the course of antigen-driven activation (14, 15). It has also been suggested that T-cells actively probe the elastic properties of the substrate (16), and global shear

forces of up to 150 pN have been observed when T-cells were activated via antigen-functionalized elastomeric pillar arrays (17) or polymer-embedded beads (18).

These and related findings prompted efforts to quantify mechanical forces directly within the immunological synapse. The use of DNA-based tension sensors suggested that forces larger than 12 pN acting on ligand-engaged TCRs when T-cells were confronted with antigenic glass-immobilized pMHCs (19). Forces amounting to up to 4.7 pN were reported for T-cells that interacti with planar glass-supported lipid bilayers (SLBs) featuring pMHCs anchored to gold particles (20). Furthermore, when an adjustable upper force limit was imposed on TCR-pMHC bonds with the use of rupturable DNA tension gauge, the recruitment of the TCR-proximal tyrosine kinase ZAP-70 to the plasma membrane, a process that is integral to early T-cell signaling, correlated well with the maximal amplitude of the allowed force (19).

However, current ensemble measurements render it challenging to assess pulling forces acting on individual TCRs. Typically, multiple bonds are simultaneously subjected to strain, and it is not clear whether the DNA-based tension sensors are arranged in parallel or zipper-like configuration (21). Furthermore, when considering the mechanism underlying the function of such sensors there is a strong dependence of the unzipping force on the loading rate (22) as well as on the duration of the applied force (23), which constitute *a priori* unknown parameters. If given sufficient time, all DNA duplexes will eventually separate when any force is applied. Last but not least, ensemble-based readout modalities fail to afford kinetic force profiles acting on individual receptor-ligand bonds which hence remain undetermined.

In this study, we report the development and application of a quantitative FRET-based force sensor for application within the immunological synapse, which operates at the single-molecule level. For a spring element, we employed a peptide derived from the spider silk protein flagelliform comprised of 25 amino acids with known elastic properties (24, 25). The spring peptide was anchored to either gel- or fluid-phase SLBs and conjugated to a single chain antibody fragment (scFv) derived from the TCRβ-reactive H57 monoclonal antibody (H57-scFv) to serve as a TCR-ligand. In its collapsed configuration the sensor spans a distance of less than 8nm and is therefore compatible with length restrictions of approximately 13nm as

they apply within the immunological synapse (26, 27). We observed 5 to 6 pN force-peaks per TCR for T-cells engaging gel-phase SLBs, yet only barely measurable forces below 2 pN for T-cells contacting fluid-phase SLBs. Due to the fact that both gel- and fluid-phase SLBs support T-cell stimulation, our measurements imply that perpendicular tensile forces smaller than 2 pN are already effective during the initiation of TCR signaling, yet are no longer *a sine qua non* at later stages within the more mature immunological synapse, at least at levels that are measurable with our FRET-based system.

RESULTS

Design of the force sensor

We designed the molecular force sensor (MFS) in a fashion that rendered it compatible with the use of protein-functionalized SLBs. The latter has been widely employed as surrogates of antigen presenting cells (1), especially with the incorporation of nickel-chelating lipids, such as 1,2-dioleoyl-sn-glycero-3-[(N-(5-amino-1-carboxypentyl)iminodiacetic acid)succinyl] (Ni^{2+}) (Ni-NTA-DGS), which supports decoration of SLBs with poly-histidine (His)-tagged proteins of choice in experimentally definable densities (8). The use of SLBs also allows for the application of Total Internal Reflection- (TIR) based imaging modalities, which give rise to sufficiently low background noise for recording single molecule traces.

Monovalent recombinant streptavidin (mSAv) was utilized in our study as anchor unit for the force sensor to be embedded in the SLB (**Fig. 1A, B**). To ensure stoichiometric anchorage we engineered the tetrameric mSAv complex to contain one "alive", i.e. biotin-binding subunit and three "dead" subunits, which were mutated to no longer bind biotin. Instead, dead subunits featured a tag comprised of six histidine residues each (mSAv-3xHis$_6$, for more details refer to Methods section) for attachment to SLBs employed in our study, which contained 2% Ni-NTA-DGS for functionalization with poly-histidine-tagged proteins. Depending on the experiment, the carrier lipids (98%) were either 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) giving rise to immobile, gel-like SLBs which allowed for shear and pulling forces, or 1-palmitoyl-2-oleoyl-glycero-3-phosphocholine (POPC), which enabled high lateral protein mobility, and were as a consequence permissive for pulling forces only.

We designed the MFS to support direct visualization and quantification of single-digit piconewton forces via FRET with a temporal resolution in the millisecond range, as they were expected to occur between receptor-ligand pairs within the immunological synapse (**Fig. 1A**). Integral to the MFS were two corresponding fluorophores acting as a FRET pair and flanking the sensor's core which were derived from the spider silk protein flagelliform and consisted of five repeats of the peptide motif GPGGA (24, 28). We

expected the spring to collapse with high FRET yields associated with the absence of force, yet upon force application to give rise to reduced FRET yields due to an increased distance separating the two fluorophores (25).

For a quantitative readout, the core module of the MFS had to (i) be amenable to site-specific modification with fluorophores of choice, (ii) afford directional conjugation of the spring module to the T-cell ligand and (iii) provide anchorage to SLBs (**Fig. 1A, B**). To fulfill these needs, we synthesized the spring peptide with an unpaired cysteine and a single lysine residue for fluorophore conjugation via maleimide and succinimidyl chemistry, respectively. The peptide's N-terminus was biotinylated to omit the N-terminal amino group and to confer binding of the peptide to SLB-resident mSAv-3xHis$_6$ (29). C-terminal incorporation of an ε-N$_3$ modified lysine residue in the sensor's core facilitated covalent attachment of site-specifically dibenzylcyclooctyne- (DBCO-) conjugated TCR-ligand via biorthogonal copper-free click chemistry (30).

As a TCR-engager we utilized the TCRβ-reactive recombinant H57-scF$_V$ (for construction see **Supplementary Fig.1A-D**), as we anticipated this to give rise to stable interactions with the TCRs, which unlike pMHC sustain the tensile forces applied in the synapse without rupture. Importantly, the N-terminal biotin residue within the MFS core allowed for stoichiometric coupling to mSAv-3xHis$_6$ present on the SLB (**Supplementary Fig. 1E**).

In some experimental settings we aimed to measure forces exerted by individual TCRs and to simultaneously maintain the antigenic stimulus (see below). To this end we diluted the fluorophore signals with an unlabeled force sensor termed MFS$_0$. In addition, a stretch-insensitive version of the sensor MFS$_C$ was constructed to serve as a control (MFS$_C$). In this instance the FRET acceptor and FRET donor fluorophores were separated by only a single amino acid (**Fig. 1B, C**).

Integrity of all employed constructs was confirmed by SDS-PAGE (**Supplementary Fig. 1E**). As shown in **Supplementary Fig. 1F, G**, SLBs charged with all constructs gave rise to robust T-cell activation as monitored via the rise in intracellular calcium.

Forces in the immunological synapse visualized at the ensemble level

As a source of T-cells we employed $CD4^+$ helper T-cells which had been isolated from 5c.c7 TCR transgenic mice and stimulated *in vitro* for one week in the presence of nominal antigen to promote differentiation into antigen-experienced T-cell blasts. These were then seeded onto SLBs functionalized with Intercellular Adhesion Molecule 1 (ICAM-1), B7-1, and MFS or MFS_C (50-100 molecules per μm^2). We first confronted T-cells with gel-phase SLBs prepared from DPPC, with associated proteins showing minimal diffusion ($D \leq 1.5 \times 10^{-4}$ μm^2 s^{-1}; **Supplementary Fig. 2A**). As shown by calcium imaging (**Supplementary Fig. 2B**) T-cells became readily activated in response to such SLBs. Of note, the high viscosity inherent to DPPC SLBs prevented the clustering of TCR-engaged SLB-resident MFS (**Supplementary Fig. 2C**).

Ensemble FRET efficiencies (E_{bulk}) were quantified via Donor Recovery After Acceptor Photobleaching (DRAAP) by comparing the average donor signal in a chosen region of interest before and after the photoablation of the acceptor fluorophore. A substantial reduction in E_{bulk} underneath the T-cell indicated the elongation of the force sensor (**Fig. 2A**) which we did not observe when applying the control construct. For the statistical analysis, we calculated the difference ΔE_{bulk} by subtracting the region-matched E_{bulk} recorded on a functionalized SLB in the absence of T-cells from E_{bulk} underneath the T-cell, yielding a reduction ΔE_{bulk} of -0.104 (median). In contrast, the control construct showed a significantly smaller FRET-reduction ΔE_{bulk} of -0.033 (**Fig. 2B**).

To account for the comparably high mobility of MHC, ICAM-1 and B7-1 present on the surface of living APCs (31), we confronted T-cells with fluid-phase SLBs which were prepared from 1-palmitoyl-2-oleoyl-

glycero-3-phosphocholine (POPC) and subsequently functionalized with proteins in the same fashion as the gel-phase SLBs. SLB-anchored MFS that underwent lateral diffusion with D equaling $0.72\pm0.02\mu m^2 s^{-1}$ (**Supplementary Fig. 2A**), were rapidly recruited into TCR-microclusters, transported to the center of the synapse (**Supplementary Fig. 2C**) and resulted in a robust calcium response (**Supplementary Fig. 2B**). However, we did not observe significant reductions in FRET within synaptic MFS (ΔE_{bulk}=0.000 median for the MFS and ΔE_{bulk}=0.005 for the MFS_C) (**Fig. 2B**).

Together, ensemble-based FRET experiments indicate the presence of tensile forces exerted by T-cells via their TCRs. Such forces however, were only detectable when T-cells interacted with gel- but not fluid-phase membranes. Our observations suggest that TCR-exerted force fields acted foremost tangentially to the T-cell surface: in such cases, gel-phase membranes provided adequate counterforces to stretch the MFS, while fluid-phase membranes would allow for MFS dragging without any registration of FRET changes.

Forces in the immunological synapse at the single molecule level

While being very informative on a qualitative level, there are a number of complications limiting the determination of quantitative force data from ensemble FRET measurements via the MFS. First and foremost, any information regarding the kinetics and amplitude of single-molecular forces is masked in the ensemble average, which includes in addition to sensors under tension also contributions from unbound force sensors or sensors bound to TCR-enriched microvesicles (32), rendering the quantification of individual TCR-exerted forces impossible. Moreover, defective sensors may contribute to observed ensemble FRET efficiencies in ways that are difficult to account for and predict.

To circumvent these limitations altogether, we carried out single molecule microscopy experiments, which allowed us to filter out defective sensors and quantify forces between individual TCRs and SLB-anchored MFSs in time and space with the use of both gel- and fluid-phase SLBs for T-cell stimulation. For this, T-cells were seeded onto SLBs functionalized with ICAM-1, B7-1, the unlabeled sensor MFS_0 (50-100 μm^{-}

^{2}), and low concentrations of the MFS or MFS$_C$ (<0.01 μm^{-2}). As shown in **Supplementary Fig. 2B**, the stimulatory potency of such SLBs was verified by monitoring the calcium response of interacting T-cells.

After low enough MFS densities were chosen to allow for the emergence of well isolated single molecule signals within the immunological synapse, we imaged the donor and acceptor channel at alternating excitation over time and calculated the FRET efficiency E via sensitized emission, taking into account the according cross-talk signals (see Materials & Methods for a detailed description, and **Fig. 3A** for exemplary traces) (33). We also determined the stoichiometry parameter S, which measures the fraction of donor molecules in donor-acceptor complexes (33). When plotting S against E for each observed data point, we found a single population centered around *S~0.5*, testifying to the robustness of the filtering steps for single and functionally intact MFS (**Fig. 3B**, see **Supplementary Figure 3** for more details on the procedure). In the absence of T-cells we observed for MFS and MFS$_C$ pronounced maxima at $E = 0.87$ and $E = 1.00$, respectively, which represented the FRET efficiencies of collapsed springs. The higher FRET efficiency recorded for the MFS$_C$ construct resulted in all likelihood from the shorter distance within the FRET donor-FRET acceptor dye pair, regardless of the spring extension.

Inspection of the immunological synapse revealed considerable differences in the spectral behavior of the two MFS constructs: while MFS$_C$ showed FRET efficiencies that were almost identical to those recorded in the absence of cells (E = 0.99); for MFS we detected a second peak at $E = 0.42$ with a weight of 32% in addition to the high-FRET peak at $E = 0.87$. This low-FRET peak corresponded to MFS subjected to force-induced elongation via ligated TCRs.

To read out the applied tensile forces from determined FRET efficiencies we factored in the elastic properties of the flagelliform force sensor, which were previously determined in combined single molecule force spectroscopy and FRET experiments on a similar construct (25). We inferred the relationship between FRET efficiency and the tensile force acting on the sensor used in our study (see Materials and Methods for a detailed description of the calibration) and calculated a pulling force F distributed around a peak of 4.6 pN for the low FRET peak at E equaling 0.42 (**Fig. 4A**, first panel). To assess the dynamic range of our

sensor, we estimated the minimum and maximum FRET efficiency of the low FRET peak that could be measured with the use of the MFS. The smallest detectable FRET value E_{min} was set by the donor-acceptor distance at maximum load, the Förster radius of the chosen FRET pair, and the signal to noise ratio of the data. As is laid out in more detail in the Methods section, we estimated E_{min} to amount to 0.11, which relates to a tensile force F_{max} of 10 pN. Furthermore, the largest detectable FRET value E_{max} was limited by the overlap with the high FRET peak at $E = 0.87$. With the observed proportions of the two peaks we estimated E_{max} to ~ 0.8, which related to a minimum detectable force F_{min} of about 0.9 pN. We therefore conclude that the observed force peaks fell well within the sensor's dynamic range, and that they corresponded to a not fully extended spring element.

In line with the results of the ensemble measurements (see above) we did not observe a high force peak for sensors within synapses when applying fluid-phase SLBs prepared from POPC (**Fig. 4A**, second panel**, Supplementary Fig. 4**, and **Supplementary Table 1**). MFS binding to the TCR was confirmed in single molecule tracking experiments, in which we found the majority of MFS (58%) to exhibit substantially reduced mobility underneath T-cells. In control experiments involving human Jurkat T-cells, which do not engage the MFS for lack of H57-scF$_V$-reactivity, we did not observe synaptic MFS immobilization (**Supplementary Fig**. **5**).

TCR-imposed forces under scanning conditions

Next, we aimed to assess the degree to which the high number of molecular bonds formed between the SLB and the T-cell disperses cellular forces over multiple TCR-ligand bonds while lowering the forces acting on individual MFS molecules and their associated TCRs. Of note, such parallel arrangement of multiple TCR-ligand bonds may very well apply to activated T-cells featuring a fully established synapse, yet to a much lesser extent to T-cells scanning APC surfaces for antigenic ligands, where isolated TCR-ligand interactions are likely subject to larger tensile forces. To discriminate these scenarios, we seeded T-cells onto ICAM-1- and B7-1-functionalized SLBs featuring MFS in low densities below the activation threshold

(<0.01 molecules per μm^2) (**Supplementary Fig. 2B**). Again, we detected a distinct high force peak amounting to a peak value of ~5.5 pN for T-cells confronted with gel-phase SLBs, which slightly exceeded the force peak measured for fully activated T-cells (**Fig. 4A**, third panel). We noticed a shoulder towards lower applied forces corresponding on average to 1.7 pN for T-cells contacting fluid-phase SLBs, (**Fig. 4A, fourth panel**). Next, we took advantage of the higher mobility of unbound sensors when compared to that of bound sensors, and specifically analyzed the force histograms for bound and unbound MFS (**Fig. 4B**). As expected, unbound, i.e. mobile MFS did not register any tensile forces, while immobilized, i.e. TCR-engaged sensors gave rise to a shoulder in the force peak corresponding to 2.0 pN, a value that was in agreement with the global distribution of the combined data and about three-fold below that measured for TCRs interacting with immobile MFS. Together, the data acquired at low MFS concentrations imply that perpendicular pulling forces exerted via the TCR are not a consequence of T-cell activation and may well precede the actual activation step.

Temporal analysis of forces acting on single TCR-bound MFS molecules

We finally determined the force kinetics of single TCR-bound MFS molecules. We exclusively analyzed MFS entities, which could be ascribed without ambiguity to the low-FRET peak, i.e. trajectories with at least five FRET efficiency data points below 0.6. The variability in the FRET signals within single traces was unexpectedly low: we observed an average standard deviation per trajectory of 0.12 (corresponding to force fluctuations of 1.5 pN), which was considerably lower than those derived from all low-FRET signals (equaling 0.22 and corresponding to 2.8 pN). This indicates that MFS-exerted forces covered a broad range but remained rather constant over the entire duration of the trajectories (on average 0.9s recorded at 50 frames per second). To investigate transitions in force amplitudes applied on individual TCRs in more detail, we conducted experiments at considerably lower frame rates with trajectories recorded for up to 71s on average. With decreasing frame rates, we observed more frequent force transitions within single trajectories. Accordingly, the dispersion of force values per single molecule trajectory increased towards

the dispersion of all recorded force data. Based on this and the representative single molecule FRET trajectories shown in **Fig. 5**, we conclude that transitions between FRET levels occurred on a time scale of seconds. Analysis of single molecule force transitions yielded an average loading rate of 1.5 pN s^{-1}, which persisted for approximately 2.5s before the accumulated force was released in a step-wise fashion. This behavior was reminiscent of load-fail events observed in traction-force microscopy (18), yet at 100-fold smaller peak force levels, and could be ascribed to rupture of the connection between the TCR and the cytoskeleton or between the TCR and the MFS.

DISCUSSION

In recent years, forces exerted on TCR-pMHC bonds have been frequently associated with T-cell antigen recognition (4-6, 34). The origin of these forces has been linked to actomyosin-based contractions in case of migrating T-cells, and actin reorganization during synapse formation as a consequence of T-cell activation (35). Indeed, in resting cells ~30% of the TCR are linked to the cytoskeleton (36). A clutch-like coupling between the cortical actin and microcluster-resident TCRs has been proposed for their directional transport to the synaptic center following T-cell activation (37). It is therefore conceivable that T-cells exert tensile forces onto TCR-pMHC bonds.

Our MFS-based findings are best explained by TCR-exerted forces which act largely tangentially with respect to the T-cell surface. In our T-cell stimulation regimen, fluid-phase SLBs provide an environment with substantially lower viscous drag to its embedded proteins than gel-phase SLBs. Of note, the drag γ experienced by a lipid-anchored protein in a fluid-phase membrane is smaller than $3x10^{-9}$ Pa s m (38). As a consequence, MFSs pulled with the velocity of a TCR microcluster migrating to the synaptic center ($v = 0.14$ μm s^{-1} (39)) would be subjected to a viscous force F that is lower than 0.4 fN (with $F = \gamma \cdot v$) and therefore negligible when compared to F_{min}. Thus, moving the MFS laterally within fluid-phase SLBs does not result in a drag that is sufficient to stretch the sensor. In contrast, the viscous drag encountered in a gel-phase membrane is large enough to arrest embedded MFSs. As a consequence, the MFS becomes invariably extended whenever the TCR places it under lateral load.

Pulling the MFS away from the SLB will result in its elongation. This is because the detectable maximum force F_{max} is substantially lower than the force required to pull six to nine lipids associated with the mSAv-3xHis$_6$ (40) out of the membrane (> 7 pN at loading rates >2 pN s^{-1} per single lipid (41) amounting to > 42 pN for six parallel anchors). Furthermore, extracting a biotin from its binding pocket requires more than 25 pN at loading rates larger than 1 pN s^{-1} (42).

Our observations are consistent with the reported occurrence of tangential forces detected via traction force microscopy (17, 18). In this study, TCR-exerted forces of 50 to 100 pN were detected per pillar with a

cross-sectional area of ~0.75μm². Considering a parallel configuration of the bonds and TCR densities of approximately 70 molecules per μm², as they are typical for most T-cells (43), traction forces per bond would amount to a low piconewton regime, similar to the results of our study. In contrast, Ma and colleagues reported TCR-imposed forces of 4.7 pN when T-cells contacted fluid-phase SLBs (20). However, in that study multiple DNA tension probes, and, as a consequence, multiple TCR molecules were cross-linked via the solid gold nanoparticles, allowing for tensile forces between the cross-linked TCRs.

When T-cells engage stimulatory and fluid-phase SLBs, TCR microclusters are formed in the synaptic periphery. They are concomitantly transported towards the synaptic center with a broad range of velocities amounting to 20 ± 8 nm s^{-1}, a process that was reported to be strongly affected by constraints built into the SLB (37, 44): TCR-microclusters encountering barriers in an orientation perpendicular to their movement were found arrested, while barriers positioned in intermediate angles decelerated transport (37). A frictional model was proposed based on transient links between TCRs and the cytoskeleton forming and breaking non-synchronously. We consider it plausible that our single molecule force traces shown in **Fig. 5** represent such transient links and that the properties of our gel-phase SLBs prevented the transport of ligated TCR along the T-cell surface in analogy to the previously described diffusion barriers. The sensor's stretching rate of 4 nm s^{-1} as calculated from the observed loading rate of 1.5 pN s^{-1} (for more details see Methods section, force sensor calibration, Eq. 3) corresponds to the lower end of the reported spectrum for microcluster transport rates, which probably reflects reduced cytoskeletal dynamics under load. After a force had accumulated for ~ 2.5 s, the connection between the TCR and cytoskeletal drag appeared to be released, resulting in an immediate drop in the force applied to the MFS. As is illustrated in **Fig. 5A** (third panel), the TCR repeatedly reconnected to the cytoskeleton on a time scale of minutes.

Of particular importance, we obtained evidence that molecular forces beyond 2 pN exerted via the TCR are neither a consequence of nor a precondition for T-cell signaling:

(i) We observed a high-force peak exclusively when T-cells had been seeded onto gel- but not fluid-phase SLBs, regardless of whether T-cells had been stimulated in the course of contact formation or not. For

scanning T-cells, we attribute such forces to the contractility of the actomyosin cytoskeleton, to which immobile plasma membrane-resident TCRs are tethered.

(ii) Furthermore, our results render it unlikely that forces larger than 2 pN and imposed on the TCR are required to trigger the TCR. After all, T-cells seeded onto functionalized fluid-phase SLBs did not give rise to changes in FRET of interacting MFS corresponding to forces higher than 2 pN even though they showed all the hallmarks of TCR-proximal signaling, such as elevated intracellular calcium levels, TCR-microcluster formation and TCR-movement towards the synaptic center.

Taken together, the gel-phase SLBs allow us to observe the entirety of mechanical forces (perpendicular and tangential component) experienced by the TCR, whereas the nature of the fluid-phase SLBs only allows for the creation of perpendicular forces. We found that TCR-mediated forces occur within the immunological synapse in the 4-9 pN regime only when TCR-ligands were embedded in a gel-phase SLB, indicating their predominantly shear force nature. It should be noted that perpendicular pulling forces in the range of 2 pN as we detected them with the use of fluid-phase SLBs may improve fidelity in T-cell recognition of specific versus non-specific pMHCs. This is because even such low forces affect the off-rate distribution, and likely yield improved specificity and sensitivity of antigen discrimination (4). While the MFS presented in this study was designed so that it persisted TCR-mediated pulling, the nominal pMHC ligand may dissociate prior to being exposed to the maximally possible force load. Given the high spatiotemporal resolution afforded, we expect the experimental system presented in this study to prove highly adequate and versatile to further studies delineating the role and characteristics of TCR-imposed forces in TCR-proximal signaling and ligand discrimination.

MATERIALS AND METHODS

Force sensor - anchor unit

Recombinant mSAv-3xHis$_6$ served as anchor unit for the force sensor and was produced as described (29). Briefly, biotin binding deficient subunits equipped with a C-terminal His$_6$-tag – "dead" – and functional subunits with C-terminal 3C protease cleaving site followed by Glu$_6$-tag – "alive" – were expressed as inclusion bodies in E.coli BL-21 using the pET expression system (Novagen). Inclusion bodies were dissolved in 6 M Guanidine hydrochloride at 10 mg/ml, mixed at a ratio of 3 "dead" to 1 "alive" subunit and refolded in 100 volumes PBS. The salt concentration was reduced to 2 mM by cycles of concentration using a 10 kDa cutoff spin concentrator (Amicon) and dilution with 20 mM TRIS pH 7.0. The mixture was loaded onto a MonoQ 5/50 column (GE Healthcare), and after washing the column with 100mM NaCl in 20 mM TRIS pH 7.0 mSAv-3xHis$_6$ was eluted at 20 mM TRIS pH 7.0, 240 mM NaCl. Human 3C protease (Pierce) was added and allowed to cleave the Glu$_6$-tag overnight. The digested protein was subjected to Superdex 200 (GE Healthcare) size exclusion chromatography in PBS to yield to pure anchor unit.

Force sensor - spring unit

The MFS peptide biotin-N-GG$\underline{C}$GS(GPGGA)$_5$GG$\underline{K}$YGGS-K(ϵ-N$_3$) (residues for fluorophore attachment are underlined), MFS$_C$ peptide biotin-N-GGGGS(GPGGA)$_5$GG$\underline{K}$Y$\underline{C}$GS-K(ϵ-N$_3$) and the MFS$_0$ peptide biotin-N-GGGGS (GPGGA)$_5$ GGKYG GS-K(ϵ-N$_3$) were purified via preparative C18 reverse phase HPLC (5 µm particle size, 250x21.2 mm, Agilent) applying a linear gradient of 0.1% triflouracetic acid (TFA) in deionized water (buffer A) and 0.1 % TFA, 9.9 % deionized water, 90 % acetonitrile (buffer B) over 80 minutes at a flow rate of 5 ml/min. The identity of the spring modules was confirmed via MALDI-TOF mass spectrometry (Bruker). 350 µg lyophilized peptide was dissolved in PBS at a final concentration of 5 mg/ml, mixed with Alexa Fluor 555 maleimide (Thermo Fisher Scientific) at a 1.5 fold molar surplus and then incubated for 2 hours followed by an incubation with 1.5-fold excess Alexa Fluor 647 succinimidyl ester (Thermo Fisher Scientific) in the presence of 330 mM NaHCO$_3$ for 2 hours at room temperature.

HPLC purification and lyophilization was performed after each step and successful conjugation was confirmed via mass spectrometry.

Force sensor - functional unit

TCRβ-reactive H57 antibody single chain fragments featuring a C-terminal unpaired cysteine residue (H57 cys scF$_V$) were expressed in E.coli BL-21 as inclusion bodies using the pET expression system (Novagen) and refolded as described (8, 36). The product was then concentrated using a 10 kDa cutoff spin concentrator (Amicon) in the presence of 25 µM tris-(2-carboxyethyl)phosphine hydrochloride (TCEP, Pierce). Monomeric H57 cys scF$_V$ was purified by Superdex 75 (GE Healthcare) size exclusion chromatography in PBS, then concentrated to 1 mg/ml in the presence of 25 µM TCEP and allowed to react with a 10-fold surplus of dibenzylcyclooctyne-maleimide (DBCO, Jena Bioscience) for 2 h at room temperature. Unreacted DBCO was removed by gel filtration and proteins were brought to a concentration of 1 mg/ml.

Force sensor assembly

Refolded from E.coli inclusion bodies H57-scF$_V$ were modified for sensor attachment with DBCO maleimide at a mutant unpaired C-terminal cysteine residue (**Supplementary Fig. 1A-D**). MFS peptides were mixed with DBCO conjugated proteins and 25 Vol % 2 M TRIS pH 8.0 at a molar ratio of at least 5:1 and incubated for 2 h at room temperature. Unreacted protein was removed with the use of a monomeric avidin column (Pierce). After elution with biotin, unconjugated spring was separated from MFS conjugates using size exclusion chromatography on a Superdex 75 column (GE Healthcare). mSAv-3xHis$_6$ gel shift assays documented successful conjugation as only a neglectable proportion of H57-scFv gave rise to dimers (**Supplementary Fig. 1E**). The existence of such entities may have yielded a small underestimation in

ensemble readouts. However, in the case of single-molecule experiments, multimeric structures could be directly identified and subsequently eliminated from the analysis.

T-cells

Antigen-experienced 5c.c7 murine T-cell blasts were obtained as previously described (45). Briefly, $7.5*10^6$/ml spleenocytes were stimulated with 2 μM HPLC-purified MCC peptide (ANERADLIAYLKQATK, Intavis Bioanalytical Instruments) in T-cell medium, which consisted of RPMI 1640 (Gibco) supplemented with 10 % FCS (Sigma), 100 U/ml penicillin/streptomycin (Gibco), 2 mM glutamate (Gibco), 1mM sodium pyruvate (Gibco), 1x non-essential amino acids (Gibco) and 50 μM ß-Mercaptoethanol (Gibco). Culture volume was doubled and also supplemented with 100 U/ml IL-2 (eBioscience) on day 2. T-cell cultures were expanded in a ratio of 1:1 on day 3 and 5. Dead cells were removed by centrifugation on a cushion of Histopaque-1119 (Sigma) on day 6. T-cells were used for experiments on day 7-9 after initial stimulation.

Ethical compliance statement

All animal experimentation (related to breeding, sacrifice for T-cell isolation) was evaluated by the ethics committee of the Medical University of Vienna and approved by the Federal Ministry of Science, Research and Economy, BMWFW (BMWFW-66.009/0378-WF/V/3b/2016). Animal husbandry and experimentation were performed under national laws (Federal Ministry of Science, Research and Economy, Vienna, Austria) and the ethics committee of the Medical University of Vienna and according to the guidelines of the Federation of Laboratory Animal Science Associations (FELASA).

Reagents

Hank's buffered salt solution (HBSS) and phosphate buffered saline (PBS) as well as fetal calf serum (FCS) were from Sigma Aldrich (Merck KGaA, Germany). DPPC (1,2-dipalmitoyl-sn-glycero-3-phosphocholine), POPC (1-palmitoyl-2-oleoyl-glycero-3-phosphocholine), and Ni-NTA-DGS (1,2-dioleoyl-sn-glycero-3-[(N-(5-amino-1-carboxypentyl)iminodiacetic acid)succinyl] (nickel salt)) were from Avanti Polar Lipids, Inc., USA. ICAM-1 and B7-1 were produced in Hi-5 insect cells using baculovirus and purified as described via size exclusion and ion exchange chromatography (8).

Preparation of Functionalized Lipid Bilayers

98 mol-% POPC or DPPC and 2 mol-% Ni-NTA-DGS, all dissolved in chloroform, were mixed and then dried under nitrogen flow for 20 min. The lipid mixture was suspended in 1 ml PBS at room temperature (POPC) or 55 °C (DPPC) and then sonicated for 10 min in an ultrasound bath (USC500TH, VWR, England) at the same temperature to form small unilamellar vesicles (SUVs). The SUV solution was finally diluted to 125 µM using PBS.

Cover slips (MENZEL-Gläser Deckgläser 24 × 60 mm #1.5) were treated in a plasma cleaner (PDC-002 Plasma Cleaner Harrick Plasma, Ithaca, NY, USA) for 10 min and glued to the bottom of an 8-well chamber (Nunc Lab-Tek, Thermo Scientific, USA) using duplicating silicone (Twinsil soft 18, picodent, Germany). SUVs were incubated for 20 min at room temperature (POPC) or 55 °C (DPPC) to form bilayers. After incubation, excess vesicles were washed away using PBS.

His-tagged proteins (for activating conditions: 10 ng mSAv-3xHis$_6$, 30 ng ICAM-1, 50 ng B7-1; for scanning conditions: 0.4 ng mSAv-3xHis$_6$, 30 ng ICAM-1) were added in 0.5 ml PBS to each well and incubated for 75 min at room temperature. Afterwards, the bilayer was washed with PBS.

Finally, the bilayer was incubated for 20 min with biotinylated MFS variants (for activating conditions: 10 ng MFS_0, $5 - 20$ pg MFS or MFS_C; for scanning conditions: $20 - 60$ pg MFS or MFS_C) in 0.5 ml PBS per well for binding to the mSAv-3xHis$_6$. Bilayers were washed again to remove excess MFS.

Immediately before adding cells, the buffer was exchanged for HBSS containing 2% FCS.

For ensemble experiments, molecular densities were determined by dividing the fluorescence signal per pixel by the single molecule brightness recorded at the same settings, considering the effective pixel width of 160 nm. In single molecule samples, densities were determined by counting the number of molecules and dividing by the area of the field of view.

Fura-2 labeling of T-cells

3 ml of HBSS + 2% FCS were added to $\sim 10^6$ T-cells. After centrifugation (300 RCF for 3 min), supernatant was removed and cells were incubated with 1 μg of Fura-2 AM dye (Thermo Scientific, USA) at 1mg/ml DMSO in a total volume of 100μl for 20 min on ice, and washed (addition of 5 ml HBSS + 2% FCS, centrifugation at 300 RCF for 3 min, removal of supernatant, resuspension in 100μl). Cells were kept on ice until seeding onto the functionalized bilayers.

Microscopy Experiments

Experiments were performed at room temperature, and carried out on two different microscopy systems.

Microscopy system #1: Fluorophores were excited with 640nm (OBIS 640, Coherent, CA) or 532 nm (Millenia Prime, Spectra Physics, CA) laser light coupled into an epifluorescence microscopy (Zeiss, Germany). All experiments were performed in objective-type total internal reflection (TIR) configuration, using a high-NA objective (α Plan-FLUAR 100x/1.45 oil, Zeiss, Germany) and a quad-band dichroic mirror (Di01-R405/488/532/635-25x36, Semrock Inc., USA) to separate excitation light from emission light. The

fluorescence emission path was split along wavelengths with the use of an emission beam splitter system (Optosplit II, Cairn Research, Kent, UK) employing a dichroic mirror with a cutoff wavelength of 640nm (FF640-FDi01-25x36, Semrock Inc., USA) and emission filters (ET570/60m and ET675/50m band pass, Chroma Technology Corp, USA). Both color channels were imaged side-by-side on the same chip of an EM-CCD camera (Andor iXon Ultra 897, Andor Technology Ltd, Belfast, UK). In addition, we used a 405 nm laser (iBeam Smart 405-S, Toptica Photonics AG, Germany) for visualizing the cell contours.

Microscopy system #2: Excitation was achieved by coupling a 532nm (OBIS LS, Coherent, Santa Clara, CA) and a 640 nm (iBeam smart, Toptica Photonics, Graefelfing, Germany) laser line into a T*i*-E inverted microscope (Nikon, Tokyo, Japan) via a dichroic mirror ZT405/488/532/640rpc (Chroma, Bellows Falls, VT) into a 100x objective (SR Apo TIRF, Nikon, Tokyo, Japan) in an objective-based TIR setting. Emission was split using an Optosplit II (Cairn Research, Kent, UK) equipped with a 640 nm dichroic mirror (ZT640rdc, Chroma, Bellows Falls, VT) and emission filters (ET575/50, ET655LP, Chroma, Bellows Falls, VT) and simultaneously imaged on an EM-CCD camera (Andor iXon Ultra 897, Andor Technology Ltd, Belfast, UK). Calcium imaging was performed using a xenon lamp (Lambda LS, Sutter Instrument Company) with excitation filters (340/26, 387/11, Semrock, Rochester, NY) for illumination via a 20x objective (S FLUOR, Nikon, Tokyo, Japan) and Fura-2 emission filter (ET525/50, Chroma, Bellows Falls, VT). Devices were controlled by MetaMorph imaging software (Molecular Devices, Downingtown, PA).

Microscopy system #1 was used for all single molecule experiments and for ensemble FRET measurements with the use of DPPC bilayers. Microscopy system #2 was employed for ensemble FRET experiments using POPC membranes and for calcium measurements.

Ensemble FRET measurements

FRET efficiencies were determined at the ensemble level using donor recovery after acceptor photobleaching (DRAAP). For this, an image of the donor channel was recorded before and after acceptor

photobleaching, and the pixel-averaged fluorescence signal f_{pre} and f_{post}, respectively, was calculated for each cell. The FRET efficiency was then given by $E_{bulk} = (f_{post} - f_{pre})/f_{post}$. For quantitation an area was selected underneath the T-cell and averaged, giving rise to E_{bulk}^{cell}. The same area was also recorded on lipid bilayers devoid of T-cells, analyzed for FRET efficiency, and averaged, yielding $E_{bulk}^{no\,cell}$. Finally, the difference was calculated via $\Delta E_{bulk} = E_{bulk}^{cell} - E_{bulk}^{no\,cell}$. Bleaching times were between 500ms and 1s, illumination times and delays were typically below 10ms. Single-molecule FRET measurements

Single-molecule FRET measurements

In a typical experiment we recorded 300 images alternating between green and red excitation, with illumination times t_{ill}=5ms (see **Supplementary Fig. 3B**). The delay between the green and the red color channel, t_{gr}, was kept as short as possible, typically t_{gr}=5-15ms. The delay between two consecutive green-red doublets, t_{delay}, was chosen in a range between 5ms and 975ms. For each time point, three images were recorded: donor emission upon donor excitation (Im_{DD}), acceptor emission upon donor excitation (Im_{DA}), and acceptor emission upon acceptor excitation (Im_{AA}). To convert camera counts into emitted photons, count values were multiplied by the manufacturer-provided camera efficiency at the chosen settings (15.7 photons per count) and divided by the EM gain (typically, 300).

To determine single-molecule FRET efficiencies, a series of seven processing steps was performed (**Supplementary Fig. 3** and **Supplementary Fig. 6**). Data analysis was implemented in Python 3, using the Numpy, SciPy (46) and Pandas packages for general numerical computations, matplotlib for plotting and path handling (47), scikit-learn for fitting of Gaussian mixture models (48), trackpy for single-molecule tracking (49), pims for reading image sequences from disk, and OpenCV for adaptive thresholding. Jupyter notebook and ipywidgets were used as a front-end to Python.

First: Image registration. Two-color images (Im_{DD} and Im_{DA}) were registered and corrected for chromatic aberrations using beads of 200nm diameter (TetraSpec, Invitrogen) as fiducial markers, by fitting an affine

transformation to the set of localization pairs. In our case, we transformed and interpolated the donor color channel to match the acceptor color channel ($T(Im_{DD})$).

Second: identification of candidates for single-molecule signals. We localized single-molecule signals on the sum image $T(Im_{DD}) + Im_{DA}$ and on the image Im_{AA} (50), and interlaced the resulting localizations according to the recording sequence. Taking the sum image ensured the detection of both low FRET and high FRET signals with high sensitivity. The resulting single-molecule position data were linked into trajectories using a single-molecule tracking algorithm (49), with a maximum search radius of 800nm between two consecutive frames. Only trajectories with a length ≥ 4 observations were considered. We allowed for gaps of up to three frames in the trajectories to account for bleaching and blinking of fluorophores as well as missed signals. These gaps were filled using linear interpolation between the corresponding positions before and after the missed events. As a result, we obtained the coordinates and according trajectories of all candidates for single-molecule signals in the coordinate system of the acceptor channel. For further processing, the coordinates corresponding to signals recorded in the donor channel were transformed back to the coordinate system of the donor channel via T^{-1}.

Third: determination of fluorescence brightness values. We first applied a Gaussian blur ($\sigma = 1$ pixel) to the three raw data images Im_{DD}, Im_{DA}, and Im_{AA}, which reduces noise while preserving the signal intensity (**Supplementary Fig. 3D**). To determine the single-molecule brightness, the camera counts in a 9px-diameter foreground circle around each signal's position, s_{ij}, were considered (**Supplementary Fig. 3C**). Local background, bg, was estimated by calculating the mean camera count in a ring around the signal circle (excluding any pixels affected by nearby signals). Fluorescence brightness was calculated according to $f = \sum_{i,j \in \text{circle}}(s_{ij} - bg)$. Finally, f was corrected for uneven excitation laser profile across the field of view $f \mapsto f(x,y)/profile(x,y)$. We determined $profile(x,y)$ for each laser from heavily stained samples, applied a Gaussian blur ($\sigma=10$px), and scaled to a maximum value of 1. For each signal, we hence obtained here three values: the fluorescence brightness at donor excitation in donor emission (f_{DD}), donor excitation in acceptor emission (f_{DA}), and acceptor excitation in acceptor emission (f_{AA}).

Fourth: initial filtering steps. We applied several filter steps to disregard signals which do not match stringent criteria for single-molecules. i) remove all signals from areas with <50% of the maximum laser intensity in the donor channel. ii) remove all trajectories that show potential overlap with other trajectories. Trajectories, where >25% of data points showed overlapping foreground circles, were completely disregarded. In all other trajectories, only single signals with overlapping foreground circles were removed. iii) remove all trajectories which are not present in the first frame. This step ensures that the full photobleaching profile of each trajectory is available in the subsequent steps. iv) single step transitions were identified independently in the time trace of f_{AA} and of $f_{DD}+f_{DA}$ (51). Only trajectories were selected for further processing where f_{AA} showed single step photobleaching. Also, trajectories showing partial photobleaching in $f_{DD}+f_{DA}$ were discarded.

Fifth: determination of stoichiometry factor and FRET efficiency. We calculated the corrected single-molecule FRET efficiency E and the FRET stoichiometry factor S according to Ref. (33) and (52):

$$E = \frac{f_{DA}-\alpha f_{DD}-\delta f_{AA}}{\gamma f_{DD}+f_{DA}-\alpha f_{DD}-\delta f_{AA}} \qquad \text{Equation 1}$$

$$S = \frac{\gamma f_{DD}+f_{DA}-\alpha f_{DD}-\delta f_{AA}}{\gamma f_{DD}+f_{DA}-\alpha f_{DD}-\delta f_{AA}+\frac{f_{AA}}{\beta}} \qquad \text{Equation 2}$$

The bleedthrough coefficient, α, and the coefficient describing direct acceptor excitation, δ, were determined from separate experiments using donor only and acceptor only. The detection efficiency, γ, was determined globally from single step photobleaching events according to Ref. (53) using MFS_C datasets without cells as they provide large intensity differences due to high FRET efficiencies. The excitation efficiency factor β scales f_{AA}; it is adjusted to reach S=0.5 in case of a calibration sample of known stoichiometry (54). In our case, we used MFS without cells as calibrations standard.

Sixth: final filtering steps. After the correction, we applied further filtering steps. i) only data were taken before the bleaching step of the acceptor (or donor, if donor bleaching happened prior to acceptor bleaching). ii) to exclude incorrectly synthesized force sensors or clusters in the data analysis, we removed

trajectories where more than 25% of the corresponding data showed S>0.6 (excluding two or more donors per acceptor) or S<0.35 (excluding two or more acceptors per donor). iii) for experiments with cells, we selected only data points underneath an adhered T-cell. We identified T-cells from Fura-2 images, which were recorded synchronously to the FRET signals, using an adaptive thresholding algorithm.

Seventh: fitting of the FRET histograms. 2D S-E data were fitted with a maximum likelihood fit, assuming a Bayesian Gaussian mixture model (48).

Calibration of the force sensor

The calibration of the force sensor is based on the study by (25), who found for the same flagelliform protein a linear force-distance relationship according to

$$F = \frac{r - b \cdot n - c}{a \cdot n} \qquad \qquad Equation\ 3$$

where r is the separation of the two fluorophores, b=0.044 nm the average length of a single amino acid in the collapsed state, c a constant linker length, and a=0.0122 nm pN^{-1} the compliance per amino acid. n=25 was the number of amino acids between the two fluorophores (in our case n=29). From the r-dependence of the FRET efficiency,

$$E = \frac{1}{1 + \left(\frac{r}{R_0}\right)^6} \qquad \qquad Equation\ 4$$

we determined the functional dependence of F on E. Here, R_0=5.1nm denotes the Förster distance for the FRET pair AlexaF647 and AlexaF555 (55). The parameter c in Eq. 3 can be determined from experiments with F=0, which were realized here by recording MFS on a supported lipid bilayer without cells; from the according FRET efficiency E, we calculated c=2.4 nm.

Determination of the loading rate

Single-molecule trajectories recorded on DPPC bilayers under activating conditions at 10fps were selected for transitions between low and high force levels, based on visual inspection (see **Fig. 5A,** second panel for a representative example). The selected trajectories were aligned along the time axis such that the peak forces coincided, and the rising phase was fitted with a linear function. For display, single-molecule forces were averaged for each time point.

Dynamic range of the force sensor

Simulations were implemented in Python 3. The NumPy package was used for general numerical operations as well as for random number generation.

To assess the dynamic range of the force sensor, we based our calculations on the reported elastic properties of the flagelliform protein (25). As a conservative estimate of E_{min}, we considered the MFS to follow a linear force-extension relationship up to 10 pN tensile forces. Using Eq.3 and Eq.4 this relates to a minimum FRET efficiency E_{min}=0.11.

E_{max} is limited by the difficulty to discriminate the low-FRET peak from the high-FRET peak. To quantify E_{max}, we simulated Gaussian-distributed random data sets for the high-FRET peak, taking a mean FRET efficiency of 0.87 and a standard deviation σ=0.12. For the low FRET peak, we simulated different values for the mean FRET efficiency and the standard deviation at a weight of 0.15 or 0.22, which reflects the experimental data shown in **Fig. 4A**. Samples consisting of 2300 data points were simulated and compared to a distribution lacking the low FRET peak, using a two-sample Kolmogorov-Smirnov test. The determined p-values are shown in **Supplementary Fig. 4** (corresponding values in **Supplementary Table 1**): regions of low p-values correspond to parameter settings which allow for identification of a low FRET peak. As expected, the smaller the weight of the low FRET becomes, the more difficult is its identification. As a conservative estimate we took for the estimate of the dynamic range a weight of 0.15 (corresponding

to the low FRET peak in **Supplementary Fig. 4**), yielding a threshold of approximately $E_{max}=0.8$, which corresponds to $F_{min}=0.9$ pN.

Simulation of Single-molecule Signals

To assess the quality of the Gauss blur (**Supplementary Fig. 3D**), we simulated Gauss-distributed signals on a grid, with the σ-width of the Gauss being identical to the pixel size. On each pixel, the obtained number was Poisson-distributed (to simulate shot noise), and camera noise was added by a normal distributed value, which was taken from recordings on our microscopy setup. The obtained signals were analyzed as described above, both with and without the Gauss blur.

Diffusion Analysis

To obtain diffusion coefficients in **Supplementary Fig. 2**, image sequences were acquired at an illumination time of $t_{ill} = 5$ ms and a delay of $t_{delay} = 15$ ms, 95 ms or 995 ms. Single-molecule positions were determined as for FRET analysis. Mean square displacement analysis was performed as described in (56) and fitted with the equation $MSD = 4Dt_{lag} + offset$, where D specifies the lateral diffusion coefficient, t_{lag} the analyzed time-lag, and *offset* the localization precision.

Analysis of the mobile fraction

To determine the immobilized molecules (**Supplementary Fig. 5**), we calculated the smallest enclosing circle for each trajectory using the Python implementation from Project Nayuki (https://www.nayuki.io/page/smallest-enclosing-circle). Since diffusion coefficients scale with the radius of this circle, r_c, over the square-root of the length of the trajectory, $\sqrt{t_d}$, we plotted histograms of $r_c/\sqrt{t_d}$,

yielding peaks for the immobilized molecules $r_c/\sqrt{t_d} < 0.35 \; \mu m \cdot s^{-0.5}$ and mobile molecules $r_c/\sqrt{t_d} \geq 0.35 \; \mu m \cdot s^{-0.5}$.

Statistical analysis

Statistical analysis was performed with two-sample Kolmogorov–Smirnoff tests to evaluate general distribution properties and Mann Whitney U tests to evaluate differences in the mean values of the distributions. P-values <0.01 were considered as significant, ≥0.01 as not significant (n.s.).

Code availability

We provide the Python code for single-molecule FRET analysis (https://github.com/schuetzgroup/force-fret-analysis) as well as the underlying Python library (https://github.com/schuetzgroup/sdt-python).

Acknowledgements

This work was supported by the Austrian Science Fund (FWF) projects P30214-N36 (LS, GJS) and P 25775-B2 (FK, JBH), the Boehringer Ingelheim Fonds (RP), Long-Term Fellowship from Federation of European Biochemical Societies (EK), Human Frontiers Science Program RGY0065/2017 (EK), and by the Vienna Science and Technology Fund (WWTF) LS13-030 (JG, FK, LS, JBH, GJS).

Author Contributions

JG and FK synthesized constructs. JG, FK and LS performed imaging experiments. LS developed single-molecule data analysis tools. HS and RP provided important insights. G.J.S, J.B.H. and E.K. conceived the

study. JG, FK, LS, JBH and GJS contributed to data analysis, interpretation of the results and wrote the manuscript.

REFERENCES

1.   A. Grakoui *et al.*, The immunological synapse: a molecular machine controlling T cell activation. *Science* **285**, 221-227 (1999).

2.   D. J. Irvine, M. A. Purbhoo, M. Krogsgaard, M. M. Davis, Direct observation of ligand recognition by T cells. *Nature* **419**, 845-849 (2002).

3.   T. W. McKeithan, Kinetic proofreading in T-cell receptor signal transduction. *Proc Natl Acad Sci U S A* **92**, 5042-5046 (1995).

4.   E. Klotzsch, Gerhard J. Schütz, Improved Ligand Discrimination by Force-Induced Unbinding of the T Cell Receptor from Peptide-MHC. *Biophys J* **104**, 1670-1675 (2013).

5.   Jun F. Allard, O. Dushek, D. Coombs, P. Anton van der Merwe, Mechanical Modulation of Receptor-Ligand Interactions at Cell-Cell Interfaces. *Biophys J* **102**, 1265-1273 (2012).

6.   C. Zhu, W. Chen, J. Lou, W. Rittase, K. Li, Mechanosensing through immunoreceptors. *Nature Immunology* **20**, 1269-1278 (2019).

7.   S. T. Kim *et al.*, The ab T cell receptor is an anisotropic mechanosensor. *J Biol Chem* **284**, 31028-31037 (2009).

8.   J. B. Huppa *et al.*, TCR-peptide-MHC interactions in situ show accelerated kinetics and increased affinity. *Nature* **463**, 963-967 (2010).

9.   B. Liu, W. Chen, Brian D. Evavold, C. Zhu, Accumulation of Dynamic Catch Bonds between TCR and Agonist Peptide-MHC Triggers T Cell Signaling. *Cell* **157**, 357-368 (2014).

10. J. Huang *et al.*, The kinetics of two-dimensional TCR and pMHC interactions determine T-cell responsiveness. *Nature* **464**, 932-936 (2010).

11. L. Limozin *et al.*, TCR–pMHC kinetics under force in a cell-free system show no intrinsic catch bond, but a minimal encounter duration before binding. *Proceedings of the National Academy of Sciences* **116**, 16943-16948 (2019).

12.   Y. C. Li *et al.*, Cutting Edge: mechanical forces acting on T cells immobilized via the TCR complex can trigger TCR signaling. *J Immunol* **184**, 5959-5963 (2010).

13.   Y. Feng *et al.*, Mechanosensing drives acuity of αβ T-cell recognition. *Proceedings of the National Academy of Sciences* **114**, E8204-E8213 (2017).

14.   J. Husson, K. Chemin, A. Bohineust, C. Hivroz, N. Henry, Force generation upon T cell receptor engagement. *PLoS ONE* **6**, e19680 (2011).

15.   K. H. Hu, M. J. Butte, T cell activation requires force generation. *The Journal of Cell Biology* **213**, 535-542 (2016).

16.   E. Judokusumo, E. Tabdanov, S. Kumari, Michael L. Dustin, Lance C. Kam, Mechanosensing in T Lymphocyte Activation. *Biophys J* **102**, L5-L7 (2012).

17.   K. T. Bashour *et al.*, CD28 and CD3 have complementary roles in T-cell traction forces. *Proc Natl Acad Sci U S A* **111**, 2241-2246 (2014).

18.   H. Colin-York *et al.*, Cytoskeletal Control of Antigen-Dependent T Cell Activation. *Cell Reports* **26**, 3369-3379.e3365 (2019).

19.   Y. Liu *et al.*, DNA-based nanoparticle tension sensors reveal that T-cell receptors transmit defined pN forces to their antigens for enhanced fidelity. *Proc. Natl. Acad. Sci* **113**, 5610-5615 (2016).

20.   V. P.-Y. Ma *et al.*, Ratiometric tension probes for mapping receptor forces and clustering at intermembrane junctions. *Nano Letters* 10.1021/acs.nanolett.6b01817 (2016).

21.   D. Isabey *et al.*, Force distribution on multiple bonds controls the kinetics of adhesion in stretched cells. *Journal of Biomechanics* **46**, 307-313 (2013).

22.   S. Cocco, R. Monasson, J. F. Marko, Force and kinetic barriers to unzipping of the DNA double helix. *Proceedings of the National Academy of Sciences of the United States of America* **98**, 8608-8613 (2001).

23.   M. Mosayebi, A. A. Louis, J. P. Doye, T. E. Ouldridge, Force-Induced Rupture of a DNA Duplex: From Fundamentals to Force Sensors. *ACS nano* **9**, 11993-12003 (2015).

24. N. Becker *et al.*, Molecular nanosprings in spider capture-silk threads. *Nat Mater* **2**, 278-283 (2003).

25. M. D. Brenner *et al.*, Spider Silk Peptide Is a Compact, Linear Nanospring Ideal for Intracellular Tension Sensing. *Nano Letters* **16**, 2096-2102 (2016).

26. E. Cai *et al.*, Visualizing dynamic microvillar search and stabilization during ligand detection by T cells. *Science* **356**, eaal3118 (2017).

27. K. Choudhuri, D. Wiseman, M. H. Brown, K. Gould, P. A. van der Merwe, T-cell receptor triggering is critically dependent on the dimensions of its peptide-MHC ligand. *Nature* **436**, 578-582 (2005).

28. C. Grashoff *et al.*, Measuring mechanical tension across vinculin reveals regulation of focal adhesion dynamics. *Nature* **466**, 263-266 (2010).

29. M. Howarth *et al.*, A monovalent streptavidin with a single femtomolar biotin binding site. *Nat Methods* **3**, 267-273 (2006).

30. J. C. Jewett, C. R. Bertozzi, Cu-free click cycloaddition reactions in chemical biology. *Chemical Society Reviews* **39**, 1272-1279 (2010).

31. D. R. Fooksman, Organizing MHC Class II Presentation. *Frontiers in Immunology* **5** (2014).

32. K. Choudhuri *et al.*, Polarized release of T-cell-receptor-enriched microvesicles at the immunological synapse. *Nature* **507**, 118–123 (2014).

33. A. N. Kapanidis *et al.*, Fluorescence-aided molecule sorting: analysis of structure and interactions by alternating-laser excitation of single molecules. *Proc Natl Acad Sci U S A* **101**, 8936-8941 (2004).

34. J. Hong *et al.*, A TCR mechanotransduction signaling loop induces negative selection in the thymus. *Nature Immunology* **19**, 1379-1390 (2018).

35. J. K. Burkhardt, E. Carrizosa, M. H. Shaffer, The Actin Cytoskeleton in T Cell Activation. *Annu Rev immunol* **26**, 233-259 (2008).

36.   M. Brameshuber *et al.*, Monomeric TCRs drive T cell antigen recognition. *Nature Immunology* **19**, 487–496 (2018).

37.   A. L. Demond, K. D. Mossman, T. Starr, M. L. Dustin, J. T. Groves, T Cell Receptor Microcluster Transport through Molecular Mazes Reveals Mechanism of Translocation. *Biophys J* **94**, 3286-3292 (2008).

38.   A. Pralle, P. Keller, E. L. Florin, K. Simons, J. K. Horber, Sphingolipid-cholesterol rafts diffuse as small entities in the plasma membrane of mammalian cells. *J Cell Biol* **148**, 997-1008 (2000).

39.   Y. Kaizuka, A. D. Douglass, R. Varma, M. L. Dustin, R. D. Vale, Mechanisms for segregating T cell receptor and adhesion molecules during immunological synapse formation in Jurkat T cells. *Proc Natl Acad Sci U S A* **104**, 20296-20301 (2007).

40.   E. Hochuli, H. Döbeli, A. Schacher, New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues. *Journal of Chromatography A* **411**, 177-184 (1987).

41.   E. Evans, F. Ludwig, Dynamic strengths of molecular anchoring and material cohesion in fluid biomembranes. *Journal of Physics: Condensed Matter* **12**, A315-A320 (2000).

42.   R. Merkel, P. Nassoy, A. Leung, K. Ritchie, E. Evans, Energy landscapes of receptor-ligand bonds explored with dynamic force spectroscopy. *Nature* **397**, 50-53 (1999).

43.   B. Rossboth *et al.*, TCRs are randomly distributed on the plasma membrane of resting antigen-experienced T cells. *Nature Immunology* **19**, 821-827 (2018).

44.   T. Yokosuka *et al.*, Newly generated T cell receptor microclusters initiate and sustain T cell activation by recruitment of Zap70 and SLP-76. *Nat Immunol* **6**, 1253-1262 (2005).

45.   J. B. Huppa, M. Gleimer, C. Sumen, M. M. Davis, Continuous T cell receptor signaling required for synapse maintenance and full effector potential. *Nat Immunol* **4**, 749-755 (2003).

46.   P. Virtanen *et al.*, SciPy 1.0--Fundamental Algorithms for Scientific Computing in Python. *arXiv e-prints*, arXiv:1907.10121-arXiv:11907.10121 (2019).

47.   J. D. Hunter, Matplotlib: A 2D graphics environment. *Computing in Science & Engineering* **9**, 90-95 (2007).

48.   F. Pedregosa *et al.*, Scikit-learn: Machine Learning in Python. *J Mach Learn Res* **12**, 2825-2830 (2011).

49.   D. B. Allan, T. Caswell, N. C. Keim, C. M. van der Wel, trackpy: Trackpy v0.4.1. *Zenodo*, http://doi.org/10.5281/zenodo.1226458 (2018).

50.   H. Babcock, Y. M. Sigal, X. Zhuang, A high-density 3D localization algorithm for stochastic optical reconstruction microscopy. *Optical Nanoscopy* **1**, 6 (2012).

51.   R. Killick, P. Fearnhead, I. A. Eckley, Optimal Detection of Changepoints With a Linear Computational Cost. *J Am Stat Assoc* **107**, 1590-1598 (2012).

52.   B. Hellenkamp *et al.*, Precision and accuracy of single-molecule FRET measurements—a multi-laboratory benchmark study. *Nature Methods* **15**, 669-676 (2018).

53.   J. J. McCann, U. B. Choi, L. Zheng, K. Weninger, M. E. Bowen, Optimizing methods to recover absolute FRET efficiency from immobilized single molecules. *Biophys J* **99**, 961-970 (2010).

54.   N. K. Lee *et al.*, Accurate FRET Measurements within Single Diffusing Biomolecules Using Alternating-Laser Excitation. *Biophys J* **88**, 2939-2953 (2005).

55.   M. Spence, I. D. Johnson, *The molecular probes handbook: a guide to fluorescent probes and labeling technologies* (Life Technologies Corporation, 2010).

56.   S. Wieser, G. J. Schütz, Tracking single molecules in the live cell plasma membrane-Do's and Don't's. *Methods* **46**, 131-140 (2008).

FIGURE LEGENDS

Figure 1: FRET-based sensor designed to detect forces on a single-molecule level.

(**A**) Schematic illustration of the force sensor. (**B**) Exertion of forces stretches the spring core module which reduces FRET yields for the molecular force sensor (MFS) but not for the control force sensor (MFS_C). (**C**) Scheme of the molecular force sensors including sequence information. A cysteine residue (red) and a lysin residue (blue) were used to conjugate the FRET donor Alexa Fluor 555 and the FRET acceptor Alexa Fluor 647, respectively. In the unlabeled molecular force sensor (MFS_0) the cysteine residue was mutated to glycine to avoid disulfide bond formation.

Figure 2: Ensemble FRET readout reveals TCR-mediated tensile forces.

(**A**) Representative images illustrating FRET efficiencies determined on a pixel-by-pixel basis via FRET Donor Recovery after FRET Acceptor Photobleaching (DRAAP) on DPPC (gel-phase) and POPC (fluid-phase) SLBs with the use of the MFS or the MFS_C construct. (**B**) Statistical analysis of ensemble FRET measurements. Each data point corresponds to a single DRAAP experiment of up to three cells in the field of view; boxes indicate the interquartile range (first quartile to third quartile), whiskers extend 1.5 times the interquartile range from the first and third quartile, the line indicates the median. A significant reduction in FRET efficiency (ΔE_{bulk}) was only observed for MFS on DPPC SLBs (n.s., not significant; two-sample Mann Whitney U test). n = 68 cells for MFS on DPPC, n = 24 for MFS on POPC, n = 27 for MFS_C on DPPC, n = 21 for MFS_C on POPC.

Figure 3: Reading out the force sensor at the single-molecule level

(**A**) Representative single-molecule signals and time traces. The left panel depicts the fluorescence signals prior to and after photobleaching of the FRET acceptor (scale bar 1 μm). The middle panel shows single-molecule brightness values of the FRET donor, the FRET signal itself and the FRET acceptor. Bleaching of the acceptor could be observed as a correlated step-wise decrease of FRET and FRET acceptor intensities and an increase in FRET donor intensity. The right panel shows the stoichiometry and FRET efficiency. (**B**) FRET efficiency – stoichiometry (*E-S*) analysis of single-molecule datasets obtained for MFS and MFS$_C$ present on DPPC SLBs in the absence and presence of T-cells. Each data point in the depicted cloud corresponds to one single-molecule FRET event. Measured FRET efficiencies *E* are indicated in histograms (right panels). A Gaussian mixture model was fitted to the *E-S* data, yielding pronounced maxima at *E* = 0.86 and *E* = 0.42 (MFS in the presence of T-cells), *E* = 0.87 (MFS in the absence of T-cells), *E* = 0.99 (MFS$_C$ in the presence of T-cells), and E = 1.00 (MFS$_C$ in the absence of T-cells). In the presence of T-cells the low FRET peak corresponds to a weight of 32%. n = 5313 from 265 movies for MFS in the presence of T-cells, n = 5313 from 90 movies for MFS in the absence of T-cells, n = 1451 from 127 movies for MFS$_C$ in the presence of T-cells, n = 1506 from 30 movies for MFS$_C$ in the absence of T-cells.

Figure 4: Single-molecule pulling forces imposed on individual TCRs

(**A**) Single-molecule forces obtained under conditions under which T-cells became fully activated or for non-activated scanning T-cells confronted with DPPC (gel-phase) or POPC (fluid-phase) SLBs. A high force peak emerged for activated or scanning T-cells confronted with DPPC SLBs at *F* = 4.6 pN and *F* = 5.5 pN, which corresponded to 32% and 14% of observed events, respectively. For T-cells scanning POPC SLBs we observed an additional high force peak at *F* = 1.7 pN with 21% weight (p<10^{-2}). n = 5313 (265 movies) for MFS on DPPC SLBs in contact with activated T-cells, n = 5313 (90 movies) for MFS on fully stimulatory DPPC without T-cells, n = 4654 (323 movies) for MFS on DPPC SLBs including scanning T-cells, n = 4169 (130 movies) for MFS on DPPC SLBs (non-stimulatory) without T-cells, n = 1161 (293 movies) for MFS on POPC interacting with activated T-cells, n = 1528 (90 movies) for MFS on stimulatory

POPC without T-cells, n = 2147 (333 movies) for MFS on POPC SLBs in contact with scanning T-cells, n = 1796 (70 movies) for MFS on non-stimulatory POPC without T-cells. (**B**) Bound MFS were distinguished from unbound MFS on POPC SLBs under scanning conditions by means of the single molecule mobility as described in **Supplementary Fig. 5**. The dashed line indicates the chosen boundary drawn between mobile and immobilized MFS. Unbound MFS (termed "mobile") gave rise to a force distribution showing no significant difference from the control scenario recorded without cells. Bound MFS (termed "immobilized") displayed a high-force shoulder similar to **Fig. 4A**, fourth panel**.**

Figure 5: Temporal analysis of TCR-applied forces observed for activated T-cells confronted with DPPC SLBs

(**A**) Representative single-molecule force traces recorded at indicated frame rates. We observed transitions within force trajectories only in traces recorded at low frame rates, whereas trajectories recorded at high frame rates showed rather constant force levels. (**B**) For quantitative readout, we determined the force fluctuations of each trajectory as standard deviation, calculated the average over all trajectories (termed "std_{local}"), and divided the number by the global standard deviation of all measured force values (termed "std_{global}"). With higher frame rates deviation from ergodic behavior was observed. Error bars are standard errors of the mean. 1880 events in 40 tracks from 265 movies in dataset (50 fps), 1823 events in 38 tracks from 368 movies in dataset (10 fps), 649 events in 10 tracks from 108 movies in dataset (1 fps). (**C**) Assessment of the loading rate for single-molecular force transitions recorded at 10fps. The panel displays pooled analysis of 470 data points taken from 7 single-molecule trajectories. A linear fit (orange line) yielded a loading rate of 1.5 pN s^{-1}.

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1: Synthesis and functional validation of the TCR-specific force sensor

(**A**) Refolded single chain antibody fragment derived from the TCRβ-reactive H57 monoclonal antibody (H57-scF$_V$) was purified by Superdex 200 10/300 GL (S200) size exclusion chromatography. The gray area corresponds to collected fractions. (**B**) Post conjugation, the surplus of DBCO was removed from monomeric H57-scF$_V$-DBCO via Superdex 75 10/300 GL (S75) gel filtration. (**C**) Unconjugated MFS was separated from H57-MFS via S75 gel filtration. (**D**) Mass spectra of HPLC purified unlabeled, Alexa Flour 555-labeled and Alexa 555 / Alexa 647 dual-labeled MFS (top to bottom) confirmed successful conjugation. The theoretical molar weight of the unconjugated peptide is 3172 Da. (**E**) SDS-PAGE gel shift analysis of the force sensor in the presence of mSAv-3xHis$_6$ performed under non-boiling and non-reducing conditions testifies to the complete conjugation of the force sensor. A second slight band corresponding to a double shift and a third barely visible band corresponding to triple shift indicated the presence of a small fraction of H57-scF$_V$ conjugated to more than one sensor peptide. (**F, G**) 5c.c7 TCR transgenic T-cell blasts were brought into contact with fluid-phase SLB decorated with 100-250 molecules μm^{-2} of force sensor bound to mSAv-3xHis$_6$ together with ICAM-1-His$_{12}$ and B7-1-His$_{12}$. Fura-2 emission was measured after 340 and 380 nm excitation. Increase of intracellular calcium levels measured by ratiometric Fura-2 analysis shows the stimulatory potency of the H57-scF$_V$-based force sensor. SLBs lacking the force sensors served as negative control (only mSAv-3xHis$_6$, ICAM-1-His$_{12}$ and B7-1-His$_{12}$). Median per time point normalized by median of negative control is shown, $n \geq 100$ cells.

Supplementary Figure 2: Diffusion analysis of the force sensor anchored to DPPC and POPC and functional response of T-cells in contact with DPPC and POPC SLB displaying the force sensor.

(**A**) Diffusion analysis of MFS on DPPC membranes yielded $D = 5.8 \cdot 10^{-5} \pm 8.6 \cdot 10^{-5}$ µm² s^{-1} in the presence and $D = 1.3 \cdot 10^{-4} \pm 6 \cdot 10^{-5}$ µm² s^{-1} in the absence of T-cells (left panel). Analogous analysis employing POPC SLBs gave rise to $D = 0.27 \pm 0.02$ µm² s^{-1} in the presence and $D = 0.72 \pm 0.02$ µm² s^{-1} in the absence of T-cells (right panel). Reduced MFS mobility in the synapse arose from the presence of bound and immobilized molecules (compare **Supplementary Figure 5**). 2945 datapoints in 90 tracks from 77 movies for DPPC underneath cells, 3867 datapoints in 102 tracks from 30 movies for DPPC without cells, 724 datapoints in 29 tracks from 109 movies for POPC underneath cells, 904 datapoints in 31 tracks from 30 movies for POPC without cells. (**B**) Ratiometric calcium imaging of T-cells confronted with gel-phase DPPC SLBs (left panel) and fluid-phase POPC SLBs (right panel). All SLBs employed featured ICAM-1 and B7-1. In addition, one of the following (combination of) constructs were added: His-tagged IE^k-MCC (blue), high density of MFS$_0$ + single-molecule density of MFS$_C$ (green), high density of MFS$_0$ + single-molecule density of MFS (red), single-molecule density of MFS only (orange), no additional proteins (black). We used between 72 and 202 cells for analysis. (**C**) TIRF imaging of MFS employed in high densities showed microcluster formation within the immunological synapse of T-cells in contact with POPC SLBs (right panel) but not of T-cells engaging DPPC SLBs (left panel).

Supplementary Figure 3: Quantitation of the single-molecule FRET signal

(**A**) Determination of the lower boundary, F_{min}, of the sensitivity towards applied forces conveyed by the MFS. We simulated Gaussian-distributed random data sets for the high-FRET peak, taking a mean FRET efficiency of 0.87 and a standard deviation $\sigma = 0.12$. For the low FRET peak, we simulated different values for the mean FRET efficiency and the standard deviation (**i**). Samples were compared to a distribution lacking the low FRET peak using a two-sample Kolmogorov-Smirnov test. Determined p-values were plotted for the different parameter settings. Panel (**ii**) shows results assuming a statistical weight of 0.15 for the low FRET peak, panel (**iii**) for a statistical weight of 0.22. Dashed lines indicate the results from the experiments shown in **Fig. 4A,** fourth panel (mean = 0.72, std = 0.28). (**B**) Illumination timing. Each image

sequence is a repetition of two consecutive images recorded with 532 nm and 640 nm excitation, yielding Im_{DD}, Im_{DA}, and Im_{AA}. The timings are indicated in the image. Prior to this, samples were illuminated with 405nm to obtain one image of the outline of cell in the Fura-2 channel. (**C**) Mask used for brightness analysis. If a molecule is located in the orange position, we calculated its brightness by summing over the yellow pixels, and subtracted the weighted blue pixels as background (top). If background of one signal overlaps with the foreground of another signal (bottom image), the corresponding affected pixels (red) were omitted in the analysis. (**D**) Gaussian blur reduces noise in brightness determination (left). For dim signals, noise is decreased by 25 % to 30 %. The blur leads to slight underestimation of the brightness (typically < 1 %, right). Since this is constant over the brightness range, derived ratiometric quantities, like E and S, are not biased.

Supplementary Figure 4: T-cells scanning for antigen embedded in fluid-phase POPC SLBs cause mild force (~ 1.7 pN) on ligand-engaged TCRs

Histograms of FRET efficiencies *E* for MFS on DPPC- or POPC-derived SLBs under conditions that led to full T-cell activation (upper panels) or caused T-cells to scan for antigen (lower panels). Data sets are identical to data sets employed in **Fig. 4**.

Supplementary Figure 5: MFS mobility on POPC SLBs is reduced after synaptic TCR binding

Mobility of MFS on POPC SLBs analyzed in a synapse involving 5c.c7 T-cells (left) or Jurkat T-cells serving as a negative control (right). Panel (**A**) shows selected trajectories, panel (**B**) a histogram of the calculated extension of the trajectory, *r*, normalized by the square root of its duration, t_d. The selection threshold is indicated as dashed line. Diffusion analysis of MFS yielded: D=0.018 ± 0.007 μm² s^{-1} and D=0.89 ± 0.17 μm² s^{-1} for the immobile and mobile fraction in a synapse with 5c.c7 T-cells, respectively,

and D=0.71 ± 0.1 μm² s^{-1} for the mobile fraction in a synapse with Jurkat T-cells. 724 events in 29 tracks from 109 movies for 5c.c7 cells, 561 events in 20 tracks from 68 movies for Jurkat cells.

Supplementary Figure: 6: FRET analysis workflow

Flow chart describing the single-molecule FRET analysis workflow. Subheadings relate to the description in the Materials and Methods section.

Supplementary Table 1: The results of a fit with a two-component Gaussian mixture model yielded the results shown in the table. n.s., not significant; two-sample Kolmogorov–Smirnoff test.

Figure 1

A

H57 (anti-TCRβ)-single chain antibody fragment (H57-scF$_V$) exhibiting a C-terminal cysteine residue

covalent linkage via click chemistry

Alexa Fluor 647 (FRET acceptor) coupled to a lysine residue

flagelliform spidersilk spring (GPGGA)$_5$ core module

Alexa Fluor 555 (FRET acceptor) coupled to a cysteine residue

N-terminal biotin

mSAv-3 x His$_6$ for SLB anchorage

B

CD4+ T-cell

force

LFA-1

TCR-CD3 complex

H57-scF$_V$

high FRET

ICAM-1

low FRET

high FRET

H57-scF$_V$-MFS: force-sensor with distal FRET donor

H57-scF$_V$-MFS$_C$: sensor with proximal FRET donor

C

H57-scF$_V$-MFS   H57-scF$_V$-MFS$_C$ H57-scF$_V$-MFS$_0$

MFS: biotin-N-GG**C**GS (GPGGA)$_5$ GG**K**YG GS-K(ε-N$_3$) + H57-scF$_V$-DBCO

MFS$_C$:   biotin-N-GGGGS (GPGGA)$_5$ GG**K**Y**C** GS-K(ε-N$_3$) + H57-scF$_V$-DBCO

MFS$_0$:   biotin-N-GGGGS (GPGGA)$_5$ GG**K**YG GS-K(ε-N$_3$) + H57-scF$_V$-DBCO

Figure 2

A

MFS   MFS$_c$ MFS   MFS$_c$
DPPC   DPPC   POPC   POPC

B

A

B

Figure 4

A

B

POPC scanning

Figure 5

A

50 FPS   10 FPS   1 FPS

B   **C**

Supplementary Figure 1

Supplementary Figure 5

A

Supplementary Figure 6

Supplementary Table 1

condition		low-FRET component			high-FRET component		
		mean	std.	weight	mean	std.	weight
DPPC activating	inside synapse	0.42	0.19	0.32	0.87	0.11	0.68
	no T-cells	0.81	0.35	0.07	0.87	0.10	0.93
POPC activating	inside synapse	0.81	0.32	0.05	0.86	0.12	0.95
	no T-cells	0.86	0.10	0.90	0.92	0.25	0.10
DPPC scanning	inside synapse	0.34	0.16	0.14	0.86	0.12	0.86
	no T-cells	0.78	0.25	0.06	0.87	0.09	0.94
POPC scanning	inside synapse	0.72	0.28	0.21	0.84	0.12	0.79
	no T-cells	0.86	0.10	0.92	0.93	0.26	0.08