Brain networks involved in empathy for positive and negative emotions in younger and older adults

Abbreviated title: Empathy and aging

Maryam Ziaei[1*], David C. Reutens[1#] & Natalie C. Ebner[2,3,4 #]

[1] Centre for Advanced Imaging, the University of Queensland, Brisbane, Australia

[2] Department of Psychology, University of Florida, Florida, USA

[3] Department of Aging and Geriatric Research, Institute on Aging, University of Florida, Florida, USA

[4] Center for Cognitive Aging and Memory, Department of Clinical and Health Psychology, University of Florida, Gainesville, FL, USA

*Corresponding Author: Maryam Ziaei; Centre for Advanced Imaging, The University of Queensland, St. Lucia, QLD 4072, Australia; Email: maryamziae@gmail.com and maryam.ziaei@cai.uq.edu.au; Tel: +61- 422 916 362

# denotes joint senior authorship

Acknowledgment

The authors would like to thank the participants for their time and acknowledge the practical support provided by the imaging staff at the Centre for Advanced Imaging. We would also like to thank Ms. Megan Campbell and Nicola Pease for their help during the data collection. This work was supported by New Staff Research Funding from the Centre for Advanced Imaging. The authors declare no competing financial interests.

Abstract

Empathy, among other social-cognitive processes, changes across adulthood. More specifically, cognitive components of empathy (understanding another's perspective) appear to decline with age, while findings for affective empathy (sharing another's emotional states) are rather mixed. Not well understood yet are the neural correlates underlying cognitive and affective empathy and the extent to which valence modulates these components of empathy in aging. To fill these research gaps, young and older participants completed a modified version of the Multifaceted Empathy Test, which measures both cognitive and affective empathy, while brain images were acquired in the MRI scanner. Multivariate analysis showed that regions of the salience network, including anterior insula and anterior cingulate, were more involved in cognitive empathy to negative emotions, in older than in younger participants. For affective empathy to positive emotions, in contrast, young and older participants recruited a similar brain network including the main nodes of the default mode network. These results provide novel insights into the neural networks subserving cognitive and affective empathy in young and older adults and highlight a modulatory role of valence on empathy in aging. Our findings highlight the importance of valence for the study of higher-order social-cognitive processes such as empathy in aging.

Keywords: empathy, aging, salience network, default network, PLS, multivariate

Introduction

Empathic response is a social-cognitive capacity and refers to the understanding of and response to others' unique experience of emotions (Decety & Jackson, 2006). Empathy contributes to successful social interaction and enhances prosociality and positive health outcomes (Beadle & de la Vega, 2019; Sun et al., 2019). The literature often distinguishes between two components of empathy: *cognitive* empathy, the ability to decode and understand another's perspective, and *affective* empathy, the affective sharing of another's emotional states (Singer & Lamm, 2009).

Empathy in aging

Mounting evidence suggests age-related decline in social-cognitive capacities including perception of eye gaze (Slessor et al., 2010; Ziaei et al., 2016), emotional facial expression (Ruffman et al., 2008), theory of mind (Henry et al., 2013), and social behavior (von Hippel & Dunlop, 2005). Compared to other components of social cognition, empathy has received considerably less attention in aging research. Furthermore, the few existing studies on age-related differences in empathic response have almost exclusively used self-report and have captured more trait-like aspects of the construct (e.g., using the Interpersonal Reactivity Index; Davis, 1983). Taken together, these previous studies largely agree that aging is associated with decline in cognitive empathy (Beadle et al., 2019; Henry et al., 2013). Evidence is less clear for affective empathy for which some studies suggest no age-related differences (Bailey et al., 2008; Beadle et al., 2012) and other studies support an increase in affective empathy with age (Grühn et al., 2008; O'Brien et al., 2012).

Even more sparse are studies measuring state empathy in aging, for example by using experimentally induced alterations in the state of affective empathy. In particular, Sze and

colleagues (2012) manipulated empathic response by showing uplifting or distressing films and reported an age-related linear increase in empathic concern and personal distress in response to both types of films. Similarly, Bailey et al., 2018) found increased emotional distress and reactivity towards another's pain, and this enhanced affective empathy predicted prosociality (i.e., willingness to help). Finally, Beadle et al., 2015) did not find evidence for age-group differences in affective empathy or personal distress in response to a note by cancer patients describing their experiences with the disease.

Thus, the current research consistently suggests that cognitive empathy is lower in older than in younger adults. Observed age effects on affective empathy, however, are more mixed with some studies reporting age-related increase, others age-related decrease, and yet others no differences in affective empathy between the age groups. Also, not yet well understood are the neural substrates subserving cognitive and affective empathy, particularly in older adults.

Neural correlates of empathy

Anterior insula, mid and dorsal anterior cingulate gyrus, and temporo-parietal junction have been identified as key areas involved in empathy (Bernhardt & Singer, 2012; Bzdok et al., 2012b; Decety et al., 2006; Lamm et al., 2011). However, to date, only two studies have investigated the neural substrates underlying empathy in older adults. In particular, Chen and colleagues (2014) asked participants to rate their feelings towards another's pain and found an age-related decrease in activation of insula and anterior cingulate during this task. Of note, while recent meta-analytic evidence supports similarities in the neural correlates of empathy in response to pain relative to non-pain stimuli (Timmers et al., 2018), processing of empathy to non-pain stimuli has not been studied in older adults, and will be addressed in the present study.

The second study that explored neural substrates underlying empathy in young and older adults measured both self-reported cognitive and affective empathy and correlated those measures with brain activity during cognitive and affective tasks (e.g., the N-back and a facial affect matching task; (Moore et al., 2015). Using a region of interest analysis, they found that higher levels of affective empathy were associated with deactivation of insula and amygdala during the performance of an N-back working memory task. Of note, Moore and colleagues measured empathic response outside the MRI scanner, and thus their study cannot speak directly to age-related differences in the neural substrates subserving cognitive and affective empathy, as specifically targeted in the present study. Also, given that empathy is a complex and multidimensional process, it is likely that empathic response activates large-scale brain networks and not just individual circumscribed regions as examined in previous studies. To fill this research gap, the present study adopted a multivariate analysis approach that allowed delineation of neural processes underlying cognitive and affective empathy among young and older adults.

Valence modulation on empathic response in aging

Robust evidence supports that emotional information has prioritized access to further processing over neutral information. This phenomenon is termed 'emotional enhancement' and affects cognitive processes such as memory and attention, generally and also specifically in aging (Charles et al., 2003; Leigland et al., 2004; Mather & Carstensen, 2003). Further qualifying the emotional enhancement effect, both neuroimaging and behavioural studies over the last two decades have reliably documented that older compared to younger adults show enhanced processing of positive over negative information, such as reflected in greater attention to, and memory for, positive over negative stimuli, termed the "positivity effect" in aging (Mather et al., 2003; Reed & Carstensen, 2012; Ziaei & Fischer, 2016; Ziaei et al., 2015).

While the phenomena of emotional enhancement and the positivity effect in aging have been extensively studied in the context of facial cue processing, there has been no investigation to date of the impact of valence on empathic response in aging. Evidence in younger adults supports the distinction of empathic response to positive vs. negative stimuli, known as positive vs. negative empathy (Morelli et al., 2015). Given the robust support for age-related differences in processing emotional information, it is reasonable to assume that the valence of stimuli modulates empathic response differentially in young and older adults. The current study investigated this hypothesis.

Current study

This study examined age-related differences in cognitive and affective empathy both on the level of behavior and of its underlying neural substrates. The study furthermore addressed the modulatory role of valence on empathic response in young and older adults. We administered a modified version of the Multifaceted Empathy Test (MET) (Dziobek et al., 2011) that measured both cognitive and affective empathy. Following Mazza et al. (2015), we presented positive, negative, and neutral face images to allow for a systematic investigation of the effects of valence on empathic response. We added an age perception condition to allow for a non-empathy control.

Predictions regarding age-related differences in cognitive and affective empathy. Based on previous studies (see Beadle & de la Vega, 2019), we predicted that older compared to younger participants would perform more poorly (reflected in longer response times) in cognitive empathy. We also predicted that reduced performance during cognitive empathy would be reflected in greater recruitment of core empathic regions such as anterior cingulate and insula (Fan et al., 2011) as well as dorsolateral and inferior frontal, posterior cingulate, and temporoparietal junction (Bzdok et al., 2012a) among older compared to younger participants.

In contrast, given the somewhat conflicting evidence regarding age-related differences in affective empathy, we expected that older participants would perform comparably to or even better than younger participants in the affective empathy condition (as reflected in equal or faster response time). Given limited evidence on brain responses during affective empathy (especially for non-pain stimuli), we entertained two alternative hypotheses: (a) age-related decreased activity of core empathy regions, such as insula during affective empathy (Fan et al., 2011); reflecting decline in internal bodily perception in older adults, impacting on affective empathy (Mendes, 2010); or (b) no age-related differences in anterior cingulate activity during affective empathy, reflecting brain maintenance in aging (Nyberg et al., 2012). In particular, given individual differences in brain changes with age, it is possible that brain integrity is preserved in some older adults, resulting in no differences in neural processing during affective empathy between young and older adults, confirming the brain maintenance hypothesis.

Predictions regarding age-related differences in valence modulation on cognitive and affective empathy. Based on robust evidence of a positivity effect in aging (Reed et al., 2014), we anticipated that empathic response would be comparable, or even better (as reflected in equal or faster response times) in older than in younger participants to positive but not to negative stimuli; we expected this effect to hold for both cognitive and affective empathy. With respect to brain patterns during processing of positive stimuli (both during cognitive and affective empathy), we predicted greater activity in brain areas such as posterior cingulate or medial prefrontal cortex for older compared to younger participants. These regions constitute hubs of the default mode network (Buckner et al., 2008) and were expected to be involved due to reduced cognitive effort and enhanced self-referential processing for positive emotions among older adults (Lin et al., 2017; Ziaei, Burianova, et al., 2016). In contrast, for negative stimuli (both during cognitive and

affective empathy), we expected that older compared to younger participants would recruit additional prefrontal and cognitive control areas, such as inferior frontal gyrus and dorsolateral prefrontal cortex, reflecting enhanced cognitive demand.

Method

Participants

Twenty-six younger and 25 older adults participated in this study. Due to large head movement (> 1.5mm), two younger participants were excluded, leaving 24 younger ($M = 21.81$, $SD = 4.06$) and 25 older ($M = 71.52$, $SD = 3.80$) participants for final data analysis. All younger participants were University of Queensland undergraduate students who were reimbursed either with course credits or AUD$20 per hour. Older participants were volunteers from the community recruited through advertising in public notice boards of different clubs, libraries, churches, and the University of Queensland's Aging Mind Initiative. Older participants were reimbursed with AUD$20 per hour.

All participants were right-handed, English speakers who had normal or corrected-to-normal vision using MRI compatible glasses, and no history of psychiatric illnesses, cardiovascular disease, head or heart surgery, or neurological impairment (e.g., epilepsy). The age groups were comparable in years of education and gender distribution (Table 1). All older participants scored above the recommended cut-off of 27 ($M = 28.76$, $SD = 1.26$) on the Mini Mental State Examination (Folstein et al., 1975).

Procedure

The experiment was approved by the Royal Brisbane and Women's Hospital and the University of Queensland Research Ethics Committees. All participants provided written informed consent prior to enrollment. The study comprised a 1-hour MRI session, followed by a

2-hour behavioral/neuropsychological assessment on the same day. Prior to the MRI, participants received verbal instruction about the MET (described next) and worked on practice trials for familiarization with the trial timing and task sequence. After the MRI, participants completed a series of background measures (described below), were debriefed, and received reimbursement.

Multifaceted Empathy Task (MET)

We used a modified version of the MET (Dziobek et al., 2011). As depicted in Figure 1, the MET consists of naturalistic face images for which participants were asked to indicate (i) what kind of emotion the depicted person feels (cognitive empathy condition), (ii) how strong the emotion they (the participants) feel about the depicted person (affective empathy condition), and (iii) how old the depicted person is (age perception control condition). Following Mazza et al. (2015), positive (happy), negative (sad, angry), or neutral face images were presented in each condition and, different from the original Dziobek et al. (2011) design, we used a block design and included three response options. Each block included seven pictures of the same valence, with each image presented for six seconds, for a total duration of 42 seconds per block. Pictures were selected from the original MET and supplemented by pictures from the International Affective Picture System (Lang et al., 2008). The list of pictures used from each valence category is presented in the supplementary material and valence and arousal ratings are as follow: Negative pictures (valence: $M = 3.14$, $SD = 1.47$; arousal: $M = 4.78$, $SD = 2.12$; e.g., homeless man, crying baby, angry man, and distress woman); positive pictures (valence: $M = 6.98$, $SD = 1.61$; arousal: $M = 4.35$, $SD = 2.17$; e.g., laughing boy, grateful girl, happy elderly woman, and pilot); and neutral pictures (valence: $M = 5.27$, $SD = 1.45$; arousal: $M = 3.57$, $SD = 1.96$; e.g., neutral faces of woman, man, and child). Equal numbers of male and female faces

were used across all conditions. Stimuli were presented in color and standardized in size to 507 x 635 pixels, against a gray background.

The task comprised three conditions: a cognitive empathy, an affective empathy, and an age perception control condition. For the cognitive empathy condition, participants were asked to identify "what kind of emotion does this person feel?" by choosing "positive", "negative", or "neutral." For the affective empathy condition, participants were asked to rate "how strong is the emotion you feel about this person?" by choosing "low", "average", and "high". For the age perception control condition, participants were asked to identify "what is the age of this person?" depicted on the picture by choosing "child", "adult", and "elderly". This condition was used to control for higher-order cognitive processes involved in processing of the specific stimuli used in this task. Responses were recorded using three keys on an MRI-compatible response box. To reduce working memory load, all response options remained on the screen.

The task was presented in three runs, each including three blocks (one block for each cognitive empathy, affective empathy, and age perception condition) of positive, negative, and neutral face images, resulting in nine blocks in each run. The order of conditions in each run was pseudo-randomized. To control for the nature of the pictures, stimuli from each valence category were counterbalanced across conditions and runs. That is, positive pictures presented in the affective empathy condition in Run 1 were presented during the cognitive and control conditions in Run 2 and 3, respectively. We ensured that each block of positive, negative, or neutral pictures was only presented once within each run. The order of presenting each run was counterbalanced between participants. To enhance design efficiency (Huettel et al., 2014), each run included two low-level blocks, presented randomly for 42 seconds during the run and consisting of a fixation cross on gray background. In addition, a jittered fixation cross was presented between each block

in each run randomly from one of the following durations: 1.5, 2, and 2.5 seconds. Each run lasted 7.7 minutes, for a total task duration of 23.1 minutes. We used Psychtoolbox for task programming, presentation, and recording of responses.

[Insert Figure 1 about here]

Background measures

In the behavioral/neuropsychological test session, participants completed a series of tasks pertaining to executive functioning: the Stroop Task (Jensen & Rohwer, 1966), the abbreviated version of the Raven's Progressive Matrices (Bilker et al., 2012), the Trail Making Test (Reitan & Wolfson, 1986), and a verbal fluency measure (Newcombe, 1969). Emotional well-being was measured with the Depression, Anxiety, Stress Scale – DASS-21 (Lovibond & Lovibond, 1995). In addition, the Empathy Quotient (Baron-Cohen & Wheelwright, 2004), the Interpersonal Reactivity Index (Davis, 1983), and the Reading the Mind in the Eye test (Baron☐Cohen et al., 2001) measured empathy and theory of mind. Table 1 provides descriptive data and inference statistics of these measures. As shown in Table 1, there were no differences between the two age groups in any of the background measures, with the exception of the three subscales of the DASS-21, for which younger participants reported higher levels than older participants, and the personal distress subscale from the IRI, for which younger participants scored higher than older participants (all $p < 0.001$).[1]

fMRI image acquisition

Functional images were acquired at the Centre for Advanced Imaging on a 3T Siemens scanner with a 32-channel head coil, using a whole-brain T2*-weighted multiband sequence (834 interleaved slices, repetition time (TR) = 612 ms, echo time (TE) = 30 ms, voxel size = 2.5 mm^3,

---

[1] Behavioral performance was analyzed using DASS-21 and personal distress as covariates. Inclusion of these covariates did not change the results.

field of view (FOV) = 190 mm, flip angle = 52º, multi-band acceleration factor = 5). High-resolution T1-weighted images were acquired with an MP2RAGE sequence (176 slices per slab, TR = 4000 ms, TE = 2.91 ms, TI = 700 ms, voxel size = 1 mm^3, FOV = 256 mm, PAT mode = GRAPPA). To minimize noise and head movement, participants were provided with earplugs and cushions around their head inside the head coil. Participants observed the task on a computer screen through a mirror mounted on top of the head coil.

Data Analysis

Behavioral data

We conducted a repeated measure analysis (ANOVA) on mean response times with the experimental conditions (cognitive empathy, affective empathy, age perception) and valence (positive, negative, neutral) as within-subject factors and age group (younger, older) as between-subject factor. While both response times and accuracy were collected during the experiment, given the block design which presented blocks of seven positive, negative, or neutral images respectively, accuracy did not vary within a block and thus was not considered in the analyses.

fMRI

Preprocessing. For functional analysis, T2*-weighted images were preprocessed with Statistical Parametric Mapping Software (SPM12; http://www.fil.ion.ucl.ac.uk/spm) implemented in MATLAB 2015b (Mathworks Inc., MA). Following realignment to a mean image for head-motion correction, images were segmented into gray and white matter. Images were spatially normalized into a standard stereotaxic space with a voxel size of 2 mm^3, using the Montreal Neurological Institute (MNI) template and spatially smoothed with a 6 mm^3 Gaussian Kernel.

Analyses. We used task Partial Least Squares (PLS; McIntosh et al., 1996; McIntosh et al., 2004), as implemented in PLS software running on MATLAB 2012b (The MathWorks Inc., MA), to determine (i) age-related differences in whole-brain activity patterns for the three experimental conditions (cognitive empathy, affective empathy, and age perception); and (ii) age-related differences in modulation of valence (positive, negative, neutral) on cognitive and affective empathy.

PLS is a model-free, multivariate analytical technique (for a detailed tutorial and review of PLS, see Krishnan et al., 2011) that allows examination of the relationship between brain activity and multiple experimental conditions simultaneously. This approach does not require multiple comparison correction (McIntosh et al., 2004). For the whole-brain analysis, we included all three experimental conditions: cognitive empathy, affective empathy, and age perception for both age groups. PLS captures the pattern of covariance between data that are unique to the data set without imposing arbitrary contrasts for experimental conditions or assumptions about conditions. PLS organizes all data from all participants and all experimental conditions into a single matrix and by using a singular value decomposition (SVD) finds a set of orthogonal latent variables (LVs) which represents linear combinations of the original variables.

Each LV delineates brain activity patterns related to the experimental conditions. Usually, the first LV accounts for the largest covariance of the data, with a progressively smaller amount of covariance for subsequent LVs. Each LV consists of a spatiotemporal pattern of brain activity (referred to as voxel saliences), a set of weights that indicates the relationship between experimental conditions with brain activity (referred to as task saliences), and the amount of covariance accounted by the LV (referred to as singular value). Each LV contains brain scores that reflect how each participant contributed to the pattern expressed in the respective LV. A

permutation test with 500 random reordering and resampling was conducted to infer the statistical significance of each LV (McIntosh et al., 1996). Additionally, the robustness of voxel saliences was assessed with a bootstrap estimation with 100 resampling iterations (Efron & Tibshirani, 1985). Peak voxels with a bootstrap ratio (i.e., salience/standard error) ≥ 2.5 were considered reliable, as this approximates $p < 0.001$ (Sampson et al., 1989).

In this study, we used a block design analysis by defining, within each of the three experimental conditions (cognitive empathy, affective empathy, age perception), the onset of each block of positive, negative, and neutral stimuli respectively. For our first aim, to determine age-related differences in brain activity patterns for cognitive and affective empathy, we included all three experimental conditions, irrespective of valence, and both age groups. For the second aim, to examine the modulatory role of valence, we included all three valences and both age groups and conducted two separate analyses: one for cognitive empathy and one for affective empathy.

Results

Behavioral performance

Age-related differences in cognitive and affective empathy and valence modulation. All three main effects were significant. The main effect for experimental condition ($F(2,96) = 89.54$, $p < 0.001$, $\eta_p^2 = 0.65$) showed that participants overall responded slower in the affective than the cognitive condition. The main effect for valence ($F(2,96) = 40.95$, $p < 0.001$, $\eta_p^2 = 0.46$) showed that participants overall responded faster to positive than negative stimuli. The main effect for age group ($F(1,48) = 13.15$, $p = 0.003$, $\eta_p^2 = 0.21$) showed that overall older participants responded more slowly than younger participants.

Qualifying these main effects, the following interactions were significant: the experimental condition by age group interaction ($F(2,96) = 8.36$, $p < 0.001$, $\eta_p^2 = 0.14$), in that older compared to younger participants responded more slowly during both empathy conditions, and particularly during cognitive empathy, but not in the age perception control condition; the valence by age group interaction ($F(2,96) = 4.34$, $p = 0.016$, $\eta_p^2 = 0.08$), in that older compared to younger participants responded more slowly to negative than to positive stimuli; the experimental condition by valence interaction ($F(4,192) = 36.53$, $p < 0.001$, $\eta_p^2 = 0.43$), in that participants responded more slowly to neutral than to emotional stimuli during cognitive empathy but not during the other conditions.

Finally the three-way interaction between experimental condition, age group, and valence was significant ($F(4,192) = 2.48$, $p = 0.04$, $\eta_p^2 = 0.05$). As summarized in Figure 2, both young and older participants responded slower to neutral than to both positive and negative stimuli in the cognitive empathy condition; this difference between the three emotions was not significant for either of the age groups in the affective empathy condition. Also, for younger participants responses to positive, negative and neutral stimuli were similar in the age perception condition whereas older participants responded faster to positive and neutral relative to negative stimuli in that condition.

[Insert Figure 2 about here]

Whole-brain activity pattern

Age-related differences in cognitive and affective empathy. This analysis resulted in two significant LVs. The first LV accounted for 48% covariance of the data ($p < 0.001$). This brain pattern included bilateral insula, bilateral parahippocampus, right superior frontal gyrus, left superior temporal gyrus, right anterior cingulate, left medial frontal gyrus, bilateral precuneus, and bilateral posterior cingulate (Figure 3A); this network was only active among older participants but similarly so during all three experimental conditions (Figure 3B).

The second LV accounted for 26% of the covariance of the data ($p < 0.001$). This mainly left-sided brain pattern included left inferior frontal gyrus, left superior gyrus, left medial frontal gyrus, left inferior parietal lobe, bilateral anterior cingulate, left supramarginal gyrus, left superior temporal gyrus, left middle temporal gyrus, and bilateral insula (Figure 3C). This network was recruited by both young and older participants during affective empathy (Figure 3D). Additionally, this LV included another pattern which corresponded to the age perception control condition in both age groups and included right pre and post central gyrus, bilateral inferior parietal lobe, and right precuneus.

Taken together, the age groups differed in that older but not younger participants engaged a brain pattern including anterior, posterior cingulate and temporal areas comparably during cognitive empathy, affective empathy, and age perception. However, both age groups similarly engaged a left-lateralized fronto-parietal network during affective empathy.

[Insert Figure 3 about here]

Age-related differences in valence modulation on cognitive or affective empathy. The analysis pertaining to cognitive empathy resulted in one significant LV that accounted for 39% of covariance of the data ($p = 0.002$). This network included bilateral inferior frontal gyrus, anterior cingulate, bilateral superior temporal gyrus, bilateral superior frontal gyrus, medial frontal gyrus, and bilateral precentral gyrus regions (Figure 4A). This network was recruited only by older participants during cognitive empathy to negative and neutral stimuli (Figure 4B).

The analysis pertaining to affective empathy also resulted in one significant LV which accounted for 51% of covariance of the data ($p < 0.001$). This network included bilateral middle and superior frontal gyrus, posterior cingulate, precuneus, lingual gyrus, inferior parietal lobe, and superior temporal gyrus (Figure 4C). This wide-spread network was activated during affective empathy to positive stimuli in both young and older participants, and also to neutral stimuli in older participants (Figure 4D).

[Insert Figure 4 about here]

Taken together, during cognitive empathy older but not younger participants engaged a pattern of brain regions resembling the salience network in response to negative and neutral stimuli. During affective empathy, in contrast, the age groups were similar in recruiting major nodes of the default mode network in response to positive stimuli; older participants also recruited this network in response to neutral simuli.

Discussion

Cognitive empathy has been consistently shown to decrease with age whereas studies measuring age-related differences in affective empathy have generated mixed findings. Only two previous studies have investigated the neural correlates of empathic response in older adults and either used only pain stimuli (Chen et al., 2014) or measured self-reported empathic response

outside the scanner (Moore et al., 2015). Our study went beyond these past approaches and generated several novel insights regarding age-related differences in cognitive and affective empathy both in behavior and in the brain, advancing understanding of the neural networks involved in empathic response in adulthood. Furthermore, our study allowed a systematic test of valence modulation of cognitive and affective empathy in young and older adults.

We found that older adults responded more slowly to empathic conditions and their response time was modulated by the emotional valence during cognitive, but not affective, empathy. In addition, supporting age-related differences in brain response, older but not young participants recruited the same network across cognitive and affective empathy (as well as during the control condition). The age groups, however, did not differ in their recruitment of a left-lateralized fronto-parietal network during affective empathy. We also found that older but not younger participants recruited regions of the salience network with cognitive empathy in response to negative emotions. In contrast, with affective empathy in response to positive emotions, both age groups recruited a similar pattern of brain regions that contained nodes of the default mode network. We next discuss the central findings.

Age-comparability of behavior and brain patterns during affective empathy

Neither our brain nor our behavioral data supported age-related differences in affective empathy. Thus, in contrast to Chen and colleagues (2014), we did not observe reduced engagement of core empathy regions including anterior insula and mid-cingulate in older participants. Rather, both young and older adults recruited these regions similarly during the performance of the affective empathy task. It is possible that differences in the type of stimuli used in our study (i.e., non-pain stimuli) compared to the pain stimuli used in Chen et al. underlie these divergent findings. Speaking against this methodological explanation, a recent meta-

analysis found similarities in empathic responses to pain and non-pain stimuli (Timmers et al., 2018); this comparability across stimulus types, however, has not been confirmed in research with older individuals yet.

Our brain findings for affective empathy do not support the notion of diminished internal bodily perception (interoceptive awareness) among older adults (Mendes, 2010), but functional sparing for affective processing in aging could offer an explanation, in line with the brain maintenance hypothesis discussed in the cognitive aging literature (Nyberg et al., 2012). This idea of functional reserve that protects older adults from decline may be particularly apposite to our data given that participants in the current study were cognitively high functioning (see background measures in Table 1). Thus, they may have had high cognitive reserve, which could have resulted in improved performance, on a level comparable with younger participants, especially with the affective empathy task which may have been easier to perform than the cognitive empathy task. Future research will be able to test this interpretation.

Valence modulation on cognitive and affective empathy

Age-related differences were observed in valence modulation during cognitive empathy. We found that older but not young participants engaged a neural network that comprised bilateral insula, inferior frontal gyrus, and anterior cingulate, core nodes of the salience network (Menon, 2015; Menon & Uddin, 2010), with cognitive empathy to negative emotions. This finding aligns with behavioral evidence that older compared to younger adults experience greater difficulty with processing negative emotions than with positive emotions (Hayes et al., 2020; Ruffman et al., 2008). Our findings also corroborate evidence of increased prefrontal cortex and insula activity for negative emotions in older adults, possibly reflecting more processing effort for

and/or greater salience when processing negative compared to positive stimuli with advanced age (Ebner et al., 2012; Ziaei, Burianova, et al., 2016).

One possibility for engagement of the salience network during cognitive empathy to negative emotions in older adults is that this network, and insula specifically, is involved in orienting attention towards relevant stimuli in the environment (Menon et al., 2010). Given the salience feature of negative emotions and their importance for survival, orienting attention towards negative emotions is crucial and is thus associated with the insular response. Insular activity in this context may be reflective of a response that is commonly expressed among various cognitive tasks to guide behavior in dynamic social contexts (Bernhardt et al., 2012), crucial for recognizing negative emotions. Our result, however, contradicts suggestions of age-related reductions in insular activation subserving interoception and the simulation of emotions in others (Mather, 2016). Further investigation is needed to determine the relationship between bodily response (such as heart rate variability and skin conductance) and insular activity during empathy, and social cognition more broadly, across adulthood.

Age-similarity in valence modulation during affective empathy

Our finding of no age-related difference in affective responding to positive stimuli is also largely in line with the socioemotional selectivity theory (SST) (Carstensen et al., 2003; Carstensen et al., 1999) which proposes that older participants preferentially process positive over negative stimuli. In other words, older adults' bias towards positive emotions may have facilitated processing of positive emotions in the affective empathy task, resulting in comparable brain activity patterns between the age groups in this condition. Thus, the present study's results suggest that the positivity effect reported in the aging literature for various cognitive and social-cognitive processes also extends to affective empathic responding to positive emotions.

However, we only found support for the positivity effect in affective but not in cognitive empathy, which warrants further investigation.

We also found involvement of the posterior cingulate cortex during affective empathy for positive emotions in both age groups. This involvement of posterior cingulate during empathizing with people who express positive emotions may reflect the relatively lower effort needed with this task. However, it is also possible that the observed posterior cingulate cortex activity with positive affective empathy reflects self-referential (affective) processing in linking one's own and another's emotional state to enable adequate empathic responding. In support of this interpretation, posterior cingulate cortex has been shown to play a role in a wide range of social-cognitive processes (Brewer et al., 2013; Sperduti et al., 2012). For example, posterior cingulate activation is involved in theory of mind and mentalizing (Frith & Frith, 2006; Mitchell, 2009; Molenberghs et al., 2016). A meta-analysis furthermore showed that posterior cingulate cortex subserves empathy (Bzdok et al., 2012a), and specifically the evaluation of how "one relates to one's experience" (Brewer et al., 2013). Our findings importantly add to previous literature by demonstrating a role for the posterior cingulate in affective empathy in older adults.

Our finding that valence modulates empathy is largely in line with Morelli and colleagues' (2015) suggestion that empathy is not a uni-dimensional construct and that empathic response to positive emotions is a different construct to general positivity or general empathy for others. Differentiating these diverse constructs and determining the respective brain processes involved across adulthood will be an interesting future direction to pursue. Also, in the present study we did not measure prosocial behavior and therefore cannot determine the link between empathic responding to positive and negative emotions and prosociality. This research is particularly promising to explore among older adults moving forward given growing evidence of a

relationship between empathic response to positive emotions and increased prosocial behavior, social closeness, and well-being among younger adults (Morelli et al., 2015).

Conclusion

This is the first study to distinguish behavioral and brain responses for empathic response towards positive vs. negative emotions among young and older adults. Older (but not young) adults engaged the salience network during cognitive empathy in response to negative emotions, which could reflect older adults' difficulty in the processing of and/or their enhanced interoception for negative emotions during cognitive empathy. Furthermore, both age groups recruited a bilateral network including nodes of the default mode network, possibly reflecting self-referential processing and/or decreased cognitive effort during affectively empathizing with positive emotions. These findings show that valence plays a critical role in empathic response both in young and older adults and therefore needs to be considered in investigations into higher-order social cognitive functions not only in the field of gerontology but also in other populations with deficits in social-cognitive function (e.g., individuals with autism spectrum disorder or neurodegenerative disorders; Henry et al., 2016). Based on our results future research should test the extent to which emotions displayed by another affect social interactions such as closeness or altruistic behavior as well as general well-being via empathic responses in both young and older adults.

References

Bailey, P. E., Brady, B., Ebner, N. C., & Ruffman, T. (2018). Effects of Age on Emotion Regulation, Emotional Empathy, and Prosocial Behavior. *J Gerontol B Psychol Sci Soc Sci*. doi:10.1093/geronb/gby084

Baron-Cohen, S., & Wheelwright, S. (2004). The empathy quotient: an investigation of adults with Asperger syndrome or high functioning autism, and normal sex differences. *J. of autism and dev. dis., 34*(2), 163-175. doi:10.1023/B:JADD.0000022607.19833.00

Baron☐Cohen, S., Wheelwright, S., Hill, J., Raste, Y., & Plumb, I. (2001). The "Reading the Mind in the Eyes" test revised version: A study with normal adults, and adults with Asperger syndrome or high☐functioning autism. *Journal of child psychology and psychiatry, 42*(2), 241-251. doi:10.1111/1469-7610.00715

Beadle, J. N., & de la Vega, C. E. (2019). Impact of Aging on Empathy: Review of Psychological and Neural Mechanisms. *Front Psychiatry, 10*, 331. doi:10.3389/fpsyt.2019.00331

Beadle, J. N., Sheehan, A. H., Dahlben, B., & Gutchess, A. H. (2015). Aging, empathy, and prosociality. *J Gerontol B Psychol Sci Soc Sci, 70*(2), 215-224. doi:10.1093/geronb/gbt091

Bernhardt, B. C., & Singer, T. (2012). The neural basis of empathy. *Annual review of neuroscience, 35*, 1-23.

Bilker, W. B., Hansen, J. A., Brensinger, C. M., Richard, J., Gur, R. E., & Gur, R. C. (2012). Development of abbreviated nine-item forms of the Raven's standard progressive matrices test. *Assessment, 19*(3), 354-369. doi:10.1177/1073191112446655

Brewer, J. A., Garrison, K. A., & Whitfield-Gabrieli, S. (2013). What about the "Self" is Processed in the Posterior Cingulate Cortex? *Front Hum Neurosci, 7*, 647. doi:10.3389/fnhum.2013.00647

Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: anatomy, function, and relevance to disease. *Ann. N. Y. Acad. Sci., 1124*, 1-38. doi:10.1196/annals.1440.011

Bzdok, D., Schilbach, L., Vogeley, K., Schneider, K., Laird, A., Langner, R., & Eickhoff, S. (2012a). Parsing the neural correlates of moral cognition: ALE meta-analysis on morality, theory of mind, and empathy. *Brain Structure and Function, 217*(4), 783-796. doi:10.1007/s00429-012-0380-y

Bzdok, D., Schilbach, L., Vogeley, K., Schneider, K., Laird, A. R., Langner, R., & Eickhoff, S. B. (2012b). Parsing the neural correlates of moral cognition: ALE meta-analysis on morality, theory of mind, and empathy. *Brain Structure and Function, 217*(4), 783-796. doi:10.1007/s00429-012-0380-y

Carstensen, L. L., Fung, H. H., & Charles, S. T. (2003). Socioemotional Selectivity Theory and the regulation of emotion in the second half of the life *Motivation and Emotion, 27*, 103-123.

Carstensen, L. L., Isaacowitz, D. M., & Charles, S. T. (1999). Taking time seriously: A theory of socioemotional selectivity. *American Psychologist, 54*(3), 165.

Charles, S. T., Mather, M., & Carstensen, L. L. (2003). Aging and emotional memory: The forgettable nature of negative images for older adults. *Journal of Experimental Psychology: General, 132*(2), 310-324. doi:10.1037/0096-3445.132.2.310

Chen, Y. C., Chen, C. C., Decety, J., & Cheng, Y. (2014). Aging is associated with changes in the neural circuits underlying empathy. *Neurobiol Aging, 35*(4), 827-836. doi:10.1016/j.neurobiolaging.2013.10.080

Davis, M. H. (1983). Measuring individual differences in empathy: Evidence for a multidimensional approach. *Journal of personality and social psychology, 44*(1), 113. doi:10.1037/0022-3514.44.1.113

Decety, J., & Jackson, P. L. (2006). A Social-Neuroscience Perspective on Empathy. *Current Directions in Psychological Science, 15*(2), 54-58.

Dziobek, I., Preissler, S., Grozdanovic, Z., Heuser, I., Heekeren, H. R., & Roepke, S. (2011). Neuronal correlates of altered empathy and social cognition in borderline personality disorder. *Neuroimage, 57*(2), 539-548. doi:10.1016/j.neuroimage.2011.05.005

Ebner, N. C., Johnson, M. K., & Fischer, H. (2012). Neural mechanisms of reading facial emotions in young and older adults. *Front Psychol, 3*, 223-242. doi:10.3389/fpsyg.2012.00223

Efron, B., & Tibshirani, R. (1985). The bootstrap method for assessing statistical accuracy. *Behaviormetrika, 12*(17), 1-35.

Fan, Y., Duncan, N. W., de Greck, M., & Northoff, G. (2011). Is there a core neural network in empathy? An fMRI based quantitative meta-analysis. *Neuroscience & Biobehavioral Reviews, 35*(3), 903-911.

Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-Mental State": a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research, 12*(3), 189-198. doi:10.1016/0022-3956(75)90026-6

Frith, C. D., & Frith, U. (2006). The neural basis of mentalizing. *Neuron, 50*(4), 531-534.

Grühn, D., Rebucal, K., Diehl, M., Lumley, M., & Labouvie-Vief, G. (2008). Empathy Across the Adult Lifespan: Longitudinal and Experience-Sampling Findings. *Emotion (Washington, D.C.), 8*(6), 753-765. doi:10.1037/a0014123

Hayes, G. S., McLennan, S. N., Henry, J. D., Phillips, L. H., Terrett, G., Rendell, P. G., . . . Labuschagne, I. (2020). Task characteristics influence facial emotion recognition age-effects: A meta-analytic review. *Psychol Aging, 35*(2), 295-315. doi:10.1037/pag0000441

Henry, J. D., Phillips, L. H., Ruffman, T., & Bailey, P. E. (2013). A meta-analytic review of age differences in theory of mind. *Psychol Aging, 28*(3), 826. doi:10.1037/a0030677

Henry, J. D., von Hippel, W., Molenberghs, P., Lee, T., & Sachdev, P. S. (2016). Clinical assessment of social cognitive function in neurological disorders. *Nat Rev Neurol, 12*(1), 28-39. doi:10.1038/nrneurol.2015.229

Huettel, S. A., Song, A. W., & McCarthy, G. (2014). *Functional Magnetic Resonance Imaging* (Vol. 3rd volume).

Jensen, A. R., & Rohwer, W. D. (1966). The Stroop color-word test: A review. *Acta psychologica, 25*, 36-93.

Krishnan, A., Williams, L. J., McIntosh, A. R., & Abdi, H. (2011). Partial Least Squares (PLS) methods for neuroimaging: A tutorial and review. *Neuroimage, 56*(2), 455-475. doi:10.1016/j.neuroimage.2010.07.034

Lamm, C., Decety, J., & Singer, T. (2011). Meta-analytic evidence for common and distinct neural networks associated with directly experienced pain and empathy for pain. *Neuroimage, 54*(3), 2492-2502. doi:10.1016/j.neuroimage.2010.10.014

Lang, P. J., Bradley, M. M., & Cuthbert, B. N. (2008). International affective picture system (IAPS): Affective ratings of pictures and instruction manual. *Technical report A-8.*

Leigland, L. A., Schulz, L. E., & Janowsky, J. S. (2004). Age related changes in emotional memory. *Neurobiol Aging, 25*(8), 1117-1124. doi:https://doi.org/10.1016/j.neurobiolaging.2003.10.015

Lin, T., Ankudowich, E., & Ebner, N. C. (2017). Greater perceived similarity between self and own-age others in older than young adults. *Psychol Aging, 32*(4), 377-387. doi:10.1037/pag0000173

Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behaviour research and therapy, 33*, 335-343. doi:10.1016/0005-7967(94)00075-U

Mather, M. (2016). The Affective Neuroscience of Aging. *Annual review of psychology, 67*, 213-238. doi:10.1146/annurev-psych-122414-033540

Mather, M., & Carstensen, L. L. (2003). Aging and Attentional Biases for Emotional Faces. *Psychological Science, 14*(5), 409-415. doi:10.1111/1467-9280.01455

McIntosh, A. R., Bookstein, F. L., Haxby, J. V., & Grady, C. L. (1996). Spatial Pattern Analysis of Functional Brain Images Using Partial Least Squares. *Neuroimage, 3*(3), 143-157. doi:10.1006/nimg.1996.0016

McIntosh, A. R., Chau, W. K., & Protzner, A. B. (2004). Spatiotemporal analysis of event-related fMRI data using partial least squares. *Neuroimage, 23*(2), 764-775. doi:10.1016/j.neuroimage.2004.05.018

Mendes, W. B. (2010). Weakened Links Between Mind and Body in Older Age: The Case for Maturational Dualism in the Experience of Emotion. *Emotion Review, 2*(3), 240-244. doi:10.1177/1754073910364149

Menon, V. (2015). Salience Network. In A. W. Toga (Ed.), *Brain Mapping: An Encyclopedic Reference* (Vol. 2, pp. 597-611). Academic Press: Elsevier.

Menon, V., & Uddin, L. Q. (2010). Saliency, switching, attention and control: a network model of insula function. *Brain Structure and Function, 214*(5-6), 655-667. doi:10.1007/s00429-010-0262-0

Mitchell, J. P. (2009). Inferences about mental states. *Philosophical Transactions of the Royal Society B: Biological Sciences, 364*(1521), 1309-1316.

Molenberghs, P., Johnson, H., Henry, J. D., & Mattingley, J. B. (2016). Understanding the minds of others: A neuroimaging meta-analysis. *Neurosci Biobehav Rev, 65*, 276-291. doi:10.1016/j.neubiorev.2016.03.020

Moore, R. C., Dev, S. I., Jeste, D. V., Dziobek, I., & Eyler, L. T. (2015). Distinct neural correlates of emotional and cognitive empathy in older adults. *Psychiatry Res, 232*(1), 42-50. doi:10.1016/j.pscychresns.2014.10.016

Morelli, S. A., Lieberman, M. D., & Zaki, J. (2015). The Emerging Study of Positive Empathy. *Social and Personality Psychology Compass, 9*(2), 57-68. doi:10.1111/spc3.12157

Newcombe, F. (1969). Missile wounds of the brain: A study of psychological deficits.

Nyberg, L., Lövdén, M., Riklund, K., Lindenberger, U., & Bäckman, L. (2012). Memory aging and brain maintenance. *Trends in Cognitive Science, 16*(5), 292-305. doi:https://doi.org/10.1016/j.tics.2012.04.005

O'Brien, E., Konrath, S. H., Grühn, D., & Hagen, A. L. (2012). Empathic Concern and Perspective Taking: Linear and Quadratic Effects of Age Across the Adult Life Span. *The Journals of Gerontology: Series B, 68*(2), 168-175. doi:10.1093/geronb/gbs055 %J The Journals of Gerontology: Series B

Reed, A. E., & Carstensen, L. L. (2012). The theory behind the age-related positivity effect. *Front Psychol, 3*, 339. doi:10.3389/fpsyg.2012.00339

Reed, A. E., Chan, L., & Mikels, J. A. (2014). Meta-analysis of the age-related positivity effect: Age differences in preferences for positive over negative information. *Psychol Aging, 29*, 1-15. doi:10.1037/a0035194

Reitan, R. M., & Wolfson, D. (1986). The Halstead-Reitan Neuropsychological Test battery and aging. *Clinical Gerontologist: The Journal of Aging and Mental Health, 5*, 39 – 61. doi:doi:10.1300/J018v05n01_03

Ruffman, T., Henry, J. D., Livingstone, V., & Phillips, L. H. (2008). A meta-analytic review of emotion recognition and aging: implications for neuropsychological models of aging. *Neurosci Biobehav Rev, 32*(4), 863-881. doi:10.1016/j.neubiorev.2008.01.001

Sampson, P. D., Streissguth, A. P., Barr, H. M., & Bookstein, F. L. (1989). Neurobehavioral effects of prenatal alcohol: Part II. Partial Least Squares analysis. *Neurotoxicology and Teratology, 11*(5), 477-491. doi:10.1016/0892-0362(89)90025-1

Singer, T., & Lamm, C. (2009). The social neuroscience of empathy. *Ann N Y Acad Sci, 1156*, 81-96. doi:10.1111/j.1749-6632.2009.04418.x

Sperduti, M., Martinelli, P., & Piolino, P. (2012). A neurocognitive model of meditation based on activation likelihood estimation (ALE) meta-analysis. *Conscious Cogn, 21*(1), 269-276. doi:https://doi.org/10.1016/j.concog.2011.09.019

Sun, R., Vuillier, L., Hui, B. P. H., & Kogan, A. (2019). Caring helps: Trait empathy is related to better coping strategies and differs in the poor versus the rich. *PLoS One, 14*(3), e0213142. doi:10.1371/journal.pone.0213142

Sze, J. A., Gyurak, A., Goodkind, M. S., & Levenson, R. W. (2012). Greater emotional empathy and prosocial behavior in late life. *Emotion, 12*(5), 1129-1140. doi:10.1037/a0025011

Timmers, I., Park, A. L., Fischer, M. D., Kronman, C. A., Heathcote, L. C., Hernandez, J. M., & Simons, L. E. (2018). Is Empathy for Pain Unique in Its Neural Correlates? A Meta-Analysis of Neuroimaging Studies of Empathy. *Frontiers in Behavioral Neuroscience, 12*(289). doi:10.3389/fnbeh.2018.00289

von Hippel, W., & Dunlop, S. M. (2005). Aging, inhibition, and social inappropriateness. *Psychol Aging, 20*, 519. doi:10.1037/0882-7974.20.3.519

Ziaei, M., Burianova, H., Von Hippel, W., Ebner, N. C., Phillips, L. H., & Henry, J. D. (2016). The impact of aging on the neural networks involved in gaze and emotional processing. *Neurobiol Aging, 48*, 182-194. doi:10.1016/j.neurobiolaging.2016.08.026

Ziaei, M., & Fischer, H. (2016). Emotion and Aging:The Impact of Emotion on Attention, Memory, and Face Recognition in Late Adulthood. In J. R. Absher & J. Cloutier (Eds.), *Neuroimaging Personality, Social Cognition, and Character* (pp. 259-278). San Diego: Academic Press: Elsevier.

Ziaei, M., von Hippel, W., Henry, J. D., & Becker, S. I. (2015). Are Age Effects in Positivity Influenced by the Valence of Distractors? *PLoS One, 10*(9), e0137604. doi:10.1371/journal.pone.0137604

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Captions

Figure 1. Experimental task. The task included three experimental conditions: cognitive empathy, affective empathy, and age perception. Each condition consisted of three blocks of either positive, negative, or neutral face images. Each condition started with a 3-second instruction, followed by presentation of 7 images, each for 6 seconds, for a total of 42 seconds per block. Following each block, a jittered fixation cross (duration: 1.5, 2, or 2.5 seconds) was presented.

Figure 2. Mean response times (in seconds) for the three experimental conditions (cognitive empathy, affective empathy, age perception) in young and older participants for positive, negative, and neutral stimuli. Error bars = +/- 1 standard error. * denotes $p < .000$.

Figure 3. Results from the whole-brain analysis for the three experimental conditions (cognitive empathy, affective empathy, age perception) in young and older participants. Panel A represents the brain activity pattern across all three experimental conditions, differentiating between young and older participants. Panel B represents brain scores associated with the brain activity pattern presented in Panel A. Panel C represents the brain activity pattern differentiating affective empathy from the other two experimental conditions, similarly for young and older participants. Panel D represents brain scores associated with the brain activity pattern presented in Panel C. Error bars represent confidence intervals at 95%. For all reported regions a bootstrap ratio of $\geq$ 2.5 and cluster size of $\geq$ 50 voxels was applied. L = left hemisphere, R = right hemisphere.

Figure 4. Results from the whole-brain analysis regarding valence modulation (positive, negtive, neutral) for cognitive and affective empathy in young and older participants. Panel A refers to cognitive empathy and represents the brain activity pattern differentiating negative and neutral from positive simuli in older participants. Panel B represents brain scores associated with the brain activity pattern presented in Panel A. Panel C refers to affective empathy and represents the brain activity pattern differentiating positive from negative and neutral stimuli in young participants and positive and neutral from negative stimuli in older participants. Panel D represents brain scores associated with the brain activity pattern presented in Panel C. Error bars represent confidence intervals at 95%. For all reported regions a bootstrap ratio of ≥ 2.5 and cluster size of ≥ 50 voxels was applied. L = left hemisphere, R = right hemisphere.

Tables

Table 1. Descriptive data (means and standard deviations) and inferential statistics for background measures in young and older participants

	Younger Participants		Older Participants		Age-Group Differences	
	M	*SD*	*M*	*SD*	*t*	*df*
Demographics						
Age (years)	21.72	3.81	71.52	3.80	46.19[**]	48
Education (years)	15.50	2.46	17.34	8.79	1.01	48
Executive Functioning						
Verbal Fluency Test	47.84	10.67	53.20	12.42	1.63	48
Stroop Interference Score	0.23	0.25	0.29	0.18	0.82	48
Task Switching Index	20.63	9.02	24.86	11.99	1.35	45
Raven IQ	6.40	1.77	5.68	1.97	1.35	48
Emotional Well-Being						
DASS-21						
Stress	6.40	6.05	1.33	2.33	3.83**	47

	M	SD	M	SD	t	df
Anxiety	5.12	3.60	1.00	1.56	5.14**	47
Depression	10.40	5.94	3.66	3.85	4.68**	47
Empathy						
IRI						
Fantasy	16.12	4.70	13.64	5.42	1.72	48
Empathic concern	17.88	3.63	18.68	3.55	0.78	48
Perspective taking	19.44	4.11	19.64	3.46	0.18	48
Personal distress	12.04	5.67	8.64	4.31	2.38*	48
Empathy Quotient	44.44	11.16	49.08	11.45	1.45	48
RMET	25.76	4.20	27.72	4.37	1.61	48

Note. * $p < 0.05$, ** $p < .001$. Verbal Fluency Test (Benton & Hamsher, 1976): total number of words for letters F, A, and S; Stroop Interference Score (Ziaei et al., 2015) = ((response time in incongruent trials – response time in neutral trials)/response time in neutral trials); Task-Switching Index (Reitan & Wolfson, 1986): Trail Making Test Part B – Trail Making Test Part A; Raven IQ (Kirchner, 1958) = total number of correct responses; DASS-21 = Depression Anxiety Stress Scale (Lovibond & Lovibond, 1995); IRI = Interpersonal Reactivity Index (Davis, 1983); RMET = Reading the Mind in the Eye Test (Baron-Cohen et al., 2001); *df* = degree of freedom; *SD* = standard deviation; *M* = mean; *t* = student t-test.

Table 2. Whole-brain results showing regions modulated by valence across younger and older participants

Regions	Hem	MNI coordinates			BSR
		X	Y	Z	

Age-related differences in valence modulation during cognitive empathy

Older participants > younger participants (negative stimuli)

Regions	Hem	MNI coordinates	BSR
Middle frontal gyrus	L	-37 52 11	-4.71
Inferior frontal gyrus	L	-49 14 11	-5.42
	R	52 18 11	-3.75
Insula	L	-44 16 7	-4.19
Insula	R	37 20 7	-3.01
Medial frontal gyrus	R	0 51 28	-3.53
Superior frontal gyrus	L	-3 46 42	-4.54
Inferior parietal lobe	L	-46 -28 50	-4.15
Superior temporal gyrus	R	56 -46 11	-3.73
Superior temporal gyrus	L	-58 -48 11	-3.40
Caudate	L	-11 13 20	-4.16

Age-related similarity in valence modulation during affective empathy

Older participants = younger participants (positive stimuli)

	Hem	MNI	BSR
Anterior cingulate gyrus	L	-4 38 16	4.43
Middle frontal gyrus	L	-24 42 32	4.40
Inferior frontal gyrus	L	-50 6 16	4.79
Superior frontal gyrus	R	30 52 32	5.82
Insula	L	-36 -31 20	5.00
	R	63 -30 20	4.39
Inferior parietal lobe	R	62 -24 32	4.42
Posterior cingulate	R	2 -63 20	4.45
Superior temporal gyrus	L	-38 -36 16	4.81
	R	52 -42 20	3.62
Parahippocampus	R	28 -44 -2	3.74

Note. Hem = hemisphere; L = left, R = right; BSR = bootstrap ratio; MNI = Montreal Neurological Institute. Brain regions with BSR

> 3.5 are reported here.