Genome-wide circadian rhythm detection methods: systematic evaluations and practical guidelines

Wenwen Mei[1,*], Zhiwen Jiang[1,*], Yang Chen[2], Li Chen[3], Aziz Sancar[4,5,#], Yuchao Jiang[1,5,6,#]

1. Department of Biostatistics, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC 27599, USA.
2. Department of Statistics and Michigan Institute for Data Science, University of Michigan, Ann Arbor, MI 48109, USA.
3. Department of Medicine and Center for Computational Biology and Bioinformatics, Indiana University School of Medicine, Indianapolis, IN 46202, USA.
4. Department of Biochemistry and Biophysics, School of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA.
5. Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC 27599, USA.
6. Department of Genetics, School of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA.

Wenwen Mei is a PhD student in the Department of Biostatistics at the University of North Carolina at Chapel Hill.

Zhiwen Jiang is a MS student in the Department of Biostatistics at the University of North Carolina at Chapel Hill.

Yang Chen is an Assistant Professor in the Department of Statistics and Research Assistant Professor at the Michigan Institute of Data Science at the University of Michigan.

Li Chen is an Assistant Professor in the Dpartment of Medicine and a member of the Center for Computational Biology and Bioinformatics at Indianna University School of Medicine.

Aziz Sancar is the Sarah Graham Kenan Professor of Biochemistry and Biophysics at the University of North Carolina School of Medicine and member of UNC Lineberger Comprehensive Cancer Center.

Yuchao Jiang is an Assistant Professor in the Department of Biostatisrtics and the Department of Genetics at Univerity of North Carolina at Chapel Hill and member of UNC Lineberger Comprehensive Cancer Center.

∗   These authors contributed equally.

To whom correspondence should be addressed. Email: aziz_sancar@med.unc.edu; yuchaoj@email.unc.edu.

ABSTRACT

Circadian rhythms are oscillations of behavior, physiology, and metabolism in many organisms. Recent advancements in omics technology make it possible for genome-wide profiling of circadian rhythms. Here, we conducted a comprehensive analysis of seven existing algorithms commonly used for circadian rhythm detection. Using gold-standard circadian and non-circadian genes, we systematically evaluated the accuracy and reproducibility of the algorithms on empirical datasets generated from various omics platforms under different experimental designs. We also carried out extensive simulation studies to test each algorithm's robustness to key variables, including sampling patterns, replicates, waveforms, signal-to-noise ratios, uneven samplings, and missing values. Furthermore, we examined the distributions of the nominal p-values under the null and raised issues with multiple testing corrections using traditional approaches. With our assessment, we provide method selection guidelines for circadian rhythm detection, which are applicable to different types of high-throughput omics data.

Key points

- Various methods have been developed for circadian rhythm detection on a genome-wide scale using omics technologies, yet there has not been a comprehensive summary and evaluation of all existing methods to date.

- Using gold-standard circadian and non-circadian genes, we systematically evaluated the accuracy and reproducibility of seven existing algorithms for circadian rhythm detection on empirical datasets generated from various omics platforms.

- We carried out extensive simulation studies to test each algorithm's robustness to key variables, including sampling patterns, replicates, waveforms, signal-to-noise ratios, uneven samplings, and missing values.

- We examined the distributions of the nominal p-values under the null and raised issues with multiple testing corrections using the Benjamini-Hochberg procedure due to gene-gene correlation and testing being overly conservative.

- We provide method selection guidelines for circadian rhythm detection, which are applicable to different types of high-throughput omics data.

BACKGROUND

Circadian rhythms are approximately 24-hour oscillations of behavior, physiology, and metabolism that exist in almost all living organisms ranging from prokaryotes to mammals [1, 2]. Circadian rhythm is regulated by the circadian system, which consists of many "clock-controlled genes" that exhibit oscillatory patterns [1]. These oscillations provide organisms with an adaptive advantage by enabling them to predict and adjust to the variations within their environments [3]. Additionally, and perhaps more importantly, disruptions of circadian rhythms have shown to contribute to numerous diseases, including metabolic disorders, heart disease, and aging [4-7]. It is, therefore, of great importance and interest to perform genome-scale analysis of biological rhythms.

Recent advances in omics technologies, including both microarrays and next-generation sequencing, offer appealing platforms to identify circadian genes on a genome-wide scale. These have, indeed, led to the proposal of multifarious methodologies adopted from various fields including mathematics, statistics, astrophysics, etc. The earliest of the selected methods is Lomb-Scargle (LS) periodogram [8], an algorithm adapted from astrophysics that detects oscillations by comparing the data to sinusoidal reference curves of varying periods and phases [9, 10]. ARSER is an algorithm that employs autoregressive spectral estimation to predict periodicity and applies a harmonic regression model to fit the time-series [11]. Unlike the model-based LS and ARSER, JTK_CYCLE is a non-parametric method that detects oscillations by comparing the ranks of the measured values to a set of prespecified symmetric reference curves [3]. Both RAIN and eJTK_CYCLE build on the strengths of JTK_CYCLE: RAIN includes an additional set of asymmetric waveforms and examines the increasing and decreasing portions of the curve separately [12]; eJTK_CYCLE improves JTK_CYCLE by explicitly calculating the null distribution such that it accounts for multiple hypothesis testing and by including non-sinusoidal reference waveforms [13]. Based on the successes of the aforementioned methods, MetaCycle proposes an ensemble framework that integrates results from three different algorithms, LS, ARSER, and JTK_CYCLE [14]. Specifically, MetaCycle detects periodicity using the best of breed methods: its p-values are generated using Fisher's method; its periods and phase estimations are integrated using arithmetic and circular means; and a new periodic model, formulated from ordinary least squares

method, is applied to recalculate the amplitude. The most recent method, BIO_CYCLE, is a deep neural network trained on both simulated and empirical circadian and noncircadian time-series [15]. More general information and characteristics of each method are summarized in Table 1.

Multiple studies [10, 16, 17] have evaluated the performance of different methods for circadian rhythm detection, showing discrepancies among the methods, whose performances depend on multiple factors including experimental designs, waveforms of interest, etc. However, there has not been, to our best knowledge, a comprehensive summary and evaluation of all existing methods to date. Here, we systematically assess the performance of the seven aforementioned algorithms for circadian rhythm detection: LS, ARSER, JTK_CYCLE, RAIN, eJTK_CYCLE, MetaCycle, and BIO_CYCLE.

Specifically, we demonstrated and benchmarked the algorithms using real datasets with gold-standard circadian and non-circadian genes. All empirical data were generated using the liver tissue from *Mus musculus* that had undergone two different experimental designs. Under the dark-dark experimental design (24-hour darkness), we focused on using data from gene expression microarrays to assess the accuracy and reproducibility of each algorithm; under the light-dark experimental design (12-hour light followed by 12-hour darkness), we adopted four different next-generation sequencing platforms and explored the robustness of each method in identifying circadian genes. Furthermore, to extend our assessment to non-transcriptomic datasets, we included a proteomic dataset in our evaluation. In addition, we carried out extensive simulation studies to study how key variables, including sampling patterns, replicates, waveforms, signal-to-noise ratios, uneven samplings, missing values, affect the performance of each method. Lastly, we point out the flaw with using the Benjamini-Hochberg procedure to control for false discovery rate. Through these, we offer guidelines on experimental designs as well as best practices and methods of choice to increase the rigor and reproducibility in the analysis of large-scale circadian rhythms. To assist with the comparison of future methods and datasets using our framework, we provide detailed vignettes on applications of existing methods and performance evaluations with source code available at https://github.com/wenwenm183/Circadian_Genes_Benchmark.

RESULTS

Performance assessment using empirical datasets with dark-dark design

We first adopted three gene expression microarray datasets from Hughes et al. [18], Hughes et al. [19], and Zhang et al. [20]. For all three studies, mouse liver samples were collected in every hour or every two-hour under the dark-dark experimental design for 48 hours. We named these three datasets after the first author's last name and the year of publication as Hughes 2009, Hughes 2012, and Zhang 2014, respectively. In addition, we generated a new downsampled dataset from the Hughes 2009 dataset by keeping the even time-points only, and named it "Downsampled Hughes 2009". Refer to Table 2A for details of the data. Figure 1 shows the scaled gene expression levels of four known circadian and four non-circadian genes. The circadian genes, including the well-studied *Clock*, *Cry1*, *Npas2*, and *Per1* [10], show oscillatory patterns that can be well reproduced across studies, while the non-circadian genes exhibit only noisy signals.

We set out to apply the seven algorithms to these four datasets to detect significantly cyclic genes and evaluate their performances using 104 circadian [10] and 113 non-circadian genes [21] from previous studies (Supplementary Table 1). The accuracy of each method in Hughes 2009, Downsampled Hughes 2009, Hughes 2012, and Zhang 2014 was first assayed with the precision and recall rates for each algorithm given three p-value thresholds, 0.000005 (Bonferroni), 0.00005, 0.0005, and one q-value threshold 0.05 (Benjamini-Hochberg). Due to the tradeoff between sensitivity and specificity, with more relaxed thresholds of significance, the precision rates of all methods decrease while the recall rates increase – the 0.05 q-value threshold achieves the lowest precision rate yet the highest recall rate for any given method (Figure 2A). While there does not exist a single method that consistently achieves the highest precision or recall rate, JTK_CYCLE and BIO_CYCLE are more effective in controlling for false positives while still detecting true circadian genes. For the other methods, however, there is a much higher variability in precision, especially in the Zhang 2014 dataset (Figure 2A). RAIN and MetaCycle tend to have the highest sensitivity/recall, but this can come with significant sacrifice on precision (Figure 2A).

In addition, we find that higher sampling frequency can significantly improve the recall rates of all methods. While MetaCycle and RAIN achieve the apparently higher

recall rate under different thresholds in dataset sampled at a lower frequency (2 h/2 days),
all methods, except for LS, produce comparable recall rates when applied to the Hughes
2009 dataset, which is sampled at 1 h/2 days (Figure 2A). Notably, when analyzing the
three datasets with lower sampling frequencies, LS failed under all circumstances with
recall rates less than 0.1 (Figure 2A). This is due to the extreme p-value distribution of
the method with a spike at one, which we will discuss in more detail under "Correlated
multiple testing and non-uniform distribution of p-values under the null".

We further computed with the receiver operating characteristic (ROC) curves with
a varying threshold on the nominal p-values returned by each method (Figure 2B). The
area under the curve (AUC) values serve as a joint measure of sensitivity and specificity
and are above 0.80 across all benchmark results, suggesting that all methods achieve
good sensitivities while controlling for false positive rates. BIO_CYCLE, the deep-
learning-based method, achieves the best performance with the highest AUC across all
datasets (Figure 2B).

Reproducibility assessment using empirical datasets with dark-dark design
Reproducibility is one of the core principles for any bioinformatic tools and yet it remains
a challenge in the field of circadian rhythm detection, which has not been fully explored.
To evaluate the reproducibility of the methods, we first compared and contrasted the
significantly cyclic genes returned by each method across the four datasets. To make the
input dimensions compatible, we selected a total of 7,570 common genes that are shared
across datasets and adopted a q-value threshold of 0.05 for significance. The Venn
diagrams in Figure 3A show the overlapping relationships of the significant genes
returned by each method. While the experimental designs are the same and the observed
gene expression measurements are highly concordant (Figure 1), significant
discrepancies of the calling results are observed. Of the seven benchmarked methods,
ARSER resulted in 721 overlapping significant genes, which is the highest. This is
followed by RAIN, eJTK_CYCLE, MetaCycle, BIO_CYCLE, JTK_CYCLE, and LS with
613, 528, 485, 296, 204, and 0 mutually identified positives, respectively. As mentioned
previously, LS failed in detecting any significant oscillations for three out of the four
datasets.

To further assess the reproducibility of the methods, we computed the Jaccard
index and the Sorensen index to measure the similarities among the results from each
method. Details of these metrics are included in the Materials and Methods section. As a
result, RAIN achieves one of the highest Jaccard indices for any pair of comparisons and
ARSER achieves the highest overall Sorensen index across all datasets (Figure 3B). On
the other hand, our results indicate that JTK_CYCLE, eJTK_CYLE, and BIO_CYCLE
produce the lowest similarity metrics across all comparisons (Figure 3B).
165

Performance assessment using empirical datasets with light-dark design
Next, we adopted four datasets that underwent light-dark experimental design using
different next-generation sequencing platforms (i.e., RNA-seq [22], Nascent-seq [22],
GRO-seq [23], and XR-seq [24]) and named each one after its sequencing protocol (Table
2B). The four datasets have much fewer numbers of time-points compared to the datasets
from the dark-dark design, yet three of the four datasets have technical replicates (Table
2B). More details of the data can be found in the Materials and Methods section. The
oscillatory patterns of known circadian genes are apparent and similar among the various
sequencing technologies (Figure 4A), indicating good data quality.
ARSER, despite its high reproducibility, cannot handle replicates, and previous
studies have shown that data should never be concatenated [17]. Therefore, we focused
on assessing the performance of the other six methods. We first examined the distribution
of the nominal p-values of the 104 gold-standard circadian genes returned by each
method, visualized as beehive plots in Figure 4B, where LS is significantly underpowered
in the detection of circadian genes compared to the other methods, given any of the
sequencing platforms. This result can be attributed to LS's inability to effectively detect
circadian rhythms in datasets with low sampling resolution, which is concordant with our
previous results. We observe that JTK_CYCLE, RAIN, eJTK_CYCLE, MetaCycle, and
BIO_CYCLE can withstand the sparse sampling and result in overall good performance.
To further assess the performance of the methods, we examined the number of
significant genes identified by each method with a false discovery rate (FDR) of 0.05. Of
the 9,481 mutual genes in the four datasets, LS did not identify any significant genes in
any of the datasets. This result aligns with the results from the previous analysis, where

we observed LS as being underpowered. JTK_CYCLE and MetaCycle detected a
relatively small number of significant genes by RNA-seq and XR-seq. eJTK_CYCLE
identified 2,623 significant genes by RNA-seq, and RAIN and BIO_CYCLE identified
2,262 and 1,970 significant genes by XR-seq, respectively. When comparing across
different sequencing platforms, we observe that the number of detected significant genes
from RNA-seq and XR-seq data is much higher than that of the GRO-seq and Nascent-
seq data. This implicates a potential deficiency in detecting gene expression rhythmicity
by measuring nascent transcripts.

With the identified significant genes, we further carried out a gene set enrichment
analysis using the DAVID web server [25, 26] with the default options. Results from the
KEGG pathway enrichment analysis are shown in Supplementary Table 2. We find that
circadian rhythm is significantly enriched by various algorithms, which are marked with
asterisks in Figure 4B. Specifically, we find that of the five methods that were able to
identify statistically significant genes from RNA-seq data, all have enriched circadian
rhythm pathway. Circadian rhythm is also enriched in the three lists of genes that were
identified by eJTK_CYCLE and RAIN as well as two of the three lists of genes identified
by BIO_CYCLE.

Performance assessment using empirical proteomic dataset of dark-dark design
To assess performance of the various methods on non-transcriptomic data, we adopted
a proteomic dataset of mouse livers under dark-dark experimental design from Robles et.
al [27]. Refer to the Materials and Methods section for details. Since this dataset consists
of replicates and missing values, only LS, JTK_CYCLE, RAIN, and MetaCycle were
directly applicable. eJTK_CYCLE was not included due to its inefficiency in handling
random missing values across different genes/proteins. We calculated the number of
significant proteins identified by each method using an FDR threshold of 0.05
(Supplementary Figure 1A). LS identified the least number of oscillatory proteins.
JTK_CYCLE and MetaCycle returned a moderate number of significant proteins. RAIN
identified the largest number of oscillatory proteins, 582, exceeding that of other methods
by more than 300. Heatmaps of scaled measurements of oscillatory proteins identified by
at least two methods are shown in Supplementary Figure 1B, where the proteins are

ordered based on their inferred phases. With the identified oscillatory proteins, we

conducted a gene set enrichment analysis using the DAVID web server. While the results

did not indicate that circadian rhythm was significantly enriched by any of the algorithms,

KEGG metabolic pathways were significantly enriched by all algorithms but LS

(Supplementary Table 3).

Performance assessment using synthetic datasets

To provide guidelines for method selection, we evaluated the performance of the seven

methods in detecting circadian rhythm by simulations with known ground truths.

Examples of waveforms generated for the simulated datasets are shown in

Supplementary Table 4. We generated six groups of simulated datasets to investigate

how key factors affect the performance, including sampling patterns, replicates,

waveforms, signal-to-noise ratios (SNRs), uneven samplings, and missing values.

Supplementary Table 5 outlines the six groups of simulations and we leave the detailed

setup in the Materials and Methods section. Within each simulation group, we repeated

each assessment with three different sampling frequencies to determine whether

increasing sampling frequency may have an effect on the aforementioned factors. The

three sampling frequencies include 4 h/1 day (six time-points), 3 h/1 day (eight time-

points), and 2 h/1 day (twelve time-points) and the results are shown in Figure 5A, 5B,

and 5C, respectively.

Sampling patterns

To determine whether increasing the sampling frequency or lengthening the time-window

is more important for each method, we first evaluated the results under the sampling

pattern of 4 h/1 day versus 8 h/2 days, 3 h/1 day versus 6 h/2 days, and 2 h/1 day versus

4 h/2 days. We did not find strikingly different results within each pair of comparison,

indicating that when the total number of data points are fixed, having a denser sampling

density and enlarging the sampling time-window tend to have similar impact on

performance. However, when we increase the number of data points, the performances

of all methods are improved, which is concordant with existing studies [16, 17].

BIO_CYCLE generally outperforms the other methods, especially in datasets with lower

sampling frequency and shorter time-window, while JTK_CYCLE is the most sensitive to
fewer observations.

Replicates

To investigate the trade-off between replicates and sampling frequency, we compared
the results of higher sampling frequency without replicates to those of lower sampling
frequency with replicates. We first compared the dataset sampled at 4 h /1 day X1 to the
dataset sampled at 8 h/1 day X2. LS, JTK_CYCLE, RAIN, eJTK_CYCLE, and MetaCycle
show better performance with replicates, while BIO_CYCLE performs significantly better
on densely sampled datasets without replicates. Similar results are seen when we applied
the methods to the dataset at 3 h/1 day without replicates and the dataset at 6 h/1 day
with replicates. As expected, further increasing the sampling resolution offsets the
existing preferences that the methods have for inclusion of replicates or higher sampling
density.

Waveforms

Supplementary Table 4 outlines the different types of periodic waveforms that we
generated *in silico* in three broad categories: stationary, non-stationary, and asymmetric
ones. Through our simulations, we find that all of the algorithms perform the best in
detecting non-stationary waveforms. Additionally, all methods, with the exception of
eJTK_CYCLE, perform better on stationary waveforms, compared to asymmetric
waveforms. eJTK_CYCLE and RAIN are the top two methods for identifying asymmetric
waveforms, which are expected due to their design. This is followed by LS, BIO_CYCLE,
MetaCycle, and ARSER. JTK_CYCLE is the least effective in identifying asymmetric
waveforms regardless of sampling frequency.

Signal-to-noise ratios (SNRs)

To test the effects of different noise levels on method performance, we generated various
datasets with signal-to-noise ratios of 3, 2, 1, and 0.5. For all methods, our results suggest
that the larger the SNRs, the higher the accuracy, as expected. LS, MetaCycle, and
BIO_CYCLE are overall the most robust to noises regardless of sampling frequency,
while JTK_CYCLE has the poorest performance given high noise levels.

Uneven samplings

To understand how well the methods deal with uneven samplings, we focus on the results

of datasets with one or more uneven time-points. Our results suggest that BIO_CYCLE

and LS/MetaCycle outperform the other two compatible methods. Under a sparse

sampling design, RAIN and eJTK_CYCLE suffer significantly from an increasing number

of uneven samplings; a dense sampling design, on the other hand, rescues the

aforementioned methods.

Missing values

We generated datasets that contain 1%, 5%, and 10% missing data, and benchmarked

the four methods that allow missing values. The performances of eJTK_CYCLE and RAIN

degrade with an increasing proportion of missing values, while the performances of LS,

JTK_CYCLE, and MetaCycle are comparably invariant, especially under dense sampling

design. We note that eJTK_CYCLE does not handle missing values efficiently, unless the

same sampling time points are missing across all genes, which reduces to uneven

sampling. When there is not a shared missing pattern across different genes, the dataset

needs to be split into multiple uneven sampling cases, and eJTK_CYCLE needs to be

applied separately, followed by results integration. Note that BIO_CYCLE can be applied

to datasets with missing values only if there are replicates and the missingness only

pertains to part of the replicates. We therefore did not include it in the benchmark.

Computational efficiency

Last but not least, we evaluated the computational efficiency across all benchmarked

methods. For dataset with low sampling resolution, the execution times among the

methods are approximately the same (Supplementary Table 6). However, when analyzing

data of larger sizes, RAIN requires significantly more time compared to the other methods.

The running time for LS, ARSER, and BIO_CYCLE does not change much with varying

sampling frequency. The running time for MetaCycle, which integrates results from LS,

JTK_CYCLE, and ARSER, is calculated as the total running time of the three methods.

Correlated multiple testing and non-uniform distribution of p-values under the null

To detect circadian rhythm across thousands of genes, multiple hypothesis testing

corrections are needed [28]. A common FDR threshold of 0.05 is recommended by most

methods and adjusted p-values (q-values) are returned by all methods except for RAIN.
In the previous sections, we adopted both Bonferroni and Benjamini-Hochberg
procedures for corrections. Here, we more carefully examine such procedures and point
out a potential drawback resulted from both correlated multiple testing and non-uniform
distributions of the nominal p-values under the null. We started with the observed
expression measurements from the Hughes 2009 dataset and generated a "null" dataset
by randomly permuting the time labels for each gene (Figure 6A). Such permutations not
only deplete each gene's rhythmic signals but also disrupts any gene-gene correlations
as observed in the raw data, which are high between genes in the same pathways (Figure
6B). As such, all genes upon permutations are under the true null and additionally all
gene-level testing is independent.

Figure 6C shows the distributions of nominal p-values for each method when
applied to the dataset before and after permutation. The "U-shaped" histograms of the p-
values for LS, JTK_CYCLE, MetaCycle, and RAIN using the original data indicate that
there is dependence among the variables in the data. This violates the underlying
assumption of uniformity and raises a red flag for using Bonferroni or FDR for error control
[28]. A few methods have been developed for p-value adjustment when the tests are
correlated [29-31] and such issue has been specifically pointed out by Hutchison and
Dinner [32] for circadian rhythm detection.

We further applied the methods to the permuted data without gene-gene
correlations. The hypothesis testing by LS, JTK_CYCLE, RAIN, and MetaCycle are still
overly conservative, while the testing procedures for ARSER and BIO_CYCLE are biased
with an overabundance of p-values around 0.3 and 0.1, respectively. eJTK_CYCLE
empirically calculates the null distribution of the p-values via permutations and its
enhanced version, booteJTK, speeds up this calculation by approximating the null
distribution of the Kendall's tau using a Gamma distribution [33]. This indeed leads to a
p-value distribution closest to the null. However, neither eJTK_CYCLE nor booteJTK
handles missing values efficiently, as explained previously. As a summary, there is still
room for method development to yield p-values that better match the underlying
assumption of a uniformly distributed p-values under the null.

DISCUSSION

Here, we propose a benchmark framework to systematically evaluate the performance of seven circadian rhythm detection methods, using high-throughput omics data. The empirical datasets that we adopted in this paper were from microarray [18-20] and RNA-seq [22] to measure gene expression, Nascent-seq [22] and GRO-seq [23] to measure nascent RNA, and XR-seq [24] to measure transcription-coupled repair. While these omics data were generated from different platforms, they focus on directly or indirectly profiling transcription. It has been well studied that biological rhythm goes beyond the transcriptomic transcript-level oscillations [34]. For example, post-translational protein acetylation has been linked to circadian rhythm via mass spectrometry [35, 36]. Moreover, it has been shown that a large number of metabolites and proteins exhibit circadian oscillations [27, 37, 38]. The methods and the evaluation procedures are not limited to transcriptomic studies, but can also be applied to acetylomic, metabolomic, and proteomic experiments.

Given the assessment results from both simulations and empirical dataset anaylsis, as well as literature review of the seven methods, we have summarized the strengths and weaknesses of each method in Table 3. In general, LS, RAIN, eJTK_CYCLE, and MetaCycle are more versatile in that they can be applied to datasets with replicates, uneven samplings, or missing values. eJTK_CYCLE and BIO_CYCLE generally outperform the other methods under most situations except for handling missing values. On the other hand, JTK is sensitive to high noise levels and low sampling resolutions, and LS cannot detect any significant genes when sampling resolution is lower than 2 h/2 days with an FDR threshold of 0.05. The best detection algorithm depends on experimental designs and characteristics of the input data. Therefore, we have created two decision trees, one for low sampling resolution and the other for high sampling resolution, that outline the recommended method(s) under different scenarios (Supplementary Figure 2).

Recent advances of high-throughput technologies enable circadian rhythm detection on the genome-wide scale. As with all genomic data, the multi-time-point omics data for circadian rhythm detection bear both technical and biological variability, which can bias the analysis if not properly accounted. Data normalization and batch effect

correction are crucial to remove technical biases and artifacts [39]. Cross-subject
variability in rhythmic profiles, especially for human subjects, is a non-negligible source
of genetic variation that needs to be adjusted [14]. This is especially important in the case-
control setting where multiple subjects are involved. While we did not particularly focus
on differential analysis since it is outside the scope of this paper, a few methods, including
LimoRhyde [40] and DODR [12] have been made available for differential rhythmicity
analysis under different conditions.

Increasingly more circadian omics data are being made available through existing
studies and databases [34, 41]. We showed, from our empirical studies, that the rhythmic
signals can be well recapitulated across different studies and/or different platforms
(Figure 1, Figure 4A). Meta-analysis and multi-omics data integration remain an open-
ended question in circadian rhythm detection [42]. In addition, transfer learning has been
applied to multiple genomic research domains in genomics [43] – to borrow information
and to transfer knowledge from existing data deposited in public repositories remain one
of the future directions. Similarly, across different methods, an ensemble framework, as
implemented by MetaCycle, can potentially boost performance. However, as we have
pointed out earlier, the instability issue needs to be addressed, especially when multiple
drastically distinct results are to be integrated.

To our best knowledge, all existing studies for circadian rhythm detection resort to
bulk-tissue omics data, which characterize an averaged profile across different cell types
in a tissue. The inherent heterogeneity can bias the analysis with reduced power and/or
inflated FDR. Single-cell sequencing circumvents the averaging artifacts associated with
traditional bulk population data and has seen rapid technological developments over the
past few years. To assess the feasibility of single-cell circadian rhythm detection, we *in
silico* generated single-cell RNA sequencing profiles by downsampling bulk RNA-seq
read counts. Gold-standard circadian and noncircadian genes were used to calculate the
associated AUC values (Supplementary Table 7). All methods suffer from low sequencing
depth – a characteristic of the single-cell data. With the decreasing cost and the
increasing popularity of single-cell omics techniques, to profile circadian rhythmicity at the
cellular level and to disentangle within tissue heterogeneity with regard to biological
rhythm can be of great impact.

MATERIALS AND METHODS

Empirical transcriptomic datasets

Three datasets under the dark-dark experimental design including Hughes 2009 [18], Hughes 2012 [19], and Zhang 2014 [20] were downloaded from GEO, and all used microarrays to profile gene expressions (Table 2A). Additionally, we obtained four datasets under the light-dark experimental design from the different sequencing platforms, including Nascent-sequencing (Nascent-seq) [22], RNA-sequencing (RNA-seq) [22], Global Run-On sequencing (GRO-seq) [23], and eXcision Repair-sequencing (XR-seq) [24] (Table 2B). Nascent-seq sequence transcribed RNAs, obtained from the nuclei without formation of the 3' end [44]. GRO-seq measures nascent RNAs by mapping, characterizing, and evaluating transcriptionally engaged polymerase [45]. GRO-seq and Nascent-seq differ from traditional RNA-seq, in which the reads map to predominantly introns, while RNA-seq mainly assays exons [44]. XR-seq profiles DNA excision repair on the genome-wide scale with single-nucleotide resolution [46]. Here, we focus on XR-seq data from the transcribed strand only – it has been shown that the transcription-coupled repair from the transcribed strand is positively correlated with expression [47].

For quality control, we removed genes that had constant gene expression measurements in all datasets and further removed genes with more than half zero gene expression values in the light-dark datasets. In cases where multiple probes got mapped to the same RefSeq loci, we averaged the gene expression of the probes using the limma package [48], available in Bioconductor. For data normalization, robust multi-array average (RMA) [49] and genechip RMA (GC-RMA) [50] were used to normalize the array data; transcript per million (TPM) and reads per kilobase per million reads (RPKM) [51] were used to normalize the transcriptomic sequencing data. We scaled the normalized data within each gene to make them compatible for visualization only, as shown in Figure 1 and Figure 4A.

Empirical proteomic dataset

A proteomic dataset of *Mus musculus* liver tissues from Robles et. al [27] was adopted to detect oscillatory proteins. Mouse liver samples were collected from a total of 64 mice

that were released into constant darkness for one day after being entrained to a 12-12
hour light-dark schedule for 10 days. Four mice were sacrificed every 3 hours for 2 days.
Then, in vivo Stable Isotope Labeling by Amino acids in Cell culture (SILAC) [52, 53] in
combination with mass spectrometry was performed to profile the proteome. For each
time point, equal amount of protein liver extracts from the four mice were mixed together
with equal amount of protein lysates, collected in anti-phase, from the liver samples of
two SILAC mice. The pooled protein extracts were measured with Orbitrap mass
spectrometer. The protein abundance was calculated by taking the ratio of the signal for
the mice and the signal for the heavy SILAC mix. After assessing quantification values, a
total of 3,132 proteins remained for downstream circadian rhythm analysis.
444

Downsampled RNA-seq dataset
We generated several downsampled RNA-seq datasets from the original RNA-seq
dataset under the light-dark design to assess the robustness of the various methods to
low sequencing depths. We obtained the raw sequencing data from GEO, performed read
alignment to the mouse reference genome (mm10) using STAR [54], carried out quality
control procedures on the aligned reads, and obtained integer-valued read counts using
featureCounts [55]. We then generated downsampled RNA-seq data by multinomial
sampling with index 5K, 10K, 50K, 100K, and 500K, and gene-specific probability
parameters calculated from the raw data. RPKM was used to normalize the downsampled
RNA-seq read counts, followed by circadian rhythm detection.
455

Evaluation metrics
To evaluate the performance of the benchmarked methods, we adopted a list of 104
circadian [10] and 113 non-circadian genes [21] in mouse liver as positive and negative
controls, respectively. See Supplementary Table 1 for a full list of these gold-standard
genes. With these gold-standard genes, we calculated metrics including the precision and
recall rates given a p-value or q-value significance threshold (Figure 2A). We further
calculated the AUC values of the ROC curves, as joint measures of sensitivity and
specificity (Figure 2B).

To assess the reproducibility of each method, we compared the results from the

four dark-dark datasets by calculating the number of overlapping genes, as well as the

Jaccard and Sorensen index as metrics for similarity (Figure 3). Venn diagrams are used

to display the number of overlapping cycling genes identified across different datasets by

each method. The Jaccard index measures the pairwise similarities of the significant

genes detected between each pair of datasets. Let A_i and A_j be the set of significant

genes from dataset i and j. The Jaccard similarity index is defined as

471
$$J(A_i, A_j) = \frac{|A_i \cap A_j|}{|A_i \cup A_j|}.$$

472   The Sorensen Index is used to characterize similarity across all datasets [56]:

473
$$S(A_i, A_j, A_k, \ldots) = \frac{T}{T-1}\left(\frac{\sum_{i<j}|A_i \cap A_j| - \sum_{i<j<k}|A_i \cap A_j \cap A_k| + \sum_{i<j<k<l}|A_i \cap A_j \cap A_k \cap A_l| - \cdots}{\sum_i |A_i|}\right)$$

where T is the number of sets compared. Larger number of overlapping genes and larger

Jaccard/Sorensen index values indicate higher reproducibility of the methods.

Simulation setup

Each simulated dataset consists of 6,000 circadian and 6,000 non-circadian gene profiles.

Stationary circadian profiles with a period of 24 hours are used in each simulation group,

as outlined below. Note that when running the methods, we set the period range from 20

to 28 h for all methods except for eJTK_CYCLE and JTK_CYCLE, which either has a

fixed period of 24 h or adjusts the period on the fly. The amplitude of the waveforms is

sampled from a uniform distribution between 1 and 6; the phase shift is sampled from a

uniform distribution between 0 and 24 h; and the noise term is sampled from a standard

normal distribution. Flat waveforms are used to generate non-circadian profiles in all

simulation groups except for testing against non-stationary waveforms where linear lines

are used.

We first aimed to investigate whether higher sampling frequency or longer

sampling time-window is more beneficial for each method. In this simulation group, we

generated two datasets with different sampling frequencies and sampling time-windows.

With six time-points, we generated one dataset at 4 h/1 day and another at 8 h/2 days;

with eight time-points, we generated one dataset at 3 h/1 day and another at 6 h/2 days;

with 12 time-points, we generated one dataset at 2 h/1 day and another at 4 h/2 days.

Next, we assessed whether the inclusion of replicates can offset the effect of low
sampling frequency in methods' ability of detecting oscillations. Replicates are defined as
multiple measurements taken at the same time-point. Specifically, we generated two
datasets consisting of the same number of observations, with or without replicates: one
at 4 h/1 day X1 and the other at 8 h/1 day X2. The sampling design of the other two pairs
of datasets are 3 h/1 day X1 v.s. 6 h/1 day X2, and 2 h/1 day X1 v.s. 4 h/1 day X2.

Since biological rhythms can take on various waveforms, we generated three types
of waveforms via simulation: stationary, non-stationary, and asymmetric curves.
Supplementary Table 4 includes models that we adopted *in silico* to generate the
corresponding waveforms. Specifically, the stationary waveforms include cosine, cosine
2, and cosine peak curves; the non-stationary waveforms include cosine damp, trend
exponential, and trend linear curves; the asymmetric subgroup consists of only the saw-
tooth waveform. We assessed the performance of the methods in identifying each
category of the circadian waveforms.

The next three groups of simulations aimed to determine which methods are more
robust to different levels of signal-to-noise ratios, uneven samplings, and missing values.
Specifically, we generated four datasets with SNRs of 0.5, 1, 2, and 3. Signal-to-noise
ratio is defined by taking the ratio of the empirical variance of cosine function and the
variance of the noise, the latter of which is fixed at one. Uneven samplings are defined
as designs whose time-points are not equally spaced. To investigate the effect of uneven
samplings on performance, we generated datasets with one, two, or four uneven
samplings. With six time-points, datasets with four uneven samplings cannot be
generated as it would only have two time-points. For missing data, we generated three
levels of missing data (1%, 5%, and 10%) at three fixed, randomly selected time-points.

Lastly, we generated three datasets with sampling patterns of 1 h/2 days, 2 h/2
days, and 4 h/2 days to compute the execution times for each method. We seek to identify
the differences in computational efficiency among the methods and to explore the effect
of increasing sampling resolution on the execution time. Each dataset consists of a total
of 6,000 genes. All execution times are reported by running on a Macbook Pro (15-inch,
2019) with 2.3 GHz 8-Core Intel Core i9 and 16 GB memory.

DATA AND SOFTWARE AVAILABILITY

MetaCycle is an open-source R package available at https://github.com/gangwug/MetaCycle and is also used for individual analysis for LS, JTK_CYCLE, and ARSER. RAIN is a Bioconductor R package available at https://bioconductor.org/packages/rain/. eJTK_CYCLE was downloaded from https://github.com/alanlhutchison/empirical-JTK_CYCLE-with-asymmetry. BIO_CYCLE was downloaded from http://circadiomics.igb.uci.edu/BIO_CYCLE. All empirical datasets were downloaded from the NCBI Gene Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/). The accession numbers for dark-dark datasets are GSE11923, GSE30411, and GSE54652, respectively. The accession numbers for light-dark datasets are GSE59486, GSE36872, GSE36871 and GSE109938, respectively. The proteomic dataset was downloaded from the BioStudies database with accession number S-EPMC3879213.

ACKNOWLEDGEMENTS

This work was supported by NIH Grants R35 GM118102 (to A.S.), R01 ES027255 (to A.S.), P01 CA142538 (to Y.J.), UL1 TR002489 (to Y.J.), and a pilot award from the UNC Computational Medicine Program (to Y.J.). We thank the Sancar Lab members and Dr. John Hogenesch for helpful discussions and feedback.

REFERENCES

1. Li J, Grant GR, Hogenesch JB, Hughes ME: **Considerations for RNA-seq analysis of circadian rhythms.** *Methods Enzymol* 2015, **551:**349-367.

2. Panda S, Antoch MP, Miller BH, Su AI, Schook AB, Straume M, Schultz PG, Kay SA, Takahashi JS, Hogenesch JB: **Coordinated transcription of key pathways in the mouse by the circadian clock.** *Cell* 2002, **109:**307-320.

3. Hughes ME, Hogenesch JB, Kornacker K: **JTK_CYCLE: an efficient nonparametric algorithm for detecting rhythmic components in genome-scale data sets.** *J Biol Rhythms* 2010, **25:**372-380.

4. Asher G, Sassone-Corsi P: **Time for food: the intimate interplay between nutrition, metabolism, and the circadian clock.** *Cell* 2015, **161:**84-92.

5. Partch CL, Green CB, Takahashi JS: **Molecular architecture of the mammalian circadian clock.** *Trends Cell Biol* 2014, **24:**90-99.

558   6.   Roenneberg T, Merrow M: **The Circadian Clock and Human Health.** *Curr Biol* 2016,
559   **26:**R432-443.

560   7.   Levi F, Schibler U: **Circadian rhythms: mechanisms and therapeutic implications.** *Annu*
561   *Rev Pharmacol Toxicol* 2007, **47:**593-628.

562   8.   Glynn EF, Chen J, Mushegian AR: **Detecting periodic patterns in unevenly spaced gene**
563   **expression time series using Lomb-Scargle periodograms.** *Bioinformatics* 2006, **22:**310-
564   316.

565   9.   Wijnen H, Naef F, Young MW: **Molecular and statistical tools for circadian transcript**
566   **profiling.** *Methods Enzymol* 2005, **393:**341-365.

567   10.   Wu G, Zhu J, Yu J, Zhou L, Huang JZ, Zhang Z: **Evaluation of five methods for genome-**
568   **wide circadian gene identification.** *J Biol Rhythms* 2014, **29:**231-242.

569   11.   Yang R, Su Z: **Analyzing circadian expression data by harmonic regression based on**
570   **autoregressive spectral estimation.** *Bioinformatics* 2010, **26:**i168-174.

571   12.   Thaben PF, Westermark PO: **Detecting rhythms in time series with RAIN.** *J Biol Rhythms*
572   2014, **29:**391-400.

573   13.   Hutchison AL, Maienschein-Cline M, Chiang AH, Tabei SM, Gudjonson H, Bahroos N,
574   Allada R, Dinner AR: **Improved statistical methods enable greater sensitivity in rhythm**
575   **detection for genome-wide data.** *PLoS Comput Biol* 2015, **11:**e1004094.

576   14.   Wu G, Anafi RC, Hughes ME, Kornacker K, Hogenesch JB: **MetaCycle: an integrated R**
577   **package to evaluate periodicity in large scale data.** *Bioinformatics* 2016, **32:**3351-3353.

578   15.   Agostinelli F, Ceglia N, Shahbaba B, Sassone-Corsi P, Baldi P: **What time is it? Deep**
579   **learning approaches for circadian rhythms.** *Bioinformatics* 2016, **32:**i8-i17.

580   16.   Deckard A, Anafi RC, Hogenesch JB, Haase SB, Harer J: **Design and analysis of large-scale**
581   **biological rhythm studies: a comparison of algorithms for detecting periodic signals in**
582   **biological data.** *Bioinformatics* 2013, **29:**3174-3180.

583   17.   Hughes ME, Abruzzi KC, Allada R, Anafi R, Arpat AB, Asher G, Baldi P, de Bekker C, Bell-
584   Pedersen D, Blau J, et al: **Guidelines for Genome-Scale Analysis of Biological Rhythms.** *J*
585   *Biol Rhythms* 2017, **32:**380-393.

586   18.   Hughes ME, DiTacchio L, Hayes KR, Vollmers C, Pulivarthy S, Baggs JE, Panda S, Hogenesch
587   JB: **Harmonics of circadian gene transcription in mammals.** *PLoS Genet* 2009, **5:**e1000442.

588   19.   Hughes ME, Hong HK, Chong JL, Indacochea AA, Lee SS, Han M, Takahashi JS, Hogenesch
589   JB: **Brain-specific rescue of Clock reveals system-driven transcriptional rhythms in**
590   **peripheral tissue.** *PLoS Genet* 2012, **8:**e1002835.

591   20.   Zhang R, Lahens NF, Ballance HI, Hughes ME, Hogenesch JB: **A circadian gene expression**
592   **atlas in mammals: implications for biology and medicine.** *Proc Natl Acad Sci U S A* 2014,
593   **111:**16219-16224.

594   21.   Wu G, Zhu J, He F, Wang W, Hu S, Yu J: **Gene and genome parameters of mammalian**
595   **liver circadian genes (LCGs).** *PLoS One* 2012, **7:**e46961.

22. Menet JS, Rodriguez J, Abruzzi KC, Rosbash M: **Nascent-Seq reveals novel features of**
mouse circadian transcriptional regulation. *Elife* 2012, **1:**e00011.

23. Fang B, Everett LJ, Jager J, Briggs E, Armour SM, Feng D, Roy A, Gerhart-Hines Z, Sun Z,
Lazar MA: **Circadian enhancers coordinate multiple phases of rhythmic gene**
transcription in vivo. *Cell* 2014, **159:**1140-1152.

24. Yang Y, Adebali O, Wu G, Selby CP, Chiou YY, Rashid N, Hu J, Hogenesch JB, Sancar A:
Cisplatin-DNA adduct repair of transcribed genes is controlled by two circadian
programs in mouse tissues. *Proc Natl Acad Sci U S A* 2018, **115:**E4777-E4785.

25. Huang da W, Sherman BT, Lempicki RA: **Systematic and integrative analysis of large gene**
lists using DAVID bioinformatics resources. *Nat Protoc* 2009, **4:**44-57.

26. Huang da W, Sherman BT, Lempicki RA: **Bioinformatics enrichment tools: paths toward**
the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 2009, **37:**1-
13.

27. Robles MS, Cox J, Mann M: **In-vivo quantitative proteomics reveals a key contribution**
of post-transcriptional mechanisms to the circadian regulation of liver metabolism.
PLoS Genet 2014, **10:**e1004047.

28. Storey JD, Tibshirani R: **Statistical significance for genomewide studies.** *Proc Natl Acad*
Sci U S A 2003, **100:**9440-9445.

29. Li J, Ji L: **Adjusting multiple testing in multilocus analyses using the eigenvalues of a**
correlation matrix. *Heredity (Edinb)* 2005, **95:**221-227.

30. Conneely KN, Boehnke M: **So many correlated tests, so little time! Rapid adjustment of**
P values for multiple correlated tests. *Am J Hum Genet* 2007, **81:**1158-1168.

31. Schwartzman A, Lin X: **The effect of correlation in false discovery rate estimation.**
Biometrika 2011, **98:**199-214.

32. Hutchison AL, Dinner AR: **Correcting for Dependent P-values in Rhythm Detection.**
BioRxiv 2017.

33. Hutchison AL, Allada R, Dinner AR: **Bootstrapping and Empirical Bayes Methods Improve**
Rhythm Detection in Sparsely Sampled Data. *J Biol Rhythms* 2018, **33:**339-349.

34. Ceglia N, Liu Y, Chen S, Agostinelli F, Eckel-Mahan K, Sassone-Corsi P, Baldi P: **CircadiOmics:**
circadian omic web portal. *Nucleic Acids Res* 2018, **46:**W157-W162.

35. Masri S, Patel VR, Eckel-Mahan KL, Peleg S, Forne I, Ladurner AG, Baldi P, Imhof A,
Sassone-Corsi P: **Circadian acetylome reveals regulation of mitochondrial metabolic**
pathways. *Proc Natl Acad Sci U S A* 2013, **110:**3339-3344.

36. Mauvoisin D, Atger F, Dayon L, Nunez Galindo A, Wang J, Martin E, Da Silva L, Montoliu I,
Collino S, Martin FP, et al: **Circadian and Feeding Rhythms Orchestrate the Diurnal Liver**
Acetylome. *Cell Rep* 2017, **20:**1729-1743.

37. Dallmann R, Viola AU, Tarokh L, Cajochen C, Brown SA: **The human circadian metabolome.**
Proc Natl Acad Sci U S A 2012, **109:**2625-2629.

38.   Feng D, Lazar MA: **Clocks, metabolism, and the epigenome.** *Mol Cell* 2012, **47:**158-167.

39.   Jiang Y, Wang R, Urrutia E, Anastopoulos IN, Nathanson KL, Zhang NR: **CODEX2: full-**
636   **spectrum copy number variation detection by high-throughput DNA sequencing.**
637   *Genome Biol* 2018, **19:**202.

638   40.   Singer JM, Fu DY, Hughey JJ: **Simphony: simulating large-scale, rhythmic data.** *PeerJ* 2019,
639   **7:**e6985.

640   41.   Li X, Shi L, Zhang K, Wei W, Liu Q, Mao F, Li J, Cai W, Chen H, Teng H, et al: **CirGRDB: a**
641   **database for the genome-wide deciphering circadian genes and regulators.** *Nucleic Acids*
642   *Res* 2018, **46:**D64-D70.

643   42.   Patel VR, Eckel-Mahan K, Sassone-Corsi P, Baldi P: **CircadiOmics: integrating circadian**
644   **genomics, transcriptomics, proteomics and metabolomics.** *Nat Methods* 2012, **9:**772-
645   773.

646   43.   Eraslan G, Avsec Z, Gagneur J, Theis FJ: **Deep learning: new computational modelling**
647   **techniques for genomics.** *Nat Rev Genet* 2019, **20:**389-403.

648   44.   Trott AJ, Menet JS: **Regulation of circadian clock transcriptional output by CLOCK:BMAL1.**
649   *PLoS Genet* 2018, **14:**e1007156.

650   45.   Core LJ, Waterfall JJ, Lis JT: **Nascent RNA sequencing reveals widespread pausing and**
651   **divergent initiation at human promoters.** *Science* 2008, **322:**1845-1848.

652   46.   Hu J, Adar S, Selby CP, Lieb JD, Sancar A: **Genome-wide analysis of human global and**
653   **transcription-coupled excision repair of UV damage at single-nucleotide resolution.**
654   *Genes Dev* 2015, **29:**948-960.

655   47.   Yimit A, Adebali O, Sancar A, Jiang Y: **Differential damage and repair of DNA-adducts**
656   **induced by anti-cancer drug cisplatin across mouse organs.** *Nat Commun* 2019, **10:**309.

657   48.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK: **limma powers differential**
658   **expression analyses for RNA-sequencing and microarray studies.** *Nucleic Acids Res* 2015,
659   **43:**e47.

660   49.   Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, Speed TP:
661   **Exploration, normalization, and summaries of high density oligonucleotide array probe**
662   **level data.** *Biostatistics* 2003, **4:**249-264.

663   50.   Wu Z, Irizarry RA, Gentleman R, Martinez-Murillo F, Spencer F: **A Model-Based**
664   **Background Adjustment for Oligonucleotide Expression Arrays.** *Journal of the American*
665   *Statistical Association* 2004, **99:**909-917.

666   51.   Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A, Szczesniak
667   MW, Gaffney DJ, Elo LL, Zhang X, Mortazavi A: **A survey of best practices for RNA-seq**
668   **data analysis.** *Genome Biol* 2016, **17:**13.

669   52.   Geiger T, Cox J, Ostasiewicz P, Wisniewski JR, Mann M: **Super-SILAC mix for quantitative**
670   **proteomics of human tumor tissue.** *Nat Methods* 2010, **7:**383-385.

671   53.   Gouw JW, Krijgsveld J, Heck AJ: **Quantitative proteomics by metabolic labeling of model**
672   **organisms.** *Mol Cell Proteomics* 2010, **9:**11-24.

673   54.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
674   TR: **STAR: ultrafast universal RNA-seq aligner.** *Bioinformatics* 2013, **29:**15-21.

675   55.   Liao Y, Smyth GK, Shi W: **featureCounts: an efficient general purpose program for**
676   **assigning sequence reads to genomic features.** *Bioinformatics* 2014, **30:**923-930.

56.   Diserud OH, Odegaard F: **A multiple-site similarity measure.** *Biol Lett* 2007, **3:**20-22.

<u>FIGURE & TABLE LEGENDS</u>

Figure 1. Examples of circadian and non-circadian benchmark gene expressions

among three datasets with dark-dark experimental design. Scaled gene expressions

from selected (A) circadian genes including *Clock*, *Cry1*, *Npas2*, and *Per1* and (B) non-

circadian genes including *Utp6*, *Mtf1*, *Cln3*, *Abcd4*.

Figure 2. Evaluation of seven methods by precision, recall rates and ROC curves.

(A) A p-value threshold of 0.000005 (Bonferroni threshold), 0.00005, 0.0005, and a q-

value threshold of 0.05 (FDR threshold) are adopted for each of the seven methods

applied to the four dark-dark empirical datasets. A more relaxed threshold results in a

higher recall rate, with FDR being the most sensitive, yet this also leads to a higher

number of false positives with a lower precision rate. (B) ROC curves and AUC values

using gold-standard circadian and non-circadian genes. Each method is evaluated across

four dark-dark empirical datasets. Sensitivity and specificity are calculated using the

nominal p-values by each method with varying threshold. BIO_CYCLE returns the highest

AUC.

Figure 3. Evaluation of method reproducibility. (A) Venn diagrams display the number

of cyclic genes that are significant by each method among the four dark-dark datasets.

(B) Jaccard index and the Sorensen index are used as metrics for reproducibility for each

method across the four datasets with the same experimental design.

Figure 4. Circadian rhythm detection under light-dark experimental design by GRO-

seq, Nascent-seq, RNA-seq, and XR-seq. (A) Gene-specific measurements of nascent

RNA, RNA, and transcription-coupled repair of four circadian benchmark genes, *Clock*,
Npas2, *Cry1*, and *Per1* by four different sequencing platforms. The solid and dotted lines
are used for the first and second replicates respectively. (B) Beehive plots of negative log
p-values of base 10 of circadian genes as positive controls. The number of significant
genes detected by each method with an FDR threshold of 0.05 are shown in parenthesis.
The asterisks denote significant GO enrichments of circadian rhythm pathway. The
nominal p-values by JTK_CYCLE, MetaCycle, and BIO_CYCLE are the most significant,
while LS and RAIN tend to be underpowered. ARS is not included in the analysis because
it cannot be applied to datasets with replicates.

Figure 5. Performance assessment via simulation studies. Seven circadian rhythm
detection methods are evaluated under different experimental designs to explore how
sampling patterns, replicates, waveforms, signal-to-noise ratios (SNRs), uneven
samplings, and missing values affect performance. Simulations under each design are
carried out with different sampling frequencies: (A) 4 h/1 day, (B) 3 h/1 day, and (C) 2 h/1
day. AUC values calculated from ground truths are used as metrics.

Figure 6. Existing methods return non-uniformly distributed p-values under the null,
partially due to non-independent testing due to gene-gene correlations. (A) Gene
expression values for the benchmark circadian gene *Cry1* before and after random
permutations of the time labels. (B) Heatmaps of pairwise correlation coefficients among
the top 200 highly variable genes from the Hughes 2009 dataset. The top illustrates the
gene-gene correlation coefficients calculated from raw data input, and the bottom shows
the gene-gene correlations after permutation. (C) The distributions of nominal p-values
for each method when applied to the dataset before and after permutation. Gene-gene
correlations, which are accounted for by eJTK_CYCLE, partially lead to the systematic
deviations from the null distributions. The hypothesis testing by LS, JTK_CYCLE, RAIN,
and MetaCycle are overly conservative, while ARSER's and BIO_CYCLE's testing
procedures are biased with an overabundance of p-values around 0.3 and 0.1,
respectively, under the null.

Table 1. Summary of seven existing methods for circadian rhythm detection. [a]
BIO_CYCLE can be applied to datasets with missing values only if there are replicates
and the missingness only pertains to part of the replicates.

Table 2. High-throughput mouse liver datasets adopted for circadian rhythm
detection. (A) Dark-dark experimental design. (B) Light-dark experimental design.

Table 3. Pros and cons of circadian rhythm detection methods.

SUPPLEMENTARY FIGURE & TABLE LEGENDS

Supplementary Table 1. Circadian and non-circadian genes in *Mus muculus* liver
as gold standard. The 104 circadian gene list is extracted from Supplementary Table 4
in Wu et al. Wu G, Zhu J, Yu J, Zhou L, Huang JZ and Zhang Z [10] and the 113 non-
circadian gene list is obtained from Supplementary Table 2 in Wu et al. Wu G, Zhu J, He
F, Wang W, Hu S and Yu J [21].

Supplementary Table 2. Pathway enrichment analysis of significantly cyclic genes
from the light-dark datasets. Functional annotations (KEGG pathway mapping) of the
significant genes (q-values ≤ 0.05) are carried out using the the DAVID Bioinformatics
Resources (https://david.ncifcrf.gov/). The list only contains significantly enriched
pathways with a 0.05 cutoff of the p-values adjusted by Benjamini Hochberg.

Supplementary Table 3. Pathway enrichment analysis of significantly cyclic
proteins. Functional annotations (KEGG pathway mapping) of the significant proteins (q-
values ≤ 0.05) are carried out using the the DAVID Bioinformatics Resources
(https://david.ncifcrf.gov/). The list only contains significantly enriched pathways with a
0.05 cutoff of the p-values adjusted by Benjamini Hochberg. KEGG metabolic pathways
were enriched by all three methods.

Supplementary Table 4. *In silico* generated periodic v.s. non-periodic gene profiles.
Three types of periodic waveforms are included: stationary, non-stationary, and

asymmetric. The stationary and non-stationary subgroups consist of three forms of cosine
curves. The asymmetric subgroup consists of a saw-tooth waveform. Flat or linear lines
are adopted to generate non-periodic waveforms. The waveforms shown are constructed
without noise. 'Amp', 'pha', and 'per' represent amplitude, phase and period, respectively.
769

Supplementary Table 5. Details of simulation setup and parameters used to _in silico_
generate periodic and non-periodic profiles. Each simulation run consists of 6,000
periodic and 6,000 non-periodic gene profiles. All simulated waveforms have a period
length of 24, a phase shift that is uniformly distributed between 0 and 24, and a noise
term with standard normal distribution. The amplitude is uniformly distributed between 1
and 6 for all groups except when testing for different signal-to-noise ratios (SNRs), which
we define as the ratios of the empirical variances of the cosine function and the variances
of the noise. Non-periodic profiles are sampled from a flat/linear function. "X 1" indicates
no replicate and "X 2" indicates two replicates.
779

Supplementary Table 6. Evaluation of computational efficiency with different
sampling rates. Each method is run on a dataset with a total of 6,000 genes. All
programs are run on a Macbook Pro (15-inch, 2019) with 2.3 GHz 8-Core Intel Core i9
and 16 GB memory. Running time for MetaCycle is the sum of the runing time for LS,
ARSER, and JTK_CYCLE. Running time for BIO_CYCLE does not include the time used
to fit the deep neural network.
786

Supplementary Table 7. Performance assessment of downsampled RNA-seq data.
AUC values of downsampled RNA-seq datasets with varying sequencing depths were
calculated. Existing methods suffer from low sequencing depths. The performance of
RAIN exceeds that of all other methods in all sequencing depths with an exception at 5K,
due to its large number of significant genes detected in general. BIO_CYCLE consistently
ranks the lowest at all but the highest sequencing depth. The performances of LS,
JTK_CYCLE, eJTK_CYCLE, and MetaCycle are comparable.
794

**Supplementary Figure 1. Circadian rhythm detection of *Mus musculus* liver
protemoic dataset.** (A) Bar plot of the number of significant proteins detected by each
method using an FDR threshold of 0.05. Only methods that are able to handle both
replicates and missing values were applied and evaluated. (B) Heatmap of scaled
measurements of oscillatory proteins identified by at least two methods. Proteins (rows)
are ordered based on their inferred phases.
801
802   **Supplementary Figure 2. Decision tree as user guidance on method selection.** The
803   decision tree has decision rules for sampling resolutions, uneven samplings, replicates,
804   and missing values.

Figure 1

Figure 2

Figure 3

(A)

(B)

Figure 4

Figure 5

Package	Method Key Words	Method Type	Reference	Availability	Language	Replicates	Missing Values	Uneven Sampling
Lomb-Scargle (LS)	Periodogram	Parametric	*Bioinformatics (2006)*	https://www.iiap.res.in/astrostat/tuts/Lomb-Scargle.html	R	✔	✔	✔
ARSER (ARS)	Harmonic Regression	Parametric	*Bioinformatics (2010)*	http://bioinformatics.cau.edu.cn/ARSER	Python & R	✘	✘	✘
JTK_CYCLE (JTK)	Kendall's Tau	Non-parametric	*J Biol Rhythms (2010)*	https://openwetware.org/wiki/HughesLab:JTK_Cycle	R	✔	✔	✘
RAIN	Asymmetric waveforms	Non-parametric	*J Biol Rhythms (2014)*	http://bioconductor.org/packages/rain	R	✔	✔	✔
eJTK_CYCLE (eJTK)	Empirical *p*-values	Non-parametric	*PLOS Comp. Bio. (2015)*	https://github.com/alanlhutchison/empirical-JTK_CYCLE-with-asymmetry	Python	✔	✔	✔
MetaCycle (MC)	Integration	Parametric	*Bioinformatics (2016)*	https://cran.r-project.org/package=MetaCycle	R	✔	✔	✔
BIO_CYCLE (BC)	Deep Neural Network	Parametric	*Bioinformatics (2016)*	http://circadiomics.igb.uci.edu	R	✔	✔/✘ [a]	✔

Table 1

Design	Name	Reference	Accession Number	Tissue Type	Sequencing Platform	Number of Time Points & Replicates	Number of Genes	Time Points
(A) Dark-Dark	Hughes et al.	*PLOS Genetics (2009)*	GSE11923	Liver	Microarray	48 x 1	13,029	CT18, 19, 20, …, 65
	Hughes et al. (downsampled)	*PLOS Genetics (2009)*	GSE11923	Liver	Microarray	24 x 1	12,506	CT18, 20, 22, …, 64
	Hughes et al.	*PLOS Genetics (2012)*	GSE30411	Liver	Microarray	24 x 1	14,413	CT0, 2, 4, …, 46
	Zhang et al.	*PNAS (2014)*	GSE54652	Liver	Microarray	24 x 1	20,307	CT18, 20, 22, …, 64
(B) Light-Dark	Fang et al.	*Cell (2014)*	GSE59486	Liver	GRO-seq	8 x 1	17,463	ZT1, 4, 7, 10, 13, 16, 19, 22
	Menet et al.	*eLIFE (2012)*	GSE36872	Liver	Nascent-seq	6 x 2	17,917	ZT0, 4, 8, 12, 16, 20
	Menet et al.	*eLIFE (2012)*	GSE36871	Liver	RNA-seq	6 x 2	17,222	ZT2, 6, 10, 14, 18, 22
	Yang et al.	*PNAS (2018)*	GSE109938	Liver	XR-seq (TS)	6 x 2	17,652	ZT0, 4, 8, 12, 16, 20

Table 2

Methods	Pros	Cons
LS	• Effective in handling missing values • Not restricted by input data structure (i.e. can be applied to datasets with replicates, uneven samplings, or missing values)	• Rapid degradation in detectability when applied to datasets with low sampling resolution • U-shaped p-values distribution • Sensitive to outliers
ARSER	• High reproducibility	• Cannot handle replicates, uneven samplings, or missing values
JTK_CYCLE	• High precision • Robust to outliers	• Incapable of detecting asymmetric waveforms • U-shaped p-values distribution • Sensitive to high level of noise • High false negative rates • Low reproducibility
RAIN	• High recall • Effective in detecting asymmetric waveforms • High reproducibility • Not restricted by input data structure	• High false positive rates • U-shaped p-values distribution • Computationally intensive with increasing sampling resolution
eJTK_CYCLE	• Uniform distribution of nominal p-values • Most effective in detecting asymmetric waveforms	• Unable to test different periods simultaneously • Inefficient in handling missing values • Sensitive to high level of uneven samplings
MetaCycle	• High recall • Not restricted by input data structure • Offset the disadvantages of one method with the other two among LS, ARSER and JTK_CYCLE • Directly return calling results from three perspective methods and perform ensemble	• P-values generated with Fisher's integration require independence assumption
BIO_CYCLE	• Most effective in controlling for false positive rates • Most robust to data with high noise, uneven samplings, and low sampling resolutions. • High precision • High computational efficiency with pre-trained model	• Require extensive time to train the DNN model • Handle missing values only if data have replicates and the missingness only pertains to part of the replicates. • Low reproducibility

Table 3