Can biomass distribution across trophic levels predict trophic cascades?

Núria Galiana[1*], Jean-François Arnoldi[2], Matthieu Barbier[1], Amandine Acloque[1], Claire de Mazancourt[1] and Michel Loreau[1]

[1] *Theoretical and Experimental Ecology Station,*
CNRS and Paul Sabatier University, 09200 Moulis, France.
[2] *Zoology department, School of Natural Sciences,*
Trinity College Dublin, The University of Dublin, Ireland.
*nuria.galianaibanez@sete.cnrs.fr

July 20, 2020

Abstract

Food chains present an interconnected set of properties. Some empirically accessible, such as the biomass distribution across trophic levels, others more elusive, such as cascading responses to perturbations. Intuitive links connect these structural and dynamical properties. Yet, do more abundant predators always entail stronger trophic cascades? Using a combination of theory and data from 31 pelagic experiments, we explore whether static features carry direct information about dynamical responses. We identify three regimes, where upward, downward or both cascades shape the biomass distribution. These regimes depend on what we call trophic dissipation, i.e. the balance between consumer mortality and primary productivity modulated by species interactions. Within-ecosystem causality translates into across-ecosystem patterns if trophic dissipation varies little among systems, as we show in experimental data. Our approach can reveal direct causal links between trophic cascades and biomass distributions, and thus guide predictions from empirical patterns.

1 Introduction

Food chains are a central concept of ecology. They provide intuitions and predictions for basic static properties of food webs, such as the distribution of biomass across the trophic hierarchy (Lindeman, 1942; Elton, 1927). They also predict dynamical behaviour, such as cascading perturbation effects known as trophic cascades (Shurin *et al.*, 2002; Borer *et al.*, 2005; Carpenter *et al.*, 1985; Polis *et al.*, 2000). Yet, studies devoted to understanding each of these properties have historically developed in isolation, generating different lines of investigation, which have seldom been connected rigorously (but see Jonsson (2017); McCauley *et al.* (2018); Barbier & Loreau (2019); Rossberg *et al.* (2019)). In particular, while trophic cascades are prevalent in nature, their strength has been shown to be highly variable across systems (Polis *et al.*, 2000; Shurin *et al.*, 2002; Frank *et al.*, 2006), prompting the widespread emergence of studies exploring the mechanisms behind their existence and magnitude (Halaj & Wise, 2001; Shurin & Seabloom, 2005; Borer *et al.*, 2005; Heath *et al.*, 2014; Ripple *et al.*, 2016).

While previous studies investigated the mechanisms that control the strength of trophic cascades and biomass distributions, there exists no consensus on whether cascade strength can be inferred from biomass distributions. A number of applied studies proposed to predict an ecosystem's "health" and dynamical response from its more accessible static biomass pyramid or size spectrum (Shin *et al.*, 2005; Cury *et al.*, 2005). Yet others claimed that these static features are not informative about the underlying food chain dynamics, due to exogenous factors such as omnivory and energetic subsidies (Trebilco *et al.*, 2016; McCauley *et al.*, 2018; Woodson *et al.*, 2018).

Even in a perfect food chain, there are several ways in which cascades and biomass distribution could be related. They could correlate across ecosystems via dependency on common parameters, as suggested by recent theory (Heath *et al.*, 2014; Barbier & Loreau, 2019). But they could also be causally related, in the sense that the observed biomass distribution may be directly shaped by cascading effects (Figure 1). A direct causal link is required to reliably predict dynamical behaviors from static snapshots alone, without additional knowledge of potential confounding factors.

This idea is at the core of the present study. We specifically ask when the static shape of the food chain, i.e. the biomass distribution across trophic levels, can be causally related to its dynamical responses, i.e. the propagation of a perturbation downward from the top of the food chain, or upward from the bottom, causing trophic cascades (Figure 1).

The simplest expectation, for any ecosystem with multiple compartments, is that the response of a compartment to a perturbation is proportional to its standing biomass (Figure 2a). This expectation will serve as a baseline:

H0 (Proportional response):
The response of any trophic level to any perturbation is proportional to its biomass.

This expectation may appear intuitive, yet many classical results on food chains do *not*, in fact, support it. On a quantitative level, deviations from strict proportionality occur in many theoretical and empirical studies, and have been used to define whether a cascade is strong or weak (Hedges *et al.*, 1999; Shurin *et al.*, 2002). These deviations are commonly interpreted as an amplification or attenuation of perturbations as they propagate along the chain (Box 1). For instance, adding nutrients at the bottom of the food chain might not increase the abundance of primary producers, but only lead to a boom in consumers (Oksanen *et al.*, 1981).

On a qualitative level, the hypothesis H0 implies that *both* upward and downward effects are directly related to the biomass distribution. Yet, it has long been noticed that these two types of effects can differ in their importance from system to system. Each has been the focus of a classic perspective on food chain structure and dynamics. The resource-based perspective posits that resources determine the biomass of higher trophic levels (Elton, 1927; Lindeman, 1942). The consumer-based perspective, however, emphasizes the role of consumers in regulating the biomass of lower trophic levels (Hairston *et al.*, 1960; Leibold, 1996; Gruner *et al.*, 2008). Thus, if a link between biomass distributions and cascades should exist, it is unclear whether it should involve upwards dynamical effects, downwards effects or both at once.

The key intuition that we explore here is that standing biomasses should reflect the dynamical responses to those ecological drivers, such as primary productivity or consumer mortality, that are *already dominant in the observed state*. However, it cannot inform us about the response to other ecological drivers that can occur as perturbations. For instance, when consumer mortality is negligible, the standing biomass distribution is entirely driven by bottom-up propagation of primary productivity, and cannot tell us whether response to a top-down perturbation (e.g. predator removal) would be weak or strong (Barbier & Loreau, 2019). Conversely, in some models such as "exploitation ecosystems" (Oksanen *et al.*, 1981), standing biomasses are driven by top-down effects, with primary productivity only controlling the number of trophic levels in the food chain.

Here we make a precise theoretical claim about which dynamical features can or cannot be predicted from biomass distributions. We show that key ecological parameters – including primary productivity, trophic interactions and self-regulation (e.g. intra-guild competition or interference) – only affect the relationship between dynamics and biomasses through a synthetic mechanism, which we call *trophic dissipation*, and its impact on both consumer and resource biomasses (Box 1, Figure 2). We then ask when this causal relationship within one ecosystem can translate into a clear statistical pattern across ecosystems using data from 31 mesocosm experiments of pelagic food webs, which monitored standing biomass and long-term responses to top-down and bottom-up perturbations (fish and nutrient addition).

Combining theory and data, we find that the experimental systems demonstrate both types of relationships between biomass and cascades, i.e. correlations driven by common factor, and direct causation (Figure 1). We conclude that understanding the entangled dependencies between biomass distribution and dynamical responses is essential to properly interpret the patterns observed in

natural communities, and enhance our capacity to predict the long-term effects of perturbations.

2 Methods

2.1 Empirical testing

2.1.1 Data collection and formatting

We searched the literature for studies analysing long-term bottom-up and top-down effects after press perturbations in food chains. We selected 21 independent studies reporting on 31 lake mesocosm experiments of pelagic food webs that performed the same experimental treatments (see Appendix S1; Hulot *et al.* (2014)). Specifically, all studies investigated community responses to nutrient enrichment, fish addition, and both perturbations together. They reported species biomass (or determined trophic levels) before and after the perturbation, which is fundamental to test whether biomass distribution across trophic levels is related to the strength of trophic cascades. Detailed information about the studies used can be found in Appendix S1.

The pelagic communities considered contain different paths of energy transmission across trophic levels (Figure 3a). To properly study biomass distribution across trophic levels and, in particular, the dynamics of trophic cascades in the community, it is fundamental to identify paths of energy flux in the food web that adequately behave as food chains (Barbier & Loreau, 2019). We thus define as *dynamically consistent chains* those where prey and predators respond in the same direction to bottom-up perturbations, and in opposite directions to top-down perturbations (Box 1). We decomposed empirical food webs into all the possible chains and analysed them independently to identify which responded to perturbations the way an isolated chain would (Figure 3a). Not all the experiments considered had information for all the food web compartments. Thus, the decomposition of the complex food webs was based on the information available (Appendix S1). To ensure that the decomposition of the food webs into multiple chains do not bias our analyses we included study system as a random effect in our statistical models (see below).

2.1.2 Data analyses

The third trophic level (fish) was viewed as a perturbation treatment in the experiments considered (only presence/absence reported). Therefore, we characterized the biomass distribution only for the first two trophic levels (phytoplankton and zooplankton), using the ratio

$$B_r = \frac{B_2^*}{B_1^*} \tag{1}$$

where B_i^*, $i = 1, 2$ corresponds to the non-perturbed biomass of trophic level 1 (phytoplankton) and level 2 (zooplankton), respectively. B_r quantifies the bottom- or top-heaviness of the biomass pyramid. In particular, if $B_r > 1$, the pyramid is inverted.

The experiments were designed to observe long-term responses to bottom-up (i.e. nutrient enrichment) and top-down (i.e. fish addition) perturbations. Therefore, we considered the biomasses reported in the various treatments as stationary. To remove the dependency on (unknown) perturbation intensity, we further based our analysis on the expectation of a linear response of biomasses to the two treatments (perturbation intensity is removed by only considering response ratios).

For each level $i = 1, 2$, we thus estimated the bottom-up change in biomass $\Delta B_{i\uparrow}$ due to nutrient enrichment, and the top-down effect $\Delta B_{i\downarrow}$ due to fish addition. We could measure these changes directly by comparing the perturbed biomasses with the unperturbed value B_i^*. But since there was also a cross-treatment with both enrichment and fish, we instead fitted a multilinear model

$$B_i = \tilde{B}_i + x_\uparrow \Delta B_{i\uparrow} + x_\downarrow \Delta B_{i\downarrow}; \quad x_\uparrow = 0, 1; \ x_\downarrow = 0, 1. \tag{2}$$

where the binary treatment variable $x_\uparrow = 0$ or 1 represents the absence or presence of nutrient enrichment, and $x_\downarrow = 0$ or 1 represents the absence or presence of fish. We thus identified the coefficients $\Delta B_{i\uparrow}$ and $\Delta B_{i\downarrow}$ as the slopes of the multilinear response, with an intercept $\tilde{B}_i$.

The goodness-of-fit informed us of whether these estimates were robust: a large fitting error indicates significant interaction between the two treatments. For instance, nutrient enrichment is realized both without and with fish addition. The associated error on response $\Delta B_{i\uparrow}$ is large when the two measurements are incompatible (Figure 3b and c). This can hint either to measurement error, non-stationary behaviour, or strong nonlinearities, which we do not consider here.

We therefore selected the fitted coefficients estimated with low errors (i.e., estimated error < coefficient value, see Figure 3b and c). From selected coefficients, we deduced upward and downward response ratios, defined respectively as

$$A_\uparrow = \frac{\Delta B_{2\uparrow}}{\Delta B_{1\uparrow}}, \text{ and } A_\downarrow = -\frac{\Delta B_{1\downarrow}}{\Delta B_{2\downarrow}} \tag{3}$$

$A_\uparrow$ measures the biomass change of level 2 relative to that of level 1 following nutrient enrichment, while $A_\downarrow$ represents the biomass change of level 1 relative to that of level 2 following fish addition. We used response ratios as measures of response to perturbations to eliminate potential differences in the magnitude of the perturbations performed across systems.

In order to focus on *dynamically consistent chains*, in addition to the filtering for coefficients estimated with low errors, we restricted ourselves to cases in which these the two trophic levels respond in the same direction to nutrient enrichment, and in opposite direction to fish addition. Discarded response ratios can be interpreted as indicating other dynamical behaviors (such as competition) that are not consistent with a simple food chain. We analysed whether using both filters (i.e. low error and expected response direction) affects the distribution of responses using Kolmogorov-Smirnov tests, which indicated that the filtering used did not bias the distribution of the responses considered (Appendix S1).

Finally, to analyse the across-system statistical relationship between biomass distribution and trophic cascades, we performed linear mixed effects models including B_r as fixed effects, study system as random effects (given that we decomposed each study system in multiple chains) and $A_\uparrow$ and $A_\downarrow$, as response variables. For bottom-up response ($A_\uparrow$) we found that the variance explained by the study system was 0.19, therefore, we included study system as a random effect. However, we found no significant differences between the model including the study system as random effect (AIC=57.9) and the model including only B_r as fixed effect (AIC=57.0). For $A_\downarrow$ the variance associated to the study system was 0.002, and the model including only B_r as a fixed effect was significantly better (AIC=58.3; linear mixed model with study system as random effect AIC=60.3). Thus, the results presented below correspond to the analyses were only B_r was included as a fixed effect in the model.

2.2 Lotka-Volterra model

To analyse the within-system dynamical relationship between biomass distribution and response to perturbations and the mechanisms driving it, we focus on a simple two-level system, and assume Lotka-Volterra interactions (i.e. type-I functional responses). However, the theoretical framework presented has no restrictions on the length of the food chain, and can be extended to different functional responses (see Appendix S2 for multi-level and nonlinear chain results). Here, long-term responses to perturbation are shifts in equilibrium biomasses. Although empirical systems are never perfectly stationary, equilibrium analysis of such dynamical models have proven successful in quantifying species interactions and responses even in data with significant temporal variance (Maynard *et al.*, 2019; Barbier *et al.*, 2020).

Let B_i, $i = 1, 2$ be the biomass density of primary producers and consumers, respectively.

$$\begin{cases} \frac{1}{B_1}\frac{dB_1}{dt} = & g - D_1 B_1 - \alpha B_2 \\ \frac{1}{B_2}\frac{dB_2}{dt} = & \varepsilon\alpha B_1 - D_2 B_2 - r \end{cases} \tag{4}$$

We denote by $g > 0$ the intrinsic growth rate of primary producers, and $r > 0$ the intrinsic death rate of consumers. The per-capita attack rate of consumers is denoted as $\alpha > 0$ (a rate per unit of biomass density). In the same units we denote by D_i $i = 1, 2$ the density-dependent mortality (or reduction of biomass productivity) of both trophic levels (i.e., self-regulation). Finally, we write $0 < \varepsilon < 1$ the efficiency of productivity transfer from primary producers to consumers. There are

two units: time and biomass density. Here we are interested only in non dimensional ratios, such as biomass ratio and response ratios. We therefore nondimensionalized the model (See Appendix S2 for details), identifying important ratios: relative attack rate $a = \frac{\alpha}{D_1}$, relative self-regulation $\psi = \frac{D_2}{D_1}$; and relative intrinsic rate $\phi = \frac{r}{g}$.

2.3   Approximate Bayesian Computation (ABC)

We can identify parameter ranges consistent with the data using Approximate Bayesian Computation (Beaumont, 2010). The idea of ABC is to leverage empirical patterns to constrain the distribution of simulation parameters. In this way, we can use across-system patterns to infer within-system dynamics. We generated predator-prey systems with different parameter values drawn from broad prior distributions, and computed their equilibrium biomass ratios B_r and top-down and bottom-up response ratios $A_\uparrow$ and $A_\downarrow$. We then retained only model realizations for which (i) the two species coexisted and (ii) the relationship between B_r and $A_\uparrow$ and between B_r and $A_\downarrow$ were consistent with those observed in our data analyses (Figure S3.1). We retained 1500 model realizations, and used the corresponding parameter values to compute a joint posterior distribution for all model parameters. All parameters were drawn uniformly on a log scale, over a span of several orders of magnitude: biomass conversion efficiency $\varepsilon \in [10^{-2}, 10^{-1}]$; relative attack rate $a = \frac{\alpha}{D_1} \in [10^1, 10^4]$; relative self-regulation $\psi = \frac{D_2}{D_1} \in [10^{-5}, 10^5]$; and relative intrinsic rate $\phi = \frac{r}{g} \in [10^{-4}, 10^4]$. Posterior distributions of parameters are presented on log scales and compared to their prior distributions, constrained on coexistence.

3   Results

3.1   Within-system dynamical relationship: theoretical predictions

To determine whether the biomass distribution within one system contains information about its dynamical features, we develop a theoretical framework that spans the various dynamical regimes previously studied in food chains (e.g. resource and consumer control). We compute the predicted quantitative relationship between the equilibrium biomass ratio and the response ratios $A_\uparrow$ and $A_\downarrow$, allowing us to understand deviations from the proportional response expectation H0.

Solving for the equilibrium of a two-species Lotka-Volterra system (see Methods and Appendix S2), we find that the biomass ratio $B_r = B_2/B_1$ can be expressed as

$$B_r = \left(\frac{\varepsilon\alpha}{D_2}\right)\frac{1-\chi}{1+\xi}; \quad \chi = \frac{D_1}{\varepsilon\alpha}\frac{r}{g}; \quad \xi = \frac{\alpha}{D_2}\frac{r}{g} \tag{5}$$

In this expression, we have isolated two terms, χ and ξ, that express the relative effect of consumer mortality r and primary productivity g on biomasses. Similar terms appear in more complex models (e.g. longer food chains and nonlinear dynamics, Appendix S2). This hints at a general notion of *trophic dissipation* (Box 1), i.e. the ratio between effects of mortality and primary productivity on standing biomass at any given trophic level.

Here, we specifically define the impact of trophic dissipation either on consumers (χ) or on resources (ξ). Indeed, χ is the ratio of predator biomass losses (at equilibrium) due to mortality r, over gains from consumption of primary productivity g. Thus, we call χ *dissipative loss*, as it represents a decrease from the maximum predator biomass allowed by primary productivity. On the other hand, ξ is the ratio of biomass gains of the prey due to predator mortality, over the contribution to standing biomass of productivity. We call ξ *dissipative release*, at it reflects the relative importance of prey release caused by consumer mortality. Both notions – which are not independent from one another– relate to what we may call trophic dissipation; seen from either the consumer's (χ) or the producer's (ξ) perspective. Importantly, because we are considering equilibrium biomasses, trophic dissipation accounts for long-term effects, and not only direct interaction strength.

We now explain how static properties – B_r, the prefactor $\varepsilon\alpha/D_2$ in Eq. (5), χ and ξ – can be associated with dynamical responses to bottom-up and top-down perturbations. To see this, we

apply a perturbation Δg of primary productivity, such as a nutrient enrichment treatment. The changes in biomass of trophic level 1 and trophic level 2 are given by, respectively

$$\Delta B_{1\uparrow} = \frac{\Delta g}{b} D_2, \text{ and } \Delta B_{2\uparrow} = \frac{\Delta g}{b} \varepsilon\alpha \tag{6}$$

where $b = D_1 D_2 + \varepsilon\alpha^2$ (but b will play no role in what follows). Similarly, changes in biomass due to a change in consumer mortality Δr (e.g. caused by fish addition), are

$$\Delta B_{1\downarrow} = \frac{\Delta r}{b}\alpha \text{ and } \Delta B_{2\downarrow} = -\frac{\Delta r}{b} D_1. \tag{7}$$

Thus, defining response ratios as in the empirical analysis Eq. (3), bottom-up and top-down responses will read, respectively

$$A_\uparrow = \frac{\varepsilon\alpha}{D_2} \text{ and } A_\downarrow = \frac{\alpha}{D_1}. \tag{8}$$

We recognize in $A_\uparrow$ the prefactor of the right hand side of Eq. (5). Furthermore, by comparing (8) with the expressions of ξ and χ in (5) we can deduce a synthetic relationship between dissipative loss, dissipative release and response ratios

$$\xi = \chi A_\uparrow A_\downarrow \tag{9}$$

This expression is not just an algebraic manipulation. It reveals the dynamical mechanism that allows us to move from the predator to the prey's perspective of trophic dissipation. This mechanism is represented by the product $A_\uparrow A_\downarrow$ of upwards and downwards growth propagation, which measures the strength of the dynamical feedback loop in the food chain (denoted λ in Barbier & Loreau, 2019).

With Eqs. (5, 9) we now have all the ingredients to propose a formal relationship between biomass ratio, response ratio and trophic dissipation (see Appendix S2 for details and generalizations)

$$B_r = \frac{A_\uparrow(1 - \chi)}{1 + \chi A_\uparrow A_\downarrow} \tag{10}$$

This expression makes it clear that three different factors determine the predator-prey biomass ratio: the two dynamical response ratios $A_\uparrow$ and $A_\downarrow$, and the additional effects of growth and mortality that enter in the dissipative loss χ. The value of dissipative release ξ (9), then determines three main qualitative regimes of Eq. (10), as shown in Figure 2 and 4. At the extremes, only upward or downward cascades are related to biomasses:

$$B_r \approx A_\uparrow(1 - \chi) \qquad\qquad \text{if} \quad \xi \ll 1,$$
$$B_r \approx \frac{1 - \chi}{\chi A_\downarrow} \qquad\qquad \text{if} \quad \xi \gg 1. \tag{11}$$

For intermediate values $\xi \sim 1$, B_r reflects an entangled contribution of upward and downward responses.

The two parameters linked to trophic dissipation are complementary: ξ determines the qualitative regime in which the system is in, while χ controls the slope of the relationship within each regime. If $\xi \ll 1$, then $A_\uparrow \approx B_r/(1 - \chi)$, the biomass distribution is entirely shaped by upward effects. Upward perturbations either follow proportional response H0 if $\chi \approx 0$, as in (Barbier & Loreau, 2019), or are amplified. On the other hand, the biomass ratio contains no a priori information about downward responses $A_\downarrow$ (Figure 4a). In the opposite limit of large ξ, $A_\downarrow$ will be inversely proportional to B_r, but with a prefactor $(1 - \chi)/\chi$ that can lead to downward cascade amplification if $\chi < 1/2$ and attenuation if $\chi > 1/2$ (Figure 4b). Proportional response is thus unlikely for top-down effects, as it requires exactly $\chi = \frac{1}{2}$ If consumer self-regulation is very weak or zero, we find $A_\uparrow, \xi \to \infty$ and only the top-down driven regime is observed, as in classical models (e.g. Oksanen et al., 1981; Rossberg et al., 2019) that have been used to showcase counter-intuitive dynamical effects in food webs.

3.2 Across-systems statistical relationship: empirical results

To determine whether across-system statistical relationships exist, we analyzed 120 potential chains, defined by all the possible paths of energy flux across trophic levels in the experiments

considered (i.e. Z-A (31), C1-A2 (17), C1-H1 (22), H1-A1 (21), H2-A2 (14), H2-A1 (14)). We found 68 responses to perturbations that were consistent with that of a dynamical food chain. That is, 33 of the considered paths showed a chain-like behaviour of similar response direction between predator and prey biomass for upward cascades, i.e. $A_\uparrow > 0$, while 35 chains showed opposite response direction between predator and prey biomass for downwards cascades, i.e. $A_\downarrow > 0$ (Table S1.2 in Appendix S1).

Paths where all zooplankton species and all phytoplankton species were grouped together to form two trophic levels (chain Z-A in Figure 3a) showed the highest percentage of chain-like responses across all experiments, together with the chain composed by small herbivores (H1) consuming highly edible phytoplankton (A1). The estimated responses to perturbations had low fit significance or wrong signs in approximately 60% of the instances analysed (61% for $\Delta B_{1\uparrow}$, 68% for $\Delta B_{2\uparrow}$, 58% for $\Delta B_{1\downarrow}$ and 60% for $\Delta B_{2\downarrow}$)(Figure S1.1 and S1.2). For instance, we observed zooplankton biomass increasing after fish addition, or phytoplankton biomass decreasing due to nutrient enrichment.

The biomass distribution was bottom-heavy in most selected paths (Figure 5)(i.e., $B_r < 1$). In the selected experimental systems, B_r was positively correlated with the upwards response ratio ($A_\uparrow$). In log scale, the slope of this relationship was close to one with an intercept close to zero, indicating that the responses to nutrient addition of each trophic level were proportional to their biomass, showing no further effect of trophic interactions in amplifying the upwards response.

In contrast, B_r was negatively correlated with downwards response ratio $A_\downarrow$ with a slope of -0.5 in log scale (Figure 5b). Yet, as we discuss below, this trend may be spurious, if top-down effects are not causally involved in shaping the biomass ratios.

3.3 Comparing theory and data

We can use our theoretical results to understand the empirical patterns and underlying mechanisms. In our ABC approach, we simulate systems in which the theoretical within-system relationship holds exactly, and infer how the variation of key ecological parameters can explain the empirical relationship across systems (Figure 6). We note that, although we did not constrain the model parameters to match the empirical biomass ratios (but only to follow the relationship of biomass and response ratios), the selected simulated data points displayed a remarkably similar distribution of biomass ratios (Figure S3.1).

Once filtered to match empirical patterns, the distribution of dissipative release ξ (i.e. prey release due to dissipation) shifted towards small values (c. 67% had $\xi < 0.1$ cf. Figure 6a). This suggests that for two thirds of systems (c. 67%), biomass ratios were shaped by bottom-up effects. The empirical trend between biomass ratio and top-down effects was allowed in our model and simulations, yet mainly driven by a weak anticorrelation between upward and downward response (Figure S3.1). Grouping simulations by dissipative release (low, intermediate and high), we found that the three classes presented qualitatively similar patterns with top-down responses (Figure S3.2), with a slight steepening of the slope for systems with strongest dissipative release.

In terms of model parameters, dissipative release reads $r/g \times \alpha/D_2$. The distribution of $\phi = r/g$ remained broad after filtering by the empirical patterns (Figure 6b). On the other hand, the ratio of attack rate and consumer self-regulation α/D_2 was more constrained by the empirical patterns: its posterior distribution shows a unimodal distribution centered around 1 (Figure 6c). Thus, the ratio r/g may be driving the across-system variation in dissipative release.

4 Discussion

Trophic cascades are widespread in nature (Pace *et al.*, 1999; Shurin *et al.*, 2002; Estes *et al.*, 2011). Changes in the abundance of a given trophic level can have further consequences along the food chain through both direct and indirect interactions, ultimately altering ecosystem structure and functioning (Paine, 1969; Carpenter *et al.*, 1985; Polis *et al.*, 2000; Duffy, 2002; Estes *et al.*, 2011). Understanding their prevalence and strength has occupied a large part of the food chain literature

(Shurin & Seabloom, 2005; Borer *et al.*, 2005; Heath *et al.*, 2014; Barbier & Loreau, 2019; Sentis *et al.*, 2020). Multiple sources of empirical data have been used as indices of trophic cascades in natural systems (Baum & Worm, 2009; Heath *et al.*, 2014). Most of these indices are based on observations of dynamical behaviour, such as the correlations between time series of interacting pairs of trophic levels (Frank *et al.*, 2007), or the response of a given trophic level to a perturbation applied elsewhere in the ecosystem (Baum & Worm, 2009; Estes *et al.*, 2011; Hulot *et al.*, 2014).

Another salient feature of food chains is the static distribution of biomass across trophic levels. It has long been studied in the form of pyramids of biomass (Elton, 1927), or as a size spectrum when, instead of discrete trophic levels, communities are ordered by organism body size (Sheldon *et al.*, 1972; Cousins, 1985; Blanchard *et al.*, 2017).

Since biomass distributions are more easily measured than dynamical properties, previous studies have discussed whether these static distributions can hint at the underlying dynamics (McCauley *et al.*, 2018; Barbier & Loreau, 2019). For instance, Rossberg *et al.* (2019) related size *domes* (i.e., high biomass density in some body-size range and depleted troughs in between) in pelagic systems to the existence of downwards trophic cascades. But a systematic and quantitative exploration of possible relations between downward and upward cascade strength and biomass distributions was still lacking.

4.1 Summary of results

We investigated the relationship between biomass structure and dynamics in food chains, using both theory and empirical data from pelagic experiments, asking under which conditions the strength of trophic cascades can have a signature in the biomass distribution across trophic levels (Figure 5).

Our framework shows that the relationship between biomass distribution and trophic cascades is determined by a combination of intrinsic rates (i.e. growth and mortality), self-regulation and attack rate. This combination can be seen from two perspectives: the consumer's or the resource's (Box 1). We define *dissipative loss* χ as the consumer's ratio of biomass losses from mortality over gains from primary productivity. Conversely, *dissipative release* ξ is the resource's ratio of biomass gains from consumer mortality over gains from primary productivity. We find that the two are connected via $\xi/\chi = A_\uparrow A_\downarrow$ the product of bottom-up and top-down responses, previously identified by Barbier & Loreau (2019) as the strength λ of predator-prey feedbacks. Thus, ξ and χ are not independent parameters, but represent a single axis of trophic dissipation in our synthetic framework.

These two notions of dissipative release ξ (resource perspective) and loss χ (consumer perspective) play complementary roles in our results: ξ determines three qualitative scenarios, while χ controls the slope of the relationship. Low dissipative release, $\xi \ll 1$, corresponds to the scenario where standing biomasses only reflect bottom-up effects. Conversely, $\xi \gg 1$ indicates that only top-down effects come into play at equilibrium. For intermediate values, $\xi \sim 1$, bottom-up and top-down effects become entangled, so that the biomass ratio does not provide precise information about either, but rather a combination of the two. By construction, many classic food chain models such as exploitation ecosystems (Oksanen *et al.*, 1981) are in the top-down limit $\xi \to \infty$. In every case, the slope of the relationship (i.e. whether it is amplified or attenuated compared to simple proportionality) depends on dissipative loss χ.

The crucial distinction between asking whether cascades are strong and whether they are causally reflected in the biomass distribution, is clearly demonstrated here: top-down cascades only shape the standing biomass distribution if natural predator losses are sufficiently high ($\xi > 1$), but the larger these losses, the weaker the cascades, since the predator population is more depleted, with less potential for further prey release from perturbations.

The existence of a pattern across multiple systems is contingent on low variation of dissipative losses χ, while the pattern's shape depends on χ's magnitude. Using Approximate Bayesian Computation (ABC), we estimated the distribution of dynamical parameters, including dissipative release, that was compatible with the observed empirical pattern. We found that experimental data was largely compatible with the bottom-up (67%) or entangled (25%) scenarios, with only 8% of systems for which top-down effects should drive biomass ratios. Nonetheless, we did observe

a trend between biomass ratios and top-down effects. This relationship mainly arose, in our ABC approach, from latent parameters that create an anticorrelation between $A_\downarrow$ and $A_\uparrow$, and subsequently B_r (Appendix S3). Only a small contribution to the trend came from the fraction ($25\% + 8\%$) of systems where top-down effects play direct, causal role. Even if we observed a similar relationship between downward responses and biomass ratios in all three groups of systems, our theory proposes that those patterns should be interpreted with differently; from a spurious correlation for low dissipation release to an actual causation for high dissipation release.

Beyond hinting at which empirical patterns are causal, our framework may be used to shed light on specific ecological parameters. The ABC analysis suggested that intra- and inter-level interactions are of roughly comparable strength in these pelagic experiments, while the balance of primary productivity and consumer mortality may be the most important axis of variation between experiments.

4.2   Limitations

Our theoretical predictions held on data pooled from multiple experimental pelagic communities. Taking systems individually, however, only 50% of the energy pathways displayed a consistent chain-like response to either bottom-up and top-down perturbations, and very few had both. This is unfortunate, as using information from both types of perturbations would have allowed us to infer relevant ecological parameters (e.g. self-regulation) in specific communities, rather than only trends across systems.

Different mechanisms could explain this high rate of discrepancies, among which omnivory is a prominent one. The presence of omnivorous links in food webs has been well documented (Polis *et al.*, 1989; Polis, 1994; Emmerson & Yearsley, 2004; Neutel *et al.*, 2007), and is known to obviate cascading effects (Diehl, 1993; Pringle & Hamazaki, 1998; Polis, 1999; McCauley *et al.*, 2018). Indeed, many 'herbivorous' zooplankton species consume phytoplankton but also a substantial amount of microzooplankton, such as rotifers and ciliates (Sprules & Bowerman, 1988; Gilbert, 1988; Brett *et al.*, 1994; Hansson *et al.*, 2004), which can explain some of the non-chain-like responses we observed. Similarly, the existence of compensation between species or functional groups within one trophic level (Gonzalez & Loreau, 2009) could also attenuate or reverse the expected dynamical response (Leibold, 1989; Hunter & Price, 1992; Strong, 1992; McCann *et al.*, 1998).

Additionally, we may be unable to correctly identify chain-like dynamics when the response of either level to a perturbation is very small, and indistinguishable from measurement noise. Taking a ratio of responses is necessary to eliminate the confounding role of perturbation magnitude, but this ratio becomes poorly defined when either response is close to zero.

We obtained the largest fraction of chain-like responses when considering all zooplankton species as a single trophic level and all phytoplankton species as another trophic level, suggesting that consistent chain dynamics may emerge as a complexity reduction of communities into few interacting levels composed by multiple species or groups of species (Ulanowicz, 1995).

4.3   Influence of primary productivity

A long-standing empirical and theoretical question, is how primary productivity can affect either biomass distribution (McNaughton *et al.*, 1989; Del Giorgio *et al.*, 1999) or trophic cascade strength (Carpenter *et al.*, 1985). Here, we showed how primary productivity can change the *relationship* between biomasses and cascades. Increasing primary productivity lowers the strength of trophic dissipation, leading to biomass pyramids shaped by bottom-up propagation of primary productivity that convey, a priori, no information about top-down cascades. Interestingly, our ABC approach suggests that the variation in dissipative release across systems was mainly driven by a variation in the ratio of intrinsic rates (consumer mortality over prey productivity) among systems. Thus, variations in primary productivity could be an essential factor explaining the diversity of observed pyramids and dynamical responses in the pelagic systems analysed.

4.4   On the importance of self-regulation

Previous research has already pointed at the importance of self-regulatory processes to modulate the strength of bottom-up and top-down effects (McCann *et al.*, 1998; Herendeen, 2004; Heath *et al.*, 2014). We provide here a precise understanding of how both resource and consumer self-regulation shape the link between static and dynamic properties in a food chain, as they determine the importance of trophic dissipation in the observed biomasses of consumers and resources.

Very few empirical studies, however, provide reliable direct estimates of consumer self-regulation (Skalski & Gilliam, 2001). Furthermore, effects from species or compartments outside of the considered chain could participate in the observed self-regulation (Loreau, 2010). This would make it an emergent property that is not directly accessible from individual behaviour such as predator interference, but could still be estimated indirectly through our framework.

Conclusion

We asked whether a food chain's biomass distribution could predict the strength of its trophic cascades. We find that a key ecological mechanism determines if the strength of upward or downward cascades is directly involved in shaping the static biomass distribution. This mechanism, that we call trophic dissipation, can be defined for each trophic level as the relative contribution of consumer mortality versus primary productivity in the standing biomass. It depends on well-studied ecological parameters, such as attack rates, but also on the elusive notion of prey and predator self-regulation.

An observed relationship between biomass distribution and upward or downward cascades can reflect this direct causal link, but could also arise from correlations driven by a common factor. In the case of direct causality, we can extrapolate this relationship to make new predictions, within or across ecosystems, without additional knowledge of confounding factors. Furthermore, the common practice of assessing cascade strength in relative terms (as amplification or attenuation compared to proportional response) is only meaningful for the type of cascades that were causally involved in shaping the standing biomasses, and misleading otherwise. Our results provide a canvas for using measurements of static properties as reliable indicators to predict a food chain's dynamical response to perturbations.

Acknowledgments

We thank Bart Haegeman for helpful discussions and review of previous versions of the manuscript. We also thank Florence Hulot for sharing experimental data. This work was supported by the TULIP Laboratory of Excellence (ANR-10-LABX-41) and by the BIOSTASES Advanced Grant, funded by the European Research Council under the European Union's Horizon 2020 research and innovation program (666971). JFA was supported by an Irish Research Council Laureate Awards 2017/2018.

Author contributions

All authors contributed to the design of the study. NG and AA assembled the empirical data and, jointly with MB, performed the empirical analyses. JFA and MB developed the theoretical framework. NG and JFA performed the numerical simulations. NG, MB and JFA wrote the first draft of the manuscript, and all the authors contributed substantially to revisions.

References

Barbier, M. & Loreau, M. (2019). Pyramids and cascades: a synthesis of food chain functioning and stability. *Ecology letters*, 22, 405–419.

Barbier, M., de Mazancourt, C., Loreau, M. & Bunin, G. (2020). Fingerprints of high-dimensional coexistence in complex ecosystems. *bioRxiv*, p. 652230.

Baum, J.K. & Worm, B. (2009). Cascading top-down effects of changing oceanic predator abundances. *Journal of animal ecology*, 78, 699–714.

Beaumont, M.A. (2010). Approximate bayesian computation in evolution and ecology. *Annual review of ecology, evolution, and systematics*, 41, 379–406.

Blanchard, J.L., Heneghan, R.F., Everett, J.D., Trebilco, R. & Richardson, A.J. (2017). From bacteria to whales: using functional size spectra to model marine ecosystems. *Trends in ecology & evolution*, 32, 174–186.

Borer, E.T., Seabloom, E.W., Shurin, J.B., Anderson, K.E., Blanchette, C.A., Broitman, B., Cooper, S.D. & Halpern, B.S. (2005). What determines the strength of a trophic cascade? *Ecology*, 86, 528–537.

Brett, M.T., Wiackowski, K., Lubnow, F.S., Mueller-Solger, A., Elser, J.J. & Goldman, C.R. (1994). Species-dependent effects of zooplankton on planktonic ecosystem processes in Castle Lake, California. *Ecology*, 75, 2243–2254.

Carpenter, S.R., Kitchell, J.F. & Hodgson, J.R. (1985). Cascading trophic interactions and lake productivity. *BioScience*, 35, 634–639.

Cousins, S. (1985). Ecologists build pyramids again. *New Scientist*, 106, 50–54.

Cury, P., Shannon, L., Roux, J., Daskalov, G., Jarre, A., Moloney, C. & Pauly, D. (2005). Trophodynamic indicators for an ecosystem approach to fisheries. *ICES Journal of Marine Science*, 62, 430–442.

Del Giorgio, P.A., Cole, J.J., Caraco, N.F. & Peters, R.H. (1999). Linking planktonic biomass and metabolism to net gas fluxes in northern temperate lakes. *Ecology*, 80, 1422–1431.

Diehl, S. (1993). Relative consumer sizes and the strengths of direct and indirect interactions in omnivorous feeding relationships. *Oikos*, pp. 151–157.

Duffy, J.E. (2002). Biodiversity and ecosystem function: the consumer connection. *Oikos*, 99, 201–219.

Elton, C.S. (1927). The animal community. *Animal ecology*, pp. 239–256.

Emmerson, M. & Yearsley, J.M. (2004). Weak interactions, omnivory and emergent food-web properties. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 271, 397–405.

Estes, J.a., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., Carpenter, S.R., Essington, T.E., Holt, R.D., Jackson, J.B.C., Marquis, R.J., Oksanen, L., Oksanen, T., Paine, R.T., Pikitch, E.K., Ripple, W.J., Sandin, S.a., Scheffer, M., Schoener, T.W., Shurin, J.B., Sinclair, A.R.E., Soulé, M.E., Virtanen, R. & Wardle, D.a. (2011). Trophic downgrading of planet Earth. *Science (New York, N.Y.)*, 333, 301–6.

Frank, K.T., Petrie, B. & Shackell, N.L. (2007). The ups and downs of trophic control in continental shelf ecosystems. *Trends in Ecology & Evolution*, 22, 236–242.

Frank, K.T., Petrie, B., Shackell, N.L. & Choi, J.S. (2006). Reconciling differences in trophic control in mid-latitude marine ecosystems. *Ecology Letters*, 9, 1096–1105.

Gilbert, J.J. (1988). Suppression of rotifer populations by Daphnia: a review of the evidence, the mechanisms, and the effects on zooplankton community structure 1. *Limnology and Oceanography*, 33, 1286–1303.

Gonzalez, A. & Loreau, M. (2009). The causes and consequences of compensatory dynamics in ecological communities. *Annu. Rev. Ecol. Evol. Syst.*, 40, 393–414.

Gruner, D.S., Smith, J.E., Seabloom, E.W., Sandin, S.A., Ngai, J.T., Hillebrand, H., Harpole, W.S., Elser, J.J., Cleland, E.E. & Bracken, M.E.S. (2008). A cross-system synthesis of consumer and nutrient resource control on producer biomass. *Ecology letters*, 11, 740–755.

Hairston, N.G., Smith, F.E. & Slobodkin, L.B. (1960). Community structure, population control, and competition. *The american naturalist*, 94, 421–425.

Halaj, J. & Wise, D.H. (2001). Terrestrial trophic cascades: how much do they trickle? *The American Naturalist*, 157, 262–281.

Hansson, L., Gyllström, M., Ståhl-Delbanco, A. & Svensson, M. (2004). Responses to fish predation and nutrients by plankton at different levels of taxonomic resolution. *Freshwater Biology*, 49, 1538–1550.

Heath, M.R., Speirs, D.C. & Steele, J.H. (2014). Understanding patterns and processes in models of trophic cascades. *Ecology letters*, 17, 101–114.

Hedges, L.V., Gurevitch, J. & Curtis, P.S. (1999). The meta-analysis of response ratios in experimental ecology. *Ecology*, 80, 1150–1156.

Herendeen, R.A. (2004). Bottom-up and top-down effects in food chains depend on functional dependence: an explicit framework. *Ecological Modelling*, 171, 21–33.

Hulot, F.D., Lacroix, G. & Loreau, M. (2014). Differential responses of size-based functional groups to bottom–up and top–down perturbations in pelagic food webs: a meta-analysis. *Oikos*, 123, 1291–1300.

Hunter, M.D. & Price, P.W. (1992). Playing chutes and ladders: heterogeneity and the relative roles of bottom-up and top-down forces in natural communities. *Ecology*, 73, 724–732.

Jonsson, T. (2017). Conditions for Eltonian pyramids in Lotka-Volterra food chains. *Scientific reports*, 7, 1–10.

Leibold, M.A. (1989). Resource edibility and the effects of predators and productivity on the outcome of trophic interactions. *The American Naturalist*, 134, 922–949.

Leibold, M.A. (1996). A graphical model of keystone predators in food webs: trophic regulation of abundance, incidence, and diversity patterns in communities. *The American Naturalist*, 147, 784–812.

Lindeman, R.L. (1942). The trophic-dynamic aspect of ecology. *Ecology*, 23, 399–417.

Loreau, M. (2010). *From populations to ecosystems: Theoretical foundations for a new ecological synthesis (MPB-46)*. vol. 46. Princeton University Press.

Maynard, D.S., Wootton, J.T., Serván, C.A. & Allesina, S. (2019). Reconciling empirical interactions and species coexistence. *Ecology letters*, 22, 1028–1037.

McCann, K.S., Hastings, A. & Strong, D.R. (1998). Trophic cascades and trophic trickles in pelagic food webs. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 265, 205–209.

McCauley, D.J., Gellner, G., Martinez, N.D., Williams, R.J., Sandin, S.A., Micheli, F., Mumby, P.J. & McCann, K.S. (2018). On the prevalence and dynamics of inverted trophic pyramids and otherwise top-heavy communities. *Ecology Letters*, 21, 439–454.

McNaughton, S.J., Oesterheld, M., Frank, D.A. & Williams, K. (1989). Ecosystem-level patterns of primary productivity and herbivory in terrestrial habitats. *Nature*, 341, 142–144.

Neutel, A.M., Heesterbeek, J.a.P., van de Koppel, J., Hoenderboom, G., Vos, A., Kaldeway, C., Berendse, F. & de Ruiter, P.C. (2007). Reconciling complexity with stability in naturally assembling food webs. *Nature*, 449, 599–602.

Oksanen, L., Fretwell, S.D., Arruda, J. & Niemela, P. (1981). Exploitation ecosystems in gradients of primary productivity. *The American Naturalist*, 118, 240–261.

Pace, M.L., Cole, J.J., Carpenter, S.R. & Kitchell, J.F. (1999). Trophic cascades revealed in diverse ecosystems. *Trends in ecology & evolution*, 14, 483–488.

Paine, R.T. (1969). A note on trophic complexity and community stability. *The American Naturalist*, 103, 91–93.

Polis, G.A. (1994). Food webs, trophic cascades and community structure. *Australian Journal of Ecology*, 19, 121–136.

Polis, G.A. (1999). Why are parts of the world green? Multiple factors control productivity and the distribution of biomass. *Oikos*, pp. 3–15.

Polis, G.A., Myers, C.A. & Holt, R.D. (1989). The ecology and evolution of intraguild predation: potential competitors that eat each other. *Annual review of ecology and systematics*, 20, 297–330.

Polis, G.A., Sears, A.L.W., Huxel, G.R., Strong, D.R. & Maron, J. (2000). When is a trophic cascade a trophic cascade? *Trends in Ecology & Evolution*, 15, 473–475.

Pringle, C.M. & Hamazaki, T. (1998). The role of omnivory in a neotropical stream: separating diurnal and nocturnal effects. *Ecology*, 79, 269–280.

Ripple, W.J., Estes, J.A., Schmitz, O.J., Constant, V., Kaylor, M.J., Lenz, A., Motley, J.L., Self, K.E., Taylor, D.S. & Wolf, C. (2016). What is a trophic cascade? *Trends in Ecology & Evolution*, 31, 842–849.

Rossberg, A.G., Gaedke, U. & Kratina, P. (2019). Dome patterns in pelagic size spectra reveal strong trophic cascades. *Nature communications*, 10, 1–11.

Sentis, A., Bertram, R., Dardenne, N., Simon, J.C., Magro, A., Pujol, B., Danchin, E. & Hemptinne, J.L. (2020). Intraspecific difference among herbivore lineages and their host-plant specialization drive the strength of trophic cascades. *bioRxiv*, p. 722140.

Sheldon, R.W., Prakash, A. & Sutcliffe Jr, W. (1972). The size distribution of particles in the ocean. *Limnology and oceanography*, 17, 327–340.

Shin, Y.J., Rochet, M.J., Jennings, S., Field, J.G. & Gislason, H. (2005). Using size-based indicators to evaluate the ecosystem effects of fishing. *ICES Journal of marine Science*, 62, 384–396.

Shurin, J.B., Borer, E.T., Seabloom, E.W., Anderson, K., Blanchette, C.A., Broitman, B., Cooper, S.D. & Halpern, B.S. (2002). A cross-ecosystem comparison of the strength of trophic cascades. *Ecology letters*, 5, 785–791.

Shurin, J.B. & Seabloom, E.W. (2005). The strength of trophic cascades across ecosystems: predictions from allometry and energetics. *Journal of animal Ecology*, 74, 1029–1038.

Skalski, G.T. & Gilliam, J.F. (2001). Functional responses with predator interference: viable alternatives to the Holling type II model. *Ecology*, 82, 3083–3092.

Sprules, W.G. & Bowerman, J.E. (1988). Omnivory and food chain length in zooplankton food webs. *Ecology*, 69, 418–426.

Strong, D.R. (1992). Are trophic cascades all wet? Differentiation and donor-control in speciose ecosystems. *Ecology*, 73, 747–754.

Trebilco, R., Dulvy, N.K., Anderson, S.C. & Salomon, A.K. (2016). The paradox of inverted biomass pyramids in kelp forest fish communities. *Proceedings of the Royal Society B: Biological Sciences*, 283, 20160816.

Ulanowicz, R.E. (1995). Ecosystem trophic foundations: Lindeman exonerata. *Complex Ecology: the part-whole relation in ecosystems. Prentice Hall, Englewood Cliffs, NJ*, pp. 549–550.

Woodson, C.B., Schramski, J.R. & Joye, S.B. (2018). A unifying theory for top-heavy ecosystem structure in the ocean. *Nature communications*, 9, 1–8.

Box 1. Definition of main concepts.

- **Upward amplification/attenuation:** stronger/weaker response of higher trophic levels relative to lower trophic levels after a perturbation at the bottom of the chain (e.g. nutrient enrichment).

- **Downward amplification/attenuation:** stronger/weaker response of lower trophic levels relative to higher trophic levels after a perturbation at the top of the chain (e.g. fish addition).

- **Self-regulation:** regulatory mechanisms that cause per capita growth rates to depend on the abundance of the species in question (e.g. intraspecific interference, cannibalism or effects from hidden compartments such as pathogens).

- **Trophic dissipation:** general concept representing, at any trophic level, the relative importance of consumer mortality versus primary productivity (i.e. energy dissipation versus basal influx) in shaping the standing biomass. This concept can be defined formally for a given trophic level, becoming dissipative loss for the consumer, and dissipative release for the resource (Eq. (5)). In every case (including longer and nonlinear food chains), it combines the base magnitude of productivity and mortality, and how they are transmitted and modulated by trophic interactions and by self-regulation.

- **Dissipative loss χ:** in the consumer's standing biomass, losses caused by consumer mortality divided by gains from primary production.

- **Dissipative release ξ:** in the ressource's standing biomass, gains caused by consumer mortality (prey release) divided by gains from primary production.

- **Dynamically consistent chains:** food chains where, in the long-term, the two levels respond in the same direction to nutrient enrichment, and in opposite directions to fish addition.

Mechanisms of cascades and biomasses

What can cause correlations between cascades and biomasses?

Figure 1: Causal relationship between biomass distribution across trophic levels and trophic cascades. Previous studies have investigated the mechanisms controlling the distribution of biomass across trophic levels and the strength of trophic cascades in isolation (top row). Yet, whether there exists a relationship between the two and whether this relationship is causal remains unexplored. There are several ways in which cascades and biomass distribution could be related. They could correlate across food chains via dependency on common parameters, but have no causal relationship between them (right bottom row). In this case, an observed relationship would reflect how common ecological parameters vary in the specific set of food chains under study. On the other hand, if cascades and biomasses are directly causally related (left bottom row), one can then deduce robust predictions from their relationship.

Figure 2: Relationship between static and dynamic properties in food chains. (a) Within one system, a simple expectation (H0) is that a trophic level's response to a perturbation is simply proportional to its biomass. Deviations from this trend indicate the amplification or attenuation of perturbations between levels (Shurin *et al.*, 2002). Food chain theory instead suggests two limiting cases where equilibrium biomasses are related either to bottom-up effects only, or to top-down effects only. Our framework connects these two limits on a single axis of trophic dissipation. From the consumer perspective we identify *dissipative loss* χ and from the resource perspective *dissipative release* ξ (see Box 1). When these two parameters are fixed, all other sources of variation affect biomasses only through trophic cascades, and a direct causal link can be established between these properties (Figure 1). Dissipative release ξ governs qualitative causality in the dynamical-structural relationship, while dissipative loss χ governs its quantitative slope. When dissipative release is small, as in (Barbier & Loreau, 2019), we can use the biomass ratio B_r to precisely predict the upward response ratio $A_\uparrow$ (red arrow), which is amplified by a factor $1/(1-\chi)$. However, we cannot predict the downward response, which could be either strong or weak. Conversely, if dissipative release is large (as in various classic models, e.g. (Oksanen *et al.*, 1981)), then the biomass ratio predicts the downward response (red arrow), amplified or attenuated by a factor $(1-\chi)/\chi$. Around the threshold $\xi \sim 1$, downward or upward responses are entangled, and the biomasses are partially correlated with both (see Figure 4). (b) Even when a causal relationship holds between biomass distribution and trophic cascades within one system, we may only be able to infer it from the statistical relationship across systems if the corresponding parameters are sufficiently similar.

Figure 3: Data analyses illustration. (a) We used data from 31 mesocosm experiments of pelagic food webs that monitored both biomass distributions and long-term responses to nutrient enrichment and fish addition, and estimated the biomass response of two trophic levels to evaluate whether we could detect any of the possible patterns in a natural system. (b) Multiple chains can be identified in these communities. We analysed all possible chains to ensure we capture the dynamics of a *consistent chain* and discarded those chains that did not follow the expected dynamics. Two examples are provided. On the left, all zooplankton species are aggregated into trophic level 2 and all phytoplankton species are aggregated into trophic level 1 (chain: Z-A). On the right, the chain analysed is composed by small herbivores (trophic level 2) consuming highly edible phytoplankton (trophic level 1) (chain: H1-A1). All chains analysed are: Z-A (31), C1-A2 (17), C1-H1 (22), H1-A1 (21), H2-A2 (14), H2-A1 (14), where the numbers in brackets indicate the number of experiments that contained information for each chain type. (c and d) Examples of response evaluation to determine if the chains considered behave as *dynamically consistent chains*. The evaluation was based on the response to both perturbations (i.e., nutrient enrichment and fish addition) of each trophic level. In dynamically consistent chains, we expect the two trophic levels to respond in the same direction to nutrient enrichment, and in opposite directions to fish addition. In (c) we illustrate a case where the system behaved as a dynamically consistent chain. Both trophic levels respond positively to nutrient enrichment (dashed arrows point towards nutrient enrichment), shown by the increase in biomass from blue (no nutrient treatment, $x_\uparrow = 0$) to green lines (nutrient enrichment, $x_\uparrow = 1$), and they show opposite responses to fish addition (coloured arrows point towards fish addition). Phytoplankton biomass increases from $x_\downarrow = 0$ (no fish addition) to $x_\downarrow = 1$ (fish addition), while zooplankton biomass decreases with fish addition. In (d) we show a case where the system do not behave as a consistent chain. First, phytoplankton biomass decreases with nutrient enrichment while zooplankton biomass increases. Second, both trophic level respond negatively to fish addition. Additionally, we can also observe the filtering of the responses based on the associated estimated error. While in (c) we observe similar slopes between the green and blue lines (i.e., low error) in (d) the error associated to the estimation of the responses is large, shown by the differences in the slope between green and blue lines.

Figure 4: The relationship between biomass distribution across trophic levels and response to perturbations, given dissipative loss χ (Box 1). (a) Within a system, if we vary responses $A_\uparrow$ and $A_\downarrow$, the threshold $1/\chi$ determines the importance of bottom-up and top-down effects. When $A_\uparrow A_\downarrow \ll 1/\chi$ (corresponding to $\xi \ll 1$ in Figure 2), the biomass distribution only reflects bottom-up effects, as shown by the vertical isolines. Above that threshold, B_r only reflects top-down effects (horizontal isolines), and close to the threshold, it entangles information about both bottom-up and top-down effects. (b) An across-system statistical relationship emerges when there is small variation in χ. Simulations of the model (4) (for the range of parameters described in the main text) showing two limit cases of Eq. (10). For low dissipation (here we selected realizations where $\xi < 10^{-1}$, and $\chi \in (0, 0.4)$), upward response ratio $A_\uparrow$ shows a proportional relationship with biomass ratio (the solid line has slope 1 in log-log scale) while there is no correlation between B_r and downward response ratio $A_\downarrow$. For large dissipation (here $\xi > 10$ and $\chi \in (0.6, 0.8)$), we observe the opposite trend: downward response ratio $A_\downarrow$ is inversely proportional to biomass ratio (solid line has slope -1 on log-log scales) while there is no correlation with $A_\uparrow$.

Figure 5: Empirical relationship between biomass ratio and response ratios. (a) Relationship between biomass ratio and bottom-up response ratio ($A_\uparrow = \frac{\Delta B_{2\uparrow}}{\Delta B_{1\uparrow}}$) due to nutrient enrichment. In (b) relationship between biomass ratio and top-down response ratio ($A_\downarrow = -\frac{\Delta B_{1\downarrow}}{\Delta B_{2\downarrow}}$) due to fish addition. Each point represents a food chain in our analyses. Blue lines represent the linear regression and shaded areas represent 95% confidence intervals. Dashed lines represent the limits of the empirical pattern used in the simulation analyses to filter the prior distribution of parameters. Coloured regions represent the simple expectation scenario where a trophic level's response to a perturbation is proportional to its biomass. In the green region perturbations are amplified through the chain while in the orange perturbations are attenuated. Thus, the interface between the two regions represents the 1:1 line where the response of each trophic level is proportional to their biomass. Notice that patterns shown in (a) and (b) are excluding the outliers, which results into an exclusion of the extremes of the biomass ratio distribution (See Figure S1.3 in Appendix S3 for results including the outliers).

Figure 6: Prior and posterior distributions of model parameters. Blue histograms represent the prior distribution of the parameters introduced into the model (conditioned on coexistence). Orange histograms represent the distribution of the parameters after filtering them to match empirical patterns. Top: distribution of $\xi = (r/g)(\alpha/D_2)$ (dissipative release) which determines the nature of within-system relationship between biomass ratio and upwards and downwards response ratios. Dissipative release is the product of two non dimensional ratios. Lower left: distribution of the ratio of predator mortality to prey growth $\phi = r/g$, Lower right: distribution of the ratio α/D_2, which represents the relative strength of trophic interactions and predator self-regulation. We see that the posterior distribution is centered close to 0 in log scale, suggesting that self-regulation rates are comparable to predation rates.

Can biomass distribution across trophic levels predict trophic cascades?

Supporting Information

Núria Galiana, Jean-François Arnoldi, Matthieu Barbier, Amandine Acloque,
Claire de Mazancourt and Michel Loreau

Appendix S1. Empirical data

Table S1.1: Experimental studies used in the analyses. We extracted 31 experimental studies analysing bottom-up and top-down effects in pelagic food chains after press perturbations. Detailed information about the studies used can be found in (Hulot *et al.*, 2014).

Article	N. of experiments
(Vanni, 1987a,b)	2
(Van De Bund & Van Donk, 2004)	2
(Lacroix & Lescher-Moutoue, 1991)	1
(Fernández-Aláez *et al.*, 2004)	2
(Stephen *et al.*, 2004)	2
(Lynch & Shapiro, 1981)	1
(Menezes *et al.*, 2010)	1
(Qin & Culver, 1995)	1
(Hansson *et al.*, 2004)	1
(Romo *et al.*, 2004)	2
(Dickman *et al.*, 2008)	1
(Acuña *et al.*, 2008)	1
(Beresford & Jones, 2010)	1
(Drenner *et al.*, 1990, 1989)	2
(Faafeng *et al.*, 1990)	2
(Hietala *et al.*, 2004)	2
(McQueen *et al.*, 1992)	1
(Proulx *et al.*, 1996)	2
(Sinistro, 2010)	1
(Riemann, 1985)	1
(Mazumder *et al.*, 1990)	1

Table S1.2: Total number of experiments analysed per chain type and number of chain-like responses for upwards amplification and downwards amplification for each chain type.

Chain type	Total number of experiments	Upwards amplification	Downwards amplification
Z-A	31	10	12
C1-A2	17	4	2
C1-H1	22	5	6
H1-A1	21	7	7
H2-A1	14	3	6
H2-A2	14	4	2

Figure S1.1: Distributions of the estimated responses to nutrient enrichment (↑) and fish addition (↓) treatments of each trophic level biomass (B_1 and B_2, for trophic level 1 and 2, respectively) relative to their original biomass in experimental mesocosms. Blue distributions represent the estimated responses of all possible chains analysed before filtering them to ensure they behave like dynamically consistent chain and before selecting the ones that presented estimated responses with low error. Orange distributions show the estimated responses of the chains that were identified as dynamically consistent chains. In dynamically consistent chains, we expect the two levels to respond in the same direction to nutrient enrichment, and in opposite directions to fish addition. More concretely, we expect responses of trophic level 1 to nutrient enrichment (a) and to fish addition (b) to be generally positive, together with the biomass responses of trophic level 2 to nutrient enrichment (c). Therefore, most of the chains selected present a positive biomass response to the perturbation. In contrast, biomass responses of trophic level 2 to fish addition (d) are expected to be generally negative. Thus, we observe that very few chains are selected when trophic level 2 presents a positive response to fish addition.

In the following lines we present the statistical results from the comparison between the filtered and non-filtered distributions using Kolmogorov-smirnov tests for each response. If the K-S statistic is small or the p-value is high, then we cannot reject the hypothesis that the two distributions are the same. a) statistic = 0.253, p-value = 0.167; b) statistic = 0.197, p-value = 0.271; c) statistic = 0.251, p-value = 0.165; d) statistic = 0.17, p-value = 0.476

Figure S1.2: Cumulative distributions of the estimated responses to nutrient enrichment (↑) and fish addition (↓) treatments of each trophic level biomass. Same information as in Figure S1.1 using cumulative distributions. Blue distributions represent the estimated responses of all possible chains analysed before filtering. Orange distributions show the estimated responses of the chains that were identified as dynamically consistent chains. As mentioned above, in dynamically consistent chains, we expect the two levels to respond in the same direction to nutrient enrichment, and in opposite directions to fish addition. Therefore, in (a, b and c) most of the chains selected present a positive biomass response to the perturbation and are accumulated after 0. In contrast, biomass responses of trophic level 2 to fish addition (d) are expected to be generally negative. Thus, we observe that very few chains are selected when trophic level 2 presents a positive response to fish addition, i.e., there is a very small accumulation of chains after 0.

Figure S1.3: Empirical relationship between biomass distribution across trophic levels and response ratio without exclusion of the outliers. (a) Relationship between biomass ratio ($B_r = \frac{B_2}{B_1}$) and bottom-up response ratio ($A_\uparrow = \frac{\Delta B_{2\uparrow}}{\Delta B_{1\uparrow}}$) due to nutrient enrichment. (b) Relationship between biomass ratio and top-down response ratio ($A_\downarrow = -\frac{\Delta B_{1\downarrow}}{\Delta B_{2\downarrow}}$) due to fish addition. Each point represents a food chain in our analyses. Blue lines represent the linear regression and shaded areas represent 95% confidence intervals.

Appensix S2. Theoretical framework

Two-level chain

Let B_i, $i = 1, 2$ be the biomass density of primary producers and consumers, respectively (e.g grams per unit of volume). We denote by $g > 0$ the intrinsic growth rate of primary producers, and $r > 0$ the intrinsic death rate of consumers. The per-capita attack rate of consumers is denoted as $\alpha > 0$ (a rate per unit of biomass density). In the same units we denote by D_i $i = 1, 2$ the density-dependent mortality (or reduction of biomass productivity) of both trophic levels (i.e., self-regulation). Finally, we write $0 < \varepsilon < 1$ the (non dimensional) efficiency of productivity transfer from primary producers to consumers. This leads us to the following expression for the growth rates $W_i = \frac{1}{B_i}\frac{dB_i}{dt}$ of both populations

$$\begin{cases} W_2 = & \varepsilon\alpha B_1 - D_2 B_2 - r \\ W_1 = & g - D_1 B_1 - \alpha B_2 \end{cases} \tag{12}$$

which define a Lotka-Volterra dynamical system (See Appendix S2 for the nonlinear chain approximation). In what follows we are interested only in non dimensional ratios, such as the biomass ratio of the two levels, and the amplification of a perturbation across the chain (i.e., effects of a perturbation on one trophic level relative to the effects on the other level in which the perturbation was directly applied). We therefore begin by adimensionalizing the problem. There are two units: time and biomass density. To remove the former we can express time relatively to the rate of growth g of primary producers. We write $\tau = gt$ (non dimensional time) and, since $\frac{d}{dt} = g\frac{d}{d\tau}$, expressing the dynamics in those units amounts to dividing by g all terms of the right hand side of (12). To remove the units of biomass, we can express all densities with respect to the carrying capacity $K_1 = g/D_1$ of primary producers. We thus define $b_i = B_i/K_1$ (non dimensional biomass). This lead us to the non dimensional system, denoting $w_i = \frac{1}{b_i}\frac{db_i}{d\tau}$ non dimensional rates, we get that

$$\begin{cases} w_2 = & \varepsilon a b_1 - \psi b_2 - \phi \\ w_1 = & 1 - b_1 - a b_2 \end{cases} ; \quad a = \frac{\alpha}{D_1}; \quad \psi = \frac{D_2}{D_1}; \quad \phi = \frac{r}{g} \tag{13}$$

At equilibrium $\frac{1}{b_i}\frac{db_i}{d\tau} = 0$ so that (if both species coexist), their respective densities satisfy

$$\begin{pmatrix} \psi & -\varepsilon a \\ a & 1 \end{pmatrix} \begin{pmatrix} b_2^* \\ b_1^* \end{pmatrix} = \begin{pmatrix} -\phi \\ 1 \end{pmatrix} \Leftrightarrow \begin{pmatrix} b_2^* \\ b_1^* \end{pmatrix} = V \begin{pmatrix} -\phi \\ 1 \end{pmatrix}$$

where the matrix V is

$$V = \begin{pmatrix} \psi & -\varepsilon a \\ a & 1 \end{pmatrix}^{-1} = \frac{1}{\psi + \varepsilon a^2} \begin{pmatrix} 1 & \varepsilon a \\ -a & \psi \end{pmatrix}$$

We deduce, in particular, the biomass ratio $B_r = b_2^*/b_1^* = B_2^*/B_1^*$ equal to

$$B_r = \frac{\varepsilon a - \phi}{\psi + a\phi}$$

where we can observe that the coexistence condition is that εa (consumption) must be larger than ϕ (mortality). We can also recognize in this expression of B_r the amplification of perturbations. This is because the matrix V that gave us the equilibrium densities also gives us their sensitivity to changes in growth rates. Indeed, consider a perturbation of the primary producer in the form of a persistent change δg of growth rate. In the non dimensional formulation, this is a perturbation $\delta w_1 = \delta g/g$ and it leads to the shift of biomass:

$$\begin{pmatrix} \delta b_2 \\ \delta b_1 \end{pmatrix} = \begin{pmatrix} \psi & -\varepsilon a \\ a & 1 \end{pmatrix}^{-1} \begin{pmatrix} 0 \\ \delta w_1 \end{pmatrix} = \frac{\delta w_1}{\psi + \varepsilon a^2} \begin{pmatrix} \varepsilon a \\ \psi \end{pmatrix}$$

where we recognize the second column of V, times δw_1. We can do the same for a perturbation δw_2 of the consumer's growth rate, which would give us the first column of V, times δw_2. Thus

$$V_{ij} = \frac{\partial b_i^*}{\partial w_j}$$

so that bottom-up amplification (i.e., perturbation applied on resources), defined as $A_\uparrow = \left| \frac{\partial b_2}{\partial w_1} / \frac{\partial b_1}{\partial w_1} \right| = |V_{21}/V_{11}|$ reads

$$A_\uparrow = \frac{\varepsilon a}{\psi}$$

Similarly, if the perturbation is applied to the consumer we get that top-down amplification $A_\downarrow = \left| \frac{\partial b_1}{\partial w_2} / \frac{\partial b_2}{\partial w_2} \right| = |V_{12}/V_{22}|$ is in fact

$$A_\downarrow = a$$

From these notations yields the following expression for the relationship between biomass ratio and trophic cascades:

$$B_r = \frac{(1-\chi)A_\uparrow}{1 + \chi A_\downarrow A_\uparrow}; \ \chi = \phi/(\varepsilon a) \tag{14}$$

Where $\chi = $ can be interpreted as a measure of dissipative loss, the ratio of biomass losses of the consumer, over gains from primary productivity. If instead we were to define $\rho = \chi A_\uparrow$ we would have had

$$B_r = \frac{A_\uparrow - \rho}{1 + A_\downarrow \rho}; \ \rho = \phi/\psi \tag{15}$$

Where $\rho = \frac{\phi}{\psi} = \frac{K_2'}{K_1}$ can be interpreted as the ratio of intraspecific characteristic densities: carrying capacity K_1 for the producer, and the characteristic density K_2' of consumer population where intraspecific competition affects the growth as much as linear mortality. In particular, constant ρ would relate the biomass ratio to the amplification factors, which signal whether cascades are weak or strong relative to the proportional baseline H0. We get that:

$$B_r = \rho \frac{a_\downarrow + 1}{a_\uparrow - 1} \tag{16}$$

This expression can be misleading, because the amplifications $a_\uparrow = A_\uparrow/B_r$ and $a_\downarrow = A_\downarrow \times B_r$, are not independent of the biomass ratio, which is in turn shaped by ρ: in particular, we know from (10) that, if $\rho = 0$, then $a_\uparrow = 1$. What this expression shows, however, is the confounding role of ρ. If this parameter varies across systems more than either amplification factor, no relationship between B_r and cascade strength will be found. If, however, ρ has low variance across ecosystems while $a_\downarrow$ varies significantly, then we can expect a relationship such as the one observed in Figure 5e.

Multi-level chain

The above reasoning lends itself to a generalization to arbitrary number levels. If $\alpha_{i,i-1}$ is the attack rate of level i on the level below, D_i is the strength of intraspecific competition and r_i is mortality, as for the the two level system, removing dimensions leads to the system

$$\begin{cases} w_N = & \varepsilon_N a_N b_{N-1} - \psi_N b_N - \phi_N \\ \vdots & \vdots \\ w_i = & \varepsilon_i a_i B_{i-1} - \psi_i B_i - a_{i+1} b_{i+1} - \phi_i; \ a_i = \frac{\alpha_{i,i-1}}{D_1}; \ \psi_i = \frac{D_i}{D_1}; \ \phi_i = \frac{r_i}{g} \\ \vdots & \vdots \\ w_1 = & 1 - b_1 - a_2 b_2 \end{cases}$$

From these we could compute the sensitivity matrix $V = (V_{ij})$ as previously defined. It is the inverse of the interaction matrix $\alpha = (\alpha_{ij})$

$$V_{ij} = \frac{\partial b_i^*}{\partial w_j} = (\alpha^{-1})_{ij} \text{ where } \alpha_{ij} = \frac{\partial w_i}{\partial b_j}$$

which takes the more explicit form here

$$\alpha = \begin{pmatrix} \psi_N & \varepsilon_N a_N & & & \\ -a_N & \psi_{N-1} & \varepsilon_{N-1}a_{N-1} & & \\ & & \ddots & & \\ & & & \psi_1 & \varepsilon_1 a_1 \\ & & & -a_1 & 1 \end{pmatrix}$$

The equilibrium densities (assuming coexistence) are then given by

$$b_i^* = V_{i1} - \sum_{j>1} V_{ij}\phi_j$$

Amplifications now read, if $j > i$ (upward amplification)

$$A_{ij}^\uparrow = \frac{V_{ji}}{V_{ii}} \tag{17}$$

and the same holds if $j < i$ (downward amplification $A_{ij}^\downarrow$). The biomass ratio between top and lowest level reads

$$B_r = \frac{b_N^*}{b_1^*} = \frac{B_N^*}{B_1^*} = \frac{\frac{V_{N1}}{V_{11}} - \sum_{j>1} \frac{V_{Nj}}{V_{11}}\phi_j}{1 - \sum_{j>1} \frac{V_{1j}}{V_{11}}\phi_j} = \frac{(1-\chi_N)A_{1N}^\uparrow - \sum_{1<j<N} \chi_j A_{1j}^\uparrow A_{jN}^\uparrow}{1 + \sum_{j>1} \chi_j A_{j1}^\downarrow A_{jN}^\uparrow} \tag{18}$$

where we defined dissipative loss at level j

$$\chi_i = \frac{V_{jj}r_j}{V_{j1}g} \tag{19}$$

thus showcasing the generality of our approach: top down responses are causally related to top-to-bottom biomass ratio if total dissipative release $\sum_j \chi_j A_{j1}^\downarrow A_{jN}^\uparrow$ is comparable (or larger) than 1. Note that having many levels is likely to extend the entangled domain, where both upwards and downwards transmission of perturbations shape the biomass ditribution.

Nonlinear chain

In fact, the above results are not specific to linear growth rates (i.e Lotka-Volterra). For any form of growth rate functions w_i we can still define the interaction matrix $\alpha_{ij} = \frac{\partial w_i}{\partial b_j}$ and the write the density independent part of those functions $-\phi_i$ for consumers (relative intrinsic mortality) and 1 for the primary producer (intrinsic growth rate, used to set a time scale). As before $V_{ij} = (\alpha^{-1})_{ij} = \frac{\partial b_i}{\partial w_j}$ and we can therefore write

$$\begin{aligned} b_i^* &= \int_0^1 \frac{\partial b_i^*}{\partial w_1}dw_1 + \sum_{j>1} \int_0^{-\phi_j} \frac{\partial b_i^*}{\partial w_j}dw_j \\ &= \int_0^1 V_{i1}dw_1 - \sum_{j>1} \int_0^{\phi_j} V_{ij}dw_j \\ &= \langle V_{i1} \rangle - \sum_{j>1} \langle V_{ij} \rangle \phi_j \end{aligned}$$

where $\langle V_{ij} \rangle$ is the average sensitivity of species i to a change in growth rate of species j, as relative mortality is varied from 0 to its actual value, ϕ_j. We now have

$$B_r = \frac{b_N^*}{b_1^*} = \frac{B_N^*}{B_1^*} = \frac{\frac{\langle V_{N1} \rangle}{\langle V_{11} \rangle} - \sum_{j>1} \frac{\langle V_{Nj} \rangle}{\langle V_{jj} \rangle} \frac{\langle V_{jj} \rangle}{\langle V_{11} \rangle}\phi_j}{1 - \sum_{j>1} \frac{\langle V_{1j} \rangle}{\langle V_{jj} \rangle} \frac{\langle V_{jj} \rangle}{\langle V_{11} \rangle}\phi_j}$$

The difference with the linear chain is that now the amplification of a weak perturbation will be (for instance)

$$A_{Nj}^\downarrow = \frac{V_{Nj}}{V_{jj}} \tag{20}$$

evaluated at the actual conditions defined by the relative intrinsic rates ϕ_i. The amplification is not exactly equal to the term

$$\mathcal{A}_{Nj}^\downarrow = \frac{\langle V_{Nj} \rangle}{\langle V_{jj} \rangle} \tag{21}$$

that appears in the above expression of biomass ratio. Nonetheless, unless the growth rates w_i are strongly nonlinear functions of b_i, $A_{Nj}^\downarrow$ and $\mathcal{A}_{Nj}^\downarrow$ can be expected to be strongly correlated. Thus if

$$\chi_i = \frac{\langle V_{ii} \rangle r_i}{\langle V_{i1} \rangle g} \tag{22}$$

we have that

$$B_r = \frac{(1-\chi_N)\mathcal{A}_{1N}^\uparrow - \sum_{1<j<N} \chi_j \mathcal{A}_{1j}^\uparrow \mathcal{A}_{jN}^\uparrow}{1 + \sum_{j>1} \chi_j \mathcal{A}_{j1}^\downarrow \mathcal{A}_{jN}^\uparrow} \tag{23}$$

This extension is useful to understand why nonlinearities can allow for some patterns in the data that would be forbidden by a the linear chain model. Indeed, if we go back to the expression of B_r in two level chain (14) we see that $A_\uparrow$ must be strictly larger than B_r. In the data this is not always the case. Slight non-linearity will remove this strong constraint since it is now $\mathcal{A}_\uparrow$ that must be larger than B_r and not $A_\uparrow$. As we show numerically, only slight non-linearity (in the form of handling time by the consumers) is enough to produce theoretical pattern consistent with the data, as shown in Figure (S1.4).

Figure S1.4: Comparison of simulations and data for linear and non-linear resource-consumer models. In the non-linear model, we use a Holling type II functional response $f(b_1) = \frac{ab_1}{1+hab_1}$ for consumption, with relative handling time h. All other parameters are the same, and drawn from the same distributions, as in the linear model described in the main text, and h is drawn uniformly on log scales in $[10^{-4}, 10^0]$. Upper row: $A_\uparrow$ vs B_r for empirical data points (orange) and simulated data points (blue). In the linear model (left), $A_\uparrow$ cannot be smaller than B_r, yet the data shows points that violate this condition. (right) slight non linearity can relax this constraint. Bottom row: distribution of handling time after filtering by the empirical patterns. We observe that only small values are selected (weak non linearity).

Appendix S3. Supplementary ABC simulation results

Figure S3.1: Experimental data versus simulated data. (a) Histograms of the biomass ratio in the experimental systems (orange) and in the simulated data (blue). Both distributions are normalised such that the area under the histogram sums to 1. (b) Relationship between bottom-up amplification ($A_\uparrow = \frac{\Delta B_{2\uparrow}}{\Delta B_{1\uparrow}}$) due to nutrient enrichment and top-down amplification ($A_\downarrow = -\frac{\Delta B_{1\downarrow}}{\Delta B_{2\downarrow}}$) due to fish addition. Orange points represent all the $A_\uparrow$ values in the empirical data. Green points represent all the $A_\downarrow$ values in the empirical data. Red points represent the empirical data for which both values of $A_\uparrow$ and $A_\downarrow$ were accessible. Blue points represent the simulated data, constrained to reproduce empirical patterns (ABC inference method). Note the slight negative correlation between $A_\uparrow$ and $A_\downarrow$. (c) Distribution of relative attack rates $a = \alpha/D_1$ after constraining for coexistence (light blue) and after constraining by empirical patterns (orange). (d) as in c) but for ratio of consumer to producer self-regulation $\psi = D_2/D_1$. The large variation of ψ may be responsible for the anticorrelation of $A_\uparrow$ and $A_\downarrow$ observed in (b). Indeed $A_\uparrow/A_\downarrow \sim \psi$. The broadest possible range of ψ is achieved if the larger $A_\uparrow$ are associated with the smallest $A_\downarrow$ and vice versa. Conversely, the larger the range in ψ the stronger the anti-correlation between the two response ratios. (e) as in (c-d) but for mortality rate over primary productivity $\phi = r/g$. (f) as in (c-e) but for conversion efficiency ε.

Figure S3.2: Relationship for $A_\downarrow$ and B_r for the three regimes of dissipative release, where the qualitatively relationship is maintained but their interpretation should differ. From a correlation for low dissipation release to an actual causation for high dissipation release.

References

Acuña, P., Vila, I. & Marín, V.H. (2008). Short-term responses of phytoplankton to nutrient enrichment and planktivorous fish predation in a temperate South American mesotrophic reservoir. *Hydrobiologia*, 600, 131–138.

Beresford, A.L. & Jones, J.I. (2010). Weedbeds and big bugs: the importance of scale in detecting the influence of nutrients and predation on macroinvertebrates in plant-dominated shallow lakes. *Freshwater Biology*, 55, 514–530.

Dickman, E.M., Newell, J.M., González, M.J. & Vanni, M.J. (2008). Light, nutrients, and food-chain length constrain planktonic energy transfer efficiency across multiple trophic levels. *Proceedings of the National Academy of Sciences*, 105, 18408–18412.

Drenner, R.W., Smith, J.D., Mummert, J.R. & Lancaster, H.F. (1990). Responses of a eutrophic pond community to separate and combined effects of N: P supply and planktivorous fish: a mesocosm experiment. *Hydrobiologia*, 208, 161–167.

Drenner, R.W., Threlkeld, S.T., Smith, J.D., Mummert, J.R. & Cantrell, P.A. (1989). Interdependence of phosphorus, fish, and site effects on phytoplankton biomass and zooplankton. *Limnology and oceanography*, 34, 1315–1321.

Faafeng, B.A., Hessen, D.O., Brabrand, Å. & Nilssen, J.P. (1990). Biomanipulation and food-web dynamics—the importance of seasonal stability. In: *Biomanipulation Tool for Water Management*. Springer, pp. 119–128.

Fernández-Aláez, M., Fernández-Aláez, C., Bécares, E., Valentín, M., Goma, J. & Castrillo, P. (2004). A 2-year experimental study on nutrient and predator influences on food web constituents in a shallow lake of north-west Spain. *Freshwater Biology*, 49, 1574–1592.

Hansson, L., Gyllström, M., Ståhl-Delbanco, A. & Svensson, M. (2004). Responses to fish predation and nutrients by plankton at different levels of taxonomic resolution. *Freshwater Biology*, 49, 1538–1550.

Hietala, J., Vakkilainen, K. & Kairesalo, T. (2004). Community resistance and change to nutrient enrichment and fish manipulation in a vegetated lake littoral. *Freshwater Biology*, 49, 1525–1537.

Lacroix, G. & Lescher-Moutoue, F. (1991). Interaction effects of nutrient loading and density of young-of-the-year cyprinids on eutrophication in a shallow lake: an experimental mesocosm study. *Mem. Ist. Ital. Idrobiol*, 48, 53–73.

Lynch, M. & Shapiro, J. (1981). Predation, enrichment, and phytoplankton community structure 1. *Limnology and oceanography*, 26, 86–102.

Mazumder, A., McQueen, D.J., Taylor, W.D., Lean, D.R.S. & Dickman, M.D. (1990). Micro-and mesozooplankton grazing on natural pico-and nanoplankton in contrasting plankton communities produced by planktivore manipulation and fertilization. *Archiv für Hydrobiologie*, 118, 257–282.

McQueen, D.J., France, R. & Kraft, C. (1992). Confounded impacts of planktivorous fish on freshwater biomanipulations. *Archiv für Hydrobiologie*, 125, 1–24.

Menezes, R.F., Attayde, J.L. & RIVERA VASCONCELOS, F. (2010). Effects of omnivorous filter-feeding fish and nutrient enrichment on the plankton community and water transparency of a tropical reservoir. *Freshwater Biology*, 55, 767–779.

Proulx, M., Pick, F.R., Mazumder, A., Hamilton, P.B. & Lean, D.R.S. (1996). Effects of nutrients and planktivorous fish on the phytoplankton of shallow and deep aquatic systems. *Ecology*, 77, 1556–1572.

Qin, J. & Culver, D.A. (1995). Separate and combined effects of larval walleye and fertilization on plankton community structure in enclosures. *Journal of Freshwater Ecology*, 10, 385–391.

Riemann, B. (1985). Potential importance of fish predation and zooplankton grazing on natural populations of freshwater bacteria. *Appl. Environ. Microbiol.*, 50, 187–193.

Romo, S., Miracle, M.R., Villena, M., Rueda, J., Ferriol, C. & Vicente, E. (2004). Mesocosm experiments on nutrient and fish effects on shallow lake food webs in a Mediterranean climate. *Freshwater Biology*, 49, 1593–1607.

Sinistro, R. (2010). Top-down and bottom-up regulation of planktonic communities in a warm temperate wetland. *Journal of Plankton Research*, 32, 209–220.

Stephen, D., Balayla, D.M., Collings, S.E. & Moss, B. (2004). Two mesocosm experiments investigating the control of summer phytoplankton growth in a small shallow lake. *Freshwater Biology*, 49, 1551–1564.

Van De Bund, W.J. & Van Donk, E. (2004). Effects of fish and nutrient additions on food-web stability in a charophyte-dominated lake. *Freshwater Biology*, 49, 1565–1573.

Vanni, M.J. (1987a). Effects of food availability and fish predation on a zooplankton community. *Ecological Monographs*, 57, 61–88.

Vanni, M.J. (1987b). Effects of nutrients and zooplankton size on the structure of a phytoplankton community. *Ecology*, 68, 624–635.