Citrus vascular proteomics highlights the role of peroxidases and serine proteases during

Huanglongbing disease progression

Jessica Y. Franco[1], Shree P. Thapa[1], Zhiqian Pang[2], Fatta B. Gurung[3], Thomas W.H. Liebrand[1],

Danielle M Stevens[1], Veronica Ancona[3], Nian Wang[2], Gitta Coaker[1]

[1]Department of Plant Pathology, University of California, Davis, USA,

[2]Citrus Research and Education Center, University of Florida, Lake Alfred, USA,

[3]Citrus Center, Texas A&M University- Kingsville, Wescalo, Texas, USA,

*Corresponding author: Gitta Coaker (glcoaker@ucdavis.edu)

Running Title: Vascular proteomics in citrus during Huanglongbing disease

Abbreviations: HLB, huanglongbing; *C*Las, *Candidatus* Liberibacter asiaticus; ACP, Asian

citrus psyllid; ABPP, activity-based protein profiling; PCA, principle component analysis.

Abstract:

Huanglongbing (HLB) is the most devastating and widespread citrus disease. All commercial citrus varieties are susceptible to the HLB-associated bacterium, *Candidatus* Liberibacter asiaticus (*C*Las), which resides in the phloem. The phloem is part of the plant vascular system and is involved in sugar transport. To investigate the plant response to *C*Las, we enriched for proteins surrounding the phloem in an HLB susceptible sweet orange variety, Washington navel (*Citrus sinensis* (L) Osbeck). Quantitative proteomics revealed global changes in the citrus proteome after *C*Las inoculation. Plant metabolism and translation were suppressed, while defense-related proteins including proteases and protease inhibitors were induced in the vasculature. Specific serine proteases were upregulated in Washington navel in the presence of *C*Las based on quantitative proteomics. Subsequent activity-based protein profiling revealed genotype-specific variation in serine protease activity in other *C. sinensis* sweet orange varieties. Plant peroxidases involved in extracellular ROS production or scavenging were induced in vascular enriched tissues based on quantitative proteomics. While peroxidase transcript accumulation was induced in *C*Las infected sweet orange varieties, peroxidase enzymatic activity varied between varieties, supporting the genotype-specific regulation of enzymatic activity. The observations in the current study highlight dynamic global reprogramming of the citrus vascular proteome and genotype-specific regulation of different enzyme classes in response to *C*Las infection. These results open an avenue for further investigation of diverse responses to HLB across different citrus genotypes.

Introduction

Currently, the citrus industry is facing a major threat from the bacterial vector-borne disease huanglongbing (HLB), also known as citrus greening. The HLB-associated pathogen, *Candidatus* Liberibacter asiaticus (*C*Las), is spread by *Diaphorina citri*, a hemipteran insect. All commercial citrus varieties are susceptible to *C*Las and current interventions such as *D. citri* control and infected tree removal have had little success in controlling the disease (1). Sweet orange varieties (*Citrus sinensis*) exhibit the fastest decline, while lemon (*C. latifolia*), mandarins (*C. reticulata*), and tangerines (*C. reticulata*) present delayed symptom progression but ultimately succumb to HLB (2, 3). In the United States, HLB was first diagnosed in 2005 in Florida and has since spread to Texas, Louisiana, South Carolina, Georgia and California (4). In California, *C*Las positive citrus was detected in 2012 and there are currently 1,788 positive plants identified by qPCR (5, 6). In Florida, it is estimated that 95% of mature trees in commercial groves are infected, resulting in a loss of over 7.8 billion dollars in revenue and 7,500 jobs since 2007 (7). For decades, Florida has been the primary citrus producer in the United States. However, due to losses from HLB, California is now the primary citrus producing area (8).

Plant pathogens can be directly deposited into the vascular system (xylem or phloem) by piercing sucking insects. *D. citri* deposits *C*Las into plant phloem sieve elements, where the bacteria proliferate (9). The phloem is rich in sugars, amino acids, and proteins (10). *C*Las infected leaf tissues exhibit increased deposition of callose, a beta 1-3 glucan, and accumulation of starch that constricts symplastic transport, which contributes to symptom development (11, 12). HLB symptoms are characterized by yellowing of the shoots and leaves as well as hard, misshapen, and bitter fruit (13, 14). Early detection efforts have been challenging to develop due

to the long latent period and the uneven distribution of CLas within its plant host (15). The

inability to culture CLas has hampered progress in studying the pathogen and its virulence

mechanisms.

One of the paramount mechanisms of bacterial pathogenesis is the presence of secretion

systems facilitating the translocation of proteins, or effectors, to the plant host (16). CLas

possesses the type I and general sec secretion system (17, 18). The sec pathway is critical for

virulence in other phloem-colonizing pathogens including mollicutes (*Ca.* Phytoplasma) (19).

Most CLas sec-dependent effectors (SDEs) were highly expressed *in planta*, indicating they are

important for pathogen proliferation (18, 20, 21). Transient expression of a conserved CLas

SDE, SDE1, induces cell death, starch accumulation, and callose deposition in *Nicotiana*

benthamiana (22). In citrus, SDE1 inhibits the activity of immune related proteases which may

serve as a strategy to dampen defense responses (23). Multiple secreted CLas proteins can

suppress immune responses (23–26). Upon pathogen perception, the production of reactive

oxygen species (ROS), including hydrogen peroxide, serve as an antimicrobial as well as a

secondary immune signal (27, 28). CLas peroxiredoxins, LasBCP and LasdPrx5, increased

oxidative stress tolerance in *Liberibacter crescens* (a culturable surrogate for CLas) by

suppressing H_2O_2 accumulation (24, 29). CLas encodes a salicylic acid hydroxylase that

degrades salicylic acid (SA), a plant defense hormone (26). Collectively, these data demonstrate

that CLas actively manipulates plant defense responses.

In order to understand the *C. sinensis* response to CLas, previous studies have

investigated dynamic changes during infection by microarray, RNAseq, and proteomics (30–32).

Microarray analyses revealed induction of genes associated with sugar metabolism, plant

defenses, phytohormones, and cell wall metabolism in infected leaf samples (33). Transcripts

encoding proteins involved in sugar and starch metabolism, stress responses, transport,

detoxification, and lipid metabolism were significantly upregulated during *C*Las infection, while

processes involved in photosynthesis were downregulated in mature leaf, stem, and root tissues

(30, 31, 34, 35). *C*Las induces plant defense responses in leaf tissue, including the transcription

of pathogenesis-related (PR) genes, chitinases, callose synthases, and defense-related WRKY

transcription factors (11, 31, 33, 35, 36). Consistent with these observations, infected *C. sinensis*

midveins exhibit callose and starch accumulation in the sieve tubes that lead to phloem plugging

and collapse (33, 34, 37). However, in roots, genes encoding callose hydrolases were induced

and callose synthases were repressed (35). These data illustrate that not all tissue types respond

ubiquitously and highlight tissue-specific responses during *C*Las infection.

Multiple studies have elucidated host responses intact leaf, root, and fruit samples(30,

31, 35, 36). However, *C*Las is limited to phloem sieve elements and there is a lack of proteomic

studies on vascular exudates. In this study, we performed comparative proteomic analyses to

investigate changes in the vascular proteome after *C*Las infection in the sweet orange variety

Washington Navel (*Citrus sinensis* (L) Osbeck). Our results highlight the accumulation of

peroxidases and proteases upon *C*Las infection, indicating the activation of plant defense

responses. Multiple citrus peroxidases were induced at the transcript and protein level during

infection across different citrus genotypes and environmental conditions. Serine protease activity

in leaf tissues also dynamically changed after *C*Las infection under greenhouse and field

conditions. This study advances our understanding of vascular changes and highlights the

important role of peroxidases and proteases in different citrus genetic backgrounds during *C*Las

infection.

Experimental Procedures

Experimental Design and Statistical Rationale

For vascular proteomics, eight Washington Naval trees grafted onto Carrizo rootstock were

grown in the UC Davis Contained Research Facility, a Biosafety level 3 greenhouse. Seven

months after grafting onto Carrizo rootstock, plants were graft-inoculated or mock-inoculated

with the *C*Las isolate, HHCA obtained from Hacienda Heights, California, USA. Plants were

grown in a completely randomized design, tested for *C*Las titers by qPCR monthly, and vascular

exudates were harvested ten months post graft-inoculation. Vascular exudates from four mock

and four *C*Las-graft inoculated Washington navel trees were extracted using the centrifugation

method (10). For mass spectrometry analyses, samples were blinded to the operator. To quantify

differentially changing proteins, label-free quantification using MaxQuant (v 1.5.1.0) was

performed. Protein identification search parameters included a decoy search database in addition

to a False Discovery Rate (FDR) of < 0.05. Statistical analyses were performed on four

individual trees per treatment (n=4). Differentially changing proteins were identified using a

two-sample student's t-test with an alpha-value of 0.05 for truncation.

Plant Growth and Inoculation Conditions

Washington Navel: Proteomic analyses were performed on citrus grown and maintained at the

University of California, Davis Contained Research Facility greenhouse. The greenhouse

temperature was maintained at 27°C. High pressure sodium lights were provided as

supplemental lighting for 16 hours. The greenhouse was covered with shade cloth (50% shade)

between mid-April and mid-October. Eight Washington navel (*Citrus sinensis* (L.) Osbeck) buds

were grafted onto Carrizo rootstock using the inverted T budding method. After seven months,

four Washington navel scions were mock or *C*Las graft-inoculated on February 2014. Samples

were harvested 10 months post-graft inoculation and tested for *C*Las presence by qPCR each

month. A random sampling of 6-8 leaves was taken from each tree and pooled for DNA

extraction. DNA was extracted using the Qiagen MagAttract plant DNA extraction kit (Qiagen,

Valencia, CA, USA). *C*Las presence was detected by qPCR using the USDA-APHIS-PPQ

protocol (38). *C*Las primers amplifying the 16s rDNA (HLBas/HLBr) were used for *C*Las

detection. The citrus cytochrome oxidase (COXf/r) primers and probe were used as an internal

control (38). All primers, gene names, and accession numbers are provided in Supplementary

Table S1.

<u>Hamlin sweet orange:</u> Hamlin sweet oranges (*Citrus sinensis* (L) Osbeck) were grown in the

greenhouse at 25°C. Trees were exposed to *C*Las positive *D. citri* for 2 months. Psyllids were

removed and plants were transferred to a psyllid-free greenhouse and sampled one year after

*C*Las transmission. *C*Las detection was performed using ethidium bromide monoazide (EMA)-

qPCR using primers that amplify the β-operon (39). Samples were collected from uninfected and

symptomatic Hamlin sweet orange, flash frozen, and sent to the Contained Research Facility on

dry ice for further processing.

<u>N-33</u>: Mature N-33 (*Citrus sinensis* (L) Osbeck) trees were grown in a commercial orchard in

Donna, TX. Leaf samples from symptomatic trees grown in the field and uninfected trees from

an adjacent screenhouse were collected and flash frozen in liquid nitrogen. *C*Las detection was

performed using qPCR with primers that amplify the 16s rDNA (HLBas/HLBr) and the *C.*

sinensis GAPDH gene was used as an internal control. Samples were sent on dry ice to the

Contained Research Facility at the University of California, Davis for further processing.

Protein extraction for Mass Spectrometry

Citrus vascular exudates were enriched by centrifugation (10). Stems were collected from 17-month-old greenhouse plants and the outer bark was manually removed. The inner part of the bark was briefly rinsed with deionized water and dried with Kimwipes. The bark was cut into 1 cm pieces using a razor blade and placed into a 0.65 mL Eppendorf tube with a small hole in the bottom. This tube was then placed into a 1.7 mL Eppendorf collection tube and centrifuged at 12,000 x g for 15 min at 25°C. Vascular extracts were flash frozen and stored at -80°C. Samples were resuspended in 2x Laemmli buffer and quantified for protein concentration using the Pierce 660nm Protein Assay Reagent (Thermo Fisher Scientific).

A total of 85 µg of protein per sample was subjected to one-dimensional SDS-PAGE using an 8-16% precise protein gradient gel (Thermo Fisher Scientific). Proteins were separated for 1 cm in the resolving gel and the entire lane was excised and cut into equal pieces. In-gel reduction was performed by adding 10 mM dithiothreitol in 50 mM ammonium bicarbonate for 30 min at 56°C. Alkylation was performed by adding 55 mM iodoacetamide in 50 mM of ammonium bicarbonate for 20 min in the dark with shaking at room temperature. In-gel tryptic digestion was performed using 300 ng of trypsin in 50 mM ammonium bicarbonate (40). The digested peptides were dried in a vacuum concentrator and then solubilized in 120 µL of 2% acetonitrile, 0.1% trifluoroacetic acid for LC-MS/MS analysis.

LC-MS/MS

Peptides were submitted to the Genomics Center Proteomics Core at the University of California,

Davis for liquid chromatography tandem mass spectrometry (LC)-MS/MS. Peptides were

analyzed by MS as described previously (23). The LC-MS/MS system configuration consisted of

a CTC Pal autosampler (LEAP Technologies) and Paradigm HPLC device (Michrom

BioResources) coupled to a QExactive hybrid quadrupole Orbitrap mass spectrometer (Thermo

Fisher Scientific) with a CaptiveSpray ionization source (Michrom BioResources). Peptides were

reconstituted in 2% acetonitrile and 0.1% formic acid and were washed on a Michrom C18 trap

then were eluted and separated on a Michrom Magic C18AQ (200 μm × 150 mm) capillary

reverse-phase column at a flow rate of 3 μL/min. A 120 min gradient was applied with a 2% to

35% B (100% acetonitrile) for 100 min, a 35% B to 80% B for 7 min and 80% B for 2 min. Then

a decrease of 80% to 5% B in 1 min followed by 98% A (0.1% formic acid) for 10 min. The

QExactive was operated in Data-Dependent Acquisition (DDA) mode with a top-15 method.

Spray voltage was set to 2.2 kV. The scan range was set to 350–1600 m/z, the maximum

injection time was 30 ms and automatic gain control was set to 1×10^6. Precursor resolution was

set to 70,000. For MS/MS, the maximum injection time was 50 ms, the isolation window was 1.6

m/z, the scan range 200–2000 m/z, automatic gain control was set to 5×104 and normalized

collision energy was 27%. The dynamic exclusion window was set to 15 s and fragment product

resolution was 17,500. An intensity threshold of 1×104 was applied and the underfill ratio was

1%.

Peptide identification, analyses and quantification

The raw MS/MS data files were imported into MaxQuant v1.5.1.0 for label-free intensity-based

quantification (41). The database search engine Andromeda (42) was used to search MS/MS

spectra against the *C. clementina* (v1.0) and *C. sinensis* (v1.0) databases downloaded from

Phytozome (43) and the *Candidatus* Liberibacter asiaticus (CLas) proteome database (44)

downloaded from Uniprot with a tolerance level of 20 ppm for the first search and 6 ppm for the

main search. Trypsin/P was set as the enzyme and two missed cleavages were allowed. Protein

N-terminal acetylation and Methionine oxidation were set as variable modifications. The

maximum number of modifications per peptide was set as five and the predefined contaminants

from MaxQuant were included in the search database. The 'match between runs' feature was

checked with a match time window of 0.7 min and an alignment time window of 20 min. The

FDR for protein level and peptide spectrum match was set to 1%. The minimum peptide length

was 6, minimum razor was changed to 0, and minimum unique peptides was set to 1. The

minimum ratio count for protein quantification was set to 2. The other MaxQuant settings were

default. The total peptide intensities for each replicate were summed and a normalization factor

was calculated for each sample (41). This normalization factor was applied based on the least

overall proteome change. Peptide ratios between samples were then calculated to obtain a pair-

wise protein ratio matrix between samples, which was subsequently used to rescale the

cumulative intensity in each sample and provides the label-free intensity (LFQ) value (41).

Parameters used in MaxQuant can be found in Supplementary Table S2. The MaxQuant output

file "Protein Groups" was imported into Perseus 1.5.5.3 (45). Contaminants, reverse hits, and

proteins identified only by modified peptides were excluded in subsequent analysis. The LFQ

intensities were log2-transformed. Proteins not consistently identified in at least two out of the

four replicates in at least one group were discarded. Missing values were imputed with values

from a normal distribution of the obtained intensities using default settings (width 0.5, downshift

1.8). Differentially changing proteins were identified using a two-tailed Student's t-test, $p<0.05$,

with a permutation-based FDR of 0.05 to correct for multiple testing. The mass spectrometry

proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (46)

partner repository with the dataset identifier PXD017480.

Comparative analyses of proteomics data

Initial data visualization was performed using Perseus (Version 1.5.5.3) software. The heatmap

was generated by hierarchical clusters for both individual protein expression and replication was

performed using Euclidean distance (47). Linkage was calculated using preprocessed k-means

with 300 clusters, 10 maximal iterations, and 1 restart. Principle component analyses was

performed using untransformed label-free values.

Comparison of vascular proteomic studies

Homologs of citrus proteins identified by mass spectrometry were identified using BLAST

analyses (version 2.2.18) against *Arabidopsis thaliana* (48, 49) and *Cucumis sativus* (50) phloem

proteins with an E-value of 1e−10 required for protein identification. The Venn diagram was

generated using the vennueler R package (51).

Gene ontology analyses were performed by using Phytozome (43). All identified protein groups

were used as background. Gene ontology enrichment for differentially changing proteins using

the Bonferroni test was used for correction with an alpha-value of 0.05. The p-values and gene

ontology terms can be found in Supplementary Tables S3-S6. A custom R script was designed to

filter redundant GO Terms and plotted in respect to the number of proteins which are

differentially regulated (Github: DanielleMStevens/Franco_2020_Proteomics_Paper) (47, 52).

Raw data and code can be found in the above repository.

Phylogenetic analyses

A total of 134 serine protease and 84 peroxidase sequences were obtained from Phytozome using

the *C. clementina* v1.0 and *C. sinensis* v1.0 databases (43, 53). Peroxidases were identified using

the RedOxiBase Database (http://peroxibase.toulouse.inra.fr/) and the pFAM domain PF00141.

Serine proteases were identified using the MEROPs peptidase database

(https://www.ebi.ac.uk/merops/index.shtml). Multiple sequence alignments were performed with

PRANK (54). A maximum likelihood approach was used to reconstruct phylogenetic trees using

RAxML (55). Bootstrapping was performed with 1,000 replicates. The resulting phylogenetic

trees were visualized using FigTree (version 1.4.3) (56).

Activity Based Protein profiling (ABPP)

Leaves from the *C. sinensis* N-33 and Hamlin varieties (n =4 trees) were ground in Tris-HCl pH

8.0 solution. The supernatant was collected and centrifuged at 5,000 x g for 10 min. Citrus

extracts were quantified using the Pierce 660 Protein assay kit (ThermoFisher Scientific). The

serine protease inhibitor cocktail (50 μM chymotrypsin, 50 μM PMSF, 50 μM of 3,4-

dichloroisocoumarin) was pre-incubated for 30 min prior to the addition of the ActivX TAMRA

fluorophosphonate (FP) serine hydrolase probe (ThermoFisher Scientific). For the no probe

control, 1 μL of dimethyl sulfoxide (DMSO) was added. The serine hydrolase FP probe was

added at a final concentration of 2 μM. Serine proteases were labeled for one hour, labeling

stopped by adding 2X Laemmli buffer, and the sample incubated at 95℃ for 5 min. Labeled

proteins were run on an 8-16% precise Tris-Glycine SDS page gel (ThermoFisher Scientific).

Serine protease activity was visualized by in-gel fluorescence using the Typhoon 8600 variable

mode Imager (Molecular Dynamics Inc, CA, USA). Significant differences in serine protease

activity were detected with a two-tailed unpaired Student's t-test alpha-value of 0.05.

Peroxidase gene expression using RT-qPCR

<u>Hamlin sweet orange:</u> Total RNA was extracted with the RNeasy Plant Mini Kit (Qiagen) and

treated with DNase I (Promega). First-strand cDNA was synthesized from purified RNA with

ImProm-II™ Reverse Transcription System (Promega) and diluted 10 times for RT-qPCR to

detect related genes with specific primers. Quantitative reverse transcription PCR reactions used

the 2xKiCqStart® SYBR® Green qPCR ReadyMix™ (Sigma-Aldrich). PCR consisted of an

initial activation step at 95°C for 3 min, followed by 40 cycles of 95°C for 15 s and 60°C for 40

s. All cDNA samples were run in triplicate. The *C. sinensis* GAPDH gene (XM_006476919.3)

was used to normalize the expression of each target gene and quantification was performed using

the ΔΔCt method. Significant differences (n=4) were detected with a two-tailed unpaired

Student's t-test.

<u>N-33 sweet orange variety:</u> Total RNA was extracted using a Trizol (Invitrogen, USA)-based

method and treated with DNase I (DNase Max Kit (50), QIAGEN). Quantitative reverse

transcription PCR reactions used Power SYBR™ Green RNA-to C_T™ 1-step kit (Applied

Biosystems-Thermo Fisher Scientific, USA). Thermocycling began with a Hold Stage (first step

at 48 °C for 30 s, second step 95 °C for 10 min), followed by a PCR stage (first step at 95°C for

15 s, second step 55 °C for 1 min over 40 cycles) in a thermal cycler (Applied Biosystems-

QuanStudio 3, Thermo Fisher Scientific, USA). The *C. sinensis* GAPDH gene

(XM_006476919.3) was used to normalize the expression of each target gene and quantification

was performed using the ΔΔCt method. Significant differences (n=3) were detected with a two-

tailed unpaired Student's t-test.

Peroxidase activity

Peroxidase activity was tested using a previously described protocol (57). Briefly, citrus

leaf punches (4 mm diameter) were excised and washed for 1 hr in 1x MS solution with

agitation. Each leaf disk was then transferred to a 96 well-microplate containing 50 μL of 1x MS

solution. Leaf disks were incubated for 20 h at 25°C with agitation. To observe peroxidase

accumulation, leaf disks were removed and 50 μL of 1mg/mL of 5-aminosalicylic acid, pH 6.0,

with 0.01% hydrogen peroxide was added and incubated for 1-3 min. The reaction was stopped

by adding 20 μM of 2 N NaOH. Optical density readings (OD_{600}) were measured using a

spectrophotometer. Uninfected samples were set to 100% peroxidase activity and changes in

peroxidase activity were calculated based on the ratio of OD_{600} readings (Infected/Uninfected).

Significant differences in peroxidase activity were detected with a two-tailed unpaired Student's

t-test.

Results

***C*Las induces global changes in the citrus vascular proteome**

In order to investigate changes occurring in citrus tissues infected with *C*Las, we isolated

protein from bark samples ten months post-inoculation and performed proteomics analyses. The

sweet orange cultivar Washington navel (*Citrus sinensis* (L) Osbeck), which is susceptible to

HLB, was graft inoculated or mock inoculated with *C*Las positive budwood. At 10 months post-

inoculation, *C*Las inoculated navels exhibited severe HLB disease symptoms and detectable

*C*Las titers using qPCR (Supplementary Table S7). Total extracted proteins were subjected to

tandem mass spectrometry (LC-MS/MS) and MS1 quantification using MaxQuant (41). To

increase total protein identifications, the protein database included the *C. sinensis* and *C.*

clementina assembled genomes (53, 58). A total of 1,354 citrus proteins were identified using

MS1 quantification for infected and uninfected samples (Supplementary Table 8, Figure 1A).

Uninfected samples had an average of 1259 protein identifications and infected samples had an

average of 1152 protein identifications across all four biological replicates (trees). 329 (24%)

proteins were significantly upregulated and 363 (27%) proteins were downregulated in infected

samples (Supplementary Table 9 and 10, respectively). Hierarchical clustering based on

Euclidean distance revealed that samples clustered according to treatment (Fig. 1A) (45).

Similarly, principle component analyses (PCA) differentiated biological samples based on their

treatment (Fig. 1B.). Taken together, these data indicate that *C*Las infection induces global and

reproducible changes in vascular protein expression patterns (Fig 1A-B).

Isolation of pure phloem proteins is extremely difficult and can only be obtained using

microneedle puncture or the stylet of a phloem-feeding insect (59, 60). To increase phloem

yields, EDTA-facilitated exudation and centrifugation are commonly used (10). However, these

approaches are not amenable to citrus exhibiting HLB symptoms due to decreased phloem

transport. Therefore, we centrifuged the outer bark to extract vascular exudates and enrich for

phloem proteins. To ascertain if we enriched for phloem proteins, we compared the overlap in

protein identifications between our study with studies performed in *Cucumis sativus* and

Arabidopsis (48–50). Citrus homologs for 62% and 64% of the *Arabidopsis* and *Cucumis*

sativus proteins were identified, respectively (Fig. 1C.). Although these plants represent diverse

families (Rutaceae, Brassicaceae and Cucurbitaceae), many of the identified proteins overlapped.

These results indicate that vascular proteins were successfully extracted from citrus and identify

universal vascular proteins between diverse plant species.

To gain insight on the biological and molecular functions altered by *C*Las infection, we

performed gene ontology analyses using Phytozome (43). Processes involved in plant growth

and development including metabolism, translation and biosynthetic processes were inhibited

after *C*Las infection (Fig. 2). For example, many ribosomal proteins and translation elongation

factors were downregulated during infection. Proteins involved in glycolysis, such as pyruvate

kinases (orange1.1g036921m, orange1.1g010627m, Ciclev10031317m, orange1.1g010449m)

were significantly repressed during infection (Supplementary Table 10). In contrast, proteins

involved in pathogen perception were upregulated in infected samples. We identified multiple

proteins with receptor-like kinase or receptor-like protein domain architecture that were

upregulated during infection, including the receptor-like kinase, Ciclev10030933m, and the

LysM domain protein orange1.1g018290m (Supplementary Table 9). Proteins with this domain

architecture are frequently implicated in pathogen perception (27, 28, 61). Multiple peroxidases

and proteases were more abundant in *C*Las infected samples (Fig 2A, 2B). Both peroxidases and

proteases have been established as key components in development and plant defense responses

(62–65) . Taken together these results implicate the tradeoff between growth and defense during

*C*Las infection.

Peroxidase transcript and protein abundance increases during *C*Las infection

In order to gain a greater understanding of citrus peroxidases, we identified and classified

peroxidases in the *C. sinensis* and *C. clementina* genomes. In plants, the heme peroxidase-

catalase superfamily (PCATS) can be further classified into three distinct classes with plants

possessing only class I and class III peroxidases (66). Both class I and class III peroxidases have

been implicated to function in growth, development, and immunity (67). Class I peroxidases

encompass intracellular enzymes such as ascorbate peroxidases and catalases. Class III

peroxidases are a large family consisting of extracellular or vacuolar localized enzymes. Class III

peroxidases can regulate H_2O_2 abundance, oxidation of toxic substrates, and defense responses

(65, 68, 69). Due to redundancy, studies to identify the *in planta* role of most peroxidases

remains elusive (69). Using PeroxiBase, 80 peroxidases (13 class I, 67 class III) were identified

in *C. sinensis.* For phylogenetic analyses, an additional 4 peroxidases (2 class I, 2 class III)

identified from *C. clementina* were included (67) (Fig 3). Maximum-likelihood phylogenetic

analyses of these peroxidases were performed using RAxML (55). The identified ascorbate

peroxidases cluster appropriately within the class I clade (Fig 3). In the vascular proteome, 22

peroxidases were identified and 10 were upregulated during *C*Las infection (3 class I and 7 class

III) (Supplemental Figure 1, Supplementary Table 11). Taken together, these data highlight the

abundance of peroxidases in *C. sinensis* and reveal an overall increase in abundance of specific

members upon *C*Las infection.

The abundance of specific enzyme classes does not always correlate with their activity.

To test peroxidase protein accumulation and enzymatic activity during *C*Las infection, we

examined the *C. sinensis* Hamlin sweet orange variety, which is the most widely grown in

Florida for early harvest (70). Healthy and symptomatic leaf tissue from Hamlin oranges were

sampled one-year after *C*Las transmission by *D. citri* (Fig 4A). To test peroxidase gene

expression in Hamlin, qPCR was performed on seven peroxidase genes that were previously

identified as induced in our proteomics analyses. Four Hamlin peroxidases were induced in

infected samples and three were expressed at the same level in infected and uninfected samples

by qPCR (Fig 4B). We investigated peroxidase activity during *C*Las infection using a

microplate-based assay to quantify the basal levels of the reactive oxygen species, H_2O_2 (57).

Overall peroxidase activity decreased in infected Hamlin samples (Fig 4C). The disconnect

between peroxidase mRNA expression and enzymatic activity may be due to *C*Las manipulation

of enzymatic activity or opposing roles of specific enzymes.

N-33 is a Navel orange variety frequently grown in Texas (71). To test peroxidase

accumulation under field conditions and with another *C. sinensis* variety, mature N-33 plants

were sampled for peroxidase accumulation by qPCR and enzymatic activity. Symptomatic leaves

and healthy leaves from an adjacent screenhouse were collected. To validate *C*Las infection,

qPCR was performed on seven peroxidases identified in our proteomics analyses as described

above (Fig 4D). The transcript abundance of all analyzed peroxidases in N-33 was higher in

infected samples, with five demonstrating a statistically significant increase (Fig 4E).

Furthermore, peroxidase enzymatic activity was significantly higher in N-33 infected leaf tissue

(Fig 4F). These data highlight the responsiveness of individual peroxidases during *C*Las

infection and indicate that the response to *C*Las can be influenced by plant genotype and/or

environmental effects.

Diverse proteases and protease inhibitors dynamically change during infection

Proteolysis is an enriched biological process and molecular function in the mass spectrometry quantification of vascular proteins from *C*Las-infected citrus (Fig 2A, Fig 2B). Plant proteases are divided into seven classes: serine, aspartic, cysteine, asparagine, threonine, glutamate, and metalloproteases (72). Proteases degrade misfolded and toxic proteins. This function plays important regulatory roles in development, reproduction and immunity (73). To regulate proteolysis, protease inhibitors target their corresponding proteases. In the proteome of vascular exudates, both proteases and protease inhibitors were present and dynamically responded to infection. A total of 22 serine, 15 aspartic, 12 cysteine and 2 metalloproteases were identified (Fig 5A).

Serine proteases are among the largest protease class in plants encompassing over 200 members and are divided into 14 families (73, 74). The classification of serine proteases depends on the catalytic mechanism and common ancestry (75). Members of the subtilases (S8), serine carboxypeptidases (S10), and lysosomal Pro-x carboxypeptidases (S28) families of serine proteases were identified in our proteomics analyses (Fig 5F, Supplementary Fig. 2). Phylogenetic analyses of 67 S8, 48 S10, and 19 S28 serine proteases from *C. sinensis* and *C. clementina* were performed (72). . Differentially accumulating proteases did not cluster phylogenetically (Supplementary Fig. 2). S10 proteases generally increased during infection except for one member, orange1.1g010508 (Fig 5F). Subtilisin and serine carboxypeptidases comprise the largest serine protease class, have had prior implications in plant immunity, and are directly inhibited by diverse pathogen classes (73–77). These results indicate that the S8, S10, and S28serine proteases families are most responsive to *C*Las infection.

In general, protease inhibitors inhibit a single catalytic class of proteases, however, some

families can inhibit multiple protease classes (78). Cystatins are simple protease inhibitors that

target papain like cysteine proteases (PLCPs) and metacaspase-like cysteine proteases. Cystatins

have dual roles as defense proteins and regulators of PLCP protein turnover (79). In this

proteomic study, five cystatins were identified, but only one decreased in abundance during

infection (Fig 5A-B). Diverse serine protease inhibitors were identified in the vasculature (Fig

5A-B). Two potato type I (Pin1) serine protease inhibitors were identified, but not changing

during CLas infection (Supplementary Table 12). Kunitz trypsin inhibitors and SERine Protease

INhibitors (SERPINs) are the most extensively studied and can inhibit diverse protease classes

(78, 80). Kunitz trypsin inhibitors are capable of binding to serine, cysteine and aspartic

proteases (78). SERPINs are active serine protease inhibitors, but can also inhibit some cysteine

proteases (78). Nine Kunitz trypsin inhibitors were identified (five induced) and two SERPINs

were identified (one induced) (Fig 5A, B). Kunitz trypsin inhibitors and SERPINs have been

demonstrated to have inhibitory effects against pathogens and insects (78). Our proteomics

analyses indicate protease activity is tightly regulated during CLas infection.

Serine protease activity dynamically changes during CLas infection

To quantify the activity of serine proteases during CLas infection, activity-based protein

profiling (ABPP) was performed. ABPP enables detection of active proteases with probes

specific to protease class (81, 82). For ABPP, we used the Hamlin and N-33 varieties grown in

the Florida greenhouse and Texas field, respectively. Proteases from four uninfected and four

infected samples were extracted from leaf tissue and subjected to ABPP (Fig 6A). A no probe

control was included to confirm probe binding to active proteases. Due to a lack of an effective

broad-spectrum serine protease inhibitor, an inhibitor cocktail containing phenylmethylsulphonyl

fluoride (PMSF), 3,4-dichloroisocoumarine and chymostatin was used to verify labeling

specificity. The inhibitor cocktail was added 30 min prior to the addition of the probe to

inactivate serine proteases. Serine protease activity is demonstrated by the presence of a band

detected by fluorescence imaging. Active serine protease bands were reduced in the presence of

the protease inhibitor cocktail, indicating that ABPP was able to specifically detect serine

protease activity (Fig 6B, D).

In Hamlin orange, three distinct protease signals changed during CLas infection (Fig 6B).

The intensity of bands 1 (~110 kD) and 2 (~100 kD) increased during infection and

quantification of the fluorescent signal across four biological replicates demonstrated this

increase in activity as statistically significant (Fig 6C). In contrast, band 3 (~30 kD) displayed a

significant decrease in protease activity during CLas infection. In N-33, bands 1-3 followed the

same trend (Fig 6D, E). Two additional groups of active serine protease bands were observed in

N-33 (Fig 6D). Due to the inability to resolve individual active serine protease bands, the

average signal was summed in the two different aggregates (bands 4 and 5). Bands 4 and 5

exhibited decreased activity (~40kD) (Fig 6E). These experiments demonstrate the induction and

activation of specific serine proteases during CLas infection is conserved across diverse

environmental conditions and sweet orange varieties. The decreased activity of specific serine

proteases during CLas infection may be due to the induction of plant serine protease inhibitors or

direct suppression by CLas.

Discussion:

HLB has caused billions of dollars in economic losses worldwide and continues to threaten the United States citrus industry. To curb HLB losses, a detailed understanding of the mechanisms of disease is essential to develop comprehensive disease management strategies. Various -omics approaches have identified molecular pathways altered during *C*Las infection in leaf, root, stem, and fruit tissues providing a heightened understanding of the responses in diverse tissue types (31, 35, 83–85). For example, proteomics coupled with nutrient profiling demonstrated that *C*Las infection reduces Ca^{+2}, Mg^{+2}, Fe^{+2}, Mn^{+4}, Zn^{+2} and Cu^{+} in grapefruit leaves (86). Consistent with these observations, diseased plants supplemented with a cocktail of Ca and Zn resulted in an increase in fruit production (87). However, these studies did not directly address responses within the phloem, the niche where *C*Las resides and propagates. In this study, we quantified proteins in vascular enriched samples to gain a greater understanding of proteomic changes where *C*Las resides.

In Washington navel, we revealed global changes occur in vascular tissue, with 51% of the total proteome dynamically changing during *C*Las infection. Upon pathogen perception, cellular resources, which would otherwise be utilized in growth-related processes, are redirected towards defense responses; this is known as the growth-defense tradeoff (88). In this study, we observed evidence for this tradeoff during *C*Las infection, with defense associated proteins induced in infected samples and processes involved in general housekeeping significantly enriched in uninfected samples. In a parallel study on Washington navel leaf tissue, processes involved in photosynthesis were downregulated while defense related processes were significantly upregulated in *C*Las infected samples (83). These data indicate that the growth-defense tradeoff also occurs in response to vascular pathogens. Washington navel is susceptible

to HLB, but clearly responded to *C*Las infection. It is unclear if HLB symptomology is due to a

lack of early and robust plant defense, or a result of a slower prolonged autoimmune response, or

possibly a combination of the two.

Reactive oxygen species (ROS) such as superoxide radicals, hydrogen peroxide and

singlet oxygen play a central role in plant pathogen defense (89). ROS is produced at the plasma

membrane by NAD(P)H oxidases and apoplastic peroxidases (89). ROS also accumulates in

different organelles including the chloroplast, mitochondria, and peroxisomes (89). Despite the

importance of ROS for defense signaling, an overabundance is cytotoxic and therefore requires

tight control (89). In melon, redox-related proteins, including peroxidases, are phloem localized

and induced during viral infection (48, 90). In this study, the transcription of seven peroxidases

that were more abundant in proteomics analyses, were also upregulated in infected samples from

different locations and citrus varieties. ROS regulating enzymes are upregulated at the protein

level in grapefruit after *C*Las infection (86, 91). Citrus undergoing thermotherapy to treat HLB

exhibited increased peroxidase accumulation (92). Additionally, bioinformatic analyses

comparing four RNA-seq studies on HLB illustrated that tolerance is linked with the induction of

redox-related genes involved in both young and mature leaf tissues (93). These studies indicate

the importance of regulating the opposing roles of redox during CLas infection.

Plant proteases are involved in pathogen perception, defense priming, signaling, and

programmed cell death (77). To ensure tight control of protease activity, plants evolved protease

inhibitors that fine-tune protease activity. Protease inhibitors can also inhibit digestive proteases

from herbaceous insects and target pathogens (60, 78, 94, 95). Multiple classes of proteases and

protease inhibitors are found in the vasculature of diverse plant species (32, 94–98). In

Arabidopsis, tomato, and maize, papain-like cysteine proteases (PLCPs) positively regulate

immunity and defense hormone accumulation (77, 99, 100). In citrus, PLCPs are localized in

phloem-enriched tissue, immune related, and targeted by the conserved *C*Las effector, SDE1

(23). In this study, proteases and their inhibitors were the largest class of dynamically changing

proteins during *C*Las infection. In Hamlin and N-33 sweet orange varieties, we observed

activation and inhibition of specific citrus serine proteases, suggesting citrus fine-tunes protease

enzymatic activity during *C*Las infection. Genotype specific effects were also observed, with

specific groups of serine proteases inhibited in the N-33 variety, but not in the Hamlin variety.

Whether this inhibition is due to the endogenous activity of plant protease inhibitors or *C*Las

inhibition of citrus proteases needs further investigation.

Comparative analyses of different citrus genotypes are beginning to reveal the

importance of genotype specific responses. RNA-seq analyses between the more HLB-tolerant

rough lemon (*Citrus jambhiri*) and susceptible sweet orange (*C. sinensis* (L) Osbeck)

demonstrated that the expression amplitude of defense related genes was much higher in the

HLB tolerant-rough lemon than HLB-susceptible sweet orange (85). In two susceptible

genotypes, Lisbon lemon and Washington navel, the root metabolome and microbiome differed

in response to *C*Las. (101). However, both genotypes exhibited a reduction of sugars, the amino

acids proline and asparagine, and in some metabolites (101). These important findings are not

entirely unexpected as sweet oranges and lemons are different species and were derived from

different admixtures (102). In this study, we observed differences in peroxidase and serine

enzymatic activity in two sweet orange varieties, Hamlin and N-33 (101). This study reveals

even closely related genotypes within the same citrus species exhibit variations in their responses

to *C*Las. Future studies investigating diversity of citrus responses to HLB across different citrus

genotypes, *C*Las isolates, and environmental conditions will provide a comprehensive

understanding of disease development.

Acknowledgements

This research was supported by a USDA National Institute of Food and Agriculture grant #2016-

70016-24833 to G.C., V.A., and N.W and grant # 2019-70016-29796 awarded to G.C., and N.W.

Additional support was provided by the Citrus Research Board (CRB) project 5200-157. J.Y.F

was supported by the USDA National Institute of Food and Agriculture grant #2017-678011-

26039. T.W.H.L. was supported by a Rubicon grant of the Netherlands Organization for

Scientific Research (NWO).

Author Contributions

G.C. and J.YF. conceived and designed the study; J.Y.F, S.P.T., Z.P, F.B.G. and performed the

experiments. S.P.T., T.W.H.L and D.M.S. performed the bioinformatic analyses. J.Y.F. and

G.C. wrote the manuscript with input from all authors.

Data Availability

Proteomic data are available via ProteomeXchange with identifier PXD017480.

Gene Ontology analysis pipeline is available via Github:

(https://github.com/DanielleMStevens/Franco_2020_Proteomics_Paper)

Competing Financial Interests

The authors declare no competing financial interest.

References

1. Li, J., Li, L., Pang, Z., Kolbasov, V. G., Ehsani, R., Carter, E. W., and Wang, N. (2019) Developing citrus huanglongbing (HLB) management strategies based on the severity of symptoms in HLB-endemic citrus-producing regions. *Phytopathology*. **109**, 582–592

2. Deng, H., Achor, D., Exteberria, E., Yu, Q., Du, D., Stanton, D., Liang, G., and Gmitter, F. G. (2019) Phloem regeneration is a mechanism for huanglongbing-tolerance of "bearss" lemon and "LB8-9" sugar belleopenspisupspi®closespisupspi mandarin. *Front. Plant Sci.* **10**, 277

3. Ramadugu, C., Keremane, M. L., Halbert, S. E., Duan, Y. P., Roose, M. L., Stover, E., and Lee, R. F. (2016) Long-term field evaluation reveals Huanglongbing resistance in Citrus relatives. *Plant Dis.* **100**, 1858–1869

4. da Graça, J. V., Douhan, G. W., Halbert, S. E., Keremane, M. L., Lee, R. F., Vidalakis, G., and Zhao, H. (2016) Huanglongbing: An overview of a complex pathosystem ravaging the world's citrus. *J. Integr. Plant Biol.* **58**, 373–387

5. Dai, Z., Wu, F., Zheng, Z., Yokomi, R., Kumagai, L., Cai, W., Rascoe, J., Polek, M., Chen, J., and Deng, X. (2019) Prophage diversity of 'candidatus liberibacter asiaticus' strains in California. *Phytopathology*. **109**, 551–559

6. McRoberts, N., Figuera, S. G., Olkowski, S., McGuire, B., Luo, W., Posny, D., and Gottwald, T. (2019) Using models to provide rapid programme support for California's efforts to suppress Huanglongbing disease of citrus. *Philos. Trans. R. Soc. B Biol. Sci.* **374**, 1–11

7. Hodges, A. W., and Spreen, T. H. (2015) Economic Impacts of Citrus Greening -(HLB) in Florida, 2006/07-2010/11. Univ Florida IFAS Ext., 7–12.

8. United States Department of Agriculture (USDA), N. A. S. S. (2017) Citrus Maturity Test Results and Fruit Size.

9. Hung, T. H., Hung, S. C., Chen, C. N., Hsu, M. H., and Su, H. J. (2004) Detection by PCR of Candidatus Liberibacter asiaticus, the bacterium causing citrus huanglongbing in vector psyllids: Application to the study of vector-pathogen relationships. *Plant Pathol.* **53**, 96–102

10. Hijaz, F., and Killiny, N. (2014) Collection and chemical composition of phloem sap from Citrus sinensis L. Osbeck (sweet orange). *PLoS One*. **9**, e101830

11. Skryhan, K., Gurrieri, L., Sparla, F., Trost, P., and Blennow, A. (2018) Redox regulation of starch metabolism. *Front. Plant Sci.* **9**, 1–8

12. Koh, E. J., Zhou, L., Williams, D. S., Park, J., Ding, N., Duan, Y. P., and Kang, B. H. (2012) Callose deposition in the phloem plasmodesmata and inhibition of phloem transport in citrus leaves infected with "Candidatus Liberibacter asiaticus." *Protoplasma*. **249**, 687–697

13. Hu, Y., Zhong, X., Liu, X., Lou, B., Zhou, C., and Wang, X. (2017) Comparative transcriptome analysis unveils the tolerance mechanisms of Citrus hystrix in response to

606  "Candidatus Liberibacter Asiaticus" infection. *PLoS One*. **12**, e0189229

607 14. Wang, N., and Trivedi, P. (2013) Citrus Huanglongbing: A Newly Relevant Disease
608   Presents Unprecedented Challenges. *Phytopathology*. **103**, 652–665

609 15. Bové, J. M. (2006) Huanglongbing: A destructive, newly-emerging, century-old disease of
610   citrus. *J. Plant Pathol.* **88**, 7–37

611 16. Dodds, P. N., and Rathjen, J. P. (2010) Plant immunity: towards an integrated view of
612   plant–pathogen interactions. *Nat. Rev. Genet.* **11**, 539–548

613 17. Cong, Q., Kinch, L. N., Kim, B. H., and Grishin, N. V. (2012) Predictive sequence
614   analysis of the Candidatus liberibacter asiaticus proteome. *PLoS One*. **7**, e41071

615 18. Yan, Q., Sreedharan, A., Wei, S., Wang, J., Pelz-Stelinski, K., Folimonova, S., and Wang,
616   N. (2013) Global gene expression changes in Candidatus Liberibacter asiaticus during the
617   transmission in distinct hosts between plant and insect. *Mol. Plant Pathol.* **14**, 391–404

618 19. Sugio, A., Kingdom, H. N., MacLean, A. M., Grieve, V. M., and Hogenhout, S. A. (2011)
619   Phytoplasma protein effector SAP11 enhances insect vector reproduction by manipulating
620   plant development and defense hormone biosynthesis. *Proc. Natl. Acad. Sci.* **108**, E1254–
621   E1263

622 20. Prasad, S., Xu, J., Zhang, Y., and Wang, N. (2016) SEC-translocon dependent
623   extracytoplasmic proteins of Candidatus liberibacter asiaticus. *Front. Microbiol.* **7**, 1–9

624 21. Thapa, S.P.;De Francesco, A; Trinh, J; Gurung, F.B.; Pang, Z; Vidalakis, G; Wang, N;
625   Ancona, V; Ma, W; Coaker, G. (2020) Genome-wide analyses of Liberibacter species
626   provides insights into evolution, phylogenetic relationships and virulence factors. *Mol.*
627   *Plant Pathol.* **00**, 1–16

628 22. Pitino, M., Armstrong, C. M., Cano, L. M., and Duan, Y. (2016) Transient expression of
629   Candidatus Liberibacter asiaticus effector induces cell death in Nicotiana benthamiana.
630   *Front Plant Sci*. **7**, 982

631 23. Clark, K., Franco, J. Y., Schwizer, S., Pang, Z., Hawara, E., Liebrand, T. W. H.,
632   Pagliaccia, D., Zeng, L., Gurung, F. B., Wang, P., Shi, J., Wang, Y., Ancona, V., Van Der
633   Hoorn, R. A. L., Wang, N., Coaker, G., and Ma, W. (2018) An effector from the
634   Huanglongbing-associated pathogen targets citrus proteases. *Nat. Commun.* **9**, 1718

635 24. Jain, M., Munoz-Bodnar, A., Zhang, S., and Gabriel, D. W. (2018) A secreted 'candidatus
636   liberibacter asiaticus' peroxiredoxin simultaneously suppresses both localized and
637   systemic innate immune responses in planta. *Mol. Plant-Microbe Interact.* **12**, 1312–1322

638 25. Wang, N., Pierson, E. A., Setubal, C., Xu, J., Levy, J. G., Zhang, Y., Li, J., Rangel, L. T.,
639   and Jr, J. M. (2017) Liberibacter – Host Interface : Insights into Pathogenesis Mechanisms
640   and Disease Control. *Annu. Rev. Phytopathol.* **55**, 1–32

641 26. Li, J., Pang, Z., Trivedi, P., Zhou, X., Ying, X., Jia, H., and Wang, N. (2017) "Candidatus
642   Liberibacter asiaticus" encodes a functional salicylic acid (SA) hydroxylase that degrades
643   SA to suppress plant defenses. *Mol. Plant-Microbe Interact.* **30**, 620–630

27. Zhang, J., and Zhou, J. M. (2010) Plant immunity triggered by microbial molecular
signatures. *Mol. Plant*. **3**, 783–793

28. Dalio, R. J. D., Magalhaes, D. M., Rodrigues, C. M., Arena, G. D., Oliveira, T. S., Souza-
Neto, R. R., Picchi, S. C., Martins, P. M. M., Santos, P. J. C., Maximo, H. J., Pacheco, I.
S., De Souza, A. A., and Machado, M. A. (2017) PAMPs, PRRs, effectors and R-genes
associated with citrus-pathogen interactions. *Ann. Bot.* **120**, 617

29. Jain, M., Fleites, L. A., and Gabriel, D. W. (2015) Prophage-encoded peroxidase in
"candidatus liberibacter asiaticus" is a secreted effector that suppresses plant defenses.
Mol. Plant-Microbe Interact. **28**, 1330–7

30. Liao, H. L., and Burns, J. K. (2012) Gene expression in Citrus sinensis fruit tissues
harvested from huanglongbing-infected trees: Comparison with girdled fruit. *J. Exp. Bot.*
63, 3307–19

31. Fan, J., Chen, C., Yu, Q., Brlansky, R. H., Li, Z. G., and Gmitter, F. G. (2011)
Comparative iTRAQ proteome and transcriptome analyses of sweet orange infected by
"Candidatus Liberibacter asiaticus." *Physiol. Plant.* **143**, 235–245

32. Albrecht, U., and Bowman, K. D. (2012) Transcriptional response of susceptible and
tolerant citrus to infection with Candidatus Liberibacter asiaticus. *Plant Sci.* **185**–**186**,
118–130

33. Albrecht, U., and Bowman, K. D. (2008) Gene expression in Citrus sinensis (L.) Osbeck
following infection with the bacterial pathogen Candidatus Liberibacter asiaticus causing
Huanglongbing in Florida. *Plant Sci.* **175**, 291–306

34. Kim, J.-S., Sagaram, U. S., Burns, J. K., Li, J.-L., and Wang, N. (2009) Response of sweet
orange (Citrus sinensis) to "Candidatus Liberibacter asiaticus" infection: microscopy and
microarray analyses. *Phytopathology*. **99**, 50–57

35. Zhong, Y., Cheng, C. Z., Jiang, N. H., Jiang, B., Zhang, Y. Y., Wu, B., Hu, M. L., Zeng, J.
W., Yan, H. X., Yi, G. J., and Zhong, G. Y. (2015) Comparative transcriptome and
iTRAQ proteome analyses of citrus root responses to candidatus liberibacter asiaticus
infection. *PLoS One*. **10**, e0126973

36. Aritua, V., Achor, D., Gmitter, F. G., Albrigo, G., and Wang, N. (2013) Transcriptional
and Microscopic Analyses of Citrus Stem and Root Responses to Candidatus Liberibacter
asiaticus Infection. *PLoS One*. **8**, e73742

37. Etxeberria, E., Gonzalez, P., Achor, D., and Albrigo, G. (2009) Anatomical distribution of
abnormally high levels of starch in HLB-affected Valencia orange trees. *Physiol. Mol.*
Plant Pathol. **74**, 76–83

38. Li, W., Hartung, J. S., and Levy, L. (2006) Quantitative real-time PCR for detection and
identification of Candidatus Liberibacter species associated with citrus huanglongbing. *J.*
Microbiol. Methods. **66**, 104–15

39. Trivedi, P., Sagaram, U. S., Kim, J. S., Brlansky, R. H., Rogers, M. E., Stelinski, L. L.,
Oswalt, C., and Wang, N. (2009) Quantification of viable Candidatus Liberibacter
asiaticus in hosts using quantitative PCR with the aid of ethidium monoazide (EMA). *Eur.*

684 *J. Plant Pathol.* **124**, 553–563

40. Shevchenko, A., Tomas, H., Havliš, J., Olsen, J. V., and Mann, M. (2006) In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nat. Protoc.* **1**, 2856–60

41. Cox, J., and Mann, M. (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* **26**, 1367–1372

42. Cox, J., Neuhauser, N., Michalski, A., Scheltema, R. A., Olsen, J. V., and Mann, M. (2011) Andromeda: A peptide search engine integrated into the MaxQuant environment. *J. Proteome Res.* **10**, 1794–1805

43. Goodstein, D. M., Shu, S., Howson, R., Neupane, R., Hayes, R. D., Fazo, J., Mitros, T., Dirks, W., Hellsten, U., Putnam, N., and Rokhsar, D. S. (2012) Phytozome: A comparative platform for green plant genomics. *Nucleic Acids Res.* **40**, D1178-86

44. Duan, Y., Zhou, L., Hall, D. G., Li, W., Doddapaneni, H., Lin, H., Liu, L., Vahling, C. M., Gabriel, D. W., Williams, K. P., Dickerman, A., Sun, Y., and Gottwald, T. (2009) Complete genome sequence of citrus huanglongbing bacterium, "Candidatus Liberibacter asiaticus" obtained through metagenomics. *Mol. Plant. Microbe. Interact.* **22**, 1011–1020

45. Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M. Y., Geiger, T., Mann, M., and Cox, J. (2016) The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat. Methods*. **13**, 731–740

46. Perez-Riverol, Y., Csordas, A., Bai, J., Bernal-Llinares, M., Hewapathirana, S., Kundu, D. J., Inuganti, A., Griss, J., Mayer, G., Eisenacher, M., Pérez, E., Uszkoreit, J., Pfeuffer, J., Sachsenberg, T., Yilmaz, Ş., Tiwary, S., Cox, J., Audain, E., Walzer, M., Jarnuczak, A. F., Ternent, T., Brazma, A., and Vizcaíno, J. A. (2019) The PRIDE database and related tools and resources in 2019: Improving support for quantification data. *Nucleic Acids Res.* **47**, D442–D450

47. Gu, Z., Eils, R., and Schlesner, M. (2016) Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics*. **32**, 2847–9

48. Carella, P., Wilson, D. C., Kempthorne, C. J., and Cameron, R. K. (2016) Vascular sap proteomics: Providing insight into long-distance signaling during stress. *Front. Plant Sci.* **7**, 1–8

49. Carella, P., Merl-Pham, J., Wilson, D. C., Dey, S., Hauck, S. M., Vlot, A. C., and Cameron, R. K. (2016) Comparative proteomics analysis of phloem exudates collected during the induction of systemic acquired resistance. *Plant Physiol.* **171**, 1495–510

50. Huang, S., Li, R., Zhang, Z., Li, L., Gu, X., Fan, W., Lucas, W. J., Wang, X., Xie, B., Ni, P., Ren, Y., Zhu, H., Li, J., Lin, K., Jin, W., Fei, Z., Li, G., Staub, J., Kilian, A., Van Der Vossen, E. A. G., Wu, Y., Guo, J., He, J., Jia, Z., Ren, Y., Tian, G., Lu, Y., Ruan, J., Qian, W., Wang, M., Huang, Q., Li, B., Xuan, Z., Cao, J., Asan, Wu, Z., Zhang, J., Cai, Q., Bai, Y., Zhao, B., Han, Y., Li, Y., Li, X., Wang, S., Shi, Q., Liu, S., Cho, W. K., Kim, J. Y., Xu, Y., Heller-Uszynska, K., Miao, H., Cheng, Z., Zhang, S., Wu, J., Yang, Y., Kang, H.,

Li, M., Liang, H., Ren, X., Shi, Z., Wen, M., Jian, M., Yang, H., Zhang, G., Yang, Z., Chen, R., Liu, S., Li, J., Ma, L., Liu, H., Zhou, Y., Zhao, J., Fang, X., Li, G., Fang, L., Li, Y., Liu, D., Zheng, H., Zhang, Y., Qin, N., Li, Z., Yang, G., Yang, S., Bolund, L., Kristiansen, K., Zheng, H., Li, S., Zhang, X., Yang, H., Wang, J., Sun, R., Zhang, B., Jiang, S., Wang, J., Du, Y., and Li, S. (2009) The genome of the cucumber, Cucumis sativus L. *Nat. Genet.* **41**, 1275–81

51. Wilkinson, L. (2012) Exact and approximate area-proportional circular venn and euler diagrams. *IEEE Trans. Vis. Comput. Graph.* **18**, 321–31

52. Gu, Z., Gu, L., Eils, R., Schlesner, M., and Brors, B. (2014) Circlize implements and enhances circular visualization in R. *Bioinformatics*. **30**, 2811–2

53. Wu, G. A., Prochnik, S., Jenkins, J., Salse, J., Hellsten, U., Murat, F., Perrier, X., Ruiz, M., Scalabrin, S., Terol, J., Takita, M. A., Labadie, K., Poulain, J., Couloux, A., Jabbari, K., Cattonaro, F., Del Fabbro, C., Pinosio, S., Zuccolo, A., Chapman, J., Grimwood, J., Tadeo, F. R., Estornell, L. H., Muñoz-Sanz, J. V., Ibanez, V., Herrero-Ortega, A., Aleza, P., Pérez-Pérez, J., Ramón, D., Brunel, D., Luro, F., Chen, C., Farmerie, W. G., Desany, B., Kodira, C., Mohiuddin, M., Harkins, T., Fredrikson, K., Burns, P., Lomsadze, A., Borodovsky, M., Reforgiato, G., Freitas-Astúa, J., Quetier, F., Navarro, L., Roose, M., Wincker, P., Schmutz, J., Morgante, M., MacHado, M. A., Talon, M., Jaillon, O., Ollitrault, P., Gmitter, F., and Rokhsar, D. (2014) Sequencing of diverse mandarin, pummelo and orange genomes reveals complex history of admixture during citrus domestication. *Nat. Biotechnol.* **32**, 656–62

54. Löytynoja, A. (2014) Phylogeny-aware alignment with PRANK. *Methods Mol. Biol.* **1079**, 155–70

55. Stamatakis, A. (2006) RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics*. **22**, 2688–90

56. Rambaut, A. (2016) FigTree v1.4.3. *Mol. Evol. phylogenetics Epidemiol.* Available at: http://tree.bio.ed.ac.uk/software/figtree/.

57. Adam Mott, G., Desveaux, D., and Guttman, D. S. (2018) A high-sensitivity, microtiter-based plate assay for plant pattern-triggered immunity. *Mol. Plant-Microbe Interact.* **31**, 499–504

58. Xu, Q., Chen, L. L., Ruan, X., Chen, D., Zhu, A., Chen, C., Bertrand, D., Jiao, W. B., Hao, B. H., Lyon, M. P., Chen, J., Gao, S., Xing, F., Lan, H., Chang, J. W., Ge, X., Lei, Y., Hu, Q., Miao, Y., Wang, L., Xiao, S., Biswas, M. K., Zeng, W., Guo, F., Cao, H., Yang, X., Xu, X. W., Cheng, Y. J., Xu, J., Liu, J. H., Luo, O. J., Tang, Z., Guo, W. W., Kuang, H., Zhang, H. Y., Roose, M. L., Nagarajan, N., Deng, X. X., and Ruan, Y. (2013) The draft genome of sweet orange (Citrus sinensis). *Nat. Genet.* **45**, 59–66

59. Kennedy, J. S., and Mittler, T. E. (1953) A Method of obtaining Phloem Sap via the Mouth-parts of Aphids. *Nature*. **171**, 528

60. Dinant, S., Bonnemain, J. L., Girousse, C., and Kehr, J. (2010) Phloem sap intricacy and interplay with aphid feeding. *Comptes Rendus - Biol.* **333**, 504–15

764   61.   Zipfel, C. (2014) Plant pattern-recognition receptors. *Trends Immunol.* **35**, 345–51

765   62.   Misas-Villamil, J. C., van der Hoorn, R. A. L., and Doehlemann, G. (2016) Papain-like
766   cysteine proteases as hubs in plant immunity. *New Phytol.* **212**, 902–907

767   63.   Jashni, M. K., Mehrabi, R., Collemare, J., Mesarich, C. H., and de Wit, P. J. G. M. (2015)
768   The battle in the apoplast: further insights into the roles of proteases and their inhibitors in
769   plant–pathogen interactions. *Front. Plant Sci.* **6**, 1–7

770   64.   Passardi, F., Cosio, C., Penel, C., and Dunand, C. (2005) Peroxidases have more functions
771   than a Swiss army knife. *Plant Cell Rep.* **24**, 255–65

772   65.   Almagro, L., Gómez Ros, L. V., Belchi-Navarro, S., Bru, R., Ros Barceló, A., and
773   Pedreño, M. A. (2009) Class III peroxidases in plant defence reactions. *J. Exp. Bot.* **60**,
774   377–90

775   66.   Pandey, V. P., Awasthi, M., Singh, S., Tiwari, S., and Dwivedi, U. N. (2017) A
776   Comprehensive Review on Function and Application of Plant Peroxidases. *Biochem.*
777   *Anal. Biochem.* **6**, 1–16

778   67.   Passardi, F., Theiler, G., Zamocky, M., Cosio, C., Rouhier, N., Teixera, F., Margis-
779   Pinheiro, M., Ioannidis, V., Penel, C., Falquet, L., and Dunand, C. (2007) PeroxiBase: The
780   peroxidase database. *Phytochemistry*. **68**, 1605–11

781   68.   Hiraga, S., Sasaki, K., Ito, H., Ohashi, Y., and Matsui, H. (2001) A large family of class
782   III plant peroxidases. *Plant Cell Physiol.* **42**, 462–8

783   69.   Cosio, C., and Dunand, C. (2009) Specific functions of individual class III peroxidase
784   genes. *J. Exp. Bot.* **60**, 391–408

785   70.   Putnam, A. H., and Hudson, M. E. (2018) Florida Citrus Statistics 2016-2017. *Florida*
786   *Department of Agriulture and Consumer Services.* Available at:
787   https://www.nass.usda.gov/

788   71.   Miller, C. S. I. D. (2017) Texas Citrus Nursery Stock Certification Manual TEXAS
789   DEPARTMENT OF AGRICULTURE. Agriculture and Consumer Protection Division,
790   Environmental and Biosecurity Programs, Citrus Program. Available at:
791   https://www.texasagriculture.gov/

792   72.   Rawlings, N. D., Barrett, A. J., Thomas, P. D., Huang, X., Bateman, A., and Finn, R. D.
793   (2018) The MEROPS database of proteolytic enzymes, their substrates and inhibitors in
794   2017 and a comparison with peptidases in the PANTHER database. *Nucleic Acids Res.* **46**,
795   D624–D632

796   73.   van der Hoorn, R. A. L. (2008) Plant Proteases: From Phenotypes to Molecular
797   Mechanisms. *Annu. Rev. Plant Biol.* **59**, 191–223

798   74.   Sueldo, D., Ahmed, A., Misas-Villamil, J., Colby, T., Tameling, W., Joosten, M. H. A. J.,
799   and Van der Hoorn, R. A. L. (2014) Dynamic hydrolase activities precede hypersensitive
800   tissue collapse in tomato seedlings. *New Phytol.* **203**, 913–925

75.   Di Cera, E. (2009) Serine proteases. *IUBMB Life*. 10.1002/iub.186

76. Figueiredo, J., Sousa Silva, M., and Figueiredo, A. (2018) Subtilisin-like proteases in
plant defence: the past, the present and beyond. *Mol. Plant Pathol.* **19**, 1017–1028

77. Thomas, E. L., and van der Hoorn, R. A. L. (2018) Ten prominent host proteases in plant-
pathogen interactions. *Int. J. Mol. Sci.* **19**, E639

78. Rustgi, S., Boex-Fontvieille, E., Reinbothe, C., von Wettstein, D., and Reinbothe, S.
(2018) The complex world of plant protease inhibitors: Insights into a Kunitz-type
cysteine protease inhibitor of Arabidopsis thaliana. *Commun. Integr. Biol.* **11**, e1368599

79. Martinez, M., Cambra, I., Carrillo, L., Diaz-Mendoza, M., and Diaz, I. (2009)
Characterization of the entire cystatin gene family in barley and their target cathepsin L-
like cysteine-proteases, partners in the hordein mobilization during seed germination.
Plant Physiol. **151**, 1531–45

80. Cohen, M., Davydov, O., and Fluhr, R. (2019) Plant serpin protease inhibitors: Specificity
and duality of function. *J. Exp. Bot.* **70**, 2077–2085

81. van der Hoorn, R. A. L. (2004) Activity Profiling of Papain-Like Cysteine Proteases in
Plants. *PLANT Physiol.* **135**, 1170–8

82. Liu, Y., Patricelli, M. P., and Cravatt, B. F. (1999) Activity-based protein profiling: The
serine hydrolases. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 14694–9

83. Chin, E. L., Ramsey, J. S., Mishchuk, D. O., Saha, S., Foster, E., Chavez, J. D., Howe, K.,
Zhong, X., Polek, M., Godfrey, K. E., Mueller, L., Bruce, J. E., Heck, M., and Slupsky, C.
M. (2019) Longitudinal transcriptomic, proteomic, and metabolomic analyses of Citrus
sinensis (L.) Osbeck graft-inoculated with 'Candidatus Liberibacter asiaticus.' *J.*
Proteome Res. **19**, 719–732

84. Wang, Y., Zhou, L., Yu, X., Stover, E., Luo, F., and Duan, Y. (2016) Transcriptome
profiling of huanglongbing (HLB) tolerant and susceptible citrus plants reveals the role of
basal resistance in HLB tolerance. *Front. Plant Sci.* **7**, 1–13

85. Yu, Q., Chen, C., Du, D., Huang, M., Yao, J., Yu, F., Brlansky, R. H., and Gmitter, F. G.
(2017) Reprogramming of a defense signaling pathway in rough lemon and sweet orange
is a critical element of the early response to Candidatus Liberibacter asiaticus. *Hortic. Res.*
4, 1–15

86. Nwugo, C. C., Lin, H., Duan, Y., and Civerolo, E. L. (2013) The effect of "Candidatus
Liberibacter asiaticus" infection on the proteomic profiles and nutritional status of pre-
symptomatic and symptomatic grapefruit (Citrus paradisi) plants. *BMC Plant Biol.* **13**, 1–
24

87. Khairulmazmi Ahmad (2011) Field assessment of calcium, copper and zinc ions on plant
recovery and disease severity following infection of Huanglongbing (HLB) disease.
African J. Microbiol. Res. **5**, 4967–4979

88. Huot, B., Yao, J., Montgomery, B. L., and He, S. Y. (2014) Growth-defense tradeoffs in
plants: A balancing act to optimize fitness. *Mol. Plant.* **7**, 1267–1287

89. Apel, K., and Hirt, H. (2004) REACTIVE OXYGEN SPECIES: Metabolism, Oxidative

Stress, and Signal Transduction. *Annu. Rev. Plant Biol.* **55**, 373–99

90. Serra-Soriano, M., Navarro, J. A., Genoves, A., and Pallás, V. (2015) Comparative
proteomic analysis of melon phloem exudates in response to viral infection. *J. Proteomics.*
124, 11–24

91. Martinelli, F., Reagan, R. L., Dolan, D., Fileccia, V., and Dandekar, A. M. (2016)
Proteomic analysis highlights the role of detoxification pathways in increased tolerance to
Huanglongbing disease. *BMC Plant Biol.* **16**, 167

92. Nwugo, C. C., Doud, M. S., Duan, Y. ping, and Lin, H. (2016) Proteomics analysis
reveals novel host molecular mechanisms associated with thermotherapy of 'Ca.
Liberibacter asiaticus'-infected citrus plants. *BMC Plant Biol.* **16**, 253

93. Balan, B., Ibáñez, A. M., Dandekar, A. M., Caruso, T., and Martinelli, F. (2018)
Identifying host molecular features strongly linked with responses to huanglongbing
disease in citrus leaves. *Front. Plant Sci.* **9**, 1–13

94. Kehr, J. (2006) Phloem sap proteins: Their identities and potential roles in the interaction
between plants and phloem-feeding insects. *J. Exp. Bot.* **57**, 767–74

95. Walz, C., Giavalisco, P., Schad, M., Juenger, M., Klose, J., and Kehr, J. (2004)
Proteomics of curcurbit phloem exudate reveals a network of defence proteins.
Phytochemistry. **65**, 1795–804

96. Petzold, H. E., Zhao, M., and Beers, E. P. (2012) Expression and functions of proteases in
vascular tissues. *Physiol. Plant.* **145**, 121–9

97. Hu, C., Ham, B. K., El-shabrawi, H. M., Alexander, D., Zhang, D., Ryals, J., and Lucas,
W. J. (2016) Proteomics and metabolomics analyses reveal the cucurbit sieve tube system
as a complex metabolic space. *Plant J.* **87**, 442–454

98. Hao, G., Pitino, M., Duan, Y., and Stover, E. (2016) Reduced Susceptibility to
Xanthomonas citri in Transgenic Citrus Expressing the FLS2 Receptor From Nicotiana
benthamiana. *Mol. Plant-Microbe Interact.* **29**, 132–42

99. Ilyas, M., Hörger, A. C., Bozkurt, T. O., Van Den Burg, H. A., Kaschani, F., Kaiser, M.,
Belhaj, K., Smoker, M., Joosten, M. H. A. J., Kamoun, S., and Van Der Hoorn, R. A. L.
(2015) Functional Divergence of Two Secreted Immune Proteases of Tomato. *Curr. Biol.*
25, 2300–2306

100. Lozano-Torres, J. L., Wilbers, R. H. P., Gawronski, P., Boshoven, J. C., Finkers-
Tomczak, A., Cordewener, J. H. G., America, A. H. P., Overmars, H. A., Van 't Klooster,
J. W., Baranowski, L., Sobczak, M., Ilyas, M., van der Hoorn, R. A. L., Schots, A., de
Wit, P. J. G. M., Bakker, J., Goverse, A., and Smant, G. (2012) Dual disease resistance
mediated by the immune receptor Cf-2 in tomato requires a common virulence target of a
fungus and a nematode. *Proc. Natl. Acad. Sci.* **109**, 10119–10124

101. Padhi, E. M. T., Maharaj, N., Lin, S. Y., Mishchuk, D. O., Chin, E., Godfrey, K., Foster,
E., Polek, M., Leveau, J. H. J., and Slupsky, C. M. (2019) Metabolome and microbiome
signatures in the roots of citrus affected by huanglongbing. *Phytopathology.* **109**, 2022–
2032

102.   Wu, G. A., Terol, J., Ibanez, V., López-García, A., Pérez-Román, E., Borredá, C.,
882   Domingo, C., Tadeo, F. R., Carbonell-Caballero, J., Alonso, R., Curk, F., Du, D.,
883   Ollitrault, P., Roose, M. L., Dopazo, J., Gmitter, F. G., Rokhsar, D. S., and Talon, M.
884   (2018) Genomics of the origin and evolution of Citrus. *Nature*. **554**, 311–316
885
886

Fig. 1. *C*Las induces global changes in protein expression in citrus vascular exudates **(A)**
Heatmap of proteins identified in citrus sap exudates. Vascular proteins were enriched and
extracted from mock and *C*Las-graft inoculated Washington navel oranges (*C. sinensis* (L)
Osbeck) and subjected to label-free MS1 quantification. Identified proteins are shown in rows,
normalized by their mean value, and the Z-score was calculated for hierarchical clustering using
Euclidean distance. Columns represent biological replicates (individual trees). Red indicates
proteins where expression was increased, and blue indicates decreased protein expression
relative to the mean. **(B)** Principal component analyses (PCA) was performed on untransformed
label free quantification values in Perseus. **(C)** Comparison of phloem proteins identified in
diverse plant species. A total of 1354 proteins from Washington navel identified from this study
were compared with previous phloem proteomics studies in *Arabidopsis thaliana* and cucumber

899 (*Cucumis sativus*) phloem proteins using BLAST with an E-value of 1e-10 as a cutoff. The venn

900 diagram indicates protein overlap and was generated using the vennueler R package.

Fig. 2. Gene ontology (GO) analyses of citrus proteomics. (A) The biological processes and **(B)** molecular function of 692 significantly changing proteins identified by proteomics analyses of enriched vascular proteins from Washington navel ten months post-graft inoculation with *C*Las compared to mock inoculated controls. GO analyses and molecular function was performed using Phytozome 12 (https://phytozome.jgi.doe.gov/). Bonferroni was used for test correction with an FDR rate of <0.05. GO term slimming was performed manually by removing redundant terms used for the same group of proteins. Blue = enriched in downregulated proteins after graft inoculation with *C*Las; Red = enriched in upregulated proteins after graft inoculation with *C*Las.

Fig. 3. Phylogenetic analyses reveal dynamic changes in specific citrus class I and III peroxidases after inoculation with *C*Las. A total of 15(13, *C. sinensis,* (*Cs*) and 2 *C. clementina,*(*Cc*)) Class I (yellow) and 78 (76 *Cs ,* 2 Cc) Class III (green) peroxidases were identified from the citrus genome using the PF00141 Pfam domain and the RedOxiBase database. Two *Arabidopsis thaliana* sequences were used to guide clustering, one class I Peroxidase (AT1G07890.3) and one class III peroxidase (AT1G05250.1). Accession numbers starting with orange1.1g indicate proteins identified from the *C. sinensis* (navel) genome while those starting with Ciclev are those identified from the *C. clementina* (clementine) genome. Redundant sequences from *Citrus clementina* were removed from the phylogeny. A maximum likelihood approach was used to generate the phylogeny with 1000 bootstrap replicates. The resulting phylogeny was visualized using FigTree. A total of 22 peroxidases were

930   identified by mass spectrometry in Washington Navel. Red indicates peroxidases that were

931   significantly induced by mass spectrometry in *C*Las inoculated Washington navel and grey

932   indicates peroxidases that were detected, but not significantly changing after inoculation.

Fig 4. Peroxidase activity and transcript accumulation depend on citrus genotype. Two sweet orange varieties were inoculated with *C*Las using *Diaphorina citri* and analyzed for peroxidase (POX) enzymatic activity and transcript abundance. **(A)** HLB symptoms on representative Hamlin leaves (top) and corresponding Ct value for *C*Las (bottom) one year after feeding with *C*Las positive and negative *D. citri*. Citrus plants were grown under greenhouse conditions. *C*Las titers were detected by qPCR using primers targeting the β-operon (GenBank: AY342001.1). **(B)** Select peroxidase (POX) transcripts increase during infection in Hamlin. Hamlin trees were inoculated as described in (A) and RNA was extracted from the leaves. POXs that were significantly induced by proteomics analyses in Washington navel after CLas inoculation were analyzed for transcriptional expression using qPCR. Transcripts were normalized to GAPDH-C (NCBI Rseq: XM_006476919.3). Graphs show means ±SD, n= 4

trees. Significant differences were detected with a two-tailed unpaired Student's t-test, alpha-value of 0.05. **(C)** POX enzymatic activity decreases during infection in Hamlin. Hamlin trees were inoculated as described in (A) and POX activity was detected on leaf punches surrounding the midrib one-year post-inoculation from the same leaves used for RNA extraction in (B). Graphs show means ±SD, n=12 samples taken from 4 individual trees. Significant differences were detected with a two-tailed unpaired Student's t-test, $p<0.01$. **(D)** HLB symptoms on representative N-33 leaves (top) and corresponding Ct value for CLas (bottom). Citrus plants were grown in a Texas field where CLas was naturally transmitted by *D. citri*. Uninfected leaf samples were taken from plants grown in an adjacent screenhouse. CLas titers were detected by qPCR primers targeting the 16s rDNA (GenBank: L22532). **(E)** Select peroxidase transcripts increase during infection in N-33. Leaf samples were taken from trees described in (D) and RNA was extracted from leaf tissue. POXs that were significantly changing in Washington navel proteomic analyses were selected and analyzed as described in (C). Graphs show mean ±SD, n=3 trees. Significant differences were detected with a two-tailed unpaired Student's t-test, Asterisks represent p values, *=<0.05, **=<0.01, ***=<0.001. Experiments were repeated three times with similar results. **(F)** POX activity increases during infection in N-33. Leaf punches from N-33 leaf samples taken from trees described in (D). Graphs show mean ±SD, n=8 taken from 3 individual trees. Significant differences were detected with a two tailed unpaired Student's t-test, $p<0.001$.

Fig 5. There is a dynamic interplay between the abundance of proteases and protease inhibitors during *C*Las infection. (A) Total number of proteases and protease inhibitors identified by proteomics analyses of enriched vascular proteins from Washington navel ten months post-graft inoculation with *C*Las compared to mock inoculated controls. Four proteases classes comprising: 22 serine proteases, 15 aspartic proteases, 12 cysteine proteases, and 2 metalloproteases were identified. Three protease inhibitor classes were identified: 5 cystatins, 12 serine protease inhibitors (9 Kunitz trypsin inhibitors and 4 SERPINs). Black indicates proteins that were significantly induced by mass spectrometry in *C*Las inoculated Washington navel and grey indicates proteins that were detected, but not significantly changing after inoculation. **(B-F)** Abundance of specific protease inhibitors and proteases that exhibit differential accumulation by mass spectrometry. Protein abundance is depicted as Log_2(MS1) intensity. Abundance in mock

977   inoculated is displayed in black and *C*Las inoculated in grey. Graphs indicate means ±SD, n=4

978   trees. Significant differences were detected by two-tailed t-test with an FDR of <0.5. * = $p<0.05$,

979   ** $p<0.01$, *** $p<0.001$.

Fig 6. The activity of serine proteases dynamically changes during *C*Las infection. (A)
Experimental Strategy. Proteases from uninfected and infected citrus trees were extracted and
subjected to activity-based protein profiling (ABPP) using the fluorescently labeled probe,
ActivX TAMRA fluorophosphonate (FP) serine hydrolase probe, to capture serine proteases.
Visualization of active serine protease bands was performed with the Typhoon 8500 scanner and
signals were quantified in Image J. Two sweet orange genotypes were inoculated with *C*Las
using *Diaphorina citri* and analyzed for serine protease enzymatic activity. **(B)** ABPP of serine
protease activity in leaves of the Hamlin variety one year after feeding with *C*Las positive and
negative *D. citri.* Active serine proteases were labeled using FP-tamara (lanes 2-10). A serine
protease inhibitor cocktail was pre-incubated 30 min prior to the addition of FP-tamara to
demonstrate probe binding specificity (lane 2 compared to lane 3). A no probe control was added

as a negative control (lane 1, band is the 70kD marker from the protein ladder). Fluorescent

signals corresponding to serine protease activity are indicated on the right with an asterisk. **(C)**

Quantification of labeled band intensity in (B) using ImageJ. Graphs indicate means ±SD, n=4

trees. Significant differences were detected by two-tailed t-test, * = alpha-value of 0.5. **(D)** ABPP

of serine protease activity in leaves of the N-33 variety grown under field conditions. Gel loading

is the same as described in (B). Signals corresponding to active serine proteases are indicated by

an asterisk. Signals unable to be resolved were grouped together and indicated by brackets. **(E)**

Quantification of labeled signal intensity in (C) using ImageJ. Graphs indicate means ±SD, n=4

trees. Significant differences were detected by two-tailed t-test, * = alpha-value of 0.05