Abundantly expressed class of non-coding RNAs conserved through the multicellular evolution of dictyostelid social amoebae

Jonas Kjellin[1], Lotta Avesson[2,3], Johan Reimegård[4], Zhen Liao[1,5], Ludwig Eichinger[6], Angelika Noegel[6], Gernot Glöckner[6], Pauline Schaap[7], and Fredrik Söderbom[1]

[1]Department of Cell and Molecular Biology, Uppsala University, Box 596 Uppsala, S-75124 Sweden,

[2]Department of Molecular Biology, Biomedical Center, Swedish University of Agricultural Sciences, Box 590, S-75124 Uppsala, Sweden, [4]Department of Cell and Molecular Biology, National Bioinformatics Infrastructure Sweden, Science for Life Laboratory, Uppsala University, Box 596 Uppsala, S-75124 Sweden, [6]Centre for Biochemistry, Institute of Biochemistry I, Medical Faculty, University of Cologne, Cologne, Germany and [7]College of Life Sciences, University of Dundee, Dundee DD1 5EH, United Kingdom.

[3]Present address: Novo Nordisk Foundation Center for Protein Research, University of Copenhagen, Blegdamsvej 3B, 2200 Copenhagen

[5]Present address: Department of Plant Biology, Swedish University of Agricultural Sciences, Box 7080 Uppsala, S-750 07 Sweden

Corresponding author. fredrik.soderbom@icm.uu.se. Phone: 46 184714901

Abstract

Background: Aggregative multicellularity has evolved multiple times in diverse groups of eukaryotes. One of the most well-studied examples is the development of dictyostelid social amoebae, e.g. *Dictyostelium discoideum*. However, it is still poorly understood why multicellularity emerged in these amoebae while the great majority of other members of Amoebozoa are unicellular. Previously a novel type of non-coding RNA, Class I RNAs, was identified in *D. discoideum* and demonstrated to be important for normal multicellular development. In this study we investigated Class I RNA evolution and its connection to multicellular development.

Results: New Class I RNA genes were identified by constructing a co-variance model combined with a scoring system based on conserved up-stream sequences. Multiple genes were predicted in representatives of each major group of Dictyostelia and expression analysis validated that our search approach can identify expressed Class I RNA genes with high accuracy and sensitivity. Further studies showed that Class I RNAs are ubiquitous in Dictyostelia and share several highly conserved structure and sequence motifs. Class I RNA genes appear to be unique to dictyostelid social amoebae since they could not be identified in searches in outgroup genomes, including the closest known relatives to Dictyostelia.

Conclusion: Our results show that Class I RNA is an ancient abundant class of ncRNAs, likely to have been present in the last common ancestor of Dictyostelia dating back at least 600 million years. Taken together, our current knowledge of Class I RNAs suggests that they may have been involved in evolution of multicellularity in Dictyostelia.

Keywords (3-10): social amoeba, *Dictyostelium discoideum*, non-coding RNA, evolution, Class I RNA, dictyostelids, multicellularity, Dictyostelia

Background

The role of RNA goes far beyond it being an intermediate transmitter of information between DNA and protein, in the form of messenger (m)RNAs. This has been appreciated for a long time for some non-coding RNAs (ncRNAs), such as transfer (t)RNAs, ribosomal (r)RNAs, small nuclear (sn)RNAs, and small nucleolar (sno)RNAs. Today, we know that ncRNAs are involved in regulating most cellular processes and the advent of high-throughput sequencing technologies have facilitated the identification of numerous different classes of ncRNAs [1]. These regulatory RNAs vary greatly in size from 21-24 nucleotides (nt), e.g. micro (mi)RNAs and small interfering (si)RNAs, to several thousands of nucleotides, such as long non-coding (lnc)RNAs. Several classes of ncRNAs are ubiquitously present in all domains of life while others are specific to certain evolutionary linages, contributing to their specific characteristics. This can be exemplified by ncRNAs in Metazoa, where an increase in number of ncRNAs, e.g. miRNAs, is associated with increased organismal complexity and is believed to be essential for the evolution of metazoan multicellularity [2]. Multicellularity in plants and animals is achieved by clonal division and development originating from a single cell. This is in contrast to aggregative multicellularity, were cells stream together to form multicellular structures upon specific environmental changes. Aggregative multicellularity has evolved independently multiple times and is found both among eukaryotes and prokaryotes [3–10]. The complexity of the aggregative multicellular life stages varies for different organisms, but they all share the transition from unicellularity to coordinated development upon environmental stress, e.g. starvation, which eventually leads to formation of fruiting bodies containing cysts or spores [3]. Probably the most well-studied aggregative multicellularity is the development of the social amoeba *Dictyostelium discoideum* belonging to the group Dictyostelia within the supergroup Amoebozoa. Dictyostelia is a monophyletic group estimated to date back at least 600 million years [11], which is similar to the age of Metazoa [12]. Dictyostelia is currently divided into four major groups (Group 1-4) where all members share the ability of transition from uni- to multicellularity upon starvation [13]. However, the complexity

of the development and the morphology of the fruiting bodies varies between different dictyostelids,

where the highest level of multicellular complexity is found among group 4 species, which includes *D.*

discoideum [14, 15]. Recently a new taxonomy was proposed for many dictyostelids [16]. As this new

taxonomy has not yet been fully adopted by the research community, we choose to use the previous

designations throughout this study (old and new names, including NCBI accession numbers, are

summarized in Additional file 1).

Well-annotated genome sequences are available for representative species of all four major groups of

Dictyostelia [11, 17–20] and multiple draft genome sequences are available for additional dictyostelids.

This has allowed for comparative genomics, which has provided information about the protein coding

genes that are important for the diversification of Dictyostelia from other amoebozoans. Comparison

between genomes has also given insight into the genes required for the evolution of the distinct

morphological characteristics, which define each group [11, 17, 18, 20, 21]. However, evolution of

complex traits such as multicellularity in Dictyostelia as well as other eukaryotic groups, cannot solely be

explained by the appearance of novel genes but also relies on an increased ability to regulate pre-existing

genes and their products so that they can function in novel genetic networks [20, 22]. This is also

supported by the major transcriptional reprogramming during multicellular development in *D. discoideum*

[23].

D. discoideum harbors several classes of developmentally regulated ncRNAs with regulatory potential, e.g.

microRNAs [24–27], long non-coding RNAs [28] and long antisense RNAs [28, 29]. In addition, a large part

of the ncRNA repertoire of *D. discoideum* is constituted by Class I RNAs, originally identified in full length

cDNA libraries [30]. So far, Class I RNAs have only been validated in *D. discoideum* [30, 31], but they have

also been computationally predicted in *Dictyostelium purpureum* [18]. Both species belong to the same

evolutionary group of Dictyostelia, i.e. group 4 [13, 16]. In *D. discoideum,* Class I RNAs are 42-65 nt long

and are expressed at high levels from a large number of genes. Members of Class I RNAs are characterized

by a short stem structure, connecting the 5' and 3' ends, and a conserved 11 nt sequence motif adjacent

to the 5´part of the stem. The remainder of the RNA is variable both in sequence and structure. Class I

RNAs mainly localize to the cytoplasm [30] where one of the RNAs has been shown to associate with four

different proteins of which at least one, the RNA recognition motifs (RRM) containing protein CIBP,

directly binds to the Class I RNA [31]. Furthermore, the Class I RNAs appear to be involved in regulating

multicellular development. This is based on the observations that Class I RNAs are developmentally

regulated and that cells where a single Class I RNA gene has been knocked out show aberrant early

development [30, 31].

In this study we investigated the prevalence of Class I RNAs within Dictyostelia as well as in other

organisms with the aim to understand if Class I RNAs are restricted to dictyostelids and perhaps required

for their aggregative multicellularity. Based on the known Class I RNAs from *D. discoideum*, we know that

the major part of the RNA is variable and hence sequence-based searches alone, such as BLAST, would

not reliably identify new genes. This was solved by constructing a Class I classifier based on a co-variance

model, which includes both sequence and structure information, combined with a scoring system for

conserved up-stream sequence motifs, e.g. promoter motifs. Using this search approach, we identified

approximately 300 Class I RNA genes predicted to be expressed in the genomes of 16 different

dictyostelids. In addition, we validate the expression of ~100 Class I RNAs from six different species, using

both northern blot and RNA-seq, which supports the high accuracy and sensitivity of our search approach

to identify expressed Class I RNAs. Comparative studies of identified Class I RNA loci show several well

conserved features, like stem forming properties connecting the 5' and 3' ends, as well as preserved

sequence motifs. Importantly, Class I RNAs appear to be specific to Dictyostelia as no Class I RNA genes

were identified in genomes of organisms outside this group of social amoebae, including their closest

known relatives or organisms exhibiting different kinds of multicellularity. Taken together, this unique

class of ncRNAs constitute a very large number of conserved highly expressed genes involved in the

evolution of Dictyostelia multicellular development.

Results

Co-variance model identifies Class I RNA genes in evolutionary distinct groups of Dictyostelia social

amoebae.

We have previously identified and characterized Class I RNAs from the social amoeba *D. discoideum*.

Members of Class I RNAs are characterized by a short stem-structure connecting the 5' and 3' ends and a

conserved 11 nt sequence motif adjacent to the 5´part of the stem. The remainder of the Class I RNAs are

highly variable both in sequence and structure (Fig. 1a). This class of ncRNA have so far only been validated

in *D. discoideum* where it is associated with development [30, 31], but has also been predicted in *D.*

purpureum [18].

The presence of Class I RNA genes in two different dictyostelids and the fact that at least one Class I RNA

member is involved in controlling early multicellular development, led us to hypothesize that this class of

ncRNA may be a general effector for early development in all members of dictyostelid social amoebae.

Hence, it may have been present in the last common ancestor of Dictyostelia, dating back approximately

600 million years in evolution. In order to investigate this, we used the complete and well-annotated

genome sequences for representatives of each major group of Dictyostelia, i.e. *D. discoideum* [17]*, D.*

purpureum [18]*, Dictyostelium lacteum* [20]*, Polysphondylium pallidum* [11]*, Acytostelium subglobosum*

[19] and *Dictyostelium fasciculatum* [11] (Fig. 1b). Class I RNAs cannot be reliably detected by sequence

searches alone due to the high sequence variability. Therefore, we constructed a co-variance model (CM)

with Infernal [32] where both sequence and secondary structure information of 34 *D. discoideum* Class I

RNAs were taken into account (see Materials and methods for details). The initial CM search of the six

Dictyostelia genomes, followed by manual inspection of the results, indicated the presence of Class I RNAs

in all major groups of Dictyostelia (Additional file 2: Fig. S1). These candidates scored ≥ 25 in the CM search

and had the potential to form a short stem similar to the *D. discoideum* Class I RNAs. In order to improve

the CM, these candidates (CM score ≥ 25 and potential to form stem) were added to the CM followed

by new genome searches. This process was repeated until no new candidates fulfilling the criteria were

identified after which a final search with increased sensitivity was performed (see Materials and methods).

In total, 126 loci distributed over all major groups of Dictyostelia were identified including 36 of the 40

published *D. discoideum* Class I RNAs [30, 31] and all the 26 previously predicted *D. purpureum* Class I loci

[18] (Additional file 2: Fig. S1).

Refining the search for Class I RNA genes using conserved promoter elements

Many *D. discoideum* ncRNA genes have an upstream putative promoter element, DUSE (*Dictyostelium*

upstream sequence element), which in most cases is situated ~60 nt from the transcriptional start site

(TSS) [33]. However, for *D. discoideum* Class I RNAs, DUSE is often found further upstream. In these cases,

a TGTG-box (AAATGTG) is located ~60 nt downstream of DUSE while the distance from the start of the

mature RNA varies. Whether the TGTG-box is an additional promotor element or the TSS of a precursor

transcript is currently not known. DUSE appears to be conserved within group 4 of Dictyostelia, as it was

also identified ~60 nt in front of the predicted *D. purpureum* Class I RNAs [18]. In order to investigate the

presence of conserved upstream motifs in the rest of Dictyostelia, we searched for enriched motifs in the

150 nt upstream sequence of all the 126 Class I RNA gene candidates identified in the CM search.

Intriguingly, DUSE like motifs could be identified ~60 nt upstream of the predicted start for the majority

of the Class I RNA gene candidates (73 of 126) in all organisms. In contrast, the TGTG-box was only found

in a subset (21 of 126) of the upstream sequences of which all belonged to *D. discoideum* Class I RNA

genes (Fig. 1c). As both sequence and distance of DUSE appeared to be conserved in all major groups of

Dictyostelia, we used this information to create a scoring system anticipating accurate prediction of

expressed Class I RNA genes. (Fig. 1d). First the score produced by the CM search (Infernal) was used and

all candidates with a score ≥ 15 were included in order to capture more divergent Class I RNA genes. Next

we scored the presence and location of DUSE and TGTG-box 150 nt upstream of the candidates identified

in the CM search based on the motif identification program FIMO [34]. Lack of DUSE and/or non-canonical

distance from predicted TSS or TGTG-box was penalized with negative scores. Taken together, a total

score of 32 could be achieved if a high-scoring DUSE was identified at the predicted distance upstream of

a candidate Class I RNA gene with the lowest allowed Infernal score (≥ 15). Based on this, all candidates

scoring 32 or higher were classified as Class I RNA loci. Using this approach, we predicted 18-39 Class I

RNAs (146 in total) for each of the six dictyostelids investigated (see below).

Class I RNAs of predicted sizes are expressed at high levels in all four groups of Dictyostelia

Based on the Class I classifier, we predicted Class I RNA genes in all dictyostelids included in the CM build.

But are all these genes really expressed and how accurate are the size predictions? From our previous

studies, we know that Class I RNAs in *D. discoideum* are expressed at high levels at vegetative growth and

are readily detected by northern blot [30, 31]. Hence, we used the same approach to validate a subset of

randomly chosen candidates that made the total score threshold in *D. purpureum* (Group 4)*, D. lacteum*

(Group 3)*, P. pallidum* (Group 2A)*, A. subglobosum* (Group 2B) and *D. fasciculatum* (Group 1). RNA was

prepared from vegetative growing amoebae and specific Class I RNAs were analyzed by northern blot (Fig.

2a). For *D. purpureum*, we probed for DpuR-7, predicted to be 54 nt long, resulting in a strong signal. In

addition, we designed two probes predicted to recognize six different 85 nt long RNAs (DpuR-X) and the

majority (24/30) of Class I RNAs (DpuR-Y), respectively. As expected, probing for DpuR-X resulted in one

band on the northern blot. For DpuR-Y, we expected signals from several Class I RNAs to overlap due to

similar/identical sizes but we could still detect several distinct bands within the expected size range. In *D.*

fasciculatum, we probed for one Class I RNA predicted to be 62 nt long (DfaR-4) while at least two

candidates were probed for in the other organisms, i.e. *D: lacteum:* DlaR-1 (61 nt) and DlaR-5 (54 nt), *P.*

pallidum: PpaR-1/2 (59/62 nt) and PpaR-9 (58 nt), and *A. subglobosum*: AsuR-13/14 (61 nt) and AsuR-10

(82 nt). Distinct bands were detected for each Class I RNA candidate and the sizes matched the predictions

well although the northern results often indicated that the RNAs were a few nt longer than predicted (Fig.

2a and see below). The larger (but much weaker) signals observed for AsuR-13/14 and PpaR-9 are likely

cross hybridizations to longer Class I RNAs. Taken together, the results confirm that Class I RNAs are

conserved and expressed in all four groups of Dictyostelia social amoebae.

Classifier accurately predicts expressed Class I RNAs in all major groups of Dictyostelia

Next, we performed RNA-seq on *D. discoideum, D. lacteum, P. pallidum* and *D. fasciculatum* representing

each major group of Dictyostelia. RNA was prepared from growing cells as well as two multicellular life-

stages, i.e. mound and slug/finger stages, to increase our chances to also detect Class I RNAs that are only

expressed at specific life stages. Expression was evaluated based on the read count and coverage over all

loci identified in the CM search (Infernal score ≥ 15) as exemplified in Additional file 2: Fig. S2. Strikingly,

expression both during vegetative growth and development could be confirmed for almost all Class I RNA

candidates identified by the classifier (total score ≥ 32) (Fig. 2b, Additional file 3). Next, we calculated

receiver operator characteristics (ROC) curves in order to evaluate the classifier performance and

investigate if it improves Class I RNA identification compared to CM search alone. ROC curves were

generated for *D. discoideum, D. lacteum, P. pallidum* and *D. fasciculatum* individually (Additional file 2:

Fig. S3) as well as for the pooled data (Fig. 2c) based on the RNA-seq validation and either CM search score

(≥ 25) or classifier score (≥ 32). Evaluation of the two search approaches show an increase in both

sensitivity and accuracy of prediction for the classifier, i.e. when the promoter (DUSE) presence and

distance were included in the classification of Class I RNA gene candidates. To summarize, the classifier

reliably detects expressed Class I RNAs in all the tested dictyostelids with almost no false positives.

Conserved features of Dictyostelia Class I RNA

The wealth of newly identified Class I RNA genes in six different and evolutionary separated dictyostelids

allowed us to construct a general/unifying picture of Class I RNAs. This will also be of importance when

searching for Class I RNA genes in other species in order to track down the birth of this class of ncRNAs in

evolution (see below).

Class I transcription is dependent on DUSE

Both the sequence of DUSE and its upstream location is highly conserved in all of the analyzed amoebae

(Fig. 3a). In the group 1 and 4 dictyostelids, DUSE contains three consecutive C residues, while two

consecutive C's are found in the majority of the DUSE of *A. subglobosum* (Group 2B) and *D. lacteum* (Group

3) and in all *P. pallidum* (Group 2A). The RNA-seq data show that the promoter element is essential for

transcription as it is found in front of all expressed Class I RNA genes. Further strengthening its importance

is the observation that high scoring Class I RNAs (both considering score produced by classifier or CM

alone) lacking DUSE at the correct up-stream location are not expressed (Additional file 3). The TGTG-box,

situated 60 nt down-stream of the DUSE and in front of the predicted TSS, is only found in *D. discoideum*

suggesting that this is a rather late addition in the evolution of Class I RNA genes. The genes are most

likely transcribed by RNA polymerase III (Pol III) as the majority of the Class I RNAs from the different

dictyostelids exhibit a stretch of at least four consecutive T-residues downstream of the predicted end of

transcription, which is a common Pol III termination signal [36].

5' and 3' ends of Class I RNAs are conserved

The RNA-seq analyses showed that the *D. discoideum* Class I RNAs started at the predicted (and previously

defined [30]) G residue while the majority of group 1-3 Class I RNAs started one to two nucleotides

upstream of the conserved G (Additional file 2: Fig. S2). When we compared all loci, we noticed that the

two nucleotides preceding the completely conserved G residue, are highly conserved A-residues in all six

species investigated. The difference in 5' ends of the mature Class I RNAs indicates that either

transcription initiation or 5´ processing of Class I RNAs differ between group 4 species and species

belonging to the other evolutionary groups of Dictyostelia. Also, the very 3' end of Class I RNAs is highly

similar with an almost perfectly conserved CTGT sequence in the genomic loci. The coverage from the

RNA-seq data indicates that these four nucleotides are transcribed so that the CUGU sequence is included

in the mature Class I RNA (Additional file 2: Fig. S2) where the C residue always have base pairing potential

with the conserved 5' G residue (Fig. 3a-b). The RNA-seq coverage agrees with the slightly longer than

predicted lengths of Class I RNAs observed by northern blot (see above).

Class I RNA GC-content, size, sequence motif, and stem-structure are conserved throughout Dictyostelia

Comparison of all identified Class I RNAs showed that both Class I RNA length (median of ~60 nt) and GC

content (32-41%) are highly conserved (Additional file 2: Fig. S4a-b). The stable GC content of Class I RNAs

is remarkable considering the variation in overall genome GC content for these six dictyostelids (Additional

file 2: Fig. S4b). In spite of these conserved features and specific sequence motifs discussed above and

later, the overall sequence variability of Class I RNAs is extensive both within and between species. Only

a few examples of loci with identical sequences are found within *D. discoideum, D. purpureum* and *P.*

pallidum, respectively (Additional file 3). No Class I genes with identical sequences were found between

these six analyzed species.

In contrast to the overall variable sequences, the 11 nt sequence motif identified among the *D. discoideum*

Class I RNAs, is highly conserved both within and between all six dictyostelids (marked in grey in Fig. 3a).

The motif is nearly perfectly conserved within group 4, while some positions of the motif are variable in

group 1-3. However, T, C, C, A, and A at position 3, 6, 9, 10 and 11 (counting from the 5´most nt of the

motif) are almost identical between all the Class I RNAs regardless of species. Other nucleotides are well

conserved in most of the evolutionary groups but not all. The sequence motif does not seem to extensively

engage in base-pairing since computational prediction indicates that the conserved motif is less structured

compared to the full-length RNA in most of the organisms (Additional file 2: Fig. S4c-d). However, the first

5´-nucleotide of the motif is often part of the stem-structure (see below), while the base pairing potential

for the remainder of the sequence drops in a pattern similar for all six dictyostelids (Additional file 2: Fig.

S4e).

Another distinct feature common to all *D. discoideum* Class I RNAs is the short (six bp) stem structure,

connecting the 5' and 3' ends of the RNA (Fig. 3a). This stem is predicted to be present in all Class I RNAs

in all six species. However, in contrast to the conserved sequence motif, the nucleotide sequence of the

stem-structure has changed substantially during Dictyostelia evolution. Nevertheless, the base pairing

potential is retained, indicating that it is the structure rather than sequence that is crucial for function

(Fig. 3b). This is further supported by the high number of compensatory mutations found in the predicted

stem of Class I RNAs within each species (Fig. 3b). Notably, in spite of the sequence variation in the stems,

the 5' most G is completely conserved within all Class I RNAs from all six dictyostelids representing each

evolutionary group of Dictyostelia. The predicted base-paired structure of the stem and the unstructured

feature of the conserved sequence motif correspond well with previous *in vitro* probing results of one

Class I RNA, DdR21, from *D. discoideum* [31]. Taken together, Class I GC-content, length, stem structure

and 11 nt motif are highly conserved in all evolutionary groups of Dictyostelia, indicating that these parts

are essential for Class I RNA function.

Class I RNAs are developmentally regulated and highly conserved throughout Dictyostelia

We knew from our previous work that *D. discoideum* Class I RNAs are developmentally regulated and that Class I knock-out cells, *DdR-21* k.o., are disturbed in early development, leading to more and smaller fruiting bodies compared to wild-type cells [31]. In order to investigate if the developmental regulation is conserved also in other dictyostelids, we performed principal component analysis (PCA) of Class I expression in *D. discoideum, P. pallidum* and *D. fasciculatum* based on the RNA-seq data. *D. lacteum* was not included in this analysis since only one replicate per timepoint was available. The PCA plots show developmental regulation of Class I RNAs in all three amoebae as the different life stages are clearly separated (Additional file 2: Fig. S5). Taken together, this suggests a role for Class I RNAs in regulating multicellular development in Dictyostelia.

If the prediction that Class I RNAs are involved in and important for multicellular development holds true, these ncRNAs should be present in all dictyostelids. To analyze this, we used the Class I classifier to investigate the presence of Class I RNA genes in ten additional social amoebae genome sequences. Class I RNA genes were detected in all species, 9-31 genes in each genome, of which the great majority passed manual curation based on the ability to form a short stem connecting the 5' and 3' end (Fig. 4). It should be noted that these are draft genome sequences of varying degree of completeness (Additional file 1). Comparison of all curated loci reinforced the previously identified conserved Class I features i.e. the high sequence conservation of the terminal residues and the 11 nt motif as well as the short stem where structure but not sequence is preserved. In addition, presence of DUSE ~60 nt up-stream of the majority of the identified Class I RNA genes strongly suggests that expressed Class I RNAs exist in all members of Dictyostelia. The TGTG-box, previously only found upstream of *D. discoideum* Class I RNA loci*,* was identified in four additional genomes all belonging to group 4 or the *P. violaceum* complex strengthening the hypothesis that this motif emerged rather late in Class I evolution. In addition, both Class I RNA lengths

and GC content are conserved also when considering all species (Additional file 2: Fig. S6). Taken together,

this proves the existence and emphasize the importance of Class I RNA genes throughout the evolution of

Dictyostelia social amoebae. We have named all curated Class I RNA genes according to the naming

convention previously defined for *D. discoideum* Class I RNA genes [30] (Additional file 4).

Genomic distribution of Class I RNA genes

In *D. discoideum*, all Class I RNA genes are located in intergenic regions and frequently found in clusters

of two or more genes [33]. Clusters of at least two (different) Class I RNA genes are present in all analyzed

Dictyostelia genomes, except for *D. citrinum* (Additional file 5). The absence of Class I RNA gene clusters

in *D. citrinum* is likely a consequence of the quality of the genome assembly (Additional file 1), which is

also reflected in the low number of identified Class I RNA genes. Clusters with a higher number of Class I

RNA genes (three or more) are only found in five genomes, where the distribution in *P. pallidum* resembles

that in *D. discoideum,* i.e. many of the genes are collected in two larger clusters on the same chromosome

(Additional file 5). Even though many Class I RNA genes cluster together, they rarely have identical

sequences. Only a few species-specific identical Class I RNAs were found in *D. polycephalum, P. violaceum,*

D. purpureum, *P. pallidum* and *D. discoideum*, where in the latter two the identical genes are located in

clusters (Additional file 5). Are there Class I RNAs that are identical between two different species? The

only examples found were two Class I RNA loci shared between the group 4 species *D. discoideum* and *D.*

firmibasis (Additional file 4).

To explore the origin of Class I RNAs further, we used the most well-annotated genomes to search for

shared synteny by identifying orthologous genes in the 10 kb region flanking each Class I RNA locus

(Materials and methods). Using this approach, we did not identify any strong evidence for shared synteny

for Class I RNA genes between the different groups of Dictyostelia. Next we investigated if shared synteny

could be detected within group 4 only by performing the same search using the genomes of *D. discoideum,*

D. purpureum and *D. firmibasis.* Almost half of the Class I RNAs in *D. firmibasis* appear to share synteny

with *D. discoideum* Class I RNAs, supported by several protein gene orthologues (Additional file 6) while

no well-supported examples were found for *D. purpureum.* For the identical Class I RNA genes in *D.*

discoideum and *D. firmibasis*, shared synteny was detected for DfiR-4 and DdR-47 (Additional file 6).

Shared synteny between the other two identical Class I RNAs, DfiR-12 and DdR-50, could not be properly

assessed due to the lack of available genome sequence surrounding DfiR-12.

Class I RNAs are unique to dictyostelid social amoebae

The omnipresence of Class I RNAs within Dictyostelia, their developmental regulation as well as the

aberrant development of *D. discoideum* cells lacking *DdR-21* [31] led us to hypothesize that this class of

ncRNAs might be involved in the evolution of Dictyostelia aggregative multicellularity. In order to

investigate this further, we searched for Class I RNA genes in genomes of unicellular amoebae and

amoebae able to form unicellular fruiting bodies. Furthermore, we explored representative genomes of

other major eukaryotic groups, i.e. archeaplastida and ophistokonta. We also chose to include the

proteobacteria *Myxococcus xhantus* as these bacteria exhibit aggregative multicellularity [37], which in

many aspects are analogous to Dictyostelia multicellularity (Fig. 4). We searched these genomes using the

same successful approaches as for Dictyostelia, i.e. using the Class I classifier, based on promoter

characteristics combined with RNA structure and sequence, as well as CM search alone. Only a few

candidates were identified with the Class I classifier. This was anticipated since we did not expect the

DUSE sequence or its distance to the TSS to be conserved outside Dictyostelia. The Infernal search resulted

in a slightly higher number of Class I RNA gene candidates. However, manual inspection revealed that the

candidates are unlikely to represent true Class I RNA genes as they were few in numbers and did not share

characteristics, such as conserved 5' and 3' ends and presence of conserved sequence motif (Fig. 4,

Additional file 7). Taken together, no Class I RNA genes were identified outside Dictyostelia, suggesting

that this class of ncRNAs is unique to dictyostelid social amoebae and important for their aggregative

multicellularity.

Conserved Class I RNA interacting proteins

Class I RNAs are conserved throughout the evolution of Dictyostelia but does this also apply to proteins

associated with this class of ncRNA? We previously identified four Class I RNA interacting proteins in *D.*

discoideum by using one specific Class I RNA, DdR-21, as bait in pull-down experiments. Two of these

proteins, GuaB and NdkC, are involved in nucleotide metabolism while the function of DDB_G0281243 is

unknown. The fourth identified Class I RNA interacting protein, CIBP (also known as Rnp1A [38]), harbors

two RNA binding motifs (RRMs) and was demonstrated to bind directly to the Class I RNA [31]. We

searched for orthologues of the four proteins in the best annotated genomes of Dictyostelia (Fig. 1b,

Additional file 1) and could identify orthologues to GuaB*,* DDB_G0281243 and CIBP in all Dictyostelia

genomes investigated. Next, we used mRNA-seq data from *D. discoideum* (Group 4) [39] as well as *D.*

lacteum (Group 3), *P. pallidum* (Group 2) and *D. fasciculatum* (Group 1) [20] to investigate expression and

developmental regulation of the genes for each of the orthologues. According to the RNA-seq data, only

CIBP were expressed in all four species. Interestingly, during early development the expression pattern of

the gene is similar to that of Class I RNAs [31], i.e. *CIBP* appears to be down-regulated during early

development in all species. However, as *D. discoideum* Class I RNAs continue to decrease in expression

throughout development, *CIBP* expression seems to increase in the final stages of development

(Additional file 2: Fig. S7a). Next, we performed gene predictions in the ten additional Dictyostelia

genomes (Fig 4) and found *CIBP* orthologues in all species except for *D. citrinum* (Additional file 2: Fig.

S7b)*.* The absence of *CIBP* in *D. citrinum* is likely due to the quality of the genome assembly (Additional

file 1). The majority of the orthologues are predicted to encode an approximately 300 amino acids (aa)

long protein with two RRM's. However, shorter orthologues were identified in *A. leptosomum* (114 aa),

A. ellipticum (61 aa) and *D. purpureum* (203 aa). Also, only one RRM could be identified in the *CIBP*

orthologues in *D. purpureum* and *A. ellipticum* (Additional file 2: Fig. S7b). Collectively, the majority of the

disctyostelids are predicted to encode/express CIBP of similar length where the N- and C- terminal

sequences contains RNA binding motifs, RRMs, while the central part of the protein is less conserved.

Despite several efforts by us and others, all attempts to generate a CIBP knock out strain in *D. discoideum*

have been unsuccessful [31, 38]. In addition, no CIBP mutants were generated in the recent Genome Wide

Dictyostelium Insertion (GWDI) project (www.remi-seq.org). Taken together, this indicates that the CIBP

protein is essential.

397
Discussion

The development of high-throughput sequencing techniques has led to the discovery of numerous

ncRNAs. In particular, it has facilitated the identification of small ncRNAs such as mi- and si- and piwi-

interacting (pi)RNAs sized ~21-31 nt and long ncRNAs that can consist of up to several thousand nt.

However, "mid-sized" ncRNA have largely been overlooked partly due to the size selection commonly

carried out before sequencing to enrich for small RNAs and to avoid abundant RNAs such as rRNAs and

tRNAs or fragment thereof. Here we used genome analyses in combination with expression validation to

prove the existence of Class I RNAs in 16 dictyostelid social amoeba. This indicates that Class I RNAs were

present in the last common ancestor of Dictyostelia, dating back at least 600 million years, and were

involved in the transition from unicellular to multicellular life.

Class I RNAs play an important role in *D. discoideum*, as suggested by e.g. the large number of highly

expressed genes and requirement for normal multicellular development [30, 31]. This class of ncRNAs was

initially discovered by sequencing cDNA libraries of full-length RNA sized 50-150 nt [30]. In the same study,

a putative promoter element, DUSE, was identified to be associated with Class I RNA genes. Later, two

approaches to predict Class I RNAs were published. Fragrep, a tool which predict ncRNAs based on

sequence motifs separated by a variable region, identified 45 Class I RNA candidate genes in *D. discoideum*

of which 34 had been previously experimentally validated [40]. In *D. purpureum,* another group 4

dictyostelid, 26 Class I genes were predicted by searching for enriched 8-mers downstream of DUSE motifs

in the genome sequence [18]. Hence, previous to the present study, Class I RNAs had only been identified

in organisms belonging to one group of Dictyostelia. We now asked if Class I RNA genes also are present

in other organisms, both within Dictyostelia characterized by aggregative multicellularity and outside this

group of social amoebae.

In order to detect Class I RNAs, we first built a covariance model (CM) which predicts candidates based on

both sequence and structure information of Class I RNAs. Based on the CM, we created a classifier which

evaluates the identified candidates based on the presence of DUSE at the correct distance from TSS or

from the TGTG-box (found upstream of many *D. discoideum* Class I RNA genes). Next, we confirmed

expression of approximately 100 Class I RNAs, predicted by the classifier, by northern blot and/or RNA-

seq. We were now confident that we could use the Class I classifier to accurately identify expressed Class

I RNA genes also in other organisms. Based on this, we show that Class I RNAs are present in all

dictyostelids with available genome sequences (total of 16). Having established the presence of Class I

RNAs in all tested disctyostelids, we turned our focus on organisms outside of Dictyostelia. Outgroups

were chosen to both represent organisms with a unicellular life style and those that go through different

kinds of multicellular development. We used both the classifier and the CM search alone to search for

candidates, the latter to avoid the constraint of having a DUSE element in front of the Class I RNA genes.

Regardless of search approach, we did not identify Class I RNAs in any organism outside Dictyostelia,

suggesting that these RNAs are restricted to dictyostelid social amoebae.

The ubiquitous presence of Class I RNA genes in Dictyostelia and absence in the closest related unicellular

amoebae suggest that these RNAs were present in the last common ancestor of Dictyostelia and played a

role in the transition from unicellular life to aggregative multicellularity. This is further supported by the

developmental regulation of Class I RNAs and that at least one member, DdR-21, is required for normal

development [30, 31]. In *D. discoideum*, transition from unicellular growth to multicellular development

is associated with large transcriptional reprogramming of protein coding genes [23] and different cell

types, i.e. prespore and prestalk cells, can be separated based on the transcriptional signatures of

individual cells [41]. Interestingly, the majority of the protein coding genes that are essential for

multicellular development in *D. discoideum* is also present in strictly unicellular amoebae [20]. Hence, it

is likely that the ability to regulate how genes are expressed have been key in the evolution of multicellular

development. Maybe Class I RNAs can rewire gene expression of genes present in unicellular organisms

to create new networks adapted for development. We have no evidence that Class I RNA directly interacts

with mRNA to regulate gene expression even though we are not excluding this mode of action. Another

possibility is that Class I RNA regulate development by binding to proteins that directly or indirectly control

development, maybe by acting as a molecular sponge, where specific proteins are sequestered by Class I

RNAs. This could buffer the action of these proteins. Perhaps the observed down regulation of Class I RNAs

during development lead to an increase in free active proteins important for multicellular development.

CIBP would be a candidate for this kind of regulation since CIBP directly interacts with Class I RNAs, at

least with the tested DdR21, in *D. discoideum* [31]. Support for CIBP as an important protein for

development is its presence in all dictyostelids. We are currently attempting to decipher the function of

Class I RNAs in *D. discoideum* by using a coupled RNA-seq and proteomics approach to investigate the

involvement of Class I RNA in early multicellular development. The tendency to differentially regulate

genes in order to create new functions is seen also in metazoan evolution, where an increase in regulatory

miRNAs is correlated with increased organismal complexity [2]. Interestingly, *D. discoideum* is one of the

few organisms outside animals and land plants were miRNAs have been identified [24–27]. If miRNAs,

similar to Class I RNAs, are present in other dictyostelids is currently being investigated.

In Dictyostelia, the most complex multicellularity is found within group 4, exemplified by regulated

proportions of specialized cell types and a migrating slug stage [3]. Interestingly, in analogy to miRNA

expansion in complex animals, the number of expressed Class I RNA genes is correlated with Dictyostelia

complexity where group four has the largest number of Class I RNA genes (Fig. 2b). In addition, RNA-seq

data indicate that Class I RNAs are expressed at higher levels in group 4 (*D. discoideum*), further

strengthening that Class I RNAs are involved in increased organismal complexity. This increased expression

appears to be connected to the TGTG-box between DUSE and the TSS. So far, the TGTG-box have only

been identified in Class I RNA loci in species belonging group 4 (except *D. purpureum*) and *P. violaceum*

complex (Figs 3a and 4 and Additional file 3). Thus, this motif is likely a rather late addition in the evolution

of Class I RNAs. It should be noted that it is currently not known if the second motif is actually a promoter

element or the TSS of a longer precursor that is processed down to the mature RNA. In either case, the

TGTG-box is associated with Class I RNA genes in social amoebae with higher levels of complexity as

compared to other dictyostelids and appears to add another layer of regulation to Class I RNA expression.

The emergence of the TGTG-box somewhere after the split of group 3 and 4 and its connection to

increased phenotypic complexity seen in group 4 dictyostelids is somewhat analogous to changes in cis-

regulatory elements, such as enhancers, and morphological evolution in animals [42].

The high number of novel Class I RNA loci identified in Dictyostelia enabled comparative studies which

provides information on their key features. The short stem connecting the 5' and 3' ends of mature Class

I RNA is conserved. Furthermore, the sequence variability of the stem between organisms but also the

high number of compensatory mutations within each species strongly suggest that it is the structure

rather than sequence that is important for function. Flanking the stem structure at both ends are highly

conserved nucleotides present in almost all Class I RNAs: AAG and CTGT at the 5' and 3'ends respectively,

where the 5'G and 3'C forms the first base-pair of the stem. Curiously, in spite of the high conservation of

the three 5' nucleotides, the mature transcript of Class I RNAs in *D. discoideum* almost always starts with

the G residue, while it is more common in the group 1-3 representatives that also the two A's are included.

In either case, the start of each mature Class I RNA is well defined with the great majority of all RNA-seq

reads sharing the same 5' end. The 3'end is also remarkably conserved. Whether these nucleotides are

part of the mature transcript is hard to assess due to the sequencing approach used, which only allowed

us to capture a small fraction of the 3' ends. However, we do find reads covering the 3' conserved CTG in

the majority of the different Class I RNAs, suggesting that these constitute the 3' end of the mature Class

I RNAs. This is also supported by the full-length cDNA libraries of small RNAs in *D. discoideum* where this

class of RNA was first discovered [30].

The comparison of Class I RNA loci also confirms the importance of the ~11 nt motif present immediately

after the 5' part of the stem structure. Although some of the nucleotides vary within and in between

organisms, several residues are nearly perfectly conserved. Yet another conserved feature is the putative

promoter motif, DUSE. Based on studies of spliceosomal RNAs in *D. discoideum*, we previously

demonstrated that DUSE is associated with genes transcribed by both RNA Pol II and RNA Pol III [43].

However, we believe that Class I RNAs are transcribed by RNA Pol III since the canonical RNA Pol III

termination signal is present downstream of most Class I RNA loci and Class I RNA reads are lacking in

poly(A) enriched RNA-seq libraries.

Based on the findings in this study, we conclude that Class I RNAs were present in the last common

ancestor of Dictyostelia. In addition, our data also strongly suggest that the putative promoter element

DUSE was present 60 nt upstream of the ancestral Class I RNA gene and that the element was required

for expression of the gene. We also conclude that the ancient Class I RNA was characterized by a short

stem structure and a 11 nt sequence motif where at least six of the positions were identical to the

corresponding nucleotides in extant Class I RNAs. Due to the high sequence and structure variability of

the region between the 11 nt motif and the start of the 3' stem in identified Class I RNAs, we cannot

resolve this part of the ancestral sequence. However, the total length of the mature RNA was probably

approximately 60 nt long (Fig. 5).

Even though some motifs are highly conserved within all Class I RNAs, conservation of complete Class I

RNA genes are rare. Only a few examples of identical loci within the same genome are found in a handful

of dictyostelids and only two identical Class I RNAs shared by two different species were identified, i.e.

the group 4 dictyostelids *D. discoideum* and *D. firmibasis* (Additional file 4). In *D. discoideum* and *P.*

pallidum many Class I RNA genes are situated in larger clusters and it is within these clusters where the

species-specific identical Class I RNA genes are found, perhaps indicating expansion of Class I RNA genes

by duplication. Interestingly, the snRNA genes in *D. discoideum* are organized in a similar way where

closely related genes often are found in pairs situated very close together [33]. However, in general the

Class I RNA genes are spread out in the genomes and this is also true for the majority of the identical Class

I RNA genes in the different genomes. Hence, the low occurrence of shared synteny, low overall sequence

conservation and different number of loci in different organisms suggest that the expansion of Class I

RNAs mainly occurred after speciation.

Conclusions

We have identified Class I RNAs in 16 different dictyostelids and validated their expression in

representatives of each major group of Dictyostelia dating back at least ~600 million years. Despite the

large evolutionary distances, Class I RNA genes share promotor motifs and the mature RNAs have several

characteristics in common, i.e. short stem, conserved sequence motif and highly conserved 5' and 3' ends.

In addition, the *D. discoideum* Class I RNA interacting protein CIBP is conserved throughout Dictyostelia.

Furthermore, the gene shares its expression profile with Class I RNAs during early development, indicating

that function and mode of action is also conserved. Although Class I RNAs are present in all dictyostelids

investigated, no evidence was found for this class of RNAs in any other organism. Taken together, our

results suggest that Class I RNAs are important for the evolution of multicellularity in Dictyostelia.

539
Materials and methods

RNA isolation and northern blot

The following strains were used for northern blot validation of Class I expression: *D. discoideum* AX2

(DBS0235521, www.dictybase.org), *D. purpureum* WS321, *P. pallidum* PN500, *D. lacteum* Konijn, *D.*

fasciculatum SH3, *A. subglobosum* LB1. All strains, except *D. discoideum,* were kindly provided by Dr Maria

Romeralo and Professor Sandra Baldauf. Total RNA was extracted with TRIzol (ThermoFisher Scientific)

from cells grown in association with *Klebsiella aerogenes* on non-nutrient agar. Northern blots were

performed as described in [30, 31]. Briefly, 10 µg total RNA was separated on 8 % PAGE/7 M Urea and

electroblotted to Hybond N+ nylon membranes (GE Healthcare). After UV crosslinking, immobilized RNA

was hybridized with ^{32}P-labeled oligonucleotides in Church buffer at 42 °C overnight. Signals were

analyzed with a Personal Molecular Imager (BIO-RAD) normally after a few hours' exposure. Membranes

analyzed more than once were stripped with 0.1×SSC/1 % SDS buffer at 95 °C for 1 h and controlled for

residual signal before reprobing. Oligonucleotide sequences are provided in Additional file 2: Table S1.

RNA-seq validation

Strain growth and RNA extraction for RNA-seq have been described previously [20]. For each strain, RNA

was prepared from growing cells and two multicellular developmental stages: aggregates and tipped

aggregates/fingers (biological duplicates except for *D. lacteum*). Truseq small RNA Sample Preparation kit

(product # RS- 200-0012, Illumina) was used to prepare sequencing libraries from 1 µg total RNA. The

library preparation was performed according to the manufacturer's protocol (#15004197 rev G) where

cDNA representing 18 nt – 70 nt RNA were isolated. Single read 50 bp sequencing was performed using

v4 sequencing chemistry on an Illumina HiSeq2500. To reduce influence of degradation products, only full

length 50 bp reads were mapped with bowtie allowing for 1 mismatch [44] and counted with feature

counts [45]. Read coverage over Class I candidate loci were calculated with BEDTools genomecov v. 2.26.0

[46]. Class I RNAs were considered to be validated by RNA-seq if the read coverage indicated a distinct 5'

end and reads specifically matched the predicted loci, i.e. did not appear to be part of a considerably

longer transcript. Principal component analyses were performed using DESeq2 [47]. ROC plot evaluation

of Class I prediction was performed using pROC package[48] in R.

Strains and genomic resources

Strain names and accession numbers for genomic sequences are listed in Additional file 1.

Identification of Class I RNAs

Of the 40 *D. discoideum* Class I RNAs annotated previous to this study, six were excluded from the model

build (r48, r53, r54, r55, r58 and r61) as they represented truncated fragments or lacked the canonical

features such as the stem or conserved sequence motif. The remaining 34 were aligned with MAFFT

v7.407 [49] using the ginsi setting. Consensus structure for the alignment was predicted with RNAalifold

2.3.3 [50] using the -T 22 option to account for the optimal growing temperature of the amoebae.

Alignment and consensus structure was combined to a Stockholm alignment file and a co-variance model

was created with Infernal 1.1.2 [32]. Infernal was then used to search the genomes of *D. discoideum, D.*

purpureum, D. lacteum, P. pallidum, A. subglobosum and *D. fasciculatum* using default settings and

candidates with a score ≥ 25 were added to the alignment using MAFFT (ginsi –add). The new alignment

was manually curated and used to predict consensus structure, create a new co-variance model, and

perform a new search in the same genomes. This procedure was iterated six time, i.e. until no new

candidates with an Infernal score ≥ 25 were identified. Enriched sequence motifs were identified up to

150 nt upstream of identified candidates with MEME v. 5.0.3 [51]. For final Class I identification, Infernal

searches were performed with increased sensitivity and all candidates scoring 15 or higher were kept

(cmsearch –nohmm –notrunc -T 15). The candidates were then evaluated based on the presence of DUSE

and TGTG-box in the 150 nt preceding the predicted start of transcription (see above) using FIMO v. 5.0.3

[34]. Infernal score, FIMO motif score, and a motif distance score (+5) were then added to a total score.

Missing DUSE or incorrect distance was penalized with -10 or -5 respectively. If a total score of 32 was

achieved, the candidate was considered likely to be an expressed true Class I RNA and kept for further

analyses. Representative sequence logos of manually curated sequence alignments (mafft --maxiterate

1000 –localpair) were created with WebLogo 3 [52].

Orthologue identification and shared synteny search

For Dictyostelia species lacking genome annotations, gene prediction was performed with Gene id v. 1.4

[53] using Dictyostelium parameter file. Orthologue identification was performed using OrthoFinder v.

2.3.3 [54]. Protein domain architectures for CIBP/Rnp1A orthologues were analyzed with hmmscan using

the HMMER web interface [55]. Orthologue information for *D. discoideum, D. firmibasis*, *D. lacteum*, *P.*

pallidum, A. subglobosum and *D. fasciculatum* was used to investigate shared synteny for Class I RNA loci.

For each Class I RNA locus, gene information within a 10 kb flanking region was retrieved. Next, we

searched for orthologues of these genes in the other organisms included in the search. If a Class I RNA

was found within 10 kb of an orthologous gene in another organism, it was manually inspected to

determine the level of shared synteny.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

Competing interest

The authors declare that they have no competing interests

Authors' contribution

JK, LA, JR, and FS participated in the design of the project; LA and ZL generated experimental data; JK

performed most of the data analysis and prepared figures; JR participated in the bioinformatic analysis;

L.E, A.N, G.G, and P.S supplied RNA for sequencing. JK and FS drafted the manuscript.

Funding

This work was supported by the Linneus Support from the Swedish Research Council to the Uppsala RNA

Research Centre; Carl Tryggers Foundation under Grant number CST 18:381.

627
628 **Acknowledgements**

629 Sequencing was performed by the SNP&SEQ Technology Platform in Uppsala. The facility is part of the

630 National Genomics Infrastructure (NGI) Sweden and Science for Life Laboratory. The SNP&SEQ Platform

631 is also supported by the Swedish Research Council and the Knut and Alice Wallenberg Foundation.

References

1. Cech TR, Steitz JA. The Noncoding RNA Revolution—Trashing Old Rules to Forge New Ones. Cell.
2014;157:77–94.

2. Gaiti F, Calcino AD, Tanurdžić M, Degnan BM. Origin and evolution of the metazoan non-coding
regulatory genome. Developmental Biology. 2017;427:193–202.

3. Kawabe Y, Du Q, Schilde C, Schaap P. Evolution of multicellularity in Dictyostelia. Int J Dev Biol.
2019;63:359–69.

4. Brown MW, Spiegel FW, Silberman JD. Phylogeny of the "forgotten" cellular slime mold, Fonticula
alba, reveals a key evolutionary branch within Opisthokonta. Mol Biol Evol. 2009;26:2699–709.

5. Brown MW, Silberman JD, Spiegel FW. A contemporary evaluation of the acrasids (Acrasidae,
Heterolobosea, Excavata). Eur J Protistol. 2012;48:103–23.

6. He D, Fiz-Palacios O, Fu C-J, Fehling J, Tsai C-C, Baldauf SL. An alternative root for the eukaryote
tree of life. Curr Biol. 2014;24:465–70.

7. Tice AK, Silberman JD, Walthall AC, Le KND, Spiegel FW, Brown MW. Sorodiplophrys stercorea:
Another Novel Lineage of Sorocarpic Multicellularity. J Eukaryot Microbiol. 2016;63:623–8.

8. Lasek-Nesselquist E, Katz LA. Phylogenetic position of Sorogena stoianovitchae and relationships
within the class Colpodea (Ciliophora) based on SSU rDNA sequences. J Eukaryot Microbiol.
2001;48:604–7.

9. Brown MW, Silberman JD, Spiegel FW. "Slime molds" among the Tubulinea (Amoebozoa): molecular
systematics and taxonomy of Copromyxa. Protist. 2011;162:277–87.

10. Brown MW, Kolisko M, Silberman JD, Roger AJ. Aggregative multicellularity evolved
independently in the eukaryotic supergroup Rhizaria. Curr Biol. 2012;22:1123–7.

11. Heidel AJ, Lawal HM, Felder M, Schilde C, Helps NR, Tunggal B, et al. Phylogeny-wide analysis of
social amoeba genomes highlights ancient origins for complex intercellular communication. Genome Res.
2011;21:1882–91.

12. dos Reis M, Thawornwattana Y, Angelis K, Telford MJ, Donoghue PCJ, Yang Z. Uncertainty in the
Timing of Origin of Animals and the Limits of Precision in Molecular Timescales. Curr Biol.
2015;25:2939–50.

13. Schilde C, Lawal HM, Kin K, Shibano-Hayakawa I, Inouye K, Schaap P. A well supported multi gene
phylogeny of 52 dictyostelia. Molecular Phylogenetics and Evolution. 2019;134:66–73.

14. Romeralo M, Skiba A, Gonzalez-Voyer A, Schilde C, Lawal H, Kedziora S, et al. Analysis of
phenotypic evolution in Dictyostelia highlights developmental plasticity as a likely consequence of
colonial multicellularity. Proceedings of the Royal Society B: Biological Sciences. 2013;280:20130976.

15. Schilde C, Skiba A, Schaap P. Evolutionary reconstruction of pattern formation in 98 Dictyostelium

species reveals that cell-type specialization by lateral inhibition is a derived trait. EvoDevo. 2014;5:34.

16. Sheikh S, Thulin M, Cavender JC, Escalante R, Kawakami S-I, Lado C, et al. A New Classification of
the Dictyostelids. Protist. 2018;169:1–28.

17. Eichinger L, Pachebat JA, Gloeckner G, Rajandream M-A, Sucgang R, Berriman M, et al. The
genome of the social amoeba Dictyostelium discoideum. Nature. 2005;435:43–57.

18. Sucgang R, Kuo A, Tian X, Salerno W, Parikh A, Feasley CL, et al. Comparative genomics of the
social amoebae Dictyostelium discoideum and Dictyostelium purpureum. Genome Biol. 2011;12:R20.

19. Urushihara H, Kuwayama H, Fukuhara K, Itoh T, Kagoshima H, Shin-I T, et al. Comparative genome
and transcriptome analyses of the social amoeba Acytostelium subglobosum that accomplishes
multicellular development without germ-soma differentiation. BMC Genomics. 2015;16:80.

20. Glöckner G, Lawal HM, Felder M, Singh R, Singer G, Weijer CJ, et al. The multicellularity genes of
dictyostelid social amoebas. Nature Communications. 2016;7:1–11.

21. Hillmann F, Forbes G, Novohradská S, Ferling I, Riege K, Groth M, et al. Multiple Roots of Fruiting
Body Formation in Amoebozoa. Genome Biol Evol. 2018;10:591–606.

22. Deline B, Greenwood JM, Clark JW, Puttick MN, Peterson KJ, Donoghue PCJ. Evolution of
metazoan morphological disparity. Proc Natl Acad Sci U S A. 2018;115:E8909–18.

23. Rosengarten RD, Santhanam B, Fuller D, Katoh-Kurasawa M, Loomis WF, Zupan B, et al. Leaps and
lulls in the developmental transcriptome of Dictyostelium discoideum. BMC Genomics. 2015;16:294.

24. Avesson L, Reimegard J, Wagner EGH, Soderbom F. MicroRNAs in Amoebozoa: Deep sequencing
of the small RNA population in the social amoeba Dictyostelium discoideum reveals developmentally
regulated microRNAs. RNA. 2012;18:1771–82.

25. Liao Z, Kjellin J, Hoeppner MP, Grabherr M, Söderbom F. Global characterization of the Dicer-like
protein DrnB roles in miRNA biogenesis in the social amoeba Dictyostelium discoideum. RNA Biol.
2018;15:937–54.

26. Hinas A, Reimegård J, Wagner EGH, Nellen W, Ambros VR, Söderbom F. The small RNA repertoire
of Dictyostelium discoideum and its regulation by components of the RNAi pathway. Nucl Acids Res.
2007;35:6714–26.

27. Meier D, Kruse J, Buttlar J, Friedrich M, Zenk F, Boesler B, et al. Analysis of the Microprocessor in
Dictyostelium: The Role of RbdB, a dsRNA Binding Protein. PLoS Genet. 2016;12:e1006057.

28. Rosengarten RD, Santhanam B, Kokosar J, Shaulsky G. The Long Noncoding RNA Transcriptome of
Dictyostelium discoideum Development. G3 (Bethesda). 2016;7:387–98.

29. Hildebrandt M, Nellen W. Differential antisense transcription from the Dictyostelium EB4 gene locus:
Implications on antisense-mediated regulation of mRNA stability. Cell. 1992;69:197–204.

30. Aspegren A, Hinas A, Larsson P, Larsson A, Söderbom F. Novel non-coding RNAs in Dictyostelium

discoideum and their expression during development. Nucl Acids Res. 2004;32:4646–56.

31. Avesson L, Schumacher HT, Fechter P, Romby P, Hellman U, Söderbom F. Abundant class of non-
coding RNA regulates development in the social amoeba Dictyostelium discoideum. RNA Biol.
2011;8:1094–104.

32. Nawrocki EP, Eddy SR. Infernal 1.1: 100-fold faster RNA homology searches. Bioinformatics.
2013;29:2933–5.

33. Hinas A, Soederbom F. Treasure hunt in an amoeba: non-coding RNAs in Dictyostelium discoideum.
Current Genetics. 2007;51:141–59.

34. Grant CE, Bailey TL, Noble WS. FIMO: scanning for occurrences of a given motif. Bioinformatics.
2011;27:1017–8.

35. Burki F, Roger AJ, Brown MW, Simpson AGB. The New Tree of Eukaryotes. Trends in Ecology &
Evolution. 2020;35:43–55.

36. Richard P, Manley JL. Transcription termination by nuclear RNA polymerases. Genes Dev.
2009;23:1247–69.

37. Muñoz-Dorado J, Marcos-Torres FJ, García-Bravo E, Moraleda-Muñoz A, Pérez J. Myxobacteria:
Moving, Killing, Feeding, and Surviving Together. Front Microbiol. 2016;7.
doi:10.3389/fmicb.2016.00781.

38. Ngo T, Miao X, Robinson DN, Zhou Q. An RNA-binding protein, RNP-1, protects microtubules from
nocodazole and localizes to the leading edge during cytokinesis and cell migration in Dictyostelium cells.
Acta Pharmacologica Sinica. 2016;37:1449–57.

39. Parikh A, Miranda ER, Katoh-Kurasawa M, Fuller D, Rot G, Zagar L, et al. Conserved developmental
transcriptomes in evolutionarily divergent species. Genome Biology. 2010;11:R35.

40. Mosig A, Sameith K, Stadler P. Fragrep: An Efficient Search Tool for Fragmented Patterns in
Genomic Sequences. Genomics Proteomics Bioinformatics. 2006;4:56–60.

41. Antolović V, Lenn T, Miermont A, Chubb JR. Transition state dynamics during a stochastic fate
choice. Development. 2019;146. doi:10.1242/dev.173740.

42. Gaunt SJ, Paul Y-L. Changes in Cis-regulatory Elements during Morphological Evolution. Biology
(Basel). 2012;1:557–74.

43. Hinas A, Larsson P, Avesson L, Kirsebom LA, Virtanen A, Söderbom F. Identification of the major
spliceosomal RNAs in Dictyostelium discoideum reveals developmentally regulated U2 variants and
polyadenylated snRNAs. Eukaryotic Cell. 2006;5:924–34.

44. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short
DNA sequences to the human genome. Genome Biol. 2009;10:R25.

45. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning

sequence reads to genomic features. Bioinformatics. 2014;30:923–30.

46. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 2010;26:841–2.

47. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. Genome Biol. 2014;15:550.

48. Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez J-C, et al. pROC: an open-source
package for R and S+ to analyze and compare ROC curves. BMC Bioinformatics. 2011;12:77.

49. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in
Performance and Usability. Mol Biol Evol. 2013;30:772–80.

50. Bernhart SH, Hofacker IL, Will S, Gruber AR, Stadler PF. RNAalifold: improved consensus structure
prediction for RNA alignments. BMC Bioinformatics. 2008;9:474.

51. Bailey TL, Elkan C. Fitting a mixture model by expectation maximization to discover motifs in
biopolymers. Proc Int Conf Intell Syst Mol Biol. 1994;2:28–36.

52. Crooks GE. WebLogo: A Sequence Logo Generator. Genome Research. 2004;14:1188–90.

53. Blanco E, Parra G, Guigó R. Using geneid to identify genes. Curr Protoc Bioinformatics.
2007;Chapter 4:Unit 4.3.

54. Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. Genome Biol. 2015;16:157.

55. Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. HMMER web server: 2018 update.
Nucleic Acids Res. 2018;46:W200–4.

755
756

     Figures

759
Figure 1. Search strategy and classification of Class I RNAs a) Schematic representation of previously

described *D. discoideum* Class I RNAs [31]. b) Schematic phylogeny showing the location of Amoebozoa,

a sister group to Obazoa (Opisthokonta, Breviates and Apusomonada) in the eukaryotic tree of life based

on [35]. Dictyostelia is represented by species belonging to each major group [13]. The genomes of

these dictyostelids were searched for Class I RNA genes and newly identified genes were used to refine

the co-variance model. c) Enriched sequence motifs identified upstream of Class I RNA gene candidates

in the different dictyostelids represented in panel b (Infernal score ≥ 25, n=126). The putative promoter

motif (DUSE) is found approximately 60 nt from the predicted start of transcription (TSS) in all organisms

(upper part). DUSE in combination with TGTG-box, only identified in *D. discoideum* (lower part). d)

Summary of scoring system used for the classifier of Dictyostelia Class I RNA based on Infernal score ≥

15, presence of DUSE, and distance between DUSE and predicted TSS or TGTG-box.

Figure 2. Expression of predicted Class I RNA genes. a) Northern blot validation of different Class I RNAs

from *D. purpureum* (DpuR), *D. lacteum* (DlaR), *D. fasciculatum* (DfaR), *A. subglobosum* (AsuR), and *P.*

pallidum (DpaR). The number after the species-specific designations indicates which Class I RNA the

probe recognizes. When two numbers are given, the probe recognize two different Class I RNAs. DpuR-X

indicates that the probe is expected to hybridize to six different Class I RNAs predicted to be 85 nt long.

DpuR-Y indicates that the probe is expected to hybridize to 24 Class I RNAs. For each organism except

for *D. fasciculatum* (DfaR), the same membrane was probed, stripped, and reprobed for the different

Class I RNAs. Radioactively labeled size marker is indicated by M and numbers to the left indicate sizes in

nucleotides. b) Number of Class I RNA genes in each species according to the classifier. RNA-seq

designate the number of expressed Class I RNA genes verified by RNA-seq. c) ROC curves based on the

RNA-seq validation and either classifier score or CM score for all Class I candidates in *D. discoideum, D.*

lacteum, P. pallidum, and *D. fasciculatum* identified in the CM search. Input data available in Additional

784 file 3. Evaluation of the classifier and CM thresholds used throughout the study are shown above the

785 plot. Individual ROC curves for each organism are found in Additional file 2: Fig. S2.

Figure 3. Conserved characteristics of Class I RNAs a) Sequence logo representing species specific

alignments of conserved features of Class I RNA loci. DUSE indicate the putative promoter element and

the 5' and 3´stem sequences of the conserved stem-structure are indicated. The conserved 11 nt sequence

motif adjacent to the 5´part of the stem is boxed in gray. The dashed vertical line denotes the predicted

start of the Class I RNAs based on the CM search. The sequence logo between DUSE and the 5' stem motif

for *D. discoideum* represent the TGTG-box. Numbers of nucleotides (nt) correspond to the distances

between indicated motifs. b) Displayed are nucleotides representing the most common base pair for each

position (numbers to the left) of the conserved stem structure for each organism (color key down right).

Numbers to the right represent less common nucleotide combinations predicted to base pair. The few

combinations of nucleotides not predicted to base pair are not shown. Schematic structure of Class I RNA

is presented to the right.

Figure 4. Class I RNAs are ubiquitous in and restricted to dictyostelid social amoebae.

Upper part: Class I RNA loci were searched for in the genomes of 16 different Dictyostelia (Additional file

1). The number of hits identified by the classifier is indicated as well as the number of these loci that

passed manual curation. The number of curated loci with DUSE and the TGTG-box at the correct

distance are shown. Bottom part: Result from searches for Class I RNA loci outside Dictyostelia. Life style

indicate unicellular or multicellular organisms. Fruiting bodies denotes if the organism life style involves

formation of fruiting bodies. CM indicates number of candidates identified with a CM score ≥ 25.

Headings Classifier and Curated as described above. Further information about outgroup Class I RNA

candidates is available in Additional file 7.

Figure 5. Ancestral Class I RNA and conserved key features

Left: Schematic representation of the ancestral Class I RNA transcript. The putative promoter element

DUSE and its distance to the Class I RNA gene is indicated below the Class I RNA structure. Strongly

conserved features, nucleotide and base paired stem, are colored black while red denotes more variable

positions (based on data presented in figure 3). The dotted part of the loop indicates highly variable

sequence. Right: Sequence logos of alignments of conserved sequence motifs from all identified and

curated Class I RNAs. Only DUSE identified at the correct distance were included in the alignment. The

11 nt motif is boxed in gray in both the ancestral Class I RNA (left) and sequence logo (right).

Description of additional files

File name: Additional file 1

File format: excel (.xlsx)

Title of data: Genomic resources used in this study.

Description: References for dictyostelid genomic sequences used in the study together with alternative

names (Synonyms) for organisms (where applicable) and basic genome sequence statistics. Genome

sequences used in the construction of the co-variance model (CM) are indicated in bold. For outgroup

genome sequences used in the study, only the GenBank or RefSeq reference are given.

File name: Additional file 2

File format: PDF (.pdf)

Title of data: Supplementary figures and tables

Description: Supplementary figures 1-7 and table 1

File name: Additional file 3

File format: excel (.xlsx)

Title of data: Spreadsheet with search summary and expression validation for Class I RNAs from selected

dictyostelids.

Description: The spreadsheet show data for all predicted Class I RNAs, from respective organism,

passing the CM score ≥ 15. Candidates considered to be true Class I RNAs have been named according to

previous naming rules. Classifier summary includes classifier score and infernal (CM) score as well as

identified DUSE sequence (DUSE_seq), the duse score obtained from FIMO (duse_score), the duse

distance from the predicted Class I RNA start (duse_dist), the TGTG-box sequence (TGTG-box_seq) and

score obtained from FIMO (TGT-box_score), TGTG distance from predicted Class I start (TGT-box_dist)

and the distance between DUSE and TGTG-box (inter_dist). Read counts denotes raw RNA-seq read

counts for each Class I RNA candidate in each sequencing library for the organisms where we performed

RNA-seq. Read counts include time points, for growning cells (0h) and time after starvation, when RNA

was collected and A and B denotes biological replicates. Expression validation is based on read counts

and coverage over each respective locus and northern blot. For Read coverage, the values 1 and 0

denotes validated and not validated, respectively. The read coverage validation in combination with

either CM score or classifier score was used to generate ROC curves in figure 2 and Additional file 2: Fig.

S3. Northern blot indicates Class I RNAs analyzed by northern blot in this study.

File name: Additional file 4

File format: excel (.xlsx)

Title of data: Spreadsheet with name, genome location, validation and score of all curated Class I genes

for all investigated dictyostelids.

Description: For each Class I RNA gene Genomic location includes chromosome/contig/scaffold

(Chromosome), start and end of predicted RNA (Start and Stop, respectively), on which DNA strand the

RNA is located (Strand), and the predicted length (Class I RNA length). The Classifier summary includes

the classifier score, the CM score the DUSE sequence and score (DUSE_seq and duse_score,

respectively)) and TGTG-box sequence and score (TGTG-box_seq and TGTG-box_score, respecively).

Columns duse_dist and TGTG-box_dist denotes the distance between the motif and predicted TSS and

inter_dist denotes the distance between the two motifs. Under Upstream motifs, columns DUSE and

TGTG-box summarize the classifier output and indicate if respective motif is found at the correct

distance (1) or not (0). Where applicable, information about RNA-seq (1 denotes expressed Class I based

on read coverage and 0 if expression could not be validated) and/or northern blot validation is given.

File name: Additional file 5

File format: PDF (.pdf)

Title of data: Genomic distribution of Class I RNA loci.

Description: Chromosome/contig/scaffold names are indicated to the left and respective lengths are normalized to the longest one (length in bp are presented to the very right of each schematic stretch of DNA). Total number of Class I RNA genes for each organism is given at the top and for each chromosome/contig/scaffold to the very right. Class I RNA genes are indicated by black arrows except for genes with identical sequences, which are indicated with colored arrows. Short vertical lines specify every 0.5 mbp. In *D. discoideum*, the duplication on chromosome 2 (DDB0232429) is indicated by grey boxes with vertical lines.

File name: Additional file 6

File format: PDF (.pdf)

Title of data: Shared synteny between *D. discoideum* and *D. firmibasis* Class I RNA loci.

Description: Representation of *D. firmibasis* (dfi) Class I RNA loci +/- 10 kb that share synteny with *D. discoideum* (ddi) Class I RNAs supported by at least two orthologous genes (connected with dashed lines). Chromosome/contig names are given to the left of the genes. Genes situated on the forward and the reverse strand are indicated above and below the chromosome/contig, respectively. Vertical grey lines are given at every 1000 nt.

File name: Additional file 7

File format: excel (.xlsx)

Title of data: Identified Class I RNA candidates in outgroups.

Description: For each candidate (classifier score ≥ 32 and/or CM score ≥ 25) the following information is

given: classifier score, CM score as well as identified DUSE sequence (DUSE_seq), the duse_score

obtained from FIMO, the distance of DUSE (duse_dist) from the predicted Class I start, the TGTG-box

sequence (TGTG-box_seq) and score (TGTG-box_score) obtained from FIMO, TGTG_distance from

predicted Class I start and the distance between DUSE and TGTG-box (inter_dist). The last column (Seq),

contains the full genomic sequence of each candidate.