1 **The order Tethyida (Porifera) in California: taxonomy, systematics, and the first**
2 **member of the family Hemiasterellidae in the Eastern Pacific**
3

4 **THOMAS L. TURNER**
5 *Ecology, Evolution, and Marine Biology Department*
6 *University of California, Santa Barbara*
7 *Santa Barbara, CA 93106*
8
9 tlturner@ucsb.edu
10
11 ORCiD: 0000-0002-1380-1099

12
13 **Abstract**
14

Sponges (phylum Porifera) are ubiquitous inhabitants of marine ecosystems and have
been shown to provide substantial ecosystem services. Despite this ecological importance,
the sponge fauna in California has received little study. Here I use a collection of kelp
forest sponges to describe the diversity of the order Tethyida in California. This order
contains the genus *Tethya*, which has been included in long-term ecological monitoring
projects. I show that *Tethya vacua* is a junior synonym of *Tethya californiana,* and thus
all *Tethya* populations known in California belong to a single species. Genetic data from
California's only Timeidae, *Timea authia,* indicate that this species is in fact in the
Tethyidae and is the third known member of the genus *Tethytimea*. I also describe the
first member of the family Hemiasterellidae from the Eastern Pacific, *Galaxia gaviotensis*
gen. nov. sp. nov. By combining field photographs, morphological taxonomy, and
phylogenetic analysis of these samples, this work will facilitate future efforts to
understand the evolution of this order and the ecological role of sponges in the California
kelp forest.
29
30 Key Words: sponges, Tethya, Timea, Tethyidae, Timeidae, Hemiasterellidae, 18S, 28S,
31 cox1

32
33

Introduction

Though little studied in California, sponges have been shown to play many important roles in other ecosystems, including habitat creation, biological disturbance, and consumption of bacterial and viral plankton (Bell 2008). Sponges in the genus *Tethya* are often called "puffball" sponges, as they grow in a roughly spherical form. Because they are common and distinctive, they are better known than most sponges in California. Indeed, *Tethya* populations – referred to as *T. aurantium* in most ecological publications – have been included in several long-term ecological monitoring efforts aimed at understanding and conserving California's marine ecosystems (Claisse *et al.* 2018; Miller *et al.* 2018). The Kelp Forest Community Monitoring project, for example, tracks the abundance of 68 diverse marine species at permanent transects in the Channel Islands National Park; *Tethya* are the only sponge included in the survey (Kushner *et al.* 2013). In the 16 sites monitored yearly since the 1980s, *Tethya* were found at every site at least once, and were continuously present in 9 of 16 sites (as of 2011; Kushner *et al*. 2013). In some places they were abundant, with densities up to 0.43 sponges per square meter (Kushner *et al.* 2013). This long-term project also collected presence vs. absence data for over 270 species of animals and algae, and *Tethya* were among the top 10% most frequently present species across all sites (Graham 2004). Though the ecological role of *Tethya* in these habitats remains to be determined, the abundance and ubiquity of these sponges suggests they are ecologically important. Moreover, a large-scale field experiment recently found that *Tethya* populations respond dramatically to kelp forest disturbance. In a spatially replicated field experiment conducted over 9 years, large plots were seasonally cleared of giant kelp (*Macrocystis pyrifera*) to mimic natural disturbance (Castorani *et al.* 2018). The abundances of 205 species of animals and algae were monitored as response variables. *Tethya* biomass increased over 7-fold in plots with increased disturbance frequency, which was the greatest fold-change of all the diverse invertebrates, fish, and algae that were monitored. The reason for this change is unknown, but it motivates additional investigation of this taxon.

Despite this extensive monitoring of *Tethya* populations, it was previously unclear if the monitored populations included one or several species. *Tethya* from Southern and Central California were examined by de Laubenfels (de Laubenfels 1932), who described them all as *T. aurantia var. californiana*, a new variety of the Atlantic species *T. aurantium* (Pallas 1766). The variety was elevated to species status as *T. californiana* in 1993 (Sarà & Corriero 1993). More recently, Austin *et al.* (2014) described a second species of *Tethya* from California, *T. vacua*. This species was described from a single collection at Santa Rosa Island, part of the Channel Islands National Marine Sanctuary in Southern California. This sample was the only one Austin *et al.* examined from Southern California, so they proposed that *T. californiana* is found from British Columbia to Central California, while *T. vacua* is found in Southern California (South of Point Conception, see figure 1). *Tethya* have also been collected from the Gulf of California on Mexico's West Coast (Sará *et al.* 2001); Austin *et al.* (2014) did not examine these samples but based on published morphological data suggested they may represent an additional yet-to-be described species of *Tethya*, which could also range into California. It was therefore unclear how many species are found in California, whether they can be distinguished in the field, and what their ranges are.

Here I improve our understanding of these sponges using data from a SCUBA-
based survey of kelp forest sponges in California. I have collected and examined the
spicules of over 330 sponge samples to date; here I will present the results for all sponges
in the order Tethyida. By combining these fresh collections with museum samples and
publicly available data, I show that all *Tethya* known in California belong to a single
species, *T. californiana*.
The only other sponge from the order Tethyida previously known from California
is a member of the family Timeidae: *Timea authia* (de Laubenfels 1932). In contrast to
Tethya, which are large, conspicuous, and distinctive, *Timea* are encrusting and
considerably less apparent. Genetic data from other species of *Timea* have previously
revealed this genus to be problematic, with genotyped *Timea* scattered across several
different families and even orders (Cruz-Barraza *et al.* 2017, 2020). I provide the first
genetic data from *T. authia* and show that it is, in fact, in the family Tethyidae, genus
Tethytimea.
The third family in the order Tethyida is the Hemiasterellidae. These sponges,
nearly all of which have massive, vase-shaped, or upright branching morphologies, are
known primarily from the South Pacific and Indian Ocean. Several species are known
from Atlantic waters as well, but to my knowledge none have been found in the Eastern
Pacific (North or South America). I discovered a new species in Southern California, here
formally described as *Galaxia gaviotensis*. Genetic data confirm its membership in the
Hemiasterellidae; as a thinly encrusting species, it is morphologically divergent from all
previously described species in the family, so I created a new genus to contain it.
102
103 **Methods**
104
Collections
Sponges were located while SCUBA diving. All but two subtidal locations were shallow
rocky reefs; the remaining two were oil platforms. Though the survey was focused on
kelp forest habitats, I also checked three intertidal sites and floating docks in four harbors,
as shown in table S1. I attempted to photograph all sponge morphotypes found at each
location, so that presence/absence data could be compared across sites. As shown in table
S1, some locations were explored more thoroughly than others. The search times listed
are the total dive time, cumulative across all dives at a site. This is only an estimate of
search effort, as some dives were spent mainly searching and photographing sponges,
while on others considerable time was spent collecting samples. Because of the stochastic
nature of the survey, these data should be used to form hypotheses, rather than strong
conclusions, regarding species distributions.
117
Collections were made by hand with a knife. Samples were placed individually in ziplock
bags while underwater, accompanied with copious seawater. These bags were put on ice
until preservation, which was generally within 2-5 hours, but sometimes up to 12 hours
later. Samples were moved directly from seawater to 95% ethanol; in most cases, the
preservative was changed with fresh 95% ethanol after 1-3 days, and sometimes changed
again if it remained cloudy. Most samples were photographed underwater with an
Olympus TG5 before collection and photographed again in the lab. These photos
accompany this paper as supplementary data, have been archived with vouchered

specimens, and are also posted as georeferenced records on the site iNaturalist.org. Seven
samples were vouchered with the California Academy of Sciences in San Francisco;
collection numbers are listed in table S2.
129
Spicules
To examine spicules, a sponge subsample was placed in a 1.5 ml microcentrofuge tube
with household bleach for several hours, until tissue appeared to be dissolved. With the
spicules settled at the bottom of the tube, the bleach was then pipetted off and replaced
with distilled water; this was repeated several times (I found that 2-3 times was sufficient
for visualizing spicules with a light microscope, but preparing samples for scanning
electron microscopy required 5 or more rinses and worked best when the final ones were
done with absolute ethanol). In some cases, samples were centrifuged at low speed to
reduce settling time between rinses, though this increased the proportion of broken
spicules.
Spicules were imaged using a compound triocular light microscope and pictures
were taken using a D3500 SLR camera (Nikon) with a NDPL-1 microscope adaptor
(Amscope). Pictures of a calibration slide were used to determine the number of pixels
per mm, and 20-30 spicules were then measured using ImageJ (Schneider *et al.* 2012).
For scanning electron microscopy, a sample of spicules in ethanol was pipetted onto a
glass microscope slide and allowed to dry. The slide was then coated with ~20 nm of
carbon. Secondary electron images were then taken with a FEI Quanta400F Mk2.
Spicule length (or diameter) was measured as the longest possible straight line from tip to
tip, even when spicules were curved or bent. The complete distribution of measured
values for each sponge is included in the supplementary material.
I also imaged the spicular architecture in cleared tissue sections. I used a razor
blade to cut perpendicular sections that were as thin as possible by hand. These sections
were prepared in one of two ways. Some were moved from 95% to a 100% ethanol bath,
then cleared using Histoclear (National Diagnostics). Other sections were digested in a
mixture of 97% Nuclei Lysis Solution (Promega; from the Wizard DNA isolation kit) and
3% 20mg/ml Proteinase K (Promega). Many additional photos of spicules and sections
are available in the supplementary data that accompanies this paper, are archived with
museum vouchers at the Cal Academy, and have been linked to georeferenced records at
iNaturalist.org.
159
Genotyping
DNA was extracted using the Wizard DNA extraction kit (Promega), the Qiagen Blood &
Tissue kit, or the Qiagen Powersoil kit. All kits performed well on these samples in the
sense that downstream PCR was highly successful with all methods. I attempted to
amplify fragments of the cox1 mitochondrial barcoding locus for many samples. I used
the Folmer primers LCO1490 (GGT CAA CAA ATC ATA AAG AYA TYG G) and
HCO2198 (TAA ACT TCA GGG TGA CCA AAR AAY CA) (Folmer *et al.* 1994),
which worked for all samples except *T. authia*. For this species, I also tried the primers
from Rot et al. (Rot *et al.* 2006), again without success. I also amplified portions of the
28S rDNA nuclear locus. For *T. authia* and *G. gaviotensis*, I amplified the C region using
primers C2 (GAA AAG AAC TTT GRA RAG AGA GT) and D2 (TCC GTG TTT CAA
GAC GGG) (Chombard *et al.* 1998). These primers amplified bands of multiple sizes for

multiple *Tethya* vouchers, but I was able to amplify a larger fragment containing the C
region in *Tethya* using primers for the D1-D2 region: Por28S-15F (GCG AGA TCA
CCY GCT GAA T) and Por28S-878R (CAC TCC TTG GTC CGT GTT TC) (Morrow *et*
al. 2012). For some samples, a portion of the 18S locus was also amplified using the
primers SP18aF (CCT GCC AGT AGT CAT ATG CTT) and 600R18S (CGA GCT TTT
TAA CTG CAA) (Redmond *et al.* 2013). All primer sequences are listed 5' to 3'.
PCR was performed in a Biorad T100 thermocycler with the following
conditions: 95C for 3 min, followed by 35 cycles of 94C for 30 sec, 52C for 30 sec, 72C
for 1 min, followed by 72C for 5 minutes. The 28S C region was amplified with a 57C
annealing temperature instead of 52C. PCR was performed in 50 μl reactions using the
following recipe: 24 μl nuclease-free water, 10 μl 5x PCR buffer (Gotaq flexi, Promega),
8 ul MgCl, 1 μl 10mM dNTPs (Promega), 2.5 μl of each primer at 10 μM, 0.75 bovine
serum albumin (10 mg/ml, final conc 0.15 mg/ml), 0.25 μl Taq (Gotaq flexi, Promega), 1
μl template. ExoSAP-IT (Applied Biosystems) was used to clean PCRs, which were then
sequenced by Functional Biosciences using Big Dye V3.1 on ABI 3730xl instruments.
All PCR products were sequenced in both directions, and a consensus sequence was
constructed using Codon Code v.9 (CodonCode Corporation). Blastn was used to verify
that the resulting traces were of sponge origin. All sequences have been deposited in
Genbank; accession numbers are listed in table S2.
191
Genetic analysis
I used the NCBI taxonomy browser to compile all data from samples identified as
belonging to the order Tethyida. I also used blastn to search for additional sequences that
appeared to fall within this order. Some accessions identified as Tethyida were then
excluded because they were more closely related to other orders of sponges or were from
a different region of the gene being analyzed.
198
Supplementary table S4 lists every Genbank accession found and explains the reasons
why any were excluded. Together, included data are from 23 different publications and
several unpublished datasets (Chombard 1998; Cruz-Barraza *et al.* 2017, 2020; Dat *et al.*
2018; Erpenbeck *et al.* 2007, 2012, 2016; Heim & Nickel 2010; Heim *et al.* 2007c; a; b;
Idan *et al.* 2018; Kober & Nichols 2007; Lavrov *et al.* 2005; Morrow *et al.* 2012; Nichols
2005; Redmond *et al.* 2013; Regueiras *et al.* 2019; Riesgo *et al.* 2013; Shaffer *et al.* 2018;
Sorokin *et al.* 2019; Thacker *et al.* 2013; Zhang *et al.* 2016). Sequence alignments were
produced in Codon Code v.9 (CodonCode Corporation). Phylogenies were estimated with
maximum likelihood using IQ-Tree (Nguyen *et al.* 2015; Trifinopoulos *et al.* 2016). I
used the Ultrafast bootstrap (Hoang *et al.* 2018) to measure node confidence. Phylogenies
were produced from the IQ-Tree files using the Interactive Tree of Life webserver
(Letunic & Bork 2019). Figures were made ready for publication using R (r-project.org)
and/or Gimp (gimp.org).
212
213   **Results**
214
I used SCUBA to investigate the sponge fauna at 47 sites in Southern and Central
California (figure 1, table S1). The spicular morphology of 334 sponges from this project
have been analyzed to date, and several were found to belong to the order Tethyida. After

follow-up collections targeting specific species and locations, a total of 13 specimens
could be assigned to the Tethyida. Below, I explain the results of the phylogentic analysis
of these samples, and what these phylogenies tell us about other taxa. This is followed by
a systematic section with additional details on each species.
222
223 *Genetic Results*
224
To perform genetic analysis of newly collected samples, I focused on the three loci most
frequently used in sponge systematics: cox1 and the large (28S) and small (18S)
ribosomal subunits. As detailed in the methods, not all loci were successfully amplified
from all species. When possible, I sequenced sponges from both Central and Southern
California for comparison. I then compiled all published records in Genbank for the order
Tethyida for comparative analysis (see methods).
Of the three loci investigated, the largest number of public records were available
at the 28S rDNA (figure 2). Considering only the previously published data, the
phylogeny is very similar to a recently published phylogeny (Sorokin *et al.* 2019), though
my interpretation of these results differ on several key points. In addition, I excluded
records that lacked the highly variable C region. Genbank records at 28S differ greatly in
read length: though phylogenetic methods can accommodate missing data, it is preferable
to have at least a single region shared by all records. Though this led to the exclusion of
several species, it also improved the stability of the phylogeny. One notable improvement
is the placement of *T. californiana*: the published phylogeny of Sorokin et al. (2019)
found that *T. californiana* nested within the European *T. hibernica/citrina/norvegica*
clade at 28S, but was in the sister group to this clade at cox1. The 28S sequence used in
their analysis did not include the C2 region. I sequenced an 824 bp region that did include
C2 from a freshly collected *T. californiana* from Southern California, and this sequence
has a phylogenetic position consistent with what is found in other gene trees.
The 28S phylogeny recovers the Family Tethyidae as monophyletic, if the
systematic placement of two taxa are revised (as detailed below). Three well supported
clades of Tethyidae are recovered. One clade includes all records from the genus *Tethya*
save one: *T. stellagrandis*. This species falls into a second well supported clade that
includes nearly all other Tethyidae with available data (eight additional genera have no
available sequence data at this locus). The third and final clade consists of *Tethytimea*
carmelita and an unidentified sample. The unidentified sample is from a voucher that is
now lost, but personal communications with the collector, Scott Nichols, indicate it was
likely from Central California. I sequenced the 28S locus in two newly collected
individuals of *Timea authia*: one from Central California (Monterey Bay) and one from
Southern California (Carpinteria Reef). Both of these sequences are identical to the
unidentified sample over the aligned 816 bp, so it is likely the unidentified sample is also
of this species. These three samples form a clade with *Tethytimea carmelita* with 95%
bootstrap support. This result, together with the genetic data from 18S and the
morphological data below, leads me to reassign *Timea authia* to the genus *Tethytimea* (as
detailed in the Systematics section).
Only one species of Tethyidae is not included in the 28S clade discussed above:
Laxotethya dampierensis, which is the type species of its genus. This species falls into the
sister clade to the Tethyidae. This sister clade is itself composed of two clades: one that

includes all species of the Hemiasterellidae with available data, and another that includes
L. dampierensis, an undescribed species identified as "Tethyidae" in Genbank, and
Galaxia gaviotensis gen. nov. sp. nov.. The "Tethyidae" sample, collected in Ireland, is in
the process of being described by others. Based on a personal communication with
Bernard Picton, the collector, it is encrusting and has some traits in common with *Adreus*
(F. Hemiasterellidae), but does not fit within any known genus. Based on the
phylogenetic position of this clade, the simplest way to resolve its higher taxonomy
would be to move *Laxotethya* to the Hemiasterellidae, making this family the sister clade
to the Tethyidae (figure 1). I formally suggest this reassignment here (though I did not
examine the *L. dampierensis* voucher that was sequenced, I did verify that it was likely to
be correctly identified; it is present at the Western Australia Museum as voucher
WAMZ11871, identified by Dr. Jane Fromont). This would place the undescribed Irish
sponge and the new California sponge in the family Hemiasterellidae as well.
277   The reassignment of *Timea authia* adds to a growing list of taxonomic problems
for the genus *Timea* (Cruz-Barraza *et al.* 2017, 2020), which is currently in its own
family (Timeidae). The only other *Timea* record at 28S that is identified to species, and
includes the C region, is *Timea lowchoyi* (AY561871). This sample is not shown on the
phylogeny, because it is nested within the order Axinellida (Sorokin *et al.* 2019). The
remaining samples, all *Timea sp.*, do not form a single clade. These results could indicate
that 1) samples have been misidentified, or 2) the family is highly polyphyletic and in
need of revision. My results for *T. authia*, and the published results of others, suggest that
the need for revision is real, even if some samples have been misidentified (Cruz-Barraza
et al. 2020).
287   The gene tree of the Folmer barcoding region of the cox1 gene is consistent with
the key results of the 28S tree (figure 3). The family Hemiasterellidae forms a single
clade that includes *G. gaviotensis* sp. nov. The genus *Tethya* again forms a well
supported clade; unfortunately, no sequences from Tethyidae outside of Tethya are
available thus far (attempts to amplify this locus from *T. authia* repeatedly failed). *Timea*
sequences from Genbank again illustrate that this genus requires revision. Two sequences
are not shown: *Timea unistellata* from Ireland (KC869427) and an unidentified *Timea sp.*
(AY561968). Both of these sequences fell within the order Poecilosclerida. This was
recently reported by Cruz-Barraza et al. (2020), who show that these two *Timea* are not
closely related to each other within that order (their figure 6). The remaining *Timea*
sequences do form a well supported clade, but the placement of this clade is different
than for the 28S locus. None of the *Timea* vouchers sequenced at cox1 have also been
sequenced at 28S, so the most likely explanation is that there are some *Timea* that are
outside of the Hemiasterellidae + Tethyidae (as shown in the 28S tree) and other *Timea*
that are more closely related to Tethyidae than Hemiasterellidae (as shown in the cox1
tree). Note that the placement of the *Timea* clade in the cox1 phylogeny is consistent with
the placement of the morphologically similar genus *Tethytimea* in the 28S phylogeny.
304   The cox1 phylogeny fails to support species status for *Tethya vacua*. Three
sequences were previously available for *Tethya californiana*; two of these are from
British Columbia, while the other is from Central California. The Central California
sample is somewhat uncertain: it is thought to be from Carmel Bay, California, which
contains the type locality for *T. californiana*. However, the voucher and collection
information are lost. I therefore collected a new sponge from Monterey Bay, about 8 km

North of the type location, and sequenced it for comparison (MT139598 in figure 3). A *T.*
vacua paratype was also sequenced, and is genetically identical to the Monterey sponge
at the 650 bp Folmer region of cox1. Three additional sponges from island and mainland
locations in Southern California were likewise identical. The British Columbia sponges
are slightly divergent from this California haplotype (0.6% absolute sequence
divergence). This divergence was previously noted vs. the Central California sponge
from Carmel, and was used to support species status for this population under the name *T.*
leysae (Heim & Nickel 2010). Austin *et al.* (2014) synonymized *T. leysae* with *T.*
californiana; they argued that this level of genetic divergence provides limited support
for species status.
The phylogeny at the 18S locus is less resolved than the other loci due to the slow
evolutionary rate at this locus (figure 4). I sequenced ~850 bp from *Tethya californiana*
from Central and Southern California, and again found that they had identical haplotypes.
I was able to obtain a homologous 600 bp from the *T. vacua* paratype and it was identical
as well. Furthermore, a 1600 bp region was previously sequenced from several *Tethya*
from Naples Reef (Southern California) and the *T. californiana* that is likely from Carmel
(Central California); one of these Southern California sponges is identical to the Carmel
sponge over this entire region, while the others are nearly identical. These data again fail
to support species status for *T. vacua,* nor do they support genetic differentiation between
Southern and Central California in this genus.
The 18S phylogeny is consistent with the placement of *T. authia* in the Tethyidae.
No other *Tethytimea* have been sequenced at this locus, but *T. authia* sequences form a
(poorly supported) clade with the only Tethyidae sequences available outside of the
genus *Tethya*. Note that that there is a published sequence that is identical to *T. authia* at
18S (EF654532): this is from the same voucher that is identical at 28S, which is likely to
have been a *T. authia*.
The 18S phylogeny also supports the placement of *G. gaviotensis* sp. nov. in the
Hemiasterellidae. The most supported tree at 18S placed all Hemiasterellidae in a single
clade, but this clade had only 33% bootstrap support, and is collapsed in figure 4. Despite
this, *G. gaviotensis* sp. nov. clusters with *Adreus fascicularis* with good support,
consistent with the 28S phylogeny.
The 18S phylogeny once again illustrates the need for revision of *Timea*. Though
all three available sequences grouped within the Tethyida, two are outside the
Hemiasterellidae + Tethyidae, while the other is weakly supported as within the
Hemiasterellidae.
Finally, these phylogenies indicate something potentially interesting regarding
Axos cliftoni. This species is placed within the Hemiasterellidae, and sequences at 18S
and 28S are consistent with this placement. Sequences at cox1, however, are not shown,
because preliminary analyses showed them to group in the order Axinellida. This was
found by Cruz-Barraza et al. (2020) as well, as shown in figure 6 of their paper. This is
perplexing, because the cox1 sequences are annotated as being from the same vouchers as
the rDNA sequences, and come from different collections in two countries. Voucher
MT5.2015 is from Vietnam and was sequenced at two loci (18S = KY947247, cox1 =
KY947263). Voucher G300111 is from Australia and was sequenced at three loci (28S =
AY626308, 18S = EF654523, cox1 = AY561974). The cox1 sequences from these two
samples are extremely similar, but group with family Stelligeridae in the order Axinellida

(Cruz-Barraza *et al.* 2020). The 18S sequences from these vouchers are similar, and form
a clade within the Tethyida (figure 4); the 28S sequence from Australia falls within the
Hemiasterellidae (figure 3). Perhaps this conflict is due to errors with sequence
annotations in Genbank, but alternatively, it may suggest horizontal gene transfer or some
other biologically interesting phenomena.
361
362 **Systematics**
363
364 Order TETHYIDA Morrow & Cárdenas, 2015
365
366 Family Tethyidae Gray, 1848
367
368 Genus *Tethya* Lamarck, 1815
369
370 *Tethya californiana* de Laubenfels, 1932
371
372 **Synonymy.**
373 *Tethya aurantia var. californiana* de Laubenfels, 1932
374 *Tethya vacua* Austin et al. 2014
375
Material Examined. CASIZ 235109, Hopkin's Marine Station, Monterey CA (34.47182,
-120.14262), 9/16/19. CASIZ 235108, Naples Reef, Santa Barbara CA (34.42212, -
119.95154), 7/31/19. CASIZ 235106, Elwood Reef, Santa Barbara CA (34.41775, -
119.90150) 4/17/19. CASIZ 235107, Santa Cruz Island, CA (33.98378, -119.63910),
4/26/19. TLT 361, Santa Rosa Island, CA (33.89966, -120.10735) 10/9/19. CASIZ
184729, Santa Rosa Island, CA (34.00000, -120.00170), collected by France & Efford,
12/24/81 (*T. vacua* paratype). CNPGG 1199, Ensenada, Bahia Topolobampo, Baja
California, Mexico, 4/8/11. CNPGG 0012, Cabo San Miguel, Gulf of California, Mexico,
3/5/85. (For tabular format, see supplemental table 2).
385
Description.
Macroscopic features. Approximately hemispherical, with broad gutters between
tubercles when relaxed. The sponge constricts upon collection, closing the gutters and
placing the tubercles in contact. The width and depth of the gutters varies greatly in field
photos, as does color (from yellow to orange, but sometimes covered in a dark red growth
of apparent algae). As this species has been well described elsewhere (Austin *et al.* 2014),
I focused on improving our understanding of the quantitative aspects of spicules that have
been shown to vary among samples.
394
395 *Spicules.* To assess the possibility of multiple species of *Tethya* within the study region, I
396 quantified two aspects of spicular morphology found to vary in previous studies (Austin
397 *et al.* 2014; Heim & Nickel 2010; Sará *et al.* 2001). The first concerns the morphology of
398 the megascleres. Austin et al. (2014) divided megascleres into three categories defined as
399 follows:
400
401 -anisostrongyles: distal end (foot) smaller diameter than proximal end (head).

402   -strongyloxeas: fusiform, distal end (foot) a point; sides narrow toward rounded end
403   (head).
404   -styles: straight sides, distal end (foot) a point; diameter of rounded end (head) equal to
405   that of shaft.
406
After examining a large number of spicules, I found strongyloxeas and styles to be part of
a continuous spectrum. I measured the width of each spicule at the head and the widest
point, and some were very fusiform (widest point more than 4 times wider than head),
some were clearly styles (widest point equal to head), but most spicules were
intermediate. I therefore grouped these two types together. Anisostrongyles were easier to
distinguish, as many spicules clearly lacked a sharp point, though once again there were
spicules that were intermediate and hard to classify. I therefore present data on all
megascleres together, and also present data on anisostrongyles separately (table 1).
Regardless of method, the distributions of megasclere dimensions broadly overlapped
among samples. ANOVA of all megascleres considered together revealed no significant
differences among samples (F=1.73, p=0.13). One sample, from the Gulf of California
(Mexico), was found to lack anisostrongyles almost completely, while anisostrongyles
comprised 27-45% of megascleres in all other samples (see remarks).
The other quantitative measure that has been found to vary among *Tethya* in this
region relates to megasters (star-shaped spicules). By dividing the ray length by the
centrum diameter, a parameter called R/C can be calculated for each aster (Austin *et al.*
2014). I did this for 40-50 asters per sponge, and used ANOVA to determine if there were
significant differences among samples (there were, F=12.14, p=1.81e-10). Tukey's honest
significant differences test was then use to determine which samples were different from
others at the 95% family-wide confidence level. Note that the *T. californiana* sample
from Hopkin's Marine Station in Central California (near the type location for *T.*
californiana) was not significantly different from the *T. vacua* paratype (table 1).
429
Remarks. Though *Tethya* are the best studied sponge in California from an ecological
perspective, their taxonomy has remained unresolved. *Tethya aurantia* var. *californiana*
de Laubenfels was described in 1932, with Carmel Bay, in Central California, as the type
location (de Laubenfels 1930). The variety was raised to species status in 1993 (Sarà &
Corriero 1993). Heim and Nickel (2010) then split this taxon, describing those from
British Columbia as a separate species, *T. leysae*. *Tethya leysae* was synonymized with *T.*
californiana by Austin *et al.* (2014), who proposed that this species ranged from British
Columbia to Central California. In the same monograph, Austin et al. erected *T. vacua* for
Southern California, based on a single collection from Santa Rosa Island (figure 1). They
also implied that the sponges described as *T. californiana* from the Gulf of California, at
the Southern range limit, were likely a different species. I wished to confirm or refute the
validity of the name *T. vacua*, and assess the possibility of an additional species in the
South.
Genetic data do not support species status for *T. vacua*. In the same monograph
where *T. vacua* was described, Austin et al. (2014) argued that 0.6% absolute sequence
divergence at cox1 was insufficient to support species status for *T. leysae*. I show that *T.*
vacua are even more similar (0% sequence divergence) to *T. californiana* at this same
locus. The same is true at the 18S nuclear locus.

The spicules of Central and Southern California *Tethya* are likewise
indistinguishable. The only spicule data used to support *T. vacua* was that the one sample
analyzed had asters with a higher R/C value than the *T. californiana* syntype. No
statistical test was used, however, and another *T. californiana* sample from Monterey Bay
(CASIZ 067731, table 14 of Austin *et al.* 2014) was very similar to the *T. vacua* sample.
Here I show that the R/C values of the *T. vacua* paratype are not significantly different
from another sponge collected in Monterey Bay, near the type location for *T. californiana.*
Though not statistically significant, average values are slightly larger for 2 of the 3
Southern California samples analyzed here vs. the Central California sample, and this is
consistent with a geographic cline in R/C value as previously reported (Austin *et al.*
2014). Austin et al. show that R/C values are shortest in cold British Columbian waters,
and increase towards the South (and again used this cline to argue against species status
for *T. leysae*). R/C values of the Ensenada sponge analyzed here have the largest values,
consistent with this cline.
The only other morphological difference reported for *T. vacua* are the vacuoles
that are its namesake. I compared tissue sections of sponges from Monterey (Central
California), Naples Reef (Southern California), and Santa Rosa Island (Southern
California, near the holotype location for *T. vacua*), to look for these vacuoles (see
supplementary data). I failed to find vacuoles in any sponge, but these negative data
should not be considered conclusive. I did not have access to the *T. vacua* holotype as a
positive control, so it is possible that my methods were inadequate to see them. In any
case, I feel that the lack of any spicular characters, combined with the lack of genetic
differentiation, clearly indicates that these taxa should be synonymized.
I was also interested in whether there is an additional species of *Tethya* at the
Southern end of the *T. californiana* range. Two sponges from Pacific Mexico were
graciously shared with me by Patricia Gómez, Porifera Curator at the National
Autonomous University of Mexico. One is from Ensenada, just South of the US/Mexico
border, where habitats are similar to California kelp forests. The spicules of this sponge
were indistinguishable from those in Southern California (table 1). The other was from
Cabo San Miguel, in the Gulf of California; this is beyond the range of the giant kelp
Macrocystis pyrifera, in an ecosystem quite different from those on California shores.
Sponges from this collection were described as having several morphological differences
compared to the *T. californiana* holotype, the most significant of which was having only
strongyloxeas instead of anisostrongyles and strongyloxeas (Sará *et al.* 2001). Sará,
Gómez, and Sára (2001) did not feel these differences warranted species status. In my
reanalysis of one of these sponges, I replicate their finding of (nearly) exclusive
strongyloxeas, and find that the R/C values of asters are somewhat anomalous compared
to the clinal expectation. I attempted to sequence fragments of cox1 and 18S from these
Mexican samples, but was unable to amplify any DNA (I also attempted to sequence
DNA from other archived samples of Mexican *Tethya* species, again without success).
Collection of fresh material for sequencing is therefore highly recommended. It remains
unconfirmed if the Gulf of California *Tethya* are *T. californiana*, but the similarities of
the Ensenada sponge to California samples makes it unlikely that this Gulf of California
morphotype range into the state of California.
492

Geographic distribution. *Tethya californiana* was previously described as ranging from
British Columbia in the North to Central California in the South (Austin *et al.* 2014).
Genetic data now extend this range to Southern California, and morphological
comparisons extend it to at least Ensenada, Mexico. Consistent with published reports,
this sponge was common in kelp forest habitats in California. I observed it at 19/47
(40%) of kelp forest locations surveyed. There are some indications it may be more
abundant in colder waters (because of the nature of the survey, these patterns should be
considered hypotheses only; see methods). Water temperatures are colder in Central
California than Southern California, and within the Santa Barbara Channel there is a
strong gradient, with colder waters in the West (Claisse *et al.* 2018). *Tethya californiana*
were present at 2 of 3 sites in Central California, and 4 of 4 sites on the Western-most
islands of San Miguel and Santa Rosa. In contrast, it was seen at only 2 of 8 location on
Anacapa Island, and 2 of 7 locations in Los Angeles County. It was not present at the two
oil platforms surveyed, nor on floating docks in the four harbors surveyed. I did not find
it in the 3 intertidal locations checked, but pictures posted by others to the site iNaturalist
show that the sponge is sometimes present in the intertidal in Northern and Central
California (and the holotype collected in Central California in 1926 was an intertidal
sponge).

Genus *Tethytimea* de Laubenfels, 1936

Diagnosis. Tethyidae with thinly encrusting body. Megasclere skeleton composed of
parallel bundles of tylostyles, with or without styles, ending in tubercles. Megasters are
oxyspherasters and tylasters, heterogeneous in size and sometimes with giant spicules.
Micrasters are tylasters. Other microscleres, occasionally present, are spheres. (Modified
from Cruz-Barraza *et al.* (2017) to include styles and tylasters among megasters.)

Tethytimea authia de Laubenfels, 1930 (comb. nov.)

Synonymy.
Timea authia de Laubenfels, 1930

Material Examined. CASIZ 235111, Perkin's Reef, Monterey CA (36.62920, -
121.92031), 11/24/19. CASIZ 235110, Carpinteria Reef, Santa Barbara CA (34.39163, -
119.54169), 7/31/19. TLT 364, Saddleback Ridge, Santa Cruz Island CA (34.03817, -
119.52470), 10/18/19.

Description.
Macroscopic features. An encrusting sponge; samples vary from 0.5 - 2.0 mm thick.
Color yellow/orange when alive (figure 5). The surface is sculpted by gutters in situ, but
the sponge contracts when collected, and gutters are no longer visible. Scattered,
mushroom-like papillae present on surface, approximately 1-3 mm in diameter. No other
sponges were located with these features in the current survey: it is therefore possible that
this species can be identified from field photos, at least within the Southern California
portion of its range. Note that these characters are difficult to see in the field, and more
apparent in macro photos.

539
Spicules.
Megascleres are tylostyles with oval heads, styles with an enlarged band near the head,
and simple, unadorned styles (figure 6). Microscleres are asters. The most common aster
has thick, non-tapering rays with enlarged tips, usually with spines at the tips (tylasters).
Others have cone-shaped rays ending in sharp tips (oxyspherasters). Intermediates are
also found, with tapering rays that are fairly cone-shaped, but ending in enlarged tips with
spines. Spicule measurements are shown in table 2, and compared to published data
(Carballo and Cruz-Barraza 2006). Carballo and Cruz-Barraza (2006) reanalyzed the
holotype and used SEM to provide more detail regarding spicule shape. Data from my
three samples are entirely consistent with their data, qualitatively and quantitatively.
550
Tethytimea are described as having two size classes of asters: larger "megasters" and
smaller "micrasters". *Timea* are described as having only "asters". I quantified the size
distribution of asters to determine whether there was more than one size class (figure 7).
One sample (TLT364) seemed to have two size classes, with "megasters" 13-30 μm and
"micrasters" 3-7 μm (though 2% of asters were intermediate). The other two sponges had
two modes, but the distributions of the size classes were overlapping. I used Hardigan's
Diptest to test for unimodality, which was rejected for all three samples (p<0.0001 for all).
The larger sized asters included both oxyspherasters and tylasters, with tylasters more
common. Only tylasters were seen among the smallest asters.
560
Microscopic features.
Perpendicular sections show tissue packed with asters (figure 6). Megascleres
(styles/tylostyles) are found in projecting bundles, tips up, that pierce the surface of the
sponge. Bundles fan-out as they rise through the sponge, resulting in bouquets of spicules
at the surface. Other megascleres are found parallel to the substrate in the lower third of
the tissue section.
567
Geographic distribution. The range of this species in the United States was previously
described as Southern California only (Lee *et al.* 2007). This range is here extended
North to Monterey Bay in Central California. To the South, samples from as far away as
Chile have been assigned to this species, though no molecular data is yet available for
confirmation (Carballo and Cruz-Barraza 2006, Desqueyroux-Faúndez 1972). The depth
range of the species was previously described as "intertidal to deep water" (Lee *et al.*
2007). All three samples collected in the current work were from the shallow subtidal, 3.5
– 11 m in depth. It was not found in the intertidal, but sampling in the intertidal was very
limited. The holotype was collected from the Southern California intertidal in 1930 (de
Laubenfels 1932).
578
Remarks. Genetic data strongly support membership for this species in the Tethyidae
and not the Timeidae. As discussed in the phylogentic results, it is currently difficult to
ascertain the differences between *Timea* and *Tethyatimea*, because the species
membership of each genus is incompletely ascertained. One proposed distinction has
been the presence of size classes of spicules among the *Tethytimea* (Cruz-Barraza *et al.*
2017, Sará 2002). I find that samples of *T. authia* have a bimodal distribution of aster

sizes, consistent with this diagnosis. The overlapping size ranges may partially explain
why this sponge has not been diagnosed as a *Tethyatimea* in the past.
587
588 Family Hemiasterellidae Lendenfeld, 1889
589
590 *Galaxia* gen. nov.
591
592 **Type species.** *Galaxia gaviotensis* sp. nov.
593
594 **Diagnosis.** Hemiasterellidae with thinly encrusting growth habit. Megascleres include
595 tylostyles, styles, and oxeas. These are found in bundles parallel to the substrate and
596 oriented vertically, piercing the sponge surface. Microscleres are asters.
597
598 **Etymology.** From the latin galaxia, referring to the milky way. Like a galaxy, the type
599 species of the genus is packed with a diversity of stars.
600
601 *Galaxia gaviotensis* sp. nov.
602
603 **Material Examined.** Holotype, CASIZ 235112, Arroyo Quemado Reef, Santa Barbara
604 County, California, United States of America (34.46775, -120.11905), 15 meters depth,
605 7/29/19.
606
607 **Etymology.** The area where this sponge was collected is known as the Gaviota Coast,
608 and is the longest remaining undeveloped part of the Southern California coastline. As a
609 common name, I suggest "Gaviota galaxy sponge".
610
611 **Description.**
612 *Macroscopic features*. The sponge formed a pale-yellow crust with surface dimensions
613 approximately 10 x 5 cm. Thickness only 200-650 μm. Field photos (figure 8) show
614 darker spots on the surface; these disappeared in sampled fragments, so were likely
615 oscules or pores that contracted when disturbed.
616
Microscopic features. Styles comprised 70% of megascleres. Some were tylostyles with
well-formed oval heads, while others had weak heads (subtylostyles), and others had
lumpy, irregular heads (fig. 9). Many also had subterminal heads or weak bands of
slightly increased width near the head of the style. About 13% were unadorned, simple
styles. Length varied greatly, from 176-1330 μm (mean = 751, N=45). Width from 2-9
μm, with a mean of 5.3 μm. The other 30% of megascleres were asymmetrical oxeas.
These were sharply pointed at both ends, and gently bent at the thickest point, which was
closer to one end. A minority (20%) of these had very weak centrylote swelling at the
point of inflection. Length was less variable than for styles, from 174-464 μm (mean =
258, N=19); width 2-5 μm, with a mean of 3.7 μm.
Microscleres consisted of abundant and varied euasters. The most common type
had thick, untapered rays ending in enlarged, spiny bouquets ("tylasters"). The ratio of
ray length to centryole size varied greatly (fig. 9). Some were also spined along the rays,
instead of merely at the tips. In some asters, the rays were greatly reduced, to

hemispherical bulges around a large centrum, sometimes bearing spines and sometimes
nearly smooth. Asters with conical rays (oxyspherasters) were also present. These ended
in spined bouquets as well, but the bouquets were sometimes very small, resulting in
nearly smooth, tapering, conical rays. Mean aster diameter was 10.6 μm (range 4 - 22 μm,
N=62); sizes were not clearly different among types.
636
Spicule arrangement. Perpendicular sections through the sponge tissue revealed that it
was packed with asters (figure 8). These formed a thick surface crust, but were also
abundant throughout; no pattern of arrangement by type could be discerned. Megascleres
were most dense in the lower half of the sponge, running parallel to the substrate; many
were found in bundles. Bundles of megascleres were also oriented vertically, points up,
piercing the surface of the sponge.
643
Geographic distribution. This sponge has been found at only a single location: Arroyo
Quemado reef, approximately 0.15 km offshore of the small community of Arroyo
Quemado (34.468801, -120.121233). This is an area of rocky reef and kelp forest that is
part of the Santa Barbara Coastal Long-Term Ecological Research Station
(sbclter.msi.ucsb.edu). It was found on "transect 3", which is a flat area 7.5 - 9.5 m deep,
strewn with cobbles and boulders. It was growing on a single boulder. I made a total of 8
collecting dives at this reef (5 hours of cumulative search time) but no additional
individuals were found. Seven months after collecting the initial sample, the transect 3
area was re-visited in an attempt to relocate the original sponge, without success. Many
of the boulders at the site were coated in a fine layer of silt at this time, making it likely
that the sponge was still present but concealed.
655
Remarks. Most encrusting sponges in the Tethyida are placed in the genus *Timea*. *Timea*
is rejected in the current case due to the presence of oxeas and genetic data that places the
species in the Hemiasterellidae. Encrusting Hemiasterellidae are known from only two
genera: *Leptosastra* and *Liosina*. *Leptosastra* is monotypic, with the unusual *L.*
constellata having acanthostyles, and tornotes; this is a very poor fit for the current
species. *Liosina* includes at least one encrusting species (though thickly encrusting, with
chimneys) and sometimes have oxeas; however, these species lack asters. *Hemiasterella*
include some species with oxeas, but these species are all massive, digitate, or vase-
shaped. Thus, this species has become the type for a new genus.
665
666 **Conclusions**
667
By combining molecular and morphological approaches, this work significantly improves
our understanding of Californian Tethyida. Taxonomic updates include the discovery and
description of *G. gaviotensis* gen nov. sp. nov., the reassignment of *Timea authia* to the
Tethyidae as *Tethytimea authia* comb. nov., and the synonymy of *Tethya californiana*
and *Tethya vacua*. Genetic data were key to these revisions, and, in my opinion, are
indispensable in poriferan taxonomy in general. It seems unwise to erect a new species
based solely on the presence of a single morphological character with unproven utility,
like the presence of vacuoles. However, it is also unfortunate that morphological data is
not publicly available for many of the sequenced vouchers in Genbank. This greatly

reduces the utility of publicly available sequence data. Here, I have provided a combination of morphological and genetic data that will be useful for ongoing efforts to revise poriferan taxonomy. My hope is that future work on the ecology and evolution of these species improves both our understanding of the near-shore California ecosystem and the evolution of these interesting animals.

Acknowledgements

I am grateful for the help and support of many people in UCSB's Marine Science Institute and Diving & Boating Program, especially Robert Miller, Clint Nelson, Christoph Pierre, Frankie Puetzer, Christian Orsini, H. Mark Page, and Alecia Dezzani. Steve Lonhart (NOAA) and Shannon Myers (UCSC) were instrumental in facilitating collections in Central California, and the Natural History Museum of Los Angeles' DISCO program facilitated collections in Los Angeles County. Christina Piotrowski (California Academy of Sciences) and Patricia Gómez (National Autonomous University of Mexico) graciously shared vouchers from their collections.

Funding declaration

Financial support was provided by UCSB and by the National Aeronautics and Space Administration Biodiversity and Ecological Forecasting Program (Grant NNX14AR62A); the Bureau of Ocean Energy Management Environmental Studies Program (BOEM Agreement MC15AC00006); the National Oceanic and Atmospheric Administration in support of the Santa Barbara Channel Marine Biodiversity Observation Network; and the U.S. National Science Foundation in support of the Santa Barbara Coastal Long Term Ecological Research program under Awards OCE-9982105, OCE-0620276, OCE-1232779, OCE-1831937. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

Austin, W.C., Ott, B.S., Reiswig, H.M., Romagosa, P. & Mcdaniel, N.G. (2014) Taxonomic review of Hadromerida (Porifera, Demospongiae) from British Columbia, Canada, and adjacent waters, with the description of nine new species. *Zootaxa* 3823, 1–84.

Bell J.J. (2008) The functional roles of marine sponges. *Estuarine, Coastal and Shelf Science* 79, 341-353.

Carballo, J.L. & Cruz-Barraza, J.A. (2006) The genus Timea (Porifera: Timeidae) in the Eastern Pacific Ocean: revision of type material, new species and intraspecific variability. *Cahiers de Biologie Marine* 47, 301–320.

Castorani, M.C.N., Reed, D.C. & Miller, R.J. (2018) Loss of foundation species: disturbance frequency outweighs severity in structuring kelp forest communities. *Ecology* 99, 2442–2454.

Chombard, C. (1998) Les Demospongiae à asters: phylogénie moléculaire et homologie
morphologique. Muséum National d'Histoire Naturelle, Paris.

Chombard, C., Boury-Esnault, N. & Tillier, S. (1998) Reassessment of homology of
morphological characters in tetractinellid sponges based on molecular data.
Systematic Biology 47, 351–366.

Claisse, J.T., Blanchette, C.A., Dugan, J.E., Williams, J.P., Freiwald, J., II, D.J.P.,
Schooler, N.K., Hubbard, D.M., Davis, K., Zahn, L.A., Williams, C.M. & Caselle,
J.E. (2018) Biogeographic patterns of communities across diverse marine
ecosystems in southern California. *Marine Ecology* 39, e12453.

Cruz-Barraza, J.A., Vega, C., Ávila, E. & Vázquez-Maldonado, L.E. (2017) Integrative
taxonomy reveals the first record and a new species for the previously monotypic
genus Tethytimea (Tethyida: Tethyidae) in the Gulf of Mexico. *Zootaxa* 4226, 113–
125.

Cruz-Barraza, J.A., Vega, C., Rützler, K. & Salas-Castañeda, M. (2020) Morphological
and molecular taxonomy of Timea (Porifera: Timeidae) from the Gulf of Mexico
with the description of a new species and re-description of T. hechteli. *Journal of the*
Marine Biological Association of the United Kingdom, 1–13.

Dat, T.T.H., Steinert, G., Thi Kim Cuc, N., Smidt, H. & Sipkema, D. (2018) Archaeal
and bacterial diversity and community composition from 18 phylogenetically
divergent sponge species in Vietnam. *PeerJ* 6, e4970.

Desqueyroux-Faúndez, R. 1972. Demospongiae (Porifera) de la costa de Chile. *Gayana*,
20: 1-71.

Erpenbeck, D., Duran, S., Rützler, K., Paul, V.J., Hooper, J.N.A. & Wörheide, G. (2007)
Towards a DNA taxonomy of Caribbean demosponges: a gene tree reconstructed
from partial mitochondrial CO1 gene sequences supports previous rDNA
phylogenies and provides a new perspective on the systematics of Demospongiae.
Journal of the Marine Biological Association of the United Kingdom 87, 1563–1570.

Erpenbeck, D., Hall, K., Alvarez, B., Büttner, G., Sacher, K., Schätzle, S., Schuster, A.,
Vargas, S., Hooper, J.N.A. & Wörheide, G. (2012) The phylogeny of halichondrid
demosponges: past and present re-visited with DNA-barcoding data. *Organisms*
Diversity & Evolution 12, 57–70.

Erpenbeck, D., Voigt, O., Al-Aidaroos, A.M., Berumen, M.L., Büttner, G., Catania, D.,
Guirguis, A.N., Paulay, G., Schätzle, S. & Wörheide, G. (2016) Molecular
biodiversity of Red Sea demosponges. *Marine Pollution Bulletin* 105, 507–514.

Folmer, O., Black, M.B., Wr, H. & Vrijenhoek, R.C. (1994) DNA primers for

amplification of mitochondrial Cytochrome C oxidase subunit I from diverse
metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3, 294–299.

Graham, M.H. (2004) Effects of local deforestation on the diversity and structure of
Southern California giant kelp forest food webs. *Ecosystems* 7, 341–357.

Heim, I., Hammel, J.U., Nickel, M. & Bruemmer, F. (2007a) Salting sponges: a reliable
non-toxic and cost-effective method to preserve poriferans in the field for
subsequent DNA-work. In: *Porifera Research: Biodiversity, Innovation and*
Sustainability. pp. 373–377.

Heim, I. & Nickel, M. (2010) Description and molecular phylogeny of Tethya leysae sp.
nov. (Porifera, Demospongiae, Hadromerida) from the Canadian Northeast Pacific
with remarks on the use of microtomography in sponge taxonomy. *Zootaxa* 2422, 1–
21.

Heim, I., Nickel, M. & Bruemmer, F. (2007b) Phylogeny of the genus Tethya (Tethyidae,
Hadromerida, Porifera): molecular and morphological aspects. *Journal of the*
Marine Biological Association of the United Kingdom 87, 1615–1627.

Heim, I., Nickel, M., Picton, B.E. & Brümmer, F. (2007c) Description and molecular
phylogeny of Tethya hibernica sp. nov. (Porifera, Demospongiae) from Northern
Ireland with remarks on the European species of the genus Tethya. *Zootaxa* 1595, 1–
15.

Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q. & Vinh, L.S. (2018)
UFBoot2: Improving the ultrafast bootstrap approximation. *Molecular Biology and*
Evolution 35, 518–522.

Idan, T., Shefer, S., Feldstein, T., Huchon, D. & Ilan, M. (2018) Shedding light on an
East-Mediterranean mesophotic sponge ground community and the regional sponge
fauna. *Mediterranean Marine Science* 19, 84–106.

Kober, K.M. & Nichols, S.A. (2007) On the phylogenetic relationships of hadromerid
and poecilosclerid sponges. *Journal of the Marine Biological Association of the UK*
87, 1585–1598.

Kushner, D.J., Rassweiler, A., McLaughlin, J.P. & Lafferty, K.D. (2013) A multi-decade
time series of kelp forest community structure at the California Channel Islands.
Ecology 94, 2655.

de Laubenfels, M. (1932) The marine and fresh-water sponges of California. *Proceedings*
of the U.S. National Museum 81, 1–140.

Lavrov, D. V., Forget, L., Kelly, M. & Lang, B.F. (2005) Mitochondrial genomes of two

demosponges provide insights into an early stage of animal evolution. *Molecular*
Biology and Evolution 22, 1231–1239.

Lee, W.L., Elvin, D.W., Reiswig, H.M. (2007) The Sponges of California: A Guide and
Key to the Marine Sponges of California. Monterey Bay Sanctuary Foundation, 130
pp

Letunic, I. & Bork, P. (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Research* 47, W256–W259.

Miller, R.J., Lafferty, K.D., Lamy, T., Kui, L., Rassweiler, A. & Reed, D.C. (2018) Giant
kelp, Macrocystis pyrifera, increases faunal diversity through physical engineering.
Proceedings of the Royal Society B 285, 20172571.

Morrow, C.C., Picton, B.E., Erpenbeck, D., Boury-Esnault, N., Maggs, C.A. & Allcock,
A.L. (2012) Congruence between nuclear and mitochondrial genes in
Demospongiae: A new hypothesis for relationships within the G4 clade (Porifera:
Demospongiae). *Molecular Phylogenetics and Evolution* 62, 174–190.

Nguyen, L.-T., Schmidt, H.A., von Haeseler, A. & Minh, B.Q. (2015) IQ-TREE: A fast
and effective stochastic algorithm for estimating maximum likelihood phylogenies.
Molecular Biology and Evolution 32, 268–274.

Nichols, S.A. (2005) An evaluation of support for order-level monophyly and
interrelationships within the class Demospongiae using partial data from the large
subunit rDNA and cytochrome oxidase subunit I. *Molecular Phylogenetics and*
Evolution 34, 81–96.

Pallas, P.S. (1766) Elenchus zoophytorum sistens generum adumbrationes generaliores et
specierum cognitarum succintas descriptiones, cum selectis auctorum synonymis.
Fransiscum Varrentrapp, Hagae.

Redmond, N.E., Morrow, C.C., Thacker, R.W., Diaz, M.C., Boury-Esnault, N., Cárdenas,
P., Hajdu, E., Lobo-Hajdu, G., Picton, B.E., Pomponi, S.A., Kayal, E. & Collins,
A.G. (2013) Phylogeny and systematics of Demospongiae in light of new small-
subunit ribosomal DNA (18S) sequences. *Integrative and Comparative Biology* 53,
388–415.

Regueiras, A., Alex, A., Costa, M.S., Pereira, S. & Vasconcelos, V. (2019) Diversity of
intertidal marine sponges from the western coast of Portugal (North-east Atlantic).
Journal of the Marine Biological Association of the United Kingdom 99, 1253–1265.

Riesgo, A., Novo, M., Sharma, P.P., Peterson, M., Maldonado, M. & Giribet, G. (2013)
Inferring the ancestral sexuality and reproductive condition in sponges (Porifera).
Zoologica Scripta 43, 101–117.

Rot, C., Goldfarb, I., Ilan, M. & Huchon, D. (2006) Putative cross-kingdom horizontal
gene transfer in sponge (Porifera) mitochondria. *BMC Evolutionary Biology* 6, 71.

Sarà, M. & Corriero, G. (1993) Redescription of Tethya californiana de Laubenfels as a
valid species for Tethya aurantia var. Californiana (Porifera, Demospongiae).
Ophelia 37, 203–211.

Sarà, M., Gomez, P. & Sará, A. (2001) East Pacific Mexican Tethya (Porifera :
Deniospongiae) with descriptions of five new species. *Proceedings of the Biological*
Society of Washington 114, 794–821.

Sarà, M. (2002) Family Tethyidae Gray, 1848. In: Hooper J.N.A. & van Soest R.W.M.
(eds) Systema Porifera: A Guide to the Classification of Sponges: 245-265. Kluwer
Academic/Plenum Publishers, New York. 928

Schneider, C.A., Rasband, W.S. & Eliceiri, K.W. (2012) NIH Image to ImageJ: 25 years
of image analysis. *Nature Methods* 9, 671–675.

Shaffer, M.R., Davy, S.K. & Bell, J.J. (2018) Hidden diversity in the genus Tethya:
comparing molecular and morphological techniques for species identification.
Heredity 122, 354–369.

Sorokin, S.J., Ekins, M.G., Yang, Q. & Cárdenas, P. (2019) A new deep-water Tethya
(Porifera, Tethyida, Tethyidae) from the Great Australian Bight and an updated
Tethyida phylogeny. *European Journal of Taxonomy* 529, 1–26.

Thacker, R.W., Hill, A.L., Hill, M.S., Redmond, N.E., Collins, A.G., Morrow, C.C.,
Spicer, L., Carmack, C.A., Zappe, M.E., Pohlmann, D., Hall, C., Diaz, M.C. &
Bangalore, P. V. (2013) Nearly complete 28S rRNA gene sequences confirm new
hypotheses of sponge evolution. *Integrative and Comparative Biology* 53, 373–387.

Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A. & Minh, B.Q. (2016) W-IQ-TREE: a
fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids*
Research 44, W232–W235.

Zhang, Y., Huang, D., Wang, D. & Ding, S. (2016) The complete mitochondrial genome
of sponge Tethya sp. (Demospongiae, Tethyida, Tethyidae). *Mitochondrial DNA*,
472–474.

853
854

855   **Figures**

857   **Figure 1.** Subtidal collection locations. Sites where *Tethya* were found (black) and not

858   found (white) are shown. The two sites away from the coastline are oil platforms. The

859   diamond indicates the type locality for *T. vacua*.

861
862
863

864   **Figure 2.** Gene tree at the large ribosomal subunit (28S). Bootstrap values are shown for
865   nodes with > 80% support; nodes with < 50% support are collapsed. Green = F.
866   Hemiasterellidae, blue = *Tethytimea*, orange = *Tethya*. Genbank accession numbers are
867   shown; those beginning with MT, shown in bold, are new. Scale bar indicates
868   substitutions per site.
869

875 **Figure 3.** Gene tree at cox1. Bootstrap values are shown for nodes with > 80% support;
876 nodes with < 50% support are collapsed. Green = F. Hemiasterellidae, orange = *Tethya*.
877 Genbank accession numbers are shown; those beginning with MT, shown in bold, are
878 new. Scale bar indicates substitutions per site.
879

880
881
882

883 **Figure 4.** Gene tree at the small ribosomal subunit (18S). Bootstrap values are shown for
884 nodes with > 80% support; nodes with < 50% support are collapsed. Orange = *Tethya*.
885 Genbank accession numbers are shown; those beginning with MT, shown in bold, are
886 new. Scale bar indicates substitutions per site.
887

888
889
890

891 **Figure 5.** *Tethyatimea authia.* A) CASIZ 235110, Carpinteria Reef; B) TLT 364,
892 Saddleback Ridge, Santa Cruz Island; C) CASIZ 235111, Perkin's Reef, Monterey Bay;
893 D) CASIZ 235111 after collection but before preservation, showing constriction of
894 gutters. Scale bar = 1 cm.
895

896
897
898

899 **Figure 6.** Microscopic features of *T. authia.* A) tylostyles, styles from voucher CASIZ
900 235111, scale bar 100 μm; B) euasters from voucher CASIZ 235111, scale bar 20 μm; C)
901 perpendicular tissue section from voucher TLT 364, scale bar = 500 μm.
902

903
904
905 **Figure 7.** Distribution of aster diameter for *T. authia* samples. From top to bottom:
906 CASIZ 235110, Carpinteria Reef; CASIZ 235111, Perkin's Reef, Monterey Bay; TLT
907 364, Saddleback Ridge, Santa Cruz Island.
908

909
910

911   **Figure 8.** *Galaxia gaviotensis*. Top: sponge in situ before collection. Perpendicular
912   sections (center, bottom) show skeletal morphology of asters and megascleres.
913

914
915

916 **Figure 9.** Morphology of *G. gaviotensis* spicules. A) Tylostyles and styles; B) oxea; C)
917 euasters. Numbers below scale bars indicate size in µm.
918

919
920
921

922   Table 1. Spicule data for *Tethya californiana* in microns.

	Hopkin's Marine Station, Monterey	Santa Rosa Island	Santa Rosa Island (T. vacua paratype)	Naples Reef, Santa Barbara	Ensenada, Baja California Norte	Cabo San Miguel, Sea of Cortez
Survey ID	TLT 354	TLT 361	n/a	TLT 119	n/a	n/a
Museum Voucher	CASIZ 235109	n/a	CASIZ 184729	CASIZ 235108	CNPGG 1199	CNPGG 0012
Collection date	9/16/19	10/9/19	12/24/81	7/31/19	4/8/11	3/5/85
All megascleres:						
Sample size	55	55	78	81	46	88
Length	449-(1405)-2590	505-(1197)-2119	492-(1408)-2073	498-(1277)-2214	404-(1334)-2280	275-(1231)-2141
Max. width	4-(22)-47	7-(15)-29	3-(20)-33	4-(17)-37	7-(24)-44	4-(19)-37
Head/max width	1.1-(1.6)-2.3	1.0-(1.5)-2.3	1.0-(1.8)-2.7	1.0-(1.9)-3.0	1.1-(1.9)-3.4	1.1-(2.1)-4.3
Proportion anisostrongyles	0.34	0.27	0.45	0.31	0.41	0.02
Anisostrongyles						
Sample size	27	15	35	25	19	2
Length	1224-(1992)-2590	971-(1418)-1890	1156-(1603)-1977	1057-(1772)-2195	520-(1573)-2260	2139-(2140)-2141
Max. width	19-(33)-47	14-(20)-28	14-(24)-33	13-(27)-37	9-(31)-42	28-(33)-37
Head/max width	1.1-(1.7)-2.1	1.2-(1.5)-1.7	1.3-(1.8)-2.6	1.3-(2.1)-2.9	1.2-(2.0)-3.4	2.0-(2.4)-2.8
Strongyloxeas/styles						
Sample size	28	40	43	56	27	86
Length	449-(839)-1597	505-(1115)-2119	492-(1249)-2073	498-(1057)-2214	404-(1166)-2280	275-(1210)-2072
Max. width	4-(11)-29	7-(13)-29	3-(16)-33	4-(13)-33	7-(20)-44	4-(19)-37
Head/max width	1.1-(1.6)-2.3	1.0-(1.5)-2.3	1.0-(1.8)-2.7	1.0-(1.8)-3.0	1.1-(1.8)-2.6	1.1-(2.1)-4.3
Megasters:						
Sample size	41	40	48	41	49	41
Diameter	23-(43)-57	12-(22)-34	25-(37)-46	22-(37)-48	29-(45)-60	21-(43)-56
R/C	0.52-(0.79)-1.04	0.48-(0.70)-1.00	0.55-(0.85)-1.27	0.59-(0.88)-1.20	0.63-(0.92)-1.54	0.47-(0.79)-1.00
Tukey HSD of R/C	A	B	A, C	A, C	C	A

924
925

926   Table 2. Spicule data for *Tethytimea authia* in microns.

	Carpinteria Reef, Santa Barbara CA	Perkin's Reef, Monterey CA	Saddleback Ridge, Santa Cruz Island, CA	Timea authia (Carballo & Cruz-Barraza 2006)
Survey ID	TLT 125	TLT 449	TLT 364	-
Museum Voucher	CASIZ 235110	CASIZ 235111	-	-
Collection date	7/31/19	11/24/19	10/18/19	-
All styles:				
Sample size	41	28	35	-
Length	110-(491)-779	309-(560)-812	176-(469)-883	-
Width	2-(5.3)-10	5-(7.1)-10	3-(5.3)-10	-
Tylostyles only:				
Sample size	29	11	19	-
Length	208-(543)-779	379-(653)-797	176-(475)-748	208-(478)-915
Width	2-(5.3)-9	5-(7.8)-10	3-(5.4)-9	4-(6.4)-15
Simple styles only:				
Sample size	11	12	16	-
Length	110-(372)-555	309-(497)-812	176-(461)-883	187-(289)-800
Width	2-(5.5)-10	5-(6.8)-9	3-(5.3)-10	3-(3.5)-5
Asters:				
Sample size	219	369	204	-
Diameter	6-(14.8)-25	7-(17.7)-26	3-(15.5)-28	3-(14.7)-24

928
929
930

931   Supplementary table 1. Sampling effort and locations. Intertidal/marina sites are listed
932   last. For all subtidal sites, the number of sampling dives is listed, and the cumulative dive
933   time as an estimate of search effort.
934

Region	Location name	Lat	Long	Dives	Max depth (m)	Cumulative search time (min)	Tethya californiana	Tethytimea authia	Galexia gaviotae
Monterey	Middle Reef, Pt. Lobos	36.52172	-121.93894	1	15	60			
Monterey	Hopkin's Marine Station	36.62084	-121.90142	1	11	74	Collected		
Monterey	Perkin's Reef	36.62920	-121.92031	1	12	46	Photo	Collected	
Santa Barbara	Arroyo Hondo Reef	34.47182	-120.14262	2	8	52			
Santa Barbara	Arroyo Quemado Reef	34.46775	-120.11905	7	11	302	Collected		Collected
Santa Barbara	Tajigus	34.46279	-120.10185	2	8	93			
Santa Barbara	Naples Reef	34.42212	-119.95154	6	16	279	Collected		
Santa Barbara	Elwood Bluffs	34.41775	-119.90150	3	14	138	Collected		
Santa Barbara	Coal Oil Point	34.40450	-119.87890	5	11	257	Collected		
Santa Barbara	Isla Vista Reef	34.40278	-119.85755	4	12	135			
Santa Barbara	Goleta Sewer Pipe	34.41420	-119.82930	1	6	22			
Santa Barbara	Mohawk Reef	34.39407	-119.72957	7	9	220	Collected		
Santa Barbara	1000 Steps	34.39472	-119.71347	1	7	30			
Santa Barbara	Leadbetter Reef	34.39895	-119.69703	1	9	14			
Santa Barbara	Oil Platform C	34.33293	-119.63173	1	23	39			
Santa Barbara	Oil Platform A	34.33189	-119.61353	1	27	35			
Santa Barbara	Carpinteria Reef	34.39163	-119.54169	2	6	61		Collected	
Santa Barbara	Elwood Junkpile	34.42687	-119.92390	1	12	34			
Santa Barbara	UCSB intake pipe	34.40297	-119.84080	1	18	34			
Los Angeles	Hawthorne reef	33.74714	-118.42090	1	24	39	Photo		
Los Angeles	Halfway reef	33.76265	-118.42560	1	23	40			
Los Angeles	Resort Wall	33.76499	-118.42815	1	21	43			
Los Angeles	Flat rock	33.79662	-118.41110	1	8	45			
Los Angeles	Redondo barge	33.83833	-118.41040	2	27	49			
San Diego	Goalpost	32.69438	-117.26860	1	15	50	Photo		
San Diego	Six Fathoms	32.71000	-117.26860	1	18	40	Photo	Photo	
San Miguel Island	Wreck of the Cuba	34.03009	-120.45575	1	10	52	Photo		
San Miguel Island	Inside of Wycoff Ledge	34.02132	-120.38710	1	19	52	Photo		
San Miguel Island	Miracle Mile, E of Crook Point	34.01367	-120.34180	2	20	108	Photo		
Santa Rosa Island	Johnson's Lee	33.89966	-120.10735	2	15	65	Collected		
Santa Cruz Island	Hazzard's, Santa Cruz Is.	34.06066	-119.82810	1	13	75			
Santa Cruz Island	Cueva Valdez, Santa Cruz Is.	34.05498	-119.81960	1	14	75			
Santa Cruz Island	Platt's Harbor, Santa Cruz Is.	34.04765	-119.73537	1	10	86			
Santa Cruz Island	Marmeda, Santa Cruz Is.	33.98378	-119.6391	1	13	35	Collected		
Santa Cruz Island	E. of Marmeda, Santa Cruz Is.	33.98380	-119.61290	1	15	38			
Santa Cruz Island	The Suburbs, Santa Cruz Isl.	34.05291	-119.58309	1	14	59			
Santa Cruz Island	Big Rock, Santa Cruz Is.	34.05220	-119.57360	5	14	299	Photo		
Santa Cruz Island	Pink Ribbon	33.98967	-119.52470	1	11	60			
Santa Cruz Island	Saddleback Ridge	34.03817	-119.52470	2	12	134		Collected	
Anacapa Island	West End, Anacapa Is.	34.01352	-119.44570	1	12	40	Collected		
Anacapa Island	Zebra Cove	34.01000	-119.44000	1	13	66	Photo		
Anacapa Island	Landslide Cove, Anacapa Is.	34.01539	-119.43560	1	11	49			
Anacapa Island	Underwater Island, Anacapa Is.	34.00720	-119.42959	2	18	102		Photo	
Anacapa Island	Cat Rock	34.00426	-119.42335	1	11	65			
Anacapa Island	Portuguese Rock, Anacapa Is.	34.01545	-119.42149	2	26	111			
Anacapa Island	Underwater Arch, Anacapa Is.	34.01399	-119.36224	1	10	45			
Anacapa Island	Landing Cove, Anacapa Is.	34.01690	-119.36090	1	15	57			
Catalina Island	Parson's Landing	33.47502	-118.55000	2	17	120	Photo		
Catalina Island	Eel Cove	33.45662	-118.51120	1	9	30			
Monterey	Point Pinos Intertidal	36.63447	-121.93923	n/a	n/a	90			
Monterey	Monterey Marina	36.60797	-121.89288	n/a	n/a	20			
Santa Barbara	COP Intertidal	34.406656	-119.8775	n/a	n/a	60			
Santa Barbara	Santa Barbara Harbor	34.405592	-119.68964	n/a	n/a	150			
Los Angeles	22nd Street Marina	33.724772	-118.28078	n/a	n/a	20			
San Diego	Point La Jolla	32.85112	-117.27298	n/a	n/a	60			
San Diego	Mission Bay Harbor	32.764209	-117.23829	n/a	n/a	40			

935
936
937

938 Supplementary table 2. Archival information for vouchered samples and samples with
939 sequence data. CASIZ: California Academy of Sciences Invertebrate Zoology.
940

Region	Location name	Lat	Long	Date	Species	Collection ID	Mueseum voucher	CO1 sequence	28S sequence	18S sequence	iNaturalist record
Santa Barbara	Arroyo Quemado Reef	34.46775	-120.11905	7/29/19	Galexia gaviotae	TLT228	CASIZ 235112	MT139597	MT137926	MT137929	inaturalist.org/observations/41378835
Santa Rosa Island	East End	34.00000	-120.00170	12/24/81	Tethya "vacua"	-	CASIZ 184729	MT139602	-	MT137930	-
Monterey	Hopkin's Marine Station	36.62084	-121.90142	9/16/19	Tethya californiana	TLT354	CASIZ 235109	MT139598	-	MT137931	inaturalist.org/observations/32893483
Santa Barbara	Naples Reef	34.42212	-119.95154	7/31/19	Tethya californiana	TLT119	CASIZ 235108	MT139599	-	MT137932	inaturalist.org/observations/30057899
Santa Barbara	Elwood Bluffs	34.41775	-119.90150	4/17/19	Tethya californiana	TLT18	CASIZ 235106	MT139601	-	-	inaturalist.org/observations/22664350
Santa Cruz Island	Marmeda, Santa Cruz Is.	33.98378	-119.63910	4/26/19	Tethya californiana	TLT50	CASIZ 235107	MT139600	-	-	-
Santa Rosa Island	Johnson's Lee	33.89966	-120.10735	10/9/19	Tethya californiana	TLT361	-	-	MT137925	-	inaturalist.org/observations/34168264
Monterey	Perkin's Reef	36.62920	-121.92031	11/24/19	Timea authia	TLT449	CASIZ 235111	-	MT137923	MT137927	inaturalist.org/observations/36383292
Santa Barbara	Carpinteria Reef	34.39163	-119.54169	7/31/19	Timea authia	TLT125	CASIZ 235110	-	MT137924	MT137928	inaturalist.org/observations/31129708

941
942
943 Supplementary table 3. Raw data on spicule dimensions for all samples.
944
945 (This data set is not easily formatted to fit into a text document but is available upon
946 request, and will be submitted to the journal upon acceptance of the manuscript.)
947
948 Supplementary table 4. Genetic metadata.
949
950 (This data set is not easily formatted to fit into a text document but is available upon
951 request, and will be submitted to the journal upon acceptance of the manuscript.)
952