POMK Regulates Dystroglycan Function via LARGE-mediated Elongation of Matriglycan

Ameya S. Walimbe, M.S.E.[1], Tiandi Yang, Ph.D.[1], Takahiro Yonekawa, M.D. Ph.D.[1], David Venzke, B.S.[1], Mary E. Anderson, B.S.[1], Silvia Torelli, Ph.D.[2], Adnan Manzur, M.D.[2], Liping Yu Ph.D.[3], Megan Devereaux, B.S.[1], Marco Cuellar, B.S.[1], Sally Prouty B.S.[1], Saul O'Campo Landa B.S.[1], Francesco Muntoni, M.D.[2, 4], and Kevin P. Campbell, Ph.D.[1*]

[1]Howard Hughes Medical Institute, Senator Paul D. Wellstone Muscular Dystrophy Specialized Research Center, Department of Molecular Physiology and Biophysics and Department of Neurology, Roy J. and Lucille A. Carver College of Medicine, The University of Iowa, Iowa City, Iowa 52242, USA

[2]Dubowitz Neuromuscular Centre, UCL Great Ormond Street Institute of Child Health & Great Ormond Street Hospital, London, United Kingdom.

[3]Medical Nuclear Magnetic Resonance Facility, University of Iowa Roy J. and Lucille A. Carver College of Medicine, B291 Carver Biomedical Research Building, 285 Newton Road, Iowa City, IA 52242, USA

[4]National Institute for Health Research Great Ormond Street Hospital Biomedical Research Centre, UCL Great Ormond Street Institute of Child Health, London, United Kingdom.

* Corresponding author

ABSTRACT:

Protein O-Mannose Kinase (POMK) is a novel muscular dystrophy gene that phosphorylates mannose of core M3 (GalNac-β1,3-GlcNac-β1,4-Man) on α-dystroglycan (α-DG). Like-acetylglucosaminyltransferase-1 (LARGE) synthesizes matriglycan, a heteropolysaccharide [-GlcA-β1,3-Xyl-α1,3-]$_n$, that enables α-DG to function as an extracellular matrix (ECM) receptor. Extension of matriglycan enhances its binding capacity for ECM ligands and is required for skeletal muscle function. However, the mechanisms which regulate LARGE-mediated extension of matriglycan during muscle differentiation are unknown. Here we show that LARGE initiates matriglycan synthesis on α-DG without POMK but requires POMK to extend matriglycan to its mature length. Furthermore, mouse and human skeletal muscle lacking POMK express a short, non-extended form of matriglycan and exhibit muscular dystrophy. Our results demonstrate a POMK-dependent mechanism that regulates matriglycan synthesis and α-DG receptor function.

ONE SENTENCE SUMMARY:

POMK Mediates Matriglycan Extension

MAIN TEXT:

POMK is a glycosylation-specific kinase *(1-3)* that phosphorylates an *O*-mannosyl

trisaccharide (GalNac-β1,3-GlcNac-β1,4-Man), termed core M3, on the high-affinity ECM

receptor α-DG (Fig. 1A). The core M3 glycan structure terminates with matriglycan, a

heteropolysaccharide [-GlcA-β1,3-Xyl-α1,3-]$_n$ that is synthesized by LARGE *(3-5)* (Fig. 1A). A

single glucuronic acid-xylose (GlcA-Xyl) disaccharide repeat binds with high affinity to ECM

ligands bearing laminin G (LG) domains *(4,8)*, and there is a direct correlation between the

number of GlcA-Xyl repeats on α-DG and its binding capacity for ECM ligands *(10).* During

skeletal muscle differentiation, LARGE elongates matriglycan, thereby enhancing the binding

capacity of matriglycan for ECM ligands *(3, 4, 10).* Defects in matriglycan synthesis lead to

dystroglycanopathies, congenital and limb-girdle muscular dystrophies that can be accompanied

by structural brain abnormalities *(7-15).* Indeed, the clinical severity of dystroglycanopathy

correlates with the degree of non-extension of matriglycan *(10).* Despite the importance of

mature, full-length matriglycan for proper muscle structure and function, little is known about

how matriglycan elongation is regulated.

To understand the mechanisms regulating matriglycan elongation, we first studied

skeletal muscle from a patient (NH13-284) with a POMK mutation and congenital muscular

dystrophy (CMD) accompanied by structural brain malformations. Whole exome sequencing of

genomic DNA demonstrated a homozygous missense mutation causing Aspartate to Asparagine

substitution at residue 204 (D204N) of POMK (Fig. 1B). D204 is located in the catalytic loop of

POMK and serves as the catalytic base in the phosphorylation reaction catalyzed by the kinase

(Fig. 1B, S1). D204 is highly conserved (Fig. 1B), and mutation of D204 to Asparagine (D204N)

or Alanine (D204A) is predicted to eliminate POMK activity (Fig. S1) *(1)*. POMK activity from

skeletal muscle and skin fibroblasts of patient NH13-284 (POMK D204N) was undetectable

when compared to control muscle and fibroblasts, respectively (Fig. 1C). B4GAT1 and LARGE

activity was present in skeletal muscle and fibroblasts, respectively, of a control patient and

patient NH13-284 (Fig. S2A, S3). Immunofluorescence analyses of POMK D204N muscle

demonstrated partial IIH6 immunoreactivity, while the transmembrane subunit of DG, β-DG,

was expressed normally in both POMK D204N and control muscle (Fig. 1D). Flow cytometry

using IIH6 also demonstrated partial immunoreactivity in POMK D204N fibroblasts (Fig. S2B).

To test the effect of the POMK mutation on ligand binding using a more sensitive method than

IIH6 immunofluorescence, we performed a laminin overlay using laminin-111 on glycoproteins

enriched from skeletal muscle cryosections with Wheat Germ Agglutinin (WGA)-agarose. We

were surprised to find laminin binding (at ~90 to 100 kDa range) by a glycoprotein in POMK

D204N skeletal muscle (Fig. 1E).

To understand the biochemical basis of the laminin binding at ~90 to 100 kDa, we

knocked out POMK using LoxP sites and Cre driven by the Muscle Creatine Kinase (MCK)

promoter, which is active in murine skeletal muscle and heart, or both the MCK promoter and

the Paired Box 7 (Pax7) promoter, which is expressed by satellite cells and embryonic muscle

(Fig. S4) (*16, 17*). Quadriceps muscle extracts of MCK-Cre; POMK$^{LoxP/LoxP}$ mice show reduced

POMK activity compared to POMK$^{LoxP/LoxP}$ muscle but have similar levels of LARGE activity

(Fig. 2B, S5). Histologic analyses of MCK-Cre; Pax7-Cre; POMK$^{LoxP/LoxP}$ (M-POMK KO)

quadriceps muscles reveal hallmarks of a mild muscular dystrophy (Fig. 2A). M-POMK KO

mice also showed reductions in 2-limb grip strength and body weight and elevations in post-

exercise creatine kinase (CK) levels compared to littermate control POMK$^{LoxP/LoxP}$ mice (Fig.

S6). Immunofluorescence analysis of M-POMK KO muscle shows that β-DG is expressed at the

skeletal-muscle sarcolemma (Fig. 2A, 2C); however, like patient NH13-284 IIH6 immunoreactivity persisted in M-POMK KO muscle, but at a reduced intensity. Biochemical analysis of control and M-POMK KO muscle showed a typical, lower molecular weight (MW) α-DG with anti-core DG antibody (Fig. 2C) like that has been previously reported for dystroglycanopathy patients (7,10); however, on laminin overlay, we observed laminin binding at 90-100 kDa (Fig 2D), similar to POMK D204N skeletal muscle (Fig. 1E). Solid-phase binding analyses of M-POMK KO and MCK-Cre; POMK$^{LoxP/LoxP}$ skeletal muscle demonstrated a reduced binding capacity (relative B_{max}) for laminin-111 compared to control muscle (Fig. S8A), but higher than that of $LARGE^{Myd}$ muscle, which lacks matriglycan due to a deletion in the $LARGE$ gene.

To determine if a GlcA-Xyl disaccharide polymer, and, therefore, a form of matriglycan, is responsible for the laminin binding at 90-100 kDa in POMK-null muscle, we treated glycoproteins enriched from skeletal muscles of M-POMK KO and MCK-Cre; POMK$^{LoxP/LoxP}$ mice with two exoglycosidases, α-Xylosidase and β-Glucuronidase, which in combination digest matriglycan (*8*). Laminin overlay and solid phase analysis showed a reduction in laminin binding from muscles of M-POMK KO and MCK-Cre; POMK$^{LoxP/LoxP}$ mice (Fig. 2E, 2F, S8B) after dual exoglycosidase treatment. This demonstrates that matriglycan is responsible for the laminin binding at 90-100 kDa in POMK-deficient muscle.

We used a *POMK* KO HAP1 cell line to determine the identity of the above laminin-binding glycoprotein. The *POMK* KO HAP1 cells had undetectable levels of POMK activity and expression (Fig. 3A, S10) (*1*). A mass spectrometry (MS)-based glycomic analysis of O-glycans carried by recombinantly-expressed DG mucin-like domain protein further indicates the absence of an MS peak at *m/z* 873.5 corresponding to phosphorylated core M3 O-glycan (Fig. 3D, 3E),

consistent with an undetectable level of POMK activity in *POMK* KO HAP1 cells. Compared to WT HAP1 cells, immunoblots of *POMK* KO HAP1 cells showed a reduction in IIH6 immunoreactivity, a decrease in MW of core α-DG, and the presence of laminin binding at ~90 kDa on laminin overlay (Fig. 3C, S10). Laminin binding on overlay was rescued only after adenoviral transduction with wild-type (WT)-POMK, and not with POMK containing D204N (POMK D204N) or D204A (POMK D204A) mutations (Fig. 3C, S10). POMK D204N also lacked POMK activity *in vitro* but showed normal B4GAT1, B3GALNT2, and LARGE activity, thus confirming the pathogenicity of the D204N mutation (Fig. 3A, 3B, S11).

The bifunctional glycosyltransferase, LARGE, synthesizes matriglycan (*3-5*). To directly test if LARGE is required for synthesis of the 90 kDa laminin-binding glycoprotein in *POMK* KO HAP1 cells, we studied *POMK/LARGE* double knockout (DKO) HAP1 cells, which bear a CRISPR/Cas9-mediated deletion in *LARGE*. *POMK/LARGE* DKO HAP1 cells demonstrated the absence of the laminin binding at 90-100 kDa (Fig. 3F, S12), indicating that LARGE is required for the synthesis of the matriglycan responsible for laminin binding at 90-100 kDa. Moreover, *POMK/DG* DKO HAP1 cells demonstrated a complete absence of laminin binding (Fig. 3F) and IIH6 immunoreactivity at 90 kDa (Fig. S13), demonstrating that α-DG is the glycoprotein that binds laminin in the absence of POMK. We, therefore, refer to this glycoprotein as POMK-null α-DG (α-DG(POMK)). The length of matriglycan correlates with its binding capacity for ECM ligands (*4, 8, 10*). We hypothesized that, given the MW of α-DG(POMK) at 90-100 kDa, the glycan must be shorter than full-length matriglycan, and therefore, have a lower B_{max} for laminin. We measured the binding capacity of HAP1 α-DG using solid-phase binding assays. B_{max} of α-DG(POMK) for laminin-111 was reduced compared to wild-type α-DG (α-DG(WT)) but was greater than that of α-DG from *LARGE* KO HAP1 cells (Fig. 3G). *POMK/DG* DKO HAP1 cells

showed a reduction in B_{max} compared to *POMK* KO HAP1 cells, similar to the low levels

observed in *LARGE* KO HAP1 cells (Fig. 3G). These data indicate that a short, non-extended

form of matriglycan is synthesized on α-DG(POMK). This non-extended form of matriglycan

has a lower binding capacity for laminin-111 and, thus, a reduced level of α-DG receptor

function.

   We next carried out a series of experiments to characterize the enzymatic requirements

for synthesis of matriglycan on α-DG(POMK). After POMK phosphorylates core M3, Fukutin

(FKTN) modifies GalNac with ribitol-phosphate for synthesis of full-length matriglycan

(Fig.1A) *(4, 18)*. Overexpression of Isoprenoid Synthase Domain-Containing (ISPD), which

synthesizes the substrate (CDP-ribitol) of FKTN (Fig.1A), in *POMK* KO HAP1 cells increases

the matriglycan (without changing its MW) responsible for laminin binding at 90-100 kDa (Fig.

4A, S14) *(19-21)*. HAP1 cells lacking both POMK and ISPD do not express matriglycan, and

adenoviral transduction of these cells with *ISPD* restores the 90-100 kDa laminin binding, (Fig.

4B, S15). FKTN overexpression in *POMK* KO HAP1 cells also increased the 90-100 kDa

laminin binding (Fig. 4C, S16). These experiments collectively support a requirement for CDP-

ribitol for synthesis of the non-extended form of matriglycan. The DG N-terminal domain

(DGN) is required for interaction of α-DG with LARGE *(22-24)*. Synthesis of the non-extended

form of matriglycan also requires this interaction, as a DG mutant lacking the DGN (DGE)

expressed in *POMK/DG* DKO HAP1 cells did not show laminin binding at 90-100 kDa (Fig.

S17). Similar experiments also indicated that synthesis of the non-extended matriglycan in HAP1

cells requires Threonine 317 of the mucin-like domain of α-DG (Fig. S17). Thus, these

experiments show that the same pathway is used for the synthesis of the non-extended form of

matriglycan.

Overexpression of LARGE has been shown to produce very high molecular weight matriglycan and rescues the defect in matriglycan synthesis in distinct forms of CMD and in *LARGE* KO HAP1 cells (Fig. 4D) *(25)*; however, overexpression of LARGE in *POMK* or *POMK/LARGE* KO HAP1 cells did not lead to synthesis of very high molecular weight forms of matriglycan (Fig. 4E, F, S18A). Only the rescue of *POMK/LARGE* KO HAP1 cells with POMK enabled LARGE to synthesize high molecular weight matriglycan (Fig. 4F, S18B). Thus, our results demonstrate that LARGE requires POMK in order to extend matriglycan on α-DG for the synthesis of mature matriglycan and very high molecular weight matriglycan.

Our study demonstrates that POMK is required for the LARGE-mediated synthesis of mature, full length forms of matriglycan (Fig 4G). In the absence of POMK, LARGE generates a short, non-extended form of matriglycan (Fig 4F), as observed by the laminin binding at 90-100 kDa. Mice and humans lacking POMK in skeletal muscle express this short, non-extended form of matriglycan, which has a reduced binding capacity of ECM ligands, however, both humans and mice develop muscular dystrophy due to a reduction in normal α-DG-ligand interactions. Studies in HAP1 cells indicated that the short, non-extended form of matriglycan is generated because LARGE can initiate matriglycan synthesis but is unable to elongate the polysaccharide due to absence of phosphorylation of core M3. Further studies demonstrated that the short, non-extended form of matriglycan requires DGN, CDP-ribitol and Threonine 317 of the mucin-like domain of α-DG. Our study, therefore, shows that LARGE requires POMK activity in order to extend matriglycan to its mature, full-length form and thereby, enable normal α-DG function. Collectively, these results indicate that phosphorylation of core M3 by POMK is a regulatory step in the synthesis of matriglycan, thus enabling full ECM binding capacity and preventing muscular dystrophy.

REFERENCES AND NOTES:

1. Zhu Q, et al. (2016). *Elife*, 5:e22238.

2. Yoshida-Moriguchi T, et al. (2013). *Science,* 341(6148):896-899.

3. Yoshida-Moriguchi T, Campbell KP. (2015). *Glycobiology*, 25(7):702-13.

4. Hohenester E. *Current Opinion in Structural Biology*, Vol: 56, Pages: 56-63 (2019).

5. Inamori K, et al. (2012). *Science,* 335(6064), 93-96.

6. Yoshida-Moriguchi T, et. al. (2010). *Science*, 327(5961):88-92.

7. Michele, DE, et. al. (2002). *Nature*, 418(6896), 417-422.

8. Briggs DC, et. al. (2016). *Nat Chem Biol.* 12(10), 810-814.

9. Ohtsubo K and Marth JD. (2006). *Cell,* 126(5): 855-67.

10. Goddeeris MM, et. al. (2013). *Nature,* 503(7474),136-40.

11. Hohenester E, et al. (1999). *Mol Cell*. 4(5), 783-92.

12. Jae LT, et. al. (2013). *Science*, 340(6131), 479-483.

13. Di Costanzo S, et al. (2014). *Hum Mol Genet.* 23(21), 5781-5792.

14. von Renesse A, et al. (2014). *J Med Genet*. 51(4), 275-82.

15. Ardicli D, et. al. (2017). *Neuromuscul Disord.* 27(3), 239-242.

16. Brüning JC, et al. (1998). *Mol Cell*. Nov;2(5):559-69.

17. Keller C, et al. (2004). *Genes Dev*. 18(21):2608-13.

18. Kanagawa, et al. (2016). *Cell Reports*, 14(9):2209-2223.

19. Willer T, et. al. (2012). *Nat Genet.* 44(5), 575-580.

20. Gerin I, et al. (2016). *Nat Comm*. 7:11534.

21. Riemersma, et al. (2015). *Chem Biol.* 22(12), 1643-52.

22. Hara Y, et. al. (2011). *Proc Natl Acad Sci U S A*, 108(42), 17426-17431.

23. Kanagawa M, et. al. (2004). *Cell*, 117(7), 953-64.

24. Hara Y, et al. (2011). *NEJM*, 364(10):939-46.

25. Barresi R, et. al. (2004). *Nat Med.* 10(7), 696-703.

SUPPLEMENTAL REFERENCES:

26. Pinkert CA. (2002). *Academic Press*, ed. 2, pp. 20-31.

27. Ervasti, J. M. & Campbell, K. P. (1991). *Cell*, 66, 1121-1131.

28. Kunz S, et al. (2001). *J. Cell Biol.* 155, 301-310.

29. Anderson RD, et al. (2000). *Gene Ther.* 7, 1034-1038.

30. Willer, T. et al. (2014). *Elife,* 3:e03941.

31. Salleh, HM, et al. (2006). *Carbohydr. Res.* 341, 49-59.

32. Moracci, M. et al. (2000). *J. Biol. Chem.* 275, 22082-22089.

33. Mayer M. & Meyer, B. (2001). *J. Am. Chem. Soc.* 123, 6108-6117.

34. Delaglio, F. et al. (1995). *J. Biomol. NMR.* 6, 277-293.

35. Johnson, B. A. & Blevins, R. A. (1994). *J. Biomol. NMR.* 4, 603-614.

36. Jang-Lee, J. et al. (2006). *Meth. Enzymol.* 415, 59–86.

37. Zhang, H. et al. (2014). *Nat. Commun.* 5, 4339.

ACKNOWLEDGMENTS

Funding

This work was supported in part by a Paul D. Wellstone Muscular Dystrophy Specialized Research Center grant (1U54NS053672 to KPC). KPC is an investigator of the Howard Hughes Medical Institute.

S. T and F.M. are supported by the National Institute for Health Research Biomedical Research Centre at Great Ormond Street Hospital for Children NHS Foundation Trust and University College London. The BRC/ MRC Centre for Neuromuscular Diseases Biobank and is acknowledged for providing patients' serum and biopsy samples. The Muscular Dystrophy UK support to the GOSH Neuromuscular Centre is also gratefully acknowledged.

A.S.W. is a student in the University of Iowa Medical Scientist Training Program, which is funded by Medical Scientist Training Program Grant by the National Institute of General Medical Sciences (NIGMS) 5 T32 GM007337.

Authors' Contributions

ASW and KPC designed experiments and wrote the manuscript. ASW, TY, TY, DV, MC, ST, MEA, LY, MD, SOL, and SJP performed experiments. AM, ST, and FM provided critical information and materials regarding the patient. FM was involved in meaningful discussions regarding the manuscript. All authors contributed to and gave feedback on the manuscript.

Competing interests

The authors declare that they have no competing interests.

Data and materials availability

All data generated or analyzed during this study are included in this published article.

Supplementary Materials

Materials and Methods

Supplementary Text

Figures S1–S24

A

B

POMK

C

Fibroblast POMK Activity

Skeletal Muscle POMK Activity

D

	H&E	β-DG	IIH6

E

Laminin Overlay

Fig. 1. Model of Matriglycan Synthesis and Characterization of a Patient with a POMK Mutation. A) Model depicting synthesis of the laminin-binding modification on α-DG and the enzymes involved. Core M3 (GalNac-β3-GlcNac-β4-Man) is synthesized by POMT1/2, POMGNT2, and B3GALNT2. POMK phosphorylates the C6 hydroxyl of mannose. The phosphorylated core M3 is elongated by FKTN, FKRP, TMEM5, and B4GAT1 in the Golgi. ISPD produces CDP-Ribitol in the cytosol, which is a substrate for FKTN and FKRP. LARGE synthesizes matriglycan, which is the ligand-binding moiety. **B)** Domain structure of POMK and location of D204N mutation. (above) Human POMK is composed of a signal peptide, transmembrane domain (TM), and a kinase domain (N-lobe and C-lobe). The kinase domain contains the catalytic loop (orange) and activation segment (green). The patient's missense mutation (D204N) is present within a conserved MCD motif. (below) Alignment of protein sequences flanking the D204N mutation. The mutation alters a highly conserved aspartate that serves as the catalytic base in the phosphorylation reaction catalyzed by POMK. **C)** POMK activity in control and patient NH13-284 (POMK D204N) fibroblasts (left) and skeletal muscle (right). For fibroblast POMK activity, n=3 experiments were performed. Triple asterisks indicate p-value<0.001. Due to the limited amount of skeletal muscle, n=1 experiment was performed. **D)** Histology and immunofluorescence staining of cryosections from control and POMK D204N skeletal muscle using IIH6, a monoclonal antibody against matriglycan, and an antibody against β-DG. (Scale bars: Control H&E- 200 µM, POMK D204N H&E 75 µM). **E)** Laminin overlay of control and POMK D204N skeletal muscle.

Fig. 2. Mice with a Muscle-Specific Loss of POMK Express Matriglycan. A)
Immunofluorescence and histologic analyses of MCK-Cre; Pax7-Cre; POMK[LoxP/LoxP] (M-POMK
KO) and POMK[LoxP/LoxP] (Control) quadriceps skeletal muscle. Sections from 4 to 6 week-old
mice were stained with antibodies against matriglycan (IIH6) and β-DG. Histology was
evaluated by hematoxylin and eosin (H&E) staining (three replicates). Scale bars: 100 µM. **B)**
POMK activity in extracts of MCK-Cre; POMK[LoxP/LoxP] and POMK[LoxP/LoxP] quadriceps skeletal
muscles. Triple asterisks indicate statistical significance using a Student's unpaired t-test with a
p-value<0.0001. **C)** Biochemical analysis of Control and M-POMK KO skeletal muscle.
Glycoproteins were enriched from quadriceps skeletal muscles of mice using wheat-germ
agglutinin (WGA)-agarose. Immunoblotting was performed with antibody AF6868, which
recognizes core α-DG and β-DG (three replicates). **D)** Laminin overlay of quadriceps muscles of
Control and M-POMK KO mice (three replicates). **E) and F)** Laminin overlay (E) and Solid
phase analysis (F) of skeletal muscles of M-POMK KO mice treated with two exoglycosidases,
α-xylosidase (Xylsa) and β-glucuronidase (Bgus) for 17 hours (three replicates).

Fig. 3. α-DG is Modified with Matriglycan in the Absence of POMK. A) POMK activity in wild-type (WT) HAP1 and *POMK* knockout (KO) HAP1 cells expressing POMK D204N, D204A, or POMK WT. Asterisks indicate P-value<0.001 compared to *POMK* KO alone using One-Way ANOVA with Dunnett's Test for Multiple Comparisons (three replicates, errors bars represent standard deviation). **B)** LARGE activity in *POMK* KO HAP1 cells with the indicated POMK mutants. **C)** Laminin overlay of *POMK* KO HAP1 cells expressing the indicated mutants was performed (three replicates). **D) and E)** Mass Spectrometry (MS)-based O-glycomic analyses of DG mucin-like domain protein (DG390TevHis) expressed in *Fukutin (FKTN)* KO HAP1 cells (D) or *POMK* KO HAP1 cells (E). The O-glycans were reductively released from the protein backbone and permethylated prior to Matrix-Assisted Laser Desorption/Ionization time-of-flight (MALDI-TOF) analyses. MS peaks at *m/z* 779.5 (779.6) correspond to a mixture of core 2 and core M3 O-glycan, and at 873.5, phosphorylated core M3 O-glycan (red). **F)** Laminin overlays of WT, *POMK* KO, and *POMK/LARGE* KO HAP1 cells (left) or WT, *POMK* KO and *POMK/DG* KO HAP1 cells (right). **G)** Solid phase analysis of WT, *POMK* KO, *POMK/DG* double KO, and *LARGE* KO HAP1 cells (three replicates).

Fig. 4. Characterization of the Synthesis of Non-Extended Matriglycan and Requirements for LARGE-Mediated Elongation. A), B), and C) Laminin overlays of *POMK* KO HAP1 cells overexpressing adenoviral ISPD (Ad-*ISPD*) (A), HAP1 cells lacking both POMK and ISPD overexpressing Ad-*ISPD* (B), and *POMK* KO HAP1 cells overexpressing adenoviral Fukutin (C). **D)** IIH6 immunoblotting of *LARGE* KO HAP1 cells expressing adenoviral LARGE (Ad-*LARGE*). **E) and F)** Laminin overlays of *POMK* KO HAP1 cells overexpressing Ad-*LARGE* (E), *POMK/LARGE* double KO HAP1 cells expressing Ad-*LARGE* or Ad-*LARGE* and Ad-*POMK* (F)*.* All experiments were performed in triplicate. **G) and H)** Model depicting matriglycan on α-DG with POMK (G) and without POMK (H). Mature matriglycan is a long polysaccharide; however, a shorter, non-extended form of matriglycan is present in the absence of POMK.