SARS-COV-2 RBD Oral Vaccine Boosted by Mucosal Immune Adjuvant LTB26 via DCs and B Cells Activation in Mice

Yongping Ma*[1], Qinlin Shi[#2], Qiujuan Wang[#1], Sijing Chen[1], Sijie Gan[1], Changyin Fang[1], Yanxi Shen[1], Min Jiang[1],Tao Lin[2], Fangzhou Song[1]

1. Department of Biochemistry & Molecular Biology, Molecular Medicine & Cancer Research Center, Chongqing Medical University, Chongqing 400016, People's Republic of China

2. Department of Urology, Children's Hospital of Chongqing Medical University, Chongqing 400014, People's Republic of China

Keywords: SARS-COV-2; mucosal adjuvant; LTB26; vaccine; immune;

*Correspondence:

Yongping MA

Department of Biochemistry and Molecular Biology, Chongqing Medical University

Yuzhong District, Yi XueYuan Road, No 1, Chongqing, 400016, China

Co-first author

Abstract

Although several SARS-COV-2 vaccines have been approved, no one oral live vaccine is available. Here, an oral SARS-COV-2 RBD live vaccine containing LTB26 adjuvant has been developed. BALB/c mice are oral vaccinated with attenuated *Salmonella typhimurium* SL7207 containing pcDNA3.1-LTB26RBD or pcDNA3.1-RBD plasmids. The result shows that the high level of RBD specific antibody is produced in pcDNA3.1-LTB26RBD treatment. The mechanism indicates that LTB26 enhances RBD antibody production by significantly upregulating the activity of MHC II$^+$ DCs and CD19$^+$CD45$^+$ B cells. LTB26 mutant is derived from heat-labile enterotoxin B subunit (LTB) wild type of *Escherichia coli* with enhanced immune adjuvanticity. Based on the pre-experiment result that SL7207 interferes the function of LTB26, the purified LTB26 was mixed with purified human rotavirus VP8 antigen to explore the mechanism of adjuvant. The results suggests that LTB26 enhances mucosal immune responses via increased of BCR and MHC II$^+$ expression. Furthermore, LTB26 promotes both Th1 and Th2 cell mediated immunity. Therefore,

LTB26 maybe a potent adjuvant for mucosal vaccine development in view of the safety of LTB26 than LT toxin.

Introduction

Since December 2019, the 2019 novel coronavirus (2019-nCoV, SARS-COV-2) has infected over 95.38 million cases worldwide with about 2.03 million deaths. Although about 120 vaccines are in different stage of clinical trials and some inactivated or mRNA vaccine candidates have been approved, little is oral vaccination. Therefore, safe and effective mucosal vaccines are still necessary to prevent infection from SARS-COV-2.(Poland, Ovsyannikova et al., 2020) SARS-COV-2 cell entry depends on ACE2-RBD interaction(Hoffmann, Kleine-Weber et al., 2020). Therefore, receptor binding domain (RBD) of SARS-COV-2 spike protein (S protein) is the first target for vaccine development. As a pathogen of mucosal infection, mucosal immunity is very important to prevent of SARS-COV-2.

LTB26 is derived from heat-labile enterotoxin B subunit (LTB) of *Escherichia coli* with enhanced mucosal immune adjuvanticity. From the perspective of clinical application, LTB and its derivatives are safe without toxicity than LT toxin(da Hora, Conceicao et al., 2011). LT belongs to the members of the AB5 bacterial toxin family and consists of five B subunit (LTB) with a single catalytically active A subunit (LTA)(Merritt, Zhang et al., 2002). LTA has toxic effects of ADP-ribosylating activity and LTB binds to the cell membrane with GM1 ganglioside (GM1) following delivery LTA into cells inducing infectious diarrhea(da Hora et al., 2011). LT has been reported as mucosal immune adjuvant in 1988(Clemens, Sack et al., 1988). However, it fails to meet the clinical requirement for its toxicity(da Hora et al., 2011, Ma, 2016). Nevertheless, LTB is also not applicable for weaker adjuvanticity comparing with LT. Therefore, the construction of non-toxic LT has been a research hotspot in recent years(Clements & Norton, 2018, Ghazali-Bina, Pourmand et al., 2019, Ma,

2016, Nashar, 2014, White, Haghighi et al., 2017). However, there is no report to develop adjuvanticity-enhanced LTB mutant.

In this study, we optimized an adjuvanticity-enhanced LTB26 mutant from four LTB mutants (Holmner, Mackenzie et al., 2011, Ma, 2016). By fusion expressing with LTB26 adjuvant, a promising live SARS-COV-2 LTB26RBD mucosal vaccine was constructed using attenuated *Salmonella typhimurium* SL7207 to boost systematic immune responses.

Results

LTB26-RBD recombinant was constructed successfully. After commercial synthesis of LTB26-RBD and RBD sequences, pcDNA3.1-LTB26RBD and pcDNA3.1-RBD plasmids were constructed, respectively. Then attenuated *Salmonella typhimurium* SL7207 (SL7207) was electrically transformed with pcDNA3.1-LTB26RBD, pcDNA3.1-RBD and pcDNA3.1 plasmids at 2.5kv and the restriction enzymes analysis confirmed the recombinants of LTB26-RBD and RBD vaccine candidates (Fig. 1A).

LTB26 boosted immune responses via upregulating MHC II+ DCs. In order to decipher the mechanism, dendritic cells (DCs) and B cells from vaccinated mice were purified and analyzed with FCM (flow cytometry). The results suggested that LTB26 activated MCH II$^+$ DCs than *Salmonella typhimurium* SL7207 (SL7207) or RBD alone. The ratio of LTB26RBD to RBD was 29.3 times, LTB26RBD to PBS was 53.1 times, and LTB26RBD to SL7207 was 23.4 times. However, The ratio of SL7207 to PBS was 22.7 times. Even though there were no significant MCH II$^+$ DCs differences between SL7207 and SL7207-RBD vaccination mice comparing with SL7207-LTB26RBD treatment. However, SL7207-RBD and SL7207 vaccinations were significantly upregulated DCs than that of PBS treatment (Fig. 1B). That meant, LTB26 adjuvant was necessary for *Salmonella*-based vaccine construction. Meanwhile, LTB26 significantly activated CD19$^+$CD45$^+$ B cells than PBS treatment and increased about 14.5% proportion (Fig. 1C). As a summary, LTB26 significantly

upregulated both DCs and B cells activity to enhance antigen presentation

and antibody production.

LTB26 enhanced mice to produce high-level IgG antibody.

After the 3[rd] boosting vaccination, BALB/c mice produced high-level RBD

specific antibody in SL7207-LTB26RBD immune group (Fig. 2, p<0.025).

The result indicated that LTB26 has promising adjuvant activity. However,

we did not continue the vaccination of SL7207-RBD single group for the

absent of LTB26 adjuvant according to the primary test (Fig. 1B, C).

The characteristic of LTB mutants. The pET32a-LTB, pET32a-

LTB26, pET32a-LTB34, pET32a-LTB57 and pET32a-LTB85 were

confirmed by sequencing and the products of the five proteins were

detected by SDS-PAGE. Then, the proteins were purified using

BeaverBeads™ His-tag protein purification kit (Beaverbio, Suzhou, China)

and were detected a band of 30 KD protein using SDS-PAGE (Fig.3A).

The protein concentration of LTB was 3.412 µg/µl, LTB26, 3.215 µg/µl,

LTB34, 3.325 µg/µl, LTB57, 3.011 µg/µl, and LTB85, 1.375 µg/µl, respectively.

Compared with LTB, LTB26 was mutated from I26T27E28L29C30 to M26S27N28 and from E32Y33H34 to T32I33N34, and meanwhile deleted L29C30 residues (Fig.3B). In turn, LTB34 was mutated from N35T36 Q37I38Y39T40I41N42D43K44 to L35S36L37S38N39S40T41I42N43 Y44 (Fig.1B), LTB57 was mutated from E57M58 to F57Y58 and from I60I61T62F63 to E60F61H62H63 (Fig.3C) and LTB85 was mutated from I86E87R88M89K90D91 to Y86V87E88F89H90H91, respectively (Fig.3C). As a summary, the mutations were located in epitope 1(EP1) of LTB (LTB26), middle EP1 and EP2 (LTB34), EP2,3,4 (LTB57) and EP5 (LTB85), respectively (Fig.3D)(Takahashi, Kiyono et al., 1996).

LTB26 significantly enhanced both systematic and mucosal immune response. Because *Salmonella* SL7207 interfered with the adjuvant activity of LTB26 as previously described (Figure 2A), we used the purified human rotavirus VP8 antigen (hRV VP8 or VP8) to study the

molecular mechanism of the adjuvant of purified LTB26 and other mutants.

The serum VP8 specific IgG levels of LTB+VP8, LTB26+VP8 and LTB34+VP8 were significantly higher than that of the VP8 alone group (Fig.4A, $p<0.01$). However, the serum IgG level of LTB34+VP8 vaccination was slightly higher than that of LTB+VP8 treatment, but lower than that of LTB26+VP8 vaccination (Fig.4A, $p<0.01$). On the contrary, the serum IgG levels of LTB57+VP8 and LTB85+VP8 vaccinations were lower than that of LTB+VP8 groups (Fig. 4A). Similarly, the variation trends of lung mucosal VP8 specific sIgA were consistent with that of serum IgG among the LTB+VP8 and the other LTB mutants vaccination on 21 d post-vaccination ($p<0.001$) (Fig. 4B). LTB26 also elicited the highest mucosal immune response compared with LTB and LTB34, respectively. Thus, LTB26 was the optimal adjuvant mutant of LTB to elicit robust systematic and mucosal immune responses in this study.

The GM1-binding affinity of LTB mutants no positive association with its adjuvant activity. Previous study suggested that LTB adjuvant

positively associated with its GM1-binding affinity(de Haan, Verweij et al., 1998, Holmner et al., 2011). Thus, the GM1-binding activity of the four LTB mutants was tested in turn. The GM1-binding affinity was decreased gradually from LTB57, LTB26, LTB34, and LTB85 to LTB (Fig. 4C, D). However, the adjuvanticity was decreased successively from LTB26, LTB34, LTB, and LTB57 to LTB85. Therefore, LTB26 obtained the highest adjuvanticity with a weaker GM1-binding affinity. Instead, LTB57 had the highest GM1-binding affinity with the lowest adjuvanticity (Fig. 4C,D).

The transcriptome analysis indicates LTB26 enhanced B cells via BCR pathway and MHC II[+] DCs. According to GM1-binding analysis and immune data, purified LTB26 and LTB57 were selected as two models of LTB mutant adjuvants to analyze the differential gene expression. Compared with the group of PBS treated mice, 375 differentiation genes were identified (240 upregulation and 135 downregulation) in the group of LTB+VP8 treatment mice, and 654 differentiations (523 upregulation and

131 downregulation) in the group of LTB26+VP8 treatment. Similarly,

725 differentiations (610 upregulation and 115 downregulation) in the

group of LTB57+VP8 treatment, and 890 differentiations (746

upregulation and 144 downregulation) in the group of VP8 alone treatment,

respectively (Table 1).

The Gene ontology (GO) pathway analysis suggested that LTB26 activated

several B cell related signal pathways, APC (antigen presenting cell)

pathways and MHC II pathway (Fig. 5A). Instead, LTB57 just upregulated

inflammatory related signal pathway (Fig. 5B). However, LTB preferred

to activate inflammatory response, upregulate TNF and IL-10 expression

(Fig. 5C). The results indicated that LTB26 acted as adjuvant via

enhancing MHC II related APCs activation and B cell related signal

pathway activation (Fig. 5A). That was consistent with the result of SL727-

LTB26RBD described above (Fig. 2A, B). Therefore, the molecular

mechanism of LTB26 adjuvant was the same, either by binding the

receptor into the cell or by expressing it directly in the cell.

Compared with the expression of immune-related cluster of differentiation (CD) genes, the LTB26 upregulated thirteen B cell associated CDs (CD79a, CD79b, CD19, CD22, CD14, CD37, CD38, CD40, CD48, CD52, CD72, CD74 and CD180), one macrophage marker (CD68) and one common lymphocyte marker (D300c2). Among the fifteen upregulated genes, about 86.7% belonged to B cell associated CDs (Table 2). Of course, LTB upregulated two T cell (and NK cell) marker (CD2, CD8a), one macrophage marker (CD68) and one monocyte, macrophage and DC common marker (CD14). However, B cell marker was no variation in LTB treatment (Table 2). Similarly, LTB57 upregulated three T cell associated CDs (CD47, CD63, and CD164), one lymphocyte homing receptor (CD44), one lymphocyte differentiation marker (CD52), one leukocyte marker (CD53), one neutrophils marker (CD177) and one mast cell marker (D300ld3), respectively (Table 2). The T cell related CDs accounted for about 37.5%. However, LTB57 lost the ability of BCR related activation.

More importantly, CD79a, CD79b and CD19 were the major component of BCR, which indicated that LTB26 functioned adjuvanticity via BCR pathway. The variation of BCR downstream genes showed that LTB26 upregulated the expression of BCR downstream genes more than 2-fold. They were following from Syk, Plc-γ, Ras (Rasa3 and Rras), ERKs (Map3k1 or ERK1), Mapk1ip1l (ERK2), Map4k1 (ERK4)) to transcriptional factors Jund and Atf6b (Table 3). However, LTB57 significantly upregulated the expression of Bcl-10 and the downstream of NFκB inhibitors (Nfκbiz, Nfκbia, and Nfκbib). Otherwise, the upregulation of Mapk6, Akt3 and two transcriptional factors (Egr-1 and Crebrf) did not contribute to adjuvant activity of LTB57 (Supplementary Table 2). Unexpectedly, LTB only upregulated Jun expression (half-level of Jund of LTB26) and significantly enhanced the expression of two NFκB inhibitors (Nfκbia and Nfκbiz) (Table 3). The result suggested that the inhabitation of NFκB activity induced adjuvanticity damage in LTB57 vaccination.

Quantitative real-time PCR (q-PCR) verifies the results of transcriptome data. In order to verify the transcriptome data and confirm the above-mentioned APCs, MHC II and BCR pathways, CD8a, CD79a, CD4, IL-1β, IL-4, TNF-α, Jun, Junb, Jund, H2-Ab1, and NFκB1 of PBMC (peripheral blood mononuclear cell) were tested by quantitative real time polymerase chain reaction PCR(q-PCR). The result indicated that ratio of CD4, CD8a, CD79a, IL-4, Jun, H2-Ab1, and NFκB1 were significantly upregulated in LTB26 treatment group compared with LTB and LTB57 treatments, respectively. The increased expression of IL-4 suggested that LTB26 activated Th2 cells response. Meanwhile, the expression of IL-1β, Jund, CD68, TNF-α and Junb were significantly down regulated in LTB26 treatment (Fig. 5D). The result confirmed the transcriptome analysis that LTB26 activated BCR and MHC II pathways and decreased inflammatory response via downregulation of IL-1β and TNF-α.

IL-10 was also commonly produced in Th2 cells and regarded as an anti-inflammatory cytokine and stimulated the proliferation of B cells(Mannino,

Zhu et al., 2015, Mocellin, Marincola et al., 2005). Meanwhile, IFN-γ made

Naive CD4[+] T cells differentiate to Th1 cells(Zhang, Zhang et al., 2014).

Therefore, IFN-γ, IL-10 and IL-21were also detected by qPCR. Compared

with LTB and LTB57 adjuvant, the LTB26 treatment mice were

significantly increased expression of IL-10 and IL-21 in lymphocytes (Fig.

3E, F, p<0.001). Because IL-21 played a critical role in T cell-dependent

B cell activation(Sondergaard & Skak, 2009, Tangye & Ma, 2020), the

result suggested that LTB26 increased the function of T, B and DCs.

Similarly, the expression of IFN-γ was significantly increased in LTB26

and LTB57 treatment mice compared with the LTB adjuvant, indicating

LTB26 also activated Th1 cells (Fig. 5G, p<0.001). However, there were

no significantly change between LTB26 and LTB57 treatment (Fig. 5G,

p>0.05). Taken together, LTB26 significantly upregulated both Th1 and

Th2 cells activation.

The characteristic of FCM of PBMCs. Peripheral blood was sampled to

confirm the function of B cell, T cell and APCs elicited by purified LTB

and the two mutants (LTB26, LTB57) at 24 h of nasal vaccination. The proportion of CD19$^+$CD45R$^+$ B cells in BPMCs were 17.82$\pm$0.61% (PBS)，24.91 $\pm$ 0.82% (LTB26+VP8), 20.08 $\pm$ 0.73% (LTB57+VP8), 21.57 $\pm$ 1.24% (LTB+VP8) and 20.32 $\pm$ 0.61% (VP8), respectively. Compared with VP8 treatment, LTB26+VP8 and LTB+VP8 treatment were significantly increased the proportion of CD19$^+$CD45R$^+$ B cells in BPMCs than that of LTB57+VP8 vaccination (Fig. 6A, p<0.05, P<0.01), respectively. Meanwhile, there were no significant differences between LTB57+VP8, and VP8 vaccination (Fig. 6A, p>0.05). However, the proportion of CD19$^+$CD45R$^+$ B cell in groups of LTB26+VP8 and LTB+VP8 were no significantly difference (Fig. 6A, p>0.05).

As to the proportion of CD3e$^+$ NK1.1$^+$ T cells in PBMCs, there were no significant variations among the groups of LTB26+VP8 (8.22$\pm$0.60%), LTB57+VP8 (7.47$\pm$0.51%), and LTB+VP8 (8.34$\pm$0.48%) compared with VP8 treatment (7.71$\pm$0.63%), respectively (Fig. 6B, p>0.05).

The proportion of CD11c$^+$CD11b$^+$IA/IE$^+$ DCs in BPMCs were 0.76 ± 0.01% (PBS), 1.23 ± 0.03% (LTB26+VP8), 0.91 ± 0.01% (LTB57+VP8), 1.08±0.06% (LTB+VP8), and 0.88±0.03% (VP8), respectively. The proportion of CD11c$^+$CD11b$^+$IA/IE$^+$ DCs were significantly increased in LTB26+VP8 treatment compared with that of LTB57+VP8, LTB+VP8, and VP8 treatments (Fig. 6C, P<0.001, P<0.01, P<0.001), respectively. That means LTB26 upregulated DCs activation than its wild type.

The F4/80 antigen as a major macrophage marker is expressed on mature macrophages and a subpopulation of DCs (APCs)(Dos Anjos Cassado, 2017). F4/80$^+$ cells were major MHC II$^+$ mature macrophages with APC function; therefore, the proportion of F4/80$^+$ cells in CD11c$^+$CD11b$^+$IA/IE$^+$ DCs was a very important functional indicator of DCs. In this study, the proportion of F4/80$^+$ cells in DCs were 1.85±0.11% (PBS), 3.61±0.37% (LTB26+VP8), 2.33±0.10% (LTB57+VP8), 2.79±0.26% (LTB+VP8), 0.28±0.11% (VP8), respectively. Compared with LTB26+VP8 treatment,

the proportion of F4/80$^+$ cells in DCs were significantly decreased in

LTB57+VP8, LTB+VP8, and VP8 treatments (Fig. 6C, P<0.01，P<0.05，

P<0.01), respectively. That meant LTB26 were significantly upregulated

MHC II$^+$ APCs (macrophages and a subpopulation of DCs) activation than

its wild type (p<0.05). Summarily, FCM analysis also confirmed the GO

data that LTB26 activated MHCII$^+$ APCs function and increased B cell

activation in line with LTB (wild type).

Immunohistochemical staining test confirms the mechanism of LTB26

adjuvanticity. IHC was performed to verify the transcriptome and FCM

data. IHC of extent and intensity (EI) score of 0-3 were considered low

expression and EI score >3 were considered high expression. The EI values

of splenic CD11b were 5.5, 2.875, 2.125, and 1.875 in groups of

LTB26+VP8, LTB57+VP8, LTB+VP8, and VP8, respectively. The result

suggested that CD11b were significantly upregulated more than 2-fold by

LTB26 than that of LTB57, LTB and VP8 treatments (Fig.7A). While the

EI values of splenic CD45 were 3, 1.125, 1.25, and 1.375 in groups of

LTB26+VP8, LTB57+VP8, LTB+VP8, and VP8, respectively. The result suggested that CD45 were significantly upregulated more than 2-fold by LTB26 than that of LTB57, LTB and VP8 treatments (Fig. 7A). Of course, the level of CD45 was slightly decreased in LTB26+VP8 treatment than that of CD11b. The EI values of splenic CD4 were 4.125，3.75，1.625, and 1.875 in groups of LTB26+VP8, LTB57+VP8, LTB+VP8, and VP8, respectively. The result suggested that CD4 were significantly upregulated more than 2-fold by LTB26 and LTB57 than that of LTB treatments (Fig. 7A). Nonetheless, the EI value of LTB26 treatment was higher than that of LTB57.

NLRP3 was one of NOD-like Receptors (NLRs) and played a crucial role in Alum-based adjuvant immune by activating of NLRP3 inflammasome and IL-1β production via MAPK signaling pathway(Sun, Ji et al., 2017). However, the expression of NLRP3 in nasal tissue of the purified LTB, LTB26 and LTB57 adjuvant mice was lower than that of in VP8 alone

treatment (Fig. 7B, p<0.01). The result indicated that LTB and its mutants

functioned mucosal adjuvant independent on NLRs activation.

Modeling the three-dimensional structure of LTB26

The wild type of LTB hit a protein named subunit of heat-labile enterotoxin

B (template serial number: 5lzi.1.A) as a reference template for this

modeling in SWISS-MODEL database. Then 3D model of LTB26, LTB34,

LTB57 and LTB85 monomer and pentamer structures were established

(Fig. 8 A, B, C, D, E, F, G, H, I, J). The result showed that LTB26 changed

the first α-helix of LTB (epitope P1, EP1) to a β–sheet and endowed

LTB26 more higher adjuvanticity than LTB with unknown mechanisms

(Fig. 3D)(Guan, Liu et al., 2015, Takahashi et al., 1996). Intriguingly, the

octapetide (M26S27N28T32I33N34) mutation of LTB26 enhanced the

GM1-binding capacity instead. It indicated that the first α-helix of LTB

hindered its adjuvanticity, not its amino acid sequences. LTB34 mutation

covered three amino acid of the C-terminal of EP1 and three amino acid of

the   N-terminal   of   EP2.   However,   the   decapeptide   mutation

(L35S36L37S38N39S40T41I42N43Y44) of LTB34 were maintained the structure of the second β–sheet and impaired neither GM1-binding affinity nor adjuvanticity (Fig. 3D). Thus, this fragment was no contribution to both of GM1-binding affinity and adjuvanticity of LTB. The residue of Y39 of LTB was reported to involve in stabilizing the GM1-binding(Holmner et al., 2011, Ma, 2016). It was suggested that Y39N mutation did not damage it stabilizing effect.

Instead, the LTB57 mutant maintained the structure of the third β–sheet similar to LTB, however, the amino acids mutation led to lose its adjuvanticity. This mutation (F57Y58E60F61H62H63) was located in the central of epitope EP3 (48-63) and EP4 (54-69) and significantly impaired the adjuvanticity (Fig. 3D)(Guan et al., 2015). However, it was separated two residues from the downstream of the typical GM1-binding residue of G33 (numbered G54 in this study) (Fig.3B, C, D)(Ma, 2016, Nashar, Webb et al., 1996, Turcanu, Hirst et al., 2002). Similarly, LTB85 (86-91 aa) mutation was also located in the central T- and B-cell epitope EP5 and

retained the second α-helix of LTB(Guan et al., 2015, Ma, 2016). Even

though the K91 of LTB57 (K91H) was a conserved GM1-binding residue,

the hexapetide mutation (Y86V87E88F89H90H91) was not impaired the

GM1 affinity, too (Holmner et al., 2011, Ma, 2016). However, both LTB57

and LTB85 lost their adjuvanticity for destroying the sequences of T- and

B-cell epitope of LTB (Fig.3C, D). Therefore, the sequences of epitopes of

EP3, EP4 and EP5 were important than that of 3D structures for

maintaining the adjuvanticity of LTB.

Discussion

COVID-19 was the most dangerous infectious disease to threat public

health since December 2019 causing more than 89 million cases infection

with more than 2.1 million death to date(Harrison, Lin et al., 2020).

Vaccines could help prevent the spread of COVID-19 and end this

pandemic. Since on Dec. 18, 2020, FDA had approved two COVID-19

vaccines, several vaccines have been approved and vaccinated in many

countries. However, the constant mutation of SARS-COV-2 was not only

challenged the effectiveness of the vaccine, but also increased its infectivity(Chen, Wang et al., 2020, Plante, Liu et al., 2020). Previous study suggested that G614 mutation might reduce the ability of vaccines in clinical trials to protect against COVID-19(Plante et al., 2020). Unfortunately, several variations (A348T, R408I, K417N, G476S, V483A, E484K, N501Y, H519Q, A520S) were found at the RBD region (328-533aa, DOI:10.1101/2020.12.24.20248822v1; DOI: 10.1101/2020.12.21. 20248640)(Gu & Chen, 2020, Lokman, Rasheduzzaman et al., 2020). N501Y mutant was recently reported with the increased virulence and pandemic worldwide(Gu & Chen, 2020). A strain, VOC 202012/01, was found in British accumulated 17 amino acid mutations including 69-70del, N501Y, P681H, and other 4 mutations in spike protein (DOI:10.1101/2020.12.24.20248822v1). The 69-70 deletion was linked to immune escape in immunocompromised patients and enhanced viral infectivity in vitro (DOI:10.1101/2020.12.24.20248822v1). Other mutants with K417N or E484K were found in South Africa.

Therefore, it remained to be monitored whether the current vaccines would be effective against these mutants in the future. Therefore, the battle against COVID-19 would continue for some time because of insufficient supplies of vaccines. However, RBD was still the effective antigen for vaccine development. We, based on this, developed an attenuated *Salmonella typhimurium*-based RBD oral vaccine fusing LTB26 as mucosal adjuvant. Then the RBD mutants' vaccine will be constructed on the basis of it in the future.

Salmonella ssp was broadly used as oral vaccine delivery(Yurina, 2018). Attenuated *Salmonella typhimurium* was the most widely used vectors. It was suitable for oral immunization and boosted both mucosal and systemic immune responses(Yurina, 2018). Therefore, as a mucosal infectious pathogen, *Salmonella*-based oral vaccine was suitable for SARS-COV-2 prevention. Another advantage of *Salmonella*-based oral vaccine was that the recombination delivery vehicle was easy to construct. Meanwhile, *Salmonella*-based oral vaccine was essentially DNA vaccine and the

antigenicity of RBD was same as the production of infectious SARS-COV-

2 spike protein. Although *Salmonella*-derived PAMPs (pathogen-

associated molecular patterns) has adjuvant activity, we found that its

adjuvanticity was weaker than LTB26 (Fig. 2)(Bivona, Cerny et al., 2016).

Therefore, we designed LTB26RBD fusion gene to enhance the immune

response in this study.

Mucosal immunity was one of the important ways to protect the body from

the mucosal invading pathogens(Lycke, 2012, Ma, 2016, Perez-Lopez,

Behnsen et al., 2016). However, mucosal adjuvant research lagged

behind vaccine development and resulted in vaccines failure in some

case(Atmar & Keitel, 2009, Young, Sadarangani et al., 2015). Therefore,

it was also important to develop safe and high effective mucosal adjuvants

to enhance the effectiveness of mucosal vaccines.

LT could regulate the differentiation of B cells and regulate the production

of T cells through mucosal immunity(da Hora et al., 2011, Holmner et al.,

2011). However, the toxicity of LT prevented its application as adjuvant in

human. In recent years, non-toxic LTB adjuvant had been widely used(Cunha, Moreira et al., 2017, Ghazali-Bina et al., 2019, Holmner et al., 2011, Liu, Ma et al., 2016, Ma, Luo et al., 2012). The binding specificity of LTB to GM1 was associated with adjuvant activity. Previous studies reported that the property of LTB adjuvant was dependent on GM1-binding affinity(de Haan et al., 1998). The LTB (G33D, numbered G54 in this study for counting the N-terminal signal peptide) mutant might lose the immune adjuvant activity for failing bind to GM1(de Haan et al., 1998). However，GM1 binding sites also existed in other epitopes of LTB such as 51E, Ile58, 58I and 91K(Holmner et al., 2011). Therefore, we had mentioned that beyond GM1-binding, G33 of LTB was a crucial antigenic determinant residue located in the B- and T-cell epitope region (residue S26 to G45)(Ma, 2016). Scientist supposed that G33 residue was crucial for the binding affinity of LTB to GM1 and LTB (G33D) mutant lost its GM1-binding affinity and oral immune adjuvant activity(Nashar et al., 1996).

Even so, we had deduced that LTB (G33D) destroyed the function of the

key residue of LTB antigenic determinant (residue S26 to G45) rather than

the GM1-binding affinity(Ma, 2016). Therefore, the integrity of LTB

epitope was more important than the GM1-binding affinity(Ma, 2016).

This hypothesis has been verified that LTB57 (57-63 aa) mutation

(F57Y58E60F61H62H63) was located in the central T- and B-cell epitope

and significantly impaired the adjuvanticity (Fig.3D)(Ma, 2016, Nashar et

al., 1996, Turcanu et al., 2002). Intriguingly, the octapetide (M26S27 N28-

-T32I33N34) mutation in LTB26 was not only impaired the GM1 affinity,

but also enhanced the GM1-binding capacity. Similarly, LTB85 (86-91 aa)

mutation was also located in the central T- and B-cell epitope(Ma, 2016),

and this hexapetide (Y86V87E88F89H90H91) mutation was not impaired

the GM1 affinity, too. Even though the K91 of LTB57 (K91H) was a

conserved GM1-binding residue(Holmner et al., 2011, Ma, 2016).

However, both LTB57 and LTB85 lost their adjuvanticity for destroying

the structure of T- and B-cell epitope of LTB (Fig.3D). Therefore, the

highlight of this work suggested that the adjuvanticity of LTB was not positively associated with GM1-binding affinity.

Adjuvant stimulated APCs and targeted the innate immune system to induce a robust immune response. In principle, adjuvant could induce host-derived damage-associated molecular patterns (DAMPs) or recognizing PAMPs to enhance antigen-specific immune responses via APCs (DCs) or macrophages[Hayashi, Momota et al., 2018]. DCs sensed and phagocytized invading pathogens and activated naïve T cells. Then they acted as a major link between innate and acquired immunity to determine the polarization of T cell responses into different effector subtypes(Tesfaye, Gudjonsson et al., 2019) (Steinman & Hemmi, 2006).

Cytokines were important immune regulators. Purified LTB26 was capable of orchestrating upregulated IL-4, IL-10 and IL-21 expression, significantly, compared with LTB and LTB57 (Fig. 5D, 5E, 5F). Generally, B cells and CD8$^+$ T cells were the responders to IL-21 by IL-21R(Sondergaard & Skak, 2009). CD4$^+$ T cells were main producers of IL-

21, while TCR stimulation increased CD4[+] T cell expression of IL-21R as a positive feedback. IL-10 stimulated cytotoxicity of CD8[+] T cells and IFN-γ expression(Oft, 2014).

IL-4 promoted humoral immunity and IFN-γ promoted cell-mediated immunity(El-Sissi, Mohamed et al., 2020). LTB, LTB26 and LTB57 significantly upregulated the expression of IFN-γ and TNF-α compared with VP8 alone treatment, but there were no differentiation among LTB, LTB26 and LTB57 (Fig. 5G). Thus, purified LTB26 promoted both Th1 and Th2 cells mediated immunity via upregulating IL-4 and IFN-γ expression, which was different from Alum-based vaccine (IL-4 only). This consistent with the data obtained from FCM and transcriptome analysis.

Together our data indicated that LTB26 adjuvant fused with SARS-COV-2 RBD or purified and mixed with hRV VP8 significantly increased the proportion of MHC II[+] DCs and CD19[+]CD45[+] B cells in peripheral blood, which indicated that LTB26 enhanced the processing of vaccine (antigen)

via DCs activation and the producing of antibody by B cell activation (Fig

1, 6). We found *Salmonella typhimurium* SL7207 was not an ideal vehicle

for oral vaccine development for its toxicity. In this study, the BALB/c

mice and SD rats were died within 24 h after intraperitoneally injection

with the same dose of oral vaccination SL7207 (data not show). Therefore,

the less toxicity *Salmonella* strain or other vehicles were necessary in

future research. Furthermore, the mucosal responses of SL7207-

LTB26RBD will be performed as a single topic in the future.

Material and Methods

Construction of pcDNA3.1-LTB26RBD plasmid. LTB26 was derived

from LTB (Genbank No. EU113252) with I26MT27SE28NL29ΔC30Δ

(L29C30 deletion) and E32TY33IH34N mutation. SARS-COV-2 RBD

(301-521 aa, Genbank No. YP_009724390) was fused with GSGSG linker

at C-terminal of LTB26. The sequence of LTB26RBD and RBD were

commercially synthesized and inserted into pcDNA3.1(+) plasmid at

*Bam*H I/*Not* I position to construct pcDNA3.1-LTB26RBD and

pcDNA3.1-RBD plasmids, respectively. Then the plasmids were electrically transformed into attenuated *Salmonella typhimurium* SL7207 (SL7207) as oral vaccine candidates (SL7207-RBD, SL7207-LTB26RBD) at 2.5kv and confirmed by sequencing.

Animal and Immunization. BALB/c mice of 3-4 weeks (male) were bred in the experimental animal center of Chongqing Medical University and divided into four groups. Six groups of mice (n=5) were oral vaccinated with *Salmonella* SL7207, SL7207-RBD, SL7207-LTB26RBD and 200 μl PBS (2.5×10^6 cfu /mouse), respectively, after treating with saturated sodium bicarbonate (0.2 ml/mouse). The mice were boosted twice on 10, and 20 day as described above. Blood was sampled from the orbital vein on day 0, 10, 20, 30.

This study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. The protocol was approved by the Committee on the

Ethics of Animal Experiments at the Chongqing Medical University (SYXK2012-0001, 2013-03-11).

Flow Cytometry. PBMCs from heparinized blood were isolated with Ficoll-Paque. The PBMC samples were stained with 5 µl of anti-mouse CD3e-PerCP/cy5.5 (Biolegend, USA, cat#100327), NK1.1-PE (Biolegend, USA, cat#108707),CD19-PE (Biolegend, USA, cat#115511), CD45R/B220-APC (Biolegend, USA, cat#103207), CD11c-APC (Biolegend, USA, cat#117309), CD11b-FITC (Biolegend, USA, cat# 101205), F4/80-PE (Biolegend, USA, cat#123109), and IA/IE-PerCP/cy5.5 (Biolegend, USA, cat#107625), respectively, for 30 min at RT in the dark. The cell were stained with 5 µl of PerCP/cy5.5-Armenian Hamster-IgG (Biolegend, USA, cat#400931), PE-Rat-IgG2a (Biolegend, USA, cat#400508), PE-Rat- IgG2b (Biolegend, USA, cat#400211), APC-Rat-IgG2a (Biolegend, USA, cat#400511), APC-Rat-IgG1 (Biolegend, USA, cat#401903), FITC-Rat-IgG2b (Biolegend, USA, cat#400634), PerCP/cy5.5-Rat IgG2b (Biolegend, USA, cat#400631), and respective

isotype control for 30 min at RT in the dark, respectively. Then washed

twice with 500 µl PBS and assessed by four-colored flow cytometry. Then

measured the percentages of CD3e⁺NK1.1⁺ NK cells, CD19⁺CD45R⁺ B

cells and CD11c⁺CD11b⁺F40⁻80⁺IA/IE⁺ DCs (dendritic cells) and the

fluorescence intensity (MFI) of cell finally.

SL7207-LTB26RBD immune assay. Blood samples were individually

collected from oral immunized mice by orbital venous bleeding on days 0,

10, 20 and 30 for the analysis of systemic RBD specific antibodies (n=5).

All the samples were treated as previously described(Liu et al., 2016). The

antibodies were analyzed with horseradish peroxidase (HRP)-labeled goat

anti-mice IgG and goat anti-mice IgA (1.0 µg/ml, Boster, Wuhan, China)

by ELISA, respectively, as previously described(Liu et al., 2016). SARS-

CoV-2 RBD-His protein was purchased from Nanjing Okay

Biotechnology Co., Ltd (Nanjing, China, cat#K1516). Endpoint titers were

determined as the dilution of each sample showing a 2.1-fold higher

absorbance level of 450 nm as compared to that of the negative control samples. Average OD_{450} values for the animals were calculated.

LTB mutants design. Full-length LTB DNA was cloned from EC44815 strain and constructed pET32a-LTB plasmid as our previously report(Ma et al., 2012). The four LTB mutants were designed to mutate some amino acids located in the B- and T-cell epitope region(Ma, 2016). The amino acid was numbered from the first M (Met) of signal peptide in LTB. The four LTB mutants were also constructed into plasmid pET32 at *Bam*H I /*Sal* I site, respectively. Full-length of hRV VP8 DNA (GenBank: L34161) was commercially synthesized (Sangon, Shanghai, China) and inserted into plasmid pET32 at *Bam*H I/*Sal* I site. The recombinants of pET32-LTB, pET32-LTB26, pET32-LTB34, pET32-LTB57, pET32-LTB85 and pET32-VP8 were expressed in *E. coli* BL21 cells and purified with BeaverBeads™ His-tag protein purification kits (Beaver, Suzhou, China, cat#70501-5). The endotoxin was removed using ToxinEraserTM resin (Genscript, Nanjing, China, cat#L00338) and the protein concentration

was measured by BCA protein assay kit (TaKaRa, Dalian, China, cat#T9300A) as previous described(Ma et al., 2012).

Animal and Immunization. Six groups of BALB/c mice (3-4 weeks old, male, n=6) were nasal vaccinated with human rotavirus VP8 (VP8), LTB+VP8, LTB26+VP8, LTB34+VP8, LTB57+VP8, LTB85+VP8, and 20 μl PBS, respectively, after anesthetizing with chloral hydrate (0.5 mL/100 g). The VP8, LTB and four LTB mutants were each added 10 μg/mouse respectively. The mice were intranasal boosted twice on 7, and 14 day with same method after first vaccination. All surgery was performed under sodium pentobarbital anesthesia, and euthanized by cervical dislocation. Then immune assay was performed as described above to detect the VP8 specific antibodies from blood, fecal and bronchial mucosal washing, respectively. The PBMCs FCM analysis was performed as previous described.

GM1-binding analysis. The GM1-binding affinity of LTB and its four mutants were determined using GM1-ELLSA (enzyme-linked

immunosorbent assay) assay[(Minke, Roach et al., 1999)]. The wells of microplate were coated with 200 µL (2.0 µg/ml and 10 µg/ml, respectively) of GM1 (Qilu pharma, Shandong, China) or PBS, respectively, at 48°C overnight. Plates were washed three times with 500 µl PBS to remove uncombined GM1. Subsequently, plates were blocked by the addition of 200 µl PBS containing 1.0 % BSA at 37°C for 30 min. Then, plates were washed again as described above. Finally, plates were incubated with 1000 ng/well LTB, LTB26, LTB34, LTB57, LTB85, and PBS, respectively, at 37°C for 2 h, followed by washing as described above. 100 µl of rabbit anti-His-tag antibody (1:1000, BioVision, USA, cat#3998) was incubated at 37°C for 2 h and washed as described above. 100 µl of HRP conjugated goat anti-mouse IgG(H+L) secondary antibody (1:1000, Boster, China, cat# BA1051) was incubated at 37°C for 2.5 h, followed by washing four times. Finally, added 100 µl of TMB (tetramethylbenzidine) to each well and incubated at 37°C for 5 min. Then the reaction was stopped by 200 µl H_2SO_4 (2.0 mol/l). The OD_{450} was read by microplate Reader.

Transcriptome analysis of peripheral lymphocyte. According to animal immune data, the peripheral lymphocytes were sampled in groups of LTB, LTB26+VP8 (the highest immune adjuvanticity with middle GM1-binding capacity), LTB57+VP8 (the highest GM1-binding affinity with lowest immune adjuvanticity), VP8 (in PBS), and PBS treatment mice after 24 hours vaccination. Total RNA was extracted using TRIzol Reagent (Invitrogen, USA) according the manufacturer manual. The RNA quality were measured using a NanoDrop 2000 spectrophotometer (Thermo Fisher, USA) at 260/280 nm and agarose gel electrophoresis. Then RNA quantity was measured using Qubit2.0 Fluorometer. mRNA sequencing was performed on Illumina Hiseq platform (Illumina, USA) with 12 G bps and 10 M reads (Genminix Informatic Ltd., Shanghai, China).

The differentially expressed genes were selected as having more than 2-fold difference between their geometrical mean expression in the compared groups and a statistically significant P-value (<0.05) by analysis of DEseq2. The GO analysis on differentially expressed genes was performed with an

R package: cluster profiler using a P<0.05 to define statistically enriched GO categories. Pathway analysis was used to determine the significant pathway of the differential genes according to Kyoto Encyclopedia of Genes and Genomes Database (http://www.genome.jp/kegg/).

Peripheral blood qPCR analysis. Total RNA was isolated as previous described. RNA reverse transcription was performed using the TaqMan RNA Reverse Transcription kit (Thermo Fisher, USA, cat#4366596). qPCR was performed using SYBR Green Supermix (Bio-Rad, USA, cat# 1708882) and the iCycler iQ Real-Time PCR system (Bio-Rad, USA). The reactions were run as follow: denaturation at 95°C for 10 min followed by 50 cycles of 95°C for 10 sec, 55°C for 10 sec and 72°C for 5 sec; 99°C for 1 sec; 59°C for 15 sec; 95°C for 1 sec; then cooling to 4°C. Relative mRNA expression was normalized against the endogenous control, GAPDH, using the 2-ΔΔCt method. The primers used in the current study were listed as table (Supplementary Table 1).

Immunohistochemistry. Spleen played an important role for lymphocyte

migration and immune response after receiving antigen stimulation. Therefore, CD4, CD11b, and CD45 of splenic lymphocyte were tested after 24h poster vaccination. Haematoxylin and eosin staining and antibody labeling were performed on 4-μm tissue sections as described previously(Liu et al., 2016). Antibodies(rabbit) against CD11b, CD45 and CD4 were incubated overnight at 4℃. Antibodies (goat) against IgG (rabbit) were used for labeling for 2 h at room temperature. Images of histology and immunohistochemistry were taken with a Nikon Eclipse E600 and Nikon DS-Ri1 camera or with a Digital Slide Scanner (20 × magnification).

The expressions of CD11b, CD45 and CD4 were quantified using a visual grading system based on the extent of staining as previously described(Marechal, Demetter et al., 2009). Briefly, percentage of positive spleen cells (extent of staining) was graded on scale from 0 to 3: 0, none; 1, 1-30%; 2, 31-60%; 3, 41-60%. The intensity of staining was graded on a scale of 0-3: 0, none; 1, weak staining; 2, moderate staining; 3, strong

staining. The combination of extent (E) and intensity (I) of staining was

obtained by the product of E times I called EI varying from 0 to 9 for each

spot. For statistical analysis, EI score of 0-3 were considered low

expression and EI score >3 were considered high expression.

Three-dimensional structure prediction

In order to understand the enhanced adjuvanticity of LTB26, the three-

dimensional structures of LTB26, LTB34, LTB57 LTB85 and LTB were

analyzed using SWISS-MODEL project

(https://swissmodel.expasy.org/interactive/HTEqUf/models/). The quality

of model was evaluated using the SWISS-MODEL scoring function of

GMQE, QMEAN4, Identity, and Method parameters. GMQE [0, 1] score

estimation results of the tertiary structure of the model accuracy, the higher

the better; QMN4 is to evaluate the global and local quality of the model

to ensure the reliability of the overall quality of the model. QSQE combines

interface protection, structural clustering, and other template features to

provide quaternary structural quality estimates that reflect the expected

accuracy of interchain contact based on a given alignment and templated

model. The higher the number, the higher the reliability, which

complements the GMQE score. Identity stands for sequence alignment

matching rate, the higher the better.

Acknowledgements

The work was funded by a research grant from the National Natural
Science Foundation of China (NSFC no. 30972585); We thank professor
Shengyong Yang (Chongqing Medical University) proofread the
manuscript and gave some constructive advises.

Contributions

Y.M. conceived of the paper; Q.S. and Q.W. contributed equally and
performed main experiments; S.C., S.G. and C.F. performed some
experiments; Y.M., Q.S., F.S. and T.L. contributed to write the paper.

Competing interests

No competing interests.

Supplementary material: Supplementary Table 1. qPCR primers used in
this study.

Reference

Atmar RL, Keitel WA (2009) Adjuvants for pandemic influenza vaccines. *Current topics in microbiology and immunology* 333: 323-44

Bivona AE, Cerny N, Alberti AS, Cazorla SI, Malchiodi EL (2016) Attenuated Salmonella sp. as a DNA Delivery System for Trypanosoma cruzi Antigens. *Methods in molecular biology (Clifton, NJ)* 1404: 683-695

Chen J, Wang R, Wang M, Wei GW (2020) Mutations Strengthened SARS-CoV-2 Infectivity. *Journal of molecular biology* 432: 5212-5226

Clemens JD, Sack DA, Harris JR, Chakraborty J, Neogy PK, Stanton B, Huda N, Khan MU, Kay BA, Khan MR, et al. (1988) Cross-protection by B subunit-whole cell

cholera vaccine against diarrhea associated with heat-labile toxin-producing enterotoxigenic Escherichia coli: results of a large-scale field trial. *J Infect Dis* 158: 372-7

Clements JD, Norton EB (2018) The Mucosal Vaccine Adjuvant LT(R192G/L211A) or dmLT. *mSphere* 3

Cunha C, Moreira CJ, Rocha A, Finger PF, Magalhaes CG, Ferreira MRA, Dellagostin OA, Moreira AN, Conceicao FR (2017) Parenteral adjuvant potential of recombinant B subunit of Escherichia coli heat-labile enterotoxin. *Memorias do Instituto Oswaldo Cruz* 112: 812-816

da Hora VP, Conceicao FR, Dellagostin OA, Doolan DL (2011) Non-toxic derivatives of LT as potent adjuvants. *Vaccine* 29: 1538-44

de Haan L, Verweij WR, Feil IK, Holtrop M, Hol WG, Agsteribbe E, Wilschut J (1998) Role of GM1 binding in the mucosal immunogenicity and adjuvant activity of the Escherichia coli heat-labile enterotoxin and its B subunit. *Immunology* 94: 424-30

Dos Anjos Cassado A (2017) F4/80 as a Major Macrophage Marker: The Case of the Peritoneum and Spleen. *Results and problems in cell differentiation* 62: 161-179

El-Sissi AF, Mohamed FH, Danial NM, Gaballah AQ, Ali KA (2020) Chitosan and chitosan nanoparticles as adjuvant in local Rift Valley Fever inactivated vaccine. 10: 88

Ghazali-Bina M, Pourmand MR, Mirshafiey A, Bakhtiari R, Khaledi A, Kazemian H, Afshar D, Getso MI, Eshraghi S (2019) Vaccine potential of LenA and LcpA proteins of Leptospira interrogans in combination with Escherichia coli heat-labile enterotoxin, B subunit (LTB). *Iranian journal of microbiology* 11: 39-47

Gu H, Chen Q (2020) Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine efficacy. 369: 1603-1607

Guan W, Liu W, Bao J, Li J, Yuan C, Tang J, Shi D (2015) Analysis and application of a neutralizing linear epitope on liable toxin B of enterotoxin Escherichia coli. *Applied microbiology and biotechnology* 99: 5985-96

Harrison AG, Lin T, Wang P (2020) Mechanisms of SARS-CoV-2 Transmission and Pathogenesis. *Trends in immunology* 41: 1100-1115

Hayashi T, Momota M, Kuroda E, Kusakabe T, Kobari S, Makisaka K, Ohno Y, Suzuki Y, Nakagawa F, Lee MSJ, Coban C, Onodera R, Higashi T, Motoyama K, Ishii KJ, Arima H (2018) DAMP-Inducing Adjuvant and PAMP Adjuvants Parallelly

Enhance Protective Type-2 and Type-1 Immune Responses to Influenza Split Vaccination. *Frontiers in immunology* 9: 2619

Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu NH, Nitsche A, Müller MA, Drosten C, Pöhlmann S (2020) SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 181: 271-280.e8

Holmner A, Mackenzie A, Okvist M, Jansson L, Lebens M, Teneberg S, Krengel U (2011) Crystal structures exploring the origins of the broader specificity of escherichia coli heat-labile enterotoxin compared to cholera toxin. *Journal of molecular biology* 406: 387-402

Liu L, Ma Y, Zhou H, Wu M (2016) Quantitative Proteomic Analysis of Escherichia coli Heat-Labile Toxin B Subunit (LTB) with Enterovirus 71 (EV71) Subunit VP1. *International journal of molecular sciences* 17

Lokman SM, Rasheduzzaman M, Salauddin A, Barua R, Tanzina AY, Rumi MH, Hossain MI, Siddiki A, Mannan A, Hasan MM (2020) Exploring the genomic and proteomic variations of SARS-CoV-2 spike glycoprotein: A computational biology approach. *Infection, genetics and evolution : journal of molecular epidemiology and evolutionary genetics in infectious diseases* 84: 104389

Lycke N (2012) Recent progress in mucosal vaccine development: potential and limitations. *Nature reviews Immunology* 12: 592-605

Ma Y (2016) Recent advances in nontoxic Escherichia coli heat-labile toxin and its derivative adjuvants. *Expert review of vaccines* 15: 1361-1371

Ma Y, Luo Y, Huang X, Song F, Liu G (2012) Construction of Bifidobacterium infantis as a live oral vaccine that expresses antigens of the major fimbrial subunit (CfaB) and the B subunit of heat-labile enterotoxin (LTB) from enterotoxigenic Escherichia coli. *Microbiology (Reading, England)* 158: 498-504

Mannino MH, Zhu Z, Xiao H, Bai Q, Wakefield MR, Fang Y (2015) The paradoxical role of IL-10 in immunity and cancer. *Cancer letters* 367: 103-7

Marechal R, Demetter P, Nagy N, Berton A, Decaestecker C, Polus M, Closset J, Deviere J, Salmon I, Van Laethem JL (2009) High expression of CXCR4 may predict poor survival in resected pancreatic adenocarcinoma. *British journal of cancer* 100: 1444-51

Merritt EA, Zhang Z, Pickens JC, Ahn M, Hol WG, Fan E (2002) Characterization and crystal structure of a high-affinity pentavalent receptor-binding inhibitor for cholera toxin and E. coli heat-labile enterotoxin. *Journal of the American Chemical Society* 124: 8818-24

Minke WE, Roach C, Hol WG, Verlinde CL (1999) Structure-based exploration of the ganglioside GM1 binding sites of Escherichia coli heat-labile enterotoxin and cholera toxin for the discovery of receptor antagonists. *Biochemistry* 38: 5684-92

Mocellin S, Marincola FM, Young HA (2005) Interleukin-10 and the immune response against cancer: a counterpoint. *Journal of leukocyte biology* 78: 1043-51

Nashar TO (2014) The Quest for an HIV-1 Vaccine Adjuvant: Bacterial Toxins as New Potential Platforms. *Journal of clinical & cellular immunology* 5

Nashar TO, Webb HM, Eaglestone S, Williams NA, Hirst TR (1996) Potent immunogenicity of the B subunits of Escherichia coli heat-labile enterotoxin: receptor binding is essential and induces differential modulation of lymphocyte subsets. *Proceedings of the National Academy of Sciences of the United States of America* 93: 226-30

Oft M (2014) IL-10: master switch from tumor-promoting inflammation to antitumor immunity. *Cancer immunology research* 2: 194-9

Perez-Lopez A, Behnsen J, Nuccio SP, Raffatellu M (2016) Mucosal immunity to pathogenic intestinal bacteria. *Nature reviews Immunology* 16: 135-48

Plante JA, Liu Y, Liu J, Xia H (2020) Spike mutation D614G alters SARS-CoV-2 fitness.

Poland GA, Ovsyannikova IG, Crooke SN, Kennedy RB (2020) SARS-CoV-2 Vaccine Development: Current Status. *Mayo Clinic proceedings* 95: 2172-2188

Sondergaard H, Skak K (2009) IL-21: roles in immunopathology and cancer therapy. *Tissue antigens* 74: 467-79

Steinman RM, Hemmi H (2006) Dendritic cells: translating innate to adaptive immunity. *Current topics in microbiology and immunology* 311: 17-58

Sun B, Ji Z, Liao YP, Chang CH, Wang X, Ku J, Xue C, Mirshafiee V, Xia T (2017) Enhanced Immune Adjuvant Activity of Aluminum Oxyhydroxide Nanorods through Cationic Surface Functionalization. 9: 21697-21705

Takahashi I, Kiyono H, Jackson RJ, Fujihashi K, Staats HF, Hamada S, Clements JD, Bost KL, McGhee JR (1996) Epitope maps of the Escherichia coli heat-labile toxin B

subunit for development of a synthetic oral vaccine. *Infection and immunity* 64: 1290-8

Tangye SG, Ma CS (2020) Regulation of the germinal center and humoral immunity by interleukin-21. *The Journal of experimental medicine* 217

Tesfaye DY, Gudjonsson A, Bogen B, Fossum E (2019) Targeting Conventional Dendritic Cells to Fine-Tune Antibody Responses. *Frontiers in immunology* 10: 1529

Turcanu V, Hirst TR, Williams NA (2002) Modulation of human monocytes by Escherichia coli heat-labile enterotoxin B-subunit; altered cytokine production and its functional consequences. *Immunology* 106: 316-25

White JA, Haghighi C, Brunner J, Estrada M, Lal M, Chen D (2017) Preformulation studies with the Escherichia coli double mutant heat-labile toxin adjuvant for use in an oral vaccine. *Journal of immunological methods* 451: 83-89

Young BE, Sadarangani SP, Leo YS (2015) The avian influenza vaccine Emerflu. Why did it fail? *Expert review of vaccines* 14: 1125-34

Yurina V (2018) Live Bacterial Vectors-A Promising DNA Vaccine Delivery System. *Medical sciences (Basel, Switzerland)* 6

Zhang Y, Zhang Y, Gu W, Sun B (2014) TH1/TH2 cell differentiation and molecular signals. *Advances in experimental medicine and biology* 841: 15-44

Table 1. The profiles of differential genes summary

Groups	Upregulation gene numbers	downregulation gene numbers	Total variation gene numbers
LTB+VP8 vs PBS	240	135	375
LTB26+VP8 vs PBS	523	131	654
LTB57 +VP8 vs PBS	610	115	725
VP8 vs PBS	746	144	890

Table 2. The variation of CDs expression in LTB26 and other groups

CDs	LTB+VP8	LTB26+VP8	LTB57+VP8	Adjuvant Significant	Cell type
CD2	2.0110114	N	2.235971319	-	T cell, NK cell
CD14	2.1038286	N	2.825551073	-	monocyte, macrophage, dendritic cell
CD8a	2.1593616	N	n	-	T cell, NK cell
CD68	2.0262211	2.666606105	n	+	macrophage
CD79b	n	4.669013871	n	+	B cell antigen receptor (BCR)
CD79a	n	2.166643886	n	+	BCR
CD48	n	2.404601005	n	+	T cell
CD40	n	2.243108146	n	+	B cell
CD180	n	2.771753281	n	+	B cell
CD74	n	2.811559737	n	+	B cell
CD72	n	2.546204783	n	+	B cell
CD38	n	2.521370117	n	+	B cell
CD22	n	2.982325847	n	+	Mature B cell
CD19	n	3.040386195	n	+	primitive & mature B cell
CD37	n	2.566581649	n	+	B cell
CD300c2	n	2.458761325	n	-	monocytes, neutrophils, partial T & B cell
CD52	n	2.525736598	2.242208379	-	B cell
CD44	n	N	2.058549419	-	B1 cell, IgM>IgG
CD164	n	N	2.137254166	-	T cell
CD63	n	N	2.115720638	-	T cell
CD53		N	2.25028684	-	leukocyte
CD177		N	7.451954307	-	neutrophils
CD47		N	2.041258354	-	T cell
CD300ld3		N	2.117187532	-	mast cells

*:n, no differentiate;   +, LTB26 vs LTB and LTB57 with significant adjuvanticity;

-, no significant adjuvanticity;

Table 3. The variation of BCR downstream genes

genes	LTB+VP8	LTB26+VP8	LTB57+VP8	Adjuvant Significant
Syk	n	2.9179075	2.8384351	-
Plcg2	n	**2.7476057**	n	+
Cdc42ep5	n	4.1154716	7.770395047	-
Rras	n	**2.357993**	n	+
Rasa3	n	**2.2090269**	n	+
Map3k1	n	**2.115917**	n	+
Mapk1ip1l	n	**2.1197696**	n	+
Map4k1	n	**2.2946296**	n	+
Mapk6	n	n	2.2102702	-
Jun	2.2490814	n	n	+
Jund	n	**5.183886**	n	+
Bcl-10	n	n	2.0637835	-
Nfkbia	3.8505929	n	5.6544579	-
Nfkbib	n	n	2.0815464	-
Nfkbiz	3.3298047	n	6.6704407	-
Rasgrp3	n	n	2.0318061	-
Atf1	n	n	2.1382357	-
Atf6b	n	**2.0878157**	n	+
Crebrf	n	n	2.2254203	-
Egr1	n	n	3.2136266	-
Foxo1	n	**2.5975225**	n	+
Akt3	n	n	2.4481639	-

*:n, no difference;   +, LTB26 vs LTB and LTB26 vs LTB57 with significant adjuvanticity;   -, no significant adjuvanticity;

Figures legends

SSC

IA/IE-Percp/Cy5.5

CD11c-APC

CD11b-FITC

F4-80(MHC II)-PE

Figure 1. FCM analysis of dendritic cells (DCs) and B cells.

(A) pcDNA3.1-LTB26-RBD analyzed with restriction enzymes. Lane 1: DNA marker,

lane 2: pcDNA3.1- RBD, lane 2: pcDNA3.1-LTB26-RBD.

(B) LTB26 significantly activated MHC II$^+$ DCs and increased about 29.3% proportion

than that of RBD alone vaccination. There were no significant differences between

Salmonella and Salmonella-RBD vaccination mice. (C) LTB26 significantly activated

CD19$^+$CD45$^+$ B cells and increased about 14.5% proportion than that of RBD alone

vaccination.

Figure 2. Immune responses analysis

Serum IgG titer on post-vaccination of 10 d, 20 d and 30 d, respectively; *, $p < 0.02$.

PBS treatment was used as control group.

Figure 3. The construction and analysis of LTB mutants. (A) LTB and the mutants

were purified and detected in SDS-PAGE. Lane 1: standard proteins; lane 2-6: LTB,

LTB26, LTB34, LTB57, and LTB85; (B) Alignment of LTB26 and LTB34 with LTB;

(C) alignment of LTB57 and LTB85 with LTB. The mutations were highlighted with

black background; (D) B cell and T cell epitopes of LTB and the LTB mutants regions

located in LTB epitopes.

Figure 4. Immune responses and GM1-binding affinity analysis

(A) Serum IgG titer on post-vaccination of 7 d, 14 d and 21 d, respectively; (B) Lung

mucosal sIgA titer on the third robust post-vaccination; (C) 1000 ng/well LTB, LTB26,

LTB34, LTB57, LTB85, and VP8 were incubated with 2.0 µg/ml GM1. LTB57 had the

highest GM1-binding affinity with the lowest adjuvanticity; (D) 1000 ng/well LTB,

LTB26, LTB34, LTB57, LTB85, and VP8 were incubated with 10 µg/ml GM1. LTB57

had the highest GM1-binding affinity with the lowest adjuvanticity; ***, p<0.001; *, p<0.01; #, p>0.05.

Figure 5. GO analysis the adjuvant related signal pathways and qPCR test.

(A)Compared with LTB57, LTB26 significantly activated APCs (MHC II), B cell proliferation, BCR activation and immunoglobulin mediated immune responses, respectively.　(B) Compared with LTB, LTB26 significantly activated Th1 immune response, TNF and IL-10 expression, and immune response. (C) LTB 57 did not active APCs (MHC II) and B cell related signal pathways, but upregulated inflammatory response. (D) Cd79a, H2-Ab1, IL-4, CD4, NκkB1, Cd8a, and Jun were significantly upregulated by LTB26. IL-1β and TNFα were significantly downregulated by LTB26. S26: LTB26; S57: LTB57. (E-G) LTB26 significantly upregulated IL-10 and IL-21expression, respectively. LTB26 and LTB57 were significantly upregulated IFN-γ expression than that of VP8, but there were no differences each other. ****, p<0.001; ***, p<0.01; #, p>0.05.

Figure 6. FCM analysis of B cells and NK cells. (A) LTB26 significantly activated

$CD19^+$ $CD45R^+$ B cells. (B) LTB26 significantly activated $CD19^+$ $CD45R^+$ B cells. (B)

NK1.1 cells were no variations among LTB, LTB26, LTB57 and VP8 alone treatment.

(C) $CD11c^+CD11b^+IA/IE^+$ DCs were significantly increased in LTB26+VP8 treatment

compared with that of LTB57+VP8, LTB+VP8, and VP8 treatments. **, $p<0.001$; *,

$p<0.01$;   #, $p>0.05$.

Figure 7. IHC analysis of CDs and NLRP3. (A) CD11b, CD45, CD4 were

significantly upregulated more than 2-fold by LTB26. (B) NLRP3 was significantly

downregulated by LTB26.****, p<0.001, **, p<0.01; #, p>0.05.

Figure 8. Prediction of LTB and LTB mutants 3D structure.

(A) 3D structure of LTB monomer. (B) 3D structure of LTB pentamer. (C) 3D structure of LTB26 and LTB fitting monomer. The first α-helix of LTB was replaced with β-sheet in LTB26 (red box) (D) 3D structure of LTB26 pentamer. Arrow showed the β-sheet. (E) 3D structure of LTB34 and LTB fitting monomer. The 3D structure was intact. (F) 3D structure of LTB34 pentamer. Arrow showed the first α-helix. (G) 3D structure of LTB57 and LTB fitting monomer. The 3D structure was intact. (H) 3D structure of LTB57 pentamer. Arrow showed the first α-helix. (I) 3D structure of LTB85 and LTB fitting monomer. The 3D structure was intact. (J) 3D structure of LTB85 pentamer. Arrow showed the first α-helix.