1 **An *in vitro* ovarian explant culture system to examine sex change in a**
2 **hermaphroditic fish**
3

4 Goikoetxea Alexander[1], Damsteegt Erin L[2], Todd Erica V[1], McNaughton Andrew[1],
5 Gemmell Neil J[1], Lokman P Mark[2].
6

7 [1]Department of Anatomy, University of Otago, Dunedin, New Zealand
8 [2]Department of Zoology, University of Otago, Dunedin, New Zealand
9

10 **Corresponding author:**
11 Alexander Goikoetxea
12 +34 647 539 500
13 alexandergoikoetxea@gmail.com
14

15 **Keywords**: Organ culture, sex change, previtellogenic oocyte, cortisol, spotty
16 wrasse.

18 **Abstract**
19

Many teleost fishes undergo natural sex change, and elucidating the physiological
and molecular controls of this process offers unique opportunities not only to
develop methods of controlling sex in aquaculture settings, but to better
understand vertebrate sexual development more broadly. Induction of sex change
in some sequentially hermaphroditic or gonochoristic fish can be achieved *in vivo*
through social manipulation, inhibition of aromatase activity, and steroid
treatment. However, the induction of sex change *in vitro* has been largely
unexplored. In this study, we established an *in vitro* culture system for ovarian
explants in serum-free medium for a model sequential hermaphrodite, the New
Zealand spotty wrasse (*Notolabrus celidotus*). This culture technique enabled
evaluating the effect of various treatments with 17β-estradiol (E$_2$), 11-
ketotestosterone (11KT) or cortisol (CORT) on spotty wrasse ovarian architecture

for 21 days. A quantitative approach to measuring the degree of ovarian atresia
within histological images was also developed, using pixel-based machine learning
software. Ovarian atresia likely due to culture was observed across all treatments
including no-hormone controls, but was minimised with treatment of at least 10
ng/mL E_2. Neither 11KT nor CORT administration induced proliferation of
spermatogonia (i.e. sex change) in the cultured ovaries indicating culture beyond
21 days may be needed to induce sex change *in vitro*. The *in vitro* gonadal culture
and analysis systems established here enable future studies investigating the
paracrine role of sex steroids, glucocorticoids and a variety of other factors during
gonadal sex change in fish.

Introduction

In most vertebrates, sex is fixed after birth and remains the same throughout life (1). Some teleost fishes, however, develop as one sex but remain able to change to the other during adulthood (2). *In vitro* culture manipulations enable precise monitoring of the effects of biotic and abiotic factors (e.g. temperature, pH, microplastics, etc.) on gonadal development, but are underutilised in aquaculture and fisheries research. *Ex vivo* approaches show particular potential to study responses of gonadal tissue to sex steroids and other hormones (e.g. glucocorticoids). Testicular and ovarian organ culture has been experimentally achieved with a handful of gonochoristic fish species (3–7). However, with the exception of a conference abstract on the effects of several hormones on the gonadal architecture of sex-changing three-spot wrasse (*Halichoeres trimaculatus* (8)), explant culture systems for the study of sex change in fish remain unreported. Gonadal remodelling in sex-changing fishes occurs in adulthood and, therefore, the study of sex-determining mechanisms is not limited to egg or larval stages as in other vertebrates. Tissue and organoid *in vitro* culture allows evaluating multiple factors in a large number of replicates in well-controlled and standardised experimental conditions, while reducing animal usage and handling compared with *in vivo* manipulations (9–11).

The aim of this study was to develop an organ culture system for a sequential hermaphrodite to establish a new model for investigating the molecular and cellular basis of sex reversal in the gonad. We chose the New Zealand spotty wrasse (*Notolabrus celidotus*), a common and widespread marine species, to evaluate the effects of a suite of hormonal factors on gonadal architecture. In this diandric protogynous hermaphroditic species, sex change is socially regulated, whereby the removal of the dominant male from the social group induces sex change in a resident female (12). Spotty wrasses exhibit sexual dimorphism with alternative male phenotypes. Initial phase (IP) individuals consist of females and primary males which develop directly from a juvenile stage with a bipotential

gonad (13). Both females and IP males can sex or role change, respectively, to become terminal phase (TP) males (13).

While natural sex change involves modifications at the behavioural, physiological, molecular and anatomical levels (1), herein, the term 'sex change' is used to refer to the development of testicular tissue in female gonads. The effect of different doses of steroid hormones 17β-estradiol (E_2), 11-ketestosterone (11KT) or cortisol (CORT) on spotty wrasse ovary explants was evaluated *in vitro* to elucidate their effect on gonadal sex change in this species. Controls (C, no steroid) were used to determine the viability of the culture system and the potential triggering of gonadal sex change in the absence of exogenous factors. Treatment with E_2 was used to evaluate its role in the successful maintenance of spotty wrasse ovaries *in vitro*. 11KT, the most potent androgen in fish (14), was employed to explore its potential to induce sex change (i.e. proliferation of spermatogonia) in cultured spotty wrasse ovaries. In addition, treatment with the stress hormone CORT was predicted to induce oocyte degeneration through the inhibition of aromatase (15), and it was hypothesised that this effect alone may trigger the female-to-male gonadal transformation. A strong effect of CORT, the main glucocorticoid in fish, has been linked to fish sex determination and sex change in several studies (16–20). This is the first study to evaluate the effects of CORT in an *in vitro* gonadal culture system of a sex-changing fish.

Materials and Methods

Sample collection

Five IP spotty wrasse individuals ranging from 160 – 230 mm total length were captured by hook and line at the Portobello Marine Laboratory, Department of Marine Science, University of Otago, New Zealand during May 2019. Live fish were immediately transported to cell culture facilities at the Department of Zoology, University of Otago, where fish were euthanised by overdose in benzocaine (0.3 g/L) and gonads dissected within 1.5 h of capture. At this time,

gonadal fragments from each fish were preserved for histological analysis as day
0 reference tissues (D0). Gonads fixed in 4% PFA were processed for routine
embedding in paraffin (Otago Histology Services Unit, Department of Pathology,
University of Otago). Fish were captured and manipulated with approval from the
University of Otago Animal Ethics Committee, and in accordance with New
Zealand National Animal Ethics Advisory Committee guidelines.

Ovarian organ culture technique

In vitro culture of spotty wrasse ovary was carried out using the floating tissue
culture method originally described by (6) with minor modifications. For each of 5
IP fish, freshly removed ovaries were cut into pieces of approximately $2 - 3$ mm
and ovarian fragments were placed on floats of 1.5% agarose covered with a
nitrocellulose membrane in a 24-well plastic tissue culture dish. The basal
medium consisted of Leibovitz L-15 medium supplemented with 0.5% bovine
serum albumin, 10,000 U/L penicillin, 10 mg/L streptomycin and 1 µg/mL porcine
insulin adjusted to pH 7.4. Ovarian fragments were exposed to E_2, 11KT or CORT
for 21 days at 16 °C. The effect of various concentrations of each hormone (1, 10 or
100 ng/mL) was examined. Steroids were first dissolved in EtOH and then diluted
with the medium (0.1% EtOH), which was changed every 7 days. Controls (no
steroid, 0.1% EtOH) were included. At the end of the experiment, tissues were
fixed in 4% PFA and processed for routine paraffin embedding.

Image analysis

An open source pixel-based machine learning trainable classifier, Ilastik (v. 1.3.2)
(https://www.ilastik.org) (21), was employed to discriminate cell types in the
gonadal images captured using brightfield light microscopy. Approximately 85
previtellogenic oocytes (PVO), from a subset of 5 images, were used to train Ilastik.
Atretic oocytes, stromal tissue and background were labelled separately with a
paintbrush-style tool using the Pixel Classification module. The training process
was refined iteratively, updating pixel classifications until object types were

accurately discriminated in the segmentation maps, as described in (22). The
Ilastik classifier files were extracted in HDF5 (Hierarchical Data Format 5) format
and loaded into open source image processing software ImageJ (v. 2.0.0-rc-
69/1.52p), where the different object types were each assigned a different
threshold. ImageJ 'Analyse particles' function was used to calculate the total
relative surface area occupied by non-atretic PVO and measure the total surface
of the tissue section in each image. 'Remove outliers' function was used to remove
any outlier < 10 pixels from analysis and 'Fill holes' function was used to mark
PVO nuclei in the exceptional occasions when Ilastik failed to correctly recognise
these.

Statistical analysis

Generalised linear mixed models were used to assess the effect of steroid (i.e. E_2,
11KT or CORT) treatments on the proportion of total tissue surface occupied by
non-atretic PVO ('PVO_area') in comparison to the total area of each ovarian
section ('total_area'), using individual fish as a random effect. Models followed a
negative binomial distribution and were performed using the following command
line in R (v. 3.6.1): 'glmmPQL (PVO_area/total_area ~ Treatment, random =
~1 | Fish, weights = total_area, family = negative.binomial (theta = 1), data = dat)'.
If significant differences between treatments were found, Tukey's multiple
comparison tests were performed to determine where the significance lay between
treatments. All statistical analyses were performed using R (23). Results are
expressed as means ± SEM of 5 biological replicates, except for the control (n = 15)
or unless stated otherwise.

Results and Discussion

Histological analysis of D0 reference tissues confirmed that all 5 IP spotty wrasse
captured were female. Visual inspection of gonadal sections under light
microscopy revealed that oocyte degeneration was present in all experimental
groups, including controls, but not the D0 reference tissues. However, initiation of

spermatogenesis was not observed in any group, and no male structures were

identified.

Atresia of PVO appeared to be a background effect of the *in vitro* culture (8). This

was reflected in reduced oocyte surface area in controls compared to D0 reference

tissues, with the proportion of total tissue area filled with non-atretic PVO being

significantly higher in D0 (73.6 ± 2.4 %) versus control ovaries (31.8 ± 3.7 %) (p-

value < 0.001) (Figure 2A). Estrogen treatment reduced the incidence of oocyte

degeneration. A greater surface area covered by non-atretic PVO was observed in

ovaries treated with the two highest E_2 doses (10 ng/mL, 51.1 ± 5.1 %; 100 ng/mL,

50.7 ± 9.8 %) (Figure 2A). Differences between the control ovaries and those

treated with 100 ng/mL of E_2 were significant (p-value < 0.05). Therefore,

exogenous E_2 administration to cultured ovaries can effectively, albeit partially,

prevent oocyte degeneration (i.e. increased surface area occupied by non-atretic

PVO) associated with explant culture. This reinforces observations in female coho

salmon (*Oncorhynchus kisutch*), in which *in vivo* fadrozole treatment (an

aromatase inhibitor) induced reductions in E_2 and a higher incidence of associated

atresia (24). In future culture systems, the addition of growth factors or the use of

an artificial extracellular matrix (25) could be investigated to further mitigate

oocyte degeneration as a consequence of culture.

Unexpectedly, there was no histological evidence that treatment with 11KT

induced sex change in spotty wrasse ovaries *in vitro*. The proportion of non-atretic

PVO found in the ovarian tissues was not dose-dependent and did not significantly

vary between any of the 11KT-treated ovaries. Although all 11KT-treated ovary

explants showed a much lower percentage of PVO compared to total tissue area

than D0 ovaries, values were comparable to the control (Figure 2B). The

proportion of non-atretic PVO in 11KT-treated ovarian tissues was also unaffected

by dose (1 ng/mL, 24.0 ± 10.7 %; 10 ng/mL, 25.6 ± 9.1 %; 100 ng/mL 21.6 ± 9.7 %).

Successful induction of spermatogenesis via *in vitro* androgen treatment in

cultured testes has been demonstrated in Japanese eel (*Anguilla japonica*), in

which Sertoli cells were activated by 11KT to stimulate the spermatogenic cascade (3,6). The role of 11KT as a promoter of spermatogenesis in ovaries of sex-changing fish has also been demonstrated (8,26). In the protogynous honeycomb grouper (*Epinephelus merra*), *in vivo* implantation of 11KT into pre-spawning females caused masculinisation in all fish (26). Likewise, *in vitro* 11KT treatment of ovarian explants from protogynous three-spot wrasse induced the formation of presumed spermatogenic crypts and sperm within $2-3$ weeks (8). However, three-spot wrasses live in tropical waters, and sex change in captive populations has been observed in under 40 days (27), around half that required in spotty wrasse $(60-70$ days (12)). In the current study, it may be that spotty wrasse ovaries were not cultured long enough for 11KT to promote male germ cell proliferation, or that 11KT alone is insufficient to induce gonadal transformation in this species. It has been hypothesised that endogenous E_2 levels need to decrease below a physiological threshold for oocyte maintenance to become jeopardised and androgen levels to effectively induce proliferation of bipotential gonial stem cells in the ovary (26). Co-treatment of 11KT with an aromatase inhibitor (e.g. fadrozole) should be performed in the future to further characterise the triggering of *in vitro* gonadal restructuring, potentially establishing this technique as a model for sex change research.

It was hypothesised that CORT treatment would promote oocyte degeneration, following evidence that CORT may mediate sex change in fish by inhibiting aromatase transcription and promoting androgen production (14,15). However, no significant differences were observed in the proportion of non-atretic PVO between the controls and ovaries treated with 1 and 10 ng/mL of CORT (43.3 ± 7.7 % and 42.9 ± 6.2 %, respectively) (Figure 2C). While a significant increase in the percentage area of PVO was detected between the controls and 100 ng/mL CORT treatment group (39.5 ± 18.0 %) (p-value < 0.005), this may reflect variability between fish and the effect of smaller sample size (n=4, as one biological replicate was lost during histological processing due to technical issues) on the statistical analysis. Therefore, these differences were not deemed a direct effect of CORT administration to the ovaries. Cortisol has been observed to negatively impact

ovarian development by suppressing E_2 secretion in cultured rainbow trout
(*Oncorhynchus mykiss*) follicles (7). However, in that same study, Carragher &
Sumpter (7) argued that there may be a time lag for CORT suppressive effects to
become evident during *in vitro* ovarian culture. Longer-term culture of spotty
wrasse ovaries may be needed to observe any effects of CORT on ovarian structure,
a finding in keeping with our findings for 11KT.

Cortisol also failed to initiate obvious sex change in cultured spotty wrasse ovaries,
with no evidence of male tissues observed in our ovarian sections. It may be that
CORT alone is ineffective in inducing spotty wrasse sex change, or that the
experiment needs technical adjustments (e.g. longer duration) for the successful
induction of sex change in spotty wrasse. Low doses of CORT (i.e. 0.01 – 10 ng/mL)
were reported to promote testicular development in Japanese eel and pejerrey
(*Odontesthes bonariensis*) by increasing the production of 11KT, which, in turn,
induced spermatogonial proliferation (3,5). Interestingly, higher doses of CORT
(i.e. 100 ng/mL) had the opposite effect and inhibited 11KT synthesis in these
species (3,5). Although measurement of 11KT concentration in the cultured media
should be performed to evaluate such effects on spotty wrasse ovaries, absence of
spermatogonia in the cultured ovaries suggests 11KT levels were not significantly
affected. The possible metabolic conversion of CORT into 11KT and other 11-
oxygenated androgens has also been suggested in several teleosts (3,28), as there
is cross-talk between the glucocorticoid and androgen synthesis pathways (14,29).
However, the mechanisms underlying 11KT synthesis are still not fully
understood and a potential CORT-to-11KT direct conversion remains to be
clarified (14,30). This work constitutes the first-ever *in vitro* organ culture system
in a temperate sex-changing teleost, and opens doors for future work in a wide
variety of hermaphroditic fish.

Although the current experiment could not induce any observable female-to-male
restructuring of spotty wrasse gonads in culture, ovarian tissues were successfully
maintained in culture for 21 days. Such long-term culture of fish organ explants
has only been accomplished in a few instances (3–7). Moreover, this work validates

the use of open-source machine learning software Ilastik to reliably recognise cell

types in fish gonads, with enormous potential in future studies. The image

analysis strategy presented here can be adapted and applied to other studies

where accurate cell segmentation is required.

Conclusion

This study successfully developed an organ culture system for gonadal tissue

explants in New Zealand spotty wrasse, providing an *in vitro* system for

investigating control and perturbation of gonadal sex change in this and other

species. Although culture was associated with a degree of PVO degeneration, it

was shown that administration of at least 10 ng/mL E_2 can partially mitigate

tissue degeneration during culture. However, neither 11KT nor CORT treatment

successfully accelerated ovarian atresia or induced spermatogenic proliferation.

Future *in vitro* manipulations should explore longer cultures and/or co-treatment

of multiple steroids to analyse potential antagonistic or synergistic effects. This

culture system will facilitate future investigations of ovarian and testicular trans-

differentiation in sex-changing fish. For example, an *in vitro* system would be

especially amenable to the use of small interfering RNAs to knock down expression

of specific candidate genes or application of methylation inhibitors to further

investigate the genetic and epigenetic cascade underlying sex change. *In vitro*

approaches also offer valuable ethical benefits over experiments with live fish,

enabling the application of the 3Rs guiding principles: reduce, replace and refine

(31).

Abbreviations

11KT: 11-ketestosterone; C: control (no steroid); D0: day 0 reference tissues;

CORT: cortisol; E_2: 17ß-estradiol; HDF5: hierarchical data format 5; IP: initial

phase; PVO: previtellogenic oocytes; TP: terminal phase.

Acknowledgements

We thank Sam Karelitz for assistance with bioinformatic analysis and support collecting samples. We are grateful to Helen Taylor for her help with sampling, the Portobello Marine Laboratory staff for kindly allowing us to use their facilities, and members of the Lokman and Gemmell Labs for support. Sheri Johnson, Clare Holleley and Takashi Todo provided critical feedback on an earlier draft. AG was supported by a scholarship from the Department of Anatomy, University of Otago. EVT held a Health Sciences Career Postdoctoral Fellowship and Rutherford Postdoctoral Fellowship that supported this work. This work was further supported by a University of Otago Research Grant (ORG 0117-0318) to NJG and EVT, and a Marsden Grant (UOO1308) to NJG and a Research Enhancement Grant (PBRF-ML63) to PML.

Competing Interests

The authors have declared that no competing interest exists.

References

1. Todd E V, Liu H, Muncaster S, Gemmell NJ. Bending genders: the biology of natural sex change in fish. Sex Dev. 2016;10(5–6):223–41.

2. Gemmell NJ, Todd E V, Goikoetxea A, Ortega-Recalde O, Hore TA. Natural sex change in fish. In: Current Topics in Developmental Biology. 2019. p. 71–117.

3. Ozaki Y, Higuchi M, Miura C, Yamaguchi S, Tozawa Y, Miura T. Roles of 11β-Hydroxysteroid dehydrogenase in fish spermatogenesis. Endocrinology. 2006 Nov;147(11):5139–46.

4. Ozaki Y, Damsteegt EL, Setiawan AN, Miura T, Lokman PM. Expressional regulation of gonadotropin receptor genes and androgen receptor genes in the eel testis. Gen Comp Endocrinol. 2019 Sep;280:123–33.

5. Fernandino JI, Hattori RS, Kishii A, Strüssmann CA, Somoza GM. The cortisol and androgen pathways cross talk in high temperature-induced masculinization: the 11β-hydroxysteroid dehydrogenase as a key enzyme. Endocrinology. 2012

330   Dec;153(12):6003–11.

6.   Miura T, Yamauchi K, Takahashi H, Nagahama Y. Hormonal induction of all stages of spermatogenesis in vitro in the male Japanese eel (*Anguilla japonica*). Proc Natl Acad Sci USA. 1991 Jul 1;88(13):5774–8.

7.   Carragher JF, Sumpter JP. The effect of cortisol on the secretion of sex steroids from cultured ovarian follicles of rainbow trout. Gen Comp Endocrinol. 1990 Mar;77(3):403–7.

8.   Todo T, Sato M, Ashida M, Yamaguchi N, Kobayashi Y, Adachi S, et al. In vitro induction of gonadal sex change in a protogynous fish, three-spotted wrasse (*Halichoeres trimaculatus*). Cybium Int J Ichthyol. 2008;32:106.

9.   Schulz RW, de França LR, Lareyre J-J, LeGac F, Chiarini-Garcia H, Nobrega RH, et al. Spermatogenesis in fish. Gen Comp Endocrinol. 2010 Feb;165(3):390–411.

10.   Ortega-Recalde O, Goikoetxea A, Hore TA, Todd E V, Gemmell NJ. The genetics and epigenetics of sex change in fish. Annu Rev Anim Biosci. 2020 Feb 15;8(1):annurev-animal-021419-083634.

11.   Rak-Raszewska A, Hauser P V., Vainio S. Organ *in vitro* culture: what have we learned about early kidney development? Stem Cells Int. 2015;2015:1–16.

12.   Goikoetxea A. Stress and sex change in New Zealand spotty wrasse (*Notolabrus celidotus*). University of Otago; 2020.

13.   Choat JH. Sexual dimorphism in the labrid fish *Pseudolabrus celidotus* (Bloch and Schneider) 1801. Pacific Sci. 1965;XIX(19):451–7.

14.   Goikoetxea A, Todd E V, Gemmell NJ. Stress and sex: does cortisol mediate sex change in fish? Reproduction. 2017 Dec;154(6):R149–60.

15.   Fernandino JI, Hattori RS, Moreno Acosta OD, Strüssmann CA, Somoza GM. Environmental stress-induced testis differentiation: androgen as a by-product of cortisol inactivation. Gen Comp Endocrinol. 2013 Oct;192:36–44.

16.   Hayashi Y, Kobira H, Yamaguchi T, Shiraishi E, Yazawa T, Hirai T, et al. High temperature causes masculinization of genetically female medaka by elevation of cortisol. Mol Reprod Dev. 2010 May 25;77(8):679–86.

17.   Kitano T, Hayashi Y, Shiraishi E, Kamei Y. Estrogen rescues masculinization of genetically female medaka by exposure to cortisol or high temperature. Mol Reprod Dev. 2012 Oct;79(10):719–26.

18.   Yamaguchi T, Yoshinaga N, Yazawa T, Gen K, Kitano T. Cortisol is involved in temperature-dependent sex determination in the Japanese flounder. Endocrinology. 2010 Aug;151(8):3900–8.

19. Nozu R, Nakamura M. Cortisol administration induces sex change from ovary to
testis in the protogynous wrasse, *Halichoeres trimaculatus*. Sex Dev. 2015 Feb
11;9(2):118–24.
20. Miller KA, Kenter LW, Breton TS, Berlinsky DL. The effects of stress, cortisol
administration and cortisol inhibition on black sea bass (*Centropristis striata*) sex
differentiation. Comp Biochem Physiol Part A Mol Integr Physiol. 2019
Jan;227:154–60.
21. Sommer C, Straehle C, Kothe U, Hamprecht FA. Ilastik: interactive learning and
segmentation toolkit. In: 2011 IEEE International Symposium on Biomedical
Imaging: From Nano to Macro. IEEE; 2011. p. 230–3.
22. Logan DJ, Shan J, Bhatia SN, Carpenter AE. Quantifying co-cultured cell
phenotypes in high-throughput using pixel-based classification. Methods. 2016
Mar;96:6–11.
23. Core Team R. R: A language and environment for statistical computing. 2013;
24. Afonso LOB, Iwama GK, Smith J, Donaldson EM. Effects of the aromatase inhibitor
fadrozole on plasma sex steroid secretion and oocyte maturation in female coho
salmon (*Oncorhynchus kisutch*) during vitellogenesis. Fish Physiol Biochem.
1999;20(3):231–41.
25. Kim B-S, Park I-K, Hoshiba T, Jiang H-L, Choi Y-J, Akaike T, et al. Design of
artificial extracellular matrices for tissue engineering. Prog Polym Sci. 2011
Feb;36(2):238–68.
26. Bhandari RK, Alam MA, Soyano K, Nakamura M. Induction of female-to–male sex
change in the honeycomb grouper (*Epinephelus merra*) by 11-ketotestosterone
treatments. Zoolog Sci. 2006 Jan;23(1):65–9.
27. Kuwamura T, Suzuki S, Tanaka N, Ouchi E, Karino K, Nakashima Y. Sex change
of primary males in a diandric labrid *Halichoeres trimaculatus*: coexistence of
protandry and protogyny within a species. J Fish Biol. 2007 Jun;70(6):1898–906.
28. Kime DE. The hepatic catabolism of cortisol in teleost fish—adrenal origin of 11-
oxotestosterone precursors. Gen Comp Endocrinol. 1978 Jul;35(3):322–8.
29. Arterbery AS, Deitcher DL, Bass AH. Divergent expression of 11β-hydroxysteroid
dehydrogenase and 11β-hydroxylase genes between male morphs in the central
nervous system, sonic muscle and testis of a vocal fish. Gen Comp Endocrinol. 2010
May;167(1):44–50.
30. Schulz RW. *In vitro* metabolism of steroid hormones in the liver and in blood cells
of male rainbow trout (*Salmo gairdneri* Richardson). Gen Comp Endocrinol. 1986

Nov;64(2):312–9.

31.   Russell WMS, Burch RL. The principles of humane experimental technique.

Methuen; 1959.

Figure legends

Figure 1. Pixel-based trainable classifier Ilastik workflow. Example of spotty wrasse (*Notolabrus celidotus*) ovarian tissue treated with 100 ng/mL of 17β-estradiol processed for histology and visualised under light microscopy (A). Ilastik was manually trained to discriminate previtellogenic oocytes (yellow), atretic oocytes (turquoise), stromal/connective tissue (red) and background (blue) (B). Segmentation map with different object types created with Ilastik (C). Scale bar: 100 μm.

Figure 2. Histological measurement of area covered by non-atretic previtellogenic oocytes (PVO) in ovaries of spotty wrasse (*Notolabrus celidotus*) treated with E_2 (A), 11KT (B) or CORT (C) (day 21). Bars show the surface area of all non-atretic PVO over total tissue area (expressed as %). Same day 0 reference tissues and controls (day 21) were used for all treatments. Results are shown as the mean ± SEM. Letters denote a significant difference between groups. Samples sizes: D0 n = 5; C (no steroid) n = 15; E_2, 1 ng/mL n = 5, 10 ng/mL n = 5, 100 ng/mL n = 5; 11KT, 1 ng/mL n = 5, 10 ng/mL n = 5, 100 ng/mL n = 5; CORT, 1 ng/mL n = 5, 10 ng/mL n = 5, 100 ng/mL n = 4. Abbreviations: 11-ketotestosterone (11KT), control (C), cortisol (CORT), day 0 reference tissues (D0), 17β-estradiol (E_2).

Authors' biography

Alexander Goikoetxea obtained his BSc in Biotechnology and MSc in Aquaculture from the *Universitat of Barcelona* and completed his PhD at the University of Otago. His PhD thesis focuses on the elucidation of the genetic basis underlying sex change in the protogynous New Zealand spotty wrasse (*Notolabrus celidotus*), in particular, on the role of stress during this astonishing transformation.

Erin Damsteegt completed a BSc (Hons) in Zoology and a PhD at the University of Otago. After completing a Post-Doctoral research position combined between King's College London and The University of Roehampton (United Kingdom), she

436 is currently working as a Research Fellow in the Department of Zoology,
437 University of Otago, New Zealand.
438

439 Erica Todd received her PhD from James Cook University, Australia. Her Post-
440 Doctoral research at the University of Otago, New Zealand, focused on the
441 molecular basis of female-to-male sex change in wrasses using transcriptomic,
442 epigenomic and comparative genomic approaches. Currently, she is a lecturer in
443 Marine Genomics at Deakin University, Australia.
444

445 Neil Gemmell is a Professor at the University of Otago. His research broadly spans
446 reproduction and genomics, with the genetic basis of sex determination and
447 differentiation a long-standing topic of interest.
448

449 Mark Lokman is an academic at the University of Otago. His research interests
450 centre around effects of androgens on ovarian physiology and around endocrine
451 control of oogenesis in fish, chiefly using anguillid eels as research models.

452   Figure 1

A

B

C

456   Figure 2

458