Is aggression costly? Acute aggressive behavior increases oxidative stress independently of testosterone

Lucia Mentesana[1*] & Nicolas M. Adreani[1]

[1] Max Planck Institute for Ornithology, Seewiesen, Germany

* Author for correspondence (lmentesana@orn.mpg.de)

Authors' contributions

LM and NMA contributed equally. Both conceived the study, designed it, analyzed the data and wrote the manuscript.

Abstract

Aerobically demanding activities like aggression can lead to an elevated oxidative metabolism influencing the balance between pro-oxidant and antioxidant compounds and result in an overall increase in oxidative stress. Aggression may also generate oxidative stress indirectly through an increase in testosterone levels and in metabolic rate. Given that oxidative stress could represent a physiological cost of aggression we tested the hypothesis that acute aggressive behavior can increase oxidative stress and evaluated the role of testosterone as a potential mediator. To achieve this, we experimentally manipulated the aggressive behavior of wild male and female birds and measured pro-oxidants, enzymatic- and non-enzymatic antioxidants, and testosterone concentrations. After 20 minutes of aggressive behavior, both sexes had higher levels of oxidative stress levels than control individuals. This effect was independent of testosterone levels and much stronger in females. However, only in females but not in males being more aggressive comes at the expense of lower antioxidant capacity. We provide the first experimental evidence that acute aggressive behaviors can generate oxidative stress independently of testosterone in a wild vertebrate, with potential ecological and evolutionary implications given the role of oxidative stress in shaping life-history traits.

Keywords: female aggression, male aggression, oxidative stress, testosterone, rufous hornero, territorial birds.

1. Introduction

Animals frequently engage in aggressive interactions to obtain and secure limited resources that can enhance their reproductive success, like their breeding territory [1-6]. However, aggressive interactions entail costs (e.g., territory loss or injuries), which could partially explain why individuals differ in their degree of aggressiveness. A particularly relevant but understudied cost associated with this behavior concerns the oxidative status of an individual [7,8]. Oxidative status is the balance between pro-oxidants (i.e., reactive oxygen species, ROS) and antioxidants [i.e., non-enzymatic and enzymatic compounds; 9-14]. Oxidative stress, defined as a disturbance in the balance in favour of ROS, can lead to damage of biomolecules such as lipids, proteins and DNA [reviewed by 12,13]. This damage can shape life-history decisions of individuals as well as life history traits such as longevity [e.g., 10], this translating into ecological and evolutionary consequences.

Activities that elevate metabolic rate, such as aggression, can lead to elevated production of free radicals since a fixed percentage of oxygen usage in mitochondrial metabolism contributes to ROS formation [1,12,15-17; but see 18]. Thus, aggressive individuals, who might have an elevated metabolic rate compared to non-aggressive individuals, have to cope with increased costs associated with oxidative stress. Although our understanding of aggression contributing to a state of oxidative stress is still in its early stages, there is some empirical evidence for such costs in selected lines of mice (*Mus musculus domesticus*), where aggressive males had lower levels of antioxidants than non-aggressive males [7]. In contrast, aggressive males of wild-caught white skinks (*Egernia whitii)* had higher levels of non-enzymatic antioxidants than less aggressive males, whereas in females no such relationship was apparent [8]. In addition, aggressive behaviors such as territorial fights could incur an oxidative challenge as a consequence of the behavior's aerobic demand. Yet, whether this is the case or not remains unknown.

Aggression can also increase oxidative stress indirectly through an increase in testosterone levels [19,20]. Testosterone is assumed to be the key hormone related to resource-defence aggression and can increase during aggressive interactions in males [21-23]. Testosterone has not only been postulated to have intrinsic oxidative properties [e.g., 24-26] but it can also enhance the metabolic rate of an individual [e.g., 27-29]. Therefore, an increase in aggressive behavior after a territorial intrusion can potentially lead to an increase in testosterone levels [e.g., 30-32], with testosterone levels influencing the balance between pro-oxidant and antioxidants compounds and resulting in an overall increase in oxidative stress. To date, only one study has looked at the relationship between aggression, oxidative status and testosterone levels in a wild-caught vertebrate [8]: the aggressive phenotype of male white skinks, but not testosterone levels, was positively related to the oxidative condition of individuals. Thus, these findings do not support the hypothesis of testosterone affecting the

oxidative status of an individual during aggressive interactions. However, experimental
studies directly testing the causal relationship of these patterns are needed.

The main goal of our study was to test the hypothesis that an acute increase in
aggressive behavior could impair the oxidative status of wild rufous horneros (Aves:
Furnarius rufus, hereafter termed hornero) and to test whether this was related to testosterone
or not. To elicit an aggressive response, we challenged male and female birds with 20 minutes
of simulated territorial intrusions (STI) during the nest-building period. Immediately after the
STI we collected blood samples to measure three oxidative status markers and testosterone
concentrations. We then compared these parameters with those of birds that were not exposed
to acute aggressive behaviors. The hornero represents an excellent model system to study the
effects of acute aggression across sexes. They are seasonal breeders and both sexes are
involved in territorial defense throughout the year, with males being more aggressive than
females [33,34]. Further, horneros are sexually monomorphic in both body size [i.e., there is
no difference in body condition between sexes; 34] and plumage, and all breeding behaviors
studied so far, except for aggressive interactions, are shared and coordinated between the
sexes [35,36].

2. Materials and methods

83
2.1 Field site and simulated territorial intrusion (STI)
The study was conducted between August 22nd and September 28th of 2016 on the campus
of INIA 'Las Brujas' (National Institute of Agricultural Research), department of Canelones,
Uruguay (34°40' S, 56°20' W; 0-35 m a.s.l.). During this period horneros were building their
nests and females were close to egg laying. In total, we collected behavioral and physiological
data from birds defending 51 territories that were subjected to the experimental treatments
between 7 am - 1 pm.

We experimentally manipulated hornero aggressive behavior by performing
simulated territorial intrusions (STI). STIs are a common method to investigate the aggressive
and physiological responses of a territory holder in response to an intruder [e.g., 37-40], and
STIs are known to be effective in horneros [41]. In particular, a stuffed decoy of a male
hornero, a speaker and folded mist nets were placed 5 m away from the nest of focal pairs and
solo songs and duets of conspecifics were played (recordings were extracted from the
database www.xeno-canto.org and the amplitude of every recording was normalized). For
ethical reasons we could not use more than one decoy (i.e., the visual component of the STIs
was the same for every territory). However, horneros are sexually indistinguishable in
morphological traits (i.e., mass, tarsus and bill) and plumage coloration [i.e., assessed in
relation to the avian visual system; 34]. Further, in a comparable STI study done in European

great tits (*Parus major*) the decoys used (n=15) did not explain any significant variation in aggressive behavior [42]; thus, suggesting that the use of only one decoy did not bias our results. On the other hand, given the lack of information on the function of the different vocal signals in horneros, we decided to form a pool of ten male solo songs and ten duets to make sure that both males and females responded to the STI. We also added ten 'silence' files of 7-15 sec duration to have unpredictable gaps between the different stimuli. The auditory stimuli for each STI were randomly selected from this pool in order to avoid pseudo-replication of the acoustic component across territories [e.g. 39]. Horneros are suboscines and do not learn their vocalizations [43]. Thus, compared to oscines, the acoustic variability of songs and duets across individuals is low [43]. Thus, even though the auditory stimuli were randomly chosen from the pool, it is likely that all playbacks elicited a comparable behavioral response across territories. For each territory, the treatment (i.e., control or STI) was randomly determined, and both members of the couple were subjected to same treatment.

Challenged birds (hereafter 'STI birds', N_{males}, $N_{females}$ = 18, 9) were exposed to an STI for 20 minutes. During this time, two observers recorded the aggressive behavior of the territorial couple in digital voice recorders (Philips VoiceTraicer DVT1200 and Olympus Digital Recorder VN-733 PC). Each observer followed one member of the focal STI pair, which was decided randomly before the playback begun (to avoid that one observer always recorded the behavior of one specific sex). Observers did not differ in their quantification skills for the behaviors scored (Figure S1 & Table S1; see Supplementary Material section for a detailed description of the inter-observer reliability test). The following parameters were recorded: 1) response latency (time between start of playback and first approach), 2) time spent within 5 m of the decoy, 3) time spent on the nest, 4) number of solo songs, 5) number of duets, and 6) number of flights over the decoy, which represents both attacks and attempted attacks towards the decoy. After 20 minutes, mist nets were unfolded for ten additional minutes or until capturing both individuals of the pair (mean capture time ± sd = 5.94 ± 4.68 mins for males and 3.11 ± 2.33 mins for females). During a duet, the contributions of each sex are acoustically distinct from each other [44]. Thus, sex identification in the field was done based on the singing behavior and was later confirmed by molecular sexing [for details see 41]. Control individuals (N_{males}, $N_{females}$ = 24, 7) were attracted with playback and presentation of the decoy, but because mist nets were open from the beginning of the playback, control birds were caught as soon as they approached the experimental set-up (i.e., they did not perform aggressive behaviors; mean capture time ± sd = 4.50 ± 3.75 mins for males and 4.57 ± 2.97 mins for females). From all captured birds 80 µl of blood were taken from the wing vein using heparinized capillaries (mean sampling time ± sd = 4.75 ± 1.48 mins). In addition, morphometric measurements (i.e. mass, tarsus length and wing length) were recorded. Control and STI individuals did not differ in body condition index (Supplementary materials

Table S2). Before releasing them back in their territories, each individual was marked with a

numbered aluminum ring and a unique combination of three colored plastic split rings.

Both sexes showed clear behavioural responses towards our treatment. Both control

and STIs birds flew immediately towards the stimulus and approached the decoy (N.M.

Adreani, personal observations). Further, the simulated territorial intrusions elicited similar

behavioural responses as intrusions by real conspecifics or territorial conflicts with

neighbours [36,41]. Finally, it is important to note that although both sexes responded to the

territorial intrusion, once the male was captured the female would immediately quit the

territorial behavior and start alarming. This is the main reason why we caught more than

twice the number of males compared to females during our experiments.

2.3 Analysis of blood oxidative status parameters

The redox state of the captured individuals was determined from both non-enzymatic and

enzymatic antioxidants present in plasma and red blood cells respectively, and from reactive

oxygen species present in plasma. Antioxidant plasma concentrations (i.e., OXY) were

measured by the OXY-Adsorbent test (Diacron International SRL, Grosseto, Italy) following

the protocol described by Costantini *et al.* [45]. Measurements are expressed as mm of HOCl

neutralized. The activity of glutathione peroxidase (GPX) present in red blood cells was

determined using the Ransel assay (Randox Laboratories, Germany) following Costantini *et*

al. [46]. Measurements are expressed as U/ml of red blood cells. Oxidative damage was

quantified by measuring the levels of reactive oxygen metabolites (dROMs) with the d-ROM

test (Diacron International SRL, Grosseto, Italy) following the protocol described by

Costantini *et al.* [45]. Reactive oxygen metabolites are expressed as mm of H_2O_2 equivalents.

The absorbance was read using a spectrophotometer (Thermo Scientific Multiskan GO) at a

wavelength of 546 nm for OXY and dROMs (endpoint mode) and 340 nm for GPX (kinetic

modality).

2.4 Analysis of blood testosterone levels

Testosterone was extracted from plasma and concentrations were determined by

radioimmunoassay following Goymann *et al.* [47]. Mean recovery following extraction of the

samples was 89 ± 2% (mean ± sd). The lower detection limit of the hormone assay was at

0.39 pg/tube. All samples used in our analyses were above these detection limits and the intra-

assay variation was 7.5%. For males, we followed the same procedure as the one previously

described. Testosterone data and analyses are already published [for details see 41]. It is

important to note that although males and females were analyzed in two separate assays -

because the samples collected for males were also part of a complementary study - mean

recoveries were similar in both assays (mean recovery ± sd: for males = 87 ± 1.9 % and for

females = 89±2%). It is therefore unlikely that differences in testosterone levels obtained for

males and females are due to samples measured in separate assays.

2.5 Statistical analysis

All our analyses were performed using the R-packages "lme4" and "arm" in R-3.3.3 [48] in a

Bayesian framework with non-informative priors. For all models, we assumed a Gaussian

error distribution, and model fit was assessed by visual inspection of the residuals. We used

the "*sim*" function to simulate posterior distributions of the model parameters. Based on

10,000 simulations, we extracted the mean value and 95% credible intervals (CrI) [49].

Assessment of statistical support was obtained from the posterior distribution of each

parameter [50]. We considered an effect to be statistically meaningful when the posterior

probability of the mean difference (hereafter termed '*p(dif.)*') between compared estimates

was higher than 95% or when the estimated CrI did not include zero. For details on this

approach see Korner-Nievergelt *et al.* [50].

First, we ran four linear models fitting OXY, GPX, dROMs, and testosterone as response

variables to i) compare the oxidative status and testosterone levels between male and female

control birds (i.e., individuals that better represent the baseline status), and to ii) study the

effect of 20 minutes of aggressive behavior on the oxidative status and testosterone levels of

male and female horneros. To do so, sex (female vs male), and treatment (control vs STI)

were included as fixed effects, as well as the interaction between both factors. Although from

Adreani *et al.* [41] we know that there is no difference between control and STI birds in

capture time and handling time we included them as covariates to account for the variation

that they might add. Preliminary exploration of our data showed that time of the day as well

as observer ID did not explain significant variance, thus they were not included in the final

model.

Second, we ran three linear models to investigate the relationship between oxidative

status and testosterone levels in control individuals. OXY, GPX, and dROMs were fitted as

response variable, and in each model, testosterone was fitted as a covariate. Sex and the

interaction of sex with testosterone levels were included as fixed factors. Since males and

females differ in their testosterone concentrations (See Figure 1), for each sex we normalized

testosterone levels between 0 and 1. This allowed us to compare results between sexes.

Finally, within the STI birds, we studied the effect that aggressive behavior had on the

oxidative status and testosterone levels. To do so, we fitted OXY, GPX, dROMs and

testosterone concentrations as the response variable in a linear model. While GPX and

dROMS were square-root transformed for a better fit of each model, testosterone was log-

transformed for the same purpose. Although we measured 6 behavioral responses as proxies

of aggressiveness (detailed above), to avoid multiple testing, here we only used "flights over

the decoy" as a covariate in the model since this is the best proxy for aggression during a
territorial intrusion in horneros [41, Adreani *et al.* under review). In addition, since males and
females differ in their aggressive behaviors [i.e., males are more aggressive than females; 36],
for each sex we normalized the behavioral variables between 0 and 1 in order to capture the
aggression gradient in each sex and to be able to compare the effects between the sexes in one
model. Therefore, we tested the effect of aggression on the different physiological parameters
by including "normalized flights over the decoy" as a covariate in the model. Sex, together
with the interaction between aggressive behavior and sex, were fitted as fixed factors. Given
that we had a limited sample size and that neither capture nor handling time had an effect on
any of the oxidative parameters or testosterone levels (see Table S3), these variables were not
included in the final models.

3. Results

3.1 Sex differences in oxidative status and testosterone levels of control birds
Control males had a higher non-enzymatic antioxidant capacity (OXY) than control females
(Fig. 1A; *p(dif.)* > 99.99%). Both sexes had similar enzymatic antioxidant concentrations
(GPX; Fig. 1B; *p(dif.)* = 79.03%). We found a trend for control males having lower levels of
pro-oxidants than females (dROMs; Fig. 1C; *p(dif.)* = 94.67%). Control males had higher
testosterone levels than control females (Fig. 1D; *p(dif.)* = 99.86%). For details on model
estimates see Table S3. There was no relationship between any oxidative marker and
testosterone level in either sex (Fig. S2, see Table S4 for the model estimates).

3.2 Effect of acute aggressive behavior on oxidative status and testosterone levels
Both sexes had lower levels of OXY after 20 minutes of simulated territorial intrusion (Fig.
1A; $p(dif._{females}$ > 99.99%; $p(dif.)_{males}$ > 99.99%) compared to control individuals. However,
the treatment affected males and females differently: the difference in OXY concentrations
between control and STI females was much stronger than that in males (females' OXY
differed by 39.62% between treatments, while in males this difference was 6.18%; *p(dif.)* =
99.50%). No difference in GPX ($p(dif.)_{females}$ = 76.62%; $p(dif.)_{males}$ = 44.85%), dROMs
($p(dif.)_{females}$ = 31.35%; $p(dif.)_{males}$ = 44.14%) or testosterone ($p(dif.)_{females}$ = 45.56%; $p(dif.)_{males}$
= 79.65%) concentration was observed between groups (Fig. 1 B-D). For details on model
estimates see Table S3.

Fig. 1. Concentrations of testosterone and three oxidative markers of female (♀) and male (♂) horneros that were exposed to simulated territorial aggression (STI) and controls (CNT). (A) Non-enzymatic antioxidant concentrations (OXY); (B) enzymatic antioxidant concentrations (GPX); (C) pro-oxidant concentrations (dROMs) and (D) testosterone levels. Grey circles and triangles show the individual data for females and males respectively. Mean estimates of the model (black symbols) and 95% credible intervals (colored vertical bars) are also shown. Statistically meaningful differences are marked with an asterisk. Testosterone data of males were already published by Adreani *et al.* [41].

3.3 Relationship between aggression and oxidative status

In males, aggressive behavior and OXY concentrations were unrelated (Figure 2A; posterior probability for the slope being < 0: 52.42%), but more aggressive females had lower concentrations of antioxidants than less aggressive females (Fig. 2A; posterior probability for the slope being < 0: 99.64%). Aggression levels were unrelated to testosterone levels (Fig. 3B), GPX or dROMs in either sex (the posterior probability for the slope being < 0 was smaller than 90 % for each). For details in these models' estimates see Table S5.

Fig. 2. Relationship between aggressive behavior and non-enzymatic antioxidant concentration (OXY) and testosterone in female (♀; circles; red line) and male (♂; triangles; blue line) horneros. (A) More aggressive females had lower OXY concentrations than less aggressive females; no relationship was found in males. (B) No relationship between aggression and testosterone in male or female birds. Triangles and circles show the individual data for males (♂) and females (♀) respectively. Mean estimates of the model (solid lines) and 95% credible intervals (shaded area) are also shown. Testosterone data from the males were already published by Adreani *et al.* [41].

4. Discussion

We experimentally tested whether aggressive behavior can induce a change in oxidative stress and whether this effect is mediated by testosterone in a wild bird. The results provide the first evidence that an acute increase in aggressive behavior (i.e., for 20 mins) elicits an increase in oxidative stress in males and females independently of testosterone concentrations. This effect of STI was much stronger in females than males. Moreover, being more aggressive during the nest-building period was related to higher levels of oxidative stress in females, but not in males.

4.1. Sex differences in the oxidative status and testosterone concentrations in control birds

Control females had higher levels of oxidative stress (i.e., lower concentrations of non-enzymatic antioxidants and a trend for higher concentrations of reactive oxygen species) than control males. In both sexes, we found no relationship between testosterone and any of the markers of oxidative stress measured. These results were unexpected given that female horneros are less aggressive [36], have lower levels of testosterone and have similar body

condition compared to males (Supplementary Materials Table S2). Also, estrogens, which are

generally present at higher concentrations in females close to egg laying compared to males,

can increase antioxidant defenses [51]. One possible explanation for the sex difference in

oxidative status might be that males and females face different reproductive challenges during

the early reproductive period, when females are about to lay eggs. In birds, the resting

metabolic rate can increase by up to 27% during the period of egg production [52]. Such an

increase in metabolic rate could potentially explain the lower levels of non-enzymatic

antioxidant protection (OXY) and their tendency to have higher levels of reactive oxygen

species ($p(dif.)$ = 94.75%; Fig. 1C) compared to males (Fig. 1A). The egg laying period is

also physiologically demanding for females since some components mothers transmit into

their eggs are limited in their availability [53-55]. Antioxidants, such as vitamin E and

carotenoids, cannot be synthesized *de novo* [e.g., 56] and so are of dietary origin. Therefore,

another possible and non-exclusive explanation for the difference in the oxidative status

between sexes, could be associated with females allocating non-enzymatic antioxidant

components into the egg yolk [57,58] to protect the developing embryo [55].

4.2. Effect of aggressive behavior on oxidative status

An increase in aggressive behavior and in associated activity patterns can lead to an increase

of free radicals as a consequence of the aerobic demand [1,15,16]. However, there was no

difference in pro-oxidant concentrations (dROMs) between control and STI individuals in

either sex, while STI individuals had lower levels of OXY. Further, the concentration of

enzymatic antioxidants (i.e., GPX) was not different between control and STI birds. The latter

could be explained because GPX plays an important role in cellular detoxification of dROMs

[59,60], which did not change after the STI, or by the fact that an up- or down- regulation of

enzymatic defenses takes more than 20 minutes. On the other hand, aggressive behaviors can

increase oxidative stress indirectly through an increase in testosterone levels [8,19,20].

However, STI horneros had similar testosterone levels compared to control birds indicating

that the difference in oxidative status observed between experimental groups is not mediated

by this steroid hormone. In line with our results, the relationship between aggressive

phenotypes and oxidative status was also independent of plasma testosterone levels in both

male and female wild-caught lizards [8].

The effect that acute aggressive behaviors had on OXY concentrations was much

stronger in females than in males (i.e., STI females had 39.62% lower OXY concentrations

than control females while STI males had only 6.18% lower OXY concentrations than

controls). This sex difference in sensitivity could be given by a differential re-allocation of

antioxidants to: protect gametes [61], the developing embryo [see above; 19,56,62], and/or

those tissues with the greatest oxygen consumption [e.g., muscles; 59]. Sex differences could

also be explained by other physiological pathways besides an increase in activity levels (see below for further discussion). The implications of these sex differences will depend on the life-history stage that the individuals are in [63]. Repeated aggressive interactions experienced by females during egg laying may come at a cost to the deposition of substances into the egg that could affect her fitness. For example, an exercised group of female zebra finches (*Taeniopygia guttata*) that had higher oxidative stress than non-exercised birds after flight training, decreased the deposition of antioxidants (i.e., lutein) into the egg yolk [64]. However, birds can also recover their circulating non-enzymatic antioxidant capacity by consuming antioxidant-rich foods [e.g., 14] to avoid negative fitness consequences. Thus, to understand if acute aggressive interactions have long-lasting and sex-specific effects, future studies should ideally measure the oxidative condition of an individual not only immediately after facing aggressive interactions but also over repeated interactions, in relation to fitness traits and across different life-history stages.

To the best of our knowledge this is the first study that demonstrates an effect of an acute increase in aggressive behavior (i.e., 20 minutes) on the oxidative status of wild animals. To date, only two other studies investigated the relationship between aggressive behavior and oxidative status by looking at the correlation between the personality trait "aggression" (i.e., over a long-term scale) and the redox status of a selected line of mice [7] and white skinks [8]. Interestingly, these two studies support our findings: they also found a relationship between aggression and OXY concentration, but not with dROMs as previously suggested. Altogether, this suggests that aggression has both short- and long-term consequences on the oxidative status of an individual, and that this occurs through a change in non-enzymatic antioxidants. On a broader scale, these studies show that the relationship between aggression and oxidative status is present across different vertebrate groups (i.e., reptiles, mammals and birds). The oxidative status of an individual might therefore be a key physiological mechanism underlying different life-history strategies (e.g., longevity) in individuals that show different levels of aggression.

4.3. Relationship between aggression and oxidative status

Whereas more aggressive females had lower levels of OXY than less aggressive females, no such relationship was found in males. There was also no relationship between testosterone levels and aggressive behavior in either sex. Together with the previous results, this indicates that mechanisms other than testosterone determine the levels of oxidative stress observed in males and females after aggressive interactions. Aggression is often regulated by other hormones, such as progesterone [e.g., 47] or glucocorticoids [e.g., 23,66]. Progesterone has indeed been linked with the generation of oxidative stress in studies done *in vitro* [e.g., 25,67]. To test for the potential effect that this hormone might have on the oxidative status of

control and STI horneros in this study we measured, besides testosterone, plasma
progesterone levels (see Supplementary Materials). In horneros, progesterone (as
testosterone) seems to be decoupled from aggression-induced oxidative stress (Sup. Table 6-
8, Supplementary Material). We did not include these results in the main text given the lack
of clear hypotheses about the relationship between progesterone and oxidative status in living
organisms.

Engaging in aggressive interactions during the nest-building period could also have
led to an increase in glucocorticoid hormones (i.e., through an activation of the hypothalamic-
pituitary-adrenal axis) to help an individual recover and survive an immediate challenge [68].
An increase of glucocorticoids in the blood stream after aggressive interactions has been
reported both in seasonally territorial [e.g., 69] and year-round territorial vertebrates [70,71].
Further, a direct link between glucocorticoids and oxidative stress has previously been
reported [72-75], and females seem to be more susceptible to the effects of glucocorticoids on
oxidative stress than males [reviewed by 73, but see 76]. Thus, the interaction between
glucocorticoids and oxidative status could provide an alternative explanation of our results
that we cannot resolve at the moment: aggression may lower OXY concentrations directly as
a consequence of an increased metabolism or indirectly via an increase in glucocorticoid
concentration, or both.

5. Conclusions
Until now, the effect of aggression on oxidative stress was investigated in the framework of
personality traits [7,8]. Our study demonstrates, for the first time, that acute increases in
aggression can increase oxidative stress, and suggests that the mechanisms mediating
oxidative stress during agonistic encounters differ between the sexes. Altogether, the three
studies suggest that aggressive behaviors performed over both short- and long-term have
direct consequences on the redox balance of an individual. Remarkably, by generating a
change in non-enzymatic antioxidants and not in the concentration of pro-oxidants as
previously suggested. Given that oxidative stress can shape central life history traits such as
longevity, the oxidative status of an individual might be a key physiological mechanism
underlying different life-history strategies in individuals that show different levels of
aggression.

Ethical statement
The experimental procedures were approved by the Ethics Committee of Animal
Experimentation (CEUA) of the Facultad de Ciencias, Universidad de la República, Uruguay
(University of the Republic of Uruguay). Protocol number 186, file 2400-11000090-16.

Competing interests

We declare no competing interests.

Funding

We acknowledge funding and support from the International Max Planck Research School (IMPRS) for Organismal Biology, and by Idea Wild that provided field equipment.

Acknowledgements

We thank Enzo Cavalli and Ernesto Guedes for their assistance in the field, and Monika Trappschuh for conducting the hormone analysis. We also thank the members from the Ethology Lab at the Universidad de la República in Montevideo and especially to Bettina Tassino for their support. Also, we thank all the "INIA Las Brujas" staff for providing us with accommodation and equipment during fieldwork. Furthermore, we are thankful to Juan Carlos Reboreda and the Ecology and Behaviour lab (LEyCA) from the University of Buenos Aires for the logistic support. We also thank Michaela Hau and Manfred Gahr for their valuable support. We thank Laurie O'Neil and Glenn Cockburn for helping us with the English proof of the manuscript. We are grateful to Stefania Casagrande, Michaela Hau, Wolfgang Goymann and Scott McWilliams for valuable discussions about the study and constructive criticism on previous versions of the manuscript.

References

1. Beckman, K. B., & Ames, B. N. (1998). The Free Radical Theory of Aging Matures. Physiological Reviews, 78(2).
2. Clutton-Brock, T. (2009). Sexual selection in females. Animal Behaviour, 77(1), 3–11.
3. Demas, G. E.., Cooper, M. A., Albers, H.E., & Soma, K.K. (2007). Novel mechanisms underlying neuroendocrine regulation of aggression: a synthesis of rodent, avian, and primate studies.
4. Duckworth, R. A. (2006). Aggressive behaviour affects selection on morphology by influencing settlement patterns in a passerine bird. Proceedings of the Royal Society of London B: Biological Sciences, 273(1595).
5. Rosvall, K. A. (2008). Sexual selection on aggressiveness in females: evidence from an experimental test with tree swallows. Animal Behaviour, 75(5), 1603–1610.
6. Smith, B. R., & Blumstein, D. T. (2008). Fitness consequences of personality: a meta-analysis. Behavioral Ecology, 19(2), 448–455.
7. Costantini, D., Carere, C., Caramaschi, D., & Koolhaas, J. M. (2008). Aggressive and non-aggressive personalities differ in oxidative status in selected lines of mice (*Mus musculus*). Biology Letters, 4(1).
8. Isaksson, C., While, G. M., McEvoy, J., van de Crommenacker, J., Olsson, M., Groothuis, T. G.G., Komdeur, J., & Wapstra, E. (2011). Aggression, but not testosterone, is associated to oxidative status in a free-living vertebrate. Behaviour, 148(5), 713–731.
9. Sies, H. (1991). Oxidative stress: From basic research to clinical application. The American Journal of Medicine, 91(3), S31–S38.
10. Finkel, T., & Holbrook, N. J. (2000). Oxidants, oxidative stress and the biology of ageing. Nature, 408(6809), 239–247.
11. Halliwell, B., & Gutteridge, J. M. C. (2007). Free radicals in biology and medicine.
12. Costantini, D. (2008). Oxidative stress in ecology and evolution: lessons from avian studies. Ecology Letters, 11(11), 1238–1251.
13. Monaghan, P., Metcalfe, N. B., & Torres, R. (2009). Oxidative stress as a mediator of life history trade-offs: mechanisms, measurements and interpretation. Ecology Letters, 12(1), 75–92.
14. Skrip, M. M., & McWilliams, S. R. (2016). Oxidative balance in birds: an atoms-to-organisms-to-ecology primer for ornithologists. Journal of Field Ornithology, 87(1), 1–20.
15. Leeuwenburgh, C., & Heinecke, J. (2001). Oxidative Stress and Antioxidants in Exercise. Current Medicinal Chemistry, 8(7), 829–838.

16. Speakman, J. R. (2003). Oxidative phosphorylation, mitochondrial proton cycling, free-
radical production and aging. 14, 35–68.

17. Powers, S. K., & Jackson, M. J. (2008). Exercise-Induced Oxidative Stress: Cellular
Mechanisms and Impact on Muscle Force Production. Physiological Reviews, 88(4).

18. Salin, K., Auer, S. K., Rudolf, A. M., Anderson, G. J., Cairns, A. G., Mullen, W., Hartley,
R. C., Selman, C., & Metcalfe, N. B. (2015). Individuals with higher metabolic rates have
lower levels of reactive oxygen species in vivo. Biology Letters, 11:20150538.

19. Schantz, T. von, Bensch, S., Grahn, M., Hasselquist, D., & Wittzell, H. (1999). Good
genes, oxidative stress and condition–dependent sexual signals. Proceedings of the Royal
Society of London B: Biological Sciences, 266(1414).

20. Alonso-Alvarez, C., Bertrand, S., Faivre, B., Chastel, O., & Sorci, G. (2007).
Testosterone and oxidative stress: the oxidation handicap hypothesis. Proceedings of the
Royal Society of London B: Biological Sciences, 274(1611).

21. Goymann, W., Landys, M. M., & Wingfield, J. C. (2007). Distinguishing seasonal
androgen responses from male-male androgen responsiveness-revisiting the Challenge
Hypothesis. Hormones and Behavior, 51(4), 463–476.

22. Hirschenhauser, K., & Oliveira, R. F. (2006). Social modulation of androgens in male
vertebrates: meta-analyses of the challenge hypothesis. Animal Behaviour, 71(2), 265–
277.

23. Wingfield, J. C., Hegner, R. E., Dufty, A. M., & Ball, G. F. (1990). The "Challenge
Hypothesis": Theoretical Implications for Patterns of Testosterone Secretion, Mating
Systems, and Breeding Strategies. The American Naturalist, 136(6), 829–846.

24. Chainy, G. B. N., Samantaray, S., & Samanta, L. (2009). Testosterone-induced changes
in testicula antioxidant system. Andrologia, 29, 343–9.

25. Zhu, X. D., Bonet, B., Knopp, R. H. (1997). 17β-estradiol, progesterone, and testosterone
inversely modulate low- density lipoprotein oxidation and cytotoxicity in cultured
placental trophoblast and macrophages. American Journal of Obstetrics and Gynecology,
177, 196–209.

26. Casagrande S., Dijkstra C., Tagliavini J., Goerlich V. C.& Groothuis T. G. G. G. (2011).
Differential effects of testosterone, dihydrotestosterone and estradiol on carotenoid
deposition in an avian sexually selected signal. Journal of Comparative Physiology. 197,
1–13.

27. Marler C. A. & Moore M (1989). Time and Energy Costs of Aggression in Testosterone-
Implanted Free-Living Male Mountain Spiny Lizards (*Sceloporus jarrovi*). Physiological
Zoology, 62(6), 1334-1350.

28. Welle S., Jozefowicz R., Forbes G, Griggs R. C. (1992). Effect of testosterone on
metabolic rate and body composition in normal men and men with muscular dystrophy,
The Journal of Clinical Endocrinology & Metabolism, 74(2), 332–335,

29. Wikelski, M., Lynn, S., Breuner, J. C., Wingfield, J. C., & Kenagy, G. J. (1999). Energy
metabolism, testosterone and corticosterone in white-crowned sparrows. Journal of
Comparative Physiology A: Sensory, Neural, and Behavioral Physiology, 185(5), 463–
470.

30. Wingfield, J., & Wada, M. (1989). Changes in plasma levels of testosterone during male-
male interactions in the song sparrow, *Melospiza melodia*: time course and specificity of
response. Journal of Comparative Physiology A, 166(2), 189–194.

31. Wingfield, J. C., & Hahn, T. P. (1994). Testosterone and territorial behaviour in sedentary
and migratory sparrows. Animal Behaviour, 47(1), 77–89.

32. McGlothlin, J. W., Jawor, J. M., Greives, T. J., Casto, J. M., Phillips, J. L., & Ketterson,
E. D. (2008). Hormones and honest signals: males with larger ornaments elevate
testosterone more when challenged. Journal of Evolutionary Biology, 21(1), 39–48.

33. Fraga, R. M. (1980). The breeding of Rufous Horneros (*Furnarius rufus*). The Condor,
82(1), 58–68.

34. Diniz, P., Ribeiro, P. H. L., Rech, G. S., & Macedo, R. H. (2016). Monochromatism,
cryptic sexual dimorphism and lack of assortative mating in the Rufous Hornero,
Furnarius rufus albogularis. Emu - Austral Ornithology, 116(3), 294–300.

35. Massoni, V., Reboreda, J. C., López, G. C., & Aldatz, M. F. (2012). High coordination
and equitable parental effort in the Rufous Hornero. The Condor, 114(3), 564–570.

36. Diniz, P., da Silva Júnior, E. F., Webster, M. S., & Macedo, R. H. (2018). Duetting
behavior in a Neotropical ovenbird: sexual and seasonal variation and adaptive signaling
functions. Journal of Avian Biology.

37. Wingfield, J. C. (1994). Regulation of Territorial Behavior in the Sedentary Song
Sparrow, *Melospiza melodia morphna*. Hormones and Behavior, 28(1), 1–15.

38. Gill, S. A., Alfson, E. D., & Hau, M. (2007). "Context matters: female aggression and
testosterone in a year-round territorial neotropical songbird (*Thryothorus leucotis*)."
Proceedings of the Royal Society B: Biological Sciences 274(1622): 2187-2194.

39. Apfelbeck, B., & Goymann, W. (2011). Ignoring the challenge? Male black redstarts
(*Phoenicurus ochruros*) do not increase testosterone levels during territorial conflicts but
they do so in response to gonadotropin-releasing hormone. Proceedings. Biological
Sciences, 278(1722), 3233–3242.

40. Villavicencio, C. P., Blas, J., & Goymann, W. (2014). The number of life-history stages
does not influence the androgen responsiveness to male–male interactions: Sedentary and
migratory black redstarts (*Phoenicurus ochruros*) do not elevate testosterone in response

to simulated territorial intrusions. General and Comparative Endocrinology, 205, 159–
165.

41. Adreani, N.M., Goymann, W. & Mentesana, L. (2018) Not one hormone or another:
Aggression differentially affects progesterone and testosterone in a South American
ovenbird. Hormones and Behavior, 105, 104-109.

42. Araya-Ajoy, Y.G. & Dingemanse, N.J. (2013) Characterizing behavioural "characters":
an
evolutionary framework. Proceedings of the Royal Society of London B: Biological
Sciences, 281, 20132645–20132645.

43. Freeman, B. G., Montgomery, G. A., & Schluter, D. (2017). Evolution and plasticity:
Divergence of song discrimination is faster in birds with innate song than in song learners
in Neotropical passerine birds. Evolution, 71(9), 2230-2242.

44. Laje, R., & Mindlin, G. B. (2003). Highly structured duets in the song of the South
American hornero. Physical review letters, 91(25), 258104.

45. Costantini, D., Casagrande, S., De Filippis, S., Brambilla, G., Fanfani, A., Tagliavini, J.,
& Dell'Omo, G. (2006). Correlates of oxidative stress in wild kestrel nestlings (*Falco*
tinnunculus). Journal of Comparative Physiology B, 176(4), 329–337.

46. Costantini, D., Monaghan, P., & Metcalfe, N. B. (2011). Biochemical integration of blood
redox state in captive zebra finches (*Taeniopygia guttata*). The Journal of Experimental
Biology, 214(7), 1148–1152.

47. Goymann, W., Wittenzellner, A., Schwabl, I., & Makomba, M. (2008). Progesterone
modulates aggression in sex-role reversed female African black coucals. Proceedings of
the Royal Society of London B: Biological Sciences, 275(1638).

48. R Core Team 2013. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.
in press.

49. Gelman, A., & Hill, J. (2007). Data analysis using regression and multilevel/hierarchical
models.

50. Korner-Nievergelt, F., Roth, T., von Felten, S., Guélat, J., Almasi, B., & Korner-
Nievergelt, P. (2015). Bayesian data analysis in ecology using linear models with R,
BUGS, and Stan.

51. Halifeoglu, I., Karatas, F., Canatan, H., Colak, R., & Karadas, E. (2003). Investigation of
antioxidant vitamins (A, E and C) and selenium levels in chickens receiving estrogen or
testosterone. Cell Biochemistry and Function, 21(2), 133–136.

52. Nilsson, J.-Å., & Råberg, L. (2001). The resting metabolic cost of egg laying and nestling
feeding in great tits. Oecologia, 128(2), 187–192.

53. Hulbert, A. J., & Abbott, S. K. (2011). Nutritional ecology of essential fatty acids: an
evolutionary perspective. Australian Journal of Zoology, 59(6), 369.

54. Møller, A. P., Biard, C., Blount, J. D., Houston, D. C., Ninni, P., Saino, N., & Surai, P.
(2000). Carotenoid-dependent signals: indicators of foraging efficiency,
immunocompetence or detoxification ability? Avian Poultry Biol. Rev., 11, 137–159.

55. Surai, P. F. (2002). Natural antioxidants in avian nutrition and reproduction. British
Library Cataloguing in Publication Data Poultry Environment Problems: A Guide to
Solutions.

56. Surai, P. F., Speake, B. K., & Sparks, N. H. C. (2001). Carotenoids in avian nutrition and
embryonic development. 1. Absorption, availability and levels in plasma and egg yolk.
The Journal of Poultry Science, 38(1), 1–27.

57. Royle, N. J., Surai, P. F., Mccartney, R. J., & Speake, B. K. (1999). Parental investment
and egg yolk lipid composition in gulls. Functional Ecology, 13(3), 298–306.

58. Mentesana, L., Isaksson, C., Goymann, W., Andersson, M. N., Trappschuh, M., and Hau,
M. 2019. Female variation in allocation of steroid hormones, antioxidants and fatty acids:
a multilevel analysis in a wild passerine bird. Journal of Avian Biology, 50(1).

59. Ji, L. L. (2008). Modulation of skeletal muscle antioxidant defense by exercise: Role of
redox signaling. Free Radical Biology and Medicine, 44(2), 142–152.

60. Casagrande, S. & Hau, M. (2018). Enzymatic antioxidants but not baseline
glucocorticoids mediate the reproduction–survival trade-off in a wild bird. Proceedings of
the Royal Society of London B: Biological Sciences, 285: 20182141.

61. Rojas Mora, A., Meniri, M., Gning, O., Glause,r G., Vallat, A., & Helfenstein, F. (2017).
Antioxidant allocation modulates sperm quality across changing social environments.
PLoS ONE, 12(5): e0176385.

62. Parolini, M., Khoriauli, L., Possenti, C. D., Colombo, G., Caprioli, M., Santagostino, M.,
Nergadze, S. G., Milzani, A., Giulotto, E. & Saino, N. (2017). Yolk vitamin E prevents
oxidative damage in gull hatchlings. Royal Society Open Science, 4.

63. Speakman, J. R., Blount, J. D., Bronikowski, A. M., Buffenstein, R., Isaksson, C.,
Kirkwood, T. B. L., et al. (2015). Oxidative stress and life histories: unresolved issues
and current needs. Ecology and Evolution. 5, 5745–5757.

64. Skrip, M. M., Seeram, N. P., Yuan, T., Ma, H., & McWilliams, S. R. (2016). Dietary
antioxidants and flight exercise in female birds affect allocation of nutrients to eggs: how
carry-over effects work. Journal of Experimental Biology, 219(17).

65. Goymann, W., Wittenzellner, A., Schwabl, I. & Makomba, M. (2008) Progesterone
modulates aggression in sex-role reversed female African black coucals. Proceedings of
the Royal Society of London B: Biological Sciences, 275.

66. Woodley, S. K., & Moore, M. C. (1999). Female territorial aggression and steroid
hormones in mountain spiny lizards. Animal Behaviour, 57(5), 1083–1089.

67. Itagaki, T., Shimizu, I., Cheng, X., Yuan, Y., Oshio, A., Tamaki, K., Fukuno, H., Honda,
H., Okamura, Y., & Ito, S. (2005). Opposing effects of oestradiol and progesterone on
intracellular pathways and activation processes in the oxidative stress-induced activation
of cultured rat hepatic stellate cells. Gut, 12:1782-1789.

68. Sapolsky, R. M., Romero, L. M., & Munck, A. U. (2000). How do glucocorticoids
influence stress responses? Integrating permissive, suppressive, stimulatory, and
preparative actions. Endocrine Reviews, 21(1), 55–89.

69. Charlier, T. D., Underhill, C., Hammond, G. L., & Soma, K. K. (2009). Effects of
aggressive encounters on plasma corticosteroid-binding globulin and its ligands in white-
crowned sparrows. Hormones and Behavior, 56, 339–47.

70. Landys, M. M., Goymann, W., Schwabl, I., Trapschuh, M., & Slagsvold, T. (2010).
Impact of season and social challenge on testosterone and corticosterone levels in a year-
round territorial bird. Hormones and Behavior, 58(2),317-325.

71. Ros, A. F. H., Vullioud, P., Bruintjes, R., Vallat, A., Bshary, R. (2014). Intra- and
interspecific challenges modulate cortisol but not androgen levels in a year-round
territorial damselfish. Journal of Experimental Biology, 217,1768–74.

72. Haussmann, M. F., & Marchetto, N. M. (2010). Telomeres: Linking stress and survival,
ecology and evolution. Current Zoology, 56(6), 714–727.

73. Costantini, D., Marasco, V., & Møller, A. P. (2011). A meta-analysis of glucocorticoids
as modulators of oxidative stress in vertebrates. Journal of Comparative Physiology B,
181(4), 447–456.

74. Haussmann, M. F., Longenecker, A. S., Marchetto, N. M., Juliano, S. A., & Bowden, R.
M. (2012). Embryonic exposure to corticosterone modifies the juvenile stress response,
oxidative stress and telomere length. Proceedings of the Royal Society of London B:
Biological Sciences, 279(1732).

75. Majer, A. D., Fasanello, V. J., Tindle, K., Frenz, B. J., Ziur, A. D., Fischer, C. P.,
Fletcher, K. L., Seecof, K. L., Gronsky, S., Vasallo, B. G., Reed, W. L., Paitz, R. T.,
Stier, A., & Haussmann, M. F. (2019). Is there an oxidative cost of acute stress?
Characterization, implication of glucocorticoids and modulation by prior stress
experience. Proceedings of the Royal Society of London B: Biological Sciences, 286.

76. Ouyang, J. Q., Lendvai, Á. Z., Moore, I. T., Bonier, F. & Haussmann, M .F. (2016). Do
Hormones, Telomere Lengths, and Oxidative Stress form an Integrated Phenotype? A
Case Study in Free-Living Tree Swallows. Integrative and Comparative Biology, 56,
138–145.