Hybrid genome assembly and evidence-based annotation of the egg parasitoid and biological control agent *Trichogramma brassicae*

K. B. Ferguson[*], T. Kursch-Metz[†,‡], E. C. Verhulst[§], and B. A. Pannebakker[*]

* Wageningen University, Laboratory of Genetics, Wageningen, The Netherlands

† Technische Universität Darmstadt, Department of Biology, Darmstadt, Germany

‡ AMW Nützlinge GmbH, Pfungstadt, Germany

§ Wageningen University, Laboratory of Entomology, Wageningen, The Netherlands

ABSTRACT

Trichogramma brassicae (Bezdenko) are egg parasitoids that are used

throughout the world as biological control agents and in laboratories as model

species. Despite this ubiquity, few genetic resources exist beyond COI, ITS2, and

RAPD markers. Aided by a *Wolbachia* infection, a wild-caught strain from

Germany was reared for low heterozygosity and sequenced in a hybrid *de*

novo strategy, after which several assembling strategies were evaluated. The

best assembly, derived from a DBG2OLC-based pipeline, yielded a genome of

235 Mbp made up of 1,572 contigs with an N50 of 556,663 bp. Following a

rigorous *ab initio-*, homology-, and evidence-based annotation, 16,905 genes were annotated and functionally described. As an example of the utility of the genome, a simple ortholog cluster analysis was performed with sister species *T. pretiosum*, revealing over 6000 shared clusters and under 400 clusters unique to each species. The genome and transcriptome presented here provides an essential resource for comparative genomics of the commercially relevant genus *Trichogramma*, but also for research into molecular evolution, ecology, and breeding of *T. brassicae*.

INTRODUCTION

The chalcidoid *Trichogramma brassicae* (Bezdenko) (Hymenoptera: Trichogrammatidae) is a minute parasitoid wasp (~0.5 mm in length) that develops within the eggs of other insects (Smith, 1996). For over 50 years, it has been in use world-wide as a biological control agent as many lepidopteran pests of different crops are suitable hosts (Polaszek, 2009). The most common application of *T. brassicae* in Europe is against *Ostrinia nubilalis* (Hubner) (Lepidoptera: Pyralidae), the European corn borer. For example, in 2003 alone, over 11000 ha of maize in Germany was treated with *T. brassicae* (Zimmermann, 2004). It is also released against lepidopteran pests in spinach fields as well as in greenhouses (e.g. tomato, pepper, and cucumber) (Klug and Meyhöfer, 2009). With its wide application in biological control, *T. brassicae* is a well-studied species. Field trials have been conducted on several aspects, such as host location and dispersal behaviour (Suverkropp et al., 2010, 2009), overwintering ability (Babendreier et al., 2003), while other biological control related studies considered issues related to low temperature storage (Lessard

and Boivin, 2013), reaction to insecticides (Delpuech and Delahaye, 2013;

Ghorbani et al., 2016; Jamshidnia et al., 2018; Liu and Zhang, 2012; Thubru et

al., 2018), or risk assessment (Kuske et al., 2004).

Next to its application as a biological control agent, this tiny parasitoid has

been used in other research, both in genetic studies (Cruaud et al., 2018;

Laurent et al., 1998; Wajnberg, 1993) and ecological studies (Cusumano et al.,

2015; Fatouros and Huigens, 2012; Huigens et al., 2009). In addition, several

initiatives investigate the infection of *T. brassicae* with *Wolbachia* bacteria

(Ivezić et al., 2018; Poorjavad et al., 2012) and the consequences of such an

infection (Farrokhi et al., 2010; Poorjavad et al., 2018; Rahimi-Kaldeh et al.,

2018). As *T. brassicae* is a cryptic species with several other congenerics,

misidentification and misclassification is a known issue (Polaszek, 2009). In

response, molecular identification of trichogrammatids is well studied and

established (Ivezić et al., 2018; Rugman-Jones and Stouthamer, 2017;

Stouthamer et al., 1999; Sumer et al., 2009). Recently, several RADseq libraries

were constructed from single *T. brassicae* wasps to aide in resolving the

aforementioned phylogenetic issues within *Trichogramma* (Cruaud et al.,

2018). Otherwise, the genomics of *T. brassicae* have largely been neglected

even though a well annotated genome would allow researchers and

biological control practitioners access to a wealth of information and open

new avenues for comparative genomics and transcriptomics for evolutionary,

ecological, and applied research.

Here, we report the whole-genome sequencing and annotation of a *T.*

brassicae strain infected by *Wolbachia* that had thelytokous reproduction, in

which females arise from unfertilized eggs. A hybrid *de novo* sequencing

strategy was chosen to address two common issues: we used long PacBio

Sequel reads to bridge the large segments of repetitive sequences often found

in Hymenoptera, while countering the error bias of long read technology with

the accuracy of Illumina short reads. A similar strategy was recently applied to

improve the *Apis mellifera* genome, where the long PacBio reads were the

backbone that boosted the overall contiguity of the genome, alongside the

incorporation of repetitive regions (Wallberg et al., 2019).

In this report, we present the hybrid *de novo* genome of *T. brassicae*. Three

different assemblers were evaluated, and the most complete genome

assembly was used for decontamination and *ab initio*-, homology-, and

evidence-based annotation. The resulting annotation was functionally

described using gene ontology analysis. Finally, a heterozygosity comparison

and simple ortholog cluster analysis with the congeneric *T. pretiosum* was

performed, which can be considered a starting point for future comparative

genomics of the commercially important genus *Trichogramma*.

METHODS

Species origin and description:

Individuals of *Trichogramma brassicae* were acquired by AMW Nützlinge

GmbH (Pfungstadt, Germany). The strain was baited in May 2013 in an apple

orchard near Eberstadt, Germany. The orchard was surrounded by blackberry

hedges, forest, and other orchards. For baiting, the eggs of *Sitotroga cerealella*

(Olivier) (Lepidoptera: Gelechiidae) (Mega Corn Ltd., Bulgaria) were glued on

paper cards (AMW Nützlinge GmbH, Germany), usually used for releasing

Trichogramma sp. in corn fields and households. These cards were placed directly into the trees, approximately two meters above ground. After five days in the field, baiting cards were collected and incubated together at 25°C. Following emergence, individuals were kept together, offered *S. cerealella* eggs, and reared in a climate chamber (27±2°C, L:D=24:0h for four days, then transferred to16±2°C, L:D=0:24h until emergence).

In 2016, the offspring of twenty isolated females were transferred to Wageningen University (The Netherlands) to be reared for low heterozygosity. The resulting offspring were reared in a single general population on irradiated *Ephestia kuehniella* (Zeller) (Lepidoptera: Pyralidae) eggs as factitious hosts under laboratory conditions in a climate chamber (20 ± 5°C, RH 50 ± 5%, L:D=12:12 h). *Wolbachia* presence was determined following the PCR amplification protocol of Zhou et al 1998 in a presence/absence assessment with known positive and negative control samples (Zhou et al., 1998). Natural *Wolbachia* infections have previously been detected in Iranian populations of *T. brassicae* (Farrokhi et al., 2010), but none of the Eurasian populations have been known to support this symbiosis (Stouthamer, 1997; Stouthamer and Huigens, 2003). Sequence coverage was calculated using the previously established genome size estimate for *T. brassicae* of 246 Mbp (Johnston et al., 2004).

Isofemale line:

Following confirmation of *Wolbachia* infection (Supplementary materials S1.1.1), a single female from the general population was isolated (generation 0, G0), and given eggs *ad libitum*. In the resulting generation (G1), unmated

females were isolated and reared with eggs *ad libitum*. Offspring of the initial

isolations G0 and G1 were confirmed to be entirely female, suggesting

thelytokous parthenogenetic reproduction. Combined with isolating single

females, this maximizes genetic similarity of the following generation (G2) of

these G1 females. One of these G2 strains, S301, was boosted for multiple

generations over the period of one year. By the time of collection for

sequencing, both the S301 and general population no longer harboured

Wolbachia at detectable levels (Supplementary materials S1.1.2).

gDNA extraction:

Three separate extractions were prepared in 1.5 mL safelock tubes with each

several hundred *Trichogramma brassicae*. The tubes were frozen in liquid

nitrogen with approximately six 1-mm glass beads and shaken for 30 s in a

Silamat S6 shaker (Ivoclar Vivadent, Schaan, Liechtenstein). DNA was then

extracted using the Qiagen MagAttract Kit (Qiagen, Hilden, Germany).

Following an overnight lysis step with Buffer ATL and proteinase K at 56°C,

extraction was performed according to the MagAttract Kit protocol. Elutions

were performed in two steps with Buffer AE (Tris-EDTA) each time (first 60 µL,

then 40 µL), yielding 100 µL. The two extractions yielding the largest amount of

DNA (5.49 µg and 8.24 µg) were combined for long-read sequencing, while the

remaining extraction (1.67 µg) was used for short-read sequencing. DNA

concentration was measured with an Invitrogen Qubit 2.0 fluorometer using the

dsDNA HS Assay Kit (Thermo Fisher Scientific, Waltham, USA) while fragment

length was confirmed on gel.

Library preparation and sequencing:

Library preparation and sequencing was performed by Novogene

Bioinformatics Technology Co., Ltd., (Beijing, China). For Illumina sequencing,

gDNA was used to construct one paired-end (PE) library according to the

standard protocol for Illumina with an average insert size of 150 bp and was

sequenced using an Illumina HiSeq 2000 (Illumina, San Diego, USA). For Single

Molecule Real Time (SMRT) sequencing, gDNA was selected for optimal size

using a Blue Pippin size selection system (Sage Science, Beverley, USA) following

a standard library preparation. The library was then sequenced on a PacBio

Sequel (Pacific Biosciences, Menlo Park, USA) with 16 SMRT cells.

Assembly and decontamination:

Prior to assembly, Illumina reads were assessed for quality using *FASTQC*

(Andrews et al., 2015), then trimmed for quality in *CLC Genomics Workbench*

11 using default settings (Qiagen). Trimmed Illumina reads were paired for

subsequent analysis.

In order to achieve the best possible assembly, three assembly pipelines were

evaluated: one for PacBio-only reads and two hybrid assemblers. The PacBio-

only were assembled with *Canu* (v1.6) with modifications based on PacBio

Sequel reads (`correctedErrorRate=0.085 corMhapSensitivity=normal`

`alongside corMhapSensivity=normal`) (Koren et al., 2017). This is assembly

version v1.0 in the subsequent discussion.

The first hybrid assembly pipeline using both long and short sequencing read

sets was *SPAdes* (v3.11.1) (Bankevich et al., 2012). The *SPAdes* genome toolkit

supports hybrid assemblies with the *hybridSPAdes* algorithm (Antipov et al.,

2016). Three iterations of the *SPAdes* pipeline were run with varying k-mer sizes

resulting in three different assembly versions: 21, 33, 55 (default, v2.1); k-mer sizes

21, 33, 55, 77 (v2.2); and a single k-mer size of 127 (v2.3).

The second hybrid assembly pipeline was *DBG2OLC* (Ye et al., 2016). The

DBG2OLC pipeline can be readily tweaked with other programs depending on

the job (Chakraborty et al., 2016). Following the *DBG2OLC* pipeline, de Bruijn

graph contigs were generated using *SparseAssembler* using default settings

and setting the expected genome size to 750 Mbp to ensure a genome size

output that is unrestricted (Ye et al., 2012). Contigs were transformed into read

overlaps using *DBG2OLC* with settings suggested for large genomes and

PacBio Sequel data (`k=17; AdaptiveTh=0.01; KmerCovTh=2;`

`MinOverlap=20; RemoveChimera=1`), according to the *DBG2OLC* manual

(https://github.com/yechengxi/DBG2OLC). This creates an assembly

backbone of the best overlaps between the short-read de Bruijn contigs and

the long reads. *minimap2* (v2.9) and *Racon* (v1.0.2) were used for consensus

calling remaining overlaps to the assembly backbone (Li, 2018; Vaser et al.,

2017). The resulting consensus assembly was polished twice using the Illumina

reads with *Pilon* (v1.22) (Walker et al., 2014). This final assembly is v3.0 in

subsequent discussion.

The best of the five assemblies generated was determined on the basis of N50,

genome size, and completeness (Table 1). Genome statistics such as N50,

number of contig, and genome size were determined using *Quast* (Gurevich

et al., 2013). Assembly completeness was assessed using *BUSCO* (v3.0.2) with

the insect_odb9 ortholog set and the fly training parameter (Simão et al., 2015).

Based on these characteristics, the decision was made to move forward with

assembly v3.0, which was then decontaminated for microbial sequences using

NCBI *BLASTn* (v2.2.31+) against the NCBI nucleotide collection (nr).

***Wolbachia* contamination:**

Two contigs contained a large amount of *Wolbachia* content, with over 80%

of the scaffold containing material with 75% or higher homology to *Wolbachia*.

These contigs were assessed for homology against the NCBI nucleotide

collection (nr) and removed from the assembly (Supplementary material S1.2).

Post-decontamination, the assembly is referred to as v3.5.

RNA extraction, library construction, and sequencing:

T. brassicae wasps from the S301 line were collected for RNAseq for evidence-

based annotation. Hundreds of adult individuals (male and female) were killed

by freezing at -80˚C, then frozen in liquid nitrogen in a single 1.5 mL safelock

tube with approximately six 1-mm glass beads and shaken for 30 s in a Silamat

S6 shaker (Ivoclar Vivadent). The RNeasy Blood and Tissue Kit (Qiagen) was

used according to manufacturer's instructions, and final column elution was

achieved using 60 µL sterilized water. The sample was measured for quality and

RNA quantity using an Invitrogen Qubit 2.0 fluorometer and the RNA BR Assay

Kit (Thermo Fisher Scientific). The RNA sample was then processed by

Novogene Bioinformatics Technology Co., Ltd., (Beijing, China) using poly(A)

selection followed by cDNA synthesis with random hexamers and library

construction with an insert size of 300 bp. Paired-end sequencing was

performed on an Illumina HiSeq 4000 according to manufacturer's instruction.

Quality filtering was applied to remove adapters, reads with more than 10%

undetermined bases, and reads of low quality for more than 50% of the total

bases (Qscore less than or equal to 5).

Ab initio gene finding, transcriptome assembly, and annotation:

For the _ab initio_ gene finding, a training set was established using the reference

genome of _Drosophila melanogaster_ (Meigen) (Diptera: Drosophilidae)

(Genbank: GCA_000001215.4; Release 6 plus ISO1 MT) and the associated

annotation (Adams et al., 2000; Dos Santos et al., 2015). The training parameters

were used by _GlimmerHMM_ (v3.0.1) for gene finding in the _T. brassicae_ genome

assembly v3.5 (Majoros et al., 2004). For homology-based gene prediction,

GeMoMa v1.6 was used with the _D. melanogaster_ reference genome

alongside our RNAseq data as evidence for splice site prediction (Keilwagen

et al., 2016). For evidence-based gene finding, the pooled RNAseq data was

mapped to the to the _T. brassicae_ genome separately with _TopHat_ (v2.0.14)

with default settings (Trapnell et al., 2009). After mapping, _Cufflinks_ (v2.2.1) was

used to assemble transcripts (Trapnell et al., 2010). _CodingQuarry_ (v1.2) was

used for gene finding in the genome using the assembled transcripts, with the

strandness setting set to 'unstranded' (Testa et al., 2015).

The tool _EVidenceModeler (EVM)_ (v1.1.1) was used to combine the _ab initio_,

homology-based, and evidence-based information, with evidence-based

weighted 1, _ab initio_ weighted 2, and homology-based weighted 3 (Haas et al.

2008). We annotated the predicted proteins with _BLASTp_ (v2.2.31+) on a

custom database containing all SwissProt and Refseq genes of _D._

melanogaster (Acland et al., 2014; Boutet et al., 2008; Camacho et al., 2009),

followed by an additional search in the NCBI non-redundant protein database

(nr) to obtain additional homology data.

GO term analysis:

A list of genes was constructed for Gene Ontology (GO) term classification by

deduplicating the annotated proteins and removing the non-annotated

proteins. These accession IDs were converted into UniProtKB accession IDs using

the UniProt ID mapping feature and deduplicated a final time (Boutet et al.,

2008). These UniProtKB accession IDs were in turn used with the *DAVID 6.8*

Functional Annotation Tool to assign GO terms to each accession ID with the

D. melanogaster background and generate initial functional analyses (Huang

et al., 2009a, 2009b) (see supplementary S1.3 for *DAVID* input list).

Heterozygosity estimates:

The heterozygosity of the S301 line was assessed using sequence reads and k-

mer counting, and compared to the congeneric *Trichogramma pretiosum*

(Riley) (Hymenoptera: Trichogrammatidae), for which sequence data exists for

both a thelytokous (asexual) *Wolbachia*-infected strain as well as an inbred

arrhenotokous (sexual) line (Lindsey et al., 2018). Using *jellyfish* (v2.3.0) to count

k-mers, the same trimmed and paired Illumina reads used for assembly were

assessed using the default k-mer size of 21 (m=21), with results exported to a

histogram (Marçais and Kingsford, 2011). This histogram file was then used with

GenomeScope (v1.0) to estimate heterozygosity of the reads based on a

statistical model, where a Poisson distribution is expected for a homozygous

sample while a bimodal distribution is expected for a homozygous distribution

(Vurture et al., 2017). This genome profiling gives a reliable estimate for

heterozygosity as well as estimates of repetitive content. The same *jellyfish* and

GenomeScope analyses were performed on *T. pretiosum* short-read sequence

data for the thelytokous strain (NCBI SRA database, SRR1191749) and the

arrhenotokous line (SRR6447489), with adaptions for reported insert sizes

(Lindsey et al., 2018).

Ortholog cluster analysis:

The complete gene set of *T. brassicae* was compared to that of *T. pretiosum*

(Lindsey et al., 2018), which was retrieved from the i5K Workspace (Poelchau

et al., 2016). An ortholog cluster analysis was performed on both gene sets via

OrthoVenn2 with the default settings of E-values of 1e-5 and an inflation value

of 1.5 (Xu et al., 2019). For *T. brassicae* protein set, see supplementary materials

S1.5.

Data availability:

All sequence data are available at the EMBL-ENA database under BioProject

PRJEB35413, including assembly (CADCXV010000000.1). An additional,

complete annotation file (.gff) is also available (Ferguson, 2020). Additional

data, such as gel images, the *Wolbachia* contaminated contigs, input gene list

for *DAVID*, *GenomeScope* images, and complete protein set are available via

the supplementary materials.

RESULTS AND DISCUSSION

Sequencing, assembly, and decontamination:

Sequencing of the Illumina 150 bp paired-end library yielded 80,489,816 reads.

After quality filtering and trimming, 80,483,128 paired-end reads were retained.

Sequencing the PacBio Sequel library yielded 2,500,204 subreads with an

average length of 6377 bp. The genome size estimate for *T. brassicae* is 246

Mbp (Johnston et al., 2004) indicating that short-read coverage was 98x while

long-read coverage was 64x, resulting in a total coverage of 162x. Three

assembly pipelines were used, resulting in five potential assemblies where one,

v3.0, was eventually selected for further use. Results of these assemblies are

detailed in Table 1.

The first draft assembly generated with Canu with the altered settings for PacBio

Sequel data resulted in an assembly of approximately 70 Mbp in size, drastically

smaller than the 246 Mbp expected, and contained a total of 3,007 contigs

with an N50 of 27,303. The longest contig was 126,800 bp in size.

The second assembly strategy relied on hybrid assembly pipelines, and *SPAdes*

was used with the default k-mer settings, which resulted in an assembly of

approximately 227 Mbp in size with an N50 of 36,870 and a *BUSCO*

completeness of 96.8%. Three different assembly runs were done with differing

k-mer sizes: the default k-mer sizes of 21, 33, 55 (v2.1); default k-mer sizes plus 77

(v2.2); or the highest possible k-mer size of 127 (v2.3). Increasing the k-mer size

only improved N50 scores to a point, along with decreasing the number of

contigs, and stable *BUSCO* scores, however, the assembled genome size drops

dramatically with the third attempt shrinking down to 211 Mbp. Based on

BUSCO scores and N50 alone, the second *SPAdes* attempt, v2.2, would be the

best of the three, though all three are similar in most measures.

The third assembly strategy used the *DGB2OLC+Racon+Pilon* pipeline, which

resulted in assembly v3.0. Here, there is a large difference compared to the

previous *SPAdes* assemblies. Particularly, the number of contigs is reduced

dramatically from the 70,000 to 280,000 range of the *SPAdes* output down to a

mere 1,572. Meanwhile, the assembled genome size is now 235 Mbp and with

an N50 of 556,663 and a BUSCO score of 95.5%. The full completeness score for

this assembly, using the 1658 BUSCO groups within the insect_od09 BUSCO set,

returned 1531 (92.3%) complete and single-copy BUSCOs, 53 (3.2%) complete

and duplicated BUSCOs, 22 (1.3%) fragmented BUSCOs, and 52 (3.2%) missing

BUSCOs (Simão et al., 2015).

While the PacBio-only assembly in *Canu* could have been improved using

different settings or additional tools, we decided to focus on using the

additional sequence information of the Illumina reads in the subsequent hybrid

assembly strategies. The *SPAdes* assemblies (v2.1-3) were already decent but

could have been further improved using *Pilon*, a tool that improves assemblies

at the base pair level using high quality Illumina data. However, the v3.0

assembly was by far the best assembly based on assembled genome size, N50,

and BUSCO scores and therefore we chose this strategy for our *T. brassicae*

genome assembly.

Decontamination of this assembly (v3.0) resulted in the removal of two contigs

as the homology analysis using *BLASTn* with the NCBI nr database indicated

that both contigs were confirmed to be largely composed of *Wolbachia*

genomic content. Contig "Backbone_1176" is 9,448 bp in length and two areas

of the contig, representing over 80% of its length, showed high homology to

Wolbachia. Similarly, contig "Backbone_1392" is 17,350 bp and three separate

areas representing over 80% showed similar levels of homology to *Wolbachia*

After decontamination this final assembly (v3.5) was used for annotation.

Ab initio gene finding, transcriptome assembly, and annotation:

In our RNA sequencing experiment, we generated 26,479,830 150bp paired-

end cDNA reads. Filtering the reads for quality retained 99.3% of these reads to

be used for evidence-based gene finding via transcriptome assembly.

The annotations from the evidence-based gene finding were used alongside

homology-based findings and _ab initio_ annotations in a weighted model,

resulting in a complete annotation for the assembly. In 865 mRNA tracks,

representing approximately 5.1% of the official gene set, a gene model could

not be annotated via the SwissProt database, and these tracks are named

"No_blast_hit." The majority of tracks are annotated with reference to SwissProt

or GenBank accession number of the top _BLASTp_ hit.

Transcriptome assembly and mapping resulted in 45,876,158 mapped

transcripts (48,327,134 total). _CodingQuarry_ predicted 45,454 evidence-based

genes from these mapped transcripts, while _ab initio_ gene finding using

GlimmerHMM resulted in 16,877 genes and homology-based gene finding with

GeMoMa resulted in 6,675 genes. The final complete gene set was created

using _EVidenceModeler_, where a weighted model using all three inputs

resulted in a complete gene set of 16,905 genes.

GO term analysis:

The complete gene set of 16,905 genes was deduplicated and genes with no

correlating _BLASTp_ hit were removed from this analysis. The remaining 9,373

genes were subjected to UniProtKB ID mapping, resulting in 8,247 genes with a

matching ID after another round of deduplication (828 duplicates found). The

remaining 755 accession IDs were not able to be matched, half of which are

obsolete proteins within the UniParc database (377).

The *DAVID Functional Annotation Tool* used 6,585 genes for the analysis and

showed that 80.8% (5,320) contribute to 530 biological processes, 77.5% (5,104)

contribute to 115 different cellular component categories, and 74.2% (4,889)

contribute to 93 molecular functions (genes can code to multiple GO terms).

The remaining 1,662 genes are uncategorised.

Heterozygosity estimates:

Using short-read data and k-mer counting, heterozygosity was estimated for

our isofemale S301 line and compared to both a parthenogenesis inducing

Wolbachia-infected strain and an arrhenotokous line of *T. pretiosum* (Lindsey et

al., 2018). The average estimated heterozygosity for our S301 *T. brassicae* line is

0.0332% with approximately 0.608% repetitive content (for full details, see Table

2). This is similar to the thelytokous *T. pretiosum* line, which has a slightly lower

estimated heterozygosity (0.0289%) and a lower amount of repetitive content

(0.482%). Both have a very distinct Poisson distribution, indicating a low

heterozygosity (Figure S1.4.1-2). The arrhenotokous *T. pretiosum* showed a

higher estimated heterozygosity (0.863%), a larger amount of repetitive content

(2.64%), and a slightly bimodal distribution (Figure S1.4.3). The fact that both

thelytokous *Trichogramma* species have a similar low level of heterozygosity

when compared to the arrhenotokous *T. pretiosum* suggests that in both cases

Wolbachia infection had a severe effect on genetic diversity. As the canonical

mechanism of parthenogenesis-induction in other *Wolbachia* infected

thelytokous *Trichogramma* species is gamete duplication (Pannebakker et al.,

2004; Stouthamer and Kazmer, 1994), in which unfertilized eggs are diploidized

and results in fully homozygous progeny in a single generation, the low genomic

heterozygosity rate suggests a similar mechanism for *Wolbachia*-induced

parthenogenesis in *T. brassicae*. However, the involvement of *Wolbachia* in

causing all-female offspring in this *T. brassicae* strain and the presence and

mechanisms of *Wolbachia* in other thelytokous *T. brassicae* strains (Farrokhi et

al., 2010; Poorjavad et al., 2018, 2012) does require further investigation.

Ortholog cluster analysis:

The complete gene set of *T. brassicae* was compared to that of *T. pretiosum*

using *OrthoVenn2* (full output in Table 3). Both species have a similar range of

proteins (16,905 in *T. brassicae* and 13,200 in *T. pretiosum*) that form a similar

number of clusters (6,537 in *T. brassicae* and 6,489 in *T. pretiosum*). The two

species share 6,158 clusters (of 16,899 proteins), while *T. brassicae* has 379

unique clusters (1,726 proteins) and *T. pretiosum* has 331 unique clusters (1,005

proteins), as shown in Figure 1. These unique clusters account for approximately

5% of the entire cluster set for both species, and may both indicate true areas

of differentiation, or result from differences in the annotation strategies. There is

a similar amount of singleton clusters (proteins that do not cluster with others) in

T. brassicae (5,291) and *T. pretiosum* (5,184). Both the unique clusters and the

unique single-copy genes could be novel proteins, regions of contamination,

evidence of unique horizontal gene transfer, or pseudogenes. More

investigation into these protein clusters in addition to a more comprehensive

manual annotation should shed some light on the differences between these

closely related yet geographically distinct parasitoid wasps.

CONCLUSIONS AND PERSPECTIVES

Here, we present the genome of biological control agent *Trichogramma*

brassicae, a chalcidoid wasp used throughout the world for augmentative

biological control as well as genetic and ecological research. This unique strain

hosted a parthenogenesis-inducing *Wolbachia* infection and is the first

European *Trichogramma* genome to be published, allowing for comparative

analyses with other *Trichogramma* genomes, as we have shown. Our genomic

data also illuminates the possible mechanism of parthenogenesis-induction by

Wolbachia in this strain. Furthermore, the variety of genomic and transcriptomic

data generated for this genome provide much-need resources to bring *T.*

brassicae into the -omics era of biological research.

A hybrid approach was used, resulting in a highly contiguous assembly of 1,572

contigs and 16,905 genes based on *ab initio*, homology-based, and evidence-

based annotation, for a total assembly size of 235 Mbp. Two scaffolds were

identified that were of *Wolbachia* origin and removed. Ortholog cluster analysis

showed 379 unique protein clusters containing 1,726 proteins. Future studies are

needed to show whether these clusters are truly unique. This genome and

annotation provides the basis for future, more in-depth comparative studies

into the genetics, evolution, ecology, and biological control use of

Trichogramma species.

ACKNOWLEDGEMENTS AND FUNDING

We would like to thank Bernd Wührer (AMW Nützlinge) for providing access to

specimens; Gabrielle Bukovinszkine Kiss, José van de Belt, Frank Becker

(Wageningen University), and Lorraine Latchoumane for assistance with

rearing, DNA, and RNA extraction; Richard Stouthamer and Nina Fatouros for

sharing their *Trichogramma* knowledge; and Sophie Chattington, Andra Thiel

(University of Bremen), and Bas Zwaan (Wageningen University) for their

assistance in this project. GenomeScan B. V. performed decontamination and

annotation analysis. This work has received funding from the European Union

Horizon 2020 research and innovation programme under the Marie

Skłodowska-Curie grant agreement no. 641456.

SUPPLEMENTARY MATERIALS

Additional supplementary material from this study is available in an attached

document, with some material found on the DANS EASY Repository,

https://doi.org/10.17026/dans-23w-a9tn (explanation within supporting

document).

T. brassicae   *T. pretiosum*

429   Figure 1 Ortholog clusters analysis between *Trichogramma brassicae* and *T.*

430   *pretiosum* using OrthoVenn2 (Xu et al., 2019). The number of clusters shared

431   between the two organisms is in bold.

Table 1. Statistics for five assemblies of *Trichogramma brassicae*. The first strategy was PacBio-only in *Canu*, while three hybrid assembly strategies were based on *SPAdes* and modulating k-mer sizes, and an additional hybrid assembly was based on an adapted *DBG2OLC+Racon+Pilon* protocol. BUSCO score is based on the insect_db09 dataset (Simão et al., 2015).

Assembler	Version	Size (bp)	Contigs	Longest contig (bp)	N50 (bp)	BUSCO (Complete %)
Canu	v1.0	69,522,446	3,007	126,800	27,303	18.7
SPAdes (k=21, 33, 55)	v 2.1	227,096,967	282,988	474,998	36,870	96.8
SPAdes (k=21,33, 55, 77)	v 2.2	226,864,253	189,696	548,753	49,096	97.1
SPAdes (k=127)	v 2.3	211,402,326	73,567	537,817	63,558	96.4
DBG2OLC+ Racon+Pilon	v 3.0	235,413,774	1,572	2,953,580	556,663	95.5

Table 2. Heterozygosity and repetitive content analysis of *Trichogramma brassicae* (thelytokous), *Trichogramma pretiosum* (thelytokous), and *T. pretiosum* (arrhenotokous) lines based on sequence data.

	Heterozygosity (%)	Repetitive content (%)	Source of sequence data
T. brassicae, thelytokous S301 line	0.0332	0.608	This publication
T. pretiosum, thelytokous *Wolbachia* line	0.0289	0.482	Lindsey et al., 2018
T. pretiosum, arrhenotokous inbred line	0.863	2.64	Lindsey et al., 2018

Table 3. Output of OrthoVenn2 ortholog cluster analysis of *Trichogramma brassicae* and *Trichogramma pretiosum*.

Species	Proteins	Clusters	Singletons	Source of gene set
T. brassicae	16,905	6,537	5,291	This work (S1.5)
T. pretiosum	13,200	6,489	5,184	Lindsey et al., 2018; Poelchau et al., 2015

439   LITERATURE CITED

440   Acland, A., Agarwala, R., Barrett, T., Beck, J., Benson, D.A., Bollin, C., Bolton, E.,

441   Bryant, S.H., Canese, K., Church, D.M., Clark, K., Dicuccio, M.,

442   Dondoshansky, I., Federhen, S., Feolo, M., Geer, L.Y., Gorelenkov, V.,

443   Hoeppner, M., Johnson, M., Kelly, C., Khotomlianski, V., Kimchi, A.,

444   Kimelman, M., Kitts, P., Krasnov, S., Kuznetsov, A., Landsman, D., Lipman,

445   D.J., Lu, Z., Madden, T.L., Madej, T., Maglott, D.R., Marchler-Bauer, A.,

446   Karsch-Mizrachi, I., Murphy, T., Ostell, J., O'Sullivan, C., Panchenko, A.,

447   Phan, L., Pruitt, D.P.K.D., Rubinstein, W., Sayers, E.W., Schneider, V., Schuler,

448   G.D., Sequeira, E., Sherry, S.T., Shumway, M., Sirotkin, K., Siyan, K., Slotta, D.,

449   Soboleva, A., Soussov, V., Starchenko, G., Tatusova, T.A., Trawick, B.W.,

450   Vakatov, D., Wang, Y., Ward, M., John Wilbur, W., Yaschenko, E., Zbicz, K.,

451   2014. Database resources of the National Center for Biotechnology

452   Information. Nucleic Acids Res. 42, 8–13.

453   https://doi.org/10.1093/nar/gkt1146

454   Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D.,

455   Amanatides, P.G., Scherer, S.E., Li, P.W., Hoskins, R.A., Galle, R.F., George,

456   R.A., Lewis, S.E., Richards, S., Ashburner, M., Henderson, S.N., Sutton, G.G.,

457   Wortman, J.R., Yandell, M.D., Zhang, Q., Chen, L.X., Brandon, R.C., Rogers,

458   Y.H.C., Blazej, R.G., Champe, M., Pfeiffer, B.D., Wan, K.H., Doyle, C.,

459   Baxter, E.G., Helt, G., Nelson, C.R., Gabor Miklos, G.L., Abril, J.F., Agbayani,

460   A., An, H.J., Andrews-Pfannkoch, C., Baldwin, D., Ballew, R.M., Basu, A.,

461   Baxendale, J., Bayraktaroglu, L., Beasley, E.M., Beeson, K.Y., Benos, P. V.,

462   Berman, B.P., Bhandari, D., Bolshakov, S., Borkova, D., Botchan, M.R.,

463 Bouck, J., Brokstein, P., Brottier, P., Burtis, K.C., Busam, D.A., Butler, H.,

464 Cadieu, E., Center, A., Chandra, I., Michael Cherry, J., Cawley, S., Dahlke,

465 C., Davenport, L.B., Davies, P., de Pablos, B., Delcher, A., Deng, Z.,

466 Deslattes Mays, A., Dew, I., Dietz, S.M., Dodson, K., Doup, L.E., Downes, M.,

467 Dugan-Rocha, S., Dunkov, B.C., Dunn, P., Durbin, K.J., Evangelista, C.C.,

468 Ferraz, C., Ferriera, S., Fleischmann, W., Fosler, C., Gabrielian, A.E., Garg,

469 N.S., Gelbart, W.M., Glasser, K., Glodek, A., Gong, F., Harley Gorrell, J., Gu,

470 Z., Guan, P., Harris, M., Harris, N.L., Harvey, D., Heiman, T.J., Hernandez,

471 J.R., Houck, J., Hostin, D., Houston, K.A., Howland, T.J., Wei, M.H.,

472 Ibegwam, C., Jalali, M., Kalush, F., Karpen, G.H., Ke, Z., Kennison, J.A.,

473 Ketchum, K.A., Kimmel, B.E., Kodira, C.D., Kraft, C., Kravitz, S., Kulp, D., Lai,

474 Z., Lasko, P., Lei, Y., Levitsky, A.A., Li, J., Li, Z., Liang, Y., Lin, X., Liu, X., Mattei,

475 B., McIntosh, T.C., McLeod, M.P., McPherson, D., Merkulov, G., Milshina, N.

476 V., Mobarry, C., Morris, J., Moshrefi, A., Mount, S.M., Moy, M., Murphy, B.,

477 Murphy, L., Muzny, D.M., Nelson, D.L., Nelson, D.R., Nelson, K.A., Nixon, K.,

478 Nusskern, D.R., Pacleb, J.M., Palazzolo, M., Pittman, G.S., Pan, S., Pollard,

479 J., Puri, V., Reese, M.G., Reinert, K., Remington, K., Saunders, R.D.C.,

480 Scheeler, F., Shen, H., Christopher Shue, B., Siden-Kiamos, I., Simpson, M.,

481 Skupski, M.P., Smith, T., Spier, E., Spradling, A.C., Stapleton, M., Strong, R.,

482 Sun, E., Svirskas, R., Tector, C., Turner, R., Venter, E., Wang, A.H., Wang, X.,

483 Wang, Z.Y., Wassarman, D.A., Weinstock, G.M., Weissenbach, J., Williams,

484 S.M., Woodage, T., Worley, K.C., Wu, D., Yang, S., Alison Yao, Q., Ye, J.,

485 Yeh, R.F., Zaveri, J.S., Zhan, M., Zhang, G., Zhao, Q., Zheng, L., Zheng, X.H.,

486 Zhong, F.N., Zhong, W., Zhou, X., Zhu, S., Zhu, X., Smith, H.O., Gibbs, R.A.,

Myers, E.W., Rubin, G.M., Craig Venter, J., 2000. The genome sequence of

Drosophila melanogaster. Science (80-.). 287, 2185–2195.

https://doi.org/10.1126/science.287.5461.2185

Andrews, S., Krueger, F., Seconds-Pichon, A., Biggins, F., Wingett, S., 2015.

FastQC. A quality control tool for high throughput sequence data.

Babraham Bioinformatics. Babraham Inst.

Antipov, D., Korobeynikov, A., McLean, J.S., Pevzner, P.A., 2016. HybridSPAdes:

An algorithm for hybrid assembly of short and long reads. Bioinformatics

32, 1009–1015. https://doi.org/10.1093/bioinformatics/btv688

Babendreier, D., Rostas, M., Höfte, M.C.J., Kuske, S., Bigler, F., 2003. Effects of

mass releases of Trichogramma brassicae on predatory insects in maize.

Entomol. Exp. Appl. 108, 115–124. https://doi.org/10.1046/j.1570-

7458.2003.00075.x

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S.,

Lesin, V.M., Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A. V, Sirotkin,

A. V, Vyahhi, N., Tesler, G., Alekseyev, M.A., Pevzner, P.A., 2012. SPAdes: a

new genome assembly algorithm and its applications to single-cell

sequencing. J. Comput. Biol. 19, 455–77.

https://doi.org/10.1089/cmb.2012.0021

Boutet, E., Lieberherr, D., Tognolli, M., Schneider, M., Bairoch, A., 2008.

UniProtKB/Swiss-Prot: The manually annotated section of the UniProt

KnowledgeBase. Methods Mol. Biol. 406, 89–112.

https://doi.org/10.1007/978-1-59745-535-0

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer,

K., Madden, T.L., 2009. BLAST+: Architecture and applications. BMC

Bioinformatics 10, 421. https://doi.org/10.1186/1471-2105-10-421

Chakraborty, M., Baldwin-Brown, J.G., Long, A.D., Emerson, J.J., 2016.

Contiguous and accurate de novo assembly of metazoan genomes with

modest long read coverage. Nucleic Acids Res. 44, 1–12.

https://doi.org/10.1093/nar/gkw654

Cruaud, A., Groussier, G., Genson, G., Sauné, L., Polaszek, A., Rasplus, J.-Y.,

2018. Pushing the limits of whole genome amplification: successful

sequencing of RADseq library from a single microhymenopteran

(Chalcidoidea, *Trichogramma*). PeerJ 6, e5640.

https://doi.org/10.7717/peerj.5640

Cusumano, A., Weldegergis, B.T., Colazza, S., Dicke, M., Fatouros, N.E., 2015.

Attraction of egg-killing parasitoids toward induced plant volatiles in a

multi-herbivore context. Oecologia 179, 163–174.

https://doi.org/10.1007/s00442-015-3325-3

Delpuech, J.M., Delahaye, M., 2013. The sublethal effects of deltamethrin on

Trichogramma behaviors during the exploitation of host patches. Sci. Total

Environ. 447, 274–279. https://doi.org/10.1016/j.scitotenv.2012.12.096

Dos Santos, G., Schroeder, A.J., Goodman, J.L., Strelets, V.B., Crosby, M.A.,

Thurmond, J., Emmert, D.B., Gelbart, W.M., Brown, N.H., Kaufman, T.,

Werner-Washburne, M., Cripps, R., Broll, K., Gramates, L.S., Falls, K.,

Matthews, B.B., Russo, S., Zhou, P., Zytkovicz, M., Adryan, B., Attrill, H.,

Costa, M., Marygold, S., McQuilton, P., Millburn, G., Ponting, L., Stefancsik,

R., Tweedie, S., Grumbling, G., 2015. FlyBase: Introduction of the

Drosophila melanogaster Release 6 reference genome assembly and

large-scale migration of genome annotations. Nucleic Acids Res. 43,

D690–D697. https://doi.org/10.1093/nar/gku1099

Farrokhi, S., Ashouri, A., Shirazi, J., Allahvari, H., Huigens, M.E.E., 2010. A

comparative study on the functional response of Wolbachia-infected

and uninfected forms of the parasitoid wasp Trichogramma brassicae. J.

Insect Sci. 10, 167. https://doi.org/10.1673/031.010.14127

Fatouros, N.E., Huigens, M.E., 2012. Phoresy in the field: natural occurrence of

Trichogramma egg parasitoids on butterflies and moths. BioControl 57,

493–502. https://doi.org/10.1007/s10526-011-9427-x

Ferguson, K., 2020 Trichogramma brassicae hybrid genome annotation file.

https://doi.org/10.6084/m9.figshare.12073833.v1

Ghorbani, M., Saber, M., Bagheri, M., Vaez, N., 2016. Effects of diazinon and

fipronil on different developmental stages of Trichogramma brassicae

Bezdenko (Hym.; Trichogrammatidae). J. Agric. Sci. Technol. 18, 1267–

1278.

Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G., 2013. QUAST: Quality

assessment tool for genome assemblies. Bioinformatics 29, 1072–1075.

https://doi.org/10.1093/bioinformatics/btt086

Huang, D.W., Sherman, B.T., Lempicki, R.A., 2009a. Bioinformatics enrichment

tools: Paths toward the comprehensive functional analysis of large gene

lists. Nucleic Acids Res. 37, 1–13. https://doi.org/10.1093/nar/gkn923

Huang, D.W., Sherman, B.T., Lempicki, R.A., 2009b. Systematic and integrative

analysis of large gene lists using DAVID bioinformatics resources. Nat.

Protoc. 4, 44–57. https://doi.org/10.1038/nprot.2008.211

Huigens, M.E., Pashalidou, F.G., Qian, M.-H., Bukovinszky, T., Smid, H.M., van

Loon, J.J.A., Dicke, M., Fatouros, N.E., 2009. Hitch-hiking parasitic wasp

learns to exploit butterfly antiaphrodisiac. Proc. Natl. Acad. Sci. U. S. A.

106, 820–825. https://doi.org/10.1073/pnas.0812277106

Ivezić, A., Rugman-Jones, P., Stouthamer, R., Ignjatović-Ćupina, A., 2018.

Molecular identification of Trichogramma egg parasitoids of Ostrinia

nubilalis in northeastern Serbia. Arch. Biol. Sci. 70, 425–432.

https://doi.org/10.2298/ABS171103002I

Jamshidnia, A., Abdoli, S., Farrokhi, S., Sadeghi, R., 2018. Efficiency of spinosad,

Bacillus thuringiensis and Trichogramma brassicae against the tomato

leafminer in greenhouse. BioControl 63, 619–627.

https://doi.org/10.1007/s10526-018-9893-5

Johnston, J.S., Ross, L.D., Beani, L., Hughes, D.P., Kathirithamby, J., 2004. Tiny

genomes and endoreduplication in Strepsiptera. Insect Mol. Biol. 13, 581–

585. https://doi.org/10.1111/j.0962-1075.2004.00514.x

Keilwagen, J., Wenk, M., Erickson, J.L., Schattat, M.H., Grau, J., Hartung, F.,

2016. Using intron position conservation for homology-based gene

prediction. Nucleic Acids Res. 44. https://doi.org/10.1093/nar/gkw092

Klug, T., Meyhöfer, R., 2009. Performance of two Trichogramma brassicae

strains under greenhouse and field conditions for biocontrol of the silver Y

moth in spinach cultures. J. Pest Sci. (2004). 82, 73–79.

https://doi.org/10.1007/s10340-008-0224-y

Koren, S., Walenz, B.P., Berlin, K., Miller, J.R., Bergman, N.H., Phillippy, A.M.,

2017. Canu: scalable and accurate long-read assembly via adaptive k-

mer weighting and repeat separation. Genome Res. 27, 722–736.

https://doi.org/10.1101/gr.215087.116

Kuske, S., Babendreier, D., Edwards, P.J., Turlings, T.C.J., Bigler, F., 2004.

Parasitism of non-target lepidoptera by mass released Trichogramma

brassicae and its implication for the larval parasitoid Lydella thompsoni.

BioControl 49, 1–19. https://doi.org/10.1023/B:BICO.0000009379.13685.47

Laurent, V., Wajnberg, E., Mangin, B., Schiex, T., Gaspin, C., Vanlerberghe-

Masutti, F., 1998. A composite genetic map of the parasitoid wasp

Trichogramma brassicae based on RAPD markers. Genetics 150, 275–82.

Lessard, E., Boivin, G., 2013. Effect of low temperature on emergence,

fecundity, longevity and host-feeding by Trichogramma brassicae.

BioControl 58, 319–329. https://doi.org/10.1007/s10526-012-9493-8

Li, H., 2018. Minimap2: Pairwise alignment for nucleotide sequences.

Bioinformatics 34, 3094–3100.

https://doi.org/10.1093/bioinformatics/bty191

Lindsey, A.R.I., Kelkar, Y.D., Wu, X., Sun, D., Martinson, E.O., Yan, Z., Rugman-

Jones, P.F., Hughes, D.S.T., Murali, S.C., Qu, J., Dugan, S., Lee, S.L., Chao,

H., Dinh, H., Han, Y., Doddapaneni, H.V., Worley, K.C., Muzny, D.M., Ye, G.,

Gibbs, R.A., Richards, S., Yi, S. V., Stouthamer, R., Werren, J.H., 2018.

Comparative genomics of the miniature wasp and pest control agent

Trichogramma pretiosum. BMC Biol. 16, 1–20.

https://doi.org/10.1186/s12915-018-0520-9

Liu, T.X., Zhang, Y., 2012. Side effects of two reduceδ-risk insecticides,

607 indoxacarb and spinosad, on two species of Trichogramma

608 (Hymenoptera: Trichogrammatidae) on cabbage. Ecotoxicology 21,

609 2254–2263. https://doi.org/10.1007/s10646-012-0981-5

610 Majoros, W.H., Pertea, M., Salzberg, S.L., 2004. TigrScan and GlimmerHMM: Two

611 open source ab initio eukaryotic gene-finders. Bioinformatics 20, 2878–

612 2879. https://doi.org/10.1093/bioinformatics/bth315

613 Marçais, G., Kingsford, C., 2011. A fast, lock-free approach for efficient

614 parallel counting of occurrences of k-mers. Bioinformatics 27, 764–770.

615 https://doi.org/10.1093/bioinformatics/btr011

616 Pannebakker, B.A., Pijnacker, L.P., Zwaan, B.J., Beukeboom, L.W., 2004.

617 Cytology of *Wolbachia*-induced parthenogenesis in *Leptopilina clavipes*

618 (Hymenoptera: Figitidae). Genome 303, 299–303.

619 https://doi.org/10.1139/G03-137

620 Poelchau, M., Childers, C., Moore, G., Tsavatapalli, V., Evans, J., Lee, C.Y., Lin,

621 H., Lin, J.W., Hackett, K., 2015. The i5k Workspace@NAL-enabling genomic

622 data access, visualization and curation of arthropod genomes. Nucleic

623 Acids Res. 43, D714–D719. https://doi.org/10.1093/nar/gku983

624 Poelchau, M.F., Coates, B.S., Childers, C.P., Peréz De León, A.A., Evans, J.D.,

625 Hackett, K., Shoemaker, D.W., 2016. Agricultural applications of insect

626 ecological genomics. Curr. Opin. Insect Sci. 13, 61–69.

627 https://doi.org/10.1016/j.cois.2015.12.002

628 Polaszek, A., 2009. Species Diversity and Host Associations of Trichogramma in

629 Eurasia, in: Consoli, F.L., Parra, J.R.P., Zucchi, R.A. (Eds.), Egg Parasitoids in

630 Agroecosystems with Emphasis on Trichogramma. Springer Netherlands,

Dordrecht, pp. 237–266. https://doi.org/10.1007/978-1-4020-9110-0_9

Poorjavad, N., Goldansaz, S.H., Machtelinckx, T., Tirry, L., Stouthamer, R., van

Leeuwen, T., 2012. Iranian Trichogramma: ITS2 DNA characterization and

natural Wolbachia infection. BioControl 57, 361–374.

https://doi.org/10.1007/s10526-011-9397-z

Poorjavad, N., Goldansaz, S.H., van Leeuwen, T., 2018. Fertility life table

parameters, coi sequences and wolbachia infection in populations of

trichogramma brassicae collected from chilo suppressalis. Bull.

Insectology 71, 89–96.

Rahimi-Kaldeh, S., Ashouri, A., Bandani, A., 2018. Does Wolbachia Infection

Change the Overwintering Ability of Trichogramma brassicae

(Hymenoptera: Trichogrammatidae)? Neotrop. Entomol. 47, 583–590.

https://doi.org/10.1007/s13744-017-0549-4

Rugman-Jones, P.F., Stouthamer, R., 2017. High-resolution melt analysis without

DNA extraction affords rapid genotype resolution and species

identification. Mol. Ecol. Resour. 17, 598–607. https://doi.org/10.1111/1755-

0998.12599

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E. V., Zdobnov, E.M.,

2015. BUSCO: Assessing genome assembly and annotation completeness

with single-copy orthologs. Bioinformatics 31, 3210–3212.

https://doi.org/10.1093/bioinformatics/btv351

Smith, S.M., 1996. Biological control with Trichogramma: advances, successes,

and potential of their use. Annu. Rev. Entomol. 41, 375–406.

https://doi.org/10.1146/annurev.ento.41.1.375

Stouthamer, R., 1997. Wolbachia-induced parthenogenesis, in: O'Neill, S.L.,

Hoffmann, A.A., Werren, J.H. (Eds.), Influential Passengers: Inherited

Microorganisms and Arthropod Reproduction. Oxford University Press,

Oxford, UK, pp. 102–124.

Stouthamer, R., Hu, J., Van Kan, F.J.P.M.P.M., Platner, G.R., Pinto, J.D., 1999.

The utility of internally transcribed spacer 2 DNA sequences of the nuclear

ribosomal gene for distinguishing sibling species of Trichogramma.

BioControl 43, 421–440. https://doi.org/10.1023/A:1009937108715

Stouthamer, R., Huigens, M., 2003. Parthenogenesis Associated With

Wolbachia, in: Bourtzis, K., Miller, T.A. (Eds.), Insect Symbiosis. CRC Press,

Boca Raton, pp. 247–266. https://doi.org/10.1201/9780203009918.ch15

Stouthamer, R., Kazmer, D.J., 1994. Cytogenetics of microbe-associated

parthenogenesis and its consequences for gene flow in Trichogramma

wasps. Heredity (Edinb). 73, 317–327. https://doi.org/10.1038/hdy.1994.139

Sumer, F., Tuncbilek, A.S., Oztemiz, S., Pintureau, B., Rugman-Jones, P.,

Stouthamer, R., 2009. A molecular key to the common species of

Trichogramma of the Mediterranean region. BioControl 54, 617–624.

https://doi.org/10.1007/s10526-009-9219-8

Suverkropp, B.P., Bigler, F., van Lenteren, J.C., 2010. Movement and host

finding of Trichogramma brassicae on maize plants. Bull. Insectology 63,

115–127.

Suverkropp, B.P., Bigler, F., van Lenteren, J.C., 2009. Dispersal behaviour of

Trichogramma brassicae in maize fields. Bull. Insectology 62, 113–120.

Testa, A.C., Hane, J.K., Ellwood, S.R., Oliver, R.P., 2015. CodingQuarry: Highly

accurate hidden Markov model gene prediction in fungal genomes using

RNA-seq transcripts. BMC Genomics 16, 1–12.

https://doi.org/10.1186/s12864-015-1344-4

Thubru, D.P., Firake, D.M., Behere, G.T., 2018. Assessing risks of pesticides

targeting lepidopteran pests in cruciferous ecosystems to eggs parasitoid,

Trichogramma brassicae (Bezdenko). Saudi J. Biol. Sci. 25, 680–688.

https://doi.org/10.1016/j.sjbs.2016.04.007

Trapnell, C., Pachter, L., Salzberg, S.L., 2009. TopHat: Discovering splice

junctions with RNA-Seq. Bioinformatics 25, 1105–1111.

https://doi.org/10.1093/bioinformatics/btp120

Trapnell, C., Williams, B.A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M.J.,

Salzberg, S.L., Wold, B.J., Pachter, L., 2010. Transcript assembly and

quantification by RNA-Seq reveals unannotated transcripts and isoform

switching during cell differentiation. Nat. Biotechnol. 28, 511–5.

https://doi.org/10.1038/nbt.1621

Vaser, R., Sovic, I., Nagarajan, N., Sikic, M., 2017. Fast and accurate de novo

genome assembly from long uncorrected reads. Genome Res. 27, 737–

746. https://doi.org/10.1101/068122

Vurture, G.W., Sedlazeck, F.J., Nattestad, M., Schatz, M.C., Gurtowski, J.,

Underwood, C.J., Vurture, G.W., Fang, H., 2017. GenomeScope: fast

reference-free genome profiling from short reads. Bioinformatics 33, 2202–

2204. https://doi.org/10.1093/bioinformatics/btx153

Wajnberg, E., 1993. Genetic variation in sex allocation in a parasitic wasp:

variation in sex pattern within sequences of oviposition. Entomol. Exp.

703   Appl. 69, 221–229. https://doi.org/10.1111/j.1570-7458.1993.tb01745.x

704 Walker, B.J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S.,

705   Cuomo, C.A., Zeng, Q., Wortman, J., Young, S.K., Earl, A.M., 2014. Pilon: An

706   integrated tool for comprehensive microbial variant detection and

707   genome assembly improvement. PLoS One 9.

708   https://doi.org/10.1371/journal.pone.0112963

709 Wallberg, A., Bunikis, I., Pettersson, O.V., Mosbech, M.-B., Childers, A.K., Evans,

710   J.D., Mikheyev, A.S., Robertson, H.M., Robinson, G.E., Webster, M.T., 2019.

711   A hybrid de novo genome assembly of the honeybee, Apis mellifera, with

712   chromosome-length scaffolds. BMC Genomics 20, 275.

713   https://doi.org/10.1186/s12864-019-5642-0

714 Xu, L., Dong, Z., Fang, L., Luo, Y., Wei, Z., Guo, H., Zhang, G., Gu, Y.Q.,

715   Coleman-Derr, D., Xia, Q., Wang, Y., 2019. OrthoVenn2: A web server for

716   genome wide comparison and annotation of orthologous clusters across

717   multiple species. Nucleic Acids Res. 47, W52–W58.

718   https://doi.org/10.1093/nar/gkz333

719 Ye, C., Hill, C.M., Wu, S., Ruan, J., Ma, Z., 2016. DBG2OLC: Efficient assembly of

720   large genomes using long erroneous reads of the third generation

721   sequencing technologies. Sci. Rep. 6, 1–9.

722   https://doi.org/10.1038/srep31900

723 Ye, C., Ma, Z.S., Cannon, C.H., Pop, M., Yu, D.W., 2012. Exploiting sparseness in

724   de novo genome assembly. BMC Bioinformatics 13 Suppl 6, S1.

725   https://doi.org/10.1186/1471-2105-13-S6-S1

726 Zhou, W., Rousset, F., O'Neill, S., 1998. Phylogeny and PCR-based classification

of Wolbachia strains using wsp gene sequences. Proc. R. Soc. B Biol. Sci.

265, 509–515. https://doi.org/10.1098/rspb.1998.0324

Zimmermann, O., 2004. Der Einsatz von Trichogramma-Schlupfwespen in

Deutschland: Zum aktuellen Stand der Forschung und Nutzung von

Eiparasitoiden gegen Schadlepidopteren im biologischen Pflanzen- und

Vorratsschutz. Gesunde Pflanz. 56, 157–166.

https://doi.org/10.1007/s10343-004-0031-1