The evolution of red colour vision is linked to coordinated rhodopsin tuning in lycaenid butterflies

Marjorie A. Liénard[a,b,*], Gary D. Bernard[c], Andrew A. Allen[a], Jean-Marc Lassance[b], Siliang Song[b], Richard Rabideau Childers[b], Nanfang Yu[d], Dajia Ye[b], Adriana Stephenson[b], Wendy Valencia-Montoya[b], Shayla Salzman[b], Melissa R.L. Whitaker[b], Michael Calonje [e], Naomi E. Pierce[b,*]

[a] The Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA.

[b] Department of Organismic and Evolutionary Biology and Museum of Comparative Zoology, Harvard University, Cambridge, MA 02138, USA.

[c] Department of Electrical & Computer Engineering, University of Washington, Seattle, WA, USA.

[d] Department of Applied Physics and Applied Mathematics, Columbia University, New York, NY 10027, USA

[e] Montgomery Botanical Center, 11901 Old Cutter Road, Miami, FL 33156, USA.

Present addresses:

Department of Biology, Lund University, Lund, Sweden (MAL), Department of Ecology and Evolutionary Biology, University of Michigan, 500 S. State Street, Ann Arbor, MI 48109, USA. (SiSo), Department of Biology, University of Pennsylvania, PA 19104, USA (DY); Gene therapy program, University of Pennsylvania, PA 19104, USA (AS); School of Integrative Plant Sciences, Section of Plant Biology, Cornell University, 502 Mann Library, Ithaca, NY 14853, USA (ShSa); Entomological Collection, Department of Environmental Systems Science, ETH Zürich, Zürich, Switzerland (MRLW)

[*]Corresponding authors:

Marjorie A. Liénard, Biology Department, Lund University, Sölvegatan 35B, 22362 Lund, Sweden, marjorie.lienard@biol.lu.se

Naomi E. Pierce, Department of Organismic and Evolutionary Biology, MCZ laboratories 4[th] floor, Harvard University, 26 Oxford Street, Cambridge, MA 02138, USA, npierce@oeb.harvard.edu

ORCIDs in alphabetical order by last name:

GDB: 0000-0001-7460-5123, RRC: 0000-0002-7137-3192, JML: 0000-0002-3675-6956, MAL: 0000-0003-3193-3666, WVM: 0000-0001-9246-2330, NEP: 0000-0003-3366-1625, SSa: 0000-0001-6808-7542 MRLW: 0000-0002-8774-8518.

Classification

Biological Sciences, Evolution, Ecology/Molecular Biology

This PDF file includes:

Main Text, Figures 1 to 7, Supplementary appendix (Figures S1 to S5, datafiles S1-S2)

Abstract

Colour vision is largely mediated by changes in number, expression, and spectral properties of rhodopsins, but the genetic mechanisms underlying adaptive shifts in spectral sensitivity remain largely unexplored. Using *in vivo* photochemistry, optophysiology, and *in vitro* functional assays, we link variation in eye spectral sensitivity at long wavelengths to species-specific absorbance spectra for LW opsins in lycaenid butterflies. In addition to loci specifying an ancestral green-absorbing rhodopsin with maximum spectral sensitivity (λ_{max}) at 520-530 nm in *Callophrys sheridanii* and *Celastrina ladon,* we find a novel form of red-shifted LW rhodopsin at λ_{max} = 565-570 nm in *Arhopala japonica* and *Eumaeus atala.* Furthermore, we show that *Ca. sheridanii* and *Ce. ladon* exhibit a smaller bathochromic shift at BRh2 (480-489 nm), and with the ancestral LW rhodopsin, cannot perceive visible red light beyond 600 nm. In contrast, molecular variation at the LW opsin in *A. japonica* and *E. atala* is coordinated with tuning of the blue opsin that also shifts sensitivity to longer wavelengths enabling colour discrimination up to 617 nm. We then use *E. atala* as a model to examine the interplay between red and blue spectral sensitivity. Owing to blue duplicate expression, the spatial distribution of opsin mRNAs within an ommatidium defines an expanded retinal stochastic mosaic of at least six opsin-based photoreceptor classes. Our mutagenesis *in vitro* assays with BRh1 (λ_{max} = 435 nm) chimeric blue rhodopsins reveal four main residues contributing to the 65 nm bathochromic shift towards BRh2 (λ_{max} = 500 nm). Adaptations in this four-opsin visual system are relevant for discrimination of conspecific reflectance spectra in *E. atala*. Together, these findings illustrate how functional changes at multiple rhodopsins contribute to the evolution of a broader spectral sensitivity and adaptation in visual performance.

Keywords

molecular evolution, ecological adaptation, visual system/vision, rhodopsin, spectral sensitivity, insects, Lepidoptera

Significance Statement

Rhodopsins are photosensitive protein molecules that absorb specific wavelengths of incoming light and convey colour information in the visual system. We show that molecular evolution in a green insect opsin gene resulted in a shift in its maximal absorbance peak, enabling some lycaenid butterflies to use spectral energy of longer wavelengths (LW) to discriminate colours in the red spectrum better than relatives bearing ancestral green LW rhodopsins. Lycaenids also evolved a duplicate blue opsin

gene, and we illustrate an example where species equipped with red LW rhodopsins shifted their blue sensitivity peak to longer wavelengths due to changes in several blue-tuning residues that have evolved repeatedly in different insect lineages. We demonstrate how changes at multiple vision genes in the insect eye effectively create a coordinated mechanism expanding spectral sensitivity for visually guided behaviours such as selecting host plants and mates.

Introduction

Multiple studies have demonstrated the important contributions of gene duplication and protein-coding changes in the evolution of novelty in lineage-specific phenotypic traits (1-3). For instance, gene duplication combined with modifications leading to functional divergence has contributed to gene family expansion, ultimately increasing transcriptional and functional diversity across lineages (2, 4). Recent examples show that specialization of ancestral functions (5-7) or cooperation between encoded products of duplication (8) are common mechanisms resolving constraints in existing eukaryotic gene networks (9). Genomic segmental duplications have also been shown to promote repeated allelic fixation, linking adaptive duplicated loci to convergent evolutionary pathways (10).

However, structural, functional, or gene network constraints can impose evolutionary trade-offs if there is limited variation for alternative biochemically stable encoded products among duplicated loci within multigene families (9, 11). Characterizing the molecular patterns of evolution within multigene families, and identifying the functional role of mutations at lower taxonomic scales is needed to distinguish the multiple sources of molecular variation. This includes the relative roles of gene duplication, expression changes, protein sequence divergence and convergence, or cellular relocalization of encoded gene products that contribute to new or convergent phenotypes within lineages (2, 7, 8, 12, 13).

Opsins belong to a diverse multigene family of G protein-coupled receptors, offering a robust framework to study how molecular changes can ultimately cause changes in behaviour and favour diversification (14). Opsins bind to a small non-protein retinal moiety derived from vitamin A to form photosensitive rhodopsins and enable vision across animals (15-18). The evolution of vision across the animal kingdom has long been linked to independent opsin gene gains and losses (19-23), genetic variation across opsins (15, 24-26), and spectral tuning mutations within opsins (27, 28). These molecular mechanisms , together with alterations in visual regulatory networks (29), have been shown to contribute to rhodopsin adaptation and the diversification of spectral sensitivity phenotypes in insects (18, 30, 31) and vertebrates including fish, birds, bats and primates (32-36). Thus multiple levels of organization, including the evolutionary

74 diversification of opsin subfamilies, their functional properties and regulatory networks
75 constitute the mechanistic basis for organisms to discriminate light sources of varying
76 wavelengths and ultimately interpret them as colours, yet how multiple changes across
77 opsin paralog repertoires impact spectral sensitivity at small taxonomic scales has been
78 understudied.

79  The evolution of insect colour vision in particular shows how a complex sensory
80 trait can play a central role in adaptations involving signalling and mate communication
81 (18, 31, 37, 38). Whereas the ancestral repertoire of insects involved three types of light-
82 absorbing rhodopsin genes: ultraviolet (UVRh, 350nm), blue (BRh, 440 nm) and green,
83 also called long-wavelength (LWRh, 530nm) (18), today's genomes harbor smaller or
84 larger rhodopsin repertoires with strong experimental evidence for repeated functional
85 convergence toward UV, Blue and LW spectral sensitivities across lineages. For
86 example, beetles lost their ancestral blue opsin gene 300 million years ago, and
87 compensated for the loss of blue sensitivity via either UV or LW gene duplication across
88 lineages (23, 39). Blue opsin duplications occurred independently in pierid and lycaenid
89 butterflies (27, 30, 40-42); and extend photosensitivity into the UV/blue in *Heliconius*
90 spp. with λ_{max} = 355 nm and 398 nm (21) and into the violet/blue, in *Pieris rapae* with λ_{max}
91 = 420 and 450 nm (27). UV and LW duplications occurred in butterflies, hemipterans and
92 dragonflies (20, 22, 26, 30, 43-45). In butterflies, LW opsin duplications have been
93 identified in two papilionids, *Papilio xuthus* (31) and *Graphium sarpedon* (46) as well as
94 a riodinid *(Apodemia mormo)* (20, 47), and contribute to extend spectral sensitivity into
95 the far red.

96  Although duplicated LW opsins have never been detected in the families
97 Nymphalidae, Pieridae or Lycaenidae (25, 40, 41), photoreceptor types with sensitivity
98 peaks in the red have been identified in these groups (40, 41). This supports the
99 hypothesis that additional mechanisms such as lateral filtering and/ or molecular
100 variation of ancestral LW opsin genes also contribute to modify long-wavelength
101 sensitivity.

102  For example, photostable lateral filtering pigments are relatively widespread
103 across butterfly lineages (e.g. *Heliconius* (21)*, Papilio (45), Pieris* (48), *Colias erato* (49),
104 and some moths (*Adoxophyes orana,* (50)*; Paysandisia archon* (51)). These pigments

act as long-pass filters, absorbing short wavelengths and pushing the sensitivity peak of

LW photoreceptors into the red, to create distinct spectral types that can contribute to

colour vision (19, 31, 40, 48, 52, 53). Lateral filtering can shift peak sensitivity to longer

wavelengths while reducing peak amplitude, but cannot *extend* photoreceptor sensitivity

into the far red beyond the exponentially decaying long-wavelength rhodopsin

absorbance spectrum (52). Molecular variation of ancestral LW opsin genes could

potentially extend photoreceptor sensitivity, but this mechanism has remained difficult to

disentangle from the effects of filtering granules using classical electrophysiological

approaches (31, 53, 54).

The molecular underpinnings of mammalian MWS/ LWS spectral tuning has

been studied using mutagenesis experiments and led to the identification of critical

amino acid replacements and their interactions at key residues (55, 56). Most

mammalian lineages possess long-wavelength sensitive (LWS) cone opsins that specify

Medium (M, λ_{max} 510-540 nm) and Long (L, λ_{max} > 540 nm) rhodopsins. In humans,

trichromatic vision is conferred through the use of Short (S, λ_{max} = 414 nm), and tandem

duplicate M (λ_{max} = 530 nm) and L (λ_{max} = 560 nm) cone rhodopsins (34), that collectively

allow us to discriminate longer wavelengths of light as green-red colours (57, 58). In

birds, tetrachromatic vision is based on two SWS cone opsins, together with a green M

opsin (λ_{max} = 497-514 nm) and a red-sensitive LWS opsin (λ_{max} = 543-571 nm) (reviewed

in 59). In insects, the evolution of red receptors occurred independently multiple times

(30) but the possible contribution of molecular variation to insect LW opsin gene

diversification has remained elusive, notably due to difficulties in expressing LW opsins

in vitro (27, 60, 61, but see 62). Red receptors are intriguingly very common in butterflies

compared to other insect groups such as bees or beetles (30), raising the possibility that

perception of longer wavelengths plays an important role in the context of foraging (31,

46, 63), oviposition (64, 65) and mate recognition (25) for species equipped with them.

Lycaenids comprise the second largest family of butterflies, representing almost

thirty percent of all species, and exhibiting considerable ecological and morphological

diversity (66, 67). Pioneering work showed that species of Lycaenidae in the genera

Lycaena and *Polyommatus* have expanded spectral sensitivity at long wavelengths, and

this has been postulated to arise from filtering pigments, modified opsins or both (19, 40,

41). Here, by combining physiological, molecular and functional approaches, we identify

additional lycaenid species with red photoreceptors and elevated spectral sensitivity at

long wavelengths, and show that their LW opsin locus specifies a novel type of LW

rhodopsin with red-shifted maximal absorbance. We focus on the Atala hairstreak

(*Eumaeus atala*) as a suitable model to show the interplay between regulatory and

adaptive changes at multiple opsin loci in the evolution of red spectral sensitivity and link

the evolution of finely tuned four-opsin vision to relevant visually guided behaviours in

these butterflies.

Results

Hairstreak butterflies with elevated sensitivity at long wavelengths express red-shifted rhodopsin receptors

A photographic series at distinct elevations in the eye revealed dramatic differences in eyeshine colouration between two pairs of lycaenids (Fig. 1 insets and Fig. S1). The Spring Azure, *Celastrina ladon* (subfamily Polyommatinae) and Sheridan's green hairstreak, *Callophrys sheridanii* (subfamily Theclinae), present homogeneous green and orange eyeshine in their dorsal eyes at 30° elevation, with a relatively small number of red ommatidia (*Ce. ladon*, 5%; *Ca. sheridanii*, 16%) (Fig. S1A-B). By contrast, the Japanese Oakblue, *Arhopala japonica,* and the Atala Hairstreak, *Eumaeus atala* (both subfamily Theclinae), share high levels of saturated red eyeshine in the dorso-equatorial region, due at least in part to species-specific differences in rhodopsin content (Fig. 1, Fig. S1B).

We partially bleached eyeshine of *A. japonica* and *E. atala* using repeated white-light flashes to reveal two types of ommatidia. Some ommatidia were resistant to bleaching and maintained their red eyeshine owing to lateral filtering by cherry-red pigment granules located distally within the photoreceptor cells of those ommatidia. The rest of the ommatidia were bleached, suggesting the presence of red sensitive opsin-based photoreceptors in these two species (Fig. S1C).

To test for the presence of red sensitive receptors, we performed analyses of *in vivo* photochemical rhodopsin bleaching measurements of adult butterflies. These experiments revealed long wavelength (LW) spectral sensitivities in dark-adapted eyes of *A. japonica* with λ_{max} ± standard error at 571 ± 2.45 nm ($CI_{95\%}$ = 566 to 576 nm) (Fig. 1A) and *E. atala* with λ_{max} at 563 nm ± 0.9 nm ($CI_{95\%}$ = 561 to 566 nm), respectively (Fig. 1B). We found that the two other lycaenid species we studied, *Ca. sheridanii* and *Ce. ladon,* were difficult subjects for our *in vivo* methods. In *Ca. sheridanii,* a noisy difference spectra obtained by photochemistry provided an estimate at LW λ_{max} = 518 nm ± 3.7 nm for ($CI_{95\%}$ = 511 to 526 nm) (Fig. 1C). Similarly, in *Ce. ladon,* partial LW bleaches were not measurable due to low LW rhodopsin densities,.

We chose to examine in detail the *in vivo* and *in vitro* contributions of all
rhodopsin pigments in *E. atala*, a multi-brooded and abundant hairstreak butterfly
naturally occurring throughout the year in Florida (USA) (68). Unlike most lycaenids, *E.*
atala larvae are extreme specialist herbivores on New World cycads in the genus *Zamia*
and show an unusually bright aposematic colouration advertising toxins they sequester
from their hosts (69). Conversely, the adult is a striking velvety black butterfly with bright
blue and red colours on the wings and the abdomen (Fig. 1) and can collect nectar from
43 species across 20 plant families (69)

We first analyzed epi-microspectrophotometric difference spectra obtained after
intense ommatidial flashing from a series of interference filters, which in addition to
identifying R565 (LW rhodopsin, Fig. 1B) allowed us to narrow down a blue rhodopsin at
R440 (Fig. 2A). Optophysiology measurements confirmed that neither the R565 LW nor
the R440 blue sensitive rhodopsins could be responsible for the high sensitivity around
360 nm, which instead was due to UV-receptors   (Fig. 2B, Table S2). Finally, a
densitometric analysis of male and female eyes (Fig 2C-D, Tables S3-S4), showed that
these physiological data are best fit using least-squares regression by a model in which
four rhodopsins are present in the eye with λ_{max} values matching the LWRh rhodopsin as
well as UVRh 360 nm, BRh1 441.3 nm ± 4.7 nm ($CI_{95\%}$ = 431.7 to 450.9 nm, Table S2)
and a second blue rhodopsin BRh2 494.2 ± 1.2 nm ($CI_{95\%}$ = 492 to 497 nm) (Fig. 2E-F,
Tables S3-4).

Red sensitivity is due to functional variation at long-wavelength opsin loci

Transcriptomic mRNA profiling of *E. atala* yielded only a single LW opsin (Table
S6), which is in line with earlier molecular evidence showing that lycaenid species
possess one LW opsin gene (30, 40, 53). Accordingly, eye cDNA library screening using
degenerated oligonucleotide primers followed by RACE cDNA amplification led to the
characterization of single orthologous LW cDNA sequences in *Ce. ladon*, *Ca. sheridanii*
and *A. japonica* (Fig. 3). We optimized an *in vitro* HEKT293 cell culture assay to
reconstitute heterologous rhodopsin pigments (27, 61, 70, 71) by using a newly
engineered heterologous vector to increase expression levels, and improving purification

procedures to obtain higher yields of actively reconstituted LW rhodopsins. Our
expression cassette derives from the pcDNA5 plasmid and expresses coding opsin
sequences tagged with a C-terminal FLAG epitope under a strong CMV promoter.
Directly downstream the FLAG epitope, a short peptide linker plus a T2A cleavage
cassette enable co-transcription of a fused cytoplasmic fluorescent mRuby2 coding
sequence (Fig. 3A). This expression cassette proved to be efficient at detecting
monomeric units of ultraviolet, blue and long-wavelength (LW) heterologous *E. atala*
opsin proteins *in vitro* (Fig. S2A). It is also successful at reconstituting and purifying
active LW rhodopsins, *i.e.* in *A. japonica, E. atala, Ca. sheridanii* and *Ce. ladon* lycaenid
butterflies (Fig. 3B-F, Fig. S2B, Table S5).

When purified from large-scale HEK293T cell cultures and reconstituted *in vitro* in
the dark in the presence of 11-*cis*-retinal, we found that the LW rhodopsin from *A.*
japonica absorbs maximally at λ_{max}= 574 ± 4 nm ($CI_{95\%}$ = 570-586 nm) (Fig. 3C) whereas
that of *E. atala* absorbs maximally at λ_{max}= 569 ± 2 nm ($CI_{95\%}$ = 565-573 nm) (Fig. 3D).
The peak of absorbance of purified LW rhodopsin measurements is within confidence
intervals of the best fit for LW linear absorbance estimates *in vivo* and supports the
hypothesis that the LW rhodopsin limb of absorbance in these two species pushes red
sensitivity above 600 nm. The LW rhodopsin pigment from *Ca. sheridanii* absorbs
maximally at λ_{max} = 519.2 nm ± 1.1 ($CI_{95\%}$ = 517-521 nm) (Fig. 3E), which is in
accordance with photochemical measurements (Fig. 1C), whereas *Ce. ladon* LW
rhodopsin absorbs maximally at λ_{max}= 531.7 ± 1.5 nm ($CI_{95\%}$ = 529-535 nm) (Fig. 3F,
Table S5).

In summary, these findings indicate that our expression system can be used
successfully to assess the functionality of LW rhodopsins outside of the complex eye
environment and demonstrate that at least some hairstreaks (Theclinae) express a new
functional type of far-red shifted visual opsin.

Blue opsin mRNA expression patterns lead to enhanced eye spectral richness

Following photochemical and densitometric evidence that *E. atala* butterflies likely
possess four rhodopsins including contributions from two blue rhodopsins, we identified

two differentially expressed blue opsin-like transcripts from the eye transcriptome,
namely BRh1 and BRh2 (Table S6). We then examined their respective mRNA
expression patterns in photoreceptor cells, aided by the histological reconstruction of
photoreceptor organization in a typical ommatidium (Fig. 4).

We found that *E. atala* hairstreaks have a straight, 480-micron long rhabdom
composed of eight longitudinal photoreceptor cells (Fig. 4A-B (a-g)) and a ninth cell
close to the basement membrane (Fig. 4A-B (h)). The two most distal R1-R2
photoreceptor cells contribute the majority of microvillar extensions from 0 to 160
microns, whereas R3-R4 distal cells contribute a majority of microvilli from 140 to 300
microns, thereby overlapping partially with R1-R2 in the distal rhabdom tier. The
proximal R5-R8 cells contribute most microvilli in the last rhabdom tier up to 440
microns, a depth where the photoreceptor cells no longer bear any microvilli and the
ninth cell becomes visible (Fig. 4A-B (i)).

This rhabdomeric analysis provided the spatial morphological insights necessary
to subsequently identify photoreceptor cells expressing each opsin mRNA across
different eye regions. Using double fluorescent *in situ* hybridization experiments in
transverse and longitudinal eye sections of males and females of *E. atala*, we showed
that LWRh is expressed in all ommatidia in the six photoreceptor cells R3 to R8 (Fig. 5),
which is typical of many butterfly species (25, 29). No fluorescent signal was detected for
the long-wavelength opsin in R1-R2 cells (Fig. 5D-F). We next examined the cellular
localization of the short UV and medium blue opsin mRNAs in transverse sections in the
dorsal eye. Using probes targeting LWRh in combination with cRNA probes for UVRh,
BRh1 or BRh2 mRNAs, our data provide evidence that the latter rhodopsin mRNAs can
be expressed in either or both R1-R2 receptor cells forming single ommatidia (Fig 5A-F).

We also assessed the possibility that these opsins are co-expressed in R1 and
R2 cells using cRNA probes for UVRh and BRh1 (Fig. 5G), UVRh and BRh2 (Fig. 5H) or
BRh1 and BRh2 (Fig. 5I). We find that R1 and R2 follow a one-cell one-opsin regulation
pattern, with mutually exclusive expression of UVRh, BRh1 and BRh2 mRNAs (Fig. 5J).
Females exhibit a dorso-ventral expression gradient in which BRh2-expressing cells are
sparse in the dorsal eye region. In the ventral retina however, all three opsins (UVRh,
BRh1, BRh2) are found in addition to LWRh (R3-R8) (Fig. S3).

Cellular expression data therefore show that both of the duplicated blue opsin
mRNAs encode functional transcripts. Overall, the cellular localization of blue opsin
duplicates in adjacent photoreceptor cells creates multiple ways in which rhodopsins are
distributed within individual ommatidia, forming a local stochastic rhodopsin mosaic of at
least six opsin-based photoreceptor classes (UV-UV, B1-B1, B2-B2, UV-B1, UV-B2, B1-
B2, Fig. 5j). Additional structural features of the eye may also contribute to the diversity
of spectral sensitivity functions of individual photoreceptors, including the spectral
influence from lateral filtering granules found in distal rhabdomeres that add to the
spectral sensitivity of some photoreceptor cells.

BRh1 and Brh2 opsin duplicate loci encode blue and green-shifted rhodopsins

We reconstituted active BRh1 and BRh2 rhodopsins *in vitro* and measured their
spectral sensitivities via spectroscopy as detailed in the methods. We determined that
BRh1 λ_{max} = 435 nm ± 2 nm and BRh2 λ_{max} = 500 nm ± 2 nm (Fig. 6A-B, Table S7),
thereby confirming that they encode the rhodopsins conferring the expanded blue to
green spectral sensitivity in *E. atala*. In addition to its LWRh red-shifted opsin (Fig. 3D),
E. atala uses a fourth rhodopsin that absorbs UV wavelengths *in vitro* at λ_{max}= 352 nm ±
3.5 nm ($CI_{95\%}$ = 345-360 nm) (Fig. 3, Fig. S2d, Table S7).

Four spectral tuning sites and epistasis leads to bathochromic shifts between
blue duplicates

In order to understand the proximate mechanisms driving the 65 nm spectral shift
between blue opsin duplicates, we performed homology modeling against the
invertebrate squid rhodopsin (72). Along with their 380-aa long opsin protein sequences,
Eat-BRh1 and Eat-BRh2 exhibit 101 amino acid residue differences (corresponding to
73% similarity in aa sequence). Six variant sites were identified among 21 homologous
residues located within 5Å of any carbon forming the *cis*-retinal binding pocket (Fig.
6A,C). Two variant sites (S1 and S4) are shared with a blue/violet opsin duplication that
occurred independently in pierid butterflies, causing a UV-shift (VRh λ_{max} = 420 nm) from

the ancestral blue rhodopsin (BRh λ_{max} = 450 nm) (27), whereas four residues are

unique to duplicated blue opsins found in lycaenids (S2, S3,S5,S6).

Given this combination of shared and unique yet limited number of variant sites,

we decided to test their possible involvement in blue spectral tuning by site-targeted

mutagenesis. Specifically, we first modified individual residues S1 to S4 and measured

their spectral tuning effect *in vitro* (Fig. 6E-G, Fig. S4). Chimeric opsins showing

bathochromic shifts were used sequentially to substitute adjacent variant sites following

two candidate evolutionary trajectories. We observed an additive bathochromic shift

totaling 40 nm (λ_{max} = 475 nm) by substituting A116S together with I120F and Y177F

(S1, S2 and S4, Fig. 6E). In a second tuning trajectory, we observed that Y177F alone

conferred a 79 nm bathochromic shift (λ_{max} = 514 nm), which could then be

compensated by a 6 nm hypsochromic shift (λ_{max} = 508 nm) by combining G175S and

A116S (Fig. 6G). A third evolutionary trajectory explored the contribution of two lycaenid-

specific cysteine substitutions (I106C, F207C) in helix 5 (Fig. S4). These changes

caused a strong green-wave spectral shift that was not compensated for by additional

candidate interacting residues in the quintuple mutant (S1,2,4,5,6; Fig. S5, Table S8).

G165S (S3) tested alone or in various double mutant combinations caused

bathochromic shifts, but we did not obtain the sextuple chimeric construct bearing

G175S (S3) and cannot therefore exclude the possibility that it also plays a role in this

case. The available results from all variants bearing cysteines in S5 and S6, however,

suggest that both cysteine residues on helix 5, which are absent in Pieridae blue opsins,

are unlikely to be the main evolutionary drivers of blue spectral shifts in lycaenid BRh2

loci.

Finally, our optophysiological density analyses showed that blue spectral

sensitivity differs between species equipped with green or red LW rhodopsins. Whereas

all species possess a blue photoreceptor with conserved absorbance with λ_{max} = 435-

440 nm, the second blue sensitivity peak in *A. japonica* and *E. atala* is at λ_{max} = 500 nm

(Fig. 2F, S5A) unlike λ_{max} = 489 in *C. ladon* (Fig. S6B, Table S9). These data suggest a

role for coordinated shifts between blue and LW rhodopsins in the evolution of red

spectral sensitivity.

Shifted Blue and LW rhodopsins tune colour vision in the context of conspecific
recognition

In order to investigate possible correspondence between *E. atala* visual spectral
sensitivities and colour traits that might be important in signalling and sexual selection,
we measured the reflectance spectra associated with specific wing patches (Fig. 7, SI
Dataset 2). For the butterfly to interpret colours, it must i) possess at least two spectral
types of receptors sensitive to the reflectance spectrum of incident visible light
illuminating the coloured area, and ii) be able to compare individual receptor responses
neurally to create an output chromatic signal (73). Although we did not take this analysis
to the second step of recording responses of different individual receptors to reflectance
spectra, we investigated in detail the first necessary condition by measuring and
comparing reflectance spectra of body and wing patches from males and females
including blue scales on the abdomen and thorax, as well as black, blue and red scales
on forewings and hindwings (Fig. 7A). In males, dorsal forewings are bright iridescent
blue in summer, whereas scales appear more generally green/ teal in winter generations
(68). Female dorsal wings, on the other hand, display a darker royal blue colour along
the edge of their upper forewings. Both sexes also have conserved wing and body
patterns, including regularly spaced rows of blue spots visible on folded and unfolded
hindwings, and a bright red abdomen with a large bright red spot on the mid-caudal
hindwing area that falls precisely along the abdomen in folded wings (Fig. 1A, inset).

First, epi-microspectrophotometry measurements from dark areas of male and
female wings (Fig. 7B) showed that black scales are only 1 % as reflective as adjacent
cyan scales in the blue/green band at 450 nm, with a reflectance maximum of 0.053
(female, *f*) and 0.041 (male, *m*) (SI datafile 2) compared to reflectance maxima at 510
nm of ventral hindwing blue scales 1.53 (*f*) and 1.76 (*m*) (Fig. 7B). This indicates that
brightness – which is a function of reflectance - in black regions decreases 100 fold
compared to adjacent coloured scales in both sexes.

Blue scales on the dorsal forewings have a maximal reflectance peak at 490 nm
in females and 510 nm in males, and blue scales on the ventral hindwings have a

maximal reflectance peak at 510 nm (*f*) and 530 nm (*m*) (Fig. 7B). Thorax scale reflectance is 1.86 times and 2.88 times higher than dorsal wing reflectance in females and males, respectively.

Red scales on male and female hindwings reflect maximally in the far red (750 nm), similarly to abdominal scales (Fig. 7B-C). Blue scales on the thorax have a maximal reflectance peak at 490 nm in both sexes, overall indicating that male and female reflectance spectra are similar for blue scales on the body, but not for blue scales on the wings. We noted, too, that leaf surfaces of the butterfly's primary host cycad, *Zamia integrifolia*, have a peak of reflectance at 550 nm and a red edge inflection point around 700 nm with high reflectance in the near infrared region (Fig. 7D). Altogether, our analyses support that the butterfly's photoreceptor spectral sensitivities (Fig. 7E) can efficiently discriminate between its host plant (for oviposition), the colours of male and female conspecifics, and colour variation between sexes.

Discussion

Red opsin receptors contribute to far-red spectral sensitivity

It has remained challenging to identify the contribution of different genetic mechanisms that affect phenotypic variation amongst complex traits in nature, and ultimately an organism's fitness. Opsins represent a robust system to link molecular changes to phenotypic changes in animal colour vision as they are the first elements in phototransduction cascade and have been shown to directly modulate the visual spectral sensitivity of insects, primates and other vertebrates. .We have optimized a functional assay to disentangle the contribution of the long-wavelength (LW) rhodopsin from filtering pigment granules and variable eye reflectance properties in the eyes of lycaenid butterflies. We discovered a novel functional form of LW rhodopsin with a red-shifted maximal absorbance spectrum (λ_{max}) between 565 and 575 nm in two lycaenid species, *A. japonica* and *E. atala,* that is further supported by microspectrophotometry (MSP) and optophysiology data (Figs. 1-3, Fig S5). The red-shifted rhodopsins have a longer-wavelength limb of absorbance than lycaenid green LW rhodopsins due to a bathochromic shift in their λ_{max} (Figs. 1, 3), which elevates the photoreceptor response at long wavelengths (Fig. 2F, S5A). Consequently, those two species have much greater optophysiological sensitivity in the far red than the lycaenid species with photoreceptors expressing green LW rhodopsins, whether or not surrounded by lateral filtering pigments.

Red spectral sensitivity has previously been examined *in vivo* notably in a species of cabbage white butterfly, *Pieris rapae*, where light perception above 600 nm results from ommatidia expressing a single LW opsin together with two types of pigment granules that confer three types of red photoreceptors (48). Methods employing cAMP-dependent heterologous spectroscopy identified maximal peak absorptions at 540 nm and 560nm for *Papilio* PxRh1 and PxRh3 LW opsins respectively (62), whereas earlier electrophysiological λ_{max} estimates placed PxRh1 at 525 nm and PxRh3 at 575 nm (45). Our assay provides quantifiable expression and yields to obtain accurate λ_{max} *in vitro* outside of the complexity of the eye structure itself which circumvents potential inaccuracy from optophysiological estimates measured *in vivo* in certain species due to interfering lateral pigments or other properties of the eye itself. It also has the advantage

of functionally studying variation in orthologous rhodopsin genes independently from inferences based on sequence data alone. Our study reconciles *in vivo* and *in vitro* approaches to substantiate that higher visual performance in the far red is achieved by modifying absorbance properties of long-wavelength rhodopsins.

Duplication, spectral tuning and adjacent cellular localization provide expanded photoreceptor types

Opsin evolution has undergone recurrent events of gene duplication and loss across animals including insects (16). From ancestral insect trichromatic colour vision (UV, blue, green), insects that have lost one of these three rhodopsins are often found to have compensated from that loss by recruiting a duplicate gene copy that has undergone spectral tuning to shift the peak sensitivity, as seen in true bugs and beetles (23, 39). By contrast, tetrachromatic insects that activate an additional rhodopsin are also known to have acquired increased spectral sensitivity in ranges of visible light, as seen in *Heliconius* duplicated UV opsins (21), or *Pieris* duplicated blue opsins (27). Lycaenid species examined thus far have two blue opsins for which epi-microspectrophotometric estimates previously yielded sensitivities of λ_{max} in the range 420-440 nm and 480-500 nm (40, 41).

Our results show that blue opsin loci in *E. atala* specify blue and green-shifted rhodopsins (Figs 2,6) similarly to species of *Lycaena* and *Polyommatus* (40, 41). The most striking functional insight in the evolution of blue spectral tuning in *E. atala* comes from chimeric BRh1 variants bearing mutations A116S, G175S and Y177F, which confer a 73-nm bathochromic shift (λ_{max} = 508 nm) that most closely recapitulates the spectral properties of Eat-BRh2 (λ_{max} = 500 nm) compared to other tested variants (Fig. 5G, Fig S4). Intermediate adaptive phenotypes can also be revealed via gradual evolutionary trajectories. Eat-BRh1 variant A116S causes a +5 nm shift alone, but together with I120F, shifts maximal absorbance by an additional +15 nm to λ_{max} = 455 nm. Since these two residue substitutions are conserved across all characterized lycaenid BRh1 loci (Fig. 3A), these results show how adjacent sites in helix 3 can contribute to intermediate blue absorbance spectra across lycaenids.

The third tuning residue, Tyr177Phe is a key spectral tuning mutation in *E. atala,*
since the triple BRh1 variant (A116S/I120F/Y177F) displays a 30 nm bathochromic shift
(λ_{max} = 475 nm) compared to its native rhodopsin, and illustrates the multiple ways that
gradual spectral tuning can evolve, at least in this species. Two of the reverse tuning
substitutions are the same sites responsible for hypsochromic spectral shifts both in a
blue-shifted LW rhodopsin of a *Limenitis* butterfly (Y177F, -5 nm) (61) and in a violet-
shifted blue rhodopsin of a *Pieris* butterfly (S116A, -13 nm; F177Y -4 nm) (27), stressing
the importance of tuning residues lying on the ionone ring portion of the chromophore
binding pocket. Spectral tuning modulation in blue-rhodopsin duplicate functions has
therefore involved conserved biochemical constraints along independent evolutionary
trajectories that selected for partial spectral tuning sites in rhodopsin G-coupled
receptors. Reverse mutations are not functionally equivalent in their absolute
magnitudes ($\Delta\lambda_{max}$), therefore underscoring the role of epistatic interactions with
neighbouring sites resulting in distinct λ_{max} shifts across butterfly lineages.

Molecular and phylogenetic studies of opsin evolution in vertebrates have also
shown that homologous tyrosine residues in the ionone ring portion of the chromophore-
binding pocket (e.g. Y262) in the human blue cone opsin (λ_{max} = 414 nm) are responsible
for a 10 nm bathochromic spectral tuning when mutated to Tryptophan (Trp) (74). This
illustrates that distant opsin loci meet similar structural and biochemical constraints as
those observed in the evolution of vision genes in other vertebrates (11, 14). However,
Y177 is unique to *E. atala*, and other BRh1 loci at this position readily possess the F177
similarly to Eat-BRh2 (Fig. 3A). Not all BRh2 loci have S175 but instead keep G175 in
both blue loci, suggesting an additional yet unknown role for adjacent residues in
positions R176K and I178V. The latter residues do not differ highly in hydrophobicity but
may still provide the necessary molecular interactions with the ionine ring portion of the
chromophore necessary to modulate variable BRh2 spectral phenotypes. Human
ancestors also achieved blue sensitivity gradually and almost exclusively via epistasis at
seven amino acid residues (55). By studying independent opsin gene duplicates in the
butterfly family Lycaenidae, our *in vitro* assays refine our understanding of the molecular
basis of convergent colour vision phenotypes and help to identify key determinants of
genotype-phenotype relationships across insect blue opsins, although these are only a
snapshot of all possible chimeric blue rhodopsin variants along at least three possible

evolutionary trajectories. Testing the non-additive interactions at co-evolving BRh2
adjacent sites (175-178) in lycaenids lacking red sensitivity will help to fully recapitulate
intermediate phenotypes across derived blue-shifted rhodopsin duplicates in addition to
those generated by A116S and I120F.

In theory, a typical tetrachromat can achieve better wavelength discrimination than a
typical trichromat (55, 73) because of the interplay between additional gene copies and
the coevolution of spectral tuning across rhodopsins decreases the minimal wavelength
difference ($\Delta\lambda$) that an insect can discriminate. *Papilio* butterflies, for instance, have
some of the most complex retinal mosaics known, with three ommatidial types
expressing various combinations of five rhodopsin proteins (UV, B, 3 LW), as well as
filtering pigments that produce diverse ommatidial spectral sensitivities (29, 45). Amino
acid residues in helix 3 have been shown to mediate an absorbance shift between
duplicates LW PxRh1 and PxRh3 in *Papilio (62)*, but lycaenid LW opsins possess a
highly conserved helix 3 that does not exhibit any of the spectral substitutions found
across Papilionidae, supporting the hypothesis that distinct spectral tuning mechanisms
have evolved independently in invertebrate LW opsins. Future work will be able to
disentangle the spectral tuning mechanisms of LW opsins *in vitro* using the recombinant
purification system described here.

In *E. atala*, ommatidial density and expression profiling showed that
photoreceptor cells containing BRh2 are less abundant (~25%) compared to those
containing UV (~50%) and BRh1 (75%) rhodopsins. In spite of its low density across the
eye, the derived green-shifted blue rhodopsin improves signal resolution in the green
spectrum and provides direct adaptive benefits for light perception in both males and
females. Compared to trichromats, the expanded six-ommatidial types primarily derive
from the non-overlapping cellular expression of BRh1 and BRh2 rhodopsin mRNAs in
distal rhabdomeric cells. Together, the stochastic rhodopsin mosaic offers remarkable
properties that could not be achieved with the limited sensitivity of a single blue opsin
gene.

In summary, our results support the scenario of a sensory system in which gene duplication generates new opsin paralogs, cis-regulatory changes generate new photoreceptor subtypes via gene-specific expression patterns, and coding mutations tune the spectral sensitivity maxima of two of the four rhodopsins (BRh2, LWRh). All of these changes have contributed to the diversification of visual spectral sensitivities, thereby enhancing colour discrimination.

Co-evolving shifts at blue and red opsins tune intraspecific visual signalling

Additional structural features of the eye can contribute to spectral sensitivity of individual photoreceptors at long wavelengths, including lateral filtering granules found in distal rhabdomeres. However, here the primary mechanism pushing red sensitivity is through functional divergence at the LW rhodopsin locus. This remarkable novelty provides a two-factor interaction (Blue-LW) to modulate green-red sensitivity among closely related taxa.

Building on earlier butterfly microspectrophotometry work (19, 40, 41, 53), our study shows that lycaenid species with red-shifted LW opsins have their second maximum sensitivity peak and associated blue rhodopsin shifted to green maximal absorbance with λ_{max} 495-500 nm. By contrast species with a LW ancestral-green opsin with λ_{max} 520-530 nm have their second maximum sensitivity peak at blue wavelengths λ_{max} = 480-490 nm (Fig. 1C, S6A), suggesting modulatory benefits for coevolving spectral shifts at multiple rhodopsins. Within Lycaenidae, reflectance patterns of wing scales have been investigated in the genera *Celastrina* and *Callophrys* (75, 76) and these studies support a scenario of co-evolution of spectral shifts across opsins. The ventral wings visible at rest in Green Hairstreak lycaenids (genus *Callophrys*) typically appear green and consist of yellow and bluish scales with an omnidirectional reflectance peak in the green slightly above 550 nm (76). *Celastrina* species tend to display bright blue coloured wings, which predominantly reflect light in the UV-blue band (75) and can thus be perceived by comparing neuronally the spectral input from light activating retinal photoreceptors expressing both UV/ blue rhodopsins without requiring expanded red sensitivity.

These observations suggest a role for molecular tuning of blue and LW opsin
genes in driving the dynamic evolution of green-red spectral sensitivity. Extant human
trichromatic colour vision is similarly thought to have evolved via spectral tuning of
existing short wavelength-sensitive opsins in concert with molecular variation at
duplicated opsin loci leading to middle and long-wavelength sensitivities (55). Ultimately,
mechanistic and functional studies of spectral tuning evolution in different orthologs in
additional species will enable us to infer the ancestral states and better resolve the
evolutionary time scale involved in stages of spectral tuning of blue and red sensitivities
across lycaenids.

Colour tuning improves intraspecific signal detection

Two effects of molecular variation in visual pigments in *E. atala* are that its eyes have
expanded spectral sensitivity at longer wavelengths as well as increased spectral
discrimination in the blue band. In order to investigate the evolutionary consequences of
molecular changes in butterfly opsins in the broader context of behaviours requiring
colour vision, such as finding oviposition sites, and intraspecific recognition (37, 77-80),
we need to analyze the degree of overlap of spectral sensitivities (73) and evaluate this
against the spectral composition of the reflected light of the background and other
objects, such as that produced by foliage and conspecifics.

The limiting wavelength at which only the human L cone opsin (λ_{max} = 559-563
nm) is sensitive corresponds to approximately 625 nm (81), which is the wavelength at
which orange-red can be discriminated from pure red in human colour perception. The
human neighboring M cone rhodopsin is at λ_{max} = 530 nm, which at 625 nm, has a
sensitivity of 0.05 (81). By extension, for *E. atala*, BRh2 has a sensitivity of 0.05 at 613
nm (Fig. 2, Fig. 6), indicating that R565 would perceive 'pure red' wavelengths of light
from 613 nm onwards. Alternatively, we can compute the relative sensitivity of the LW
rhodopsin at 700nm (3%, Fig. 1, Fig. 3), which is generally accepted as the limit for
photopic vision, and then identify the corresponding wavelength at which BRh2 reaches
that same sensitivity, which is 619 nm (Fig. 2). Conservatively, we can conclude that the
red rhodopsin together with the green-shifted blue rhodopsin BRh2 of *E. atala* contribute

to the perception of colours at long-wavelengths up to 613 nm. The LW rhodopsin sensitivity in *A. japonica* is similar to *E. atala*, and the green-red visual spectral sensitivities of *Arhopala* butterflies would capture the dorsal wing secondary reflectance peak between 500-600 nm, which is absent in *Celastrina* or *Callophrys* wings (75). Dorsal *Arhopala* wings also have velvety black edges filled with blue regions that primarily reflect light at 370 nm in males and 400 nm in females (82) and further benefit from the expanded blue sensitivity seen across lycaenid butterflies. Similarly to super-black plumage colouration in birds (83), peacock spiders (84), and recent examples in papilionids and nymphalid butterflies (85), the adjacency and sex-specific regions covered with structural velvety black scales in the lycaenid *E. atala* likely enhance the perceived brightness of nearby colour signals.

Our wing scale analysis indicates that red colour discrimination is important to see conspecifics due to their striking reflectance above 600 nm (Fig. 7A). Reflectance spectra from the red abdomen and the ventral hindwing are similar and overlap spectral sensitivities of two rhodopsins (LWRh and BRh2), meaning the butterfly can not only detect brightness but can also derive a colour signal from these patterns, and separate these input signals from those of environmental colour cues such as signals produced by foliage (Fig. 7D) (86). Abdominal reflectance in the range 613-619 nm is roughly a third of its maximal reflectance, but there is substantial overlap of the abdomen's spectrum with BRh2 spectral sensitivity, which suggests that the Atala hairstreak's colour perception of the abdominal colour of conspecifics is not pure red, but something more like Orange-Red. All *Eumaeus* species have this orange-red colouration and the red scale reflectance spectra are similar between the sexes, suggesting that it evolved at the base of the *Eumaeus* genera, as an aposematic signal driven by the association with cycads. Whereas our results show a clear sex-specific reflectance difference on dorsal wings, it is unclear whether the red warning colouration patterns may have been coopted as a conspecific signal, as in other unpalatable butterflies (79, 87) , in which case it could be a by-product of the evolution of a signal used primarily to avoid predation. Hence, the L cones of most birds have red sensitivity from about 550 nm to 700 nm (59, 88), and the bright red abdominal/ hindwing reflectance in the near-infrared spectrum would stimulate mostly bird LWS /red (without MWS/ green) to perceive a colour close to "pure red".

Lycaenid butterflies with four rhodopsins can see more colour hues in the blue-
green range than mammals that have either di or trichromatic vision (34). Sex-specific
scale reflectance spectra on dorsal wings convey colour information that is readily
distinguishable by the visual system. Our study, by showing the expanded spectral
sensitivity exhibited by lycaenid butterflies through molecular variation and functional
changes in their 4-rhodopsin visual system, highlights the importance of peripheral
sensory genes in driving the adaptive evolution of multi-modal communication.

Materials and Methods

Butterflies

Pupae of *Eumaeus atala* were collected from host plants of *Zamia integrifolia* (locally known as "coontie") at the Montgomery Botanical Garden, Miami, FL, USA and reared at 22°C in an insectary in the MCZ laboratories under a 12:12 L:D cycle until emergence. *Callophrys sheridanii* and *Celastrina ladon* were collected on Badger Mountain, Wenatchee WA, USA and Wahclella Falls, OR, USA respectively. Eggs and young larvae of *Arhopala narathura japonica* were collected feeding on oak trees (*Quercus glauca*) from field sites near Ginoza, Okinawa, Japan and reared in the laboratory in Cambridge, MA, USA until they eclosed as adults.

Epi-Microspectrophotometry

Quantitative epi-microspectrophotometry (epi-MSP) was used to determine absorption spectra of butterfly rhodopsins by measuring eyeshine reflectance spectra after photoconversion of the rhodopsin to its metarhodopsin product (54, 89).

Compound eyes of most adult butterfly species exhibit eyeshine, a property that allows measuring rhodopsin absorbance spectra as well as spectral sensitivity of photoreceptor pupillary responses in eyes of intact butterflies. When subjected to repeated bright white flashes under incident-light microscope, 1) there is a photochemical effect, i.e. the coloration of eyeshine changes during each flash owing to changes in absorbance spectra that accompany photo-isomerization of rhodopsins to their metarhodopsin photoproducts (89); and 2) there is a pupillary response mediated by intracellular migration of pigment granules within photoreceptor cells, causing the intensity of eyeshine to decrease rapidly with time during each flash (90).

Three MSP methods (*in vivo* Photochemistry, Optophysiology, Retinal densitometry) are critical to study rhodopsin properties. The methods are presented here, whereas the rationale upon which the three methods are based and the general procedures for using eyeshine to make photochemical measurements from butterfly eyes are described in detail in SI methods.

In vivo Photochemistry

When rhodopsin is photo-isomerized to become metarhodopsin, it undergoes a spectral

shift in the absorbance spectrum. For LW rhodopsins, the shift is to shorter wavelengths;

the metarhodopsin peak is usually between 490 nm and 500 nm. For both UV-absorbing

and blue-absorbing rhodopsins, the shift is to longer wavelengths, typically to 475 nm -

490 nm. These photochemical changes are observable in eyeshine reflectance spectra,

as increased reflectance caused by loss of rhodopsin, and decreased reflectance

caused by the metarhodopsin. The computed absorbance-difference spectrum (DS),

therefore, has a positive peak caused by accumulation of metarhodopsin (M), and a

negative peak caused by loss of rhodopsin (R) (Fig. 1, 2A).

The absorbance difference spectrum relaxes with time in the dark, but changes

shape in doing so; the positive peak relaxes to zero much faster than the negative peak.

The entire temporal evolution of difference spectra can be reproduced quantitatively by

assuming different kinetics for the dark-processes of metarhodopsin decay and

rhodopsin recovery. Metarhodopsin decay is well approximated by a single exponential

process, but the time constant is a strong function of temperature. Rhodopsin recovery

is considerably slower than metarhodopsin decay, making it possible to create a partial

bleach using repeated episodes of bright flashes followed by dark periods during which

metarhodopsin decays totally from the rhabdom. The difference spectrum for that partial

bleach is a direct measurement of the absorbance spectrum of the LW rhodopsin (Fig. 1)

(89). Similar experiments with photoconversion of the other spectral types of rhodopsin

are more complicated because a bright blue flash designed to efficiently photo-isomerize

blue-absorbing rhodopsin will also convert some LW rhodopsins to their

metarhodopsins, although with less efficiency. However if the LW rhodopsins are first

bleached, then difference spectra for the blue rhodopsin are measurable (Fig. 2A).

Photochemical measurements of the *E. atala* blue rhodopsin were done with a

male oriented similarly to the female at Elevation 0° and Azimuth 10°. Before

photoconversion of the blue rhodopsin, the LW rhodopsins were partially bleached by

flashing with 20s exposure to RG645 (20s, 2s/55s). After resting in the dark for 24 min

for metarhodopsins to decay, the eye was flashed with 12s RG430 (2s/60s), which

converted blue rhodopsin to its metarhodopsin M505. A difference spectrum for R440

was computed from reflectance spectra measured before and 9 min after the series of

bright blue flashes.

Optophysiology

Photoreceptor cells in butterfly eyes contain intracellular pigment granules that move

centripetally in response to bright illumination and deplete light from the rhabdom by

scattering and absorption. This process creates an effective pupillary response

observable as a decrease in eyeshine reflectance (90). This intracellular pigment

migration is mediated exclusively by photo-isomerization of the rhodopsin contained

within the same cell's rhabdomere and is not influenced by physiological responses of

neighboring ommatidia. Thus, the pupillary pigment granules can be used as an optically

measured intracellular probe of physiological responses to light from that cell (91).

A double-beam Epi-MSP apparatus was used for optophysiological

measurements of eyeshine. One beam is deep-red filtered (e.g., 710 nm), that monitors

continuously the reflectance of eyeshine but does not itself cause a pupillary response.

The second beam delivers a monochromatic flash that evokes a pupillary response,

measured by the first beam as a decrease in deep-red eyeshine reflectance. At each

stimulating wavelength, the flash intensity is adjusted with computer-controlled neutral-

density wheels to produce a criterion decrease in reflectance (usually 3% - 5%).

Wavelength sequence is randomized. Both flash duration and inter-stimulus interval are

held constant. After completing an experimental series, the butterfly is replaced by a

factory calibrated Hamamatsu S1226 photodiode and quantum flux Q measured for

every criterion combination of wavelength and wheel setting. Spectral sensitivity is

computed as $S(\lambda) = 1 / Q(\lambda)$.

Retinal Densitometry

The visual pigment content of rhabdoms can be estimated quantitatively when the

tapetal reflectance spectrum is constant (white) for wavelengths shorter than about 600

nm (25, 92). A computational model based on an electron micrograph of the tracheolar

tapetum shows that this "chirped" set of layers functions as a broadband reflecting
interference filter exhibiting a computed reflectance greater than 90% for wavelengths
between 320 and 680 nm, thereby justifying the assumption of a white reflectance
spectrum in that band. This property of wideband white tapetal reflectance can be
exploited to computationally estimate rhodopsin contents. It is most valuable when
applied after λ_{max} of UV, blue, and LW rhodopsins have been determined from *in vivo*
photochemistry (Fig. 1; Fig. 2A) and optophysiology (Fig. 2B). R360, R440, and R565
have been determined in *E. atala* (Fig. 2C-D).

The procedure is sequential. First, a reflectance spectrum was measured after all
metarhodopsins had decayed from the rhabdoms, e.g., after overnight dark-adaptation
(black filled circles). Next, the dark spectrum was stripped of round-trip optical density
(OD) 1.50 (*male, m)* or 1.00 (fe*male, f)* of R565 rhodopsin so that the residual spectra
(red lines) were flat from 570 nm out to the 690 nm roll-off of tapetal reflectance. Next,
those residual spectra were stripped of OD 0.36 (*m*) and 0.22 (*f*) of R440 (cyan lines),
leaving large dips in the blue-green around 500 nm that were poorly fit by a single blue
opsin. However, stripping of OD 0.72 (*m)* and 0.55 (*f*) λ_{max} 395 (Retinal Binding Protein,
RBP) left UV residues (magenta lines) well fit by density of 0.55 (*m*) and 0.55 (*f*) of R360.
The remaining residuals (blue lines) were subjected to least-squares fitting to the
rhodopsin template (Fig. 2E), which produced excellent R495 fits of 0.39 (*f*) and 0.70
(*m*), supporting the presence of the fourth rhodopsin, R495.

The spectral sensitivities of *Arholopa, Celastrina and Callophrys* rhodopsins were
investigated using the same techniques as described in SI methods.

De novo E. atala *eye transcriptome*

The heads of 10 adult males (1 day old) were dissected under ambient light with their
antennae and palpi removed prior to flash freezing in liquid nitrogen. Total RNA was
extracted using the Direct-zol RNA extraction kit (Zymoresearch, CA, USA). Illumina
paired-end libraries were constructed using the Ultra II RNA Directional kit (New England
Biolabs, USA) and sequenced with an Illumina HiSeq v4. Adaptors were removed using
Trimgalore (93) and low quality reads were filtered out prior to generating a *de novo*

assembly reference transcriptome for all libraries in the Trinity sequence assembly and
analysis pipeline (94). This assembly resulted in 301656 transcripts with a contig N50 of
2519 bp, and average contig lengths of 1029 bp. Small fragments were filtered out
based on expression values using Kallisto(95). Opsin sequences from two other
lycaenids (41) were used as queries to identify all opsin mRNAs across tissues in the
final assembly using BLAST(96). To confirm their identity, the candidate opsin
sequences were blasted back to the NCBI non-redundant database.

Lycaenid Blue and LW opsin cDNA characterization

RNA was extracted from eye tissue preserved in RNA shield reagent (Zymo Research)
for *A. japonica, Ce. ladon and Ca. sheridanii*. Samples were first removed from the
storage solution with sterile forceps, briefly blot dried on a sterile Kimwipes paper, flash
frozen in a mortar containing liquid nitrogen and finely ground using a cold pestle.
Following RNA purification, we quantified RNA using a Quant-iT RNA kit and a Qubit
fluorometer (Invitrogen). From purified RNA, we synthesized cDNA using the The
GoScript™ Reverse Transcription System (Promega) and amplified a central region of
the long-wavelength opsin using degenerated oligonucleotide primers 5'-
TTGAAGCTTCARTTYCCNCCNATGAAYCC-3' (forward) and 5'-CGAATTCGTCAT
RTTNCCYTCIGGNACRTA-3' (reverse) (48). Single bands of expected sizes were
obtained, and the PCR products were purified with Exo-SAP, following Sanger
sequencing. We thereafter used the SMARTer RACE cDNA Amplification kit (Clontech)
to prepare 5'- and 3'- RACE cDNA for each species. We carried out RACE PCRs, and to
increase the specificity of RACE reactions, we performed nested PCRs for each cDNA
and obtained single-band PCR products, which were gel-purified using a Qiaquick Gel
Extraction Kit (Qiagen), ligated into PCR2.1 Vector kit (Invitrogen) and transformed into
competent TOPO10 cells (Invitrogen). Single bacterial clones were purified, and plasmid
DNAs were sequenced using M13F and M13R primers at the Harvard DF/HCC DNA
Resource Core. In total we obtained sequences from 5 to 10 opsin clones for each
RACE cDNA. Based on the 5'- and 3'-UTR information, gene-specific primers were
designed and used in combination with respective eye cDNAs to confirm the integrity of
each full-length LW opsin coding frame sequence. Opsin subfamily phylogenetic
placements were confirmed by aligning selected lepidopteran opsin genes extracted

from Genbank using the MAFFT package as implemented in Geneious(97), and a
Neighbor-Joining (NJ) tree of the aligned dataset was constructed using RAxML(98).

Cloning and protein expression

The coding region of each opsin transcript was amplified from eye cDNA and subcloned
in a modified pFRT-TO expression vector cassette derived from pcDNA5 and containing
the human cytomegalovirus (CMV) immediate early promoter (Invitrogen, USA). The
expression plasmid was modified to include a C-terminal tag by the monoclonal antibody
FLAG epitope sequence (DYKDDDDK), followed by a Ser-Gly-Ser linker peptide, a T2A
peptide sequence (EGRGSLLTCGDVEENPG) and the fluorescent marker protein
mRuby2. Plasmid DNAs were verified by Sanger sequencing and purified with the endo-
free ZymoPURE™ II Plasmid Midiprep Kit (Zymo Research, USA). Two micrograms of
plasmid DNAs were used to transfect small-scale HEKT293 cultures and optimize
expression conditions both via mRuby2 visualization and western blot analysis. Cells
were plated at a density of 0.6×10^6 cells in a 6-well culture dish containing DMEM
medium (Gibco), and transient transfection was achieved after 48h when reaching 80%
confluency in a 1:3 ratio DNA (µg): PEI (µL) (Polysciences, USA) at 1mg/mL in
molecular grade water, filter-sterilized at 0.22 µm. The transfected cells were harvested
in cold D-PBS (Sigma-Aldrich) after 2 days, centrifuged at 4°C for 5 min at 4000 rpm,
and resuspended in 50 µL Ripa lysis buffer (Invitrogen) supplemented with 1% n-
Dodecyl β-D-maltoside (Sigma-Aldrich). Cell membranes were lysed for 1h at 4°C with
gentle rotation on a sample homogenizer, and cell debris collected by centrifugation at
4°C for 15 min at 13,000 rpm. The crude protein lysate concentration was quantified by
BSA (Sigma-Aldrich) and 25 µg crude extract was loaded on NuPAGE™ 3-8% Tris-
Acetate gels (ThermoFisher) and transferred to a polyvinylidene difluoride membrane on
a TurboBlotTransfer system (Biorad Laboratories). The membranes were blocked with
1% milk (Biorad) in phosphate-buffered saline containing 0.1% Tween 20 (PBS-T,
Biorad) and incubated overnight with primary antibodies (aFLAG 1:2,500, aHSP90
1:50,000, GE Healthcare) containing 0.01% Sodium azide (Sigma Aldrich) on a gently
rocking platform at 4°C. After washing with TPBS the membranes incubated with aFLAG
and aHSP90 were respectively incubated with HRP Conjugated ECL anti-mouse and

ECL anti-rabbit (Amersham, USA), revealed using the SuperSignal West Femto (Thermo
Scientific) and imaged on a ChemiDoc system (Biorad Laboratories).

Transient expression, purification of expressed rhodopsins and spectroscopy

High-expressing clones from GPCR opsin cDNAs were transiently expressed in
HEKT293 cells prior to *in vitro* purification. For each construct, cells were seeded at a
density of 1.0×10^6 cells on day 0 in fifteen tissue culture dishes (10 cm diameter, ref
25382-166, VWR) in DMEM High Glucose, GlutaMAX (Life Technologies) supplemented
with 10% FBS (Seradigm Premium, VWR, USA). Lipid complexes containing 24 µg
DNA: 72 µL PEI (1mg/mL) diluted in Opti-MEM I Reduced Serum (Life Technologies)
were added 48h later to cells reaching 75-85% confluency. Six-hours post-transfection,
the culture medium was exchanged with new medium containing 5 $\mu.mol^{-1}$ 11-*cis*-retinal
(2mg/mL stock in 95% Ethanol) and under dim red illumination. The *cis*-retinal
absorption peak at 380 nm was confirmed using a NanoDrop™ 2000/2000c UV-VIS
Spectrophotometer (Thermo Fisher) prior to each experiment using a 1:100 dilution in
ethanol. Culture plates supplemented with cis-retinal were wrapped in aluminium foil and
cells were incubated in the dark. Forty-eight hours post-transfection, the medium was
decanted under dim red light illumination. Cells were scraped from the plates in cold
filter-sterilized HEPES wash buffer (3mM $MgCl_2$, 140mM NaCl, 50mM HEPES pH6.6-8.5
depending on protein isoelectric point) containing complete EDTA-free protein inhibitors
(Sigma-Aldrich), centrifuged for 10 min at 1,620 rcf at 4°C, and resuspended in 10 mL
wash buffer for two consecutive washes. After the second wash, cell pellets were gently
resuspended in 10mL cold wash buffer containing 40 µM 11-*cis*-retinal. Cells expressing
opsin-membrane proteins were incubated in the dark during 1h at 4°C on a nutating
mixer (VWR) to increase active rhodopsin complexes, and cells were then collected by
centrifugation at 21,500 rpm for 25 min at 4°C on a Sorvall WX Ultra 80 Series equipped
with an AH-629 Swinging Bucket Rotor (Thermo Scientific).

Transmembrane proteins were gently extracted by pipetting in 10 mL ice-cold extraction
buffer (3mM $MgCl_2$, 140mM NaCl, 50mM HEPES, 20% Glycerol v/v, 1% n-dodecyl β-D-
maltoside, complete EDTA-free protein inhibitors) and incubated for 1h at 4°C prior to

centrifugation at 21,500 rpm for 25 min at 4°C. The 10mL crude extract supernatant

containing solubilized rhodopsin complexes was added to 1mL Pierce™ Anti-

DYKDDDDK Affinity Resin (Thermo Scientific, USA) and incubated overnight at 4°C in a

15mL falcon on a nutating mixer. Samples were loaded on Pierce™ Centrifuge Columns

(ref 89897, Thermo Scientific, USA) and after 3 washes of the resin-bound FLAG-

epitope rhodopsin complexes with 3-column reservoir volumes of elution buffer (3mM

$MgCl_2$, 140mM NaCl, 50mM HEPES, 20% Glycerol v/v, 0.1% n-dodecyl β-D-maltoside),

the rhodopsin was eluted in 2mL elution buffer containing 1.25 mg (265 µM) Pierce™ 3x

DYKDDDDK Peptide (Thermo Scientific, USA). The eluate was concentrated using an

Amicon Ultra-2 Centrifugal Filter Unit with Ultracel-10 membrane (Millipore, USA), for 35

min at 4°C and 3,500rpm. The concentrated eluate (~350µL) was aliquoted in Amber

light-sensitive tubes (VWR, USA) and kept on ice in the dark. Ultraviolet-visible

absorption spectra (200-800nm) of dark-adapted purified proteins were measured in the

dark from 1.5 uL aliquots using a NanoDrop™ 2000/2000c UV-VIS spectrophotometer

(Thermo Fisher). Opsin purification yields were estimated following BSA analysis (Table

S23, SI methods). Spectroscopic analysis was performed from the mean value of 4-6

independent spectral measurements. Raw absorbance data were fitted to a visual

template (99) and polynomial functions analyses performed in R (V.0.99.486) (100)to

determine the opsin maximal absorption peaks.

Preparation of RNA probes and RNA in situ *hybridization*

We created *in vitro* transcription templates from UVRh, BRh1, BRh2, LWRh opsin

complementary DNA cloned in approximately 700-base-pair (bp) segments to pCRII-

TOPO (Invitrogen). Antisense cRNA probes were synthesized using T7 or Sp6

polymerases using either digoxigenin (DIG) or fluorescein (FITC) labelling mix (Sigma-

Aldrich) from gel-purified PCR templates. The synthesized cRNA probes were ethanol-

precipitated with NH_4OAc 7.5M and 1 µL glycogen, spun down at 4°C for 30 min, re-

dissolved in pure water and stored at -80°C. These RNA probes were first used to test

mRNA expression for each opsin receptor gene. We then tested the probes by dual

colour *in situ* hybridization using combinations of DIG and FITC probes to map opsin

receptor expression patterns.

For *in situ* hybridization, *E. atala* compound eyes were dissected and immersed in 1.5mL

eppendorf tubes containing freshly-made 4% formaldehyde (FisherScientific)/1x PBS for

2h at room temperature for fixation, then immersed successively in increasing sucrose

gradient solutions (10%, 20%, 30% in PBS) for 1h each, stored in 30% sucrose solution

overnight at 4°C, briefly transferred in OCT:sucrose 1:1, embedded in OCT (Tissue-Tek)

and frozen in dry ice. Tangential and longitudinal eye sections (12 µm) were obtained

using a cryostat (Leica), mounted on VWR Superfrost Plus Micro slides and used for

RNA *in situ* hybridization following a procedure described in details previously (101).

Double fluorescence *in situ* hybridization was performed using 100 µl hybridization

solution (pre-hybridization buffer supplemented with 4% Dextran sulfate (Sigma)

containing a combination of two opsin cRNA probes, one labeled with DIG and one

labeled with FITC (at 1ng.µl^{-1} for UVRh and BRh2, and 0.5ng.µl^{-1} for BRh1 and LWRh).

Eye anatomy

Each *E. atala* eye was immersed for prefixation in 2.5% Glutaraldehyde/2%

paraformaldehyde in 0.1M Sodium Cacodylate buffer (pH 7.4) (Electron Microscopy

Sciences, PA, USA) for 2h at room temperature, then stored at 4°C for 12-14h prior to

fixation, embedding, ultrathin sectioning and mounting on copper grids for TEM analysis

as described in SI methods.

Homology modeling and targeted mutagenesis

To investigate the molecular basis of spectral tuning differences between the duplicated

blue visual opsins, we carried out site-directed mutagenesis of amino acid substitutions

that could contribute to possible changes in the maximal absorption spectra of dark-

adapted rhodopsins. First, the Eat-B1 (λ_{max} = 435nm) and Eat-B2 (λ_{max} = 500nm) opsin

amino acid sequences were uploaded to the SWISS-MODEL protein recognition engine

(102) to generate templates aligned against the invertebrate squid rhodopsin crystal

structure (PDB2Z73) (72). The predicted homology model for each blue opsin was

analyzed in Pymol (103) to identify homologous binding sites in the *cis*-retinal binding

pocket within a range of 5Å from any carbon in the retinal polyene chain. Of the 101

amino acid substitutions that differ between the duplicate opsins, 21 residues were

predicted to interact with the *cis*-retinal chromophore, with 6 variant sites between both
opsin sequences.

Amino acid sequences from blue opsins were retrieved from Genbank and aligned using
the MAFFT package followed by NJ tree inference and support analysis derived from
1,500 bootstrap replicates in Geneious (97) prior to visualization in EvolView (98) with
squid rhodopsin as the outgroup. The phylogeny was used to identify functionally
convergent amino acid replacements repeatedly associated with similar shifts in
absorption spectra between blue opsin duplicates. We identified amino acid positions
that were likely to reside within the chromophore binding pocket of the opsin protein, and
that also had diverging biochemical properties (charge and/or polarity).. A BRh1 plasmid
DNA construct was modified to incorporate variant positions found in BRh2, namely
A116S (S1), I120F (S2), G175S (S3), Y177F (S4), I206C (S5), and F207C (S6).

Chimeric BRh1 rhodopsin constructs bearing single variant sites located on helix 3 (S1,
S2) and the β-strand located between helices 3 and 4 (S3, S4) were purified and
analyzed by spectroscopy in the native dark state. Since only S1 and S4 had tuning
effects in the range of interest, we combined coevolving adjacent site variants S1/S2,
and S3/S4 and followed two distinct routes by successively adding variant sites creating
triple mutants. Starting from a green-shifted BRh1 variant carrying S5 and S6, a third
trajectory was studied where variant sites S1, S2, S4 were also successively added.

Wing reflectance

Reflectance spectra were measured from leaves of *Zamia integrifolia* leaves (Zamiacae)
collected at the Montgomery Botanical Garden (Miami, FL, USA), and from *E. atala*
discrete wing, thorax, and abdominal patches of coloured-scales from both males and
females ($N_{individuals}$=3-5, 2 to 4 measurements per scale type, SI datafile 2), and in a Leitz
Ortholux-Pol microscope equipped with a Leitz MPV-1 photometer with epi-illumination
block, fitted with a Leitz 5.6X/0.15P objective. The illuminator filled the back focal plane
of the objective with axial incident light. The photometer measured reflected light from
the full aperture of the objective from a spot in the front focal plane that was 210 μm in
diameter. Reflectance data were corrected for stray light by subtracting data measured

from the MSP objective viewing a light-dump comprised of substantially out-of-focus
black velvet cloth. Corrected reflectance data were normalized against the same
normalization constant of 0.179 to preserve relative brightnesses among all measured
body patches. Normalized reflectance data were analyzed in R (V.0.99.486) (100).
898

Data accessibility

Our sequencing data have been deposited in the GenBank database under accession
numbers MN831881-MN831887 and under SRA Bioproject XXXXX. Data for
photochemistry and optophysiological measurements, opsin absorbance spectra,
reflectance spectra and transcriptomics expression are available as supplementary files.

Competing interests

We declare no competing interests

Acknowledgements

We thank Masaru Hojo for providing eggs, larvae and pupae of *Arhopala japonica*, Maria
Eriksson and Peg Coughin for assistance with electron microscopy, Sara Jones and
Jonathan Schmidt-Burk for advice with cell culture, Ian Slaymaker for advice with protein
purification procedures, Rachel Gaudet and José Velilla for guidance using Pymol and
homology modeling, Rosalie Crouch for providing the 11-cis-retinal, Jeanne Serb and
Davide Faggionato for advice with cis-retinal delivery, Almut Kelber, Mandyam
Veerambudi Srinivasan, Mary Caswell Stoddard, Julien Ayroles, and Riannon Macrae for
helpful discussions and Richard Belliveau, Maggie Starvish and Rachel Hawkins for
providing logistical support. This work was supported by a Mind Brain Behavior
Interfaculty grant to NEP and MAL, an Alice and Knut Wallenberg Postdoctoral
fellowship at the Broad Institute of MIT and Harvard to MAL, a NSF Graduate
Research Fellowship to SSa, personal funds of GDB, and NSF DEB- 1541560 to NEP
and PHY-1411445 to NEP and NY.

    Figures and Tables

    Figure 1. Long-wavelength photochemical difference spectra from lycaenid butterflies.

    Difference spectra (DS) were obtained following partial bleaches of long wavelength rhodopsins

    from dorsal retina of intact butterflies. Datapoints represent absorbance differences between

    amounts of Rhodopsin (R, green or red) and its Metarhodopsin photoproduct (M, blue) product

    after a dark-period that followed photoconversion. Each black curve represents a computed

    difference spectrum for least square fits estimates at *(A)* R570 of *Arhopala japonica*, *(B)* R562 of

    Eumaeus atala and *(C)* R518 of *Callophrys sheridanii*. In *E. atala* the difference spectrum was

    acquired upon complete degradation of the M photoproduct and with small contributions from

    R440 and RBP395. In *Celastrina ladon*, λ_{max} of M and λ_{max} of R are very close, which pushed the

    DS negative peak to the right. Photographs represent respective butterflies (left) and butterfly

    eyeshine (right) in the dorsal retina (see also Fig. S1).

    Figure 2. *In vivo* **evidence for four rhodopsins in the Atala hairstreak butterfly.** *(A)*

    Photochemical analysis of a butterfly male eyeshine using an epi-microspectrophotometer. Log-

    reflectance difference absorbance spectra (DS, Difference Spectrum, filled circles) and fitted

    curves (solid line) measured from dark-adapted eyes via partial-bleaching experiments of R440.

    (B) Optophysiological analyses designed to measure pupillary sensitivity in the UV. The least

    squares fit analysis provides confident estimates for a UV rhodopsin with λ_{max} at 360nm.

    Comparison to sensitivities in the blue, driven by R440 and in the red, driven by LW R565

    rhodopsins, show that those two opsins make no contributions in the UV. *(C-D)* Densitometric

    analysis of an epi-microspectrophotometric reflectance spectrum of a male *(C)* and female *(D).*

    Completely dark-adapted *E. atala* eyeshine is used to confirm the contribution of each estimated

    visual pigment in the eye. The black dots plot the log-reflectance spectrum. The red curve is the

    spectrum after having computationally stripped optical density (OD) 1.50 of rhodopsin with λ_{max}

    565 nm so that the residual spectrum is flat from 565 nm to 660 nm; cyan curve, the log-

    reflectance spectrum after having stripped OD 0.36 of R440 to produce a residual flat from 450

    nm. The black curve is the log-reflectance spectrum after stripping OD 0.72 for R395 and 0.55

    R360, leaving a residual spectrum fit by a fourth-rhodopsin R495. The densitometric analysis of

    female eyes *(D)* is qualitatively identical, with stripping densities of 1.55 R565, 0.40 R440, 0.80

    RBP395 and 0.55 R350, leaving a residual indicative of a fourth pigment contribution, fit by

    rhodopsin R495. Stripping the R495 residual leaves the curve plotted in open squares that is the

    putative average log-reflectance spectrum of the tapetum, that is flat from the UV out to 660 nm.

    (E) Computational analysis of male residual reflectance spectra supports the contribution of a

    fourth opsin pigment with λ_{max} peaking around 495 nm. The residual fit for females is shown in the

inset. *(F)* Sensitivity data from an optophysiological threshold experiment measured at
wavelengths close to λ_{max} values (360 nm, 440 nm, 495 nm, 562 nm) of the four rhodopsins. This
shows that sensitivities at 440 nm and 495 nm cannot possibly be driven by R565.

Figure 3. Functional characterization of red-shifted long wavelength lycaenid butterfly
opsins. *(A)* Schematic of modified pCDNA expression vector cassette and workflow for functional
characterization of rhodopsin complexes with cis-retinal chromophore. All steps following addition
of cis-retinal are performed under dim red-light illumination. *(B)* Neighbour-Joining (NJ) tree of
butterfly long wavelength opsins. Bootstrap node support is as follows: 50-74%, white circle; 75-
94%, grey circle; ≥ 95%, black circle. Previously published λ_{max} physiological data are indicated in
parentheses when known (20, 27, 40, 41, 45). Corresponding family names are labeled on the
right. L, T and P correspond to lycaenid subfamilies Lycaeninae, Theclinae and Polyommatinae,
respectively. *(C-F)* Dark spectra of long wavelength rhodopsin (LWRh) expressed using an
HEKT293 transient cell culture system and purified via FLAG-epitope. LWRh rhodopsin
absorbance spectra are indicated with black dots (Table S7, Dataset S1), and a rhodopsin
template data (99) was computed to obtain the best estimates of λ_{max} fitting the data. (C)
Arhopala japonica purified LW opsin with λ_{max} = 570 nm, (D) *Eumaeus atala* purified LW opsin
with λ_{max} = 567 nm, *(E) Callophrys sheridanii* purified LW opsin with λ_{max} = 525 nm, (F) *Celastrina*
ladon purified LW opsin with λ_{max} = 525 nm.

Figure 4. **Anatomical overview of a typical ommatidium in *Eumaeus atala*.** *(A)* Diagram
illustrating a typical ommatidium and the relative contribution of nine photoreceptor cells R1-R9 at
different depths along the rhabdom (480 µm long). Photoreceptors R1 and R2 are proximal cells
in region *a* (126 µm) that contribute microvillar structures containing UVRh, BRh1 and BRh2
rhodopsins. Photoreceptors R3-R4 are proximal receptors containing LWRh and Brh1
rhodopsins. Photoreceptors R5-R8 are distal cells exclusively expressing the LWRh rhodopsin
(see also Fig 4, Fig S2). The basal cell, R9 is restricted to the region immediately proximal to the
basement membrane (460 µm, h). The expressed mRNA opsin type was not investigated in this
cell. *(B)* Scanning electron micrographs from a male dorsal eye region at 30° elevation across the
rhabdom. Scale bars, 2 µm. *(C)* Distal microvilli in regions *a- b* are exclusively oriented parallel to
the R1-R2 axis (left panel). As R3-R4 cells expand and contribute the majority of microvilli from
regions *c* to *e*, together with novel contributions from proximal receptors R5 to R8, three
ommatidia subtypes are formed that exhibit microvillar contributions parallel to (left panel) but
also intertwisted (middle panel) and perpendicular to the R1-R2 axis (right panel). Scale bars, 500

nm. *bm,* basement membrane*; rh,* rhabdom*; nc,* nucleus*; tp,* tapetum*; vc,* vacuole; 1-9,
photoreceptor cells R1 to R9.

Figure 5. Adjacent photoreceptor localization between duplicate blue opsin mRNAs drives
retinal mosaic expansion. Double fluorescent *in situ* hybridization shows six ommatidial types in
E. atala compared to ancestral butterfly eyes. *(A-F)* Exclusive one-cell one-mRNA expression
pattern in distal photoreceptors R1 and R2, showing for *(A)* UVRh (inset in *(D)***,** *(B)* BRh1 mRNA
(inset in *(E)*) and *(C)* BRh2 mRNA (inset in *(F)*). Circles highlight the four distinct opsin expression
patterns in R1 and R2. *(G-I)* UVRh, BRh1 and BRh2 mRNAs do not coexpress in R1-R2 cells.
Photoreceptor cells R3-R8 express the LWRh opsin (red). Males and females show dorso-ventral
dimorphism (Fig. S4). *(J)* Schematic representation of opsin-based photoreceptor classes in
males and females of the six ommatidial types as follows (R1-R2-R3/R8): UV-UV-LW, UV-B1-
LW, UV-B2-LW, B1-B1-LW, B1-B2-LW, B2-B2-LW.

Figure 6. Residues responsible for spectral tuning shifts in blue rhodopsins. *(A)* NJ tree of
selected lepidopteran blue opsin amino acid sequences. The squid rhodopsin, *Todarodes*
pacificus (acc. nr. CAA49906) is used as outgroup. Bootstrap node support is as follows: 50-74%,
white circle; 75-94%, grey circle; ≥ 95%, black circle. Dots above the partial multiple sequence
alignment shows the 21 amino acid residues residing within 5 Å of any carbon atom in the retinal
polyene chain. *Blue* dots identify the six positions where amino acid residues differ between *E.*
atala BRh1 and BRh2, blue rectangles highlight variants at these positions in lycaenids. Residue
numbering is based on residue position in the squid rhodopsin. Residues are coloured according
to their physicochemical properties in Jalview v2 (104). Grey arrows indicate β-strands forming
the binding pocket. *(B)* Blue opsin absorbance spectra (dots) fitted to the visual template (cyan
and blue line functions), respectively. *(C)* Predicted structure for Eat-BRh1 based on homology
modeling with the squid rhodopsin with variant sites Ala116, Iso120, GLy175, Phe177, Iso206.
(D) Native Brh1 rhodopsin bearing S1 (A116), S2 (I120) and S4 (Y177). *(E)* Substituting residues
116, 120 and 177 leads to a 30-nm partial blue-shift in rhodopsin absorbance λ_{max} in the triple
mutant (see also Figure S2). This tuning shift may be mediated by several possible mechanisms
including additive effects caused by novel hydrogen bond formation at the coevolving adjacent
sites 116 and 120 *(F)* and with nearby conserved residues G115 and G121. *(G)* An alternative
evolutionary route involves substituting G175S (S3) which partially compensates the green tuning
shift of double mutant A116S/Y177F (Fig. S6, Table S8) and tunes the absorbance spectrum
near 500nm. These alternative trajectories highlight additive and epistatic interactions between
four residues at sites 1-4 *(F, H)* in the acquisition of the blue-shifted function.

1024

Figure 7. Wing and body spectral reflectance in *Eumaeus atala*. *(A)* Female (F) and male (M) dorsal forewing (DFW), ventral hindwing (VHW), thorax and abdominal scales. The photographs below each wing are magnified views of representative coloured wing scales in patches measured by epi-microspectrophotometry (MSP). The field of view for each photograph is 210 µm in diameter. Photographs were acquired using an Olympus TG-1 camera set at zoom 2.8. *(B-C)* Graphs of mean reflectance spectra ± standard errors of the mean of wing scale patches *(B)* and body scales *(C)* (N=3-5 individuals per sex). *(D)* Leaf reflectance spectra (4-5 measurements per leaf surface) (SI Datafile 2). *(E) E. atala* rhodopsin absorbance spectra. Male and female wing reflectance spectra are not exclusive, and spectral shapes are highly consistent indicating that male and female spectra are similar in the visible. In females, hindwing cyan spots show a reflectance peak at 540 nm *(B)*. Blue scales on the thorax and dorsal forewings have a reflectance peak at 490-500 nm. Hindwing *(B)* and abdominal red patches *(C)* have identical high reflectance spectra at long wavelengths in both sexes. MSP analyses from black areas of ventral hindwings *(B)* showed that black scales are only 1 % as reflective as adjacent cyan scales in the blue/green band. Brightness in black regions decreases 100-fold compared to adjacent cyan scales in both sexes. All reflectance curves overlap at contributing wavelengths for at least two rhodopsins, thereby efficiently distinguishing foliage and sex-specific wing colour patterns.

References

References

1.   M. Lynch, J. S. Conery, The evolutionary fate and consequences of duplicate genes.
Science **290**, 1151-1155 (2000).

2.   H. Kaessmann, Origins, evolution, and phenotypic impact of new genes. *Genome*
research **20**, 1313-1326 (2010).

3.   M. Long, K. Thornton, Gene duplication and Evolution. *Nature* **293**, 1551a (2001).

4.   J. Zhang, Evolution by gene duplication: an update. *Trends in Ecology & Evolution* **18**,
292-298 (2003).

5.   M. A. Liénard, H.-L. Wang, J.-M. Lassance, C. Löfstedt, Sex pheromone biosynthetic
pathways are conserved between moths and the butterfly *Bicyclus anynana*. *Nat Comms* **5:3957**
(2014).

6.   R. Feuda, F. Marlétaz, M. A. Bentley, P. W. H. Holland, Conservation, Duplication, and
Divergence of Five Opsin Genes in Insect Evolution. *Genome biology and evolution* **8**, 579-587
(2016).

7.   P. Brand *et al.*, The evolution of sexual signaling is linked to odorant receptor tuning in
perfume-collecting orchid bees. *Nature Communications* **11**, 244 (2020).

8.   M. Chapal, S. Mintzer, S. Brodsky, M. Carmi, N. Barkai, Resolving noise–control conflict
by gene duplication. *PLOS Biology* **17**, e3000289 (2019).

9.   M. Soskine, D. S. Tawfik, Mutational effects and the evolution of new protein functions.
Nature Reviews Genetics **11**, 572-582 (2010).

10.   J. Zhang, Y. Zhang, H. Rosenberg, Adaptive evolution of a duplicated pancreatic
ribonuclease gene in a leaf-eating monkey. *Nature Genetics* **30**, 411-415 (2002).

11.   G. M. Castiglione, B. S. W. Chang, Functional trade-offs and environmental variation
shaped ancient trajectories in the evolution of dim-light vision. *eLife* **7**, e35957 (2018).

12.   M. Manceau, V. Domingues, C. Linnen, E. Rosenblum, H. Hoekstra, Convergence in
pigmentation at multiple levels: mutations, genes and function. *Philos Trans R Soc Lond B Biol*
Sci **365**, 2439-2450 (2010).

13.   S. B. Carroll, Evolution at Two Levels: On Genes and Form. *PLOS Biology* **3**, e245
(2005).

14.   F. Hauser, B. Chang, Insights into visual pigment adaptation and diversity from model
ecological and evolutionary systems. *Current opinion in genetics & development* **47**, 110-120
(2017).

15.   A. Terakita, The opsins. *Protein family review* **6**, 213 (2005).

16.   S. Yokoyama, Evolution of Dim-Light and Color Vision Pigments. *Annu Rev Genom Hum*
G **9**, 259-282 (2008).

17.   D. E. Nilsson, M. F. Land, J. Howard, Optics of the butterfly eye. *Journal of Comparative*
Physiology A **162**, 341-366 (1988).

18.   A. D. Briscoe, L. Chittka, The evolution of color vision in insects. *Ann Rev Entomol* **46**,
471-510 (2001).

19.   G. Bernard, C. Remington, Color vision in *Lycaena* butterflies: spectral tuning of receptor
arrays in relation to behavioral ecology. *Proc Nat Acad Sci USA* **88**, 2783-2787 (1991).

20.   F. Frentiu, G. Bernard, M. Sison-Mangus, A. Van Zandt Brower, A. Briscoe, Gene
Duplication Is an Evolutionary Mechanism for Expanding Spectral Diversity in the Long-
Wavelength Photopigments of Butterflies. *Mol Biol Evol* **24**, 2016-2028 (2007).

21.   A. D. Briscoe *et al.*, Positive selection of a duplicated UV-sensitive visual pigment
coincides with wing pigment evolution in Heliconius butterflies. *PNAS* **107**, 3628-3633 (2010).

22.   D. Armisén *et al.*, The genome of the water strider Gerris buenoi reveals expansions of
gene repertoires associated with adaptations to life on the water. *BMC Genomics* **19**, 832 (2018).

23.   C. R. Sharkey *et al.*, Overcoming the loss of blue sensitivity through opsin duplication in
the largest animal group, beetles. *Scientific Reports* **7**, 8 (2017).

24.   H. Koshitaka, M. Kinoshita, M. Vorobyev, K. Arikawa, Tetrachromacy in a butterfly that
has eight varieties of spectral receptors. *Proc Biol Sci* **275**, 947-954 (2008).

25.   F. D. Frentiu *et al.*, Adaptive evolution of color vision as seen through the eyes of
butterflies. *PNAS* **104**, 8634-8640 (2007).

26.   R. Futahashi *et al.*, Extraordinary diversity of visual opsin genes in dragonflies. *PNAS*
112, E1247 (2015).

27.   M. Wakakuwa *et al.*, Evolution and Mechanism of Spectral Tuning of Blue-Absorbing
Visual Pigments in Butterflies. *PLOS ONE* **5**, e15015 (2010).

28.   F. E. Hauser, I. v. Hazel, B. S. W. Chang, Spectral tuning in vertebrate short wavelength-
sensitive 1 (SWS1) visual pigments: Can wavelength sensitivity be inferred from sequence data?
J Exp Zool **322**, 529-539 (2014).

29.   M. Perry *et al.*, Molecular logic behind the three-way stochastic choices that expand
butterfly colour vision. *Nature* **535**, 280-284 (2016).

30.   A. D. Briscoe, Reconstructing the ancestral butterfly eye: focus on the opsins. *J Exp Biol*
211, 1805-1813 (2008).

31.   K. Arikawa, The eyes and vision of butterflies. *J Physiol* **595**, 5457-5464 (2017).

32.   G. H. Jacobs, Primate photopigments and primate color vision. *Proceedings of the*
National Academy of Sciences **93**, 577 (1996).

33.   M. Luehrmann *et al.*, Short-term colour vision plasticity on the reef: changes in opsin
expression under varying light conditions differ between ecologically distinct fish species. *The*
Journal of Experimental Biology **221**, jeb175281 (2018).

34.   G. H. Jacobs, Evolution of colour vision in mammals. *Philos Trans R Soc B* **364**, 2957-
2967 (2009).

35.   Y. Shichida, T. Matsuyama, Evolution of opsins and phototransduction. *Philos Trans R
Soc B* **364**, 2881 (2009).

36.   A. Gutierrez Eduardo de *et al.*, The role of ecological factors in shaping bat cone opsin
evolution. *Proc Biol Sci* **285** (2018).

37.   A. Monteiro, Origin, Development, and Evolution of Butterfly Eyespots. *Annual Review of
Entomology* **60**, 253-271 (2015).

38.   K. Dasmahapatra *et al.*, Butterfly genome reveals promiscuous exchange of mimicry
adaptations among species. *Nature* **487**, 94-98 (2012).

39.   N. P. Lord *et al.*, A cure for the blues: opsin duplication and subfunctionalization for short-
wavelength sensitivity in jewel beetles (Coleoptera: Buprestidae). *BMC Evolutionary Biology* **16**,
107 (2016).

40.   M. P. Sison-Mangus, A. D. Briscoe, G. Zaccardi, H. Knüttel, A. Kelber, The lycaenid
butterfly Polyommatus icarus; uses a duplicated blue opsin to see green. *Journal of Experimental
Biology* **211**, 361 (2008).

41.   M. P. Sison-Mangus, G. D. Bernard, J. Lampel, A. D. Briscoe, Beauty in the eye of the
beholder: the two blue opsins of lycaenid butterflies and the opsin gene-driven evolution of
sexually dimorphic eyes. *J Exp Biol* **209**, 3079-3090 (2006).

42.   H. Awata, M. Wakakuwa, K. Arikawa, Evolution of color vision in pierid butterflies: blue
opsin duplication, ommatidial heterogeneity and eye regionalization in Colias erate. *Journal of
Comparative Physiology A* **195**, 401-408 (2009).

43.   P. Chauhan *et al.*, De novo transcriptome of Ischnura elegans provides insights into
sensory biology, colour and vision genes. *BMC Genomics* **15**, 808 (2014).

44.   A. Briscoe, Six opsins from the butterfly *Papilio glaucus*: molecular phylogenetic evidence
for paralogous origins of red-sensitive visual pigments in insects. *Journal of Molecular Evolution*
51, 110-121 (2000).

45.   K. Arikawa, Spectral organization of the eye of a butterfly, *Papilio*. *J Comp Physiol* **189**,
791-800 (2003).

46.   P.-J. Chen, H. Awata, A. Matsushita, E.-C. Yang, K. Arikawa, Extreme Spectral Richness
in the Eye of the Common Bluebottle Butterfly, Graphium sarpedon. *Front Ecol Evol* **4**, 18 (2016).

47.   G. Pratt, J. Emmel, G. D. Bernard, The Buckwheat Metalmarks. *Am Butterflies* **19**, 4-31
(2011).

48.   M. Wakakuwa, D. G. Stavenga, M. Kurasawa, K. Arikawa, A unique visual pigment
expressed in green, red and deep-red receptors in the eye of the small white butterfly, *Pieris
rapae crucivora*. *J Exp Biol* **207**, 2803-2810 (2004).

49.   Y. Ogawa, M. Kinoshita, D. G. Stavenga, K. Arikawa, Sex-specific retinal pigmentation
results in sexually dimorphic long-wavelength-sensitive photoreceptors in the eastern pale

clouded yellow butterfly, Colias erate. *The Journal of Experimental Biology* **216**, 1916 (2013).

50.   A. Satoh *et al.*, Red-shift of spectral sensitivity due to screening pigment migration in the eyes of

a moth, Adoxophyes orana. *Zool Lett* **3**, 14 (2017).

51.   P. Pirih *et al.*, The giant butterfly-moth Paysandisia archon has spectrally rich apposition eyes with unique light-dependent photoreceptor dynamics. *Journal of Comparative Physiology A* **204**, 639-651 (2018).

52.   G. Bernard, Red-absorbing Visual Pigment of Butterflies. *Science* **203**, 1125-1127 (1979).

53.   G. Zaccardi, A. Kelber, M. P. Sison-Mangus, A. D. Briscoe, Color discrimination in the red range with only one long-wavelength sensitive opsin. *J Exp Biol* **209**, 1944-1955 (2006).

54.   G. D. Bernard, Bleaching of rhabdoms in eyes of intact butterflies. *Science* **219**, 69-71 (1983).

55.   S. Yokoyama *et al.*, Epistatic Adaptive Evolution of Human Color Vision. *PLOS Genetics* **10**, e1004884 (2014).

56.   M. Neitz, J. Neitz, G. H. Jacobs, Spectral tuning of pigments underlying red-green color vision. *Science* **252**, 971 (1991).

57.   Dulai KS, v. D. M, M. JD, H. DM, *Genome Res* **9**, 629-638 (1999).

58.   S. Kawamura, Kawamura S. Color vision diversity and significance in primates inferred from genetic and field studies. Genes Genomics. 2016;38:779–791. doi:10.1007/s13258-016-0448-9. *Genes Genomics* **38**, 779-791 (2016).

59.   A. Kelber, Bird colour vision - from cones to perception. *Current opinion in Behavioral Sciences* **30**, 34-40 (2019).

60.   A. Terakita *et al.*, Expression and comparative characterization of Gq-coupled invertebrate visual pigments and melanopsin. *J Neurochem* **105**, 883-890 (2008).

61.   F. Frentiu *et al.*, Clines in butterflies suggest novel roles for insect photopigments. *Mol Biol Evol* **32**, 368-379 (2014).

62.   T. Saito *et al.*, Spectral tuning mediated by helix III in butterfly long wavelength-sensitive visual opsins revealed by heterologous action spectroscopy. *Zoological Letters* **5**, 35 (2019).

63.   M. Wakakuwa, F. Stewart, Y. Matsumoto, S. Matsunaga, K. Arikawa, Physiological basis of phototaxis to near-infrared light in *Nephotettix cincticeps*. *Journal of Comparative Physiology A* **200**, 527-536 (2014).

64.   A. Kelber, Ovipositing butterflies use a red receptor to see green. *J Exp Biol* **202**, 2619-2630 (1999).

65. R. Cannon, *Courtship and mating in butterflies: reproduction, mating, behaviour and*
sexual conflicts. R. J. C. Cannon, Ed. (Severn, Gloucester, UK, 2020), pp. 376.

66. M. Heikkilä, L. Kaila, M. Mutanen, C. Pena, N. Wahlberg, Cretaceous origin and repeated
tertiary diversification of the redefined butterflies. *Proceedings of the Royal Society of London,*
Series B: Biological Sciences **279**, 1093-1099 (2012).

67. M. Espeland *et al.*, A Comprehensive and Dated Phylogenomic Analysis of Butterflies.
Cur Biol **28**, 770-778.e775 (2018).

68. S. Koi, D. Hall, Atala Butterfly, Atala Hairstreak, Coontie Hairstreak, *Eumaeus atala* Poey
1832 (Insecta: Lepidoptera: Lycaenidae). (2016).

69. A. Myrberg, M. Mohler, J. Catala, Sound production by males of a coral reef fish
(*Pomacentrus partitus*): its significance to females. *Animal Behaviour* **34** (1986).

70. D. Faggionato, J. M. Serb, Strategy to Identify and Test Putative Light Sensitive Non-
Opsin G-Protein-Coupled Receptors: A Case Study. *Biol. Bull.* **233**, 70-82 (2017).

71. J. Morrow, B. Chang, The p1D4-hrGFP II expression vector: A tool for expressing

and purifying visual pigments and other G protein-coupled receptors. *Plasmid* **64**, 162-169
(2010).

72. M. Murakami, T. Kouyama, Crystal structure of squid rhodopsin. *Nature* **453**, 363 (2008).

73. A. Kelber, L. S. V. Roth, Nocturnal colour vision – not as rare as we might think. *J Exp*
Biol **209**, 781 (2006).

74. J. Fasick, N. Lee, D. Oprian, Spectral tuning in the human blue cone pigment.
Biochemistry **38**, 11593-11596 (1999).

75. B. Wilts, H. Leertouwer, D. G. Stavenga, Imaging scatterometry and
microspectrophotometry of lycaenid butterfly wing scales with perforated multilayers. *J R Soc*
Interface **6**, S185-S192 (2009).

76. D. G. Stavenga, H. Leertouwer, P. Pirih, M. Wehling, Imaging scatterometry of butterfly
wing scales. *Optics Express* **17**, 193-202 (2009).

77. R. Vane-Wright, M. Boppre, Visual and chemical signalling in butterflies: functional and
phylogenetic perspectives. *Proceedings of the Royal Society B Biological Sciences* **340**, 197-205
(1993).

78. J. A. Fordyce, C. C. Nice, M. L. Forister, A. M. Shapiro, The significance of wing pattern
diversity in the Lycaenidae: mate discrimination by two recently diverged species. *Journal of*
Evolutionary Biology **15**, 871-879 (2002).

79. C. Jiggins, R. Naisbit, R. Coe, J. Mallet, Reproductive isolation caused by colour pattern
mimicry. *Nature* **411**, 302-305 (2001).

80. L. Wilkins, D. Osorio, Object colours, material properties and animal signals. *The Journal*
of Experimental Biology 10.1242/jeb.204487, jeb.204487 (2019).

81.   J. Schnapf, T. Kraft, D. Baylor, Spectral sensitivity of human cone photoreceptors. *Nature*
325, 439-441 (1987).

82.   M. Imafuku, I. Shimizu, H. Imai, Y. Shichida, Sexual difference in color sense in a
lycaenid butterfly, *Narathura japonica*. *Zoolog Sci* **24**, 611-613 (2007).

83.   D. McCoy, R. Prum, Convergent evolution of super black plumage near bright color in 15
bird families. *J Exp Biol* **222**, Pt 18 (2019).

84.   D. McCoy *et al.*, Structurally assisted super black in colourful peacock spiders. *Proc Biol*
Sci **286**, 20190589 (2019).

85.   A. L. Davis, H. F. Nijhout, S. Johnsen, Diverse nanostructures underlie thin ultra-black
scales in butterflies. *Nature Communications* **11**, 1294 (2020).

86.   O. Liew, P. Chong, B. Li, A. Asundi, Signature Optical Cues: Emerging Technologies for
Monitoring Plant Health. *Sensors* **8**, 3205-3239 (2008).

87.   M. Rossi *et al.*, Neural processing genes are linked to divergence in visual mate
preference in sympatric Heliconius butterflies. *bioRxiv*
10.1101/2020.03.22.002121, 2020.2003.2022.002121 (2020).

88.   D. Osorio, M. Vorobyev, A review of the evolution of animal colour vision and visual
communication signals. *Vision Research* **48**, 2042-2051 (2008).

89.   G. D. Bernard, Dark-Processes Following Photoconversion of Butterfly Rhodopsins.
Biophysics of Structure and Mechanism **9**, 277-286 (1983).

90.   D. G. Stavenga, J. Numan, J. Tinbergen, J. Kuiper, Insect pupil mechanisms. II. Pigment
migration in retinula cells of butterflies. *J Comp Phys A* **113**, 73-93 (1977).

91.   G. D. Bernard, "Noninvasive optical techniques for probing insect photoreceptors" in
Biomembranes Part H: Visual pigments and purple membranes I, Methods in Enzymology*, P. L,
Ed. (Academic Press, New York, 1982), vol. 81, pp. 752-759.

92.   A. Briscoe, G. D. Bernard, A. Szeto, L. Nagy, R. White, Not all butterfly eyes are created
equal: Rhodopsin absorption spectra, molecular identification, and localization of ultraviolet-, blue-
, and green-sensitive rhodopsin-encoding mRNAs in the retina of Vanessa cardui. *J Comp Neurol*
458, 334-349 (2003).

93.   F. Krueger, "Trim galore." A wrapper tool around Cutadapt and FastQC to consistently
apply quality and adapter trimming to FastQ files. **516**, 517 (2015).

94.   B. J. Haas *et al.*, De novo transcript sequence reconstruction from RNA-seq using the
Trinity platform for reference generation and analysis. *Nat Prot* **8**, 1494-1512 (2013).

95.   N. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq
quantification. *Nature Biotechnology* **34**, 525-527 (2016).

96.   S. F. Altschul *et al.*, Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Research* **25**, 3389-3402 (1997).

97.   M. Kearse *et al.*, Geneious Basic: An integrated and extendable desktop software
platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647-1649
(2012).

98.   A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).

99.   A. Palacios, T. Goldsmith, G. D. Bernard, Sensitivity of cones from a cyprinid fish (Danio
aequipinnatus) to ultraviolet and visible light. *Vis Neurosci* **13**, 411-421 (1996).

100.   R. C. Team, R: A language and environment for statistical computing. R Foundation for
Statistical

Computing, Vienna, Austria. URL https://www.R-project.org/. (2015).

101.   Y. Isogai *et al.*, Molecular organization of vomeronasal chemoreception. *Nature* **478**, 241-
245 (2011).

102.   A. Waterhouse *et al.*, SWISS-MODEL: homology modelling of protein structures and
complexes. *Nucleic acids research* **46**, W296-W303 (2018).

103.   Anonymous (The PyMOL Molecular Graphics System, Version 1.8, Schrödinger, LLC.

104.   A. M. Waterhouse, J. B. Procter, D. M. A. Martin, M. Clamp, G. J. Barton, Jalview version
2: A Multiple Sequence Alignment and Analysis Workbench. *Bioinformatics* **25**, 1189-1191
(2009).

1277

A

M485
R570
DS

B

RBP395
R440
R562
DS

C

M508
R519
DS

A

CMV | Opsin | Linker | FLAG | Linker | T2A | mRuby2 | Amp R* | Hygro R*

1. Transfect high-expression *opsin-FLAG-T2A-mRuby2* plasmids

2. Change medium, Add *cis*-retinal — 6h

3. Harvest cells, nutate rhodopsin, solubilize membrane fraction — 36h

4. Purify rhodopsin via resin affinity, measure UV-VIS absorption — FLAG-resin, o/n

B

Eurema hecabe
Colias erate CeL
Pieris rapae (563 nm)
Anthocharis scolymus   Pieridae

Lycaena helloides   L
Lycaena rubidus (565 nm)
Lycaena heteronea
Lycaena nivalis
Arhopala japonica   T
Callophrys sheridanii
Eumaeus atala
Satyrium behrii
Celestrina ladon   P
Agriades glandon
Polyommatus icarus   Lycaenidae

Apodemia mormo LW2
Apodemia mormo LW1   Riodinidae

Heliconius melpomene
Heliconius erato
Agraulis vanillae
Eueides lineata
Dryas iulia
Limenitis archippus (514 nm)
Limenitis lorquini (530 nm)
Limenitis weidemeyerii (530 nm)
Limenitis astyanax (545 nm)
Vanessa cardui
Bicyclus anynana
Danaus plexippus   Nymphalidae

Troides aeacus TaL2
Papilio xuthus Rh2 (515 nm)
Papilio machaon PmL2
Graphium sarpedon GsL2
Graphium sarpedon GsL3b
Troides aeacus TaL3
Papilio xuthus Rh3 (560 nm)
Papilio machaon PmL3
Graphium sarpedon GsL3a
Papilio machaon PmL1
Papilio xuthus Rh1 (540 nm)   Papilionidae

C A.ja-LW (λ_{max}= 574nm)

Absorbance vs Wavelength (nm)

D E.at-LW (λ_{max}= 569nm)

Absorbance vs Wavelength (nm)

E Ca.sh-LW (λ_{max}= 521nm)

Absorbance vs Wavelength (nm)

F Ce.la-LW (λ_{max}= 530nm)

Absorbance vs Wavelength (nm)

UVRh   BRh1   BRh2   LWRh