TITLE

Effects of visual inputs on neural dynamics for coding of location and running speed in medial

entorhinal cortex.

AUTHORS

Holger Dannenberg[1], Hallie Lazaro[1], Pranav Nambiar[1], Alec Hoyland[1], Michael E. Hasselmo[1]

INSTITUTIONAL AFFILIATIONS

[1]Center for Systems Neuroscience, Boston University, Boston, MA 02215, USA
10

ABSTRACT

Neuronal representations of spatial location and movement speed are important for a broad

range of cognitive functions, including spatial self-localization and memory-guided navigation.

Two possible speed signals, by theta frequency or by firing rate, have been hypothesized to

provide the velocity signal needed for generating the spatially periodic grid cell firing pattern.

However, which of these speed signals is utilized by the brain remains unknown. By

manipulating visual inputs and analyzing the time-courses of evoked changes, we demonstrate

that changes in spatial stability of grid cell firing correlate in time with changes in speed coding

by local field potential theta frequency. In contrast, visual inputs do not affect speed coding by

firing rate even if baseline firing rates are changed. Moreover, grid cells maintain a spatially

periodic firing pattern, though less stable, in complete darkness. These data suggest that mice

use an oscillatory speed signal to perform path integration.

INTRODUCTION

The spatially periodic firing of grid cells in the medial entorhinal cortex (MEC) (Fyhn et al., 2004,

2008; Hafting et al., 2005) is widely recognized as a neuronal correlate of memory-guided

and/or self-motion-based spatial navigation (McNaughton et al., 2006). Based on that idea,

computational models of grid cells including attractor network models (Burak & Fiete, 2009;

Fuhs & Touretzky, 2006), oscillatory interference models (Blair et al., 2008; Burgess, 2008;

Burgess et al., 2007; Hasselmo, 2008; Zilli & Hasselmo, 2010), hybrid models of attractor

networks and oscillatory interference (Bush & Burgess, 2014), or probabilistic learning models

(Cheung, 2016) suggest that grid cells integrate running direction and traveled distance to

maintain a spatially periodic firing pattern. However, the source of both the direction signal and

the speed signal are unclear. The direction signal has been attributed to the head direction

signal terminating in the MEC (Sargolini et al., 2006; Taube et al., 1990; Taube et al., 1990;

Taube, 2007), but this does not match movement direction (Raudies et al., 2015). The analysis

presented here focuses on the question of the dominant source of the speed signal which is

also still under debate. One potential candidate for the speed signal is the linear modulation of

firing rates in a subpopulation of context-invariant "speed cells" in the MEC (Kropff et al., 2015).

However, such a speed signal by firing rate has been shown to be more heterogeneous than

previously assumed (Hardcastle et al., 2017) and not accurate on sub-second-long time scales

(Dannenberg et al., 2019). An alternative or complementary oscillatory speed signal is provided

by the positive correlation between the local field potential (LFP) theta frequency and the

running speed of an animal and/or the positive correlation between theta rhythmic firing

frequency of theta modulated MEC neurons and the running speed of an animal (Hinman et al.,

2016). Both the firing rate and rhythmicity running speed signals depend on self-motion cues,

which include vestibular and proprioceptive signals (Chen et al., 2019; Tennant et al., 2018) and

also moving sensory features such as optic flow (Campbell et al., 2018; Raudies et al., 2012;

Raudies & Hasselmo, 2015). However, any path integration mechanism ultimately suffers from

the accumulation of error over distance traveled or elapsed time (Kraus et al., 2015;

Sreenivasan & Fiete, 2011) necessitating error correction by static environmental sensory inputs

(Evans et al., 2016; Giocomo, 2016) such as visual landmark cues (Hafting et al., 2005) or

environmental boundaries (Barry et al., 2007; Hardcastle et al., 2015) to maintain an accurate

representation of relative self-location within a spatial environment. Experimental data from

studies on rats support the hypothesized role of grid cells in performing self-motion based path

integration and error correction by sensory cues: Self-motion-based signals such as head

direction are necessary for grid cell firing (Winter et al., 2015), and grid cell firing persists during

darkness with reduced spatial selectivity (Hafting et al., 2005) reflecting the loss of error

correction by visual cues. In contrast, two more recent studies on mice reported a nearly

complete loss of spatially periodic grid cell firing during complete darkness (Chen et al., 2016;

Pérez-Escobar et al., 2016) suggesting a critical role for visual inputs on grid cell firing and

presenting a challenge for current path integration models. However, visual cues appear to have

been the only available source of sensory information in those experiments leaving the question

open if and how visual inputs affect grid cell firing in mice under standard conditions which

include other landmark features such as walls, auditory, and olfactory cues still available to the

animal. Moreover, little is known about the time courses of changes in coding of location by grid

cells and running speed by theta oscillations or firing rate (for instance during changes from light

to darkness), although changes in grid cell firing, theta rhythmic dynamics, and speed

modulation of firing rates may underlie many cognitive functions on a behavioral time scale

ranging from seconds to minutes. Moreover, we propose that testing for temporal correlations

between grid cell spatial stability and the two proposed speed signals—by firing rate or theta

oscillations—will help to identify the speed signal with the greatest influence on grid cell firing.

Data presented in this study are relevant both for models of path integration and for our

understanding of how visual inputs modulate the coding of location and running speed in the

entorhinal cortex.

RESULTS

To study changes in neural dynamics after transitioning from foraging in an illuminated

environment (light) to foraging in complete darkness (dark) and vice versa, we recorded single

unit and local field potential (LFP) activity in mice freely exploring an open field environment

during alternating light and dark epochs (**Fig. 1A**). We first studied if the absence of visual

inputs in the dark condition changed the speed signal by LFP theta frequency. We found that

the slope of the LFP theta frequency vs. running speed relationship was changed between light

and dark conditions (t-statistic of interaction effect: 18.045, $p < 0.0000$, n = 15 mice) (**Fig. 1B**).

In contrast, the theta amplitude vs. running speed relationship did not change significantly (t-

statistic of interaction effect: 1.7343, $p = 0.0832$, n = 15 mice) (**Fig. 1C**). Average running speed

during light was 11.6 ± 0.5 cm/s which slightly decreased after transition to dark by 0.89 ± 0.31

cm/s (7.7%) to 10.72 ± 1.4 cm/s (mean ± sem, $p = 0.0117$, n = 15 mice, Wilcoxon's signed rank

test) (**Fig. 1D and Supplementary Fig. 1A**). Similarly, running acceleration also slightly

decreased from 20.8 ± 1.9 cm/s^2 in the light condition by 1.7 ± 0.7 cm/s^2 (8.0%) to 19.1 ± 2.3

cm/s^2 in the dark condition ($p = 0.0070$, n = 15 mice, Wilcoxon's signed rank test) (**Fig. 1E**). We

therefore examined if the observed decrease in the slope of the LFP theta frequency vs. running

speed relationship might result from the decrease in running *acceleration* during darkness. If the

decrease in slope is due to a decrease in running acceleration, there should be no change in

the y-intercept of the LFP theta frequency vs. running *acceleration* between light and dark

conditions. However, the y-intercept of the LFP theta frequency vs. running acceleration

relationship did change significantly in darkness (t-statistic: 114.67, $p < 0.0000$, n = 15 mice,

Student's paired t-test) (**Fig. 1F**)**.** Furthermore, if a decrease in acceleration causes the change,

then there should be no change in the slope of the LFP theta frequency vs. running speed

relationship between light and dark conditions if we restrict our analysis of speed tuning to time

points where the running acceleration is constant. However, testing the relationship between

LFP theta frequency vs. running speed only at time points when running acceleration was near

zero (0 ± 5 cm/s^2) resulted in speed tuning curves of LFP theta frequency (**Fig. 1G**) that looked

nearly identical to the ones obtained without that restriction (Fig. 1B). These data demonstrate

that the LFP theta frequency is directly modulated by running speed and that the observed

change in the slope of the LFP theta frequency vs. running speed relationship in darkness is not

an artifact of an underlying change in running behavior but instead directly reflects a change of

visual inputs.

The LFP theta frequency vs. running speed relationship has been hypothesized to serve as a

speed signal important for coding of location during self-motion-based navigation including path

integration. Having established that the removal of visual inputs changes the speed signal by

LFP theta frequency, we were further interested in the time-course of that change. We therefore

analyzed sequential 10-sec blocks after transitioning from light to dark and vice versa.

Averaging data from 73 sessions from 13 mice, we found that the time course of the change in

LFP theta frequency vs. running speed relationship showed two components: an instantaneous

change and a further exponential change at a time scale of tens of seconds (**Fig. 1H, I, K**). After

removal of all visual inputs, LFP theta frequency shows an instantaneous decrease followed by

a further exponential decay. Once the environment is illuminated again, LFP theta frequency

shows an instantaneous increase followed by a further saturating exponential increase over

tens of seconds. Analyzing the slopes of the LFP theta frequency speed tuning curves for

sequential 10-sec blocks revealed that the observed exponential change in theta frequency over

time was largely driven by a change in the slope of the theta frequency vs. running speed

relationship (**Fig. 1J, L**), while the y-intercepts were only slightly—though significantly—lower

during darkness ($t = 4.90$, $p = 3.6 \times 10^{-5}$, $n = 15$ time bins, Student's paired t-test)

(**Supplementary Fig. 1B & C**).

Local field potential activity in the MEC primarily reflects the summed activity of synaptic inputs

to the MEC, which is likely to be correlated with firing patterns of MEC neurons. Many neurons

in the MEC show a theta rhythmic firing pattern and—similarly to the LFP theta frequency—the

frequency of theta rhythmic firing increases with running speed (Hinman et al., 2016). In our

data set, we identified a total of $n = 342$ neurons from 14 mice as theta modulated based on

their autocorrelograms. We asked if and how theta rhythmic firing in these theta modulated

MEC neurons is affected by removing and reinstating all visual inputs. Towards that aim we

used an MLE approach (Climer et al., 2015) to fit a model of theta spiking rhythmicity to the

observed spike train autocorrelations and identify the frequency and magnitude of theta

rhythmic firing in theta modulated MEC neurons in the light and dark conditions (see **Fig. 2A** for

one example neuron). We first analyzed the relationship between the frequency of theta

rhythmic firing and running speed. The results of this analysis show that the frequency of theta

rhythmic firing increases with running speed in an illuminated environment (n = 340 neurons,

Fig. 2B) as previously shown in rats (Hinman et al., 2016). Similar to the observed decrease in

slope of the LFP theta frequency vs. running speed relationship, the slope of the frequency of

theta rhythmic firing vs. running speed relationship decreased in complete darkness (**Fig. 2B**).

To study the time-course of changes in theta rhythmic firing, we next analyzed theta rhythmic

firing in sequential 10-sec blocks after transitioning from light to dark and vice versa. In contrast

to the exponential changes observed in the LFP theta frequency, the frequency of theta

rhythmic firing decreased instantaneously with removal of all visual inputs and increased again

instantaneously with reinstating all visual inputs (**Figs. 2C-E**). Interestingly, the theta rhythmic

magnitude of the average autocorrelogram model across all 342 theta modulated neurons

appeared higher during complete darkness compared to the light condition (**Figs. 2C & E**). The

increased amplitude in the average across autocorrelogram models could either reflect an

increase of theta rhythmic magnitude on the single cell level or an increased synchrony across

neurons on the population level. To distinguish between these two possibilities, we analyzed

changes in theta rhythmic firing on the single cell and population level for different running

speeds (immobility defined as running speed < 3 cm/s, 5-10 cm/s, 10-15 cm/s, 15-20 cm/s, 20-

25 cm/s) (n = 339 neurons, **Fig. 2F**). To our surprise, we found that neurons continued to fire

rhythmically during immobility periods in both light and dark conditions. However, theta rhythmic

frequencies varied largely across neurons within a wide range of theta frequencies (4-12 Hz)

during immobility. As soon as the mouse started running, however, theta rhythmic firing

synchronized across neurons around a narrower theta frequency range of 6-10 Hz.

Interestingly, theta dynamics showed a different development between light and dark conditions

(**Fig. 2F-I**). The magnitude of theta rhythmic firing increased with running speed during the light

condition, but decreased with running speed during the dark condition (notice brighter yellow

peaks in **Fig. 2F**). Similarly, the frequency of theta theta rhythmic firing increased with running

speed during the light condition, but less so during the dark condition (**Fig. 2F**).

In order to quantify these changes in theta rhythmic firing on the population level, we identified

individual neurons with statistically significant rhythmic modulation of firing per running speed

condition (alpha = 0.1). In the light condition, we found 206 out of 339 neurons with statistically

significant rhythmic firing during immobility. This number sharply increased to 298 with the onset

of running, but then steadily decreased to 270, 256, and 241 for higher running speeds. In the

dark condition, the number of rhythmically firing neurons showed a similar sharp increase with

the onset of running (from 187 to 313 neurons). In contrast to the light condition, however, the

number of rhythmic neurons during darkness further increased to 322, 326, and 328 out of 339

neurons with higher running speeds (**Fig. 2G**). These numbers demonstrate that the ensemble

of theta rhythmic firing neurons decreased with running speed in the light condition, but

increased to ceiling during the dark condition (**Fig. 2G**) contributing to the stronger synchrony of

theta rhythmic firing observed on the population level (**Fig. 2F**).

We next quantified the changes in theta rhythmicity magnitude and theta rhythmic frequency in

the subset of rhythmically firing neurons per running speed condition and comparing light vs.

dark conditions. Theta rhythmicity magnitude increased with running speed during light, but

decreased with running speed during dark resulting in larger differences between light and dark

conditions for higher running speeds. Consequently, these differences in theta rhythmicity

magnitude reached statistical significance for running speeds higher than 15 cm/s (**Fig. 2H,**

Kolmogorov-Smirnov test, $p < 0.001$ for running speeds > 15 cm/s, see Fig. 2H for exact p-

values). Notably, the amplitudes of the average autocorrelogram models across all cells were

very similar between light and dark conditions (**Fig. 2F,** third row) despite the observed

differences in the magnitudes of theta rhythmicity on the single cell level (**Fig. 2H**). These data

suggest that—during darkness— higher synchrony across MEC neurons on the population level

counterbalances the lower magnitude of theta rhythmicity on the single cell level resulting in little

change on the LFP level. In contrast, changes in theta rhythmic frequencies with higher running

speed correlated with changes in LFP theta frequency in both light and dark conditions (**Figs.**

1B & 2I); theta rhythmic frequencies increased with running speed and more so in the light

compared to the dark condition (**Fig. 2I**). Consequently, the differences between light and dark

became more prominent and statistically more significant with higher running speeds (**Fig. 2I**,

Kolmogorov-Smirnov test, $p < 0.05$ for immobility periods, $p < 0.0001$ for running speeds > 5

cm/s, and $p < 0.00001$ for running speeds > 10 cm/s, see Fig. 2I for exact p-values).

Motivated by the differences in LFP activity during light and dark conditions, we asked if these

changes were accompanied by changes in firing rates of MEC neurons. We therefore first asked

if and how the removal and reinstating of visual inputs affected baseline firing rates in MEC

neurons. We found striking differences in baseline firing rates between light and dark epochs

among a wide variety of MEC neurons including speed cells, head direction cells, grid cells,

boundary-vector cells and interneurons **(Fig. 3A & B; Supplementary Fig. 2)**. In general,

changes in firing rates of MEC neurons could be positive or negative with positive changes

being observed ~4 times more frequently than negative changes (139 out of 1205 or 11.5%

positively modulated neurons; 32 out of 1205 or 2.7% negatively modulated neurons at

significance level alpha = 0.05; data from 15 mice). In both cases, changes in firing rate after

transition from light to dark or from dark to light occurred instantaneously (**Fig. 3C & D**).

One possible explanation for the observed changes in firing rates can be a binary change in the

visibility of landmark cues, i.e. the presence or absence of visual information about the animal's

spatial surroundings. Alternatively, changes in firing rates could be driven by changes in net

excitatory drive from areas in the visual cortex due to changes in retinal stimulation independent

of any gain or loss of information about the animal's spatial surroundings. To distinguish

between these two alternative hypotheses, we analyzed data from optogenetic control

experiments in a subset of two mice, in which mice carried a light fiber attached to their

headstage with the fiber tip approximately 1cm above the animal's head. Green laser light

(561nm, 1mW) was emitted from the tip of that light fiber and illuminated the head and

immediate surroundings of the animal's head. While being visible to the animal as an

egocentrically stationary spot of light, it was too weak to illuminate the open field environment

and thus did not contribute to the visibility of distal visual cues useful for navigational purposes

(e.g. walls or the cue card of the open field arena). Nevertheless, we found strong effects of that

kind of visual point stimulation on firing rates of five MEC neurons from two mice during foraging

under the light condition (**Fig. 3E & F**). Four out of those five neurons showed positive

modulation by the egocentrically stationary spot of laser light, and one showed negative

modulation. The effects of presenting an egocentrically stationary visual stimulus during the light

condition on baseline firing rates of MEC neurons resembled the effects of transitioning from the

dark to the light condition. Overall, increased baseline firing rates due to stronger retinal

stimulation were ~4 times more likely to be observed than decreases in firing rates. In summary,

these data demonstrate that the changes in retinal stimulation due to changes in the amount of

spatially non-informative light hitting the retina after transitioning from dark to light and vice

versa are sufficient to explain the observed changes in firing rates of MEC neurons.

We next turned to the analysis of the hypothesized speed signal by firing rate, in particular

"speed cells" in the MEC (Kropff et al., 2015) and asked if the speed modulation of firing rates in

MEC neurons is changed in complete darkness. Speed cell firing has previously been reported
to be affected by removal of visual inputs and this change in speed cell firing has been
interpreted as a change in the hypothesized speed signal by speed cells, which has further
been indicated to underlie changes in grid cell firing and path integration during darkness (Chen
et al., 2016; Pérez-Escobar et al., 2016). As shown above, many neurons change baseline firing
rates during darkness and many speed cells show conjunctive coding properties which may be
changed in complete darkness (Hardcastle et al., 2017; Hinman et al., 2016). To our knowledge,
previous analyses of firing rate modulation by running speed have not been corrected for
changes in baseline firing rates or changes in conjunctive coding properties. In this study, we
therefore 1) corrected for effects on baseline firing rates by random spike deletion to match the
number of spikes across light and dark conditions, 2) applied a linear-nonlinear (LN) model
(Hardcastle et al., 2017) to identify speed modulated neurons and correct for conjunctive coding
properties, such as conjunctive coding for position, head direction, and LFP theta phase. The
LN-model provides an unbiased method for identification of significantly speed modulated
neurons and for quantification of their speed response curves (Hardcastle et al., 2017). The
firing rates of n = 142 neurons out of 425 cells from 14 mice showed significant modulation by
running speed in either the light or the dark condition. Consistent with previous reports, speed
modulation of firing rates was heterogenous (**Fig. 4A**) (Dannenberg et al., 2019; Hardcastle et
al., 2017; Hinman et al., 2016). Hierarchical clustering of speed cells resulted in two main
clusters of 119 positively and 23 negatively modulated neurons (**Fig. 4B**). We did not find
biologically significant differences in the speed tuning of firing rates comparing light vs. dark
conditions in positively or negatively modulated MEC neurons (**Fig. 4C**). Likewise, speed scores
(Pearson's correlation coefficient between firing rate and running speed) of all speed modulated
neurons were not significantly changed between light and dark conditions on the population
level (p = 0.6722, n = 142, Kolmogorov-Smirnov test) (**Fig. 4D**) and were highly correlated
between light and dark conditions on the single cell level (R = 0.87) **(Fig. 4E)**. Moreover,
conjunctive coding of position, head direction and theta phase was unchanged (**Fig. 4F**). In
summary, our data on speed modulation of firing rates in MEC neurons demonstrate that a
potential speed signal by speed cells in the MEC depends neither on visual inputs nor visibility
of landmark cues. Furthermore, speed modulation of firing rates is indifferent to changes in
baseline firing rates or changes in conjunctive coding properties.
270
While grid cells have been reported to show stable grid cell firing during complete darkness in
rats (Hafting et al., 2005), two recent studies in mice showed a strong reduction in the

hexagonal symmetry of grid cell firing during complete darkness (Chen et al., 2016; Pérez-
Escobar et al., 2016) challenging computational models of grid cells and path integration. To
test if this discrepancy is due to a difference between species or due to differences in the
experimental settings, the focus in this study was on selectively eliminating visual inputs only,
while leaving other environmental and/or sensory cues such as walls or odors and auditory cues
potentially present in the testing arena intact. We first tested if grid cell firing can emerge in
complete darkness. We therefore introduced mice to the familiar open field arena during
complete darkness (the experimenter was wearing night-vision goggles) and recorded grid cells
during the initial dark session and a following light and dark session (10-20 min duration each).
We found that spatial periodicity of grid cell firing was low—but not completely eliminated—in
the initial dark session and then strongly increased during the first light session **(Fig. 5A & B)**.
Interestingly, after this initial orientation, spatial periodicity of grid cell firing decreased again
after removal of visual inputs, but remained significantly higher than in the initial dark session
(Chi-square = 10.89, p = 0.0043, n = 9 cells from two mice, Friedman test; post hoc tests
revealed p < 0.05 comparing initial dark and light, and initial dark and second dark sessions).
We next aimed to study the change in grid cell spatial firing patterns after transitioning from light
to dark in more detail. Towards that aim, we recorded grid cells during alternating light/dark
epochs, always beginning with a light epoch followed by a dark epoch. 27 cells from six mice
were identified as grid cells based on their grid scores (see Methods). To compare spatial firing
properties of grid cells between light and dark conditions, we concatenated all light and dark
sessions to compute firing rate maps for light and dark conditions. We found that grid cell field
centers remained stable, but grid fields appeared wider during complete darkness (**Fig. 5C**).
Quantification of spatial periodicity and spatial precision of grid cell firing by computing grid
scores and spatial information (see Methods) revealed a significant decrease in both spatial
periodicity and spatial precision of spike locations (**Fig. 5D & E**). However, firing rate maps are
the results of integrating firing rates of grid cells over long time periods, thereby measuring only
spatial variability, but not spatio*temporal* variability of grid cell firing. This distinction becomes
important when considering two alternative hypotheses with regard to the underlying cause for
the observed broadening of grid fields in complete darkness. Grid fields could appear wider
because grid cells fire over a longer distance throughout a single grid field traversal.
Alternatively, grid fields could appear wider because grid cell firing becomes less stable at the
level of single grid field traversals. To distinguish between these two alternative hypotheses, we
computed the spatiotemporal correlation between the observed firing rate over time and the
firing rate predicted by the animal's location at any given point in time and the associated value

in the firing rate map at that location. The results show a significant reduction in spatiotemporal

correlation during complete darkness (**Fig. 5F**) supporting the hypothesis that grid fields in the

firing rate appear broader because the spatiotemporal accuracy of grid cell firing is reduced.

Our data set included data on n = 7 grid cells from three mice recorded over an exceptionally

long time period across multiple days. These data provided sufficient statistical power and

sufficient sampling of the environment to analyze the time courses of the observed reductions in

spatial periodicity, spatial selectivity and spatiotemporal stability of grid cell firing after transitions

from light to dark and vice versa. Towards this goal, we first computed firing rate maps for each

of the 10-sec bins of activity to analyze the time course of spatial periodicity quantified by the

grid score (**Fig. 6A & B**). Second, we computed the spatial correlations between the firing rate

maps of the 10-sec binned data and the overall firing rate map for the light condition (**Fig. 6C**).

Third, we computed the spatiotemporal correlation for each of the 10-sec bins (**Fig. 6C**). We

found an exponential change in spatial periodicity of grid cell firing (**Fig. 6B**), spatial correlation

(**Fig. 6C**), and spatiotemporal correlation (**Fig. 6D & E**) over tens of seconds after transitions

from light to dark and vice versa. Intriguingly, the time course of changes in spatiotemporal

correlation showed a remarkable similarity to the time course of changes in the slope of the LFP

theta frequency vs. running speed relationship (**Fig. 6F**). Taken together, these data

demonstrate that the representation of space by grid cell firing is correlated to the speed signal

by LFP theta frequency and that both of these signals undergo plastic changes at a behavioral

time scale of tens of seconds.

DISCUSSION

Data presented in this study show that visual inputs affect neural dynamics underlying the

coding of location and running speed in the MEC. These effects include changes in LFP theta

rhythmicity as well as changes in baseline firing rates, changes in theta rhythmic firing, and

changes in spatial periodicity of grid cell firing in response to visual inputs. By analyzing the time

courses of changes in those variables, we found that spatial periodicity of grid cell firing and

LFP theta frequency showed an instantaneous component and a slower exponential component

of change. In contrast, changes in baseline firing rates and theta rhythmic firing only showed the

instantaneous component. Furthermore, speed modulation of firing rates showed no change

after transitions between light and dark conditions. Our data demonstrate that changes in the

speed signal from LFP theta frequency is in concert with changes in the spatial stability of grid

cells. Moreover, coding of location by grid cells and running speed by LFP theta frequency

showed changes at a behavioral time scale of tens of seconds. Such changes in response to
changes in sensory input are longer than previously addressed by theoretical and computational
models of grid cell firing. Lastly, we report changes in the running speed-dependent population
dynamics of theta rhythmic MEC neurons upon changes in visual inputs. In both the light and
dark condition, theta rhythmic neurons become more synchronized with the onset of running.
However, synchronization across neurons is stronger during darkness, while frequency and
magnitude of theta rhythmic firing of individual neurons increase significantly more with running
speed during the light compared to the dark condition.
349
It has previously been proposed that disruption of speed signals may cause disruption of
spatially periodic grid cell firing in the MEC. Computational models of grid cell firing and path
integration require a linear speed signal (Burak & Fiete, 2009; Burgess, 2008; Burgess et al.,
2007; Fuhs & Touretzky, 2006; McNaughton et al., 2006). Two candidates for such a speed
signal terminating in the MEC have been proposed: a speed signal by firing rate, most clearly
present in a subpopulation of "speed cells" showing context-invariant positive and linear speed
tuning curves (Kropff et al., 2015); and an oscillatory speed signal by changes in LFP theta
frequency or by changes in the theta rhythmic firing frequency of individual MEC neurons
(Burgess, 2008; Hinman et al., 2016). Both speed signals can change in response to changes of
the size and shape of the environment (Munn et al., 2020), as do grid cells (Barry et al., 2007;
Krupic et al., 2015). Both speed signals may arise from the medial septum/diagonal band of
Broca (MSDB) (Fuhrmann et al., 2015; Hinman et al., 2016; Justus et al., 2016) and contribute
to self-motion-based linear distance estimation in the MEC (Jacob et al., 2017). However, speed
modulation of firing rates is inaccurate at sub-second long time scales (Dannenberg et al., 2019)
and the use of longer integration time windows has been shown to be insufficient for spatial
periodicity of grid cell firing in a computational model of grid cell firing (Dannenberg et al., 2019).
366
The theta oscillatory speed signal also depends on the MSDB (Chrobak et al., 1989;
Dannenberg et al., 2015; Givens & Olton, 1994; Hinman et al., 2016; Lawson & Bland, 1993)
and pharmacological inactivation of MSDB neurons strongly reduces both theta oscillations and
spatial periodicity of grid cell firing in the MEC (Brandon et al., 2011; Koenig et al., 2011).
Furthermore, passive transportation of rats reduces theta oscillatory power and abolishes both
spatially periodic grid cell firing and the speed signal by LFP theta frequency (Winter et al.,
2015). Likewise, pharmacological inactivation of the vestibular system or preventing
linear/translational head movements in a virtual reality setup abolish the modulation of LFP

theta frequency by running speed and are associated with a decrease in the spatial information

score and an increase in grid scale (Chen et al., 2018; Jacob et al., 2014). Interestingly, the

increase in grid scale observed during path integration on a circular track is associated with a

change in LFP theta oscillations, but not with a change in the firing patterns of speed and head

direction cells (Jacob et al., 2019). Taken together, these data suggest a dominant role of theta

frequency modulation by running speed in generating the spatially periodic grid cell firing

pattern.

However, the relative importance of the speed signal by firing rate on the one hand and the

oscillatory speed signal by theta oscillations on the other hand for grid cell firing has not been

explicitly tested so far. In order to test the relative importance of both speed signals, we took

advantage of the disruption of spatially periodic grid cell firing caused by the absence of visual

inputs and included the time domain into our analysis. Similar to the reasoning underlying the

concept of Granger causality, we argued that a change in the speed signal driving path

integration of grid cell firing should correlate in time with the changes in grid cell firing. However,

if changes in the speed signal in question are uncorrelated to changes in grid cell firing, we can

exclude a causal contribution of that potential speed signal to grid cell firing. In order to analyze

the time course of grid cell stability over time after transitions between light and dark, we took

advantage of long-term recordings of the same grid cells over multiple days providing us with

sufficient statistical power. Intriguingly, changes in spatial stability of grid cell firing showed an

instantaneous component and a slow exponential component at a behavioral time scale of tens

of seconds after transitions between light and dark conditions. Notably, the observed

exponential component in changes of grid cell spatially periodic firing is validated by an earlier

report by Pérez-Escobar et al. (2016) although they reached different conclusions. We next

tested if a change in the potential speed signal by firing rate may underlie the observed change

in spatial stability of grid cell firing. However, the speed modulation of firing rates during

complete darkness may show heterogeneous and multiplexed speed modulation of firing rate

(Hardcastle et al., 2017) as well as changes in baseline firing rates due to loss of retinal

stimulation and excitatory inputs from visual cortex (Campbell & Giocomo, 2018). We therefore

corrected for differences in the number of spikes between light and dark epochs. We further

corrected for multiplexed or conjunctive coding properties by computing speed response curves

using a linear-nonlinear model (Hardcastle et al., 2017). After correcting for changes in baseline

firing rates and conjunctive coding properties, we found that visual inputs had no impact on the

speed modulation of firing rates in MEC neurons.

We next turned to the analysis of the time courses of changes in the two oscillatory speed

signals, namely the speed signal by LFP theta frequency and the speed signal by theta rhythmic

firing frequency. Intriguingly, the time course of changes in LFP theta frequency matched the

time course of changes in grid cell spatial stability (Fig. 6F). In contrast, changes in theta

rhythmic firing frequency showed only the instantaneous component of change (Fig. 2). These

data on the parallel time course of changes in LFP theta and grid cell stability suggest that grid

cells may depend more on the mean attractor state of rhythmic firing in the MEC reflected by the

LFP theta signal than the rhythmic firing of individual neurons. Alternately, the exponential

change in LFP theta may reflect a slow change in an oscillatory speed signal arising from other

cortical structures such as the medial septum which has been demonstrated to be critical for

both LFP theta and grid cell firing (Brandon et al., 2011; Koenig et al., 2011). Future studies are

needed to address the question how theta rhythmic firing of MEC neurons can change more

rapidly than the overall mean response in the LFP signal. Notably, the average frequency of

theta rhythmic firing is higher than the average LFP theta frequency and we found variability in

the range of theta rhythmic firing frequencies across neurons indicating that rhythmic firing of

MEC neurons is only loosely coupled to the LFP theta frequency. Moreover—in contrast to LFP

theta rhythmicity—theta rhythmic firing persists during immobility, though significantly less

synchronous across neurons. The observed changes in theta rhythmic firing frequency at the

single cell and population levels also provide a possible explanation for why the LFP theta

amplitude vs. running speed relationship is not changed during complete darkness. During

darkness, movement-related theta rhythmicity magnitude is decreased on the single cell level,

while theta synchronicity across neurons is increased on the population level. These two effects

would influence the LFP theta amplitude in opposite directions and thus may cancel each other

out.

The observed changes in coding of running speed and location during complete darkness may

be attributed to either a loss of visual cues for navigational purposes or alternatively to a net

decrease in excitatory drive from areas in the visual cortex due to the lack of retinal stimulation

(Campbell & Giocomo, 2018). In order to distinguish between these two possibilities, we took

advantage of control data from an optogenetic study (Dannenberg et al., 2019), where a visible

bright spot of light was moving geostationary to the animal without further illuminating the

environment. During this manipulation, the amount of light hitting the retina increased without

changing the visibility of static visual landmark cues. This perturbation resulted in very similar

changes of firing rates in MEC neurons as observed during the light vs. dark paradigm (Fig. 3).
Importantly, changes of firing rates and theta rhythmic firing frequency occurred instantaneously
upon changes in visual inputs (Figs. 2D and 3B & D). These data suggest that changes in firing
rates and theta rhythmic firing frequency are primarily driven by changes in net excitatory input
from visual cortex and do not reflect changes in the perception of the spatial aspects of the
environment. We conclude that a lack of visual cues contributes to error accumulation in the
path integration system. However, our data oppose previous proposals that changes in baseline
firing rates of speed cells reflect the lack of information from visual cues. Instead, our data
suggest that changes in baseline firing rates of speed cells are driven by changes in retinal
stimulation or levels of brightness in the environment, while the modulation of firing rates by
running speed—the actual speed signal by firing rate—is unaltered by changes in illumination or
presence or absence of visual cues. Taken together, data presented in this study argue against
a role of speed modulation of firing rates as speed signals for path integration and/or grid cell
firing. In contrast, we found a striking correlation of changes in the LFP theta frequency vs.
running speed relationship with changes in spatiotemporal accuracy of grid cell firing and
hexagonal symmetry of grid cell firing. These data argue in favor of computational models of
grid cell firing and path integration using an oscillatory speed signal by changes in LFP theta
frequency (Burgess, 2008; Burgess et al., 2007; Hasselmo, 2008). This also suggests that
individual neuron spiking may respond more rapidly than a slower mean population response of
attractor dynamics underlying grid cell firing changes during the transitions.
463
Two recent studies on freely moving mice reported a major disruption of spatially periodic grid
cell firing in the absence of visual cues (Chen et al., 2016; Pérez-Escobar et al., 2016). These
findings questioned the role of grid cells for path integration-based navigation (Gil et al., 2018;
Jacob et al., 2019) and are at odds with previous reports on grid cells in rats (Hafting et al.,
2005) as well as results from studies using virtual reality demonstrating that physical motion has
a greater influence on grid cell firing in mice than visual inputs (Campbell et al., 2018; Chen et
al., 2019; Tennant et al., 2018). It remains controversial, however, if those discrepancies reflect
a species difference between mice and rats or a difference in the experimental conditions.
Critical to the interpretation of the studies carried out by Chen et al. (2016) and Pérez-Escobar
et al. (2016), their studies carefully removed olfactory, auditory, and tactile cues in addition to
the manipulation of visual cues to leave the animals completely disoriented, and impeding the
selective study of visual inputs on spatially periodic grid cell firing. In this study, we therefore
selectively manipulated visual inputs leaving other static environmental and sensory cues such

as walls, auditory cues, and potential olfactory cues unaltered. During this more selective

manipulation, grid cells in the MEC persisted during complete darkness but showed reduced

spatial selectivity. Our results further show that grid cell spatial periodic firing was reduced

significantly more when mice were introduced to the test environment in complete darkness.

These data suggest that grid cell firing needs to be initially anchored to a cognitive map and that

this initial anchoring or orientation under standard testing conditions largely depends on the

presence of static visual cues. However, once the animals have oriented themselves in the

environment and grid cells have been anchored to a cognitive map, grid cell firing can be

maintained by path integration even in the absence of visual information—though with

decreasing spatial stability due to the absence of error correction by visual cues. Taken

together, these data suggest that vestibular signals and proprioceptive feedback as well as

external geometrical cues are dominant factors maintaining the orientation, scale, and phase of

grid cell firing fields, while visual cues are used for refining a self-motion-based cognitive

representation of self-location.

In summary, the time courses of changes in grid cell hexagonal symmetry, spatially selective

firing, and spatiotemporal correlation caused by changes in visual inputs strongly resembled the

time course of changes in speed tuning of LFP theta frequency (Fig. 6F). In contrast, neither

changes in baseline firing rate, nor changes in theta rhythmic firing frequency showed the slow

exponential component of change after transitioning from light to dark or vice versa.

Furthermore, speed modulation of firing rates remained unaltered between light and dark

conditions. These data suggest that the hypothesized oscillatory speed signal by LFP theta

frequency may be more important for grid cell firing than a hypothesized speed signal by firing

rate. Theta signals within the hippocampal formation have been suggested to not only reflect

alterations in sensorimotor integration and/or the flow of sensory input, but also distinct cognitive

operations (Hinman et al., 2011). Interestingly, we observed not only a slow exponential *decay*

of spatial stability of grid cell firing upon removal of visual inputs, but also a slow saturating

exponential *increase* in grid cell stability after reinstating illumination of the environment. Such a

slow component at a time scale of tens of seconds after each transition between light and dark

conditions suggest that the representation of space in a cognitive map is updated continuously

at a behavioral time scale and more malleable than previously acknowledged.

METHODS

Subjects. Data were collected from a total of 17 adult 3-6 month old male mice (*mus musculus*)

of wildtype and transgenic backgrounds (Wildtype, C57Bl/6J; ChAT-IRES-Cre, B6;129S6-

Chat[tm2(cre)Lowl]/J; PV-IRES-Cre, B6;129P2-Pvalb[tm1(cre)Arbr]/J). Experimental PV-IRES-Cre mice

were homozygous and experimental ChAT-IRES-Cre mice were heterozygous. All experimental

procedures were approved by the Institutional Animal Care and Use Committee for the Charles

River Campus at Boston University. Mice underwent surgery for implantation of 2 movable

tetrodes targeting the superficial layers of the MEC in the left hemisphere. For the purpose of

additional experiments not reported in this study, mice also underwent rAAV injections of

Archaerhodopsin T or GCaMP6f into the medial septum and light fiber implantation targeting the

medial septum (Dannenberg et al., 2019). No differences were observed between wildtype or

transgenic mice. Prior to surgery, mice were housed in Plexiglas cages together with their

siblings. After surgery, mice were separated for individual housing and maintained on a

reversed 12-h light/12-h dark cycle. Mice were habituated to the experimenter and testing room

before the start of all experimental procedures.

Surgery for microdrive implantation in mice. Mice were injected with buprenorphine (0.1-

mg/kg, s.c.) and atropine (0.1-mg/kg, i.p.), and chronically implanted under isoflurane

anesthesia with 8-channel Axona microdrives (Axona, St. Albans, UK) carrying two movable

tetrodes targeting the superficial layers of the dorsal MEC in the left hemisphere. One ground

screw was implanted above the cerebellum, five anchoring screws were positioned across the

skull, and a craniotomy was performed for implantation of the microdrive, 0.5-mm anterior to the

edge of the transverse sinus and 3.4-mm lateral to the midline. Tetrodes were then implanted

with a 6° polar angle and lowered 1-mm into the brain. The microdrive was secured to the skull

with dental cement. Animals were given buprenorphine (0.1-mg/kg, i.p.), enrofloxacin (7.5-

mg/kg, i.p.), and ketoprofen (3-mg/kg, i.p.) during a 5-day postsurgical care period and allowed

one week in total to fully recover after surgery before the start of recordings.

In vivo electrophysiological recordings. Before implantation, impedance of tetrodes (17-μm

platinum wires, California Fine Wires) was adjusted by electroplating to < 150-kOhm. Tetrodes

were lowered daily by 50-μm until the abundance of spikes and the presence of theta rhythmic

cells in test recordings indicated that tetrode tips were located near or within the superficial cell

layers of the MEC. Neural signals were pre-amplified by unity-gain operational amplifiers

located on each animal's head (head stage) and AC-wire-coupled to an Axona recording system

(Axona Limited, St. Albans, UK). Spikes were threshold-detected and recorded at 48-kHz. The

LFP signal was recorded single-ended with the ground screw as reference at 4800-Hz. Position and head direction of mice were tracked with headstage-mounted invisible infrared LEDs at 50-Hz. Spikes were clustered into single units using spike clustering software (DacqUSB/Tint, Axona, St. Albans, UK). Further offline analyses were performed using MATLAB (MathWorks, Natick, MA, USA) and RStudio software packages.

Behavior during light and dark conditions. All recordings were performed while animals foraged for randomly scattered small pieces of Froot Loops (Kellog Company, Battle Creek, MI, USA) in a 1m square open-field arena with 60-cm high black walls and a white letter size cue card at one wall. Recordings were performed during alternating epochs of open field illumination (light) and complete darkness (dark). Alternating epochs of recordings in light and darkness could vary in duration between 30 seconds and 30 min; most experiments used 3-min alternating epochs allowing high statistical power due to multiple repetitions from light to darkness in each session. A typical recording session lasted 40-60 min. A ceiling light provided full illumination of the open field during the light condition. For complete darkness the ceiling light was turned off. All other possible light sources in the room had been eliminated and the open field was fully shielded from any remaining light (e.g. door slits) by a black curtain. Tracking of position and head direction was achieved by using invisible infrared LEDs. Experiments with alternating epochs of light and dark began with the light condition. Experiments with a dark-light-dark paradigm started in complete darkness. For those experiments, mice were brought into the room, plugged in and introduced into the arena during complete darkness by an experimenter wearing night vision goggles.

Histology. After the end of electrophysiological data collection, animals were deeply anaesthetized by isoflurane or i.p. injection of Euthasol (390-mg/kg), followed by electro-lesions (20-µA, 16s) through one channel of each tetrode and subsequent transcardial perfusion with saline followed by 10% buffered formalin (SF100-4, ThermoFisher Scientifc). Brains were extracted and stored in fixative for one day. Cresyl violet staining was performed on 50-µm sagittal slices of MEC to visualize tetrode tracks and confirm positions of recording sites in the MEC.

Data analysis.

   Computation of running speed and firing rate signals. The animal's running speed was estimated by applying a Kalman filter to the positional data obtained from the video tracking

sampled at 50-Hz (Fyhn et al., 2004; Hinman et al., 2016; Kropff et al., 2015). To obtain an

instantaneous firing rate signal matching the sampling rate of the behavioral tracking, we

calculated the firing rate as the number of spikes per video frame period (20-ms) and smoothed

this signal with a 125-ms Gaussian filter.

Speed score. The speed score was calculated as the Pearson's product moment

correlation between instantaneous firing rate and running speed (Kropff et al., 2015).

LFP theta rhythm. LFP data were down-sampled to 600-Hz and power spectral density

analyses were performed using custom written MATLAB scripts and the FieldTrip toolbox

(Oostenveld et al., 2011). A theta frequency range was determined for each session based on

the range of the theta peak in the power spectral density plot (Dannenberg et al., 2019). The

LFP signal was then band-pass-filtered at this theta frequency range with a third order

Butterworth filter, and the instantaneous theta amplitude and phase were determined via a

Hilbert transform. The instantaneous theta frequency could then be inferred from the

instantaneous phase. To analyze changes in the LFP theta frequency vs. running speed

relationship over time, we first down-sampled the LFP theta frequency to 50-Hz in order to

match the temporal binning of the behavioral tracking. We next sectioned data into 10-sec

blocks after transitions between light and dark and pooled data from all sessions and all mice

(Fig. 1I). For each of those 10-sec blocks of data, a speed tuning curve of LFP theta frequency

was computed at 1-cm/s resolution and the slope and y-intercept of the linear regression curve

were determined (Fig. 1J). The same analysis as described above was applied to 10-sec blocks

around the transition points from light to dark or from dark to light (Figs. 1K, L).

Theta rhythmic firing. We used maximum likelihood estimation (MLE) of a parametric

model of the lags within the spike time autocorrelogram to analyze theta rhythmic firing, which

has been proven a more sensitive and statistically rigorous method than traditional measures of

spike time autocorrelogram features (Climer et al., 2015). In particular, this model allowed us to

statistically test significance of theta modulation in MEC neurons and compare theta rhythmic

firing frequency and theta rhythmicity magnitude on a single cell level between light and dark

conditions. The model tested for rhythmic firing in the frequency range between 1-13 Hz. For the

analysis of the relationship between theta rhythmic firing and running speed (Fig. 2), we first

identified continuous temporal epochs larger than 400-ms, in which the running speed of the

animal varied less than ± 4-cm/s. We next sorted these epochs according to the average

running speed during the epochs into 1-cm/s bins from 1-cm/s to 29-cm/s and used the MLE

approach to fit an autocorrelogram model to each of those epochs. For each neuron and

running speed bin, an average autocorrelogram was computed across autocorrelograms for

each epoch. The maximal run time for the theta rhythmic firing vs. running speed analysis on

the research computing cluster was set to two weeks resulting in the successful analysis of 340

out of 342 neurons. For the analysis of the time course of changes in theta rhythmic firing after

transition from light to dark or from dark to light, the first two minutes after the transition were

sectioned into 10-sec blocks (n = 342 neurons, Fig. 2D). Alternately, 10-sec blocks around the

transition points from light to dark or from dark to light were analyzed (n = 342 neurons, Fig.

2E). For each neuron, an autocorrelogram model was fitted to each of those 10-sec blocks of

data using the MLE approach. For each of those 10-sec blocks of data, the average theta

rhythmic firing frequency across neurons could then be computed from the peaks of the average

autocorrelogram model across neurons by taking the inverse of the first peak in the

autocorrelogram model. To analyze running speed-dependent theta rhythmic firing on the

population level, we first identified continuous epochs of running speed within the ranges of 0-3

cm/s (immobility), 5-10 cm/s, 10-15 cm/s, 15-20 cm/s, and 20-25 cm/s. We next used the MLE

approach to model an autocorrelogram for each running speed bin of each neuron. Neurons

with sufficient number of spikes to compute an MLE-modeled autocorrelogram in at least one of

the running speed bins were included into the analysis (n = 339 neurons).

Computation of firing rate maps. Occupancy-normalized firing rate maps were

generated by dividing the open field environment into 3-cm spatial bins. For each spatial bin, an

occupancy-normalized firing rate was computed by dividing the total number of spikes falling

into that bin by the total time the animal spent in that bin. Firing rate maps were then smoothed

by a 3-cm wide two-dimensional Gaussian kernel.

Modulation of baseline firing rates by visual inputs.

In order to identify neurons, whose firing rates are modulated by the illumination of the

environment (consistently either higher or lower firing rates during light or darkness), we

computed the average firing rate for each light and dark epoch and tested if the differences

between light epochs and the following dark epochs and the differences between dark epochs

and the following light epochs reached statistical significance at alpha = 0.05 using Student's

paired t-test. To analyze the time-course of changes in baseline firing rates after transition from

light to dark or from dark to light, we first computed the peristimulus time histogram with 5-s

binning around the transition point and then computed the average of z-scored firing rates

across neurons.

Identification and classification of speed-modulated neurons. To identify speed

modulated neurons in an unbiased way, we used a linear non-linear Poisson (LN) model

(Hardcastle et al., 2017) which tested for significant modulation of firing rates by position, head

direction, LFP theta phase, and/or running speed (Fig. 4). For a robust analysis, we first

screened for cells which showed stable firing across the whole recording session. Cells, whose

mean firing rates were below 1-Hz or whose mean firing rates differed more than by a factor of

two between the first and second halves of the recording session, were excluded from the

analysis. Critically, we next adjusted for differences in the number of spikes between the light

and dark conditions by randomly deleting spikes for the condition with more spikes. Such

random deletion of spikes ensured that the statistical power to detect modulation of firing rates

by running speed, position, head direction and/or theta phase was identical for the light and dark

condition, and further ensured that differences in baseline firing rate between light and dark

conditions did not affect the further analysis shown in Fig. 4. To compare speed modulation of

firing rates on the population level, we clustered neurons into positively and negatively speed-

modulated neurons using hierarchical clustering (clustergram, bioinformatics toolbox, MATLAB).

Grid score. The grid score was used as a measure of hexagonal symmetry of grid

fields. The grid score was calculated as described in detail in Brandon et al. (2011) (gridness

measure 3). In short, we computed the two-dimensional rate map autocorrelogram and

computed correlations between this and rotated versions of itself after correction of elliptical

eccentricity. If hexagonal symmetry is present, those correlations will peak at multiples of 60

degrees. The difference between the smallest peak and the largest trough was then defined as

the grid score. Grid cells were defined as cells with a grid score > 0.85 in either light or dark

condition.

Spatial information. We defined the spatial information as the delta entropy $\Delta H =$

$H_{max} - H$. H_{max} is the maximal amount of randomness given by a completely random

distribution of spike locations in the open field environment defined by $H_{max} = -\sum_{i=1}^{n} p * \log_2 p$,

with $p = \frac{1}{n}$, where n is the number of spatial bins. H is the observed amount of randomness in

the distribution of spike locations given by $H(X|spike) = -\sum_{i=1}^{n} p(x_i|spike) * \log_2 p(x_i|spike)$,

where X is the animal's location, x_i is the ith spatial bin, n is the number of spatial bins and

$p(x_i|spike)$ is the probability that $X = x_i$ when a spike is observed. ΔH can then be understood

as an increase in the amount of order in the distribution of spike locations relative to a

completely random (uniform) distribution of spike locations.

Spatial correlation. Firing rate map similarities were quantified by the spatial correlation

between two firing rate maps. To compute the spatial correlation, the spatial bins of the firing

rate maps were transformed into a single vector. The spatial correlation was then computed as

the Pearson's correlation coefficient between two of those spatially binned firing rate vectors.

Spatiotemporal correlation. To analyze spatial stability of grid cell firing in the time

domain, we used the firing rate map of the light condition to compute the expected firing rate for

each of the 20-msec time bins of the animal's trajectory. We next computed the Pearson's

correlation coefficient between the observed 20-msec binned firing rate and the expected 20-

msec binned firing rate.

Experimental design and statistical analysis. Details on statistical analyses on each

experiment are reported in the Results section. Statistical tests used in this study include

Student's paired t-test and nonparametric statistical tests for repeated measures, namely

Wilcoxon's signed rank test and Friedman's test. Posthoc testing was performed if the Friedman

test yielded a significant result at alpha = 0.05.

Code accessibility. Custom MATLAB code used for data analysis can be found at

https://github.com/hasselmonians or can be made available upon request. Violin plots were

created using R software (R Core Team, 2019; R: A language and environment for statistical

computing; R Foundation for Statistical Computing, Vienna, Austria).

ACKNOWLEDGMENTS

This work was supported by the National Institute of Mental Health, grant numbers R01

MH60013 and R01 MH120073, by the National Institute of Neurological Disorders and Stroke,

grant number K99NS116129, and by the Office of Naval Research MURI N00014-16-1-2832

and N00014-19-1-2571, and Office of Naval Research DURIP N00014-17-1-2304.

COMPETING INTERESTS

None

REFERENCES

Barry, C., Hayman, R., Burgess, N., & Jeffery, K. J. (2007). Experience-dependent rescaling of
entorhinal grids. *Nature Neuroscience*, *10*(6), 682–684. https://doi.org/10.1038/nn1905
Blair, H. T., Gupta, K., & Zhang, K. (2008). Conversion of a phase- to a rate-coded position
signal by a three-stage model of theta cells, grid cells, and place cells. *Hippocampus*,
18(12), 1239–1255. https://doi.org/10.1002/hipo.20509
Brandon, M. P., Bogaard, A. R., Libby, C. P., Connerney, M. A., Gupta, K., & Hasselmo, M. E.
(2011). Reduction of theta rhythm dissociates grid cell spatial periodicity from directional
tuning. *Science (New York, N.Y.)*, *332*(6029), 595–599.
https://doi.org/10.1126/science.1201652
Burak, Y., & Fiete, I. R. (2009). Accurate path integration in continuous attractor network models
of grid cells. *PLoS Computational Biology*, *5*(2), e1000291.
https://doi.org/10.1371/journal.pcbi.1000291

Burgess, N. (2008). Grid cells and theta as oscillatory interference: Theory and predictions. *Hippocampus*, *18*(12), 1157–1174. https://doi.org/10.1002/hipo.20518

Burgess, N., Barry, C., & O'Keefe, J. (2007). An oscillatory interference model of grid cell firing. *Hippocampus*, *17*(9), 801–812. https://doi.org/10.1002/hipo.20327

Bush, D., & Burgess, N. (2014). A Hybrid Oscillatory Interference/Continuous Attractor Network Model of Grid Cell Firing. *Journal of Neuroscience*, *34*(14), 5065–5079. https://doi.org/10.1523/JNEUROSCI.4017-13.2014

Campbell, M. G., & Giocomo, L. M. (2018). Self-motion processing in visual and entorhinal cortices: Inputs, integration, and implications for position coding. *Journal of Neurophysiology*. https://doi.org/10.1152/jn.00686.2017

Campbell, M. G., Ocko, S. A., Mallory, C. S., Low, I. I. C., Ganguli, S., & Giocomo, L. M. (2018). Principles governing the integration of landmark and self-motion cues in entorhinal cortical codes for navigation. *Nature Neuroscience*, *21*(8), 1096–1106. https://doi.org/10.1038/s41593-018-0189-y

Chen, G., King, J. A., Lu, Y., Cacucci, F., & Burgess, N. (2018). Spatial cell firing during virtual navigation of open arenas by head-restrained mice. *ELife*, *7*. https://doi.org/10.7554/eLife.34789

Chen, G., Lu, Y., King, J. A., Cacucci, F., & Burgess, N. (2019). Differential influences of environment and self-motion on place and grid cell firing. *Nature Communications*, *10*(1), 630. https://doi.org/10.1038/s41467-019-08550-1

Chen, G., Manson, D., Cacucci, F., & Wills, T. J. (2016). Absence of Visual Input Results in the Disruption of Grid Cell Firing in the Mouse. *Current Biology: CB*. https://doi.org/10.1016/j.cub.2016.06.043

Cheung, A. (2016). Probabilistic Learning by Rodent Grid Cells. *PLoS Computational Biology*, *12*(10), e1005165. https://doi.org/10.1371/journal.pcbi.1005165

Chrobak, J. J., Stackman, R. W., & Walsh, T. J. (1989). Intraseptal administration of muscimol produces dose-dependent memory impairments in the rat. *Behavioral and Neural Biology*, *52*(3), 357–369.

Climer, J. R., DiTullio, R., Newman, E. L., Hasselmo, M. E., & Eden, U. T. (2015). Examination of rhythmicity of extracellularly recorded neurons in the entorhinal cortex. *Hippocampus*, *25*(4), 460–473. https://doi.org/10.1002/hipo.22383

Dannenberg, H., Kelley, C., Hoyland, A., Monaghan, C. K., & Hasselmo, M. E. (2019). The Firing Rate Speed Code of Entorhinal Speed Cells Differs across Behaviorally Relevant Time Scales and Does Not Depend on Medial Septum Inputs. *The Journal of Neuroscience*, *39*(18), 3434–3453. https://doi.org/10.1523/JNEUROSCI.1450-18.2019

Dannenberg, H., Pabst, M., Braganza, O., Schoch, S., Niediek, J., Bayraktar, M., Mormann, F., & Beck, H. (2015). Synergy of direct and indirect cholinergic septo-hippocampal pathways coordinates firing in hippocampal networks. *The Journal of Neuroscience*, *35*(22), 8394–8410. https://doi.org/10.1523/JNEUROSCI.4460-14.2015

Evans, T., Bicanski, A., Bush, D., & Burgess, N. (2016). How environment and self-motion combine in neural representations of space. *The Journal of Physiology*, *594*(22), 6535–6546. https://doi.org/10.1113/JP270666

Fuhrmann, F., Justus, D., Sosulina, L., Kaneko, H., Beutel, T., Friedrichs, D., Schoch, S., Schwarz, M. K., Fuhrmann, M., & Remy, S. (2015). Locomotion, Theta Oscillations, and the Speed-Correlated Firing of Hippocampal Neurons Are Controlled by a Medial Septal Glutamatergic Circuit. *Neuron*, *86*(5), 1253–1264. https://doi.org/10.1016/j.neuron.2015.05.001

Fuhs, M. C., & Touretzky, D. S. (2006). A spin glass model of path integration in rat medial entorhinal cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *26*(16), 4266–4276. https://doi.org/10.1523/JNEUROSCI.4353-05.2006

Fyhn, M., Hafting, T., Witter, M. P., Moser, E. I., & Moser, M.-B. (2008). Grid cells in mice. *Hippocampus*, *18*(12), 1230–1238. http://dx.doi.org/10.1002/hipo.20472

Fyhn, M., Molden, S., Witter, M. P., Moser, E. I., & Moser, M.-B. (2004). Spatial representation in the entorhinal cortex. *Science*, *305*(5688), 1258–1264. http://dx.doi.org/10.1126/science.1099901

Gil, M., Ancau, M., Schlesiger, M. I., Neitz, A., Allen, K., De Marco, R. J., & Monyer, H. (2018). Impaired path integration in mice with disrupted grid cell firing. *Nature Neuroscience*, *21*(1), 81–91. https://doi.org/10.1038/s41593-017-0039-3

Giocomo, L. M. (2016). Environmental boundaries as a mechanism for correcting and anchoring spatial maps. *The Journal of Physiology*, *594*(22), 6501–6511. https://doi.org/10.1113/JP270624

Givens, B., & Olton, D. S. (1994). Local modulation of basal forebrain: Effects on working and reference memory. *The Journal of Neuroscience*, *14*(6), 3578–3587.

Hafting, T., Fyhn, M., Molden, S., Moser, M.-B., & Moser, E. I. (2005). Microstructure of a spatial map in the entorhinal cortex. *Nature*, *436*(7052), 801–806. http://dx.doi.org/10.1038/nature03721

Hardcastle, K., Ganguli, S., & Giocomo, L. M. (2015). Environmental Boundaries as an Error Correction Mechanism for Grid Cells. *Neuron*, *86*(3), 827–839. https://doi.org/10.1016/j.neuron.2015.03.039

Hardcastle, K., Maheswaranathan, N., Ganguli, S., & Giocomo, L. M. (2017). A Multiplexed, Heterogeneous, and Adaptive Code for Navigation in Medial Entorhinal Cortex. *Neuron*, *94*(2), 375-387.e7. https://doi.org/10.1016/j.neuron.2017.03.025

Hasselmo, M. E. (2008). Grid cell mechanisms and function: Contributions of entorhinal persistent spiking and phase resetting. *Hippocampus*, *18*(12), 1213–1229. https://doi.org/10.1002/hipo.20512

Hinman, J. R., Brandon, M. P., Climer, J. R., Chapman, G. W., & Hasselmo, M. E. (2016). Multiple Running Speed Signals in Medial Entorhinal Cortex. *Neuron*, *91*(3), 666–679. https://doi.org/10.1016/j.neuron.2016.06.027

Hinman, J. R., Penley, S. C., Long, L. L., Escabí, M. A., & Chrobak, J. J. (2011). Septotemporal variation in dynamics of theta: Speed and habituation. *Journal of Neurophysiology*, *105*(6), 2675–2686. https://doi.org/10.1152/jn.00837.2010

Jacob, P.-Y., Capitano, F., Poucet, B., Save, E., & Sargolini, F. (2019). Path integration maintains spatial periodicity of grid cell firing in a 1D circular track. *Nature Communications*, *10*(1), 1–13. https://doi.org/10.1038/s41467-019-08795-w

Jacob, P.-Y., Gordillo-Salas, M., Facchini, J., Poucet, B., Save, E., & Sargolini, F. (2017). Medial entorhinal cortex and medial septum contribute to self-motion-based linear distance estimation. *Brain Structure & Function*, *222*(6), 2727–2742. https://doi.org/10.1007/s00429-017-1368-4

Jacob, P.-Y., Poucet, B., Liberge, M., Save, E., & Sargolini, F. (2014). Vestibular control of entorhinal cortex activity in spatial navigation. *Frontiers in Integrative Neuroscience*, *8*, 38. https://doi.org/10.3389/fnint.2014.00038

Justus, D., Dalügge, D., Bothe, S., Fuhrmann, F., Hannes, C., Kaneko, H., Friedrichs, D., Sosulina, L., Schwarz, I., Elliott, D. A., Schoch, S., Bradke, F., Schwarz, M. K., & Remy, S. (2016). Glutamatergic synaptic integration of locomotion speed via septoentorhinal projections. *Nature Neuroscience*. https://doi.org/10.1038/nn.4447

Koenig, J., Linder, A. N., Leutgeb, J. K., & Leutgeb, S. (2011). The spatial periodicity of grid cells is not sustained during reduced theta oscillations. *Science (New York, N.Y.)*, *332*(6029), 592–595. https://doi.org/10.1126/science.1201685

Kraus, B. J., Brandon, M. P., Robinson, R. J., Connerney, M. A., Hasselmo, M. E., & Eichenbaum, H. (2015). During Running in Place, Grid Cells Integrate Elapsed Time and Distance Run. *Neuron*, *88*(3), 578–589. https://doi.org/10.1016/j.neuron.2015.09.031

Kropff, E., Carmichael, J. E., Moser, M.-B., & Moser, E. I. (2015). Speed cells in the medial entorhinal cortex. *Nature*, *523*(7561), 419–424. https://doi.org/10.1038/nature14622

Krupic, J., Bauza, M., Burton, S., Barry, C., & O'Keefe, J. (2015). Grid cell symmetry is shaped by environmental geometry. *Nature*, *518*(7538), 232–235. https://doi.org/10.1038/nature14153

Lawson, V. H., & Bland, B. H. (1993). The role of the septohippocampal pathway in the regulation of hippocampal field activity and behavior: Analysis by the intraseptal microinfusion of carbachol, atropine, and procaine. *Experimental Neurology*, *120*(1), 132–144. https://doi.org/10.1006/exnr.1993.1047

McNaughton, B. L., Battaglia, F. P., Jensen, O., Moser, E. I., & Moser, M.-B. (2006). Path integration and the neural basis of the "cognitive map". *Nat Rev Neurosci*, *7*(8), 663–678. http://dx.doi.org/10.1038/nrn1932

Munn, R. G. K., Mallory, C. S., Hardcastle, K., Chetkovich, D. M., & Giocomo, L. M. (2020). Entorhinal velocity signals reflect environmental geometry. *Nature Neuroscience*, *23*(2), 239–251. https://doi.org/10.1038/s41593-019-0562-5

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, *2011*, 156869. https://doi.org/10.1155/2011/156869

Pérez-Escobar, J. A., Kornienko, O., Latuske, P., Kohler, L., & Allen, K. (2016). Visual landmarks sharpen grid cell metric and confer context specificity to neurons of the medial entorhinal cortex. *ELife*, *5*. https://doi.org/10.7554/eLife.16937

Raudies, F., Brandon, M. P., Chapman, G. W., & Hasselmo, M. E. (2015). Head direction is coded more strongly than movement direction in a population of entorhinal neurons. *Brain Research*, *1621*, 355–367. https://doi.org/10.1016/j.brainres.2014.10.053

Raudies, F., & Hasselmo, M. E. (2015). Differences in Visual-Spatial Input May Underlie Different Compression Properties of Firing Fields for Grid Cell Modules in Medial Entorhinal Cortex. *PLoS Computational Biology*, *11*(11), e1004596. https://doi.org/10.1371/journal.pcbi.1004596

Raudies, F., Mingolla, E., & Hasselmo, M. E. (2012). Modeling the influence of optic flow on grid cell firing in the absence of other cues1. *Journal of Computational Neuroscience*, *33*(3), 475–493. https://doi.org/10.1007/s10827-012-0396-6

Sargolini, F., Fyhn, M., Hafting, T., McNaughton, B. L., Witter, M. P., Moser, M.-B., & Moser, E. I. (2006). Conjunctive representation of position, direction, and velocity in entorhinal cortex. *Science (New York, N.Y.)*, *312*(5774), 758–762. https://doi.org/10.1126/science.1125572

Sreenivasan, S., & Fiete, I. (2011). Grid cells generate an analog error-correcting code for singularly precise neural computation. *Nature Neuroscience*, *14*(10), 1330–1337. https://doi.org/10.1038/nn.2901

Taube, J. S. (2007). The Head Direction Signal: Origins and Sensory-Motor Integration. *Annual Review of Neuroscience*, *30*(1), 181–207. https://doi.org/10.1146/annurev.neuro.29.051605.112854

Taube, J. S., Muller, R. U., & Ranck, J. B. (1990a). Head-direction cells recorded from the postsubiculum in freely moving rats. II. Effects of environmental manipulations. *The Journal of Neuroscience*, *10*(2), 436–447.

Taube, J. S., Muller, R. U., & Ranck, J. B. (1990b). Head-direction cells recorded from the postsubiculum in freely moving rats. II. Effects of environmental manipulations. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *10*(2), 436–447.

Tennant, S. A., Fischer, L., Garden, D. L. F., Gerlei, K. Z., Martinez-Gonzalez, C., McClure, C., Wood, E. R., & Nolan, M. F. (2018). Stellate Cells in the Medial Entorhinal Cortex Are

868   Required for Spatial Learning. *Cell Reports*, *22*(5), 1313–1324.
869   https://doi.org/10.1016/j.celrep.2018.01.005
870 Winter, S. S., Mehlman, M. L., Clark, B. J., & Taube, J. S. (2015). Passive Transport Disrupts
871   Grid Signals in the Parahippocampal Cortex. *Current Biology: CB*, *25*(19), 2493–2502.
872   https://doi.org/10.1016/j.cub.2015.08.034
873 Zilli, E. A., & Hasselmo, M. E. (2010). Coupled Noisy Spiking Neurons as Velocity-Controlled
874   Oscillators in a Model of Grid Cell Spatial Firing. *Journal of Neuroscience*, *30*(41),
875   13850–13860. https://doi.org/10.1523/JNEUROSCI.0547-10.2010
876

FIGURES

Figure 1

Figure 1. Speed signal by local field potential theta frequency decreases in darkness. A,
Experimental paradigm. Local field potential (LFP) theta oscillatory activity was recorded in mice
freely foraging in a 1-m square open field environment containing a visual cue card during
alternating epochs of illumination by a ceiling light (light) and complete darkness (dark). A
typical recording lasted 42-min with 3-min alternating light and dark epochs. **B,** Plot of theta
frequency vs. running speed shows that the slope of the relationship decreases during
darkness. Solid lines show the mean, shaded area show the s.e.m. for n = 15 mice. Data on the
light condition are shown in red, data on the dark condition are shown in black and grey. **C,**
Same as in B, but for theta amplitude. **D,** Running speed decreases slightly during darkness.
Each data point shows the mean value for one mouse, box plots show mean, 25^{th} and 75^{th}
percentiles, grey lines connect data from the same mouse; t = 2,90, df = 14, p = 0.0117,
Student's paired t-test. **E,** Running acceleration decreases slightly during darkness. Data
visualization as in D; t = 3.16, df = 14, p = 0.0070, Student's paired t-test. **F,** LFP theta
frequency vs. running acceleration relationship is reduced for all acceleration values in complete
darkness. Grey background highlights the interval of acceleration values between -5 cm/s^2 and
5 cm/s^2. **G,** LFP theta frequency vs. running speed relationship based on the subset of time
points where the running acceleration is near zero in the interval shown in F between -5 cm/s^2
and 5 cm/s^2. **H-L,** Figure panels show sequential 10-sec bins of LFP data after transitions
between light and dark. Each data point shows the mean across all epochs of the same
condition (light or dark) from 73 recording sessions from 13 mice. Yellow background indicates
the light condition; grey background indicates the dark condition. **H,** Theta frequency speed
tuning curves for subsequent 10-sec blocks after transitions between light and dark. Straight
lines are the linear regression lines to the data. **I,** Data points (with errorbars) show the mean (±
s.e.m.) of theta frequency for the first three minutes in light and dark summed over data from all
mice in 10-sec bins (errorbars are within the line width). **J,** Solid and dashed lines show the
means and 95% confidence intervals of the slopes of the theta frequency vs. running speed
relationships for sequential 10-sec bins for the first three minutes in light and dark; same
visualization scheme as in I. **K,** Left and right panels show 10-sec binned mean ± s.e.m of theta
frequency around the transition points from light to dark and from dark to light, respectively
(errorbars are within the line width). **L,** Left and right panels show 10-sec binned peristimulus
time histograms of the slopes of the theta frequency vs. running speed relationships around the
transition points from light to dark and dark to light, respectively. Solid and dashed lines show
means and 95% confidence intervals. Same visualization scheme as in K.

Figure 2

Figure 2. Changes in theta rhythmic firing with running speed are differentially affected

by visual inputs. Red color and/or yellow background indicate data on the light condition, black

and grey color and/or grey background indicate data on the dark condition. Figures show data

from theta rhythmic neurons from a total of 14 mice. **A,** Autocorrelogram of one example neuron

showing theta rhythmic firing in light and dark. Solid lines show the MLE model of the data (see

Methods). Dashed lines indicate the positions of the peaks in the autocorrelogram. Larger lags

between peaks translate into a lower frequency of theta rhythmic firing. **B,** Figure shows color-

coded mean autocorrelogram fits across neurons per running speed bin. Speed bins range from

1 ±4 cm/s cm/s to 29 ±4 cm/s (see Methods). Upper panel shows data for the light condition,

lower panel shows data for the dark condition. Dashed white line connects the first peaks in the

mean autocorrelogram models. n = 340 neurons. **C,** Left panel shows the color-coded mean

autocorrelogram models averaged across neurons for sequential 10-sec time bins of data for all

light epochs and dark epochs. Solid white lines show the peaks and troughs in the

autocorrelogram models. Data on the light condition are duplicated for better visualization. Right

panel shows the averages of autocorrelogram models across neurons and across 10-sec time

bins comparing light vs. dark. n = 342 neurons. **D,** Time course of the mean theta rhythmic firing

frequency across neurons for sequential 10-sec time bins in the light and dark conditions. Data

are plotted as mean ± s.e.m; n = 342 neurons. **E,** Left and right panels show data on the

transition periods from light to dark and dark to light, respectively. Upper panels show color-

coded mean autocorrelogram models per 10-sec bins before and after the transition point.

White lines mark the peaks and troughs in the autocorrelogram models. Lower panels show the

theta rhythmic firing frequency averaged across neurons for sequential 10-sec time bins before

and after the transition point. n = 342 neurons. **F,** Upper panels show the color coded MLE

autocorrelogram models for each neuron and different running speeds. Immobility was defined

as running speed < 3 cm/s. Neurons are sorted per condition in order of their theta rhythmic

firing frequency. White points indicate the first peak of the autocorrelogram fit in neurons

showing statistically significant theta modulation in the corresponding condition at alpha = 0.1.

Lower row shows the average across autocorrelogram models per condition. Note that higher

synchrony across neurons results in higher rhythmicity magnitude in the average across

neurons, n = 339. **G**, Number of neurons shown in F with statistically significant spiking

rhythmicity at alpha = 0.1 determined by the MLE model (see Methods) for different running

speeds in light and dark. **H,** Violin plots show the distribution of theta rhythmicity magnitudes of

neurons in G for different running speeds comparing light and dark conditions. Numbers show

the p-values for Kolmogorov-Smirnov tests comparing data between light and dark. See Results

for number of neurons per condition. **I,** Same as H, but for theta rhythmic firing frequency.

Figure 3

Figure 3. **Visual inputs alter baseline firing rates of neurons in the medial entorhinal cortex. A-D,** Yellow background indicates duration of light epochs, grey background indicates duration of dark epochs. **A,** Data on firing rate from one example neuron in the medial entorhinal cortex (MEC) with its baseline firing rate negatively modulated by darkness. Black lines show the firing rate at 1-s resolution. Blue lines show the average firing rate per light or dark epoch. **B,** Data on firing rate from one example neuron in the MEC, whose baseline firing rate is positively modulated by darkness; same visualization scheme as in A. **C,** Left and right panels show the mean z-scored firing rate across n = 139 neurons with significantly negative modulation of firing rates by darkness (see Methods) in 5-s bins around the transition point from light to dark and dark to light, respectively. Red color indicates data from the light condition, black and grey color indicate data from the dark condition. Solid lines and shaded areas show mean and SD. **D,** Same as C, but for neurons with positive modulation of baseline firing rates by darkness; n = 32. **E, F,** Yellow background indicates light condition, green background indicates the presence of an egocentrically stable spot of green laser light above the animal's head during the standard light condition. **E,** Data on one example neuron in the MEC showing positive modulation of baseline firing rate in presence of the green laser light spot. Same visualization scheme as in A and C. **F,** As in E, but for one example neuron with negative modulation of baseline firing rate.

Figure 4

Figure 4. Speed modulation of firing rates is maintained in the absence of visual inputs.

A, Left and right panels show speed response curves for the n = 142 neurons passing statistical significance of speed modulation in the LN-model (see Methods) for either the light or dark condition, respectively. For comparison of speed modulation across neurons and between light and dark conditions, firing rates were normalized and cells are plotted in the same order for light and dark. Cells were sorted by the location of peak firing on the running speed axis averaged across light and dark conditions. **B,** Clustergram visualizing the results of the hierarchical clustering of neurons by similarity of speed modulation of firing rates. The clustering was performed on the neurons' average speed response curves across light and dark conditions. **C,**

Left and right panels show speed response curves of negatively (n = 19) and positively (n =

123) modulated neurons; solid lines and shaded areas show mean ± s.e.m across neurons. **D,**

Histogram of speed scores for all neurons showing significant speed modulation in either the

light or dark condition. Red and grey color indicates data on light and dark, respectively. **E,**

Scatter plot on same data shown in D comparing speed scores during the light with speed

scores during the dark condition. **F,** Percentage of cells showing conjunctive coding of running

speed and either position, head direction, or LFP theta phase (spike-phase locking) during the

light and dark condition.

Figure 5

Figure 5. Grid cell spatial firing is maintained, but less stable in the absence of visual inputs. A, Firing rate maps of three example neurons during sequential dark, light, and dark sessions of 10-20 min duration. In the initial dark session, the animal was introduced to the open field environment in complete darkness. **B,** Box plots of grid scores during sequential dark, light, and dark session, n = 9 neurons from two mice. Black color and grey background indicate data on dark conditions, red color and yellow background indicate data on the light condition. Solid grey lines connect data points from the same grid cell. Chi-square = 10.89, p = 0.0043, Friedman's test; post-hoc tests revealed significant differences between the initial dark and light sessions as well as the initial dark and second dark sessions at alpha = 0.05. **C,** Top, middle, and bottom rows show trajectory plots, firing rate maps, and spatial autocorrelations of firing rate maps for three example grid cells recorded over multiple days during alternating light and dark epochs. Solid black lines in the trajectory plot show the trajectory of the animal, red dots mark the location of spikes during that trajectory. **D-E,** Box plots of grid scores, spatial information, and spatiotemporal correlation during light and dark; grey solid lines connect data points from the same grid cells; p = 7.4 x 10^{-5}, p = 9.8 x 10^{-6}, and p = 2.4 x 10^{-5}; n = 27, Wilcoxon's signed rank test.

Figure 6

Figure 6. Changes in grid cell spatial stability due to changes in visual inputs has a slow exponential component at a behavioral time scale and correlates with changes in the speed signal by local field potential theta frequency. A, Firing rate maps for sequential 10-sec bins of data after transitioning from dark to light (left side, yellow background) and light to dark (right side, grey background) for one example neuron. **B-D,** Time course measured in 10-sec bins of grid scores (B), spatial correlation (C), and spatiotemporal correlation (D) during the first three minutes of light and dark sessions. Solid lines show mean firing rates, shaded area indicates s.e.m.; n = 27 cells from three mice. **E,** Left and right panels show 10-sec binned data on spatiotemporal correlation around the transition points from light to dark and dark to light, respectively. Solid lines and shaded area show mean ± s.e.m.; n =27 grid cells from three mice. **F,** Overlay of data shown in Figures 6D and 1J showing the temporal alignment of changes in the slope of the theta frequency vs. running speed relationship (blue colors) and the spatial stability of grid cell firing measured by the spatiotemporal correlation (ocher colors). Left and right panels show data on first three minutes in the light and dark conditions, respectively.