TITLE

Effects of visual inputs on neural dynamics for coding of location and running speed in medial entorhinal cortex.

AUTHORS

Holger Dannenberg[1], Hallie Lazaro[1], Pranav Nambiar[1], Alec Hoyland[1], Michael E. Hasselmo[1]

INSTITUTIONAL AFFILIATIONS

[1]Center for Systems Neuroscience, Department of Psychological and Brain Sciences, Boston University, Boston, MA 02215, USA

ABSTRACT

Neuronal representations of spatial location and movement speed in the medial entorhinal cortex during the "active" theta state of the brain are important for memory-guided navigation and rely on visual inputs. However, little is known about how visual inputs change neural dynamics as a function of running speed and time. By manipulating visual inputs in mice, we demonstrate that changes in spatial stability of grid cell firing as a function of time correlate with changes in a proposed speed signal by local field potential theta frequency. In contrast, visual inputs do not affect the speed modulation of firing rates. Moreover, we provide evidence that sensory inputs other than visual inputs can support grid cell firing, though less accurately, in complete darkness. Finally, changes in spatial accuracy of grid cell firing on a 10-s time scale suggest that grid cell firing is a function of velocity signals integrated over past time.

INTRODUCTION

The spatially periodic firing of grid cells in the medial entorhinal cortex (MEC) (Fyhn et al., 2004, 2008; Hafting et al., 2005) is widely recognized as a neuronal correlate of memory-guided and/or self-motion-based spatial navigation (McNaughton et al., 2006). Based on that idea, computational models of grid cells including attractor network models (Burak & Fiete, 2009; Fuhs & Touretzky, 2006), oscillatory interference models (Blair et al., 2008; Burgess, 2008; Burgess et al., 2007; Hasselmo, 2008; Zilli & Hasselmo, 2010), hybrid models of attractor networks and oscillatory interference (Bush & Burgess, 2014), or probabilistic learning models (Cheung, 2016) suggest that grid cells integrate running direction and traveled distance to maintain a spatially periodic firing pattern during path integration (Gil et al., 2018; Jacob et al., 2019). Path integration depends on a velocity signal, which depends on vestibular and proprioceptive signals (Chen et al., 2019; Tennant et al., 2018) and also moving sensory

features such as optic flow (Campbell et al., 2018; Raudies et al., 2012; Raudies & Hasselmo,
2015). However, any path integration mechanism ultimately suffers from the accumulation of
error over distance traveled or elapsed time (Kraus et al., 2015; Sreenivasan & Fiete, 2011)
necessitating error correction by static environmental sensory inputs (Evans et al., 2016;
Giocomo, 2016) such as visual landmark cues (Campbell et al., 2018; Hafting et al., 2005) or
environmental boundaries (C. Barry et al., 2007; Hardcastle et al., 2015) to maintain an
accurate representation of relative self-location within a spatial environment. Experimental data
from studies on rats support the hypothesized role of grid cells in performing self-motion based
path integration and error correction by sensory cues: Self-motion-based signals such as head
direction are necessary for grid cell firing (Campbell et al., 2018; Winter, Mehlman, et al., 2015),
and grid cell firing persists during darkness with reduced spatial selectivity (Allen et al., 2014;
Hafting et al., 2005) reflecting the loss of error correction by visual cues. In contrast, two more
recent studies on mice reported a rapid and nearly complete loss of spatially periodic grid cell
firing during complete darkness (Chen et al., 2016; Pérez-Escobar et al., 2016) suggesting that
spatially periodic grid cell firing and path integration depend on error correction by visual cues.
However, motion-related interoceptive representations, such as proprioceptive, vestibular, and
motor efference copy signals, can combine with visual representations to form a single
multimodal representation supporting path integration (Tcheang et al., 2011) suggesting that
sensory cues other than visual cues may be sufficient to stabilize grid cells and support path
integration. In this study, we therefore investigated if and how the selective removal of visual
inputs affect grid cell firing in mice under conditions which leave other landmark features such
as walls, auditory, and olfactory cues still available to the animal.
58
Errors in path integration originate primarily in the velocity signal (Stangl et al., 2020), which
combines information about movement direction and movement speed. However, the source of
both the directional signal and the speed signal are still under debate. The direction signal has
been attributed to the head direction signal terminating in the MEC (Sargolini et al., 2006; Taube
et al., 1990; Taube et al., 1990; Taube, 2007), but head direction does not always match
movement direction (Raudies et al., 2015). Likewise, the dominant source of the speed signal is
still under debate. One potential candidate for the speed signal is the linear modulation of firing
rates in a subpopulation of "speed cells" in the MEC (Kropff et al., 2015). However, such a
speed signal by firing rate has been shown to be more heterogeneous than previously assumed
(Hardcastle et al., 2017) and not accurate on sub-second-long time scales (Dannenberg et al.,
2019) as would be required for updating grid cell firing in real time. An alternative or

complementary oscillatory speed signal is provided by the positive correlation between the theta
rhythmic firing frequency of theta modulated MEC neurons and the running speed of an animal
and/or the positive correlation between the local field potential (LFP) theta frequency and the
running speed of an animal (Hinman et al., 2016; Jeewajee, Barry, et al., 2008; McFarland et
al., 1975; Rivas et al., 1996; Shin & Talnov, 2001; Sławińska & Kasicki, 1998; Whishaw &
Vanderwolf, 1973). The slope of the LFP theta frequency vs. running speed relationship in
particular has been linked to spatial cognition and locomotion in both hippocampus and MEC
(Korotkova et al., 2018; Monaghan et al., 2017; Wells et al., 2013). Interestingly, environmental
novelty selectively reduces the slope of the LFP theta vs. running speed relationship
(Jeewajee, Lever, et al., 2008; Wells et al., 2013) and also results in an increase in grid scale
(C. Barry et al., 2012) as predicted by oscillatory interference models of grid cell firing (Burgess,
2008). In contrast, the y-intercept of the LFP theta vs. running speed relationship has been
linked to emotion-sensitivity, in particular anxiolysis (Korotkova et al., 2018; Monaghan et al.,
2017; Wells et al., 2013). Moreover, little is known about the time courses of changes in coding
of location by grid cells and running speed by theta oscillations or firing rate (for instance during
changes from light to darkness), although changes in grid cell firing, theta rhythmic dynamics,
and speed modulation of firing rates may underlie many cognitive functions on a behavioral time
scale ranging from seconds to minutes. By testing the temporal correlations between grid cell
spatial stability and the two proposed speed signals—by firing rate or theta oscillations—we test
if observed changes in grid cell firing can be explained by changes in the speed signal by firing
rate or changes in the speed signal by LFP theta frequency. We further test if changes in the
representation of a spatial metric by grid cells in the MEC is accompanied by similar changes in
the tuning of head direction cells, border cells, or both. Data presented in this study characterize
the visual input-evoked changes in neural dynamics in the MEC. These data are relevant for
models of path integration and for our understanding of how visual inputs modulate the coding
of location and running speed in the entorhinal cortex.
96

RESULTS

To study changes in neural dynamics after transitioning from foraging in an illuminated
environment (light) to foraging in complete darkness (dark) and vice versa, we recorded single
unit and LFP activity in mice freely exploring an open field environment during alternating light
and dark epochs (**Figure 1A**). We first studied if the absence of visual inputs in the dark
condition changed the correlation between LFP theta frequency and running speed. We found a
large change in the slope and a smaller change in the y-intercept of the LFP theta frequency vs.

running speed relationship during the dark condition (t-statistic of interaction effect (slope):

18.045, df = 806, p = 2×10^{-61}; t-statistic of change in y-intercept: -3.676, df = 806, p = 0.0003; n

= 15 mice) (**Figure 1B**). In contrast, the theta amplitude vs. running speed relationship did not

change significantly (t-statistic of interaction effect (slope): 1.7343, df = 806, p = 0.0832; t-

statistic of change in y-intercept: -1.764, p = 0.0781; n = 15 mice) (**Figure 1C**). Average running

speed during light was 11.6 ± 0.5 cm/s which slightly decreased after transition to dark by 0.89 ±

0.31 cm/s (7.7%) to 10.72 ± 1.4 cm/s (mean ± s.e.m., p = 0.0117, n = 15 mice, Wilcoxon's

signed rank test) (**Figure 1D** and **Figure 1—figure supplement 1A**). Similarly, running

acceleration also slightly decreased from 20.8 ± 1.9 cm/s^2 in the light condition by 1.7 ± 0.7

cm/s^2 (8.0%) to 19.1 ± 2.3 cm/s^2 in the dark condition (p = 0.0070, n = 15 mice, Wilcoxon's

signed rank test) (**Figure 1E**). We therefore examined if the observed decrease in the slope of

the LFP theta frequency vs. running speed relationship might result from the decrease in

running *acceleration* during darkness. If the decrease in slope was due to a decrease in running

acceleration, there should be no change in the y-intercept of the correlation between LFP theta

frequency and running *acceleration* during the dark condition. However, the y-intercept of the

LFP theta frequency vs. running acceleration relationship did change significantly in darkness (t-

statistic: 8.115, p =1.16×10^{-6}, n = 15 mice, Student's paired t-test) (**Figure 1F**)**.** Furthermore,

restricting our analysis to time points when running acceleration was near zero (0 ± 5 cm/s^2)

resulted in speed tuning curves of LFP theta frequency (**Figure 1G**) that looked nearly identical

to the ones obtained without that restriction (**Figure 1B**). These data demonstrate that the LFP

theta frequency is directly modulated by running speed and that the observed change in the

slope of the LFP theta frequency vs. running speed relationship in darkness is not an artifact of

an underlying change in running behavior but instead directly reflects a change of visual inputs.

The LFP theta frequency vs. running speed relationship has been hypothesized to serve as a

speed signal important for coding of location during self-motion-based navigation including path

integration. Having established that the removal of visual inputs changes the speed signal by

LFP theta frequency, we were further interested in the time-course of that change. We therefore

analyzed sequential 10-s blocks after the start of light and dark epochs. Averaging data from 78

sessions from 14 mice, we found that the LFP theta frequency vs. running speed relationships

in light and dark became more different with elapsed time after start of light and dark epochs at

a time scale of tens of seconds (**Figure 1H** and **Figure 1—figure supplement 1B**). This

divergence appeared to be largely driven by a change in the slope of the theta frequency vs.

running speed relationship as opposed to changes in the y-intercept. We therefore quantified

the time courses of those changes in slope and y-intercept during light and dark conditions. This
analysis revealed a slow increase in slope over time during the light condition (Pearson's R =
0.79, p = 0.0001) and a slow decrease in slope over time during the dark condition (Pearson's R
= -0.79, p = 0.0001) (**Figure 1I**). The increase in slope during the light condition was
accompanied by a small, yet significant, decrease in the y-intercept (Pearson's R = -0.77, p =
0.0002) (**Figure 1J**). In contrast, the y-intercept did not change significantly over time during the
dark condition (Pearson's R = -0.16, p = 0.5251) (**Figure 1J**). In addition to those slow
components of change, analyzing the transition points between light and dark conditions, also
revealed fast components of changes in slope and y-intercept (**Figs. 1K & L**). Taken together,
transitioning from light to dark resulted in an initial fast decrease in slope followed by a further
slow decrease (**Figure 1K**). Conversely, transitioning back from dark to light resulted in a fast
increase in slope followed by a further slow increase (**Figure 1K**). Similarly, the y-intercept
showed a fast decrease with the transition from light to dark and a fast increase when
transitioning from dark to light.
152
Local field potential activity in the MEC primarily reflects the summed activity of synaptic inputs
to the MEC, which is likely to be correlated with firing patterns of MEC neurons. Many neurons
in the MEC show a theta rhythmic firing pattern and—similarly to the LFP theta frequency—the
frequency of theta rhythmic firing increases with running speed (Hinman et al., 2016). In our
data set, we identified a total of n = 342 neurons from 14 mice as theta modulated based on
their autocorrelograms. We asked if and how theta rhythmic firing in these theta modulated
MEC neurons is affected by removing and reinstating all visual inputs. Towards that aim we
used an MLE approach (Climer et al., 2015) to fit a model of theta spiking rhythmicity to the
observed spike train autocorrelations and identify the frequency and magnitude of theta
rhythmic firing in theta modulated MEC neurons in light and dark conditions (see **Figure 2A** for
one example neuron). We observed a decrease in theta rhythmic firing frequency during the
dark condition (**Figure 2B**). Similar to the observed changes in the LFP theta frequency, the
decrease in theta rhythmic firing frequency was largely attributed to a decrease in the slope of
its relationship to running speed (**Figure 2C**) as previously shown in rats (Hinman et al., 2016).
167
However, averaging data across cells can hide differences between cells and important
characteristics of the population response. We therefore analyzed changes in theta rhythmic
firing on the single cell and population level for different running speeds (immobility, defined as
running speed < 3 cm/s; 5-10 cm/s; 10-15 cm/s; 15-20 cm/s; 20-25 cm/s) (n = 339 neurons,

Figure 2D). To our surprise, we found that neurons continued to fire rhythmically during

immobility periods in both light and dark conditions. However, theta rhythmic frequencies varied

largely across neurons within a wide range of theta frequencies (4-12 Hz) during immobility. As

soon as the mouse started running, theta rhythmic firing synchronized across neurons around a

narrower theta frequency range of 6-10 Hz. Interestingly, running speed-dependent theta

dynamics differed between light and dark conditions (**Figure 2D-G**). In order to quantify the

changes in theta rhythmic firing dynamics as a function of running speed, we identified

individual neurons with statistically significant rhythmic modulation of firing per running speed

condition (alpha = 0.1). In the light condition, we found 181 out of 339 neurons with statistically

significant rhythmic firing during immobility. This number sharply increased to 287 with the onset

of running, but then steadily decreased to 242, 215, and 186 for higher running speeds. In the

dark condition, the number of rhythmically firing neurons showed a similar sharp increase with

the onset of running (from 157 to 312 neurons). In contrast to the light condition, however, the

number of rhythmic neurons during darkness further increased to 322, 326, and 328 out of 339

neurons with higher running speeds (**Figure 2E**). A linear regression model for running speeds

in the range from 5 cm/s to 25 cm/s revealed a significant difference in the number of theta

rhythmic firing neurons as function of running speed between light and dark conditions (p =

0.0003). These data demonstrate that the ensemble of theta rhythmic neurons decreased with

running speed in the light condition, but increased to ceiling during the dark condition (**Figure

2E**) contributing to the stronger synchrony of theta rhythmic firing observed on the population

level (**Figure 2D**). These results were consistent across significance levels for classification of

theta rhythmic neurons between alpha = 0.001 and alpha = 0.1 (**Figure 2—figure supplement

1**). We next identified the cells which remained theta rhythmic across running speed bins up to

25 cm/s and quantified changes in the frequency and magnitude of theta rhythmic firing as a

function of running speed during light and dark conditions. The frequency of theta rhythmic firing

increased with running speed during the light condition (p = 4.2 x 10^{-8}, linear mixed effects

model), but not during the dark condition (p = 0.8085, linear mixed effects model; significant

difference in slopes between light and dark, p = 6.8 x 10^{-6}) (**Figure 2F**). Similarly, the magnitude

of theta rhythmic firing increased with running speed during the light condition (p = 0.0002,

linear mixed effects model), but not during the dark condition (p = 0.0845, linear mixed effects

model; difference in slopes between light and dark, p = 9.1 x 10^{-6}) (**Figure 2G**). Similar results

were obtained when cells had to pass a stricter threshold for being classified as theta rhythmic

(**Figure 2—figure supplement 1**). Notably, the amplitudes of the average autocorrelogram

models across all cells were very similar between light and dark conditions (**Figure 2D,** third

row) despite the observed differences in the magnitudes of theta rhythmic firing on the single
cell level (**Figure 2G**). These data suggest that—during darkness— higher synchrony across
MEC neurons on the population level counterbalances the lower magnitude of theta rhythmicity
on the single cell level which would explain that we did not observe a change in LFP theta
amplitude in the dark condition (**Figure 1C**).
211
Population characteristics of running speed-dependent theta rhythmic firing and their changes
with removal of visual inputs may differ between pyramidal neurons and interneurons. We
therefore classified all theta rhythmic neurons in our data set (n = 342) into putative pyramidal
neurons (PYRs, n = 273) and putative interneurons (INTs, n = 69) based on the spike width and
firing rate of these neurons (**Figure 2—figure supplement 2E**). In general, both the frequency
and magnitude of theta rhythmic firing was higher in PYRs compared to INTs (**Figure 2—figure**
supplements 2C and 3C). In the light condition, the frequency of theta rhythmic firing showed a
similar increase as a function of running speed for both PYRs and INTs but was consistently
higher in PYRs compared to INTs. In contrast, the magnitude of theta rhythmic firing increased
as a function of running speed in INTs but not in PYRs, which showed high theta rhythmic
magnitudes at all running speeds. In the dark condition, frequencies of theta rhythmic firing
differed less between PYRs and INTs compared to the light condition but showed an opposite
change as a function of running speed with values slightly increasing in INTs but slightly
decreasing in PYRs. In contrast, the magnitude of theta rhythmic firing was significantly higher
in PYRs compared to INTs at all running speeds. Statistical results on differences in theta
rhythmic firing between PYRs and INTs are reported in **Tables 1 and 2**. To study the time-
course of changes in theta rhythmic firing, we next analyzed theta rhythmic firing in sequential
10-s blocks after the start of light and dark epochs (**Figs. 2H**). The frequency of theta rhythmic
firing was significantly reduced in the dark condition (effect size: –0.381 Hz, t-statistic: –8.598,
df: 4316, p = 1×10^{-17}, CI: [–0.468 –0.294]). Similar to the fast-initial changes observed in the
slope of the LFP theta frequency vs. running speed relationship, we observed a fast decrease in
the frequency of theta rhythmic firing when transitioning from light to dark and a fast increase
when transitioning from dark to light (**Figure 2I**). In contrast to the observed changes in the
slope of the LFP theta frequency vs. running speed relationship, the frequency of theta rhythmic
firing did not further change on a time scale of tens of seconds during the light or dark epochs.
However, differentiating between PYRs and INTs revealed a decrease in frequency of theta
rhythmic firing as a function of time in both light and dark conditions in INTs but not PYRs
(**Figure 2—figure supplements 2F and 3E**).

Theta-frequency phase-locking of single neuron spiking is a prominent feature of the
hippocampal and entorhinal network (Klausberger et al., 2003; Mizuseki et al., 2009). Increased
phase locking has been associated with dynamical switching between encoding and retrieval
dynamics (Hasselmo et al., 2002; Siegle & Wilson, 2014), memory performance (Liebe et al.,
2012; Rutishauser et al., 2010; Schomburg et al., 2014), and spatial cognitive demand (J. M.
Barry et al., 2016). We therefore asked if the larger synchronization across neurons observed in
the dark condition was associated with an increase of theta-frequency phase locking of theta
rhythmic spiking activity. To compare phase locking of firing between light and dark conditions,
we computed the mean resultant length (MRL) of theta phase firing rate maps for the light and
dark condition (**Figure 3A**). Interestingly, phase locking of theta rhythmic neurons increased
during the dark condition (median MRL (Light) = 0.29, median MRL (Dark) = 0.37, p = 1x10^{-32},
Wilcoxon signed-rank test; **Figure 3B**). Such an increase is consistent with the increased
numbers of theta rhythmic neurons during the dark condition as shown in Figure 2E. No
significant differences were observed between PYRs and INTs (**Figure 3—figure supplement
1**). Notably, each neuron's preferred phase of firing remained unchanged between light and
dark conditions (**Figure 3C & D**). We next investigated the time course of those changes in
phase locking (**Figure 3E**). The time course of changes in theta-frequency phase locking was
similar to the changes in the frequency of theta rhythmic spiking, showing a fast-initial change
but no further slow component of change. No differences were observed between PYRs and
INTs.

Motivated by the differences in LFP activity during light and dark conditions, we asked if these
changes were accompanied by changes in firing rates of MEC neurons. We therefore first asked
if and how the removal and reinstating of visual inputs affected firing rates in MEC neurons.
Consistent with a previous study (Pérez-Escobar et al., 2016), we found striking differences in
firing rates between light and dark epochs among a wide variety of MEC neurons including
speed cells, head direction cells, grid cells, border cells and interneurons **(Figure 4A & B;
Figure 4—figure supplement 1)**. In general, changes in firing rates of MEC neurons could be
positively or negatively correlated to visual inputs. Positive modulation by visual inputs was
observed ~4 times more frequently than negative modulation (139 out of 842 or 16.5%
positively modulated neurons; 32 out of 842 or 3.8% negatively modulated neurons at
significance level alpha = 0.05; data from 15 mice). In both cases, changes in firing rate after
transitions between light and dark conditions occurred almost instantaneously (**Figure 4C & D**).

One possible explanation for the observed changes in firing rates can be a binary change in the

visibility of landmark cues, i.e. the presence or absence of visual information about the animal's

spatial surroundings. Alternatively, changes in firing rates could be driven by changes in net

excitatory drive from areas in the visual cortex due to changes in retinal stimulation independent

of any gain or loss of information about the animal's spatial surroundings. To distinguish

between these two alternative hypotheses, we analyzed data from optogenetic control

experiments in a subset of two mice, in which mice carried a light fiber attached to their

headstage with the fiber tip approximately 1 cm above the animal's head. Green laser light (561

nm, 1 mW) was emitted from the tip of that light fiber and illuminated the head and immediate

surroundings of the animal's head in alternating epochs. Importantly, this experiment was

performed with the ceiling lights turned on during the whole recording session with the green

laser light being turned on and off in alternating epochs during that session. While being visible

to the animal as an egocentrically stationary spot of light, the laser light was too weak to

illuminate the open field environment and thus did not change the visibility of distal visual cues

useful for navigational purposes (e.g. walls or the cue card of the open field arena).

Nevertheless, we found strong effects of that kind of visual point stimulation on firing rates of

five MEC neurons from two mice during foraging under the light condition (**Figure 4E & F**). Four

out of those five neurons showed positive modulation by the egocentrically stationary spot of

laser light, and one showed negative modulation. The effects of presenting an egocentrically

stationary visual stimulus during the light condition on baseline firing rates of MEC neurons

resembled the effects of transitioning from the dark to the light condition. Overall, increased

baseline firing rates due to stronger retinal stimulation were ~4 times more likely to be observed

than decreases in firing rates. In summary, these data demonstrate that the changes in retinal

stimulation due to changes in the amount of spatially non-informative light hitting the retina after

transitioning from dark to light and vice versa are sufficient to explain the observed changes in

firing rates of MEC neurons.

We next turned to the analysis of the hypothesized speed signal by firing rate, in particular

"speed cells" in the MEC (Kropff et al., 2015) and asked if the speed modulation of firing rates in

MEC neurons is changed in complete darkness. Speed cell firing has previously been reported

to be affected by removal of visual inputs and this change in speed cell firing has been

interpreted as a change in the hypothesized speed signal by speed cells, which has further

been indicated to underlie changes in grid cell firing and path integration during darkness (Chen

et al., 2016; Pérez-Escobar et al., 2016). Importantly, previous studies have compared speed

tuning between conditions after normalization *across* conditions. An alternative approach,

however, compares speed tuning between conditions after normalization *within* conditions. It is

important to understand the difference between these two alternative approaches and to

understand the implications of these definitions for our interpretation of changes in speed tuning

curves. When speed modulation is analyzed after normalization across conditions, any change

in external or internal parameters affecting a neuron's mean firing rate will inevitably also affect

the speed tuning curve. For instance, a decrease in a neuron's mean firing rate will result in a

decrease of both the y-intercept and the slope of that neuron's speed tuning curve. In other

words, any change in firing rate will be interpreted as a change in speed modulation. In contrast,

analyzing speed modulation after normalization *within* conditions allows analyzing the speed

modulation of firing rate independently from changes in a neuron's mean firing rate. A change in

speed modulation would then present itself as a change in the slope of the speed tuning curve.

In this study, we do not present evidence for the validity of either approach but argue in favor of

the second approach—analyzing changes in speed modulation after normalization within

conditions—on the grounds that it is standard in the field to compare tuning of other parameters

such as head directional tuning by head direction cells or spatial tuning by place cells, grid cells,

or border cells after normalization within conditions. As shown above, the mean firing rates of

many neurons are responsive to changes in visual inputs. In addition, many speed cells show

conjunctive coding properties which may be changed in complete darkness (Hardcastle et al.,

2017; Hinman et al., 2016). To our knowledge, previous analyses of changes in firing rate

modulation by running speed between two experimental conditions, such as light and dark

environments, have not been corrected for possible confounding factors, such as changes in

mean firing rate or conjunctive coding properties. In particular, data presented in this study show

that changes in visual inputs cause changes in the firing rates of neurons and those changes

may be independent from possible additional changes in the firing rate modulation by running

speed. The same logic applies for changes in conjunctive coding properties, which may be

misinterpreted as changes in the running speed modulation of firing rate if not properly

controlled for. To account for possible changes in conjunctive coding properties, such as

conjunctive coding for position, head direction, and LFP theta phase, we applied a linear-

nonlinear (LN) model (Hardcastle et al., 2017) which provides a method for identification of

significantly speed modulated neurons without restricting the analysis to linear modulation of

firing rates and computes speed response curves instead of speed tuning curves. Speed

response curves are speed tuning curves corrected for the influence of conjunctive coding

properties (Hardcastle et al., 2017). The firing rates of n = 164 neurons out of 495 cells from 15

mice showed significant modulation by running speed in either the light or the dark condition.

Consistent with previous reports, speed modulation of firing rates was heterogeneous (**Figure**

5A) (Dannenberg et al., 2019; Hardcastle et al., 2017; Hinman et al., 2016). Furthermore, we

noticed a distribution of peaks in the speed response curves covering the full range of running

speeds (**Figure 5A**). Hierarchical clustering of speed cells resulted in two main clusters of 138

positively and 26 negatively modulated neurons (**Figure 5B**). Consistent with previous results

(Chen et al., 2016; Pérez-Escobar et al., 2016) we observed a significant decrease in the slope

of speed tuning curves of positively speed-modulated neurons during the dark condition if speed

response curves were normalized to the mean firing rate across conditions (**Figure 5—figure**

supplement 1 and Table 3). 153 out of the 164 speed modulated neurons could also be

examined for potential co-modulation of firing rates by visual inputs. 34 neurons out of those

153 (22.2 %) speed cells showed significant modulation of mean firing rate by visual inputs. This

proportion is very similar to the proportion of 171 out of 842 (20.3%) neurons with significant

modulation of mean firing rates by visual inputs observed in the whole data set. Notably, this is

a conservative estimate because the firing rate of many more neurons may be modulated by

visual inputs without that modulation being detected as significant with the given statistical

power and an alpha level of 0.05. To address the question if the observed changes in the slopes

of speed response curves are caused by a change in speed modulation of firing rates or if those

changes are instead an artefact of changes in the mean firing rates of neurons, we compared

speed response curves between light and dark conditions after normalization *within* conditions

as opposed to normalization *across* conditions. Note that normalization within conditions

corrects for changes in mean firing rates between conditions, whereas normalization across

conditions does not. Interestingly, we found that speed response curves appeared very similar

between light and dark conditions after correcting for changes in mean firing rate and

conjunctive coding properties (**Figure 5C**). In particular, no significant difference was observed

between speed responses of positively speed-modulated neurons in the dark condition

compared to the light condition (**Table 4**). With respect to negatively speed-modulated neurons,

we even observed a small increase in speed modulation during the dark condition (**Figure 5C**

and Table 4). Likewise, speed scores (Pearson's correlation coefficient between firing rate and

running speed) of all speed modulated neurons were not significantly changed between light

and dark conditions on the population level (p = 0.912, n = 164, Kolmogorov-Smirnov test)

(**Figure 5D**) and were highly correlated between light and dark conditions on the single cell level

(R = 0.86) (**Figure 5E**). Moreover, the numbers of speed modulated cells with conjunctive

coding of position, head direction and theta phase were unchanged in the dark condition

(**Figure 5F**). In summary, these data demonstrate that the speed modulation of firing rates in

the MEC remains intact in the absence of visual inputs, even if mean firing rates change

substantially.

While in earlier studies grid cells have been reported to show stable grid cell firing during

complete darkness in rats and mice (Allen et al., 2014; Hafting et al., 2005), two more recent

studies in mice showed a strong reduction in the hexagonal symmetry of grid cell firing during

complete darkness (Chen et al., 2016; Pérez-Escobar et al., 2016). To test if this discrepancy is

due to differences in the experimental settings, the focus in this study was on selectively

eliminating visual inputs only, while leaving other environmental and/or sensory cues such as

walls or odors and auditory cues potentially present in the testing arena intact. We first tested if

grid cell firing can emerge in complete darkness. We therefore introduced mice to the familiar

open field arena during complete darkness (the experimenter was wearing night-vision goggles)

and recorded grid cells during the initial dark session and a following light and dark session (10-

20 min duration each). We found that spatial periodicity of grid cell firing was low—but not

completely eliminated—in the initial dark session and then strongly increased during the first

light session **(Figure 6A & B)**. Interestingly, after this initial orientation, spatial periodicity of grid

cell firing decreased again after removal of visual inputs, but remained significantly higher than

in the initial dark session (Chi-square = 10.89, p = 0.0043, n = 9 cells from two mice, Friedman

test; post hoc tests revealed p < 0.05 comparing initial dark and light, and initial dark and

second dark sessions). We next aimed to study the changes in grid cell spatial firing patterns

after transitioning from light to dark in more detail. Towards that aim, we recorded grid cells

during alternating light/dark epochs, always beginning with a light epoch followed by a dark

epoch. 28 cells from six mice were identified as grid cells based on their grid scores (see

Methods). To compare spatial firing properties of grid cells between light and dark conditions,

we concatenated all light and dark sessions to compute firing rate maps for light and dark

conditions. We found that grid cell field centers remained stable, but grid fields appeared wider

during complete darkness (**Figure 6C**). Quantification of spatial periodicity and spatial precision

of grid cell firing by computing grid scores and spatial information (see Methods) revealed a

significant decrease in both spatial periodicity and spatial precision of spike locations (**Figure**

6D & E). However, firing rate maps are the results of integrating firing rates of grid cells over

long time periods, thereby measuring only spatial variability, but not spatio*temporal* variability of

grid cell firing. This distinction becomes important when considering two alternative hypotheses

with regard to the underlying cause for the observed broadening of grid fields in complete

darkness. Grid fields could appear wider because grid cells fire over a longer distance

throughout a single grid field traversal. Alternatively, grid fields could appear wider because grid

cell firing becomes less stable at the level of single grid field traversals. To distinguish between

these two alternative hypotheses, we used two complementary approaches, First, we computed

the spatiotemporal correlation between the observed firing rate over time and the firing rate

predicted by the animal's location at any given point in time and the associated value in the

firing rate map at that location. The results show a significant reduction in spatiotemporal

correlation during complete darkness (**Figure 6F**). Second, we computed spike-triggered firing

rate maps of grid cells for the light and dark condition. Spike-triggered firing rate maps are not

affected by slow translational shifts of the grid pattern. Nevertheless, we observed a significant

decrease in the grid scores computed from spike-triggered firing rate maps ($p = 0.0354$, $n = 15$

cells, Wilcoxon signed-rank test) (**Figure 6—figure supplements 1 and 2A**) suggesting that

the phase of each grid map remained stable in darkness. Furthermore, we did not observe a

ring-like pattern in the spike-triggered firing rate maps suggesting that the orientation of each

grid map remained stable in darkness too. Lastly, we measured the grid spacing in each of the

spike-triggered firing rate maps. Grid spacing remained unchanged in darkness (median grid

spacing = 42 cm in light and dark, $p = 1$, Wilcoxon signed-rank test, Bayes factor = 0.2005)

(**Figure 6—figure supplement 2B**). Taken together, these data support the hypothesis that grid

fields in the firing rate map appear broader because the spatiotemporal accuracy of grid cell

firing is reduced without changes in the phase, orientation, or spacing of grid maps.

Grid cell firing may rely on an intact head direction signal, in particular on the firing of head

direction cells (Winter, Clark, et al., 2015). We therefore investigated whether the reduction in

spatial accuracy of grid cell firing was accompanied by changes in the head direction tuning of

head direction cells in the MEC. We identified 67 cells with significant unidirectional head

direction tuning, which were classified as head direction cells (Sargolini et al., 2006; Taube et

al., 1990a, 1990b). Interestingly, head direction cells maintained their head direction preference

in the dark condition (**Figure 7A & B and Figure 7—figure supplement 1**) suggesting that

animals remained oriented within the environment during complete darkness. However, the

mean resultant lengths of head directional firing rate maps were decreased in the dark condition

(Cohen's d = 0.238, $p = 0.0337$, Wilcoxon signed-rank test) suggesting that the accuracy of

head directional tuning was reduced in darkness (**Figure 7C**). In addition to grid cells, border or

boundary vector cells (Lever et al., 2010; Solstad et al., 2008) are hypothesized to contribute to

a representation of space in the MEC. We identified 27 cells border cells (**Figure 7D and**

Figure 7—figure supplement 2) and compared border scores (Solstad et al., 2008) of these

cells between light and dark conditions. Border scores were significantly decreased during

darkness (Cohen's d = –0.4975, p = 0.0325, Wilcoxon signed-rank test) (**Figure 7E**) suggesting

that representations of boundaries in the open field environment were less accurate during

darkness.

Our data set included data on n = 7 grid cells from three mice recorded over an exceptionally

long time period across multiple days. These data provided sufficient statistical power and

sufficient sampling of the environment to analyze the time courses of the observed changes in

spatial periodicity, spatial selectivity and spatiotemporal stability of grid cell firing during light and

dark conditions. Towards this goal, we first computed firing rate maps for each of the 10-s bins

of activity to analyze the time course of spatial periodicity quantified by the grid score (**Figure**

8A & B). Second, we computed the spatial correlations between the firing rate maps of the 10-s

binned data and the overall firing rate map for the light condition (**Figure 8C**). Third, we

computed the spatiotemporal correlation for each of the 10-s bins (**Figure 8D**). We found a

saturating exponential increase in spatial periodicity of grid cell firing (**Figure 8B**), spatial

correlation (**Figure 8C**), and spatiotemporal correlation (**Figure 8D & E**) during the light

condition. During darkness, we observed a similar though weaker trend in the opposite

direction. Intriguingly, grid cell stability measured as spatiotemporal correlation of grid cell firing

was strongly correlated with the slope of the LFP theta frequency vs. running speed relationship

(**Figure 8F and Figure 8—figure supplement 1**). Taken together, these data demonstrate that

the representation of space by grid cell firing is correlated to the speed signal by LFP theta

frequency and that both of these signals undergo plastic changes at a behavioral time scale of

tens of seconds.

DISCUSSION

Data presented in this study show how changes in visual inputs affect firing rates and theta

rhythmic firing dynamics in the MEC. Furthermore, these changes correlate with changes in the

representation of location by grid cells, representation of environmental boundaries by border

cells, and representation of head direction by head direction cells in the MEC. Intriguingly, a

proposed speed signal by firing rate—the modulation of firing rate by running speed—remains

unaltered if the analysis is adjusted for changes in mean firing rate. However, an alternative

speed signal by LFP theta frequency—the linear correlation between theta frequency and

running speed—is strongly dependent on visual inputs and lost in darkness. Intriguingly,

changes in the slope of the LFP theta frequency vs. running speed relationship are strongly

correlated to the changes in spatial stability of grid cell firing. While changes in firing rates of

MEC neurons and changes in theta rhythmic firing properties of MEC neurons occur almost

instantaneously during the transitions between light and dark, changes in the proposed speed

signal by LFP theta frequency and changes in grid cell stability show a bimodal change with a

fast component and a further slow saturating exponential component at a 10-s time scale. Such

changes in response to changes in sensory input are longer than previously addressed by

theoretical and computational models of grid cell firing. Lastly, we report changes in the running

speed-dependent population dynamics of theta rhythmic MEC neurons upon changes in visual

inputs. In both the light and dark condition, theta rhythmic neurons become more synchronized

with the onset of running. This increase in synchronization across neurons is accompanied by

an increase in theta frequency phase locking of spiking activity during darkness (Figure 3).

However, while synchronization across neurons is stronger during darkness, the frequency and

magnitude of theta rhythmic firing of individual neurons increase significantly more as a function

of running speed during the light compared to the dark condition.

It has previously been proposed that disruption of speed signals may cause disruption of

spatially periodic grid cell firing in the MEC. Current computational models of grid cell firing and

path integration require a linear speed signal (Burak & Fiete, 2009; Burgess, 2008; Burgess et

al., 2007; Fuhs & Touretzky, 2006; McNaughton et al., 2006). Two candidates for such a speed

signal terminating in the MEC have been proposed: a speed signal by firing rate, most clearly

present in a subpopulation of "speed cells" showing positive and linear speed tuning curves

(Kropff et al., 2015); and an oscillatory speed signal by changes in LFP theta frequency or by

changes in the theta rhythmic firing frequency of individual MEC neurons (Burgess, 2008;

Hinman et al., 2016). Both speed signals can change in response to changes of the size and

shape of the environment (Munn et al., 2020), as do grid cells (C. Barry et al., 2007; Krupic et

al., 2015). In this study, we examined how changes in spatial stability of grid cells relate to

changes in the potential speed signals by firing rate and LFP theta frequency as a function of

time.

In order to analyze the time-course of grid cell stability over time after transitions between light

and dark, we took advantage of long-term recordings of the same grid cells over multiple days

providing us with sufficient statistical power. Intriguingly, changes in spatial stability of grid cell

firing showed an instantaneous component and a slow saturating exponential component at a

behavioral time scale of tens of seconds after transitions between light and dark conditions.

Notably, the observed saturating exponential component in changes of grid cell spatially

periodic firing supports an earlier report by Pérez-Escobar et al. (2016). We next tested if the

observed change in spatial stability of grid cell firing could be explained by a change in the

potential speed signal by firing rate. In order to distinguish the effect of visual inputs on the

mean firing rate of neurons from the effect of visual inputs on the speed modulation of firing

rates, we adjusted for changes in mean firing rates between light and dark conditions before

analyzing the speed modulation of firing rates within light and dark conditions. We further

corrected for multiplexed or conjunctive coding properties by computing speed response curves

using a linear-nonlinear model (Hardcastle et al., 2017). After adjusting for changes in mean

firing rates and conjunctive coding properties, we found that visual inputs had no impact on the

speed modulation (slope) of firing rates in MEC neurons. We therefore conclude that changes in

the potential speed signal by firing rate cannot explain the change in spatially periodic grid cell

firing.

We next turned to the analysis of the time courses of changes in the two oscillatory speed

signals, namely the speed signal by LFP theta frequency and the speed signal by theta rhythmic

firing frequency. Notably, the neuronal distribution of theta rhythmic firing frequencies covers a

broad range of theta (4–12 Hz) indicating that rhythmic firing of MEC neurons is only loosely

coupled to the LFP theta frequency. Moreover—in contrast to LFP theta rhythmicity—theta

rhythmic firing persisted during immobility, though significantly less synchronous across

neurons. The observed changes in theta rhythmic firing frequency at the single cell and

population levels provide a possible explanation for why the LFP theta amplitude vs. running

speed relationship is not changed during complete darkness. During darkness, movement-

related theta rhythmicity magnitude is decreased on the single cell level, while theta

synchronicity across neurons is increased on the population level. These two effects would

influence the LFP theta amplitude in opposite directions and thus may cancel each other out.

Intriguingly, the time course of changes in the slope of the LFP theta frequency vs. running

speed signal matched the time course of changes in grid cell spatial stability (Figure 8F). In

contrast, changes in theta rhythmic firing frequency showed only the fast component of change

(Figs. 2H & I). These data on the parallel time course of changes in the oscillatory speed signal

by LFP theta frequency and grid cell stability suggest that grid cell firing may be more strongly

associated with changes in the mean attractor state of rhythmic firing in the MEC reflected by

changes in the LFP theta signal than with changes in the rhythmic firing of individual neurons.

Alternately, the exponential change in LFP theta may reflect a slow change in an oscillatory

speed signal arising from other cortical structures such as the medial septum.

Other data support the role of LFP oscillatory dynamics in the generation of the grid cell spatial

code. In contrast to the speed signal by firing rate (Dannenberg et al., 2019; Hinman et al.,

2016), the theta oscillatory speed signal depends on the MSDB (Chrobak et al., 1989;

Dannenberg et al., 2015; Givens & Olton, 1994; Hinman et al., 2016; Lawson & Bland, 1993),

and pharmacological inactivation of MSDB neurons strongly reduces both theta oscillations and

spatial periodicity of grid cell firing in the MEC (Brandon et al., 2011; Koenig et al., 2011).

Pharmacological inactivation of the medial septum also impairs self-motion-based linear

distance estimation (Jacob et al., 2017). Furthermore, passive transportation of rats reduces

theta oscillatory power and abolishes both spatially periodic grid cell firing and the speed signal

by LFP theta frequency (Winter, Mehlman, et al., 2015). Likewise, pharmacological inactivation

of the vestibular system or preventing linear/translational head movements in a virtual reality

setup abolish the modulation of LFP theta frequency by running speed and are associated with

a decrease in the spatial information score and an increase in grid scale (Chen et al., 2018;

Jacob et al., 2014). Interestingly, the increase in grid scale observed during path integration on

a circular track is associated with a change in LFP theta oscillations, but not with a change in

the firing patterns of speed and head direction cells (Jacob et al., 2019). The slope of the LFP

theta frequency vs. running speed relationship in particular has been linked to spatial cognition

and grid cell firing (Korotkova et al., 2018). In addition, speed modulation of hippocampal theta

frequency has been shown to correlate with spatial memory performance (Richard et al., 2013).

Moreover, a recent study demonstrated that artificial low-theta stimulation can affect an animal's

estimation of linear distance during approaches to a goal zone (Mouchati et al., 2020) further

strengthening a potential link between LFP theta frequency and path integration. In summary,

one possible explanation for the correlation between changes in spatial accuracy of grid cell

firing and the slope of the LFP theta frequency is that the latter serves as a speed signal in the

path integration process generating the spatially periodic grid cell firing pattern as suggested by

oscillatory interference models of grid cell firing (Burgess, 2008; Burgess et al., 2007;

Hasselmo, 2008).

However, our data also show that the reduction in spatial accuracy of grid cell firing is

accompanied by a reduction of accuracy in head directional tuning and a reduction of spatial

accuracy of border cell firing (Figure 7). An alternative explanation for the observed reduction of

spatial accuracy in grid cell firing during darkness is therefore a reduction of accuracy in head

directional signals (Winter, Clark, et al., 2015), boundary signals, or both due to the loss of

visual information. Another possibility is that the frequency of theta rhythmic firing of MEC

neurons serves as a speed signal that is integrated over longer time scales than previously

accounted for by computational models of grid cells (Dannenberg et al., 2019). In that scenario,

both the slope of the LFP theta frequency and spatial periodicity of grid cell firing would depend

on the history of a velocity signal, which would be consistent with the observed slow saturating

exponential component of change. Such an integration over seconds has previously been

shown to optimize the proposed speed code by firing rate (Dannenberg et al., 2019). Notably,

the slow component of change in spatial accuracy of grid cell firing cannot be explained by error

accumulation alone. While error accumulation could account for the observed slow *decrease* in

spatial stability after removal of visual inputs, error accumulation cannot be the reason for the

observed slow *increase* in spatial accuracy after reinstating visual inputs. If signals required for

grid cell firing are integrated over time, however, such a slow increase is expected. Notably, a

similar increase in spatial accuracy of grid cell firing was also reported recently by Weber &

Sprekeler (2019). Future experimental and computational modeling studies are needed to

address the questions, if and how longer integration time windows could support grid cell firing.

The observed changes in neural correlates of running speed and location during complete

darkness may be attributed to either a loss of visual cues for navigational purposes or

alternatively to a net decrease in excitatory drive from areas in the visual cortex due to the lack

of retinal stimulation (Campbell & Giocomo, 2018). In order to distinguish between these two

possibilities, we took advantage of control data from an optogenetic study (Dannenberg et al.,

2019), where a visible bright spot of light was moving geostationary to the animal without further

illuminating the environment. During this manipulation, the amount of light hitting the retina

increased without changing the visibility of static visual landmark cues. This perturbation

resulted in very similar changes of firing rates in MEC neurons as observed during the light vs.

dark paradigm (Figure 4). Importantly, changes of firing rates, theta rhythmic firing frequency,

and theta phase coupling of rhythmic firing occurred almost instantaneously upon changes in

visual inputs (Figs. 4C & D and Figure 2I). Taken together, these data suggest that changes in

firing rates and theta rhythmic firing are primarily driven by changes in net excitatory input from

visual cortex and do not reflect changes in the perception of the spatial aspects of the

environment. Instead, our data suggest that changes in firing rates and theta rhythmic firing are

driven by changes in retinal stimulation. In contrast, the modulation of firing rates by running

speed is unaltered by changes in illumination or presence or absence of visual cues.

Two recent studies on freely moving mice reported a major disruption of spatially periodic grid

cell firing in the absence of visual cues (Chen et al., 2016; Pérez-Escobar et al., 2016).

However, grid cell firing is intrinsically linked to path integration-based navigation (Gil et al.,

2018; Jacob et al., 2019) and results from studies using virtual reality demonstrate that physical

motion has a greater influence on grid cell firing in mice than visual inputs (Campbell et al.,

2018; Chen et al., 2019; Tennant et al., 2018). While the recording environment in the Chen et

al. study did not differ significantly from the recording environment used in this study, the

majority of light and dark trials performed in the Chen et al. study was discontinuous and, in

most trials, mice were introduced to the recording environment in complete darkness. Likewise,

we also found a significantly stronger disruption in spatially periodic grid cell firing when mice

were introduced to the recording environment in complete darkness (Figs. 6A and B).

Conversely, Chen et al. found a less severe disruption in grid cell firing during darkness in the

few trials where mice were introduced to a familiar environment with the room lights turned on.

Interestingly, Chen et al. reported changes in the preferred head direction angles of head

direction cells suggesting that mice were frequently disoriented. The study by Pérez-Escobar et

al. used a circular maze environment surrounded by water with no walls to eliminate tactile and

olfactory cues. In addition, a white noise generator was used to mask auditory cues. Finally,

mice underwent an explicit disorientation procedure before the start of recordings. These

recording settings resulted in an almost complete loss of head directional tuning and very likely

left animals disoriented. In contrast to those two previous studies, this study investigates the

specific contribution of visual inputs on spatially periodic grid cell firing and theta oscillatory

network dynamics in the MEC by selectively manipulating visual inputs on alternating epochs

but leaving other static environmental and sensory cues such as walls, auditory cues, and

potential olfactory cues unaltered. During this more selective manipulation, grid cells in the MEC

persisted during complete darkness as shown by unaltered orientation, phase, and spacing of

grid cell firing fields. However, accuracy of grid cell firing was reduced during darkness. This

reduction of accuracy in grid cell firing was accompanied by a reduction in the accuracy of the

representation of head direction and environmental boundaries in the MEC. While the preferred

angle of head directional firing in head direction cells was not altered during darkness, the

variability in head directional firing increased during darkness (Figure 7A–C). Likewise, border

cells continued to fire in their respective place fields yet with less spatial accuracy (Figure 7D &

E). Taken together, these data suggest that animals need an initial orientation in the
environment supported by static visual cues but can then use sensory cues other than visual
cues, including wall encounters, to roughly maintain that orientation over long time periods. For
maintaining a spatial representation in a completely dark environment, vestibular signals and
proprioceptive feedback as well as external geometrical cues may be sufficient to maintain the
orientation, scale, and phase of grid cell firing fields. During the light condition, visual cues may
be used instead for refining the self-motion-based cognitive representation of self-location.
655
Theta signals within the hippocampal formation have been suggested to not only reflect
alterations in sensorimotor integration and/or the flow of sensory input, but also distinct cognitive
operations (Hinman et al., 2011). Data presented in this study show that spatial accuracy of grid
cell firing is intrinsically linked to the slope of the LFP theta frequency vs. running speed
relationship. Slow changes in both the slope of the LFP theta frequency vs. running speed
relationship and the spatial periodicity of grid cells after transitions between light and dark
conditions suggest that the representation of space in a cognitive map may integrate velocity
signals over past time and may therefore be history-dependent and updated at a slow
behavioral time scale. The result of such an integration would be a representation of space
which is malleable by internal memory processes.
666

METHODS
Subjects. Data were collected from a total of 17 adult 3-6-month-old male mice (*mus musculus*)
of wildtype and transgenic backgrounds (Wildtype, C57Bl/6J; ChAT-IRES-Cre, B6;129S6-
Chat$^{tm2(cre)Lowl}$/J; PV-IRES-Cre, B6;129P2-Pvalb$^{tm1(cre)Arbr}$/J). Experimental PV-IRES-Cre mice
were homozygous and experimental ChAT-IRES-Cre mice were heterozygous. All experimental
procedures were approved by the Institutional Animal Care and Use Committee for the Charles
River Campus at Boston University. Mice underwent surgery for implantation of 2 movable
tetrodes targeting the superficial layers of the MEC in the left hemisphere. After surgery, mice
were separated for individual housing and maintained on a reversed 12-h light/12-h dark cycle.
Mice were habituated to the experimenter and testing room before the start of all experimental
procedures. For the purpose of additional experiments not reported in this study, mice also
underwent rAAV injections of Archaerhodopsin T or GCaMP6f into the medial septum and light
fiber implantation targeting the medial septum (Dannenberg et al., 2019). Light delivery of ~1–
15mW into the medial septum during one or two test recording sessions did not show any
effects and we did not continue light delivery for the rest of the study. We did not notice any

differences between data recorded before those test stimulations and after those test

stimulations and also did not observe any differences between those mice which had and had

not been stimulated. No differences were observed between wildtype or transgenic mice. Prior

to surgery, mice were housed in Plexiglas cages together with their siblings.

Surgery for microdrive implantation in mice. Mice were injected with buprenorphine (0.1-

mg/kg, s.c.) and atropine (0.1-mg/kg, i.p.), and chronically implanted under isoflurane

anesthesia with 8-channel Axona microdrives (Axona, St. Albans, UK) carrying two movable

tetrodes targeting the superficial layers of the dorsal MEC in the left hemisphere. One ground

screw was implanted above the cerebellum, five anchoring screws were positioned across the

skull, and a craniotomy was performed for implantation of the microdrive, 0.5-mm anterior to the

edge of the transverse sinus and 3.4-mm lateral to the midline. Tetrodes were then implanted

with a 6° polar angle and lowered 1-mm into the brain. The microdrive was secured to the skull

with dental cement. Animals were given buprenorphine (0.1-mg/kg, i.p.), enrofloxacin (7.5-

mg/kg, i.p.), and ketoprofen (3-mg/kg, i.p.) during a 5-day postsurgical care period and allowed

one week in total to fully recover after surgery before the start of recordings.

In vivo electrophysiological recordings. Before implantation, impedance of tetrodes (17-µm

platinum wires, California Fine Wires) was adjusted by electroplating to < 150-kOhm. Tetrodes

were lowered daily by 50-µm until the abundance of spikes and the presence of theta rhythmic

cells in test recordings indicated that tetrode tips were located near or within the superficial cell

layers of the MEC. Neural signals were pre-amplified by unity-gain operational amplifiers

located on each animal's head (head stage) and AC-wire-coupled to an Axona recording system

(Axona Limited, St. Albans, UK). Spikes were threshold-detected and recorded at 48-kHz. The

LFP signal was recorded single-ended with the ground screw as reference at 4800-Hz. Position

and head direction of mice were tracked with headstage-mounted invisible infrared LEDs at 50-

Hz. Spikes were clustered into single units using spike clustering software (DacqUSB/Tint,

Axona, St. Albans, UK). Further offline analyses were performed using Matlab (MathWorks,

Natick, MA, USA), the circular statistics toolbox (Berens, 2009) for Matlab, and R (R Foundation

for Statistical Computing, Vienna, Austria).

Behavior during light and dark conditions. All recordings were performed while animals

foraged for randomly scattered small pieces of Froot Loops (Kellogg Company, Battle Creek,

MI, USA) in a 1-m square open-field arena with 60-cm high black walls and a white letter size

cue card at one wall. Recordings were performed during alternating epochs of open field

illumination (light) and darkness (dark). The dark condition was completely dark from the

perspective of the human experimenter after adaptation to the dark condition for at least five

minutes. A ceiling light provided full illumination of the open field during the light condition.

Turning that ceiling light off resulted in complete darkness, which was achieved by the following

measures. The open field environment was located at the end of the recording room surrounded

by three walls with no doors or windows and a thick laser-proof black curtain covering the front

side. All LED lights of all electrical equipment were completely shielded with black duck-tape.

Tracking of position and head direction of the mouse was achieved by using invisible infrared

LEDs. These infrared LEDs were additionally covered with tape to minimize their brightness and

preventing a potential spectral bleed into the visible spectrum. The slits around the door to the

recording room were covered with black shower curtain material to minimize light leaking into

the space within the recording room outside the laser-proof curtain. The computer monitors

outside of the curtain were also covered with black shower curtains. Importantly, the

experimenter frequently tested for any potential light sources within or light leaks into the space

around the open field environment. Recordings only began when it was completely dark from

the perspective of the human eye after having adopted to the dark condition for at least 5-min.

Alternating epochs of recordings in light and darkness could vary in duration between 30

seconds and 30 min; most experiments used 3-min alternating epochs allowing high statistical

power due to multiple repetitions from light to darkness in each session. A typical recording

session lasted 40-60 min.

Experiments with alternating epochs of light and dark began with the light condition.

Experiments with a dark-light-dark paradigm started in complete darkness. For those

experiments, mice were brought into the room with their cage covered by a black curtain, and

mice were plugged in and introduced into the arena during complete darkness by an

experimenter wearing night vision goggles. The experimenter then left the arena side of the

curtain before recordings began.

Histology. After the end of electrophysiological data collection, animals were deeply

anaesthetized by isoflurane or i.p. injection of Euthasol (390-mg/kg), followed by electro-lesions

(20-µA, 16s) through one channel of each tetrode and subsequent transcardial perfusion with

saline followed by 10% buffered formalin (SF100-4, ThermoFisher Scientifc). Brains were

extracted and stored in fixative for one day. Cresyl violet staining was performed on 50-µm

sagittal slices of MEC to visualize tetrode tracks and confirm positions of recording sites in the

MEC.

Data analysis.

753   ***Computation of running speed and firing rate signals.*** The animal's running speed

754   was estimated by applying a Kalman filter to the positional data obtained from the video tracking

755   sampled at 50-Hz (Fyhn et al., 2004; Hinman et al., 2016; Kropff et al., 2015). To obtain an

756   instantaneous firing rate signal matching the sampling rate of the behavioral tracking, we

757   calculated the firing rate as the number of spikes per video frame period (20-ms) and smoothed

758   this signal with a 125-ms Gaussian filter.

759   ***Modulation of mean firing rates by visual inputs.*** In order to identify neurons, whose

firing rates are modulated by the illumination of the environment (consistently either higher or

lower firing rates during light or darkness), we computed the average firing rate for each light

and dark epoch and tested if the differences between light epochs and the following dark

epochs and the differences between dark epochs and the following light epochs reached

statistical significance at alpha = 0.05 using Student's paired t-test. To analyze the time-course

of changes in mean firing rates after transition from light to dark or from dark to light, we first

computed the peristimulus time histogram with 5-s binning around the transition point and then

computed the average of z-scored firing rates across neurons. Only cells, which showed spiking

activity across three light-to-dark and three dark-to-light epoch pairs were included into this

analysis.

770   ***LFP theta rhythm.*** LFP data were down-sampled to 600-Hz and power spectral density

analyses were performed using custom written MATLAB scripts and the FieldTrip toolbox

(Oostenveld et al., 2011). A theta frequency range was determined for each session based on

the range of the theta peak in the power spectral density plot (Dannenberg et al., 2019). The

LFP signal was then band-pass-filtered at this theta frequency range with a third order

Butterworth filter, and the instantaneous theta amplitude and phase were determined via a

Hilbert transform. The instantaneous theta frequency could then be inferred from the

instantaneous phase. To analyze changes in the LFP theta frequency vs. running speed

relationship over time, we first down-sampled the LFP theta frequency signal to 50-Hz in order

to match the temporal binning of the behavioral tracking. We next sectioned data into 10-s

blocks after transitions between light and dark and pooled data from all sessions and all mice

(Figs. 1 H-L). For each of those 10-s blocks of data, the slope and y-intercept of the correlation

between LFP theta frequency and running speed were computed using a linear model. The

same analysis as described above was applied to 10-s blocks around the transition points from
light to dark or from dark to light (Figs. 1K & L).

Theta rhythmic firing. We used maximum likelihood estimation (MLE) of a parametric
model of the interspike intervals underlying the spike time autocorrelogram to analyze theta
rhythmic firing, which has been proven a more sensitive and statistically rigorous method than
traditional measures of spike time autocorrelogram features (Climer et al., 2015). In particular,
this model allowed us to statistically test significance of theta modulation in MEC neurons and
compare theta rhythmic firing frequency and theta rhythmicity magnitude on a single cell level
between light and dark conditions. The model tested for rhythmic firing in the frequency range
between 1-13 Hz. For the analysis of the relationship between theta rhythmic firing and running
speed (Figure 2), we first identified continuous temporal epochs larger than 400-ms, in which
the running speed of the animal varied less than ± 4-cm/s. We next sorted these epochs
according to the average running speed during the epochs into 1-cm/s bins from 1-cm/s to 29-
cm/s and used the MLE approach to fit an autocorrelogram model to each of those epochs. The
minimum number of interspike intervals which need to be present in the autocorrelogram to
compute an accurate model is 100 (Climer et al., 2015), and no autocorrelogram model was
computed if the number of interspike intervals was smaller than 100 for the epoch of interest.
For each neuron and running speed bin, an average autocorrelogram was computed across the
autocorrelograms for each epoch. The maximal run time for the theta rhythmic firing vs. running
speed analysis on the research computing cluster was set to two weeks resulting in the
successful analysis of 340 out of 342 neurons. For the analysis of the time course of changes in
theta rhythmic firing after transition from light to dark or from dark to light, the first two minutes
after the transition were sectioned into 10-s blocks (n = 342 neurons, Figure 2H). Alternately,
10-s blocks around the transition points from light to dark or from dark to light were analyzed (n
= 342 neurons, Figure 2I). For each neuron, an autocorrelogram model was fitted to each of
those 10-s blocks of data using the MLE approach. For each of those 10-s blocks of data, the
average theta rhythmic firing frequency across neurons could then be computed from the peaks
of the average autocorrelogram model across neurons by taking the inverse of the first peak in
the autocorrelogram model. To analyze running speed-dependent theta rhythmic firing on the
population level, we first identified continuous epochs of running speed within the ranges of 0-3
cm/s (immobility), 5-10 cm/s, 10-15 cm/s, 15-20 cm/s, and 20-25 cm/s. We next used the MLE
approach to model an autocorrelogram for each running speed bin of each neuron. Neurons
with sufficient number of spikes resulting in at least 100 interspike intervals to compute an MLE-

modeled autocorrelogram in at least one of the running speed bins were included into the

analysis (n = 339 neurons).

Analysis of theta-frequency phase locking. The analysis of theta-frequency phase

locking is only valid when LFP theta rhythm is present and a sufficiently large number of spikes

is present to estimate the phase distribution. We therefore restricted our analysis to time points

where the running speed of the mouse was larger than 5 cm/s and where the sample size was

larger than 30 spikes. LFP theta phase and the mean resultant length were computed as

described in detail in Dannenberg et al. (2015). In short, the instantaneous LFP theta phase was

computed from the Hilbert transform of the LFP signal filtered within the theta-range (6–10 Hz).

For higher accuracy, the boundaries of this theta range were adjusted separately for each

animal as described in detail in Dannenberg et al. (2019). We next interpolated the LFP theta

phase signal and assigned each spike of a theta rhythmic neuron to the instantaneous LFP

theta phase at the time of spiking.

Classification of putative pyramidal neurons and putative interneurons. For

classification of single units into putative pyramidal or interneurons, we computed the spike

width and firing rate for each unit. The spike width was defined as the time between the peak

and the trough of the spike waveform. The distribution of spike widths was bimodal with a

minimum between 0.29 ms and 0.34 ms. Units with a spike width < 0.29 ms or a firing rate > 10

Hz were classified as putative interneurons, the remaining units were classified as putative

pyramidal neurons.

Computation of spatial firing rate maps. Occupancy-normalized spatial firing rate

maps were generated by dividing the open field environment into 3-cm spatial bins. For each

spatial bin, an occupancy-normalized firing rate was computed by dividing the total number of

spikes falling into that bin by the total time the animal spent in that bin. Spatial rate maps were

then smoothed by a 3-cm wide two-dimensional Gaussian kernel.

Grid score and grid cells. The grid score was used as a measure of hexagonal

symmetry of grid fields. The grid score was calculated as follows. We first computed the two-

dimensional autocorrelogram of the spatial rate map, extracted a ring that encased the six

peaks closest to the center peak excluding the central peak and computed correlations between

this modified spatial autocorrelogram and rotated versions of itself after correction of elliptical

eccentricity as described in detail in the supplementary methods for computation of gridness

measure 3 in Brandon et al. (2011). If hexagonal symmetry is present, those correlations will

peak at multiples of 60 degrees and repeat themselves infinitively at a frequency of 6 peaks per

rotational cycle. We therefore can compute the power spectrum of those correlations as a

function of rotational angle. The power at 60 degree was then defined as the grid score.

Compared to previously used grid scores, this grid score largely reduced the number of false-

positive grid cells, while reliably assigning high grid scores to clear grid cell examples. This grid

score is defined as a measure between 0 and 1, with 0 meaning no hexagonal symmetry of

firing fields in the spatial rate map or simply "gridness" and 1 meaning perfect gridness. Grid

cells were defined as cells with a grid score > 0.85 in either light or dark condition.

Spatial information. We defined the spatial information as the delta entropy $\Delta H =$

$H_{max} - H$. H_{max} is the maximal amount of randomness given by a completely random

distribution of spike locations in the open field environment defined by $H_{max} = -\sum_{i=1}^{n} p * \log_2 p$,

with $p = \frac{1}{n}$, where n is the number of spatial bins. H is the observed amount of randomness in

the distribution of spike locations given by $H(X|spike) = -\sum_{i=1}^{n} p(x_i|spike) * \log_2 p(x_i|spike)$,

where X is the animal's location, x_i is the ith spatial bin, n is the number of spatial bins and

$p(x_i|spike)$ is the probability that $X = x_i$ when a spike is observed. ΔH can then be understood

as an increase in the amount of order in the distribution of spike locations relative to a

completely random (uniform) distribution of spike locations.

Spatial correlation. Firing rate map similarities were quantified by the spatial correlation

between two firing rate maps. To compute the spatial correlation, the spatial bins of the firing

rate maps were transformed into a single vector. The spatial correlation was then computed as

the Pearson's correlation coefficient between two of those spatially binned firing rate vectors.

Spatiotemporal correlation. To analyze spatial stability of grid cell firing in the time

domain, we used the firing rate map of the light condition to compute the expected firing rate for

each of the 20-msec time bins of the animal's trajectory. We next computed the Pearson's

correlation coefficient between the observed 20-msec binned firing rate and the expected 20-

msec binned firing rate.

Spike-triggered firing rate maps. Spike-triggered firing rate maps (Chen et al., 2016;

Pérez-Escobar et al., 2016) were constructed as follows. For each spike, occupancy-normalized

firing rates were computed from the next 10 s of data and centered to the spatial bin at the time

of the spike. All those firing rate maps were then averaged and smoothed with a Gaussian

kernel (SD = 3 cm) to create the spike-triggered firing rate map. To compute the occupancy-

normalized firing rate as a function of radial distance, the circumferences of circles with

increasing radii ranging from zero to 95 cm were overlaid with the spike-triggered firing rate

map. For each of those circles, the spatial firing rates of all bins falling on the circle's

circumference were then averaged to compute the spatial firing as a function of radial distance.

The location of the first peak in the plot of the firing rate as function of radial distance defined

the spacing of the grid cell.

885   ***Border score and border cells.*** A border score was computed as described in

detail by Solstad et al. (2008). Border scores ranged from –1 for cells with central firing fields to

+1 for cells with firing fields that perfectly aligned with at least one entire wall of the open field

environment. Cells with a border score larger than 0.68 were defined as border cells.

Computation of head direction firing rate maps. Head directional firing rate maps

were generated by dividing the head directional parameter space into six-degree bins and

computing the occupancy-normalized firing rate as the total number of spikes falling into that bin

divided by the total time the animal spent in that bin.

Head directional tuning and head direction cells. Head direction cells were classified

based on the criteria reported in Peyrache et al. (2015) and Viejo & Peyrache (2020). Only cells

with at least 10 spikes in each of the light and dark conditions were included in the analysis.

Cells were classified as head direction cells if each of the following was true. First, the inverse of

the circular standard deviation was greater than 1 in either the light or the dark condition;

second, the p-value of the Rayleigh test was smaller than 0.001 in either the light or the dark

condition; third, the peak firing rate in the head directional field was larger than 1 Hz in either the

light or the dark condition.

Identification and classification of speed-modulated neurons. To identify speed

modulated neurons, we used a linear non-linear Poisson (LN) model (Hardcastle et al., 2017)

which tested for significant modulation of firing rates by position, head direction, LFP theta

phase, and/or running speed (Figure 5). For a robust analysis, we first screened for cells which

showed stable firing across the whole recording session. Cells, whose mean firing rates were

below 1-Hz or whose mean firing rates differed more than by a factor of two between the first

and second halves of the recording session, were excluded from the analysis. To compare

speed modulation of firing rates on the population level, we fitted the data with a quadratic

polynomial curve (poly2 fit, MATLAB) and clustered neurons into positively and negatively

speed-modulated neurons using hierarchical clustering (clustergram, bioinformatics toolbox,

MATLAB).

Speed score. The speed score was calculated as the Pearson's product moment

correlation between instantaneous firing rate and running speed (Kropff et al., 2015).

Experimental design and statistical analysis. Details on statistical analyses on each

experiment are reported in the Results section. Statistical tests used in this study include

Student's paired t-test and nonparametric statistical tests for repeated measures, namely

Wilcoxon's signed rank test and Friedman's test. Post hoc testing was performed if the

Friedman test yielded a significant result at alpha = 0.05. The dataset used for analysis of

differences in neural dynamics between light and dark conditions was obtained from recordings

in a total of 15 mice. This dataset included LFP data from 14 out of those 15 mice and 1267

single units from all 15 mice. From the data set of 1267 single units, 495 single units with stable

firing across the recording session and mean firing rates larger than 1 Hz were included in the

LN-model analysis of firing rate modulation by running speed; 842 single units, which showed

spiking activity over at least three light-to-dark and three dark-to-light epoch pairs, were included

in the analysis of firing rate modulation by visual inputs. Data with green laser light stimulation

(related to Figure 4) were recorded in two additional mice.

Code accessibility. Custom MATLAB code used for data analysis can be found at

https://github.com/hasselmonians or can be made available upon request. Violin plots were

created using R software (R Core Team, 2019; R: A language and environment for statistical

computing; R Foundation for Statistical Computing, Vienna, Austria).

ACKNOWLEDGMENTS

This work was supported by the National Institute of Mental Health, grant numbers R01

MH60013,R01 MH120073 and R01 MH052090, by the National Institute of Neurological

Disorders and Stroke, grant number K99NS116129, and by the Office of Naval Research MURI

N00014-16-1-2832 and MURI N00014-19-1-2571, and Office of Naval Research DURIP

N00014-17-1-2304.

COMPETING INTERESTS

None

REFERENCES

Allen, K., Gil, M., Resnik, E., Toader, O., Seeburg, P., & Monyer, H. (2014). Impaired path integration and

grid cell spatial periodicity in mice lacking GluA1-containing AMPA receptors. *The Journal of*

Neuroscience: The Official Journal of the Society for Neuroscience, *34*(18), 6245–6259.

https://doi.org/10.1523/JNEUROSCI.4330-13.2014

Barry, C., Ginzberg, L. L., O'Keefe, J., & Burgess, N. (2012). Grid cell firing patterns signal environmental

novelty by expansion. *Proceedings of the National Academy of Sciences of the United States of*

America, *109*(43), 17687–17692. https://doi.org/10.1073/pnas.1209918109

Barry, C., Hayman, R., Burgess, N., & Jeffery, K. J. (2007). Experience-dependent rescaling of entorhinal

grids. *Nature Neuroscience*, *10*(6), 682–684. https://doi.org/10.1038/nn1905

Barry, J. M., Sakkaki, S., Barriere, S. J., Patterson, K. P., Lenck-Santini, P. P., Scott, R. C., Baram, T. Z., &

Holmes, G. L. (2016). Temporal Coordination of Hippocampal Neurons Reflects Cognitive

Outcome Post-febrile Status Epilepticus. *EBioMedicine*, *7*, 175–190.

https://doi.org/10.1016/j.ebiom.2016.03.039

Berens, P. (2009). CircStat: A MATLAB Toolbox for Circular Statistics. *Journal of Statistical Software*,

31(1), 1–21. https://doi.org/10.18637/jss.v031.i10

Blair, H. T., Gupta, K., & Zhang, K. (2008). Conversion of a phase- to a rate-coded position signal by a

three-stage model of theta cells, grid cells, and place cells. *Hippocampus*, *18*(12), 1239–1255.

https://doi.org/10.1002/hipo.20509

Brandon, M. P., Bogaard, A. R., Libby, C. P., Connerney, M. A., Gupta, K., & Hasselmo, M. E. (2011).

Reduction of theta rhythm dissociates grid cell spatial periodicity from directional tuning.

Science (New York, N.Y.), *332*(6029), 595–599. https://doi.org/10.1126/science.1201652

Burak, Y., & Fiete, I. R. (2009). Accurate path integration in continuous attractor network models of grid

cells. *PLoS Computational Biology*, *5*(2), e1000291.

https://doi.org/10.1371/journal.pcbi.1000291

Burgess, N. (2008). Grid cells and theta as oscillatory interference: Theory and predictions.

Hippocampus, *18*(12), 1157–1174. https://doi.org/10.1002/hipo.20518

Burgess, N., Barry, C., & O'Keefe, J. (2007). An oscillatory interference model of grid cell firing.

Hippocampus, *17*(9), 801–812. https://doi.org/10.1002/hipo.20327

Bush, D., & Burgess, N. (2014). A Hybrid Oscillatory Interference/Continuous Attractor Network Model

of Grid Cell Firing. *Journal of Neuroscience*, *34*(14), 5065–5079.

https://doi.org/10.1523/JNEUROSCI.4017-13.2014

Campbell, M. G., & Giocomo, L. M. (2018). Self-motion processing in visual and entorhinal cortices:

Inputs, integration, and implications for position coding. *Journal of Neurophysiology*.

https://doi.org/10.1152/jn.00686.2017

Campbell, M. G., Ocko, S. A., Mallory, C. S., Low, I. I. C., Ganguli, S., & Giocomo, L. M. (2018). Principles

governing the integration of landmark and self-motion cues in entorhinal cortical codes for

navigation. *Nature Neuroscience*, *21*(8), 1096–1106. https://doi.org/10.1038/s41593-018-0189-y

Chen, G., King, J. A., Lu, Y., Cacucci, F., & Burgess, N. (2018). Spatial cell firing during virtual navigation of

open arenas by head-restrained mice. *ELife*, *7*. https://doi.org/10.7554/eLife.34789

Chen, G., Lu, Y., King, J. A., Cacucci, F., & Burgess, N. (2019). Differential influences of environment and

self-motion on place and grid cell firing. *Nature Communications*, *10*(1), 630.

https://doi.org/10.1038/s41467-019-08550-1

Chen, G., Manson, D., Cacucci, F., & Wills, T. J. (2016). Absence of Visual Input Results in the Disruption

of Grid Cell Firing in the Mouse. *Current Biology: CB*. https://doi.org/10.1016/j.cub.2016.06.043

Cheung, A. (2016). Probabilistic Learning by Rodent Grid Cells. *PLoS Computational Biology*, *12*(10),

e1005165. https://doi.org/10.1371/journal.pcbi.1005165

Chrobak, J. J., Stackman, R. W., & Walsh, T. J. (1989). Intraseptal administration of muscimol produces

dose-dependent memory impairments in the rat. *Behavioral and Neural Biology*, *52*(3), 357–

369.

Climer, J. R., DiTullio, R., Newman, E. L., Hasselmo, M. E., & Eden, U. T. (2015). Examination of

rhythmicity of extracellularly recorded neurons in the entorhinal cortex. *Hippocampus*, *25*(4),

460–473. https://doi.org/10.1002/hipo.22383

Dannenberg, H., Kelley, C., Hoyland, A., Monaghan, C. K., & Hasselmo, M. E. (2019). The Firing Rate

Speed Code of Entorhinal Speed Cells Differs across Behaviorally Relevant Time Scales and Does

Not Depend on Medial Septum Inputs. *The Journal of Neuroscience*, *39*(18), 3434–3453.

https://doi.org/10.1523/JNEUROSCI.1450-18.2019

Dannenberg, H., Pabst, M., Braganza, O., Schoch, S., Niediek, J., Bayraktar, M., Mormann, F., & Beck, H.

(2015). Synergy of direct and indirect cholinergic septo-hippocampal pathways coordinates

firing in hippocampal networks. *The Journal of Neuroscience*, *35*(22), 8394–8410.

https://doi.org/10.1523/JNEUROSCI.4460-14.2015

Evans, T., Bicanski, A., Bush, D., & Burgess, N. (2016). How environment and self-motion combine in

neural representations of space. *The Journal of Physiology*, *594*(22), 6535–6546.

https://doi.org/10.1113/JP270666

Fuhs, M. C., & Touretzky, D. S. (2006). A spin glass model of path integration in rat medial entorhinal

cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *26*(16),

4266–4276. https://doi.org/10.1523/JNEUROSCI.4353-05.2006

Fyhn, M., Hafting, T., Witter, M. P., Moser, E. I., & Moser, M.-B. (2008). Grid cells in mice. *Hippocampus*,

18(12), 1230–1238. http://dx.doi.org/10.1002/hipo.20472

Fyhn, M., Molden, S., Witter, M. P., Moser, E. I., & Moser, M.-B. (2004). Spatial representation in the

entorhinal cortex. *Science*, *305*(5688), 1258–1264. http://dx.doi.org/10.1126/science.1099901

Gil, M., Ancau, M., Schlesiger, M. I., Neitz, A., Allen, K., De Marco, R. J., & Monyer, H. (2018). Impaired

path integration in mice with disrupted grid cell firing. *Nature Neuroscience*, *21*(1), 81–91.

https://doi.org/10.1038/s41593-017-0039-3

Giocomo, L. M. (2016). Environmental boundaries as a mechanism for correcting and anchoring spatial

maps. *The Journal of Physiology*, *594*(22), 6501–6511. https://doi.org/10.1113/JP270624

Givens, B., & Olton, D. S. (1994). Local modulation of basal forebrain: Effects on working and reference

1019   memory. *The Journal of Neuroscience*, *14*(6), 3578–3587.

1020   Hafting, T., Fyhn, M., Molden, S., Moser, M.-B., & Moser, E. I. (2005). Microstructure of a spatial map in

1021   the entorhinal cortex. *Nature*, *436*(7052), 801–806. http://dx.doi.org/10.1038/nature03721

1022   Hardcastle, K., Ganguli, S., & Giocomo, L. M. (2015). Environmental boundaries as an error correction

1023   mechanism for grid cells. *Neuron*, *86*(3), 827–839.

1024   https://doi.org/10.1016/j.neuron.2015.03.039

1025   Hardcastle, K., Maheswaranathan, N., Ganguli, S., & Giocomo, L. M. (2017). A Multiplexed,

1026   Heterogeneous, and Adaptive Code for Navigation in Medial Entorhinal Cortex. *Neuron*, *94*(2),

1027   375-387.e7. https://doi.org/10.1016/j.neuron.2017.03.025

1028   Hasselmo, M. E. (2008). Grid cell mechanisms and function: Contributions of entorhinal persistent

1029   spiking and phase resetting. *Hippocampus*, *18*(12), 1213–1229.

1030   https://doi.org/10.1002/hipo.20512

1031   Hasselmo, M. E., Bodelón, C., & Wyble, B. P. (2002). A proposed function for hippocampal theta rhythm:

1032   Separate phases of encoding and retrieval enhance reversal of prior learning. *Neural*

1033   *Computation*, *14*(4), 793–817. https://doi.org/10.1162/089976602317318965

1034   Hinman, J. R., Brandon, M. P., Climer, J. R., Chapman, G. W., & Hasselmo, M. E. (2016). Multiple Running

1035   Speed Signals in Medial Entorhinal Cortex. *Neuron*, *91*(3), 666–679.

1036   https://doi.org/10.1016/j.neuron.2016.06.027

1037   Hinman, J. R., Penley, S. C., Long, L. L., Escabí, M. A., & Chrobak, J. J. (2011). Septotemporal variation in

1038   dynamics of theta: Speed and habituation. *Journal of Neurophysiology*, *105*(6), 2675–2686.

1039   https://doi.org/10.1152/jn.00837.2010

Jacob, P.-Y., Capitano, F., Poucet, B., Save, E., & Sargolini, F. (2019). Path integration maintains spatial

periodicity of grid cell firing in a 1D circular track. *Nature Communications*, *10*(1), 1–13.

https://doi.org/10.1038/s41467-019-08795-w

Jacob, P.-Y., Gordillo-Salas, M., Facchini, J., Poucet, B., Save, E., & Sargolini, F. (2017). Medial entorhinal

cortex and medial septum contribute to self-motion-based linear distance estimation. *Brain*

Structure & Function, *222*(6), 2727–2742. https://doi.org/10.1007/s00429-017-1368-4

Jacob, P.-Y., Poucet, B., Liberge, M., Save, E., & Sargolini, F. (2014). Vestibular control of entorhinal

cortex activity in spatial navigation. *Frontiers in Integrative Neuroscience*, *8*, 38.

https://doi.org/10.3389/fnint.2014.00038

Jeewajee, A., Barry, C., O'Keefe, J., & Burgess, N. (2008). Grid cells and theta as oscillatory interference:

Electrophysiological data from freely moving rats. *Hippocampus*, *18*(12), 1175–1185.

https://doi.org/10.1002/hipo.20510

Jeewajee, A., Lever, C., Burton, S., O'Keefe, J., & Burgess, N. (2008). Environmental novelty is signaled by

reduction of the hippocampal theta frequency. *Hippocampus*, *18*(4), 340–348.

https://doi.org/10.1002/hipo.20394

Klausberger, T., Magill, P. J., Márton, L. F., Roberts, J. D. B., Cobden, P. M., Buzsáki, G., & Somogyi, P.

(2003). Brain-state- and cell-type-specific firing of hippocampal interneurons in vivo. *Nature*,

421(6925), 844–848. https://doi.org/10.1038/nature01374

Koenig, J., Linder, A. N., Leutgeb, J. K., & Leutgeb, S. (2011). The spatial periodicity of grid cells is not

sustained during reduced theta oscillations. *Science (New York, N.Y.)*, *332*(6029), 592–595.

https://doi.org/10.1126/science.1201685

Korotkova, T., Ponomarenko, A., Monaghan, C. K., Poulter, S. L., Cacucci, F., Wills, T., Hasselmo, M. E., &

Lever, C. (2018). Reconciling the different faces of hippocampal theta: The role of theta

oscillations in cognitive, emotional and innate behaviors. *Neuroscience and Biobehavioral Reviews*, *85*, 65–80. https://doi.org/10.1016/j.neubiorev.2017.09.004

Kraus, B. J., Brandon, M. P., Robinson, R. J., Connerney, M. A., Hasselmo, M. E., & Eichenbaum, H. (2015). During Running in Place, Grid Cells Integrate Elapsed Time and Distance Run. *Neuron*, *88*(3), 578–589. https://doi.org/10.1016/j.neuron.2015.09.031

Kropff, E., Carmichael, J. E., Moser, M.-B., & Moser, E. I. (2015). Speed cells in the medial entorhinal cortex. *Nature*, *523*(7561), 419–424. https://doi.org/10.1038/nature14622

Krupic, J., Bauza, M., Burton, S., Barry, C., & O'Keefe, J. (2015). Grid cell symmetry is shaped by environmental geometry. *Nature*, *518*(7538), 232–235. https://doi.org/10.1038/nature14153

Lawson, V. H., & Bland, B. H. (1993). The role of the septohippocampal pathway in the regulation of hippocampal field activity and behavior: Analysis by the intraseptal microinfusion of carbachol, atropine, and procaine. *Experimental Neurology*, *120*(1), 132–144. https://doi.org/10.1006/exnr.1993.1047

Lever, C., Burton, S., Jeewajee, A., Wills, T. J., Cacucci, F., Burgess, N., & O'Keefe, J. (2010). Environmental novelty elicits a later theta phase of firing in CA1 but not subiculum. *Hippocampus*, *20*(2), 229–234. https://doi.org/10.1002/hipo.20671

Liebe, S., Hoerzer, G. M., Logothetis, N. K., & Rainer, G. (2012). Theta coupling between V4 and prefrontal cortex predicts visual short-term memory performance. *Nature Neuroscience*, *15*(3), 456–462, S1-2. https://doi.org/10.1038/nn.3038

McFarland, W. L., Teitelbaum, H., & Hedges, E. K. (1975). Relationship between hippocampal theta activity and running speed in the rat. *Journal of Comparative and Physiological Psychology*, *88*(1), 324–328.

McNaughton, B. L., Battaglia, F. P., Jensen, O., Moser, E. I., & Moser, M.-B. (2006). Path integration and

the neural basis of the "cognitive map". *Nat Rev Neurosci*, *7*(8), 663–678.

http://dx.doi.org/10.1038/nrn1932

Mizuseki, K., Sirota, A., Pastalkova, E., & Buzsáki, G. (2009). Theta oscillations provide temporal windows

for local circuit computation in the entorhinal-hippocampal loop. *Neuron*, *64*(2), 267–280.

https://doi.org/10.1016/j.neuron.2009.08.037

Monaghan, C. K., Chapman, G. W., & Hasselmo, M. E. (2017). Systemic administration of two different

anxiolytic drugs decreases local field potential theta frequency in the medial entorhinal cortex

without affecting grid cell firing fields. *Neuroscience*, *364*, 60–70.

https://doi.org/10.1016/j.neuroscience.2017.08.056

Mouchati, P. R., Kloc, M. L., Holmes, G. L., White, S. L., & Barry, J. M. (n.d.). Optogenetic "low-theta"

pacing of the septohippocampal circuit is sufficient for spatial goal finding and is influenced by

behavioral state and cognitive demand. *Hippocampus*, *n/a*(n/a).

https://doi.org/10.1002/hipo.23248

Munn, R. G. K., Mallory, C. S., Hardcastle, K., Chetkovich, D. M., & Giocomo, L. M. (2020). Entorhinal

velocity signals reflect environmental geometry. *Nature Neuroscience*, *23*(2), 239–251.

https://doi.org/10.1038/s41593-019-0562-5

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source software for

advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational*

Intelligence and Neuroscience, *2011*, 156869. https://doi.org/10.1155/2011/156869

Pérez-Escobar, J. A., Kornienko, O., Latuske, P., Kohler, L., & Allen, K. (2016). Visual landmarks sharpen

grid cell metric and confer context specificity to neurons of the medial entorhinal cortex. *ELife*,

5. https://doi.org/10.7554/eLife.16937

Peyrache, A., Lacroix, M. M., Petersen, P. C., & Buzsáki, G. (2015). Internally organized mechanisms of

the head direction sense. *Nature Neuroscience*, *18*(4), 569–575.

https://doi.org/10.1038/nn.3968

Raudies, F., Brandon, M. P., Chapman, G. W., & Hasselmo, M. E. (2015). Head direction is coded more

strongly than movement direction in a population of entorhinal neurons. *Brain Research*, *1621*,

355–367. https://doi.org/10.1016/j.brainres.2014.10.053

Raudies, F., & Hasselmo, M. E. (2015). Differences in Visual-Spatial Input May Underlie Different

Compression Properties of Firing Fields for Grid Cell Modules in Medial Entorhinal Cortex. *PLoS*

Computational Biology, *11*(11), e1004596. https://doi.org/10.1371/journal.pcbi.1004596

Raudies, F., Mingolla, E., & Hasselmo, M. E. (2012). Modeling the influence of optic flow on grid cell

firing in the absence of other cues1. *Journal of Computational Neuroscience*, *33*(3), 475–493.

https://doi.org/10.1007/s10827-012-0396-6

Richard, G. R., Titiz, A., Tyler, A., Holmes, G. L., Scott, R. C., & Lenck-Santini, P.-P. (2013). Speed

modulation of hippocampal theta frequency correlates with spatial memory performance.

Hippocampus, *23*(12), 1269–1279. https://doi.org/10.1002/hipo.22164

Rivas, J., Gaztelu, J. M., & García-Austt, E. (1996). Changes in hippocampal cell discharge patterns and

theta rhythm spectral properties as a function of walking velocity in the guinea pig. *Experimental*

Brain Research, *108*(1), 113–118. https://doi.org/10.1007/BF00242908

Rutishauser, U., Ross, I. B., Mamelak, A. N., & Schuman, E. M. (2010). Human memory strength is

predicted by theta-frequency phase-locking of single neurons. *Nature*, *464*(7290), 903–907.

http://dx.doi.org/10.1038/nature08860

Sargolini, F., Fyhn, M., Hafting, T., McNaughton, B. L., Witter, M. P., Moser, M.-B., & Moser, E. I. (2006).

Conjunctive representation of position, direction, and velocity in entorhinal cortex. *Science (New*

York, N.Y.), *312*(5774), 758–762. https://doi.org/10.1126/science.1125572

Schomburg, E. W., Fernández-Ruiz, A., Mizuseki, K., Berényi, A., Anastassiou, C. A., Koch, C., & Buzsáki, G.

(2014). Theta phase segregation of input-specific gamma patterns in entorhinal-hippocampal

networks. *Neuron*, *84*(2), 470–485. https://doi.org/10.1016/j.neuron.2014.08.051

Shin, J., & Talnov, A. (2001). A single trial analysis of hippocampal theta frequency during nonsteady

wheel running in rats. *Brain Research*, *897*(1–2), 217–221. https://doi.org/10.1016/s0006-

8993(01)02015-7

Siegle, J. H., & Wilson, M. A. (2014). Enhancement of encoding and retrieval functions through theta

phase-specific manipulation of hippocampus. *ELife*, *3*, e03061.

Sławińska, U., & Kasicki, S. (1998). The frequency of rat's hippocampal theta rhythm is related to the

speed of locomotion. *Brain Research*, *796*(1–2), 327–331.

Solstad, T., Boccara, C. N., Kropff, E., Moser, M.-B., & Moser, E. I. (2008). Representation of geometric

borders in the entorhinal cortex. *Science*, *322*(5909), 1865–1868.

http://dx.doi.org/10.1126/science.1166466

Sreenivasan, S., & Fiete, I. (2011). Grid cells generate an analog error-correcting code for singularly

precise neural computation. *Nature Neuroscience*, *14*(10), 1330–1337.

https://doi.org/10.1038/nn.2901

Stangl, M., Kanitscheider, I., Riemer, M., Fiete, I., & Wolbers, T. (2020). Sources of path integration error

in young and aging humans. *Nature Communications*, *11*(1), 2626.

https://doi.org/10.1038/s41467-020-15805-9

Taube, J. S. (2007). The Head Direction Signal: Origins and Sensory-Motor Integration. *Annual Review of*

Neuroscience, *30*(1), 181–207. https://doi.org/10.1146/annurev.neuro.29.051605.112854

Taube, J. S., Muller, R. U., & Ranck, J. B. (1990a). Head-direction cells recorded from the postsubiculum

in freely moving rats. I. Description and quantitative analysis. *The Journal of Neuroscience: The*

Official Journal of the Society for Neuroscience, *10*(2), 420–435.

Taube, J. S., Muller, R. U., & Ranck, J. B. (1990b). Head-direction cells recorded from the postsubiculum

in freely moving rats. II. Effects of environmental manipulations. *The Journal of Neuroscience:*

The Official Journal of the Society for Neuroscience, *10*(2), 436–447.

Tcheang, L., Bülthoff, H. H., & Burgess, N. (2011). Visual influence on path integration in darkness

indicates a multimodal representation of large-scale space. *Proceedings of the National*

Academy of Sciences, *108*(3), 1152–1157. https://doi.org/10.1073/pnas.1011843108

Tennant, S. A., Fischer, L., Garden, D. L. F., Gerlei, K. Z., Martinez-Gonzalez, C., McClure, C., Wood, E. R.,

& Nolan, M. F. (2018). Stellate Cells in the Medial Entorhinal Cortex Are Required for Spatial

Learning. *Cell Reports*, *22*(5), 1313–1324. https://doi.org/10.1016/j.celrep.2018.01.005

Viejo, G., & Peyrache, A. (2020). Precise coupling of the thalamic head-direction system to hippocampal

ripples. *Nature Communications*, *11*(1), 2524. https://doi.org/10.1038/s41467-020-15842-4

Weber, S. N., & Sprekeler, H. (2019). A local measure of symmetry and orientation for individual spikes

of grid cells. *PLOS Computational Biology*, *15*(2), e1006804.

https://doi.org/10.1371/journal.pcbi.1006804

Wells, C. E., Amos, D. P., Jeewajee, A., Douchamps, V., Rodgers, J., O'Keefe, J., Burgess, N., & Lever, C.

(2013). Novelty and anxiolytic drugs dissociate two components of hippocampal theta in

behaving rats. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*,

33(20), 8650–8667. https://doi.org/10.1523/JNEUROSCI.5040-12.2013

Whishaw, I. Q., & Vanderwolf, C. H. (1973). Hippocampal EEG and behavior: Changes in amplitude and

frequency of RSA (theta rhythm) associated with spontaneous and learned movement patterns

in rats and cats. *Behavioral Biology*, *8*(4), 461–484. https://doi.org/10.1016/s0091-

6773(73)80041-0

Winter, S. S., Clark, B. J., & Taube, J. S. (2015). Spatial navigation. Disruption of the head direction cell

network impairs the parahippocampal grid cell signal. *Science (New York, N.Y.)*, *347*(6224), 870–

874. https://doi.org/10.1126/science.1259591

Winter, S. S., Mehlman, M. L., Clark, B. J., & Taube, J. S. (2015). Passive Transport Disrupts Grid Signals in

the Parahippocampal Cortex. *Current Biology: CB*, *25*(19), 2493–2502.

https://doi.org/10.1016/j.cub.2015.08.034

Zilli, E. A., & Hasselmo, M. E. (2010). Coupled Noisy Spiking Neurons as Velocity-Controlled Oscillators in

a Model of Grid Cell Spatial Firing. *Journal of Neuroscience*, *30*(41), 13850–13860.

https://doi.org/10.1523/JNEUROSCI.0547-10.2010

1187

FIGURES INCLUDING SUPPLEMENTS AND LEGENDS

1189 **Figure 1**

1190

1191

1192 **Figure 1. Speed signal by local field potential theta frequency decreases in darkness. A,**

1193 Experimental paradigm. Local field potential (LFP) theta oscillatory activity was recorded in mice

freely foraging in a 1-m square open field environment containing a visual cue card during alternating epochs of illumination by a ceiling light (light) and complete darkness (dark). A typical recording lasted 42-min with 3-min alternating light and dark epochs. **B,** Plot of theta frequency vs. running speed shows that the slope of the relationship decreases during darkness. Solid lines show the mean, shaded area show the s.e.m. for n = 15 mice. Data on the light condition are shown in red, data on the dark condition are shown in black and grey. **C,** Same as in B, but for theta amplitude. **D,** Running speed decreases slightly during darkness. Each data point shows the mean value for one mouse, box plots show mean, 25th and 75th percentiles, grey lines connect data from the same mouse; t = 2,90, df = 14, p = 0.0117, Student's paired t-test. **E,** Running acceleration decreases slightly during darkness. Data visualization as in D; t = 3.16, df = 14, p = 0.0070, Student's paired t-test. **F,** LFP theta frequency vs. running acceleration relationship is reduced for all acceleration values in complete darkness. Grey background highlights the interval of acceleration values between -5 cm/s^2 and 5 cm/s^2. **G,** LFP theta frequency vs. running speed relationship based on the subset of time points where the running acceleration is near zero in the interval shown in F between -5 cm/s^2 and 5 cm/s^2. **H-L,** Each data point shows the mean across all epochs of the same condition (light or dark) from 78 recording sessions from 14 mice. **H,** LFP theta frequency speed tuning curves for sequential 10-s blocks after the start of light and dark conditions. Straight lines show the results of linear regression. **I-L,** Yellow background indicates the light condition; grey background indicates the dark condition. 95% confidence intervals are within the line width. **I, J,** Slopes (I) and y-intercepts (J) of the theta frequency vs. running speed relationships for sequential 10-s time bins for the first three minutes in light and dark.. **K, L,** Left and right panels show 10-s binned slopes (K) and y-intercepts of the theta frequency vs. running speed relationship around the transition points from light to dark and from dark to light, respectively.

1220   **Figure 1—figure supplement 1**

1221

1222

Figure 1—figure supplement 1. A, Mean running speed as a function of time around the

transition points from light to dark or from dark to light plotted in 10-s time bins. Data are shown

as mean ± SD across all data points; data from 78 sessions with 3-min alternating epochs of

light and dark from 14 mice. **B,** Speed tuning curves of local field potential (LFP) theta

frequency computed from light (red) or dark (black) epochs of different lengths from the same

animal. Note that the gap between speed tuning curves for the light and dark conditions widens

with the epochs becoming longer suggesting that speed tuning curves change over time.

Figure 2

1231

Figure 2. Changes in theta rhythmic firing with running speed are differentially affected

by visual inputs. Red color or yellow background indicate data on the light condition, black and

grey color or grey background indicate data on the dark condition. Figures show data from theta

rhythmic neurons from a total of 14 mice. **A,** Autocorrelogram of one example neuron showing

theta rhythmic firing in light and dark. Solid red and blue lines show the results of the maximum

likelihood estimate (MLE) model fit. Vertical dashed lines indicate the positions of the peaks in

the autocorrelogram. **B,** Mean probability-normalized autocorrelograms for light and dark

conditions from n = 342 neurons. s.e.m. is within line width. **C,** Figure shows color-coded mean

autocorrelogram fits across n = 340 neurons per running speed bin. Speed bins range from 1 ±

4 cm/s to 29 ± 4 cm/s. Dashed lines connect the first peaks in the mean autocorrelogram

models. **D,** Upper panels show the color coded MLE autocorrelogram models for each of the n =

339 neurons at different running speed intervals. Immobility was defined as running speed < 3

cm/s. Neurons are sorted per condition in order of their theta rhythmic firing frequency. Lower

row shows the average across autocorrelogram models per condition. Note that higher

synchrony across neurons results in higher rhythmicity magnitude in the average across

neurons. **E**, Number of neurons shown in D with statistically significant spiking rhythmicity at

alpha = 0.1 determined by the MLE model for different running speeds in light and dark. **F, G,**

Longitudinal analysis of frequency (F) and magnitude (G) of theta rhythmic firing of neurons with

significant theta rhythmic firing across all running speed bins (light: n = 102 neurons; dark: n =

150 neurons). Data show mean ± s.e.m. **H,** Theta rhythmic firing frequency plotted for

consecutive 10-s time bins after the start of light and dark conditions. **I,** Theta rhythmic firing

frequency plotted for 10-s time bins around the transition points from light to dark and from dark

to light.

1255

1256

1257 **Figure 2—figure supplement 1**

1258

Figure 2—figure supplement 1. A-D, Results with respect to the number of significant neurons

as a function of running speed (first panel) and the frequency (second panel) and magnitude

(third panel) of theta rhythmic firing as functions of running speed are consistent across different

significance levels alpha for theta rhythmic firing used as threshold criteria for classification of

theta rhythmic neurons. Data in second and third panels show mean ± s.e.m. **A,** alpha = 0.1, **B,**

alpha = 0.05, **C,** alpha = 0.01, **D,** alpha = 0.001.

1265

1266

1267 **Figure 2—figure supplement 2**

1268

1269

Figure 2—figure supplement 2. Theta rhythmic firing as a function of running speed and

visual inputs in putative pyramidal neurons. Red color or yellow background indicate data on

the light condition, black and grey color or grey background indicate data on the dark condition.

A, Upper panels show the color coded MLE autocorrelogram models for each of the n = 273

putative pyramidal neurons at different running speed intervals. Immobility was defined as

running speed < 3 cm/s. Neurons are sorted per condition in order of their theta rhythmic firing

frequency. Lower row shows the average across autocorrelogram models per condition. Note

that higher synchrony across neurons results in higher rhythmicity magnitude in the average

across neurons. **B**, Number of neurons shown in A with statistically significant spiking

rhythmicity at alpha = 0.1 determined by the MLE model for different running speeds in light and

dark. **C, D,** Longitudinal analysis of frequency (C) and magnitude (D) of theta rhythmic firing of

neurons with significant theta rhythmic firing across all running speed bins (light: n = 61

neurons; dark: n = 101 neurons). Data show mean ± s.e.m. **E,** Classification of theta rhythmic

neurons into putative pyramidal neurons (blue) and putative interneurons (green) based on the

spike width and firing rate (see Methods). **F,** Theta rhythmic firing frequency plotted for

consecutive 10-s time bins after the start of light and dark conditions. Theta rhythmic firing

frequency was not correlated with time after start of light (p = 0.3688) or dark (p = 0.8898)

conditions.

1288

Figure 2—figure supplement 3

1291 **Figure 2—figure supplement 3. Theta rhythmic firing as a function of running speed and**

1292 **visual inputs in putative interneurons.** Red color or yellow background indicate data on the

light condition, black and grey color or grey background indicate data on the dark condition. **A,** Upper panels show the color coded MLE autocorrelogram models for each of the n = 69 putative interneurons at different running speed intervals. Immobility was defined as running speed < 3 cm/s. Neurons are sorted per condition in order of their theta rhythmic firing frequency. Lower row shows the average across autocorrelogram models per condition. Note that higher synchrony across neurons results in higher rhythmicity magnitude in the average across neurons. **B**, Number of neurons shown in A with statistically significant spiking rhythmicity at alpha = 0.1 determined by the MLE model for different running speeds in light and dark. **C, D,** Longitudinal analysis of frequency (C) and magnitude (D) of theta rhythmic firing of neurons with significant theta rhythmic firing across all running speed bins (light: n = 41 neurons; dark: n = 49 neurons). Data show mean ± s.e.m. **E,** Theta rhythmic firing frequency plotted for consecutive 10-s time bins after the start of light and dark conditions. Theta rhythmic firing frequency was negatively correlated with time after start of light (p = 0.0008) or dark (p = 0.0452) conditions.

1309   **Figure 2—figure supplement 4.**

1310

1311   **Figure 2—figure supplement 4.** Spike time autocorrelograms of randomly picked 24 theta

1312   rhythmic neurons showing theta rhythmic spiking during the light and dark condition.

1313

1314

1315 **Figure 3**

1316

Figure 3. Absence of visual inputs increase LFP theta-frequency phase-locking of single

neuron spiking. A, Probability-normalized polar histograms of LFP theta phase-binned spiking

activity for two example neurons (PYR = putative pyramidal neuron, INT = putative interneuron).

Straight lines show the mean resultant vector scaled between 0 (center of polar histogram) and

1 (outer edge of polar histogram). **B,** Violin plots of theta-phase locking of n = 342 theta

rhythmic neurons during light and dark, p = 1.3 x 10^{-32}, Wilcoxon signed-rank test. **C,** Scatter

plot of preferred LFP theta phases during light and dark. Note the high circular-linear correlation.

D, Probability-normalized histograms of preferred LFP theta phases during light and dark. Note

that the majority of neurons preferred firing around the trough of pyramidal layer LFP theta

oscillations and a minority preferred firing around the peak. No difference was observed

between light and dark conditions, p = 0.9827, Kolmogorov-Smirnov test. **E,** Longitudinal

analysis of the time course of the mean resultant length (MRL) of LFP theta-frequency phase-

locking plotted in 10-s time bins after the start of light or dark conditions, n = 55 neurons. **F,**

Longitudinal analysis of MRL plotted for 10-s time bins around the transition points from light to

dark and from dark to light, n = 58 cells.

1332

1333

1334 **Figure 3—figure supplement 1**

1335

1336

Figure 3—figure supplement 1. Data for n = 273 putative pyramidal neurons (PYR) (A) and n

= 69 putative interneurons (INT) (B). Violin plots show theta-phase locking of theta rhythmic

PYRs or theta rhythmic INTs during light and dark conditions; p-values from Wilcoxon signed-

rank test. Scatter plots show preferred LFP theta phases during light and dark for PYRs (green)

or INTs (blue). Note the high circular-linear correlations for PYRs and INTs. Third panels in A

and B show probability-normalized histograms of preferred LFP theta phases during light and

dark for PYRs and INTs. Note that the majority of both PYRs and INTs preferentially fire around

the trough of pyramidal layer LFP theta oscillations, and a minority preferred firing around the

peak. No significant differences were observed between light and dark conditions in PYRs or

INTs; p-values from Kolmogorov-Smirnov test.

1347

1348   **Figure 4**

1349

Figure 4. Visual inputs alter mean firing rates of neurons in the medial entorhinal cortex.

A-D, Yellow and grey backgrounds indicate duration of light and dark epochs, respectively. **A,**

Data on firing rate from one example neuron in the medial entorhinal cortex (MEC) with its

baseline firing rate negatively modulated by darkness. Black lines show the firing rate at 1-s

resolution. Blue line connects the mean firing rate in each light or dark epoch. **B,** Data on firing

rate from one example neuron in the MEC, whose baseline firing rate is positively modulated by

darkness; same visualization scheme as in A. **C,** Left and right panels show the mean z-scored

firing rate across n = 139 neurons with significantly negative modulation of firing rates by

darkness (see Methods) in 5-s bins around the transition point from light to dark and dark to

light, respectively. Red color indicates data from the light condition, black and grey color indicate

data from the dark condition. Solid lines and shaded areas show mean and SD. **D,** Same as C,

but for neurons with positive modulation of baseline firing rates by darkness; n = 32. **E, F,**

Yellow background indicates light condition, green background indicates the presence of an

egocentrically stable spot of green laser light above the animal's head during the standard light

condition. **E,** Data on one example neuron in the MEC showing positive modulation of baseline

firing rate in presence of the green laser light spot. Same visualization scheme as in A and C. **F,**

As in E, but for one example neuron with negative modulation of baseline firing rate.

1367

Figure 4—figure supplement 1

Figure 4—figure supplement 1. Examples of functional cell types in the medial entorhinal

cortex, whose baseline firing rates are modulated by visual inputs. Example cells include (from

top to bottom) a head direction cell, a grid cell, a border cell, a theta rhythmic cell and a speed

cell. For each functional cell type example: Left most panel shows the firing rate in 1-s time bins

(black solid line) and the mean firing rate of each light or dark epoch (green solid line) for one

recording session. Grey background indicates the dark condition, white background indicates

the light condition. The remaining panels in the top row from left to right show the waveform plot,

trajectory plot, firing rate map, and spatial autocorrelation of the firing rate map. The bottom row

from left to right shows the head direction map, spike autocorrelogram, and speed tuning plot. In

the waveform plot, red solid lines show the mean waveforms of all spikes for the four tetrode

channels (each 1-ms long). Grey shading shows the s.e.m. of the waveform across all spikes. In

the trajectory plot, black lines show the trajectory of the mouse, and red dots indicate the

location of spikes on that trajectory. The firing rate map shows the occupancy-normalized firing

rate per spatial bin, blue color indicates low firing rate, yellow color indicates high firing rate; p =

peak firing rate; m = mean firing rate. Same color scheme applies for the spatial autocorrelation

of the firing rate map. The solid black line in the head direction plot shows the occupancy-

normalized firing rate per head direction bin (bin size is 15 degree) on a circular axis; numbers

next to the inner and outer circle are the circular axis tick mark labels. The solid red line shows

the mean resultant length; R = circular correlation. Black dots in the speed tuning plot show the

mean firing rate across all data points for 2-cm/s running speed bins. The red line shows the

linear regression. R = Pearson's correlation coefficient; p = p-value for linear regression.

1391

1392

 Figure 5

Figure 5. Speed modulation of firing rates is maintained in the absence of visual inputs.

A, Left and right panels show speed response curves for the n = 164 neurons passing statistical significance of speed modulation in the LN-model (see Methods) for either the light or dark condition, respectively. For comparison of speed modulation across neurons and between light and dark conditions, firing rates were normalized and cells are plotted in the same order for light and dark. Cells were sorted by the location of peak firing on the running speed axis averaged across light and dark conditions. Note that speed response curves were normalized *within* conditions, which allows a comparison of speed modulation independent of changes in mean firing rates between light and dark conditions. **B,** Clustergram visualizing the results of the hierarchical clustering of neurons by similarity of speed modulation of firing rates. The clustering

was performed on the neurons' average speed response curves across light and dark

conditions fitted by a quadratic polynomial. **C,** Left and right panels show speed response

curves of negatively (n = 26) and positively (n = 138) modulated neurons; solid lines and shaded

areas show mean ± s.e.m across neurons. **D,** Histogram of speed scores for all neurons

showing significant speed modulation in either the light or dark condition. Red and grey color

indicate data on light and dark, respectively. **E,** Scatter plot on same data shown in D comparing

speed scores during the light with speed scores during the dark condition. **F,** Percentage of cells

showing conjunctive coding of running speed and either position, head direction, or LFP theta

phase during the light and dark condition.

1414

1415

1416 **Figure 5—figure supplement 1**

1417
1418
Figure 5—figure supplement 1. Changes in mean firing rate change the slope of speed
tuning curves without changing the speed modulation of firing rates. A, Left and right
panels show normalized speed response curves for the n = 164 neurons passing statistical
significance of speed modulation in the LN-model (see Methods) for either the light or dark
condition, respectively. Cells are sorted by the location of peak firing on the running speed axis
averaged across light and dark conditions and plotted in the same order for light and dark. Note
that speed response curves were normalized *across* conditions, which allows for a comparison
of mean firing rates between light and dark conditions but precludes an analysis of speed
modulation adjusted for changes in mean firing rate. **B,** Left and right panels show speed
response curves of negatively (n = 26) and positively (n = 138) modulated neurons after
normalization across conditions; In positively modulated speed cells (Pos), the slope and y-
intercept of a neuron's speed response curve decreased significantly in darkness (p (slope) =
0.0038 and p (y-intercept) = 0.0027, linear mixed effects model). In negatively modulated speed
cells (Neg), the y-intercept increased significantly and the slope showed a trend towards a
decrease p (y-intercept) = 0.0091 and p (slope) = 0.0977, linear mixed effects model). Solid
lines and shaded areas show mean ± s.e.m across neurons. **C,** Histogram of speed scores for
all neurons showing significant speed modulation in either the light or dark condition. Red and
grey color indicate data on light and dark, respectively. The distribution of speed scores did not
change significantly between light and dark conditions (p = 0.757, Kolmogorov-Smirnov test). **D,**
Scatter plot of speed scores showing a strong correlation of scores between light and dark
conditions.

 Figure 5—figure supplement 2

1441

Figure 5—figure supplement 2. Figure shows the effect of normalizing LN-model derived
speed response curves (Hardcastle et al., 2017) and speed tuning curves across or within
sessions for ten example cells. For each example cell, the first row shows the LN-model derived
speed response curves and the second row shows the speed tuning curves; the first column
shows speed response and tuning curves with absolute firing rate values, the second column

1447   shows data after normalization across conditions, and the third column shows data after

1448   normalization within conditions. Data for the light condition are shown in red, data for the dark

1449   condition are shown in black. Each data point in the speed tuning curve shows the mean over a

1450   2-cm/s speed bin.

1451

Figure 6

Figure 6. Grid cell spatial firing is maintained but less stable in the absence of visual

inputs. A, Firing rate maps of three example neurons during sequential dark, light, and dark

sessions of 10–20 min duration. In the initial dark session, the animal was introduced to the

open field environment in complete darkness. **B,** Box plots of grid scores during sequential dark,

light, and dark session, n = 9 neurons from two mice. Black color and grey background indicate

data on dark conditions, red color and yellow background indicate data on the light condition.

Solid grey lines connect data points from the same grid cell. Chi-square = 10.89, p = 0.0043,

Friedman's test; post-hoc tests revealed significant differences between the initial dark and light

sessions as well as the initial dark and second dark sessions at alpha = 0.05. **C,** Top, middle,

and bottom rows show trajectory plots, firing rate maps, and spatial autocorrelations of firing

rate maps for three example grid cells recorded over multiple days during alternating light and

dark epochs. Solid black lines in the trajectory plot show the trajectory of the animal, red dots

mark the location of spikes during that trajectory. **D-E,** Box plots of grid scores, spatial

information, and spatiotemporal correlation during light and dark; grey solid lines connect data

points from the same grid cells; p = 2.25 x 10^{-4}, p = 7.26 x 10^{-6}, and p = 2.79 x 10^{-5}; n = 28,

Wilcoxon signed-rank test.

1471

1472

Figure 6—figure supplement 1

Figure 6—figure supplement 1. Spike-triggered firing rate maps of grid cells with a grid score
> 0.06 in either light or dark condition; grid scores computed from the spike-triggered firing rate

1477   maps. Note that grid fields in spike-triggered firing rate maps from the dark condition appear

1478   wider, but grid fields have the same spacing, orientation, and phase than during light.

1479

1480

1481 **Figure 6—figure supplement 2**

1482

1483

1484

Figure 6—figure supplement 2. A, Grid scores computed from spike-triggered firing rate maps

from the light and dark conditions of the n = 15 grid cells shown in Supplementary Figure 1

related to Figure 6. **B,** Grid spacing (first peak location in the radial distance plots shown in

Supplementary Figure 1 related to Figure 6) for the same n = 15 grid cells in the light and dark

conditions. White horizontal lines indicate the median of the distribution. Grey lines connect data

from the same cell. p-values from Wilcoxon signed-rank test.

1491

1492

1493 **Figure 7**

1494
1495

Figure 7. Visual inputs sharpen the tuning curves of head direction cells and border

cells. A, Polar histograms showing the head directional tuning of two head direction cells during

light (red) and dark (black) conditions. **B,** Scatter plot comparing the preferred head direction

angles of n = 67 head direction cells between light and dark conditions. Note the strong circular-

linear correlation. **C,** Violin plots showing the distributions of mean resultant lengths of head

directional tuning curves for light (red) and dark (grey) conditions. White horizontal lines mark

the medians. Grey horizontal lines connect data points from the same neuron. p = 0.0377,

Wilcoxon signed-rank test. **D,** Firing rate maps of four border cells for light and dark conditions.

E, Violin plots comparing the distribution of border scorers of n = 27 border cells between light

and dark conditions. p = 0.0325, Wilcoxon signed-rank test.

1506
1507

Figure 7—figure supplement 1

1510 **Figure 7—supplement 1.** Polar histograms showing the head directional tuning in light and

1511 dark conditions of the 69 head direction cells analyzed in this study. Note that in the dark

1512 condition, head direction cells maintain tuning to their preferred head direction angles, but

1513 tuning curves become slightly broader. Note also that, in addition to tuning curves becoming

1514 broader, mean firing rates are often changed between light and dark conditions.

1515

1516

1517 **Figure 7—figure supplement 2**

1518

Figure 7—figure supplement 2. Firing rate maps for the 27 border cells identified in this study.

For each cell, first panel shows the firing rate map from the whole recording session (light and

dark epochs combined), second panel shows the firing rate map from the light condition, and

third panel shows the firing rate map from the dark condition. For each cell, the same color

scale was used across conditions.

Figure 8

1525

1526

Figure 8. The time course of changes in grid cell spatial stability matches the time course

of changes in the slope of the local field potential theta frequency vs. running speed

relationship. A, Firing rate maps for consecutive 10-s bins of data after transitioning from dark

to light (left side, yellow background) and light to dark (right side, grey background) for one

example neuron. **B-D,** Time course measured in 10-s bins of grid scores (B), spatial correlation

(C), and spatiotemporal correlation (D) during the first three minutes of light and dark sessions.

Solid lines show mean firing rates, shaded area indicates s.e.m.; n = 7 cells from three mice. **E,**

Left and right panels show 10-s binned data on spatiotemporal correlation around the transition

points from light to dark and dark to light, respectively. **F,** Overlay of data shown in Figures 8D

and 1I showing the temporal alignment of changes in the slope of the theta frequency vs.

running speed relationship (blue colors) and the spatial stability of grid cell firing measured by

the spatiotemporal correlation (ocher colors). Left and right panels show data on first three

minutes in the light and dark conditions, respectively. **G,** Scatter plot showing the strong

correlation between the slope of the LFP theta frequency vs. running speed relationship and

spatial stability of grid cell firing as functions of time. Grey line shows the linear regression, R =

0.966; p = 1.83 x 10^{-21}.

Figure 8—figure supplement 1

1544

1545

Figure 8—figure supplement 1. Left side shows the slope of the local field potential (LFP)

theta frequency vs. running speed relationship as a function of 10-s time bins around the

transition points from light to dark and from dark to light for the three individual mice in which

grid cells could be recorded over multiple days and sessions. Right side shows the

corresponding spatial stability of grid cells recorded in each of those mice as a function of 10-s

time bins around the transition points from light to dark and from dark to light. The last row

shows the same data for slope and grid cell stability as mean ± s.e.m. across mice or cells,

respectively.

 TABLES

 Table 1

		Estimate	SE	t-statistic	df	p-value	CI	
Light	y-intercept	-1.34	0.36	-3.73	506	0.0002	-2.05	-0.64
	slope	0.0117	0.0169	0.69	506	0.4921	-0.0216	0.0450
Dark	y-intercept	-0.48	0.33	-1.46	746	0.1454	-1.12	0.17
	slope	0.0327	0.0142	2.31	746	0.0212	0.0049	0.0605

1557 **Table 1 related to Figure 2. Statistics on the differences in frequency of theta rhythmic firing between**
1558 **putative pyramidal neurons and interneurons.** Results from a linear mixed effects model. Values are
1559 referenced to putative pyramidal neurons. SE = standard error, df = degrees of freedom, CI = confidence
1560 intervals.

1561

1562 **Table 2**

		Estimate	SE	t-statistic	df	p-value	CI	
Light	y-intercept	-0.20	0.05	-4.23	506	2.8E-05	-0.30	-0.11
	slope	0.0060	0.0019	3.17	506	0.0016	0.0023	0.0097
Dark	y-intercept	-0.1903	0.0472	-4.03	746	6.10E-05	-0.28	-0.10
	slope	0.0004	0.0011	0.36	746	0.7163	-0.0018	0.0026

1563 **Table 2 related to Figure 2. Statistics on differences in magnitude of theta rhythmic firing between**
1564 **putative pyramidal neurons and interneurons.** Results from a linear mixed effects model. Values are
1565 referenced to putative pyramidal neurons. SE = standard error, df = degrees of freedom, CI =
1566 confidence intervals.

 Table 3

		Estimate	SE	t-statistic	df	p-value	CI	
Pos	y-intercept	-0.0527	0.017571	-2.99941	6896	0.0027	-0.08715	-0.01826
	slope	-0.00176	0.000609	-2.89551	6896	0.0038	-0.00296	-0.00057
Neg	y-intercept	-0.06784	0.025989	-2.61042	1296	0.0091	-0.11883	-0.01686
	slope	-0.00149	0.0009	-1.65711	1296	0.0977	-0.00326	0.000274

1568 **Table 3 related to Figure 5. Statistics on the differences in speed response curves of positively and**

1569 **negatively speed-modulated neurons between light and dark conditions after normalization across**

1570 **conditions.** Results from a linear mixed effects model with the light condition as reference. Pos =

1571 positively modulated, Neg = negatively modulated, SE = standard error, df = degrees of freedom, CI =

1572 confidence intervals.

1573

1574 **Table 4**

		Estimate	SE	t-statistic	df	p-value	CI	
Pos	y-intercept	4.10E-05	0.012502	0.003278	6896	0.9974	-0.02447	0.024548
	slope	-0.00023	0.000433	-0.52947	6896	0.5965	-0.00108	0.00062
Neg	y-intercept	0.067964	0.025506	2.66459	1296	0.0078	0.017926	0.118002
	slope	-0.00349	0.000884	-3.95082	1296	8.21E-05	-0.00523	-0.00176

1575 **Table 4 related to Figure 5. Statistics on the differences in speed response curves of positively and**

1576 **negatively speed-modulated neurons between light and dark conditions after correcting for changes**

1577 **in mean firing rates by normalization within conditions.** Results from a linear mixed effects model with

1578 the light condition as reference. Pos = positively modulated, Neg = negatively modulated, SE = standard

1579 error, df = degrees of freedom, CI = confidence intervals.