WORKING PAPER
Uploaded to bioRxiv

Single-dimensional human brain signals for two-dimensional economic choice options

Leo Chi U Seak*, Konstantin Volkmann*, Alexandre Pastor-Bernier, Fabian Grabenhorst and Wolfram Schultz

* These authors contributed equally to the study.

Department of Physiology
Development and Neuroscience
University of Cambridge
Cambridge CB2 3DY
United Kingdom

Corresponding author:
Wolfram Schultz
Department of Physiology, Development and Neuroscience
University of Cambridge
Cambridge CB2 3DY
United Kingdom
Email: ws234direct@gmail.com

Email addresses of all authors:
Leo Chi U Seak: chiuseak@gmail.com
Konstantin Volkmann: konstantin.volkmann@googlemail.com
Alexandre Pastor-Bernier: pastor-bernier@gmail.com
Fabian Grabenhorst: fabian.grabenhorst@googlemail.com
Wolfram Schultz: Wolfram.Schultz@protonmail.com

Running Title: Neural processing of two-component choice options

Key words: Bundles, decision-making, reward, fMRI, revealed preference, stochastic, value

40 **Abstract** (max 150 words)
41
Rewarding choice options typically contain multiple components, but neural signals in single brain
voxels are scalar and primarily vary up or down. We designed reward bundles that contained the
same two milkshakes with independently set amounts. Using psychophysics and rigorous economic
concepts, we estimated two-dimensional choice indifference curves (IC) that represented revealed
stochastic preferences for these bundles in a systematic, integrated manner. All bundles on the same
ICs were equally revealed preferred (choice indifference); bundles on higher ICs were preferred to
bundles on lower ICs. Functional magnetic resonance imaging (fMRI) demonstrated neural
responses in reward-related brain structures, including striatum, midbrain and medial orbitofrontal
cortex. These responses followed the characteristic pattern of ICs: similar responses along ICs, but
monotonic change across ICs. Thus, these brain structures integrated multiple reward components
into a scalar signal, well beyond the known subjective value coding of single-component rewards.
53
54

Introduction

In daily life, we choose between rewards that have multiple components. In a restaurant, we can get, for the same price, a small but tasty steak or a larger but less tasty steak. In that case, we trade taste against meat. The trade-off shows that our preference for a good is based on more than one component (taste and size in this case). To understand such choices, we need to know how the different components of choice options can be integrated into scalar measures that defines our preferences and the underlying neuronal processes.

Recent functional magnetic resonance imaging (fMRI) studies investigated choices between bundles with multiple-components. A number of brain regions are involved in such choices, including the striatum (Hunt et al. 2014), frontal cortex (Hunt et al. 2014; Kurtz-David et al. 2019; Busemeyer et al. 2019), cingulate cortex (Kurtz-David et al. 2019; Busemeyer et al. 2019; Fujiwara et al., 2009) and insula (Busemeyer et al. 2019). More specifically, one study showed the encoding of values of gift cards that contained an amount component and a quality component (de Berker et al. 2019). Another study used multiple bundles (each with two monetary-gamble components) along the same expected value line to study irrational choices (Kurtz-David et al. 2019). Other studies addressed irrational attraction and decoy effects (Chau et al. 2014; Gluth et al. 2017; Chung et al. 2017). Whereas these studies demonstrated neural signals for multi-component rewards, they did not address the more fundamental mechanism of how such single-dimensional (scalar) neural signals emerge from the integration of multi-dimensional vectorial choice options. To resolve this question would require to study how the increase of one reward would compensate for the decrease of the other bundle reward without changing the preference, and how such a trade-off is represented in scalar neural signals.

Exactly that trade-off mechanism constitutes the heart of indifference curves (IC) that underlie Revealed Preference Theory (Samuelson 1938). Each two-dimensional choice option is graphically represented at a specific x-y coordinate of a two-dimensional plot (Mas-Colell et al. 1995). All bundles that are equally preferred to each other (choice indifferent) are located on the same IC irrespective of their underlying variation in bundle composition. More preferred bundles are located on higher ICs (farther away from the origin). This concept is used to study consumer choice (Simonson 1989; Tversky & Simonson 1993; Rieskamp et al. 2006), animal choice (Kagel et al. 1975; Pastor-Bernier et al. 2017) and neuronal reward signals in animals (Pastor-Bernier et al. 2019). The preference scheme has been extended to stochastic choice (McFadden & Richter 1990; McFadden 2004), which is helpful for statistical analyses of human brain responses in multiple trials. Thus, the question for the current study arises: would human blood-oxygen-level-dependent (BOLD) signals follow the characteristics of ICs that define the emergence of scalar measures from vectorial bundles?

We investigated scalar BOLD signals for two-component milkshakes with sugar and fat components as these stimuli elicit subjective valuations and reliably activate neural reward structures (Grabenhorst et al. 2010; Zangemeister et al., 2016). We used three revealed preference levels (three ICs), each estimated from five equally preferred bundles (indifference points, IPs, located on same IC). Participants were presented with choice options that contained one fatty and one sugary milkshake with specific amounts. We estimated psychophysical indifference points (IP) at which a Reference bundle and a Variable bundle were chosen with equal probability. From these IPs, we estimated well-ordered and non-overlapping ICs. Using two independent general linear models, we found scalar BOLD responses in striatum, midbrain and medial orbitofrontal cortex that followed the systematic scheme of ICs that defines the orderly integration of multi-component choice options into single-dimensional measures; the BOLD responses varied monotonically across ICs but failed to vary significantly along individual ICs.

Results

Implementation of indifference curves

Participants (n = 24) chose between two visual stimuli in repeated trials. Each of the two bundle
stimuli represented a two-component bundle that contained the same two milkshakes with
independently set amounts (Fig. 1A; see Methods). Each stimulus contained two colored vertical
rectangles: the blue rectangle represented component A (low-sugar high-fat milkshake); the red
rectangle represented component B (high-sugar low-fat milkshake). In each rectangle, a vertically
positioned bar indicated the physical amount of each component milkshake, where higher was
more.
We examined choices between: (1) a pre-set Reference bundle and (2) a Variable bundle
whose component A had a fixed test amount and whose component B varied pseudorandomly. In all
24 participants, choice probabilities followed the component B monotonically. We obtained each
indifference point (IP; choice probabilities $P = 0.5$ for each bundle) from a set of six-repetition
choices using a probit choice function (Eqs. 1, 1a). We thereby obtained a two-dimensional IP that
showed the amounts of the two components of the Variable Bundle for which the participant was
indifferent against the constant Reference Bundle. We repeated this procedure, keeping the
Reference Bundle constant and increasing the amounts of component A in the Variable Bundle, thus
obtaining a set of IPs. All IPs in such a set were equally revealed preferred to the constant
Reference Bundle.
In each participant, we estimated a total of three sets of IPs (each contained 5 IPs and their
respective hyperbolic IC) by pre-setting three different amounts of component B (2 ml, 5 ml or 8 ml
with component A always 0 ml) in the Reference Bundle. We derived each IC from such a set of
five IPs by hyperbolic fitting (Eqs. 2, 2a; Fig. 1B, C). These ICs defined the trade-off of the two
components while being equally preferred (i.e. indicating how much of component B the participant
would give up in order to obtain one unit of component A). Taken together, the IPs with the
continuous ICs represented revealed preferences in a systematic manner, thus implementing the
basic concepts underlying this study.
132
Behavioral validation of indifference curves
To assess the contribution and validity of IPs (bundles) to the ICs obtained with hyperbolic fits, we
performed a leave-one-out analysis. These behavioral analyses were presented before (Pastor-
Bernier et al. 2020) and are repeated here for completeness. Briefly, we left out (removed) one IP at
a time from the five IPs within one fitted IC (except for the Reference Bundle at x = 0), and then we
refitted the IC using the remaining four IPs with the same hyperbolic equation (see Methods, Eqs. 2,
2a). We performed the same kind of leave-one-IP-out analysis separately for each IC in each
participant (4 IPs on 3 ICs in 24 participants, resulting in 288 analyses in total).
The refitted ICs resulted in consistent fits in four measures. First, there was no overlap in the
refitted IC with any refitted IC at other levels in all 24 participants; thus, the IC levels retained
separation despite one IP being left-out. Second, there was no overlap in the 72 refitted ICs with the
95% confidence intervals of other original ICs at different levels; thus, the IC levels retained
separation despite one IP being left-out. Third, most refitted ICs (92 %, 66 of 72 ICs) still within the
95% confidence intervals of the original ICs without the eft-out IPs, while the remaining curves
(8%, 6 of 72 ICs) showed only some parts of the IC that fell outside the 95% confidence intervals;
thus, individual IPs were not overweighted in the ICs. Fourth, the left-out IPs deviated only
insignificantly from the refitted ICs ($P = 0.98$ with t-test; $N = 336$; residual: 0.05 ± 0.13 ml in all
participants, mean ± standard error, SEM) (Fig. 1D); this result confirmed that individual IPs were
not overweighted in the ICs. These four validations demonstrated the robustness and consistency of
the hyperbolically fitted ICs in capturing the IPs. Thus, in all participants, the ICs provided valid
representations of the three revealed preference levels.
154
Neural responses for two-component bundles across and along ICs
During fMRI scanning, the task started with a fixation cross lasting 0.5 s (Fig. 1E). Then, a single
two-component visual stimulus appeared in the center of the computer monitor (bundle-on phase);
the stimulus predicted delivery of one of the 15 bundles (IPs) composed of two different

milkshakes. The physical amount of the milkshakes in the bundle was determined by the
participant-specific indifference point (IP) estimated from the binary choice task (see above). The
participant received the two bundle milkshakes with the respective amounts indicated by the
vertical bars on the stimulus, without choice. That presentation was either followed by a Becker-
DeGroot-Marschak (BDM) task within the trial (50% of trials, pseudorandomly selected) or
terminated (50% of trials). The BDM bidding served as a mechanism-independent measure of
utility estimation, as used before (de Berker et al. 2019; De Martino et al. 2013). In total, each
participant performed 360 trials (24 trials for each of the 15 bundles). With the fMRI data we
collected, we analyzed the various aspects of neural responses (BOLD signals) to the bundles with
several General Linear Models (GLMs) and region-of-interest (ROI) analyses.

We first used GLM1 to identify brain regions that followed the scheme of revealed
preferences represented by ICs, namely monotonic increase across increasing ICs (or decrease with
inverse coding) and insignificant change along the same ICs, as shown in Fig. 2A. To do so, the
individual contrast images (representation of BOLD signal) of each bundle in each participant were
grouped according to the IC the bundle belonged to (low, medium, high) and the position of the
bundle on each IC (1 - 5, from top left to bottom right).

We used parametric statistical tests (t-test with Flexible Factorial Design) and estimated
neuroimages of responses to each of the 15 bundles grouped into the three IC levels or five groups
along ICs (see Methods). We found that the striatum, midbrain and OFC showed significantly
increasing activation across increasing ICs (high > low IC; map threshold of $p < 0.005$; t-test) but
insignificant variations along individual ICs (exclusive mask map threshold of $p < 0.005$) (Fig. 2B).
More specifically, we found small-volume corrected significance in the striatum (peak at [10, 6, -4],
z-score = 3.27, 6 mm radius sphere, cluster-level FWE corrected $p = 0.041$), midbrain (peak at [4, -
16, -12], z-score=3.71, 6 mm radius sphere, cluster-level FWE corrected $p = 0.048$) and OFC (peak
at [22, 42, -16], z-score = 3.67, 10 mm radius sphere, cluster-level FWE corrected $p = 0.037$). (All
small-volume corrections in this study were centered on pre-defined coordinates, see Methods). In
addition, we found significant activities in other regions, including the insula and cingulate cortex
(Table 1).

To provide further evidence for neural activations following the scheme of ICs, we performed
a Spearman rank time course analysis. We first extracted BOLD signals using leave-one-subject-out
cross-validated GLM models, which should prevent potential biases with pre-selected peaks (see
Methods). Subsequently we used the BOLD signals from peak voxels in each left-out subject to
perform Spearman rank analyses. We found that the striatum, midbrain and OFC showed significant
Spearman rank correlation coefficients (Spearman's Rho) between bundles located on different ICs
at around 6s after onset of the bundle stimulus ($p < 0.05$), consistent with the standard time course
of haemodynamic response (Fig. 2C). By contrast, only insignificant ($p > 0.05$) rank coefficients
were found at 5 - 7s between bundles located along same ICs in these brain regions, as shown in the
sliding-window analysis (Fig. 2D). These time courses followed the revealed preference to bundles
across different ICs but failed to differ along the same IC, thus complying with the scheme of ICs
that represent revealed preference. Moreover, we extracted beta (slope) coefficients of the BOLD
signal at 6s with the ROI coordinates identified by GLM1 and plotted them for three revealed
preference levels in bar charts (Fig. 2E). We found a significant difference between high versus low
revealed preference level in the midbrain ($p = 0.0062$), OFC ($p = 0.0023$), and marginal significant
difference in the striatum ($p = 0.0533$). We also found a significant difference between high versus
middle revealed preference level in the OFC ($p = 6.8551 \times 10^{-4}$).

Taken together, these data showed that activations in several components of the brain's reward
system followed the basic scheme of ICs representing revealed preferences: activation across the
ICs but no activation along the same IC. Based on these results, a more stringent analysis should
test more closely how a scalar preference signal might emerge from the vectorial, multi-component
choice options.

Binary comparisons between partial physically non-dominant bundles

According to the concept of ICs, any bundle on a higher IC (farther from the origin) should be preferred to any bundle on a lower IC. Hence, a single-dimensional neural signal reflecting multi-component choice options should vary between any bundle on a higher IC and any bundle on a lower IC. To reflect the proper integration of the two bundle components irrespective of specific physical properties, the neural signal should follow the IC rank even when one component milkshake of the higher-IC bundle is lower than in the lower-IC bundle (partial physical non-dominance). To identify such differences, we used the GLM2. With pairwise comparisons, GLM2 should identify higher responses to revealed preferred bundles with partial physical non-dominance. Thus, GLM2 compared all bundle pairs that fit the following condition within each participant: bundle 1 was located on higher IC but had a lower amount of one component milkshake compared to bundle 2 that was located on a lower IC (Fig. 3A).

The GLM2 analysis demonstrated significant activations in similar regions as with GLM1, where striatum (peak at [16, 6, -6], z-score=3.8, 6 mm radius sphere, cluster-level FWE corrected p = 0.012), midbrain (peak at [4, -16, -12], z-score=2.85, 6 mm radius sphere, cluster-level FWE corrected p = 0.032) and OFC (peak at [24, 42, -16], z-score=3.99, 10 mm radius sphere, cluster-level FWE corrected p = 0.012) showed small-volume corrected significant activations (Fig. 3B). Also, we found significant activities in other regions, including insula, superior frontal gyrus and cingulate, as shown in Table 2.

We also performed ROI analyses (coordinates identified by GLM2 with leave-one-subject-out procedure) that calculated betas of partial physical non-dominance (higher revealed preference) and partial physical dominance bundles (lower revealed preference) as described in Methods. For each ROI, we computed three models, which compared bundles pairwise, with low vs. middle, middle vs. high, and low vs. high revealed preference, respectively. Neural beta regression coefficients were extracted at 6 s after the onset of the bundle stimulus, which corresponded to the canonical hemodynamic response. In regard to high vs. low revealed preference level, we found significance in the striatum ($p = 0.0459$) and OFC ($p = 0.0033$) when comparing bundles in high IC vs. low IC (Fig. 3C). We also found significance in the striatum ($p = 0.0309$) and OFC ($p = 7.6575 \times 10^{-5}$) when comparing high vs. middle IC bundles. In the midbrain, we found no significance ($p > 0.05$) in the three comparisons between bundles on low, middle and high ICs (although such a tendency existed in all three comparisons).

Taken together, these pairwise bundle comparisons demonstrated neural coding of partial physical non-dominance bundles as a necessary condition for extracting a scalar neural signal from vectorial, multi-component choice options. These results confirmed compliance with the graphic schemes of ICs demonstrated with GLM1.

Becker-DeGroot-Marschak (BDM) control of revealed preference
To validate the order of revealed preferences represented by the ICs with an independent estimation mechanism, we used a monetary Becker-DeGroot-Marschak (BDM) bidding task that estimated each participant's utility for each bundle. In 50% of trials during fMRI scanning, each participant made a monetary BDM bid (UK pence) for one of the 15 bundles, out of a fresh endowment of 20 UK pence in each trial (BDM bidding phase; Fig. 1E). The 15 bundles constituted the indifference points of the ICs that were estimated during the binary choice task with each participant (Fig. S1).

The BDM bids followed the order of revealed preference levels across ICs, as evidenced by the significant positive correlation between the three IC levels and the bid amounts for bundles (Rho = 0.5710 and $p < 1.5659 \times 10^{-32}$; Spearman Rank correlation from averages of all five bundles; $N = 24$ participants; further confirmed with binary Wilcoxon signed-rank test between the three IC levels, IC1 vs IC2: $p = 1.2802 \times 10^{-20}$; IC2 vs IC3: $p = 8.0748 \times 10^{-21}$; IC1 vs IC3: $p = 1.5954 \times 10^{-19}$) (Fig. 4A; blue, green, red). By contrast, there was no correlation between bids for the five bundles and their position along each IC (from top left to bottom right; Spearman Rho = 0.0219; $p = 0.6791$). Thus, BDM bids increased across the three IC levels but did not change monotonically with bundle position along individual ICs in the population of our participants.

In order to investigate neural mechanisms of value elicitation and BDM bidding, we compared two GLM models: (1) GLM1 to identify brain regions that encoded IC levels during bundle-on phase; (2) GLM3 to identify brain regions that encoded BDM bids (0 - 20 pence) during the bidding phase. As described above, we found significant, small-volume corrected activation in OFC in GLM1 that indicated its involvement in encoding IC levels during the bundle-on phase. This result was confirmed by an additional ROI analysis that showed significant rank correlation between OFC activation and IC levels at around 6 s after bundle onset (Fig. 4B ; bundle-on phase; $p < 0.05$; Spearman's Rho), consistent with the expected haemodynamic response function.

By contrast, analysis with GLM3 demonstrated activation in vmPFC that encoded BDM bids during the bidding phase (Fig. 4C left; peak at $[6, 44, 0]$, z-score = 4.10, whole-brain corrected with cluster-level FWE corrected $p = 0.002$), together with other brain regions (Table 3). Further ROI analysis showed significant rank correlation between vmPFC activation and BDM bids at around 6 s after BDM cue onset (Fig. 4C right; bidding phase; $p < 0.05$; Spearman's Rho), consistent with the expected haemodynamic response function. In addition, we found small-volume corrected significant activation in the dorsal striatum (Supplementary Fig. 2; Speak at $[12, 12, 0]$, z-score = 3.53, 6 mm radius sphere, cluster-level FWE corrected $p = 0.008$) encoding the BDM bids. Different from GLM1 and GLM2, such activities were not found (cluster-level FWE corrected $p > 0.1$) in the ventral striatum.

Taken together, BDM bidding provided a good validation of the estimated levels of revealed preference represented by ICs. However, and interestingly, revealed preference levels and BDM bids were encoded in different regions of the frontal cortex and striatum.

Discussion

We systematically tested characteristics of scalar neural responses to vectorial, multi-component bundles. We estimated indifference points (IPs) by asking human participants to make choices between two bundles. Our behavioral results (Pastor-Bernier et al. 2020) showed that preference relationships among multi-component choices were reliably represented by systematic ICs, as a prerequisite for testing the underlying neural mechanisms. In fMRI scans with GLM and post-hoc ROI analyses, we identified brain regions whose activations correlated with levels of revealed preference. The GLM1 and post-hoc Spearman rank analysis demonstrated activations in the ventral striatum, midbrain and OFC that reflected revealed preference levels across ICs but failed to vary along equal-preference ICs. The GLM2 specifically dissociated revealed preference from physical dominance and showed consistent results with those from GLM1. A mechanism-independent control with a Becker-DeGroot-Marschak (BDM) bidding task confirmed the validity of ICs for representing revealed preference levels. Interestingly, however, BDM bidding was associated with activations in vmPFC and dorsal striatum rather than the previously identified reward structures following IC levels. Together, these data demonstrate systematic, single-dimensional neural activations in the striatum, midbrain and OFC that reflect vectorial multi-component choice options.

In our binary choice task, we elicited revealed preferences with repeated, psychophysically controlled choices (Pastor-Bernier et al. 2020; Green, & Swets 1966). Such a multi-trial, stochastic approach is well conceptualized (McFadden & Richter 1990; McFadden 2004), fulfils the statistical requirements of neural research, corresponds to standard choice functions (Sutton, & Barto 1998) and allows direct comparison with animal neurophysiology requiring similar multi-trial schedules (Pastor-Bernier et al., 2017). These methods systematically delivered varying choice probabilities (stochastic choices) instead of single selections (deterministic choices).

Economic choice experiments often involve substantial but imaginary sizes or amounts of consumer items and money, or use random singular payouts (Simonson, 1989; Tversky & Simonson 1993; Rieskamp el al., 2006). By contrast, we tailored our payout schedule to fit the requirements of neuroimaging studies and made our participants choose tangible and consumable rewards over hundreds of trials, while also controlling for satiety effects. These behavioral choices resembled small daily activities, such as consumptions of drinks and snacks. In this way, we obtained three

well-ordered ICs for each participant that provided accurate and systematic representations of
revealed preferences for multi-component bundles, without involving imagined items or monetary
reward (Pastor-Bernier et al. 2020).
We used the BDM task as an authoritative, mechanism-independent control for eliciting
subjective values, thereby providing an additional validating mechanism for the revealed
preferences elicited in our binary choice test. The value estimating mechanism for BDM bids differs
substantially from the one for revealed preference ICs. The truthful revelations (incentive
compatibility) of BDM makes this mechanism an essential tool in experimental economics that is
becoming more popular in human decision research (Plassmann et al., 2007; Medic et al., 2014;
Zangemeister et al., 2016). The elicited BDM bids correlated well with the revealed preference
levels and thereby validated the empirically estimated IPs used during fMRI in which the
participants performed the BDM task. Previous neuroimaging studies showed activations in
ventromedial prefrontal cortex that correlated with BDM bids (Chib et al., 2009; McNamee et al.,
2013). Our experimental design dissociated value elicitation by bundles and by BDM bidding. We
confirmed the BDM activations in vmPFC and found that the two mechanistically different tasks
activated different regions in both prefrontal cortex and striatum; responses to the bundles followed
the IC scheme (different activations across but not within ICs) in OFC and dorsal striatum, whereas
BDM bidding activated vmPFC and ventral striatum.
Previous studies tested neural mechanisms of human choice of bundles with multiple
components, such as payoff amount and probability (Chau et al. 2014), quality and quantity of
goods (de Berker et al. 2019), money and time (Gluth et al. 2017), and food components (Suzuki et
al. 2017). Nevertheless, none of these studies tested bundles that were positioned along modelled
ICs (i.e. eliciting choice indifference) and thus failed to test the crucial trade-off that demonstrates
the graded and well-ordered manner of single-dimensional preferences for multi-dimensional choice
options. Without this information, we would not know how a scalar neural response may arise from
graded changes of vectorial, multi-component bundles. Our study, testing 5 bundles on each IC,
addressed this problem and identified the brain regions that showed this kind of neural response.
Although we tested the emergence of single-dimensional neural signals for multi-dimensional
bundles in a systematic and concept-driven way, there were limitations with our experimental
design. First, both bundle components had the same type of primary reward (milkshakes). It would
be interesting to study whether the same brain regions would encode different types of rewards and
follow the formalisms of ICs, including the graded trade-off. For instance, future research may
compare monetary rewards with primary nutrient rewards. Second, we only demonstrated neural
responses with the typical convex ICs. It would be interesting to study whether different brain
regions might encode preferences with different shapes of ICs. Such work may test participants'
choices with linear or concave ICs. Third, we did not test the influences of prior experience on
current decisions. Previous studies (Schultz 1998; van den Bos et al. 2013; Lopez-Persem et al.
2016) showed that choices could be influenced by previous experience and be updated by
reinforcement learning. Future research may include multi-component choice options during fMRI
scanning to study multi-component reinforcement learning. Lastly, we only demonstrated fMRI
BOLD responses, and future neurophysiology research should confirm the coding of revealed
preference at a single neuron level in human patients with intracerebrally implanted electrodes,
similar to our recently investigated neuronal encoding of revealed preference in monkey
orbitofrontal cortex (Pastor-Bernier et al. 2019). To conclude, while we showed brain activation
with bundles in a formal but standard revealed preference setting (convex ICs, primary reward), it is
desirable to know how human brains encode revealed preference in a larger variety of situations.
The reward circuit including the striatum and midbrain is known to participate in reward
anticipation and learning, including reward prediction error (Diederen et al. 2017). In monkeys,
midbrain dopamine neurons encode values for predicted rewards in economic decision tasks (Lak et
al. 2016; Schultz et al. 2017). Similar to the midbrain and striatum, previous work showed the
involvement of the human mid-OFC in valuation of primary nutrient reward (Grabenhorst et al.,
2010) and monetary reward (Kahnt et al. 2014). Remarkably, the neural activity in OFC elicited

here, in response to visual cues predicting liquid rewards with varying sugar and fat components, closely matched the coordinates observed previously (Grabenhorst et al., 2010) in a study in which subjects orally sampled very similar liquid rewards. Thus, this area of OFC seems to be involved both in reward valuation during oral consumption of primary nutrient rewards and in the economic valuation of visually cued choice options. In non-human primates, OFC neurons encode reward prediction (Tremblay & Schultz 1999; Padoa-Schioppa & Assad 2006) and follow revealed preferences for multi-component bundles (Pastor-Bernier et al. 2019). In the current study, we used a concept-driven design and found that neural responses in the striatum, midbrain and OFC integrated multiple bundle components in a way that followed the ICs scheme (changing across ICs but being similar along equal-preference ICs). Moreover, we demonstrate the involvement of the midbrain in multi-component decision making for the first time. Overall, our results show the involvement of principal reward structures of the brain in integrating the multiple components of vectorial bundles into single-dimensional neural signals that are suitable for economic decision making.

Besides the primary reward circuit (midbrain dopamine neurons, OFC, striatum, amygdala), other brain regions are also involved in economic decision making. Previous studies in multi-component decision making suggested the involvement of the cingulate, prefrontal cortex and insula in value elicitation (Kurtz-David et al. 2019; Busemeyer et al. 2019). Consistent with these studies, we also found significant activation in these regions. As shown in Table 1 and Table 2, the BOLD signals identified by GLM1 and GLM2 showed that these regions also encode bundle values during the bundle-on phase, together with the striatum, midbrain, and mid-OFC. Our results are consistent with these previous studies, suggesting that a considerable number of brain regions also play a role in multi-component decision making.

Methods

Participants

A total of 24 participants (19-36 years old with mean age 25.4 years; 11 males, 13 females) performed a binary choice task that was followed, in 50% of trials, by a Becker–DeGroot–Marschak (BDM) task inside the fMRI scanner using sugary and fatty milkshakes. All participants had known milkshake appetite, and none had diabetes or lactose intolerance. All participants provided written consent based on an information sheet. The Cambridgeshire Health Authority (Local Research Ethics Committee) approved this study. The behavioral results have been published (Pastor-Bernier et al. 2020).

Experimental design

The fundamental notion underlying this experiment posits that choice options consist of at least two components, and that preferences are revealed by observable choice. The multi-component choice options are called bundles. Decision makers prefer bundles with larger amounts of components to those with smaller amounts of components. Importantly, however, their preferences concern all components and are not directed at a single component alone. This property is manifested when participants prefer bundles in which one of the components of the preferred bundle is smaller than the same component in the non-preferred bundle (and the other component is large enough to overcompensate). At one point, participants may express equal preference for bundles in which the lower amount of one component is fully compensated by the higher amount in the other component, leading to choice indifference. We repeatedly measured choices with two options, each of which contained two milkshake components; the milkshakes constituted rewards, as shown by the voluntary consumption in all participants.

Stimuli and rewards

In each of the two bundles, we used stimuli to show the two milkshake components and their
payout amounts (Fig. 1A). In each bundle stimulus, there were two rectangles aligned vertically.
Each bundle component was indicated by the color of each rectangle. We extensively piloted
various liquidized foods and liquids, and we found that milkshakes with a controlled mixture of fat
and sugar give the most reliable across-participant behavioral performance. The presently used
milkshakes with sugar and fat components that were found in previous studies to elicit subjective
valuations and activate neural reward structures (Grabenhorst et al. 2010; Zangemeister et al.,
2016). We delivered the milkshakes separately with a 0.5 s interval (see below). As drinks
consisting of only sugar or only fat were considered as too unnatural, we used a high-fat low-sugar
milkshake (75% double cream and 25% whole milk, with no sugar) as component A (top, blue), and
a high-sugar low-fat milkshake (skimmed milk with 10% sugar) as component B (bottom, red).
Inside each rectangle, the vertical position of a bar indicated the component's physical amount
(higher was more). We delivered the milkshakes to the participants using a custom-made silicone
tubing syringe pump system (VWR International Ltd). The pump was approved for delivering
foodstuffs and was controlled by a National Instruments device (NI-USB-6009) via the Data
Acquisition Toolbox in Matlab. We displayed stimuli to participants and recorded behavioral
choices using the Psychtoolbox in Matlab running on a Windows (Dell) computer (Pastor-Bernier et
al. 2020).

Binary choice task before fMRI scanning
In the binary choice task, each participant revealed one's preference in repeated choices between
two bundle stimuli, each indicating the amounts of two milkshake components (Fig. 1A). The two
bundles (stimuli) appeared on a computer screen simultaneously in front of the participant. The left
and right positions of the bundles were fixed but pseudorandomly alternated. Each bundle stimulus
included the same two kinds of milkshakes with independent physical amounts. Both stimuli
appeared after a pseudorandomly varying interval (mean 0.5 s) after a central fixation cross. In each
trial, the participant chose between the two bundles by pressing a button once (on a computer
keyboard; left or right arrow corresponding to choosing left or right bundle). We defined reaction
time as the interval between appearance of the two bundle stimuli and the participant's button press.
We delivered the two milkshakes to the participant from the chosen bundle with a probability $P =$
0.2 using a Poisson distribution; i. e. the milkshake combination of one out of an average of five
chosen bundles was delivered, and no milkshake was delivered in the remaining trials. Component
B (high-sugar low-fat milkshake) was delivered 500 ms (constant interval) after component A
(high-fat low-sugar milkshake). We used this constant delay, instead of simultaneous delivery of
two milkshakes or a pseudo-randomly alternating milkshake sequence, to prevent uncontrolled
milkshake interactions, to maintain distinguishability of the individual milkshake rewards and to
keep temporal discounting constant. Therefore, the utility of component B derived from both
milkshake rewards and the temporal discounting specific for each milkshake. Participants were
asked not to eat or drink anything at least four hours before the task performance. However, satiety
may still be a concern given the high fat and sugar content of our milkshakes. To address this issue,
we set the probability of $P = 0.2$ payout schedule, limited each payout to 10.0 ml at most, and
delivered no more than a total of 200 ml of liquid to the participant in a session. We addressed the
issue with additional analyses and failed to find differential, sensory-specific satiety noticeable in
choice probability measures (see below; Pastor-Bernier et al. 2020).

Psychophysical assessment of indifference points (IPs)
We used a psychophysical staircase method (Pastor-Bernier et al. 2020; Green, & Swets, 1966) to
estimate the indifference points (IPs) at which, by definition, each of the two bundle options was
chosen equally frequently (i.e. probability $P = 0.5$ for each option), indicating choice indifference

for the options. We established bundles at 15 IPs for each participant and used them in the
subsequent fMRI experiment.
To start the psychophysical procedure, we first set component A to 0 ml and component B to
either 2 ml, 5 ml or 8 ml in the Reference Bundle. We then systematically varied the Variable
Bundle. In the Variable Bundle, we first set the amount of its component A to one unit higher
(mostly 0.5 ml, 1.0 ml or 2.0 ml); we thereby specified the amount of component A gained by each
participant from the choice. We then randomly selected (without replacement) one amount of
component B from a total of seven fixed amounts (multiples of 0.5 ml), which span the whole,
constant range of amounts being tested. We repeatedly selected the amounts until we tested each of
the seven amounts once. We repeated estimation for each IP six times using a sigmoid function (see
Eqs. 1, 1a below), requiring a total of 42 choices for estimating each IP. The amount of component
B in the Variable Bundle was usually lower than the one in the Reference Bundle at the IP. With
these procedures, we assessed how much of component B a participant was willing to trade-in for
an additional unit of component A.
We obtained more IPs from the participants' choices between the fixed Reference Bundle and
the Variable Bundle, in which the amount of component A was increased stepwise, at each step
varying the amount of component B to estimate the chocie indifference point at which the animal
was indifferent between the two bundles. Thus, bundle position advanced from top left to bottom
right on the two-dimensional IC (Fig. 1B). We are aware that testing with unidirectional progression
may cause particular variations in IP estimations than testing in a random sequence or in opposite
directions (Knetsch, 1989). However, our primary interest in this study was to investigate basic
neural processes in close relation to unequivocally estimated IPs and ICs rather than addressing the
more advanced features of irreversibility or hysteris in ICs.
We used three different fixed amounts of component B for the Reference Bundle (2 ml, 5 ml,
or 8 ml), to obtain three IC levels. We estimated four IPs, together with the fixed reference bundle
as an IP, at each of three indifference curves (ICs; i.e. revealed preference levels), resulting in 15
IPs, in a total of 504 choices (trials) among 84 different choice option sets in each participant (6
repetitions for 7 psychophysical amounts at each of the 12 IPs).
492
Statistical analysis
Numeric estimation of indifference points
We used a sigmoid fit to numerically estimate the choice IPs. The fit was obtained from the
systematically tested choices with a generalized linear regression. The generalized linear
regressions used the *glmfit* function in Matlab (Matlab version R2015b) with a binomial distributed
probit model, which is an inversed cumulative distribution function (G). More specifically, we
apply the link function to the generalized linear regression $y = \beta_0 + \beta_1 B_{var} + \varepsilon$ and write it as:
500
501 $$G(y) = \beta_0 + \beta_1 B_{var} + \varepsilon \qquad\qquad\qquad\qquad \text{Eq. 1}$$
502
where y represents the number of trials the Variable Bundle is chosen in each block of a six-
repetition series, β_0 represent the constant offset, β_1 represent the regression slope coefficient, B_{var}
represent the physical reward amount (ml) of component B in the Variable Bundle, and ε represent
the residual error. We used the probit model as it assumes a multivariate normal distribution of the
random errors, which makes the model attractive because the normal distribution gives a good
approximation to most of the variables. The model does not hypothesize error independence and is
frequently used in econometrics (Razzaghi, 2013). On the other hand, the logit model, which is also
commonly used in economics, is simpler to compute but has more restrictive hypotheses on error
independence. Our preliminary data had shown a similar fit for both the logit and probit model,
therefore, we used the probit model fit because of its less restrictive hypotheses. Thus, we
approximated the IPs with the probit-model sigmoid fit, which can be written as follows:
514
515 $$\text{Indifference Point} = - (\beta_0 / \beta_1) \qquad\qquad\qquad\qquad \text{Eq. 1a}$$

516
517   where β_0 and β_1 represent coefficients of the generalized linear regression (Eq. 1). We obtained
518   these coefficients from the probit analysis (Amemiya, 1981).
519
Indifference curves (ICs)
In each participant, we obtained each single IC separately from an individual set of five equally
revealed preferred IPs with differently composed bundles using a weighted least-square non-linear
regression. We used a weighted regression to account for choice variability within participant; the
weight was defined as the inverse of the standard deviation of the titrated physical amount of
component B at the corresponding IP (the IP having been estimated with the probit regression). We
estimated the best β coefficients from the least-square regression to obtain a single IC (utility level),
using the basic hyperbolic equation:
528
529      $IC = \beta_0 + \beta_1 B + \beta_2 A + \beta_3 BA + \varepsilon$ Eq. 2
530
where A and B represent physical amounts of component A and component B (ml), which refer to
the x and y axis, respectively. Note that (β_2 / β_1) is the slope coefficient and β_3 is the curvature
coefficient of the non-linear least-square regression. As IC is a constant (representing one utility
level), we merged the IC constant with the offset constant (β_0) and the error constant (ε) into a
common constant k. To draw the ICs, we calculated the amount of component B from the derived
equation as a function of the amount of component A:
537
538      $B = (k - \beta_2 A) / (\beta_1 + \beta_3 A)$ Eq. 2a
539
We graphically displayed the fitted ICs (Fig. 1B, C) by plotting the pre-set physical amount of
component A as the x coordinates, and calculated the fitted amount of component B, based on Eq.
2a, as the y coordinates. We estimated the error of the hyperbolic fit as the 95% confidence interval.
When calculating the ICs, we gave less weight to the IP with higher error. This model offered good
fits in our earlier work (Pastor-Bernier, et al. 2017; 2019; 2020). In this way, five IPs aligned to a
single fitted IC. For each participant, we fitted three ICs representing increasing revealed preference
levels (low, medium, high) farther away from the origin (Fig. 1B, C). The indifference map that
resulted from the 3 x 5 IPs was unique for each of the 24 participants (Supplementary Fig. 1).
548
Leave-one-out validation of ICs
We used a leave-one-out analysis to test the validity of the hyperbolic IC fit to the IPs. We
systemically removed one IP in each IC (excluding the initial Reference Bundle at x = 0), and then
fitted the IC again using the hyperbolic model. We then assessed the differences (deviation)
between the original IC (without IP removal) and the new IC without the one left-out IP. The
deviation was defined as the Euclidean distance of component B between the original (left-out) IP
and the IP estimated from the refitted IC:
556
557      $d = B_{IP} - B_{refit}$ Eq. 3
558
559   with d representing the difference (i.e. residual; in ml; y-axis), B_{IP} representing the physical amount
560   of component B in the left-out IP (ml), and B_{refit} representing the estimated physical amount of
561   component B in the refitted IC (ml). In this way a residual of 0 ml suggested that removal of the
562   left-out IP did not change the shape of that IC, while any residual unequal to 0 ml could quantify the
563   deviation.
564
565   *Control of alternative choice factors*
566   To assess the potential influence of other factors affecting the participants' choice, we performed a
567   logistic regression fit on choices to test whether the choices were indeed explained by the bundle

568 components. We performed a random-effect logistic regression on the choice data from each
569 participant as follows:
570
571 $y = \beta_0 + \beta_1 RefB + \beta_2 VarA + \beta_3 VarB + \beta_4 RT + \beta_5 VarPos + \beta_6 PChoice + \varepsilon$ Eq. 4
572
with y as a dummy variable (either 1 or 0, indicating choosing or not choosing the Variable Bundle),
RefB as physical amount (ml) of component B in the Reference Bundle, VarA and VarB as physical
amount (ml) of components A and B in the Variable Bundle, RT as reaction time (ms), VarPos
indicating left or right position (0 or 1) of the Variable Bundle stimuli shown on the computer
screen relative to the Reference Bundle, and PChoice representing choice of the previous trial (0 or
1). Each β coefficient was normalized by multiplying the standard deviation of the respective
independent variable and dividing by the standard deviation of the dependent variable (y). We
subsequently used a one-sample t-test against 0 to assess the statistical significance of each of the
beta (β) coefficients.
We assessed the normalized beta (β) coefficients and p-values for each individual participant
and then calculated averages across 24 participants. With the regression model, we found a negative
correlation of choosing the Variable Bundle and the amount of component B in the Reference
Bundle (RefB: $\beta = -0.43 \pm 0.16$, $P = 0.020 \pm 0.005$; mean $\pm$ SEM) (amount of component A in the
Reference Bundle was always a constant 0 ml). We also found positive correlation of choosing the
Variable Bundle and amount of both component A and component B in the Variable Bundle (VarA:
$\beta = 0.67 \pm 0.16$, $P = 0.009 \pm 0.004$; VarB: $\beta = 0.94 \pm 0.33$, $P = 0.012 \pm 0.009$). We further found
that for these three variables, the beta (β) coefficients significantly differed from 0 with one-sample
t-tests ($P = 0.012$, $P = 0.00088$ and $P = 0.00028$, respectively), confirming the robustness of these
β. Thus, we confirmed that the choices depended on the amount of reward of both Variable and
Reference Bundle. We also validated that both bundle components were important for the choices.
All remaining variables in the regression, including reaction time, left or right position of the
Reference Bundle on the computer screen and choice of the previous trial, failed to account
significantly for the participant's current choice ($P = 0.754 - 0.988 \pm 0.003 - 0.290$). We therefore
conclude that, in our experiment, the bundles with their two components, instead of other factors,
account for the revealed preference relationships.
598
Satiety control
Besides considering other components in the design, we also tested potential effects of satiety.
Satiety may have affected the preferences for the two bundle components, even if the rewards were
paid out only in one fifth of the trials on average and were limited to less than 200 ml. Differences
in devaluation between the two component milkshake might be a major factor for changing in an
uncontrolled manner the currency relationship of the two components. This kind of unequal
devaluation should result in a graded change in the instantaneous choice probability around the IPs
over the test steps of 42 trials. We used the following equation to calculate the instantaneous choice
probability:
608
609 $y = \sum$ (n=1 to 6) (CV / TS)   Eq. 5
610
611 with y representing the instantaneous probability (P ranging from 0.0 to 1.0), CV represent choice
612 or not-choice of Variable Bundle (1 or 0), and TS represent test step (repetition 1-6).
613   We found only insignificant fluctuations in choice probabilities, without any consistent
614 upward or downward trend in the 1-way repeated measures ANOVA, together with the post-hoc
615 Tukey Test (above IP: F (5, 41) = 0.28, $P > 0.05$; below IP: F (5, 41) = 1.53, $P > 0.05$).
616
Behavioral task during fMRI scanning
During scanning, we used a value elicitation task that allowed more trials in a shorter time frame. At
the beginning of each trial, one bundle was shown to the participant for 5 s (bundle-on phase in Fig.

1E) in the center of the computer monitor after the initial fixation period (500 ms). The bundle was
pseudorandomly selected from the 15 IP bundles in three ICs of each participant. Bundle
composition (amounts of the two components) was set in each participant according to performance
in the binary choice task before fMRI scanning. Hence, the 15 bundles for each participant were not
identical across participants. Subsequently, a fixation cross appeared for a pseudorandomly varying
interval (mean 2s). In 50% of the trials (pseudorandomly selected), the task was terminated after
this fixation cross.
In the other 50% of the trials, we presented the participant with a Becker-DeGroot-Marschak
(BDM) task that was akin to a second price auction (Becker, DeGroot, & Marschak, 1964). This
task served as an independent mechanism that related the estimated ICs to stated utility. In the BDM
(bidding phase in Fig. 1E), we gave the participant a fresh 20 UK pence endowment on each trial.
Using this endowment, the participant bid for a two-component bundle against a pseudorandom
computer bid (extracted from a normal distribution with replacement). To bid, the participant moved
a cursor, shown on the computer screen, horizontally with the left and right keyboard arrows. We
registered the BDM bid (position of the cursor) 5 s after presenting the bidding scale to the
participant. When bidding no less than the computer, the participant received the bundle
(milkshake) reward from both components and paid the monetary value equal to the computer bid.
By contrast, when bidding less than the computer, the participant lost the auction, paid nothing and
would not get any bundle (milkshake) reward. We showed the participant the result of the auction
immediately after having placed the bid, by displaying a respective win (green circle) or loss (red
square) stimulus on the computer monitor (Fig. 1E); when winning the bid, the participant received
the milkshake rewards in the sequence and frequency as in the binary choice task.
We first selected one bundle randomly (without replacement) from the participant-specific set
of 15 bundles (the 15 bundle IPs used to fit the 3 ICs as shown in Supplementary Fig. 1). Then we
showed the participant the selected single bundle during the bundle-on phase. We presented each of
the 15 bundles to the participant for 24 times, resulting in a total of 360 trials, which included 180
trials (50%) with BDM bidding (Fig. 1E), and we used the average of these bids as the participant's
BDM-estimated utility.
First, we assessed whether the BDM bids increased for bundles across revealed preference
levels but were similar for IP bundles on the same revealed preference level, using Spearman rank
correlation analysis and further confirmation with the Wilcoxon signed-rank test (note that this
analysis used the coordinates of the individual IPs to which the ICs had been fitted, not the IC
coordinates themselves). We also performed a generalized linear regression with a Gaussian link
function (random-effect analysis) for each participant and then averaged the β coefficients and p-
values across all participants. We used the following generalized linear regression:

$y = \beta_0 + \beta_1\ PrefLev + \beta_2\ AmBundle + \beta_3\ TrialN + \beta_4\ PrevBid + \beta_5\ Consum + \varepsilon$ Eq. 6

with y representing amount of monetary bid, PrefLev representing revealed preference level (low,
medium, high), AmBundle representing the summed amount (ml) of component A and component
B in the currency of component A (converted with Eq. 2a), TrialN representing trial number,
PrevBid representing amount of monetary bid in the previous trial, and Consum representing
accumulated consumption amount (ml) of component A and component B until that point in the
experiment. We normalized each β coefficient by multiplying the standard deviation of the
respective independent variable, and then dividing by the standard deviation of the dependent
variable y. We performed a subsequent one-sample t-test against 0 to assess the significance of each
beta (β) coefficient across all 24 participants. We found significant beta (β) coefficients of BDM
monetary bids to the preference level (PrefLev: β-coefficient difference from 0: P = 0.000026 with
one-sample t-test; mean across all 24 participants: $\beta = 0.47 \pm 0.09$, P = 0.016 ± 0.015; mean± SEM)
and bundle amount (AmBundle: P = 0.0278; $\beta = 0.15 \pm 0.13$; P = 0.020 ± 0.017), but not in trial
number (TrialN: $\beta= -0.10 \pm 0.25$; P = 0.726 ± 0.354), previous trial bid (PrevBid: $\beta= 0.12 \pm 0.11$; P
= 0.676 ± 0.427) nor consumption history (Consum: $\beta= 0.12 \pm 0.11$; P = 0.224 ± 0.185).

fMRI data acquisition

The functional neuroimaging data in this study were collected using a 3T Siemens Magnetom Skyra
Scanner at the Wolfson Brain Imaging Centre, Cambridge, UK. Echo-planar images (T2-weighted)
with blood-oxygen-level-dependent (BOLD) contrast were acquired at 3 Tesla across two days with
each participant. All images were in plane resolution 3 x 3 x 2 mm, 56 slices were acquired with 2
mm slice thinness, repetition time (TR) = 3 s, echo time (TE) = 30 ms, -90 deg flip angle and -192
mm field of view. To reduce signal dropout in medial-temporal and inferior-frontal regions during
the scanning, the acquisition plane was tilted by -30 degrees and the z-shim gradient pre-pulse was
implemented. We also applied MPRAGE sequences and co-registered to acquired high-resolution
T1 structural scans for group-level anatomical localization with 1 x 1 x 1 mm^3 voxel resolution,
slice thickness of 1 mm, 2.3 s TR, 2.98 ms TE, 9 deg flip angle and 900 ms inversion time.

fMRI data analysis

We used the Statistical Parametric Mapping package to analyze the neuroimaging data (SPM 12;
Wellcome Trust Centre for Neuroimaging, London). We pre-processed the data by realigning the
functional data to include motion correction, normalizing to the standard Montreal Neurological
Institute (MNI) coordinate, and then smoothing using a Gaussian kernel with the full width at half
maximum (FWHM) of 6 mm within data collected on the same day. We then segmented the data to
extract white matter, grey matter and cerebrospinal fluid (CSF) and followed by co-registering the
two-day data using the T1-weighted structural scans from each day. We then applied a high-pass
temporal filter to it with a 128 s cut-off period. We applied General linear models (GLMs), which
assumed first-order autoregressions, to the time course of activation. We modeled event onsets, in
the time course of activation, as single impulse response functions convolved with the canonical
hemodynamic response. We included the time derivatives in the functions set and defined linear
contrasts of parameter estimates to test the specific effect in each participant's dataset. We obtained
voxel values for each contrast in the format of a statistical parametric map with corresponding t-
statistic. We applied a standard explicit mask (mask_ICV.nii) at the first level analysis to mask out
all activations outside of the brain. To test our specific hypotheses, we used the following GLMs:

General linear model 1 (GLM1)

This GLM served to search for regions whose stimulus-induced brain activations varied across ICs
(high > low) but not along the same ICs in the bundle-stimulus-on phase (two-level t-test analysis,
Fig. 2). For each participant, we estimated a GLM with the following regressors (R) of interest:
(R1-R15) as indicator functions for each condition during the bundle-on phase (for the 15 different
bundles), at the time when participant was presented with the visual bundle cue representing the
milkshakes bundles; (R16) as indicator function for the BDM bid, at the time when the participant
made the bid; (R17) as R16 that was modulated by the response to the participant's bid (1 - 20);
(R18) as indicator function for the losing bid, at the time when the participant was presented with
visual cues showing the loss of bidding of the trial; (R19) as indicator function for the auction win
phase, at the time when the participant was presented with the visual cues representing the winning
of bidding; (R20) as indicator function for the reward phase, i.e. the times when participants
received the milkshakes; (R21) as R20 that was modulated by reward magnitude (in mL).
In the second (group random-effects) level analysis, we entered the all 24 participant-specific
linear contrasts of the first-level regressors R1-R15 (representing 5 bundles on each of the three
preference levels) into t-tests (high > low revealed preference level) using Flexible Factorial
Design, resulting in group-level statistical parametric maps. In the Flexible Factorial design matrix
(second-level analysis), the following second-level regressors were used: (R1-24) indicator
functions of participant's identifier representing participant 1 - 24 (within participant effect); (R25-
27) indicator functions of the three revealed preference levels (across ICs) (R28-32), indicator
functions of the 5 bundles representing amount of Component A in increasing magnitude or amount
of Component B in decreasing magnitude (along the same ICs). We first calculated the main

contrast image based on high>low revealed preference level (t-tests). Second, we calculated a mask
contrast based on 5 bundles of Component A in increasing magnitude (t-tests). Third, we calculated
another mask contrast based on 5 bundles of Component B in increasing magnitude (t-tests). The
final result of GLM1 was represented by the main contrast (high>low revealed preference level)
masking out (with exclusive mask) the two mask contrasts, controlling of the brain responses along
the same ICs.

General linear model 2 (GLM2)
This GLM identified regions associated with the binary comparisons of partial physical non-
dominance bundles (Fig. 3). The GLM searched for brain regions in which activations were higher
for bundles that were on a higher revealed preference level than bundles in which one component
was physically higher than in the preferred bundle (partial physical non-dominance). In the first-
level estimation, regressors were the same as in GLM1 with the 21 regressors described above. In
the second-level analysis, we entered all pairs of bundles that met the following criteria: (Bundle 1):
partial physical non-dominance bundles with higher revealed preference level, but less (with at least
0.2mL less in Components A or 0.4mL less in Component B) in one component; (Bundle 2): partial
physical dominance bundles with lower revealed preference level, but more in one component. A
third level group-level analysis (one-sample t-test) was performed with contrast images from the
second level to generate group-level statistical parametric maps across 24 participants.

General linear model 3 (GLM3)
This GLM identified brain regions in which activity correlated with the amount of BDM bid (0 - 20
pence) during the bidding phase (Fig. 4B, C). In the first-level estimation, we used the following
regressors and parametric modulators: (R1) as indicator function of bundle-on phase; (R2) as R1
modulated by amount of BDM bid; (R3) as indicator function of BDM bidding phase (50% of
trials); (R4) as R3 modulated by amount of BDM bid; (R5) as indicator function of intertrial
interval when there was no bidding phase (50% of trials); (R6) as indicator function at onset of the
loss cue, when the participant lost the BDM bidding; (R7) as indicator function at onset of the win
cue, when the participant won the BDM bidding; (R8) as indicator function at onset of milkshake
delivery; (R9) as R8 modulated by physical amount of milkshake; (R10) as contrast of win cue
onset versus loss cue onset; (R11) as contrast of loss cue onset versus win cue onset. In the second-
level analysis, a one-sample t-test analysis was performed with contrast images from the first level
to generate group-level statistical parametric maps across 24 participants.

Small volume corrections
To derive coordinates for small-volume correction in GLM1 and GLM2, we entered the term
"reward anticipation" in the Neurosynth meta-analysis database (Yarkoni et al., 2011) to obtain
MNI coordinates. The meta-analysis employed a total of 92 independent studies that showed
correlation of value elicitation with various brain regions. Our study used MNI coordinates of
ventral striatum [12, 10, -8], medial orbital frontal cortex (mid-OFC) [20, 46, -18] and midbrain [8,
-18, -14], obtained from this Neurosynth meta-analysis database.
We aimed at finding activity correlating with the BDM bid in GLM3. Therefore, for small
volume correction analysis in GLM3, we used a MNI coordinate of dorsal striatum [12, 14, 4]
found in a previous study with BDM bidding (De Martino et al., 2009).

Region-of-interest (ROI) analysis
We selected significantly activated regions from brain maps established with GLM1, GLM2 or
GLM3 for further ROI analysis. We extracted raw BOLD data from ROI coordinates based on
group clusters, which we defined independently for each participant using a leave-one-out
procedure based on the result of GLM1, GLM2 or GLM3. In the leave-one-out procedure, we re-
estimated the second-level analysis 24 times, each time leaving out one participant, to define the
ROI coordinates for the left-out participant. Following data extraction, we applied a high-pass filter

with a cut off period of 128 s. The data was then z-normalized, oversampled by a factor of 10 using
the Whittaker–Shannon interpolation formula, and separated into trials to produce a matrix of trials
against time.
A total of 3 ROI analyses were performed in this study. First, a Spearman rank analysis was
used to examine BOLD signals that changed across ICs but not along ICs (corresponding to GLM1
and GLM3). Second, a bar chart was used to illustrate the three revealed preference levels in
different ROIs (corresponding to GLM1). Third, a bar chart was used to show activation changes
between bundles with partial physical non-dominance on different revealed preference levels
(corresponding to GLM2).
785
Spearman rank
In the Spearman rank analysis, we first regressed out the motion parameters (artefact) from the
BOLD response with generalized linear models. Then we used the participant's residual BOLD
response to generate time courses of Spearman rank correlation (Rho) coefficients.
For GLM1, we tested the correlation between BOLD response (during the bundle-on phase)
and revealed preference level (across-IC analysis). We then calculated group averages and standard
errors of the mean for each time point for all participants, yielding averaged participant effect size
time courses (Fig. 2C). In the along-IC analysis, we ranked the bundles along the same IC with
individual participant's BOLD signal (Fig. 2D). A subsequent one-sample t-test against 0 served to
assess the significance of the Rho coefficients across subjects.
For GLM3, we tested the correlation of the BOLD response (BDM bidding phase) and the
amount of BDM bids. Similar to GLM1, we then calculated group averages and standard errors of
the mean of the Rho coefficients for each time point for all participants (Fig. 4C; Supplementary
Fig. 2). A subsequent one-sample t-test against 0 served to assess coefficient significance.
800
Bar chart for revealed preference level analysis
We used bars to illustrate how different IC levels were encoded in each region of the brain. To
generate an ROI bar chart, the BOLD response was first extracted using the leave-one-out
procedure described above. For each participant, we obtained three generalized linear model fits to
the BOLD signal at timepoint 6 s. In each generalized linear model fit, the identifier of one level of
revealed preference was entered as a regressor (dummy variable, e. g. 1 for bundles with high
preference level and 0 for middle or low preference level) together with motion parameter
regressors, which served to eliminate the motion artefact. We obtained beta (β) coefficients of each
level of revealed preference from the fit and then calculated the mean and standard error of the beta
coefficient. We then plotted the bar charts shown in Fig. 2E. Paired t-tests were used to compare
beta coefficients between different revealed preference levels.
812
Bar chart for partial physical non-dominance analysis
A bundle was defined as being partially physically non-dominant over another bundle if one of its
milkshake components had a physically lower amount than the same component in the dominated
bundle. Thus, the revealed preferred bundle was partially physically non-dominant. For an ROI
analysis of partial physical non-dominance, we fitted three generalized linear models to the BOLD
response with bundle identifiers, which were two dummy variables representing partially physically
dominance bundles (lower revealed preference despite larger physical amount in one milkshake)
and partial physical non-dominance bundles (higher revealed preference despite smaller physical
amount in one milkshake). Three generalized linear models were used to fit bundles in low vs.
middle, middle vs. high, and low vs. high comparisons, respectively. The domination was defined as
at least 0.2 ml more for component A or at least 0.4 ml more for component B, as in GLM2. We
calculated the mean and standard error of the averaged beta (β) coefficients across participants at
time point 6 s and plotted the bar chart as shown in Fig. 3C. Paired t-tests were used to compare
beta coefficients between partial physical dominance bundles and partial physical non-dominance

bundles. Motion parameters were also used as regressors for each participant to eliminate motion
artefacts.
829
Reward prediction errors (RPE)
The current task did not involve learning in which reward would occur in a partly unpredicted
manner and thus elicit RPEs. The only RPE could occur at the unpredicted time of the first stimulus
that explicitly and quantitatively predicted the reward amounts of the bundle components indicated
by the bundle stimulus. Conceivably, in the most simple form, the RPE would reflect the integrated
reward amounts of both bundle components relative to the prediction derived from the past trial
history. There were three levels of bundle stimulus corresponding to the three IC levels. Thus,
appearance of a given bundle stimulus would elicit an RPE relative to the past experienced bundles,
weighted by the learning coefficient. Thus, reward prediction errors would have values around -1, 0,
and +1 for bundles located on low, intermediate and high ICs, respectively, the variation depending
on the learning coefficient. For comparison, the bundle stimulus at each IC level without any RPE
would have values of 1, 2 and 3, respectively. Thus, neural responses to the RPE and to the stimulus
directly (i. e. without subtraction of prediction) would result in very similar regression slopes
(depending on the learning constant used for computing the RPE) and thus be difficult to
distinguish from each other. We modelled RPEs with various learning coefficients in the range
between 0.1 and 0.9 and for all values found high correlations between RPE and bundle stimulus
value at the three IC levels. For example, a learning coefficient of 0.2 in a Rescorla-Wagner model
resulted in a Spearman-rank correlation of 0.9337 ± 0.00085 SEM (n=15 bundles x 24 trials = 360
trials x 24 subjects pooled). For this reason, a RPE analysis would not yield new insights and will
not be further reported.
850
851 **Acknowledgements**
852

853 We thank Charles R. Plott for discussions and conceptual support, Steve Edgley for help and
854 logistic support, Simone Ferrari-Toniolo for comments on experimental economics, Jae-Chang Kim
855 and Putu Agus Khorisantono for fMRI analytical suggestions, and Arkadiusz Stasiak for computer
856 support. The Wellcome Trust supported this work (WT 095495, WT 204811).

857
858 **References**
859

Amemiya, T., (1981). Qualitative response models: A survey. Journal of economic literature, 19(4),
pp.1483-1536.
Becker, G.M., DeGroot, M.H., Marschak, J., (1964). Measuring utility by a single-response
sequential method. Behavioral science, 9(3), pp.226-232.
Busemeyer, J.R., Gluth, S., Rieskamp, J., Turner, B.M., (2019). Cognitive and neural bases of
Multi-Attribute, Multi-Alternative, Value-based decisions. Trends in cognitive sciences.
Chau, B. K. H., Kolling, N., Hunt, L. T., Walton, M. E. & Rushworth M. F. S. (2014). A neural
mechanism underlying failure of optimal choice with multiple alternatives. Nat Neurosci 17,
463-470.
Chib, V.S., Rangel, A., Shimojo, S., O'Doherty, J.P., (2009). Evidence for a common representation
of decision values for dissimilar goods in human ventromedial prefrontal cortex. J Neurosci
29(39), pp.12315-12320.
Chung, H.K., Sjöström, T., Lee, H.J., Lu, Y.T., Tsuo, F.Y., Chen, T.S., Chang, C.F., Juan, C.H., Kuo,
W.J., Huang, C.Y., (2017). Why do irrelevant alternatives matter? an fMRI-TMS study of
Context-Dependent preferences. Journal of Neuroscience, 37(48), pp.11647-11661.
de Berker, A.O., Kurth-Nelson, Z., Rutledge, R.B., Bestmann, S., Dolan, R.J., (2019). Computing
Value from Quality and Quantity in Human Decision-Making. Journal of Neuroscience, 39(1),
pp.163-176.

De Martino, B., Fleming, S.M., Garrett, N., Dolan, R.J., (2013). Confidence in value-based choice.
Nature Neuroscience, 16(1), p.105.
De Martino, B., Kumaran, D., Holt, B., Dolan, R. J. (2009). The neurobiology of reference-
dependent value computation. Journal of Neuroscience, 29(12), 3833-3842.
Diederen, K.M., Ziauddeen, H., Vestergaard, M.D., Spencer, T., Schultz, W., Fletcher, P.C., (2017.
Dopamine modulates adaptive prediction error coding in the human midbrain and striatum.
Journal of Neuroscience, 37(7), pp.1708-1720.
Fujiwara, J., Tobler, P.N., Taira, M., Iijima, T., Tsutsui, K.I., (2009). Segregated and integrated
coding of reward and punishment in the cingulate cortex. Journal of Neurophysiology, 101(6),
pp.3284-3293.
Gluth, S., Hotaling, J. M., & Rieskamp, J. (2017). The attraction effect modulates reward prediction
errors and intertemporal choices. J Neurosci 37: 371-382.
Grabenhorst, F., Rolls, E .T., Parris, B.A., D'Souza, A. (2010). How the brain represents the reward
value of fat in the mouth. Cerebral Cortex, 20: 1082-1091.
Green DM, Swets J. Signal detection theory and psychophysics: New York: Wiley, (1966).
Hunt, L.T., Dolan, R.J., Behrens, T.E., (2014). Hierarchical competitions subserving multi-attribute
choice. Nature neuroscience, 17(11), p.1613.
Kagel, J. H., Battalio, R. C., Rachlin, H., Basmann, R. L., Green, L., & Klemm, W. R. (1975).
Experimental studies of consumer demand behavior using laboratory animals. Econ Inquiry 13:
22-38.
Kahnt, T., Park, S.Q., Haynes, J.D., Tobler, P.N., (2014). Disentangling neural representations of
value and salience in the human brain. Proceedings of the National Academy of Sciences,
111(13), pp.5000-5005.
Knetsch, J. L. (1989). The endowment effect and evidence of nonreversible indifference curves. Am
Econ Rev 79: 1277-1288.
Kurtz-David, V., Persitz, D., Webb, R., Levy, D.J., (2019). The neural computation of inconsistent
choice behavior. Nature communications, 10(1), p.1583.
Lak A, Stauffer WR, Schultz W. (2016). Dopamine neurons learn relative chosen value from
probabilistic rewards. eLife 5: e18044.
Lopez-Persem, A., Domenech, P., Pessiglione, M., (2016). How prior preferences determine
decision-making frames and biases in the human brain. Elife, 5, p.e20317.
Mas-Colell, A., Whinston, M.D., Green, J.R., (1995). Microeconomic theory (Vol. 1). New York:
Oxford university press.
McFadden, D., & Richter, M. K. (1990). Stochastic rationality and revealed stochastic preference.
In: Preferences, Uncertainty, and Optimality, Essays in Honor of Leo Hurwicz, Westview
Press: Boulder, CO, 161-186.
McFadden, D. L. (2004). Revealed stochastic preference: A synthesis. *Econ Theory, 26*, 245–
264.McNamee, D., Rangel, A., O'doherty, J.P., (2013). Category-dependent and category-
independent goal-value codes in human ventromedial prefrontal cortex. Nature neuroscience,
16(4), p.479.
Medic, N., Ziauddeen, H., Vestergaard, M. D., Henning, E., Schultz, W., Farooqi, I. S., & Fletcher,
P. C. (2014). Dopamine modulates the neural representation of subjective value of food in
hungry subjects. J Neurosci 34: 16856-16864.
Padoa-Schioppa, C., Assad, J.A., (2006). Neurons in the orbitofrontal cortex encode economic
value. Nature, 441(7090), p.223.
Pastor-Bernier, A., Plott, C.R., Schultz, W., (2017). Monkeys choose as if maximizing utility
compatible with basic principles of revealed preference theory. Proceedings of the National
Academy of Sciences, 114(10), pp.E1766-E1775.
Pastor-Bernier, A., Stasiak, A., Schultz, W., (2019). Orbitofrontal signals for two-component choice
options comply with indifference curves of Revealed Preference Theory. Nature
communications, 10(1), pp.1-19.

Pastor-Bernier, A., Volkmann, K., Stasiak, A., Grabenhorst, F., Schultz, W., (2020). Experimentally
revealed stochastic preferences for multi-component choice options. J exp Psychol: Anim Lern
Cognition (in press).
Plassmann, H., O'Doherty, J., Rangel, A. (2007). Orbitofrontal cortex encodes willingness to pay in
everyday economic transactions. J Neurosci 27: 9984-9988.
Razzaghi, M. (2013). The Probit Link Function in Generalized Linear Models for Data Mining
Applications. J. Mod. Appl. Stat. Meth. 12 : Issue 1, Article 19.
Rieskamp, J., Busemeyer, J. R., & Mellers, B. A. (2006). Extending the bounds of rationality:
Evidence and theories of preferential choice. J Econ Lit 44: 631-661.
Samuelson, P.A., (1938). A note on the pure theory of consumer's behavior. Economica, 5(17),
pp.61-71.
Schultz, W., (1998). Predictive reward signal of dopamine neurons. Journal of neurophysiology,
80(1), pp.1-27.
Schultz, W., Stauffer, W.R., Lak, A., (2017). The phasic dopamine signal maturing: from reward via
behavioural activation to formal economic utility. Current opinion in neurobiology, 43, pp.139-
148.
Simonson, I. (1989). Choice based on reasons: The case of attraction and compromise effects. J
Consum Res 16: 158-174.
Sutton, R. S., Barto, A. G. (1998). Reinforcement Learning: An Introduction. Cambridge MA: MIT
Press.
Suzuki S, Cross L, O'Doherty JP (2017). Elucidating the underlying components of food valuation
in the human orbitofrontal cortex. Nature neuroscience. Dec;20(12):1780-6.
Tremblay, L., Schultz, W., (1999). Relative reward preference in primate orbitofrontal cortex.
Nature, 398(6729), p.704.
Tversky, A., & Simonson, I. (1993). Context-dependent preferences. Management Sci 39: 1179-
1189.
van den Bos, W., Talwar, A., McClure, S.M., (2013). Neural correlates of reinforcement learning
and social preferences in competitive bidding. Journal of Neuroscience, 33(5), pp.2137-2146.
Yamagata N, Ichinose T, Aso Y, Plaçais PY, Friedrich AB, Sima RJ, Preat T, Rubin GM, Tanimoto
H. Distinct dopamine neurons mediate reward signals for short-and long-term memories.
Proceedings of the National Academy of Sciences 112(2):578-83.
Yarkoni, T., Poldrack, R.A., Nichols, T.E., Van Essen, D.C., Wager, T.D., (2011). Large-scale
automated synthesis of human functional neuroimaging data. Nature Methods, 8(8), p.665.
Zangemeister L., Grabenhorst F., Schultz W. (2016). Neural basis for economic saving strategies in
human amygdala-prefrontal reward circuits. Current Biology, 26: 3004-3013.
964

966
967
Figure 1. Experimental procedure and behavior.
(A) Choice task outside the fMRI scanner. The participant chose between a reference bundle and a
varied test bundle. Each bundle consisted of two components, Component A (blue bar) and
Component B (red bar). The amount of each component was indicated to the participant by the
height of a white bar (higher was more). Component A was a low-sugar, high-fat milkshake.
Component B was a low-fat high-sugar milkshake. The two milkshakes of the chosen bundle were
delivered at the end of each trial with a probability of $P = 0.2$.
(B) Schematic diagram of three indifference curves (ICs) and five indifference points (IPs) on each
IC.
(C) Example ICs from a typical participant. Solid lines represent three ICs (hyperbolically fitted by
IPs). Dotted lines represent 95% confidence interval of the hyperbolic fit. The inset shows the
psychophysical function of one IP. The IP (black dot in the inset) was estimated by probit regression
on the test points (blue dot in the inset).
(D) Histogram of residuals between fitted ICs (with a leave-one-out procedure) and left-out IPs
across all participants. The residuals formed a normal symmetric distribution (red line).
(E) Bundle task inside the fMRI scanner. At 4 s after the bundle-on phase, the participant performed
in pseudorandomly selected 50% of trials a Becker-DeGroot-Marschak (BDM) value elicitation
task against the computer (bidding 1 - 20 UK pence). The reward was given if the participant won
the BDM (bid $\geq$ computer bid).
987

989
990
Figure 2. BOLD responses following the revealed preference scheme of two-dimensional
indifference curves (ICs).
(A) Schematic of the analysis method used in GLM1: significant BOLD signal across ICs but not
within ICs (arrows). Participants typically showed convex ICs (left) or linear ICs (right).
(B) BOLD responses discriminating bundles between ICs (map threshold p < 0.005, extent
threshold ≥ 10 voxels), but no discrimination between bundles along same ICs (map threshold p >
0.005; i.e. exclusive mask for brain response fall along the same ICs with threshold p=0.005) in a
group analysis.
(C) Across-IC Spearman rank analyses of brain activations. The Rho coefficients followed the
haemodynamic response function (HRF) across the 3 IC levels in the ROIs of the three brain
structures shown above in B. Solid blue lines represent mean Rho from 24 participants; ± SEM.
Yellow shaded boxes show analysis time window. Green asterisks p < 0.05, blue asterisks p < 0.01
for t-test of Spearman's Rho against zero. The BOLD responses (input of the Spearman rank
analyses; with motion parameters regressed out) were extracted from the peak voxels of each
participant using with a leave-one-out procedure (see Methods).
(D) Along-IC ROI activations. The Spearman rank analyses indicated hardly any significance along
same ICs in ROIs of the three brain structures shown in B.
(E) Bar charts of neural beta coefficients of GLM1 for the three IC levels in the three brain
structures shown in B in 24 participants. Bars show mean ± SEM.
1010

1011

1012
1013
Figure 3. Higher BOLD responses to more preferred (but physically partially dominated) bundles
positioned on different indifference curves (ICs).
(A) Two examples of binary bundle comparison. Each pair of black circles indicates one binary
comparison in one participant.
(B) Brain regions activated more by preferred bundles compared to alternative bundles in group
analysis with GLM2. Map threshold p < 0.005, extent threshold ≥ 10 voxels.
(C) Bar charts showing neural beta coefficients of regression in ROIs of three brain structures in the
population of 24 participants. Each group of bars (3 groups in each ROI) shows the beta coefficients
for bundles in partial physically dominating relationships on different ICs: low vs. mid; mid vs.
high and low vs. high. Orange bars represent the higher preference level and blue bars represent the
lower preference level in each comparison. The bars show the mean ± SEM.
1025

1026

1027
1028
Figure 4. Differential activation in distinct prefrontal cortex areas during value elicitation (OFC)
and BDM bidding (vmPFC).
(A) Bar chart for BDM bids for 15 bundles of 24 participants (mean ± SEM). The colors of the bars
indicate the indifference curves (IC) to which the bundle belongs (blue = low IC; green = middle
IC; red = high IC). Spearman rank correlation: across ICs: Rho = 0.5710, p = 1.5659 x 10^{-32}; within
IC: Rho = 0.0219, p = 0.6791. Wilcoxon signed rank test: IC1 vs. IC2: p = 1.2802 x 10^{-20}; IC2 vs.
IC3: p = 8.0748 x 10^{-21}; IC1 vs. IC3: p = 1.5954 x 10^{-19}.
(B) OFC activation during bundle-on phase follows IC scheme (left): significant activation across
ICs but not within ICs (GLM1: threshold p < 0.005, extent threshold ≥ 10 voxels). Spearman rank
ROI analysis (right) showed significant Rho coefficients across the 3 ICs, consistent with the
haemodynamic response function (HRF) during bundle-on phase in OFC.
(C) vmPFC activation during bidding phase (GLM3: activation correlated with the amount of BDM
bids; threshold p < 0.005, extent threshold ≥ 10 voxels). Brain map (left) shows dorsal striatum
activity during bidding phase. Spearman rank analysis (right) showed significant Rho coefficient
across bids during bidding phase in vmPFC.
1044

1045 **Table 1.** Brain regions activated across but not along indifference curves (ICs) during bundle-on
1046 phase (whole-brain analysis with GLM1).
1047

Brain region	Hemisphere	MNI peak coordinates (x,y,z)	peak z-score
Striatum*	R	10, 6, -4	3.27
Midbrain*	/	4, -16, -12	3.09
OFC*	R	22, 42, -16	3.67
Parieto-occipital transition zone/ occipital gyri	/	-12, -66, 46	7.42
Insular gyrus/ basal operculum	L	-30, 18, 2	5.80
	R	32, 26, -4	4.91
Superior frontal gyrus	/	-24, 2, 52	5.70
Middle frontal gyrus	L	-42, 2, 42	4.98
	L	-40, 34, 18	4.74
	R	44, 46, 16	4.94
Cingulate gyrus	/	-2, -24, 28	4.87
Precentral gyrus	R	40, 6, 24	4.59
Angular gyrus	L	-22, -72, 54	3.89

1048
1049 Cluster P values (P < 0.05) with family-wise error correction across the whole brain. Map threshold
1050 P < 0.005 (across ICs; high>low IC) with exclusive contrast map P > 0.005 (along ICs), extent
1051 threshold ≥ 10 voxels. *P < 0.05 with small volume correction.
1052

1053
1054 **Table 2.** Brain regions showing differences (partial physical non-dominance> partial physical
1055 dominance) in BOLD signal between partial physically dominating bundles located on different
1056 indifference curves (ICs) during bundle-on phase (whole-brain analysis with GLM2).
1057

Brain region	Hemisphere	MNI peak coordinates (x,y,z)	peak z-score
Striatum*	R	16, 6, -6	3.8
Midbrain*	/	4, -16, -12	2.85
OFC*	R	24, 42, -16	3.99
Insular gyrus/ Basal operculum	L	-30, 24, -2	5.55
	R	32, 26, -6	4.40
Angular gyrus	R	32, -68, 28	5.51
Cerebellum	L	-36, -68, -30	4.79
Superior frontal gyrus	/	22, 2, 54	4.75
Occipital gyri	L	-28, -88, 4	4.57
Middle frontal gyrus	L	-50, 40, 16	4.32
	R	46, 42, 14	3.76
Inferior frontopolar gyrus	R	18, 64, -8	4.11
Cingulate gyrus	/	-2, -24, 28	4.05

1058
1059 Cluster P values ($P < 0.05$) with family-wise error correction across the whole brain. Map threshold
1060 $P < 0.005$, extent threshold ≥ 10 voxels. *$P < 0.05$ with small volume correction.
1061

1062

1063 **Table 3.** Brain regions with BOLD responses correlating with BDM bids during the bidding phase
1064 (whole-brain analysis with GLM3).

1065

Brain region	Hemisphere	MNI peak coordinates (x,y,z)	peak z-score
vmPFC	/	6, 44, 0	4.10
Dorsal striatum*	R	12, 12, 0	3.53
Insular gyrus/ basal operculum	R	30, 24, -2	5.21
Inferior frontal gyrus, opercular part	R	46, 8, 22	4.76
Occipital gyri	L	-36, -90, -2	4.67
Superior parietal lobule	L	-30, -56, 46	4.14
Superior frontal gyrus	/	2, 26, 42	4.05
Middle frontal gyrus	R	38, 36, 18	3.93
Postcentral gyrus	R	32, -36, 48	3.62
Inferior frontopolar gyrus	R	22, 56, -4	3.53

1066

1067 Cluster P values ($P < 0.05$) with family-wise error correction across the whole brain. Map threshold
1068 $P < 0.005$, extent threshold ≥ 10 voxels. *$P < 0.05$ with small volume correction.

1069

1070

1071
1072
Supplementary Figure 1. Indifference curves (ICs) of 24 participants. In each participant, a total of
15 bundles were obtained by repeatedly comparing variable bundles with the reference bundles (see
Methods). Three ICs are shown by red (high), green (middle) and blue (low) color, with 5 bundles
(indifference points) at each level. The X-axis represents Component A (milkshake with more fat)
and the Y-axis represents Component B (milkshake with more sugar). Lines indicate mean and 95%
confidence intervals of hyperbolic fitted ICs. Error bars represent SEM of the indifference point of
the bundle.

Supplementary Figure 2. Differential activation of striatal regions during value elicitation (ventral striatum) and BDM bidding (dorsal striatum).
(A) Ventral striatum activation during bundle-on phase follows IC scheme (left): significant activation across ICs but not within ICs (GLM1: threshold $p < 0.005$, extent threshold ≥ 10 voxels). Spearman rank ROI analysis (right) showed significant Rho coefficients across the 3 ICs, consistent with the haemodynamic response function (HRF) during bundle-on phase in ventral striatum.
(B) Dorsal striatum activation during bidding phase (GLM3: activation correlated with the amount of BDM bids; threshold $p < 0.005$, extent threshold ≥ 10 voxels). Brain map (left) shows dorsal striatum activity during bidding phase. Spearman rank analysis (right) showed significant Rho coefficient across bids during bidding phase in dorsal striatum.