Mathematical modeling of genetic pest management through female lethality with independently segregating alleles

Michael R. Vella[1,2], Fred Gould[2,3], and Alun L. Lloyd[1,2,4]

[1]**Biomathematics Graduate Program, North Carolina State University, Raleigh, NC 27695, USA**

[2]**Genetic Engineering and Society Center, North Carolina State University, Raleigh, NC 27695, USA**

[3]**Department of Entomology and Plant Pathology, North Carolina State University, Raleigh, NC 27695, USA**

[4]**Department of Mathematics, North Carolina State University, Raleigh, NC 27695, USA**

Abstract

Many novel genetic approaches are under development to combat insect pests, which continue to impose substantial burdens in the forms of transmission of vector-borne disease and the reduction of agricultural yields. One category of genetic pest management aims to suppress or locally eliminate a species through large, repeated releases of fertile males that carry a dominant lethal genetic construct to render all, or exclusively female, offspring unviable. In all strains that have been tested in the field, the genetic components are in a single construct inserted at a unique location in the genome. Other conditional female lethal strains have been developed with two components at independently segregating loci. It has been suggested that males from these strains only be used for a sterile release program as the components are not linked. Here, we use a population genetics model with density-dependence to evaluate the relative effectiveness of female killing approaches based on single- and multi-locus arrangements of the components. We find that, in general, the single-locus arrangement results in slightly faster population suppression, but the two-locus arrangement can eventually cause stronger suppression and cause local extinction with a smaller release size. Based on our results, there is no a priori reason that males carrying two or more independently segregating components need to be sterilized prior to release. In some cases, a fertile release would be more efficient for population suppression.

Introduction

Insect pests remain a burden to human health and agriculture (World Health Organization 2017; Deutsch et al., 2018). Genetic pest management aims to reduce this burden by releasing engineered insects that either introduce a desired trait into a natural population or reduce the size of the population. There have historically been several large area-wide inundative releases of male insects that were rendered sterile by exposure to radiation (Gould and Schliekelman 2004). In these releases, local elimination of the target species was achieved as females increasingly mated with the sterile males rather than the wild-type males with whom they would produce viable offspring. Instead of using radiation to cause sterility, a contemporary alternative is to genetically engineer strains in which the males cause all their offspring or exclusively their daughters to die or to have low fitness (Alphey 2002). Genetically engineered strains in many different species have been cage- or field-tested with promising results (Wise de Valdez et al., 2011; Ant et al., 2012; Harris et al., 2012; Lacroix et al., 2012; Leftwich et al., 2014; Carvalho et al., 2015; Harvey-Samuel et al., 2015; Gorman et al., 2016).

One approach to developing these functionally-sterile strains involves inserting a repressible, dominant lethal gene, which can be active in both sexes or in females only (Heinrich & Scott, 2000; Thomas et al., 2000). Intuitively, modeling studies have found that female-killing (FK, also sometimes referred to as fsRIDL, or female-specific release of insects carrying dominant lethals) can be advantageous over bisex lethality (BS) because it kills females while allowing the transgene to propagate through multiple generations in heterozygote males (Schliekelman & Gould, 2000; Thomas et al., 2000). This would seem especially useful when females but not males transmit pathogens. However, heterozygous males can also serve as a reservoir for wild-type alleles, which can make FK less effective than BS under some conditions (Foster et al., 1988; Gentile et al., 2015). It should be noted that BS strains for mosquito disease vectors typically require sex-sorting because release of females is often considered unacceptable. It is also advantageous to release only males as females do not contribute to genetic suppression and tend to mate with the released males and thus reduce their efficiency (Rendón et al., 2004).

For either the BS or FK approach, in order to rear the transgenic strain in the generations prior to release, it must be possible to inactivate the dominant lethal gene. Often this is achieved through a Tet-off system where tetracycline in the diet represses the activator for a lethal gene (Gossen & Bujard, 1992). For an FK strain, only males are produced from the release generation as the diet would not contain tetracycline. Further, as the offspring of released FK or BS males would feed on a tetracycline-free diet, the lethal gene is turned on and death ensues. The full design involves two components: 1. the tetracycline-repressible transactivator (tTA) with a promoter, and 2. a lethal gene with an enhancer/promoter consisting of multiple tTA binding sites (tetO) and a core promoter. In the initial two-component systems, tTA was expressed in females by using a female-specific promoter (Heinrich & Scott, 2000; Thomas et al., 2000). The second component was a lethal gene (e.g., proapoptotic) driven by tetO enhancer-promoter. Subsequently, a simpler single component system was developed in which tTA acts as both the activator and lethal gene. The construct consists of tTA driven by the tetO enhancer-promoter. In this autoregulated system, high levels of the tTA activator cause lethality, which is possibly due to a general interference in transcription (Gong et al., 2005). FK single component strains include a sex-specifically spliced intron such as from the *transformer* or *doublesex* genes

inserted within the tTA gene (Fu et al., 2007). In these FK strains, only the female tTA transcript
encodes a functional protein. All field trials with transgenic FK or BS strains have been with
single component tTA overexpression strains. More recently, two component FK strains have
been made with an early embryo promoter driving tTA expression and a tTA-regulated lethal
gene that contains a sex-specifically spliced intron (Yan et al., 2019). An advantage of these
strains is that female lethality occurs at the embryo or early larval stages, which produces
considerable savings in larval diet costs in a mass rearing facility.
100
While the two components can be in a single construct (Yan & Scott, 2015), they are
typically at separate independently segregating loci. This is because it can be advantageous to
insert each component independently into different insect strains, identify the most effective
strains, and then bring the components together by crossing. The final transgenic insects would
have the two components located at separate loci (Schetelig & Handler, 2012; Ogaugwu et al.,
2013; Scott, 2014; Schetelig et al., 2016; Yan et al., 2019). It is often suggested that independent
inheritance of the components would require these strains to be sterilized prior to release (e.g.,
statements in Schetelig & Handler, 2012; Ogaugwu et al., 2013; Yan & Scott, 2015). However,
previous theoretical studies of such lethal systems have only modeled the components as being
inserted together on a single locus (Thomas et al., 2000; Schliekelman & Gould, 2000; Alphey et
al., 2011; Gentile et al., 2015). Our model allows for the components to be inserted together at
one-locus or independently on two, unlinked loci, and thus facilitates comparison between the
two approaches.
114
Here we evaluate the effectiveness of 1 and 2-locus FK, along with BS for context, using a
computational model parameterized for the *Aedes aegypti* mosquito that is a vector for several
human infections. We explore the release of strains with killing in either juveniles or adults. We
show that under reasonable assumptions about fitness costs of the insertions, there is not a
substantial difference between the 1 and 2-locus FK approaches, particularly when compared to
the differences between FK and BS. These results demonstrate the release potential of recently
developed 2-locus FK constructs.
122
123   **Methods**
124
Our computational model implements the genetics of FK and BS by separately tracking the
number of individuals in the population of each genotype, denoted by subscript i. For the single-
locus system (Table 1), we let the transgenic allele be represented by K and the wild-type allele
at that locus be represented by k, with a total of three possible genotypes. For the 2-locus system
(Table 2), we let A and B represent the transgenic alleles inserted at two separate loci with wild-
type alleles a and b, respectively, for a total of nine possible diploid genotypes.
131
We assume complete effectiveness of the constructs, so when there is no gene repression via
tetracycline, all individuals bearing the functional BS system and all females with the functional
FK system die (i.e. genotype viability of zero). One copy of K is assumed to be sufficient to
induce lethality in the 1-locus system, and only one copy each of A and B (i.e., tTA and lethal
gene) is required in the 2-locus system. We consider lethality acting at different points in the
lifecycle. In insects that experience strong resource competition during larval stages, having the
transgene-induced mortality occur during or shortly after the pupal stage, instead of during the

egg or larval stages, can yield stronger population suppression. This is because the transgenic
juveniles consume resources and thereby increase wild-type juvenile mortality. We model early
mortality (E) as occurring in the embryo and late mortality (L) as occurring in pupal stages or in
adults before mating, and we assume these differentiate whether the individual contributes
toward density-dependent mortality of all individuals in the population. We let $\gamma_i^{E,S}$ and $\gamma_i^{L,S}$
represent the early (embryonic) and late (adult) expected viabilities for individuals of sex S and
genotype i. Tables 1 and 2 give expected viabilities for individuals with each construct and
genotype.
147
We classify constructs into four different approaches depending on when the dominant lethal
gene is active as in Gentile et al. (2015): early bisex (EBS), late bisex (LBS), early female-killing
(EFK), and late female-killing (LFK). We assume male transgenic homozygotes are released, so
mating with wild-type females will produce offspring that are entirely heterozygous, with a copy
of each transgene. If the construct affects both sexes (BS), none of these offspring will survive to
mate and pass on their genes, making bisex 1-locus and 2-locus equivalent in terms of both
population genetics and population dynamics. Female-specific approaches (FK) allow males to
continue to propagate the transgenes, and thus inheritance differs between 1-locus and 2-locus
approaches. In all, we consider the following six approaches: EBS, LBS, 1-locus EFK (EFK 1),
2-locus EFK (EFK 2), 1-locus LFK (LFK 1), and 2-locus LFK (LFK 2).
158
Separate from the transgenic, toxin-induced lethality, we account for potential fitness costs
caused by the genetic insertion itself. We allow the fitness costs of inserting a novel genetic
element to manifest at an early stage as a reduction in the ability of a zygote to survive beyond
the egg stage, i.e., the fraction of eggs of that genotype which survive and hatch into larvae. We
let the genotype's hatching fitness, w_i^H, equal the probability of successfully entering the larval
stage, with wild-type hatching fitness $w_1^H = 1$. We also allow for transgenic fitness costs to
males in the form of reduced mating competitiveness, w_i^M, as defined below, with wild-type
mating competitiveness $w_1^M = 1$.
167
We assume that the fitness costs are equal for the homozygotes in the 1-locus and 2-locus
systems to facilitate a direct comparison between the two systems. The 2-locus system requires
two separate genetic insertions but is expected to be easier to engineer, which makes equal
fitness costs a reasonable base assumption for the purposes of this work (this assumption is
relaxed in Figure 3 and Figure S1). We let s^H and s^M be the hatching and mating
competitiveness fitness costs, respectively, to the homozygotes KK and AABB, and we allow the
two types of costs to vary independently. For simplicity, we assume the degree of dominance for
the fitness costs, h, is equal for hatching and mating competitiveness. Unless otherwise noted, we
assume costs are additive, with $h = 0.5$, such that each copy of the K allele alone contributes a
fitness cost of $0.5s^x$ for the 1-locus system (x here indicates that the fitness cost can either be
hatching or mating). For the 2-locus system, we allow for unequal fitness costs between each of
the insertions. We let two copies of the A allele contribute a fitness cost of $s^x c_A$, where c_A is the
proportion of the total 2-locus fitness accounted for by the A allele, and one copy of the A allele
contributes a fitness cost of $h \cdot s^x c_A$. A single B allele contributes a fitness cost of $h \cdot s^x (1 -$
$c_A)$, while being homozygous for B induces a cost of $s^x (1 - c_A)$. Resulting fitness expressions
for all genotypes are listed in Tables 1 and 2.
184

We model genotype counts over time using a system of ordinary differential equations
adapted from Robert et al. (2013). We let J_i^M and J_i^F be the number of juvenile (larvae and pupae)
males and females, respectively, of genotype i, and A_i^M and A_i^F be the number of viable adult
male and adult female mosquitoes in the population, respectively, of genotype i. This gives a
maximum of 12 classes of individuals to track (each with different combinations of the 3
genotypes, 2 sexes, and 2 age classes) for the 1-locus system and 36 classes for the 2-locus
system, though lethality from the genetic construct prevents survival of certain classes. For
instance, EBS only has five non-zero classes (wild-type male and female juveniles and adults,
and male adult homozygote transgenic, which are released). We also assume a 50:50 sex ratio
and equal hatching fitness costs between males and females, allowing a further reduction in the
number of unique classes for late acting approaches because $J_i^M = J_i^F$. This results in 7 classes
for LFK 1 (after removing 3 juvenile classes and 2 non-viable adult female classes) and 23
classes for LFK 2 (after removing 9 juvenile classes and 4 non-viable adult female classes).
These dimensionality reductions can be useful when finding analytical solutions, but for
simplicity, we computationally simulated all 12 (1-locus) or 36 (2-locus) equations.

Accounting for fitness costs, adult females produce juveniles of genotype i at rate

$$B_i(t) = w_i^H \sum_m A_m^F(t) \lambda \sum_n P(i|m,n) \frac{w_n^M A_n^M(t)}{\sum_g w_g^M A_g^M(t)} \tag{1}$$

where λ is the per-capita birth rate and $P(i|m,n)$ is the probability of producing juveniles of
genotype i given a mating of a female and male of genotypes m and n, respectively. The fraction
gives the probability that a randomly chosen male adult is of genotype n, weighted by mating
competitiveness. The offspring genotype probabilities are calculated assuming Mendelian
inheritance, and for the 2-locus case, independent segregation of genes at each locus. Because of
the 50:50 sex ratio, half of the hatching juveniles are male and half are female.

Juveniles of each genotype and sex emerge to adulthood at per-capita rate v. We assume
juveniles, adult males, and adult females have per-capita density-independent mortality rates of
μ_J, μ_M, and μ_F, respectively. Juveniles also undergo density-dependent mortality at a per-capita
rate $(\alpha J)^{\beta-1}$, where J is the total number of juveniles. The strength of density dependence is
adjusted by varying β, with higher β resulting in a faster return to equilibrium population size
after a small perturbation. A value of β = 2 gives the logistic model for population dynamics. By
default, we let β = 3 to model an environment which would be more difficult for successful
suppression (e.g., Hibbard et al., 2010). The equilibrium size of an entirely wild-type population
varies with α, and to keep simulations with different values of β comparable, we choose the
value of α so that the equilibrium number of wild-type females remains the same.

We assume a continuous release of homozygote engineered males at (daily) rate $u_i^M =$
$rM_1^*/7$, where r is the weekly release ratio (engineered:wild-type) based on the equilibrium
number of males prior to the release, M_1^*, such that the effective release ratio increases as the
population size decreases. The release genotype is KK for 1-locus ($i = 3$) and AABB for 2-locus
($i = 9$), and because we assume that no females are released, $u_i^F = 0$ for all i.

The resulting system of differential equations is
230

$$\frac{dJ_i^S}{dt} = \frac{1}{2}\gamma_i^{E,S}B_i(t) - J_i^S(\alpha J)^{\beta-1} - \mu_J J_i^S - v J_i^S \tag{2}$$

$$\frac{dA_i^S}{dt} = v\,\gamma_i^{L,S}J_i^S - \mu_S A_i^S + u_i^S, \tag{3}$$

231
233   for $i = 1..N$ and $S = $ F or M. All model parameters are listed in Table 3 and are based on those
234   used for *Ae. aegypti* by Robert et al. (2013). All numerical simulations of the differential
235   equations have initial conditions at the wild-type equilibrium.
236
237   In order to explore the effects of demographic stochasticity and genetic drift, we also run
238   simulations using an analogous continuous time Markov chain model (for details, see Appendix
239   1).
240
241
242   **Results**
243
Each of the FK and BS genetic strategies has the goal of causing the population to decline by
reducing the number of reproductive adult females. The strength of density-dependent mortality
moderates the reductions in population size because stronger density dependence (higher β)
causes the juvenile mortality rates to decrease more quickly as population size decreases from
equilibrium. In a system with strong density dependence, the weekly release ratio (r) must be
larger to achieve the same amount of population suppression as in a system with weak density
dependence. Large r can result in population extinction. In contrast, small r results in a new,
lower equilibrium population density, where the proportion of individuals that die due to bearing
the transgene is not high enough to outweigh the increased survival of juveniles due to decreased
density-dependent mortality in the smaller population.
254
Figure 1 demonstrates the outcome of release for each genetic approach at $r = 1$ under the
deterministic model (see Figure S1 for time series with multiple release ratios). At this release
ratio, approaches that are late-acting (i.e., mortality in pre-mated adults, shown by solid lines)
reduce the number of viable females to a lower number than approaches that are early-acting
(i.e., mortality in the embryonic life stage, shown by dashed lines). BS approaches (red) reduce
the number of females faster and to lower levels than female-specific approaches (blue/green)
with late-acting mortality, but the opposite is true for early-acting mortality. Among FK
approaches, 1-locus reduces the number of females more quickly initially, but 2-locus eventually
suppresses the population slightly more than 1-locus. Overall, this suggests that LBS is most
effective, followed by LFK, EFK, and EBS, with little difference between 1 and 2-locus FK.
Under stronger density dependence (higher values of β), releases result in weaker suppression of
the population, with none of the approaches causing local extinction of the population if β = 4.
267
268

269
270
Figure 1. Effect of transgenic releases on population size over time for various strengths of density
dependence. The number of viable adult females (relative to pre-release equilibrium) over time is plotted
for deterministic simulations with adults for each genetic approach released at a continual weekly release
ratio of 1:1 transgenic males to the pre-release equilibrium wild-type males ($r = 1$). Line type
differentiates embryo (early-acting, dashed line) and adult (late-acting, solid line) mortality, and line color
differentiates bisex (red), 1-locus female-killing (blue), and 2-locus female-killing (green) constructs.
Releases are less effective as strength of density dependence (β) increases across panels from left to right.
Simulations use fitness parameters $s^H = 0.2$, $s^M = 0.1$, and $c_A = 0.55$ and the remaining parameters as listed
in table 3.

280
In general, large release ratios result in extinction (the population goes to an equilibrium size
of zero), and small release ratios result in a suppressed but non-zero equilibrium population size.
For most sets of parameters, there is a critical release ratio, r_c, where the release is large enough
to cause the population to go extinct. For ongoing release at release ratios below r_c, there is a
non-zero stable equilibrium for the number of viable adult females, meaning that release will not
push a wild-type population to extinction. A population of zero is also stable, meaning the
release could protect against re-invasion of the wild-type if continued after population extinction,
but the system will approach the non-zero equilibrium unless starting from very low population
sizes, i.e., it is a bistable system with a low invasion threshold (as shown in Figure S2 with time
series starting from multiple initial conditions). Above r_c, population extinction is the only stable
equilibrium, meaning the release will cause extinction regardless of initial population size. (See
Appendix 2 and Figure S3 for mathematical details on this shift in qualitative behavior).

293
Figure 2 (top) shows how release ratios affect stable equilibria when β = 3, with only the
non-zero stable equilibrium plotted when the system is bistable. Each of the lines is
discontinuous, jumping from a non-zero equilibrium to a zero equilibrium at r_c. The smallest
release size required to cause extinction for each approach is $r_c^{LBS} = 0.86 < r_c^{LFK2} = 2.52 <$
$r_c^{LFK1} = 2.86 < r_c^{EFK2} = 4.74 < r_c^{EFK1} = 5.40 < r_c^{EBS} = 5.97$. This order of effectiveness matches
that of Figure 1 and mirrors the results for single locus constructs in Gentile et al. 2015.
However, when r is smaller than r_c^{LBS} and extinction does not occur, LBS has a higher
equilibrium population size than either of the late FK methods. There is a small relative

difference between the different FK approaches, with similar equilibrium sizes when *r* is below
the critical release ratios and 1-locus FK requiring a release ratio less than 15% larger than 2-
locus FK to cause extinction.
305
In settings where release causes the population to go extinct, we can consider the time it
takes to reach extinction after starting release. Figure 2 (bottom) shows the time it takes for the
number of females to reach less than 0.1% of the pre-control equilibrium in deterministic
simulations, which also reflects the time to extinction in stochastic simulations (see Figure S4).
Once release ratios are high, LBS drops the population under 0.1% of the equilibrium faster than
late FK methods. Also, both of the 1-locus FK approaches are slightly faster than the respective
2-locus FK approaches, though the differences are small for practical purposes. At *r* = 12, for
example, the times are 86 (LBS), 114 (LFK 1), 119 (LFK 2), 133 (EBS), 149 (EFK 1), and 153
(EFK 2) days. The results are similar for β = 2 (see Figure S5).

315
316
Figure 2. Release outcomes across different release ratios. Top: Long-term, stable equilibria for
number of viable adult females (relative to equilibrium) for different *r*, found by simulating the system of
differential equations until at steady-state. The asterisks indicate *r* = 1, for which the equilibrium values

correspond to the middle panel of Figure 1. Each genetic approach exhibits a critical release ratio, r_c,
above which extinction is guaranteed. Multiple lines overlap on the grey line. **Bottom:** Time until the
number of viable adult females is under 0.05% of equilibrium in deterministic simulations for different r.
Color and line type match that of Figure 1. Parameters are $\beta = 3$, $s^H = 0.2$, $s^M = 0.1$, and $c_A = 0.55$, as in
Figure 1.

Overall, FK 1 and FK 2 are quite similar. If either of the A or B alleles in 2-locus FK
becomes fixed in the population, the 2-locus approach becomes nearly identical to 1-locus FK,
where one copy of the unfixed allele causes mortality in females. For example, if all individuals
in the population already have the B allele, only one copy of the A component is additionally
necessary, just as a single copy of the K allele causes mortality. In this case, the long-term
equilibrium can be identical to 1-locus FK, though a fitness cost to the fixed allele decreases the
average fitness of the entire population and makes the population size lower for 2-locus FK than
1-locus FK. Whether one of the 2-locus FK alleles become fixed depends on the fitness costs and
release ratios of the system (see Appendix 3, Figure S6, and Figure S7).

One observation from Figure 2 (top) is that LBS, LFK 2, and LFK 1 cause a similar level of
suppression when the release ratio is near $r = 0.8$. Previous work has suggested that LFK allows
the wild-type allele to propagate in heterozygous males, making it less effective than LBS
(Gentile et al., 2015). This is true at high release ratios, but not at all release ratios. Without
fitness costs, the strategies are equally effective if the number of released males is equal to the
number of wild-type males in the population at equilibrium: heterozygotes carry both wildtype
and transgenic alleles, and therefore the survival of heterozygous males does not affect allele
frequency. At small release ratios, when LBS release results in low transgenic frequency and
thus a large equilibrium population size, survival of heterozygous males would allow the
transgenic allele to propagate further and increase in frequency, explaining why LFK has a lower
equilibrium than LBS in this narrow window of small releases.

The main difference exhibited between FK 1 and FK 2 can also be explained by their
propagation of the transgenic and wild-type alleles. When the components are separated across
two loci, the A and B alleles become unlinked, with some individuals only inheriting one allele
or the other, while having linked components guarantees inheritance of the transgenic allele and
reduces the population size more quickly initially. Eventually, however, the accumulation of
transgenic alleles in the 2-locus system causes production of a higher proportion of unviable
genotypes (Figure 2).

Results over a wide range of fitness parameters highlights the minimal differences between
FK 1 and FK 2. Figure 3 shows the times until 99.95% reduction in viable adult females
(suppression to below 1 adult female when starting from 2000) with a release ratio of $r = 7$,
which is larger than r_c for all approaches in the base parameter set, $\beta = 3$, and varying sets of
fitness parameters. Light colors show parameter sets that resulted in simulations that took longer
to reduce the population size, and dark colors show shorter times, while white areas indicate that
the population was not reduced by 99.95% within 500 days. Within each panel, which represents
a single approach (columns of panels) and degree of dominance, h, (rows of panels), there is a
similar pattern. At low fitness costs (hatching, s^H, varying across the x-axis, and mating
competitiveness, s^M, varying across the y-axis), the release reduces the population below the
threshold, and there is fairly little variation between times (colors) within these regions. As one

or both of the fitness costs increase, there is a margin of longer times (light color) separating the
successful suppression region from the unsuccessful (white) region. As evident from Figure 2,
long times to reduce the population indicate that the release ratio of $r = 7$ is only slightly greater
than r_c for those parameter values, and slightly increased fitness costs increase r_c past 7, so there
is not suppression to 0.1% of equilibrium.
372
There is little difference between FK 1 and FK 2, particularly for LFK, since the release ratio
is much higher than r_c in the region with successful release. Even with different fitness costs, FK
1 and FK 2 do not differ drastically in time until extinction; FK 2 would be substantially less
effective only if the costs become large enough for the release to become too small to cause
extinction (i.e., r becomes smaller than r_c because of the higher costs). Unpacking the differences
between all approaches, for the EBS approach, neither h nor s^H affect the length of time for
reduction; only released males experience costs so the dominance of costs does not affect results,
and all offspring that inherit a transgene are inviable so hatching costs does not affect results. For
EBS, increasing the cost to male mating competitiveness drastically decreases release efficacy
because doing so effectively reduces the release ratio (fewer of the released males successfully
mate and contribute to population reduction), and with no surviving transgenic offspring, the
release ratio is the main contributing factor to EBS efficacy. When $s^H = 1$ and $h = 1$, none of the
offspring survive for the EFK 1 and EFK 2 approaches, making the times shown identical to
EBS. Decreasing these parameters makes a difference for FK approaches because the males are
subject to fitness costs. Dominant fitness costs (bottom row) actually enable successful
population reduction for higher male mating costs. The explanation for this effect relates to the
propagation of wild-type alleles, similar to the previous description: homozygote males that are
being released are affected regardless of h, and when $h > 0$, the fitness cost also reduces the
spread of wild-type alleles in heterozygotes. This effect is also evident in other transgenic
systems, such as two-locus underdominance, where dominant transgenic allele fitness costs
prevent the wild-type from being maintained in the population at small frequencies (Dhole et al.
2018).
395
396

397
Figure 3: Effects of fitness cost variation on release efficacy for weekly release ratio $r = 7$ and $\beta = 3$.
Each column of panels shows the results for a different genetic approach, while degree of dominance, h,
varies across rows of columns. Within each individual panel, the hatching fitness cost, s^H, increases from
0 to 1 along the x-axis, and the cost to male mating competitiveness, s^M, increases from 0 to 1 along the y-
axis. For every point, a deterministic simulation was run with a unique combination of genetic approach
and fitness parameters, and color indicates the number of days until the number of viable adult females is
under 0.05% of equilibrium. Darker colors show faster times, with a minimum time of 73 days, and
lighter colors show slower times up to 500 days. White areas indicate that the number of females did not
fall below the threshold within 500 days. The colored points in the middle row correspond to the times in
Figure 2 (bottom) at $r = 7$.

Discussion

Our main interest in this study was to compare early (E) and late acting (L) female-killing
(FK) engineered into a single locus or across two, unlinked loci. We also examined bisex (BS)
approaches for context. Our overall conclusion was that 2-locus FK (FK 2) behaves similarly to
1-locus FK (FK 1) and does not present any significant disadvantages in its ability to suppress a
population. We generally made the assumption that because of the flexibility of the 2-locus
engineering approach, the total fitness cost for the two insertions would be similar to the single
insertion at 1-locus. If the total cost of the 2-locus approach was less than for the 1-locus
approach, the 2-locus approach would likely be preferred. The reverse also holds. Importantly,
based on our results, there is no a priori reason for genetic engineers to favor a 1-locus.

Assuming equal costs, FK 1 is slightly faster at initial population reduction, but FK 2 can
eventually suppress the population to lower numbers. FK 2 also has a slightly lower critical
release ratio than FK 1, meaning a smaller release size is necessary to guarantee extinction. For
many combinations of fitness costs and release ratios, one of the FK 2 alleles would be driven to
fixation, resulting in a genetic system similar to FK 1. The differences between FK 1 and FK 2

428 are much smaller than between FK and BS approaches. Comparing FK and BS approaches, our
429 results are generally the same as previous work (Gentile et al., 2015). Late acting approaches
430 cause local extinction with a lower release ratio than early acting approaches, with LBS causing
431 local extinction with a lower release ratio than LFK, and EFK causing local extinction with a
432 lower release ratio than EBS.
433
While our modeling results indicate that LBS outperforms the other methods, there are
ecological, economic and social considerations that affect which approach may be best suited in
a given scenario. In our model, parameterized for mosquitoes, density-dependent mortality
during early life-stages was an important factor and caused early-acting approaches to result in
less population reduction than late-acting approaches. In species with little density-dependent
dynamics in juveniles, the difference in effectiveness between early and late-acting would be
minor, though this is not the case for many pest species.
441
Population dynamics alone cannot determine which approach is optimal for a given scenario.
For example, rearing costs are expected to vary between approaches. With EFK, juvenile females
experience mortality before consuming food, whereas EBS, LBS and LFK require rearing of
juveniles of both sexes. Furthermore, BS approaches require sexing to remove females prior to
release, which both costs money and is often difficult to do with complete accuracy. Apart from
costs of rearing juveniles, for most agricultural pests, the juvenile stage causes damage to crops
and livestock. In the first generations of transgenic pest releases, the late acting approaches will
leave more feeding immatures in the environment. This is unlikely to be favored by farmers even
though the overall population could be decreasing rapidly, and EBS could be preferred. Finally,
even if late-acting mortality may be ideal for a given scenario, controlling the timing of mortality
at the intended life stage may not always be feasible, for example due to leaky expression of the
lethal gene.
454
The model used here has several limitations. An important factor that could affect population
genetics is spatial heterogeneity. For example, in a spatial model of FK 2, it would be possible,
particularly in small populations, for different patches to have different transgenic alleles reach
fixation. A spatial model would also be useful to determine if FK 2 has any differences in
resilience to wild-type reinvasion. The details of such a spatial model, including rates of release,
would depend on species. A species-specific model could also implement different forms of
density-dependence, age-structure, and mating parameters. Finally, given its generality, our
model does not account for any potential mechanisms for resistance development. Depending on
the mechanism of lethality, there may be advantages for having both components for lethality
inserted together.
465
466

467 **Acknowledgements**
468
469 This research was funded by the Research Training Group in Mathematical Biology, funded
470 by NSF grant RTG/DMS-1246991 (MRV and ALL), NSF IGERT grant 1068676 (MRV, FG and
471 ALL), NIH grant 1R01AI139085-01 (FG and ALL), and the NC State Drexel Endowment
472 (ALL). We thank M.J. Scott, S. Dhole, B. Hollingsworth, J. Baltzegar, and J. Sudweeks for
473 comments and helpful discussion.

References

Alphey, L. (2002). Re-engineering the sterile insect technique. *Insect biochemistry and molecular biology*, *32*(10), 1243-1247.

Alphey, N., Bonsall, M. B., & Alphey, L. (2011). Modeling resistance to genetic control of insects. *Journal of Theoretical Biology*, *270*(1), 42-55.

Ant, T., Koukidou, M., Rempoulakis, P., Gong, H. F., Economopoulos, A., Vontas, J., & Alphey, L. (2012). Control of the olive fruit fly using genetics-enhanced sterile insect technique. *BMC biology*, *10*(1), 51.

Carvalho, D. O., McKemey, A. R., Garziera, L., Lacroix, R., Donnelly, C. A., Alphey, L., ... & Capurro, M. L. (2015). Suppression of a field population of Aedes aegypti in Brazil by sustained release of transgenic male mosquitoes. *PLoS neglected tropical diseases*, *9*(7).

Deutsch, C. A., Tewksbury, J. J., Tigchelaar, M., Battisti, D. S., Merrill, S. C., Huey, R. B., & Naylor, R. L. (2018). Increase in crop losses to insect pests in a warming climate. *Science*, *361*(6405), 916-919.

Dhole, S., Vella, M. R., Lloyd, A. L., & Gould, F. (2018). Invasion and migration of spatially self-limiting gene drives: A comparative analysis. *Evolutionary applications*, *11*(5), 794-808.

Foster, G. G., Vogt, W. G., Woodburn, T. L., & Smith, P. H. (1988). Computer simulation of genetic control. Comparison of sterile males and field-female killing systems. *Theoretical and Applied Genetics*, *76*(6), 870-879.

Fouque, F., Carinci, R., Gaborit, P., Issaly, J., Bicout, D. J., & Sabatier, P. (2006). *Aedes aegypti* survival and dengue transmission patterns in French Guiana. *Journal of Vector Ecology*, *31*(2), 390-399.

Fu, G., Condon, K. C., Epton, M. J., Gong, P., Jin, L., Condon, G. C., ... & Alphey, L. (2007). Female-specific insect lethality engineered using alternative splicing. *Nature biotechnology*, *25*(3), 353-357.

Gong, P., Epton, M. J., Fu, G., Scaife, S., Hiscox, A., Condon, K. C., ... & Coleman, P. G. (2005). A dominant lethal genetic system for autocidal control of the Mediterranean fruitfly. *Nature biotechnology*, *23*(4), 453-456.

Gentile, J. E., Rund, S. S., & Madey, G. R. (2015). Modelling sterile insect technique to control the population of *Anopheles gambiae*. *Malaria journal*, *14*(1), 92.

Gorman, K., Young, J., Pineda, L., Márquez, R., Sosa, N., Bernal, D., ... & Kaiser, P. (2016). Short-term suppression of *Aedes aegypti* using genetic control does not facilitate Aedes albopictus. *Pest management science*, *72*(3), 618-628.

Gossen, M., & Bujard, H. (1992). Tight control of gene expression in mammalian cells by
tetracycline-responsive promoters. *Proceedings of the National Academy of*
Sciences, *89*(12), 5547-5551.

Gould, F., & Schliekelman, P. (2004). Population genetics of autocidal control and strain
replacement. *Annual Reviews in Entomology*, *49*(1), 193-217.

Harrington, L. C., Edman, J. D., & Scott, T. W. (2001). Why do female *Aedes aegypti* (Diptera:
Culicidae) feed preferentially and frequently on human blood? *Journal of medical*
entomology, *38*(3), 411-422.

Harris, A. F., McKemey, A. R., Nimmo, D., Curtis, Z., Black, I., Morgan, S. A., ... & Collado, A.
(2012). Successful suppression of a field mosquito population by sustained release of
engineered male mosquitoes. *Nature biotechnology*, *30*(9), 828-830.

Harvey-Samuel, T., Morrison, N. I., Walker, A. S., Marubbi, T., Yao, J., Collins, H. L., ... &
Shelton, A. M. (2015). Pest control and resistance management through release of insects
carrying a male-selecting transgene. *BMC biology*, *13*(1), 49.

Hibbard, B. E., Meihls, L. N., Ellersieck, M. R., & Onstad, D. W. (2010). Density-dependent and
density-independent mortality of the western corn rootworm: impact on dose calculations
of rootworm-resistant Bt corn. *Journal of Economic Entomology*, *103*(1), 77-84.

Lacroix, R., McKemey, A. R., Raduan, N., Wee, L. K., Ming, W. H., Ney, T. G., ... &
Angamuthu, C. (2012). Open field release of genetically engineered sterile male Aedes
aegypti in Malaysia. *PloS one*, *7*(8), e42771.

Leftwich, P. T., Koukidou, M., Rempoulakis, P., Gong, H. F., Zacharopoulou, A., Fu, G., ... &
Alphey, L. (2014). Genetic elimination of field-cage populations of Mediterranean fruit
flies. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1792), 20141372.

Muir, L. E., & Kay, B. H. (1998). *Aedes aegypti* survival and dispersal estimated by mark-
release-recapture in northern Australia. *The American journal of tropical medicine and*
hygiene, *58*(3), 277-282.

Ogaugwu, C. E., Schetelig, M. F., & Wimmer, E. A. (2013). Transgenic sexing system for
Ceratitis capitata (Diptera: Tephritidae) based on female-specific embryonic
lethality. *Insect biochemistry and molecular biology*, *43*(1), 1-8.

Rendón, P., McInnis, D., Lance, D., & Stewart, J. (2004). Medfly (Diptera: Tephritidae) genetic
sexing: large-scale field comparison of males-only and bisexual sterile fly releases in
Guatemala. *Journal of Economic Entomology*, *97*(5), 1547-1553.

Robert, M. A., Okamoto, K., Lloyd, A. L., & Gould, F. (2013). A reduce and replace strategy for
suppressing vector-borne diseases: insights from a deterministic model. *PLoS One*, *8*(9),
e73233.

567
568 Rueda, L. M., Patel, K. J., Axtell, R. C., & Stinner, R. E. (1990). Temperature-dependent
569   development and survival rates of *Culex quinquefasciatus* and *Aedes aegypti* (Diptera:
570   Culicidae). *Journal of medical entomology*, *27*(5), 892-898.
571
572 Schetelig, M. F., & Handler, A. M. (2012). A transgenic embryonic sexing system for
573   *Anastrepha suspensa* (Diptera: Tephritidae). *Insect biochemistry and molecular*
574   *biology*, *42*(10), 790-795.
575
576 Schetelig, M. F., Targovska, A., Meza, J. S., Bourtzis, K., & Handler, A. M. (2016).
577   Tetracycline-suppressible female lethality and sterility in the Mexican fruit fly,
578   *Anastrepha ludens*. *Insect molecular biology*, *25*(4), 500-508.
579
580 Schliekelman, P., & Gould, F. (2000). Pest control by the release of insects carrying a female-
581   killing allele on multiple loci. *Journal of Economic Entomology*, *93*(6), 1566-1579.
582
583 Scott, M. J. (2014). Development and evaluation of male-only strains of the Australian sheep
584   blowfly, *Lucilia cuprina*. *BMC genetics*, *15*(2), S3.
585
586 Styer, L. M., Minnick, S. L., Sun, A. K., & Scott, T. W. (2007). Mortality and reproductive
587   dynamics of *Aedes aegypti* (Diptera: Culicidae) fed human blood. *Vector-borne and*
588   *zoonotic diseases*, *7*(1), 86-98.
589
590 Thomas, D. D., Donnelly, C. A., Wood, R. J., & Alphey, L. S. (2000). Insect population control
591   using a dominant, repressible, lethal genetic system. *Science*, *287*(5462), 2474-2476.
592
593 Wise de Valdez, M. R., Nimmo, D., Betz, J., Gong, H. F., James, A. A., Alphey, L., & Black,
594   W.C. (2011). Genetic elimination of dengue vector mosquitoes. *Proceedings of the*
595   *National Academy of Sciences*, *108*(12), 4772-4775.
596
597 World Health Organization. (2017). Global vector control response 2017-2030. *Global vector*
598   *control response 2017-2030.*
599
600 Yan, Y., & Scott, M. J. (2015). A transgenic embryonic sexing system for the Australian sheep
601   blow fly *Lucilia cuprina*. *Scientific reports*, *5*, 16090.
602
603 Yan, Y., Williamson, M. E., Davis, R. J., Andere, A. A., Picard, C. J., & Scott, M. J. (2019).
604   Improved transgenic sexing strains for genetic control of the Australian sheep blow fly
605   *Lucilia cuprina* using embryo-specific gene promoters. *Molecular Genetics and*
606   *Genomics*, 1-12.

i	genotype	$\gamma_i^{a,S}$	w_i^x
1	kk	1	1
2	Kk	0	$1 - h \cdot s^x$
3	KK	0	$1 - s^x$

Table 1: 1-locus genotypes, with associated viabilities and fitnesses. Viability of genotype i, $\gamma_i^{a,S}$, takes the value listed when the approach causes loss of viability in sex S (for female killing, only when $S = F$; for bisex, when $S = F$ or $S = M$) with timing a (for early approaches, when $a = E$; for late approaches, when $a = L$), and are 1 otherwise. Fitnesses w_i^x apply for both hatching ($x = H$) and male mating competitiveness ($x = M$).

i	genotype	$\gamma_i^{a,S}$	w_i^x
1	aabb	1	1
2	aaBb	1	$1 - h \cdot s^x(1 - c_A)$
3	aaBB	1	$1 - s^x(1 - c_A)$
4	Aabb	1	$1 - h \cdot s^x c_A$
5	AaBb	0	$1 - h \cdot s^x$
6	AaBB	0	$1 - h \cdot s^x c_A - s^x(1 - c_A)$
7	AAbb	1	$1 - s^x c_A$
8	AABb	0	$1 - s^x c_A - h \cdot s^x(1 - c_A)$
9	AABB	0	$1 - s^x$

Table 2: 2-locus genotypes, with associated viabilities and fitnesses. Viability of genotype i, $\gamma_i^{a,S}$, takes the value listed when the approach causes loss of viability in sex S (for female killing, only when $S = F$; for bisex, when $S = F$ or $S = M$) with timing a (for early approaches, when $a = E$; for late approaches, when $a = L$), and are 1 otherwise. Fitnesses w_i^x apply for both hatching ($x = H$) and male mating competitiveness ($x = M$).

Parameter	Description	Value/range	Reference
μ_J	Density-independent per-capita juvenile mortality rate	0.03 day^{-1}	Rueda et al. 1990
μ_M	Adult male per-capita mortality rate	0.28 day^{-1}	Muir and Kay 1998, Fouque et al. 2006
μ_F	Adult female per-capita mortality rate	0.10 day^{-1}	Muir and Kay 1998, Fouque et al. 2006
λ	Female per-capita larval production rate	8 day^{-1}	Harrington et al. 2001, Styer et al. 2007
ν	Per-capita rate of emergence to adulthood	0.14 day^{-1}	Muir and Kay 1998
β	Strength of density dependence	2 to 4	
F_1^*	Equilibrium number of wild-type females	2000	
r	Weekly release ratio of transgenic: equilibrium wild-type	0 to 12	
α	Density dependence parameter chosen based on F_1^* and β		
w_i^H, w_i^M	Egg hatching and male mating competitiveness fitnesses of genotype i	See Tables 1 and 2	
s^H, s^M	Egg hatching and male mating competitiveness fitness costs to homozygotes	0 to 1 (0.2 and 0.1 default, respectively)	
h	Fitness cost degree of dominance (percent of s^x incurred in heterozygotes)	0 to 1 (0.5 default)	
c_A	Proportion of s contributed by A (2-locus)	0.5 to 1 (0.55 default)	
$\gamma_i^{E,S}, \gamma_i^{L,S}$	Viabilities of genotype i acting Early (E; embryonic) and late (L; adult) in sex S	See Tables 1 and 2	

639   Table 3: Model parameters.