Mathematical modeling of genetic pest management through female-specific lethality: Is one locus better than two?

Michael R. Vella[1,2], Fred Gould[2,3], and Alun L. Lloyd[1,2,4]

[1]**Biomathematics Graduate Program, North Carolina State University, Raleigh, NC 27695, USA**

[2]**Genetic Engineering and Society Center, North Carolina State University, Raleigh, NC 27695, USA**

[3]**Department of Entomology and Plant Pathology, North Carolina State University, Raleigh, NC 27695, USA**

[4]**Department of Mathematics, North Carolina State University, Raleigh, NC 27695, USA**

Abstract

Many novel genetic approaches are under development to combat insect pests. One genetic strategy aims to suppress or locally eliminate a species through large, repeated releases of genetically engineered strains that render female offspring unviable under field conditions. Strains with this female-killing (FK) characteristic have been developed either with all of the molecular components in a single construct or with the components in two constructs inserted at independently assorting loci. Strains with two constructs are typically considered to be only of value as research tools and for producing solely male offspring in rearing factories which are subsequently sterilized by radiation before release. A concern with the two-construct strains is that once released, the two constructs would become separated and therefore non-functional. The only FK strains that have been released in the field without sterilization are single-construct strains. Here, we use a population genetics model with density dependence to evaluate the relative effectiveness of female killing approaches based on single- and two-construct arrangements. We find that, in general, the single-construct arrangement results in slightly faster population suppression, but the two-construct arrangement can eventually cause stronger suppression and cause local elimination with a smaller release size. Based on our results, there is no a priori reason that males carrying two independently segregating constructs need to be sterilized prior to release. In some cases, a fertile release would be more efficient for population suppression.

Introduction

Insect pests remain a burden to human health and agriculture (World Health Organization 2017; Deutsch et al., 2018). Genetic pest management aims to reduce this burden by releasing engineered insects that either introduce a desired trait into a natural population or reduce the size of the population. There have historically been several large area-wide inundative releases of male insects that were rendered sterile by exposure to radiation (Gould and Schliekelman 2004). In these releases, local elimination of the target species was achieved as females increasingly mated with the sterile males rather than the wild-type males with whom they would produce viable offspring. Instead of using radiation to cause sterility, a contemporary alternative is to genetically engineer strains in which the males cause all their offspring or exclusively their daughters to die or to have low fitness (Alphey 2002). Genetically engineered strains in a number of species have been cage- or field-tested (Wise de Valdez et al., 2011; Ant et al., 2012; Harris et al., 2012; Lacroix et al., 2012; Leftwich et al., 2014; Carvalho et al., 2015; Harvey-Samuel et al., 2015; Gorman et al., 2016).

One approach to developing these functionally-sterile strains involves inserting a repressible, dominant lethal trait, which can be active in both sexes or in females only (Heinrich & Scott, 2000; Thomas et al., 2000). Intuitively, modeling studies have found that female-killing (FK, also sometimes referred to as fsRIDL, or female-specific release of insects carrying dominant lethals) can be advantageous over bisex lethality (BS) because it kills females while allowing the transgene to propagate through multiple generations in heterozygote males (Schliekelman & Gould, 2000; Thomas et al., 2000). This would seem especially useful when females but not males transmit pathogens. However, heterozygous males can also serve as a reservoir for wild-type alleles, which can make FK less effective than BS under some conditions (Foster et al., 1988; Gentile et al., 2015). It should be noted that BS strains for mosquito disease vectors typically require sex-sorting because release of females would be considered unacceptable. It can also be advantageous to release only males as females do not contribute to genetic suppression and tend to mate with the released males and thus reduce their efficiency (Rendón et al., 2004) except in some situations where there is age structuring in the population (Huang et al., 2009).

For either the BS or FK approach, in order to rear the transgenic strain in the generations prior to release, it must be possible to inactivate the dominant lethal gene. Often this is achieved through a Tet-off system where tetracycline in the diet represses the activator for a lethal gene (Gossen & Bujard, 1992). For an FK strain, the release generation is reared on a diet not containing tetracycline. This results in only males surviving. Further, as the offspring of released FK or BS males would feed on a tetracycline-free diet under field conditions, the lethal gene is turned on and death ensues.

The full molecular design involves two molecular components: 1. the tetracycline-repressible transactivator (tTA) with a promoter, and 2. a lethal gene with an enhancer/promoter consisting of multiple tTA binding sites (tetO) and a core promoter. In the initial two-component systems, tTA was expressed in females by using a female-specific promoter (Heinrich & Scott, 2000; Thomas et al., 2000). The second component was a lethal gene (e.g., proapoptotic) driven by a tetO enhancer-promoter. The two molecular components were built in separate constructs that were inserted independently. Subsequently, a simpler, two-component system was developed in

which tTA acts as both the activator and lethal gene. Here, a single construct includes a tTA
coding sequence driven by a tetO enhancer-promoter. In this autoregulated system, high levels of
the tTA activator cause lethality in late stage larvae or in pupae. The mortality is possibly due to
a general interference in transcription (Gong et al., 2005). FK single-construct strains have
included a sex-specifically spliced intron from the *transformer* or *doublesex* genes inserted
within the tTA gene (Fu et al., 2007). In these FK strains, only the female tTA transcript encodes
a functional protein. A different, single-construct approach for FK with *Aedes aegypti* and *Aedes*
albopictus uses a female-specific indirect flight muscle promoter from the Actin-4 gene (Fu et
al., 2010; Labbé et al., 2012). All field trials with transgenic FK or BS strains have been with
single-construct strains.

More recently, two-construct FK strains have been made with an early embryo promoter
driving tTA expression and a tTA-regulated lethal gene that contains a sex-specifically spliced
intron (Yan et al., 2019). An advantage of these strains is that female lethality occurs at the
embryo or early larval stages, which produces considerable savings in larval diet costs in a mass
rearing facility. Although it should be possible to develop any two-component systems as a
single construct (Yan & Scott, 2015), they are typically developed as independently-segregating
constructs. This is because it can be advantageous to insert each component independently into
different insect strains, identify the most effective strains, and then produce individuals bearing
both components by crossing. The final transgenic insects have the two components located at
two, separate loci (Schetelig & Handler, 2012; Ogaugwu et al., 2013; Scott, 2014; Schetelig et
al., 2016; Yan et al., 2019).

FK strains with two constructs are generally thought of as useful research tools with potential
to be used in rearing facilities so that the final generation before release would only produce
males (e.g. Schetelig & Handler, 2012; Ogaugwu et al., 2013; Yan & Scott, 2015). It has been
suggested that independent inheritance of the components would cause a breakdown in the
female killing in the second generation after release (Ogaugwu et al., 2013; Yan & Scott, 2015).
However, previous theoretical studies of FK systems have only modeled the components as
being inserted together on a single locus (Thomas et al., 2000; Schliekelman & Gould, 2000;
Alphey et al., 2011; Gentile et al., 2015).

Here we evaluate the effectiveness of 1 and 2-locus FK, along with BS for comparison. We
use a computational model parameterized for the *Aedes aegypti* mosquito that is a vector for
several human pathogens. We explore the release of strains with killing in either juveniles or
adults. We show that under reasonable assumptions about fitness costs of the insertions, there is
not a substantial difference between the 1 and 2-locus FK approaches, particularly when
compared to the differences between FK and BS. These results demonstrate the release potential
of recently developed 2-locus FK constructs.

Methods

Our computational model implements the genetics of FK and BS by separately tracking the
number of individuals in the population of each genotype, with genotype denoted by subscript *i*.
For the single-locus system (Table 1), we let the transgenic allele be represented by K and the
wild-type allele at that locus be represented by k, with a total of *N=3* possible genotypes. For the

2-locus system (Table 2), we let A and B represent the transgenic alleles (i.e., tTA and lethal
gene) inserted at two separate loci with wild-type alleles a and b, respectively, for a total of $N=9$
possible diploid genotypes.
142

i	genotype	$\gamma_i^{a,S}$	w_i^x
1	kk	1	1
2	Kk	0	$1 - h \cdot s^x$
3	KK	0	$1 - s^x$

143   Table 1: 1-locus genotypes, with associated viabilities and fitnesses. Viability of genotype i, $\gamma_i^{a,S}$,
144   takes the value listed when the approach causes loss of viability in sex S (for female killing, only
145   when $S = F$; for bisex, when $S = F$ or $S = M$) with timing a (for early approaches, when $a = E$;
146   for late approaches, when $a = L$), and are 1 otherwise. Fitnesses w_i^x apply for both hatching ($x =$
147   H) and male mating competitiveness ($x = $ M).
148
149

i	genotype	$\gamma_i^{a,S}$	w_i^x
1	aabb	1	1
2	aaBb	1	$1 - h \cdot s^x(1 - c_A)$
3	aaBB	1	$1 - s^x(1 - c_A)$
4	Aabb	1	$1 - h \cdot s^x c_A$
5	AaBb	0	$1 - h \cdot s^x$
6	AaBB	0	$1 - h \cdot s^x c_A - s^x(1 - c_A)$
7	AAbb	1	$1 - s^x c_A$
8	AABb	0	$1 - s^x c_A - h \cdot s^x(1 - c_A)$
9	AABB	0	$1 - s^x$

150   Table 2: 2-locus genotypes, with associated viabilities and fitnesses. Viability of genotype i, $\gamma_i^{a,S}$,
151   takes the value listed when the approach causes loss of viability in sex S (for female killing, only
152   when $S = F$; for bisex, when $S = F$ or $S = M$) with timing a (for early approaches, when $a = E$;
153   for late approaches, when $a = L$), and are 1 otherwise. Fitnesses w_i^x apply for both hatching ($x =$
154   H) and male mating competitiveness ($x = $ M).
155
We assume complete effectiveness of the constructs, so when there is no gene repression via
tetracycline, all individuals bearing the functional BS system and all females with the functional
FK system die (i.e. genotype viability of zero). One copy of K is assumed to be sufficient to
induce lethality in the 1-locus system, and only one copy each of A and B is required in the 2-
locus system. We consider lethality acting at different points in the lifecycle. In insects that
experience strong resource competition during larval stages, having the transgene-induced
mortality occur during or shortly after the pupal stage, instead of during the egg or larval stages,
can yield stronger population suppression. This is because the transgenic juveniles consume
resources and thereby increase wild-type juvenile mortality. We model early mortality (E) as

occurring in the embryo and late mortality (L) as occurring in pupal stages or in adults before
mating, and we assume these differentiate whether the individual contributes toward density-
dependent mortality of all individuals in the population. We let $\gamma_i^{E,S}$ and $\gamma_i^{L,S}$ represent the early
(embryonic) and late (adult) expected viabilities for individuals of sex S and genotype i. Tables 1
and 2 give expected viabilities for individuals with each construct and genotype.
170
We classify constructs into four different approaches depending on when the dominant lethal
gene is active as in Gentile et al. (2015): early bisex (EBS), late bisex (LBS), early female-killing
(EFK), and late female-killing (LFK). We assume male transgenic homozygotes are released, so
mating with wild-type females will produce offspring that are entirely heterozygous, with a copy
of each transgene. If the construct(s) affects both sexes (BS), none of these offspring will survive
to mate and pass on their genes, making bisex 1-locus and 2-locus equivalent in terms of both
population genetics and population dynamics. Female-specific approaches (FK) allow males to
continue to propagate the transgenes, and thus inheritance differs between 1-locus and 2-locus
approaches. In all, we consider the following six approaches: EBS, LBS, 1-locus EFK (EFK 1),
2-locus EFK (EFK 2), 1-locus LFK (LFK 1), and 2-locus LFK (LFK 2).
181
Separate from the transgenic, toxin-induced lethality, we account for potential fitness costs
caused by the genetic insertion itself. We allow the fitness costs of inserting a novel genetic
element to manifest at an early stage as a reduction in the ability of a zygote to survive beyond
the egg stage, i.e., the fraction of eggs of that genotype which survive and hatch into larvae. We
let the genotype's hatching fitness, w_i^H, equal the probability of successfully entering the larval
stage, with wild-type hatching fitness $w_1^H = 1$. We also allow for transgenic fitness costs to
males in the form of reduced mating competitiveness, w_i^M, as defined below, with wild-type
mating competitiveness $w_1^M = 1$.
190
We generally assume that the fitness costs are equal for the homozygotes in the 1-locus and
2-locus systems to facilitate a direct comparison between the two systems. The 2-locus system
requires two separate genetic insertions but is expected to be easier to engineer, which makes
equal fitness costs a reasonable base assumption for the purposes of this work (this assumption is
relaxed in Figure 3 and Figure S1). We let s^H and s^M be the hatching and mating
competitiveness fitness costs, respectively, to the homozygotes KK and AABB, and we allow the
two types of costs to vary independently. For simplicity, we assume the degree of dominance for
the fitness costs, h, is equal for hatching and mating competitiveness. Unless otherwise noted, we
assume costs are additive, with $h = 0.5$, such that each copy of the K allele alone contributes a
fitness cost of $0.5s^x$ for the 1-locus system (x here indicates that the fitness cost can either be
hatching or mating). For the 2-locus system, we allow for unequal fitness costs between each of
the insertions. We let two copies of the A allele contribute a fitness cost of $s^x c_A$, where c_A is the
proportion of the total 2-locus fitness cost accounted for by the A allele, and one copy of the A
allele contributes a fitness cost of $h \cdot s^x c_A$. A single B allele contributes a fitness cost of $h \cdot$
$s^x(1 - c_A)$, while being homozygous for B induces a cost of $s^x(1 - c_A)$. Resulting fitness
expressions for all genotypes are listed in Tables 1 and 2.
207
We model genotype counts over time using a system of ordinary differential equations
adapted from Robert et al. (2013). We let $J_i^M(t)$ and $J_i^F(t)$ be the number of juvenile (larvae and
pupae) males and females, respectively, of genotype i at time t, and $A_i^M(t)$ and $A_i^F(t)$ be the

number of viable adult male and adult female mosquitoes in the population, respectively, of
genotype i at time t. This gives a maximum of 12 classes of individuals to track (each with
different combinations of the 3 genotypes, 2 sexes, and 2 age classes) for the 1-locus system and
36 classes for the 2-locus system, though lethality from the genetic construct prevents survival of
certain classes. For instance, EBS only has five non-zero classes (wild-type male and female
juveniles and adults, and male adult homozygote transgenic, which are released). We also
assume a 50:50 sex ratio and equal hatching fitness costs between males and females, allowing a
further reduction in the number of unique classes for late acting approaches because $J_i^M(t) =$
$J_i^F(t)$ for all i and t. This results in 7 classes for LFK 1 (after removing 3 juvenile classes and 2
non-viable adult female classes) and 23 classes for LFK 2 (after removing 9 juvenile classes and
4 non-viable adult female classes). These dimensionality reductions can be useful when finding
analytical solutions, but for simplicity, we computationally simulated all 12 (1-locus) or 36 (2-
locus) equations.
224
225   Accounting for fitness costs, adult females produce juveniles of genotype i at time t at rate
226

227
$$B_i(t) = w_i^H \sum_m A_m^F(t) \lambda \sum_n P(i|m,n) \frac{w_n^M A_n^M(t)}{\sum_g w_g^M A_g^M(t)} \qquad (1)$$

228
where λ is the per-capita birth rate and $P(i|m,n)$ is the probability that a juvenile produced from
a mating between a female and male of genotypes m and n, respectively, will be of genotype i.
The fraction gives the probability that a randomly chosen male adult is of genotype n, weighted
by mating competitiveness w_n^M. The offspring genotype probabilities are calculated assuming
Mendelian inheritance, and for the 2-locus case, independent segregation of genes at each locus.
Because of the 50:50 sex ratio, half of the hatching juveniles are male and half are female.
235
Juveniles of each genotype and sex emerge to adulthood at per-capita rate v. We assume
juveniles, adult males, and adult females have per-capita density-independent mortality rates of
μ_J, μ_M, and μ_F, respectively. Juveniles also undergo density-dependent mortality at a per-capita
rate $(\alpha J)^{\beta-1}$, where J is the total number of juveniles. The strength of density dependence is
adjusted by varying β, with higher β resulting in a faster return to equilibrium population size
after a small perturbation. A value of β = 2 gives the logistic model for population dynamics. By
default, we let β = 3 to model an environment which would be more difficult for successful
suppression (e.g., Hibbard et al., 2010). The equilibrium size of an entirely wild-type population
varies with α, and to keep simulations with different values of β comparable, we choose the
value of α so that the equilibrium number of wild-type females remains the same.
246
We assume a continuous release of homozygote engineered males at (daily) rate
$u_i^M = r \cdot A_1^M(t = 0)/7$, where r is the weekly release ratio (engineered:wild-type) based on the
equilibrium number of males prior to the release. By maintaining a constant number of released
males, the effective release ratio increases as the population size decreases. The release genotype
is KK for 1-locus ($i = 3$) and AABB for 2-locus ($i = 9$), and because we assume that no females
are released, $u_i^F = 0$ for all i.
253

254   The resulting system of differential equations (with time dependence of $J_i^S(t)$, $A_i^S(t)$, and
255 $B_i(t)$ omitted for simplicity of notation) is
256

$$\frac{dJ_i^S}{dt} = \frac{1}{2}\gamma_i^{E,S}B_i - J_i^S(\alpha J)^{\beta-1} - \mu_J J_i^S - v J_i^S \tag{2}$$

$$\frac{dA_i^S}{dt} = v\,\gamma_i^{L,S}J_i^S - \mu_S A_i^S + u_i^S, \tag{3}$$

257
259 for $i = 1..N$ and S = F or M. All model parameters are listed in Table 3 and are based on those
260 used for *Ae. aegypti* by Robert et al. (2013). All numerical simulations of the differential
261 equations have initial conditions at the wild-type equilibrium.
262
263   In order to explore the effects of demographic stochasticity and genetic drift, we also run
264 simulations using an analogous continuous time Markov chain model (for details, see Appendix
265 1).
266

Parameter	Description	Value/range	Reference
μ_J	Density-independent per-capita juvenile mortality rate	0.03 day^{-1}	Rueda et al. 1990
μ_M	Adult male per-capita mortality rate	0.28 day^{-1}	Muir and Kay 1998, Fouque et al. 2006
μ_F	Adult female per-capita mortality rate	0.10 day^{-1}	Muir and Kay 1998, Fouque et al. 2006
λ	Female per-capita larval production rate	8 day^{-1}	Harrington et al. 2001, Styer et al. 2007
v	Per-capita rate of emergence to adulthood	0.14 day^{-1}	Muir and Kay 1998
β	Strength of density dependence	2 to 4	
$A_1^F(0)$	Equilibrium number of wild-type females	2000	
r	Weekly release ratio of transgenic: equilibrium wild-type males	0 to 12	
α	Density dependence parameter chosen based on β and $A_1^F(0)$		
w_i^H, w_i^M	Egg hatching and male mating competitiveness fitnesses of genotype i	See Tables 1 and 2	
s^H, s^M	Egg hatching and male mating competitiveness fitness costs to homozygotes	0 to 1 (0.2 and 0.1 default, respectively)	
h	Fitness cost degree of dominance (percent of s^x incurred in heterozygotes)	0 to 1 (0.5 default)	
c_A	Proportion of s contributed by A (2-locus)	0.5 to 1 (0.55 default)	
$\gamma_i^{E,S}, \gamma_i^{L,S}$	Viabilities of genotype i for early (E; embryonic) and late (L; adult) approaches in sex S	See Tables 1 and 2	

267 Table 3: Model parameters.

Results

Each of the FK and BS genetic strategies has the goal of causing the population to decline by reducing the number of reproductive adult females. The strength of density-dependent mortality moderates the reductions in population size because stronger density dependence (higher β) causes the juvenile mortality rates to decrease more quickly as population size decreases from equilibrium. In a system with strong density dependence, the weekly release ratio (r) must be larger to achieve the same amount of population suppression as in a system with weak density dependence. Large r can result in target population extinction. In contrast, small r results in a new, lower equilibrium population density, where the proportion of individuals that die due to bearing the transgene is not high enough to outweigh the increased survival of juveniles due to decreased density-dependent mortality in the smaller population.

Figure 1 demonstrates the outcome of release for each genetic approach at $r = 1$ under the deterministic model (see Figure S1 for time series with multiple release ratios). At this release ratio, approaches that are late-acting (i.e., mortality in pre-mated adults, shown by solid lines) reduce the number of viable females to a lower number than approaches that are early-acting (i.e., mortality in the embryonic life stage, shown by dashed lines). BS approaches (red) reduce the number of females faster and to lower levels than female-specific approaches (blue/green) with late-acting mortality, but the opposite is true for early-acting mortality. Among FK approaches, 1-locus reduces the number of females more quickly initially, but 2-locus eventually suppresses the population slightly more than 1-locus. Overall, this suggests that LBS is most effective, followed by LFK, EFK, and EBS, with little difference between 1 and 2-locus FK. Under stronger density dependence (higher values of β), releases result in weaker suppression of the population, with none of the approaches causing extinction of the population if β = 4.

Figure 1. Effect of transgenic releases on population size over time for various strengths of density
dependence. The number of viable adult females (relative to pre-release equilibrium) over time is plotted
for deterministic simulations with adults for each genetic approach released at a continual weekly release
ratio of 1:1 transgenic males to the pre-release equilibrium wild-type males ($r = 1$). Line type
differentiates embryo (early-acting, dashed line) and adult (late-acting, solid line) mortality, and line color
differentiates bisex (red), 1-locus female-killing (blue), and 2-locus female-killing (green) constructs.
Releases are less effective as strength of density dependence (β) increases across panels from left to right.
Simulations use fitness parameters $s^H = 0.2$, $s^M = 0.1$, and $c_A = 0.55$ and the remaining parameters as listed
in Table 3.

In general, large release ratios result in extinction (the population goes to an equilibrium size
of zero), and small release ratios result in a suppressed but non-zero equilibrium population size.
For most sets of parameters, there is a critical release ratio, r_c, above which the release is large
enough to cause the population to go extinct. With such a large number of released males,
population extinction is the only stable equilibrium, meaning the release will cause extinction
regardless of initial population size. For ongoing release at release ratios below r_c, there is a non-
zero stable equilibrium for the number of viable adult females, meaning that release will not push
a wild-type population to extinction. A population size of zero is also stable, meaning the release
could protect against re-invasion of the wild-type if continued after population extinction, but the
system will approach the non-zero equilibrium unless starting from very low population sizes,
i.e., it is a bistable system with a low invasion threshold (as shown in Figure S2 with time series
starting from multiple initial conditions). Mathematical details on the shift in qualitative behavior
at r_c and analysis using Mathematica (Wolfram Research, 2019) can be found in Appendix 2 and
Figure S3.

Figure 2A shows how release ratios affect stable equilibria when β = 3, with only the non-
zero stable equilibrium plotted when the system is bistable. Each of the lines is discontinuous,
jumping from a non-zero equilibrium to a zero equilibrium at r_c. The smallest release size
required to cause extinction for each approach is $r_c^{LBS} = 0.86 < r_c^{LFK2} = 2.52 < r_c^{LFK1} = 2.86 <$
$r_c^{EFK2} = 4.74 < r_c^{EFK1} = 5.40 < r_c^{EBS} = 5.97$. This order of effectiveness matches that of Figure 1
and mirrors the results for single locus constructs in Gentile et al. 2015. However, when r is
smaller than r_c^{LBS} and extinction does not occur, LBS has a higher equilibrium population size
than either of the late FK methods. There is a small relative difference between the different FK
approaches, with similar equilibrium sizes when r is below the critical release ratios and 1-locus
FK requiring a release ratio less than 15% larger than 2-locus FK to cause extinction.

In settings where release causes the population to go extinct, we can consider the time it
takes to reach extinction after starting release. Figure 2B shows the time it takes for the number
of females to reach less than 0.05% of the pre-control equilibrium (suppression to below 1 adult
female when starting from a pre-release equilibrium of 2000) in deterministic simulations, which
also reflects the time to extinction in stochastic simulations (see Figure S4). Once release ratios
are high, LBS drops the population under 0.05% of the equilibrium faster than late FK methods.
Also, both of the 1-locus FK approaches are slightly faster than the respective 2-locus FK
approaches, though the differences are small for practical purposes. At $r = 12$, for example, the
times are 86 (LBS), 114 (LFK 1), 119 (LFK 2), 133 (EBS), 149 (EFK 1), and 153 (EFK 2) days.
The results are similar for β = 2 (see Figure S5).

Figure 2. Release outcomes across different release ratios. A: Long-term, stable equilibria for number of viable adult females (relative to pre-release equilibrium) for different r, found by simulating the system of differential equations until at steady-state. The asterisks indicate $r = 1$, for which the equilibrium values correspond to the middle panel of Figure 1. Each genetic approach exhibits a critical release ratio, r_c, above which extinction is guaranteed. Above $r = 2.52$, two or more approaches lead to extinction of the population and hence have equilibria at zero: this is indicated using a grey line. **B:** Time until the number of viable adult females is under 0.05% of equilibrium in deterministic simulations for different r. Color and line type match that of Figure 1. Parameters are $\beta = 3$, $s^H = 0.2$, $s^M = 0.1$, and $c_A = 0.55$, as in Figure 1.

Overall, FK 1 and FK 2 are quite similar. If either of the A or B alleles in 2-locus FK becomes fixed in the population, the 2-locus approach becomes nearly identical to 1-locus FK, where one copy of the unfixed allele causes mortality in females. For example, if all individuals in the population already have the B allele, only one copy of the A component is additionally necessary, just as a single copy of the K allele causes mortality. In this case, the long-term equilibrium can be identical to 1-locus FK, though a fitness cost to the fixed allele decreases the average fitness of the entire population and makes the population size lower for 2-locus FK than 1-locus FK. Whether one of the 2-locus FK alleles become fixed depends on the fitness costs and release ratios of the system (see Appendix 3, Figure S6, and Figure S7).

An observation from Figure 2A is that LBS, LFK 2, and LFK 1 cause a similar level of suppression when the release ratio is near $r = 0.8$. Previous work has suggested that LFK allows the wild-type allele to propagate in heterozygous males, making it less effective than LBS

(Gentile et al., 2015). This is true at high release ratios, but not at all release ratios. Without
fitness costs, the strategies are equally effective if the number of released males is equal to the
number of wild-type males in the population at equilibrium: heterozygotes carry both wildtype
and transgenic alleles, and therefore the survival of heterozygous males does not affect allele
frequency. At small release ratios, when LBS release results in low transgenic frequency and
thus a large equilibrium population size, survival of heterozygous males would allow the
transgenic allele to propagate further and increase in frequency, explaining why LFK has a lower
equilibrium than LBS in this narrow window of small releases.
373
The main difference exhibited between FK 1 and FK 2 can also be explained by their
propagation of the transgenic and wild-type alleles. When the components are separated across
two loci, the A and B alleles become unlinked, with some individuals only inheriting one allele
or the other, while having linked components guarantees inheritance of the transgenic allele and
reduces the population size more quickly initially. Eventually, however, the accumulation of
transgenic alleles in the 2-locus system causes production of a higher proportion of unviable
genotypes and greater population suppression (Figure 2).
381
Results over a wide range of fitness parameters highlights the minimal differences between
FK 1 and FK 2. Figure 3 shows the times until 99.95% reduction in viable adult females with a
release ratio of $r = 7$, which is larger than r_c for all approaches in the base parameter set, $\beta = 3$,
and varying sets of fitness parameters. Light colors show parameter sets that resulted in
simulations that took longer to reduce the population size, and dark colors show shorter times,
while white areas indicate that the population was not reduced by 99.95% within 500 days.
Within each panel, which represents a single approach (columns of panels) and degree of
dominance, h, (rows of panels), there is a similar pattern. At low fitness costs (hatching, s^H,
varying across the x-axis, and mating competitiveness, s^M, varying across the y-axis), the release
reduces the population below the threshold, and there is fairly little variation between times
(colors) within these regions. As one or both of the fitness costs increase, there is a margin of
longer times (light color) separating the successful suppression region from the unsuccessful
(white) region. As evident from Figure 2, long times to reduce the population indicate that the
release ratio of $r = 7$ is only slightly greater than r_c for those parameter values, and a system with
increased fitness costs have $r_c > 7$, so there is not suppression to 0.05% of equilibrium.
397
There is little difference between FK 1 and FK 2, particularly for LFK, since the release ratio
is much higher than r_c in most of the region with successful release. Even with different fitness
costs, FK 1 and FK 2 do not differ drastically in time until extinction; FK 2 would be
substantially less effective only if the costs become large enough for the release to become too
small to cause extinction (i.e., r becomes smaller than r_c because of the higher costs). Unpacking
the differences between all approaches, for the EBS approach, neither h nor s^H affect the length
of time for reduction; only released males experience costs so the degree of dominance does not
affect results, and all offspring that inherit a transgene are inviable so hatching costs do not affect
results. For EBS, increasing the cost to male mating competitiveness drastically decreases release
efficacy because doing so effectively reduces the release ratio (fewer of the released males
successfully mate and contribute to population reduction), and with no surviving transgenic
offspring, the release ratio is the main contributing factor to EBS efficacy. When $s^H = 1$ and $h =$
1, none of the offspring survive for the EFK 1 and EFK 2 approaches, making the times shown

identical to EBS. Decreasing these parameters makes a difference for FK approaches because the
males are subject to fitness costs. Dominant fitness costs (bottom row) actually enable successful
population reduction for higher male mating costs. The explanation for this effect relates to the
propagation of wild-type alleles, similar to the previous description: homozygote males that are
being released are affected regardless of h, and when $h > 0$, the fitness cost also reduces the
spread of wild-type alleles in heterozygotes. This effect is also evident in other transgenic
systems, such as two-locus underdominance, where dominant transgenic allele fitness costs
prevent the wild-type from being maintained in the population at small frequencies (Dhole et al.
2018).
420

421
Figure 3: Effects of fitness cost variation on release efficacy for weekly release ratio $r = 7$ and $\beta = 3$.
Each column of panels shows the results for a different genetic approach, while each row of panels
depicts a different degree of dominance, h. Within each individual panel, the hatching fitness cost, s^H,
increases from 0 to 1 along the x-axis, and the cost to male mating competitiveness, s^M, increases from 0
to 1 along the y-axis. For every point, a deterministic simulation was run with a unique combination of
genetic approach and fitness parameters, and color indicates the number of days until the number of
viable adult females is under 0.05% of equilibrium. Darker colors show faster times, with a minimum
time of 73 days, and lighter colors show slower times up to 500 days. White areas indicate that the
number of females did not fall below the threshold within 500 days. The colored points in the middle row
correspond to the times in Figure 2B at $r = 7$.

432
433
434
435 **Discussion**
436
The recent literature on FK systems makes the assumption that strains built with
constructs inserted at two independent loci will not be as useful for field releases as those built
with a single construct. The assumption is that the two constructs will separate from each other
in the second generation after a release and will become non-functional. Our modeling results
demonstrate that a 2-locus FK (FK 2) should behave similarly to a 1-locus FK (FK 1) and would
not present any significant disadvantages in its ability to suppress a population. We generally

made the assumption that because of the flexibility of the 2-locus engineering approach, the total
fitness cost for the two insertions would be similar to the single insertion at 1-locus. If the total
cost of the 2-locus approach was less than for the 1-locus approach, the 2-locus approach would
likely be preferred. The reverse also holds. Importantly, based on our results, there is no a priori,
general reason for genetic engineers to favor a 1-locus system. The choice will likely depend on
specific biological and genetic characteristics of the target species.
449
Assuming equal costs, FK 1 is slightly faster at initial population reduction, but FK 2 can
eventually suppress the population to lower numbers. FK 2 also has a slightly lower critical
release ratio than FK 1, meaning a smaller release size is necessary to guarantee extinction. For
many combinations of fitness costs and release ratios, one of the FK 2 alleles would be driven to
fixation, resulting in a genetic system similar to FK 1. The differences between FK 1 and FK 2
are much smaller than between FK and BS approaches. Comparing FK and BS approaches, our
results are generally similar to previous work (Gentile et al., 2015). Late acting approaches cause
extinction with a lower release ratio than early acting approaches, with LBS causing extinction
with a lower release ratio than LFK, and EFK causing extinction with a lower release ratio than
EBS.
460
While our modeling results indicate that LBS outperforms the other methods, there are other
considerations that affect which approach may be best suited for a given scenario. In our model,
parameterized for mosquitoes, density-dependent mortality during early life-stages was an
important factor and caused early-acting approaches to result in less population reduction than
late-acting approaches. In species with little density-dependent dynamics in juveniles, the
difference in effectiveness between early and late-acting would be minor, though this is not the
case for many pest species.
468
Beyond population dynamics, there are economic and social factors that differ between
approaches. For some systems, it will be necessary to engineer constructs into lab strains and
then backcross the construct or constructs into a strain that have a genetic makeup similar to the
targeted population. In general, it should be easier to do the backcrossing with a one-locus
system. Rearing costs are also expected to vary between approaches. With EFK, juvenile females
experience mortality before consuming food, whereas EBS, LBS and LFK require rearing of
juveniles of both sexes. Furthermore, BS approaches require sexing to remove females prior to
release, which increases the total rearing costs and is often difficult to do with complete
accuracy. When releasing a species that is a disease vector, sexing accuracy is meaningful from a
social perspective as release of females could contribute to disease transmission.
479
Apart from engineering and rearing, for most agricultural pests, the juvenile stages of males
and females cause damage to crops and livestock. In the first generations of transgenic pest
releases, the late acting approaches will leave more feeding immatures in the environment. For
LFK and EFK, male immatures will still cause damage. This is unlikely to be favored by farmers
even though the overall population could be decreasing rapidly, and EBS could be preferred.
Finally, even if late-acting mortality may be ideal for a given scenario, controlling the timing of
mortality at the intended life stage may not always be feasible, for example due to leaky
expression of the lethal gene.
488

The model used here has several limitations. An important factor that could affect population
genetics is spatial heterogeneity. For example, in a spatial model of FK 2, it would be possible,
particularly in small populations, for different patches to have different transgenic alleles reach
fixation. A spatial model would also be useful to determine if FK 2 has any differences in
resilience to wild-type reinvasion. The details of such a spatial model, including rates of release,
would depend on species. A species-specific model could also implement different forms of
density-dependence, age-structure, and mating parameters. Finally, given its generality, our
model does not account for any potential mechanisms for resistance development. Depending on
the mechanism of lethality, there may be advantages for having both components for lethality
inserted together. While these areas require further investigation, our results indicate that overall,
there is little difference in the pest population suppression efficacies of 1-locus FK and 2-locus
FK.

Acknowledgements

505   This research was funded by the Research Training Group in Mathematical Biology, funded
506 by NSF grant RTG/DMS-1246991 (MRV and ALL), NSF IGERT grant 1068676 (MRV, FG and
507 ALL), NIH grant 1R01AI139085-01 (FG and ALL), and the NC State Drexel Endowment
508 (ALL). We thank M.J. Scott, S. Dhole, B. Hollingsworth, J. Baltzegar, and J. Sudweeks for
509 comments and helpful discussion.

References

Alphey, L. (2002). Re-engineering the sterile insect technique. *Insect biochemistry and molecular biology*, *32*(10), 1243-1247.

Alphey, N., Bonsall, M. B., & Alphey, L. (2011). Modeling resistance to genetic control of insects. *Journal of Theoretical Biology*, *270*(1), 42-55.

Ant, T., Koukidou, M., Rempoulakis, P., Gong, H. F., Economopoulos, A., Vontas, J., & Alphey, L. (2012). Control of the olive fruit fly using genetics-enhanced sterile insect technique. *BMC biology*, *10*(1), 51.

Carvalho, D. O., McKemey, A. R., Garziera, L., Lacroix, R., Donnelly, C. A., Alphey, L., ... & Capurro, M. L. (2015). Suppression of a field population of Aedes aegypti in Brazil by sustained release of transgenic male mosquitoes. *PLoS neglected tropical diseases*, *9*(7).

Deutsch, C. A., Tewksbury, J. J., Tigchelaar, M., Battisti, D. S., Merrill, S. C., Huey, R. B., & Naylor, R. L. (2018). Increase in crop losses to insect pests in a warming climate. *Science*, *361*(6405), 916-919.

Dhole, S., Vella, M. R., Lloyd, A. L., & Gould, F. (2018). Invasion and migration of spatially self-limiting gene drives: A comparative analysis. *Evolutionary applications*, *11*(5), 794-808.

Foster, G. G., Vogt, W. G., Woodburn, T. L., & Smith, P. H. (1988). Computer simulation of genetic control. Comparison of sterile males and field-female killing systems. *Theoretical and Applied Genetics*, *76*(6), 870-879.

Fouque, F., Carinci, R., Gaborit, P., Issaly, J., Bicout, D. J., & Sabatier, P. (2006). *Aedes aegypti* survival and dengue transmission patterns in French Guiana. *Journal of Vector Ecology*, *31*(2), 390-399.

Fu, G., Condon, K. C., Epton, M. J., Gong, P., Jin, L., Condon, G. C., ... & Alphey, L. (2007). Female-specific insect lethality engineered using alternative splicing. *Nature biotechnology*, *25*(3), 353-357.

Fu, G., Lees, R. S., Nimmo, D., Aw, D., Jin, L., Gray, P., ... & Marinotti, O. (2010). Female-specific flightless phenotype for mosquito control. *Proceedings of the National Academy of Sciences*, *107*(10), 4550-4554.

Gong, P., Epton, M. J., Fu, G., Scaife, S., Hiscox, A., Condon, K. C., ... & Coleman, P. G. (2005). A dominant lethal genetic system for autocidal control of the Mediterranean fruitfly. *Nature biotechnology*, *23*(4), 453-456.

Gentile, J. E., Rund, S. S., & Madey, G. R. (2015). Modelling sterile insect technique to control the population of *Anopheles gambiae. Malaria journal*, *14*(1), 92.

Gorman, K., Young, J., Pineda, L., Márquez, R., Sosa, N., Bernal, D., ... & Kaiser, P. (2016). Short-term suppression of *Aedes aegypti* using genetic control does not facilitate Aedes albopictus. *Pest management science*, *72*(3), 618-628.

Gossen, M., & Bujard, H. (1992). Tight control of gene expression in mammalian cells by tetracycline-responsive promoters. *Proceedings of the National Academy of Sciences*, *89*(12), 5547-5551.

Gould, F., & Schliekelman, P. (2004). Population genetics of autocidal control and strain replacement. *Annual Reviews in Entomology*, *49*(1), 193-217.

Harrington, L. C., Edman, J. D., & Scott, T. W. (2001). Why do female *Aedes aegypti* (Diptera: Culicidae) feed preferentially and frequently on human blood? *Journal of medical entomology*, *38*(3), 411-422.

Harris, A. F., McKemey, A. R., Nimmo, D., Curtis, Z., Black, I., Morgan, S. A., ... & Collado, A. (2012). Successful suppression of a field mosquito population by sustained release of engineered male mosquitoes. *Nature biotechnology*, *30*(9), 828-830.

Harvey-Samuel, T., Morrison, N. I., Walker, A. S., Marubbi, T., Yao, J., Collins, H. L., ... & Shelton, A. M. (2015). Pest control and resistance management through release of insects carrying a male-selecting transgene. *BMC biology*, *13*(1), 49.

581 Hibbard, B. E., Meihls, L. N., Ellersieck, M. R., & Onstad, D. W. (2010). Density-dependent and
582   density-independent mortality of the western corn rootworm: impact on dose calculations
583   of rootworm-resistant Bt corn. *Journal of Economic Entomology*, *103*(1), 77-84.
584
585 Huang, Y., Lloyd, A. L., Legros, M., & Gould, F. (2009). Gene-drive in age-structured insect
586   populations. *Evolutionary Applications*, *2*(2), 143-159.
587
588 Labbé, G. M., Scaife, S., Morgan, S. A., Curtis, Z. H., & Alphey, L. (2012). Female-specific
589   flightless (fsRIDL) phenotype for control of Aedes albopictus. *PLoS neglected tropical*
590   *diseases*, *6*(7).
591
592 Lacroix, R., McKemey, A. R., Raduan, N., Wee, L. K., Ming, W. H., Ney, T. G., ... &
593   Angamuthu, C. (2012). Open field release of genetically engineered sterile male Aedes
594   aegypti in Malaysia. *PloS one*, *7*(8), e42771.
595
596 Leftwich, P. T., Koukidou, M., Rempoulakis, P., Gong, H. F., Zacharopoulou, A., Fu, G., ... &
597   Alphey, L. (2014). Genetic elimination of field-cage populations of Mediterranean fruit
598   flies. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1792), 20141372.
599
600 Muir, L. E., & Kay, B. H. (1998). *Aedes aegypti* survival and dispersal estimated by mark-
601   release-recapture in northern Australia. *The American journal of tropical medicine and*
602   *hygiene*, *58*(3), 277-282.
603
604 Ogaugwu, C. E., Schetelig, M. F., & Wimmer, E. A. (2013). Transgenic sexing system for
605   *Ceratitis capitata* (Diptera: Tephritidae) based on female-specific embryonic
606   lethality. *Insect biochemistry and molecular biology*, *43*(1), 1-8.
607
608 Rendón, P., McInnis, D., Lance, D., & Stewart, J. (2004). Medfly (Diptera: Tephritidae) genetic
609   sexing: large-scale field comparison of males-only and bisexual sterile fly releases in
610   Guatemala. *Journal of Economic Entomology*, *97*(5), 1547-1553.
611
612 Robert, M. A., Okamoto, K., Lloyd, A. L., & Gould, F. (2013). A reduce and replace strategy for
613   suppressing vector-borne diseases: insights from a deterministic model. *PLoS One*, *8*(9),
614   e73233.
615
616 Rueda, L. M., Patel, K. J., Axtell, R. C., & Stinner, R. E. (1990). Temperature-dependent
617   development and survival rates of *Culex quinquefasciatus* and *Aedes aegypti* (Diptera:
618   Culicidae). *Journal of medical entomology*, *27*(5), 892-898.
619
620 Schetelig, M. F., & Handler, A. M. (2012). A transgenic embryonic sexing system for
621   *Anastrepha suspensa* (Diptera: Tephritidae). *Insect biochemistry and molecular*
622   *biology*, *42*(10), 790-795.
623
Schetelig, M. F., Targovska, A., Meza, J. S., Bourtzis, K., & Handler, A. M. (2016).
Tetracycline-suppressible female lethality and sterility in the Mexican fruit fly,
Anastrepha ludens. *Insect molecular biology*, *25*(4), 500-508.

Schliekelman, P., & Gould, F. (2000). Pest control by the release of insects carrying a female-
killing allele on multiple loci. *Journal of Economic Entomology*, *93*(6), 1566-1579.

Scott, M. J. (2014). Development and evaluation of male-only strains of the Australian sheep
blowfly, *Lucilia cuprina*. *BMC genetics*, *15*(2), S3.

Styer, L. M., Minnick, S. L., Sun, A. K., & Scott, T. W. (2007). Mortality and reproductive
dynamics of *Aedes aegypti* (Diptera: Culicidae) fed human blood. *Vector-borne and*
zoonotic diseases, *7*(1), 86-98.

Thomas, D. D., Donnelly, C. A., Wood, R. J., & Alphey, L. S. (2000). Insect population control
using a dominant, repressible, lethal genetic system. *Science*, *287*(5462), 2474-2476.

Wise de Valdez, M. R., Nimmo, D., Betz, J., Gong, H. F., James, A. A., Alphey, L., & Black,
W.C. (2011). Genetic elimination of dengue vector mosquitoes. *Proceedings of the*
National Academy of Sciences, *108*(12), 4772-4775.

Wolfram Research, Inc., Mathematica, Version 12.0, Champaign, IL (2019).

World Health Organization. (2017). Global vector control response 2017-2030. *Global vector*
control response 2017-2030.

Yan, Y., & Scott, M. J. (2015). A transgenic embryonic sexing system for the Australian sheep
blow fly *Lucilia cuprina*. *Scientific reports*, *5*, 16090.

Yan, Y., Williamson, M. E., Davis, R. J., Andere, A. A., Picard, C. J., & Scott, M. J. (2019).
Improved transgenic sexing strains for genetic control of the Australian sheep blow fly
Lucilia cuprina using embryo-specific gene promoters. *Molecular Genetics and*
Genomics, 1-12.