Similar gut bacterial microbiota in two fruit-feeding moth pests collected from different host species and locations

Qiang Gong [1, 3, 4, #], Li-Jun Cao [1, #], Jin-Cui Chen [1], Ya-Jun Gong [1], De-Qiang Pu [4], Qiong Huang [3],

Ary Anthony Hoffmann [2], Shu-Jun Wei [1, *]

1. Institute of Plant and Environmental Protection, Beijing Academy of Agricultural and

Forestry Sciences, 9 Shuguanghuayuan Middle Road, Haidian District, Beijing 100097, China

2. Pest and Environmental Adaptation Research Group, School of BioSciences, Bio21

Institute, University of Melbourne, Parkville, Victoria, Australia

3. College of Forestry, Sichuan Agricultural University, Wenjiang, Sichuan 611130, China

4. Institute of Plant Protection, Sichuan Academy of Agricultural Sciences, Chengdu 610066,

China

# these authors contributed equally.

* Corresponding author: Shu-Jun Wei, Institute of Plant and Environmental Protection,

Beijing Academy of Agricultural and Forestry Sciences, 9 Shuguanghuayuan Middle Road,

Haidian District, Beijing 100097, China; Tel: +86 10 51503439; E-mail: shujun268@163.com

Running title: Gut microbiota in two co-occurring fruit moths

Abstract

Numerous gut microbes are associated with insects, but their composition remains largely unknown for many insect groups, along with factors influencing their composition. Here, we compared gut bacterial microbiota of two co-occurring agricultural pests, the peach fruit moth (PFM) and the oriental fruit moth (OFM), collected from different orchards and host plant species. Gut microbiota of both species was mainly composed of bacteria from Proteobacteria, followed by Firmicutes. The two species shared bacteria from the genera *Pseudomonas*, *Gluconobacter*, *Acetobacter,* and *Pantoea*, although endosymbiotic *Wolbachia* was the most abundant genus in PFM and *Lactobacillus* was the most abundant in OFM. PFM tended to have lower diversity and richness of gut microbiota than OFM; however, only some of the comparisons were statistically significant. Orchards can influence gut microbiota in terms of richness, particularly for PFM, but not so much for diversity parameters. Functional prediction of gut microbiota showed that the top pathways are amino acid metabolism, translation, and membrane transport in both species, but their abundance varied between the two moth species. These results show that two fruit moths share many features of gut microbiota, and the bacterial species are relatively stable within moth species even when they use different host plants. Our study suggests that fruit-feeding behavior may play a role in shaping gut microbiota of the two fruit moths, which may provide microbial targets for pest control.

Importance

Understanding the associated microbes with insects can point to new targets for pest control. Here we compared bacterial community in the gut of two co-occurring agricultural pests, the peach fruit moth (PFM) and the oriental fruit moth (OFM), collected from different orchards and host plant species. We found that the bacterial genera *Pseudomonas*, *Gluconobacter*, *Acetobacter,* and *Pantoea* are abundant and shared in two moths. The composition of the bacterial species is relatively stable within moth species even when they use different host plants, indicating that the gut microbiota community in the PFM and OFM is likely to be related to their fruit-feeding behavior. The findings have implications for developing novel pest control approaches by targeting gut microbes associated with the two moths.

Keywords: *Carposina sasakii*; *Grapholita molesta*; gut microbiome; host; orchard; *Wolbachia*; endosymbiont

Introduction

Many microorganisms have become adapted to their insect hosts, forming close mutualistic relationships (1, 2). These microbes play important roles for their hosts, such as in the digestion and nutrient absorption of host food, protection against pathogens, and enhancement of immunity (3-5). The study of insect microorganisms can point to new approaches for the control of agricultural pests and human disease vectors as well as increasing the value of economically important insects, particularly by modifying the symbiotic relationship between symbionts and their hosts (6, 7).

The community of microorganisms living in insects can be affected by many environmental factors (1, 8, 9). In particular, gut bacteria of different insects can vary greatly in number, composition, distribution, and function for species adapted to different hosts and living in different habitats (10). Moreover, there can be a dynamic interaction between bacteria living in the gut and the environment as indicated by the acquisition and loss of Se*rratia* symbiotica strains in aphids (11)

Moths include some of the most damaging agricultural and forest pests from the order Lepidoptera. Being holometabolous, moths are characterized by different life stages and can vary in their gut microbiota during development (12-14). Many moths are polyphagous, having a wide range of diets, which represent one of the factors impacting bacterial communities in this group (13, 15). However, while moths represent useful model species to understand the determinants of gut microbiota across life stages (16), there is limited information on variation in their microbiota.

Here we focus on the peach fruit moth (PFM), *Carposina sasakii*, and the oriental fruit

moth (OFM), *Grapholita molesta*, common agricultural moth pests damaging many

economically important fruit crops, such as apple, pear, and peach (17-20). Larvae of both

these species bore into and feed on fruit, while OFM can also bore into tree shoots prior to

pupation. These species usually co-occur in the same orchard and sometimes on the same

fruit (21-23). The concealed lifestyle and wide range of host plant species used by these

species make them useful to understand factors affecting their gut microbiota. Previous

studies found that larvae of these two moths harbor a high diversity and richness of bacteria

(24, 25), but it is not yet clear the two species are more likely to share the same gut

microbores when they live in the same orchard and on the same host plant species.

We examined gut bacterial microbiota in co-occurring PFM and OFM collected from the

same host plant species, with the microbiota characterized using the V3-V4 variable region

of the 16S rRNA gene. We aimed to examine the relative contribution of moth species, host

plant, and other factors related to variation among orchards to microbial composition.

Results

Community composition of the gut microbiota in PFM and OFM

The average number of sequencing reads for each sample was 4927 after filtering (**Table S1**).

Rarefaction curves from both the original sequencing data sets and randomly subsampled

data sets showed that the curves of all samples tended to be flat, indicating that the amount

of sequencing data is enough to reflect most of the microbial diversity information in the

samples (**Fig. S1**). In total, 294 OTU were clustered, attributed to 13 phyla and 176 genera

and 234 species for both hosts, among which 234 OTUs belonging to 203 species and 284

OTUs belonging to 228 species were identified for PFM and OFM respectively (**Table S2**).

At the phylum level, OTUs in both species were mainly attributed to Proteobacteria

(98.4% in PFM, 89.2% in OFM), followed by Firmicutes (1.06% in PFM, 8.87% in OFM) (**Fig.**

1a). At the genus level, OTUs of PFM were mainly annotated to *Wolbachia* (62.06%),

Pseudomonas (19.09%), *Gluconobacter* (6.98%), *Acetobacter* (4.05%), and *Pantoea* (3.59%),

while OTUs of OFM were mainly annotated to *Pseudomonas* (49.96%), *Gluconobacter*

(12.53%), *Pantoea* (10.70%), *Lactobacillus* (7.65%), *Acetobacter* (6.61%) (**Fig. 1b** and **Fig. S2**).

Similar patterns were found at the species level where these could be identified (**Fig. 1c,**

Table S3).

The core bacterial community at the genus level for each species was identified by

comparing individuals from different orchards (**Table S4**). For PFM, 24 core genera were

identified from the four orchards sampled (**Fig. 2a**), the most common of which were

Wolbachia (67.07%), followed by *Pseudomonas* (20.63%), *Gluconobacter* (7.54%) and

Pantoea (3.88%) (**Fig. 2b**); for OFM, 33 core genera were identified from five orchards (**Fig.**

2c), the most common of which were *Pseudomonas* (59.96%), followed by *Gluconobacter*

(15.03%), *Pantoea* (12.84%), and *Lactobacillus* (9.18%) (**Fig. 2d**).

In summary, Proteobacteria was the most abundant phylum for both host species.

There were many common bacteria from the genus *Pseudomonas*, *Gluconobacter*,

Acetobacter and *Pantoea*, although in PFM *Wolbachia* was the most abundant genus

followed by *Pseudomonas*. *Pseudomonas* was the most abundant genus in OFM, and

Lactobacillus and *Dickeya* were abundant in OFM but not in found in PFM (**Figs. 1d, 2b, 2d**).

Comparison on gut microbiota between PFM and OFM

When gut bacterial microbiota was compared between all samples of PFM and OFM, in

terms of alpha diversity, there was no significant difference in OTU richness between PFM

and OFM (P_{ace} = 0.57, P_{chao} = 0.121, P_{sobs} = 0.014) (**Fig. 3a-b**, **Tables S5,** and **S6**) but

significantly lower diversity in PFM than in OFM ($P_{shannon}$ = 0.002, $P_{simpson}$ = 0.004) (**Fig. 3c-d**,

Tables S5 and **S6**). In terms of beta diversity, PFM and OFM individuals divided into two

groups in the PCoA analysis, although outlier samples were identified (**Fig. 3e, Table S7**).

We then compared gut bacterial microbiota between three pairs of PFM and OFM

populations collected from the same host species and the same orchard. In terms of alpha

diversity, OFM usually had higher richness and diversity except for one paired richness

comparison collected from apple (**Fig. S3a, Table S5**). For pear, patterns were consistent, but

only one of the three comparisons was statistically different in diversity (**Fig. S3h**). In terms

of beta diversity, individuals of PFM and OFM from the same habitat could be clustered into

different groups in the PCoA analysis (**Fig. S3c, f, i**), with individuals collected from pear

showing the clearest separation (**Fig. S3i, Table S7**).

Influence of orchard on gut microbiota within species

First, we compared the gut microbiota of the same insect species collected from different

host plant species and different orchards to examine the effect of orchard but relaxing the

host plant species. PFM individuals from four orchards and OFM individuals from five

orchards were analyzed. For PFM, four of six pairs of orchard comparisons had significant

differences in richness, while one of the six pairs showed difference in diversity (**Fig. S4**). For

OFM, two of 10 pairs of orchard comparisons were significantly different for richness, but

none were significant for diversity (**Fig. S5**). For overall comparison, there was no significant

difference in any measure of richness or diversity in either species (χ^2 = 18/18/18/14.36/18,

df = 18/18/18/14/18, p > 0.4231 for Ace, Shannon, Simpson, Sobs and Chao in PFM; χ^2 = 24,

df = 24/24/24/23/24, p > 0.4038 for Ace, Shannon, Simpson, Sobs and Chao in OFM).

Second, we compared the gut microbiota of the same species and host plant from different

orchards to test the effect of the orchard by fixing the host plant. Two pairs of PFMs from

pear and apple and two pairs of OFM from pear and peach shoot hosts were used for

analysis. In terms of alpha diversity (Ace), a significant difference in richness was found in

both paired PFM comparisons (**Figs. S6a, S6g**) and one of the two OFM comparisons (**Fig.**

S6d), while a significant difference in Shannon's index was found in one of the two PFM

comparisons (**Fig. S6b**) but not in the OFM comparisons (**Figs. S6h, S6k**). In terms of beta

diversity, individuals from different orchards of the same species were not clearly separated

in the PCoA analyses (**Fig. 5, Figs. S6c, f, i, l**).

While these results suggest that orchard can affect the composition of gut microbiota in

PFM and OFM, the effect is relatively small, particularly as shown in the beta diversity

analysis. Orchard had a higher impact on richness than on diversity, and PFM tended to vary

more among orchards with the same host than OFM.

Function prediction of gut microbiota

At level 1, functions of the gut microbiota were mainly annotated to pathways of

metabolism, genetic information processing, environmental information processing, and

cellular processing. At level 2, the top pathways were amino acid metabolism, translation

and membrane transport (**Table 2**). It can be seen in the COG (Clusters of Orthologous

Groups) function annotation that the functions of gut microbiota of OFM and PFM were

annotated to the same pathway, but the abundance of the same pathway was different (**Fig.**

6a). Among the top 10 functions in KEGG (Kyoto Encyclopedia of Genes and Genomes)

annotations pathway level 3, there was a significant difference between OFM and PFM ($p <$

0.0001). ABC transporters and two-component system were significantly higher in OFM than

PFM, and the remaining eight were higher in PFM than OFM (**Fig. 6b**).

Discussion

Comparison of gut microbiota from two fruit borers

In this study, we found that the gut microbiota of PFM and OFM was dominated by

Proteobacteria and Firmicutes, which is similar to the situation found in Y. Liu et al. (25) and

Y. Li et al. (24), and in other lepidopterans such as *Lymantria dispar*, *Helicoverpa armigera*,

and *Bombyx mori* (26-29). However, there was a difference between PFM and OFM and

other lepidopterans at the genus level. OTUs from both PFM and OFM was dominated by

Pseudomonas, *Gluconobacter*, *Acetobacter,* and *Pantoea*. In contrast, in silkworms,

Aureimonas, *Methylobacterium*, *Rhizobium*, *Sphingomonas*, *Propionibacterium*,

Pseudomonas, and *Microbacterium* were the most common genera (29). The results suggest

that PFM and OFM gut microbes had a similar composition, but they are different from

those of the *Bombyx mori*, which has a different diet. Our results support the notion that

dietary adaptation has led to different intestinal microorganisms and symbiotic interactions

(30), although more moth species with different hosts (fruit, leaf tissue, and so on) need to

be included in such comparisons.

We also found some differences between the two species examined here, where OTUs

of PFM were dominated by *Wolbachia*, and OTUs of OFM were dominated by *Lactobacillus*.

When we focused on the gut microbes of PFM and OFM from the same host and the same

orchard, this pattern was also found: *Wolbachia* was unique to PFM, while *Lactobacillus* was

abundant in OFM and rare in PFM. Perhaps this difference in species might generate

phenotypic differences among the species for traits such as pesticide resistance. For instance,

insecticide-treated resistant strains of the diamondback moth *Plutella xylostella* had more

Lactobacillales and the less common taxa *Pseudomonadales* and *Xanthomonadales* as well

as fewer *Enterobacteriales* compared with a susceptible strain (31). The OFM microbiota

might contribute to resistance, although living in fruit they would be less affected by

pesticides than *Plutella xylostella* larvae feeding on leaves. The comparison of microbes of

PFM and OFM in three orchards showed that there was no large difference in microbial

richness and diversity between PFM and OFM, but the PCA analysis highlighted differences

in species composition, with host type clearly being a major determinant of gut

microorganisms.

Influence of orchard and host species on gut microbiota

Microbial communities can vary among host locations, both in terms of community diversity

and community structure (32). In our study, there were differences in microbial richness in

larvae from the same species collected from different orchards with the same type of fruit

(PLPR/PKPR, PDAE/PGAE, OLPR/OKPR, Table S5), which suggests an impact of orchard

habitat on microbial richness. Differences in microbial diversity have also been noted in

studies on other insects, such as in comparisons of *Drosophila* between indoor and wild

environments (33). However, the gut microorganisms in neither PFM nor OFM could be

clearly separated by orchard or fruit type in the PCoA analysis, suggesting that host species

rather than location plays a more important role in microbial community composition.

Wolbachia in PFM

Wolbachia is an intracellular endosymbiont rather than a gut bacterium, but it can be found

in the gut wall of species (34). The role of *Wolbachia* in PFM is unclear; it is common in

Lepidoptera (35) where its effects have mostly not been characterized in species although in

Lepidoptera it can cause a variety of effects on host reproduction including cytoplasmic

incompatibility, feminization and male-killing (36-38) and increases the susceptibility

of its host to baculovirus (39). These effects have not yet been investigated in PFM and

require a comparison of *Wolbachia* infected and uninfected individuals for fitness as well as

crosses to establish reproductive effects.

Of particular interest from the perspective of the current study is whether *Wolbachia*

might influence the gut microbiota. *Wolbachia* may lead to decreased microbial diversity

due to competitive behavior (40), which may contribute to the lower diversity of gut

microbiota in PFM than that of OFM. In *Drosophila melanogaster*, *Wolbachia* can reduce the

richness of *Acetobacter* (41), but this group was not at a low abundance in PFM. Whether

Wolbachia in PFM influences, other microbiota requires a comparison of *Wolbachia* infected

and *Wolbachia* free lines, which might be generated through antibiotic treatment or by

taking advantage of natural polymorphism in infection status within natural populations

(42).

Implications for pest management

The insect-associated microbes provide new targets for developing novel pest control

methods (6, 16, 43, 44). The first step to find the potential bacterial targets is to investigate

the bacterial community, its impact on the pests, and its stability. We found that the

community of the gut microbiota were relatively stable within moth species in spite of host

fruit differences for microbes such as *Pseudomonas, Pantoea, Lactobacillus, Gluconobacter,*

and *Acetobacter.* Functional analysis showed that three of the ten most abundant functions

were environmental signaling processes, and others involve metabolism, genetic information

processing, and cellular processes. These functional classes suggest that gut bacteria have a

clear interaction with host processes in the intestinal environment. Among the abundant

bacteria taxa, *Pseudomonas brenneri* plays a prominent role in the removal of heavy metals

(45). This species is significantly more abundant in OFM than PFM and may contribute to the

different ratios of ABC transporters and the Two-component system. *Gluconobacter cerinus*

was another species present in PFM and OFM, which may have a beneficial role as in the

case of fruit flies where it can affect reproduction (46). *Pantoea* is a highly diverse genus that

can cause plant diseases and human diseases but also have functions in habitat restoration

and pesticide degradation (47). Functional studies of these bacteria may help to identify

potential targets for developing control methods of these two fruit moths.

We also note that the two fruit moths share many gut bacterial taxa. The similar

composition of gut bacterial microbiota indicates functions related to the common biology

of both species, particularly in terms of the fruit-feeding larvae. These larvae bore into fruit

or shoots soon after egg hatching, reducing their likelihood of exposure to environmental

bacteria when compared to the leaf-feeding moths. In the fruit-feeding spotted wing

drosophila, *Drosophila suzukii*, the gut microbiota provides nutrition by providing protein for

their hosts (48). Larvae of fruit moths often feed on immature fruits, which are rich in

compounds such as organic acids and tannins. The tannins are endogenous inhibitors of the

growth of numerous species of pests by negatively effecting the metabolism of insects (49).

We found that the most abundant function of the gut microbiota in both species were

metabolic processes. There are examples of gut microbiota in lepidopteran hosts helping to

detoxify host toxins (50, 51), but whether the fruit moths need microbes to help them to

detoxify defensive chemicals is unclear. Nevertheless, the gut microbiota community in the

PFM and OFM is likely to be related to their fruit-feeding behavior, and further tests of such

hypotheses may provide insights into the development of novel control approaches.

Materials and methods

Sample collection and DNA extraction

We sampled three pairs of PFM and OFM populations from the same host plant and orchard,

as well as one PFM population from another apple orchard, and two OFM populations from

two peach orchards infesting tree shoots (**Table 2**). We collected potentially infested pears

and apples and peach shoots from the field and kept them in the conditioned laboratory

under 25 ± 1 ℃, 60% ± 5% humidity, and a photoperiod of 16 h light: 8 h dark. Fifth instar

larvae were collected when they came out from the collected hosts. Species were identified

by morphology and kept in a clean 1.5 ml tube for 24 hours to clean out the feces by

starvation. Then, larvae were frozen in liquid nitrogen and stored in a -80 ℃ refrigerator

prior to usage. We examined the gut microbiota of 19 PFM and 25 OFM individuals (**Table 2**).

Prior to DNA extraction, larvae were washed three times, with 75% alcohol, and then

washed three times with sterile water. The whole gut tissue was dissected and homogenized

in a 1.5 ml tube by grinding manually. Total DNA was extracted from single samples using the

E.Z.N.A.® Bacterial DNA Kit (Omega Bio-tek, GA, U.S.) according to manufacturer's protocol.

The concentration and quality of the extracted DNA were determined by a NanoDrop 2000

UV-vis spectrophotometer (Thermo Scientific, Wilmington, USA) and gel electrophoresis on

1% agarose.

16S rRNA gene amplification and sequencing

We used the V3-V4 hypervariable regions of the bacterial 16S ribosomal RNA (rRNA) gene to

examine the gut microbiota of PFM and OFM. A 468-bp target gene segment was amplified

by primer pair of 338F (5'-ACTCCTACGGGAGGCAGCAG-3') and 806R

(5'-GGACTACHVGGGTWTCTAAT-3') (52). For PCR reaction, 20 μL of the mixture was prepared,

including 5 x FastPfu reaction buffer, 250 μM dNTPs 1 U FastPfu Polymerase (Transgene,

Beijing, China), 200 nM of each prime (Majorbio, Shanghai, China), 1 μL of template DNA

and DNA-free water. The PCR reaction involved a single denaturation step at 95 °C for 3 min,

followed by 27 cycles of 95 °C for 30 s, 55 °C for 30 s, 72 °C for 45 s, and finished after a final

extension at 72 °C for 10 min. The PCR products were run on a 2% (w/v) agarose gel and

those with correct size were excised and purified with a AxyPrep DNA gel extraction kit

(Axygen Biosciences, Union City, CA, USA). Illumina Miseq sequencing libraries were

constructed using the TruSeqTM DNA Sample Prep Kit (San Diego, CA, USA) for the purified

16S PCR products, and sequenced on an Illumina MiSeq (San Diego, CA, USA) to obtain

300-bp paired-end reads.

Quality control and OTU identification

Raw data from Illumina MiSeq sequencing were demultiplexed to obtain sequencing data for

each sample. The quality of raw data was checked by FASTQC version 0.19.6 (53); low-quality

data were trimmed and filtered by Trimmomatic version 0.36 (54). Paired-end reads were

merged by FLASH version 1.2.11 (55) to generate unpaired longer reads with the following

criteria: (i) the reads were truncated at any site receiving an average quality score < 20 over

a 50 bp sliding window; (ii) primers were exactly matched allowing two nucleotide

mismatching, and reads containing ambiguous bases were removed; (iii) only paired-end

reads whose overlap longer than 10 bp were merged.

Operational taxonomic units (OTUs) were clustered with a 97% similarity threshold

using UPARSE version 7.0.1090 (56), and chimeric sequences were identified and removed

using UCHIME algorithm in USEARCH version 7.0 (57). The taxonomy of each 16S rRNA gene

sequence was analyzed by a naïve Bayesian classifier of Ribosomal Database Project version

2.11 (58) against the SILVA rRNA database (59). To avoid the influence of sequencing depth

in samples, sequences from difference samples were rarefied to the same depth. Sample

sequence extraction and species screening of OTU were conducted in accordance with the

following conditions: (i) removal of mitochondrial and chloroplast sequences; (ii) retention

of only OTUs with sequence depth greater than or equal to five in at least three samples in

subsequent analyses.

Diversity analysis

For alpha diversity, community richness indexes (sobs, chao, and ace) and community

diversity indexes (Shannon, Simpson, and Pd) were estimated. The software Mothur (60) was

used to calculate the alpha diversity index under different random sampling, and the *ggplot2*

R package was used to draw the rarefaction curves. The Wilcoxon rank-sum test was used to

compare statistical differences between different groups, while the Kruskal-Wallis rank sum

test was used for overall comparison to examine the species, host, and orchard effects. In

the beta diversity analysis, principal coordinates analysis (PCoA) was conducted based on a

Bray-Curtis dissimilarity matrix computed from the samples. For group comparisons, a

non-parametric multivariate statistical test, permutational multivariate analysis of variance

(PERMANOVA), was conducted based on the Bray-Curtis dissimilarity matrix, using Qiime

and the R package *vegan* (61).

Functional analysis

We used PICRUSt version 1.1.4 (62) to predict the function of the gut microbiota from PFM

and OFM. The OTU abundance table was first normalized by removing the effect of the 16S

rRNA gene copy numbers (GCNs). The COG (Clusters of Orthologous Groups) family

information was obtained according to the Greengene id version gg_13_5 (63)

corresponding to each OTU. The description information of each COG and its function

information was parsed based on the eggNOG (evolutionary genealogy of genes:

Non-supervised Orthologous Groups) database v5.0 (64). The 16S rRNA taxonomic lineage

based on the SILVA rRNA database (59) was transformed into the taxonomic lineage of

prokaryotes in the KEGG (Kyoto Encyclopedia of Genes and Genomes) database Release 92.0

(65) through Tax4Fun (66), and the 16S rRNA gene sequence was functionally annotated. A

Wilcoxon rank-sum test was used to compare the statistical difference between OFM and

PFM for the 10 most abundant pathways at level 3.

Acknowledgements

Funding for this study was provided jointly by the National Key Research and Development

Program of China (2019YFD1002102), the Beijing Key Laboratory of Environmentally Friendly

Pest Management on Northern Fruits (BZ0432) and BAAFS-UOM Joint Laboratory on Pest

Control Research. SJW conceived and designed research. DQP, YJG, JCC and QH collected the

samples. QG and LJC conducted experiments. QG, SJW and LJC analyzed data. QG, SJW and

AAH wrote the manuscript and discussed the results.

References

1. Shapira M. 2016. Gut microbiotas and host evolution: scaling up symbiosis. Trends in
Ecology and Evolution 31:539-549.
2. Dillon R, Dillon V. 2004. The gut bacteria of insects: nonpathogenic interactions. Annual
Reviews of Entomology 49:71-92.
3. Engel P, Moran NA. 2013. The gut microbiota of insects–diversity in structure and function.
FEMS Microbiology Reviews 37:699-735.
4. Douglas AE. 2015. Multiorganismal insects: diversity and function of resident
microorganisms. Annual Review of Entomology 60:17-34.
5. Dicke M, Cusumano A, Poelman EH. 2020. Microbial symbionts of parasitoids. Annual
Review Entomology 65:171-190.
6. Berasategui A, Shukla S, Salem H, Kaltenpoth M. 2016. Potential applications of insect
symbionts in biotechnology. Applied Microbiology and Biotechnology
100:1567-1577.
7. Walker T, Johnson PH, Moreira LA, Iturbe-Ormaetxe I, Frentiu FD, McMeniman CJ, Leong
YS, Dong Y, Axford J, Kriesner P, Lloyd AL, Ritchie SA, O'Neill SL, Hoffmann AA. 2011.
The wMel *Wolbachia* strain blocks dengue and invades caged *Aedes aegypti*
populations. Nature 476:450-U101.
8. Hansen AK, Moran NA. 2014. The impact of microbial symbionts on host plant utilization
by herbivorous insects. Molecular Ecology 23:1473-1496.
9. Adair KL, Douglas AE. 2017. Making a microbiome: the many determinants of
host-associated microbial community composition. Current Opinion in Microbiology
35:23-29.
10. Priya NG, Ojha A, Kajla MK, Raj A, Rajagopal R. 2012. Host plant induced variation in gut
bacteria of *Helicoverpa armigera*. PLoS ONE 7:e30768.
11. Renoz F, Pons I, Vanderpoorten A, Bataille G, Noel C, Foray V, Pierson V, Hance T. 2019.
Evidence for gut-associated *Serratia* symbiotica in wild aphids and ants provides new
perspectives on the evolution of bacterial mutualism in insects. Microbial Ecology
78:159-169.
12. Mereghetti V, Chouaia B, Limonta L, Locatelli DP, Montagna M. 2019. Evidence for a
conserved microbiota across the different developmental stages of *Plodia*

375   *interpunctella*. Insect Science 26:466-478.

376   13. Staudacher H, Kaltenpoth M, Breeuwer JA, Menken SB, Heckel DG, Groot AT. 2016.

377   Variability of bacterial communities in the moth *Heliothis virescens* indicates

378   transient association with the host. PLoS ONE 11:e0154514.

379   14. Lin XL, Pan QJ, Tian HG, Douglas AE, Liu TX. 2015. Bacteria abundance and diversity of

380   different life stages of *Plutella xylostella* (Lepidoptera: Plutellidae), revealed by

381   bacteria culture-dependent and PCR-DGGE methods. Insect Science 22:375-385.

382   15. Montagna M, Mereghetti V, Gargari G, Guglielmetti S, Faoro F, Lozzia G, Locatelli D,

383   Limonta L. 2016. Evidence of a bacterial core in the stored products pest *Plodia*

384   *interpunctella*: the influence of different diets. Environmental Microbiology

385   18:4961-4973.

386   16. Mereghetti V, Chouaia B, Montagna M. 2017. New insights into the microbiota of moth

387   pests. International Journal of Molecular Sciences 18:2450.

388   17. Ma G, Tian BL, Zhao F, Wei GS, Hoffmann AA, Ma CS. 2017. Soil moisture conditions

389   determine phenology and success of larval escape in the peach fruit moth, *Carposina*

390   *sasakii* (Lepidoptera, Carposinidae): Implications for predicting drought effects on a

391   diapausing insect. Applied Soil Ecology 110:65-72.

392   18. Quan LF, Qiu GS, Zhang HJ, Sun LN, Li YY, Yan WT. 2016. Sublethal concentration of

393   beta-cypermethrin influences fecundity and mating behavior of *Carposina sasakii*

394   (Lepidoptera: Carposinidae) adults. Journal of Economic Entomology 109:2196-2204.

395   19. Rothschild GHL, Vickers RA. 1991. Biology, ecology and control of the oriental fruit moth,

396   p 389-412. *In* van der Geest L, Evenhuis H (ed), World Crop Pests Tortricid pests: their

397   biology, natural enemies and control, vol 5. Elsevier, Amsterdam.

398   20. Wang YZ, Li BY, Hoffmann AA, Cao LJ, Gong YJ, Song W, Zhu JY, Wei SJ. 2017. Patterns of

399   genetic variation among geographic and host-plant associated populations of the

400   peach fruit moth *Carposina sasakii* (Lepidoptera: Carposinidae). BMC Evolutionary

401   Biology 17:265.

402   21. Kwon D, Kwon H, Kim D, Yang C. 2018. Larval species composition and genetic structures

403   of *Carposina sasakii*, *Grapholita dimorpha*, and *Grapholita molesta* from Korea.

404   Bulletin of entomological research 108:241-252.

405   22. Kwon DH, Kim DH, Kim HH, Lee SH, Yang CY. 2017. Genetic diversity and structure in

406   apple-infesting pests of *Carposina sasakii*, *Grapholita dimorpha* and *Grapholita*

407   *molesta* in Korea. Journal of Asia-Pacific Entomology 20:13-16.

408   23. Sarker S, Lim UT. 2019. Development and fecundity performance of *Grapholita molesta*

409   and *Grapholita dimorpha* (Lepidoptera: Tortricidae) on different immature fruits.

410   PLoS ONE 14:e0217492.

411   24. Li Y, Sun L, Tian Z, Haibin H, Zhang H, Guisheng Q, Yan W, Yue Q. 2018. Gut bacterial

412   community diversity in *Carposina sasakii* and *Grapholitha molesta*. Chinese Journal

of Applied Ecology 29:293-300.

25. Liu Y, Shen Z, Yu J, Li Z, Liu X, Xu H. 2020. Comparison of gut bacterial communities and
their associations with host diets in four fruit borers. Pest Management Science
76:1353-1362.

26. Yun JH, Roh SW, Whon TW, Jung MJ, Kim MS, Park DS, Yoon C, Nam YD, Kim YJ, Choi JH,
Kim JY, Shin NR, Kim SH, Lee WJ, Bae JW. 2014. Insect gut bacterial diversity
determined by environmental habitat, diet, developmental stage, and phylogeny of
host. Applied and Environmental Microbiology 80:5254-64.

27. Broderick NA, Raffa KF, Goodman RM, Handelsman J. 2004. Census of the bacterial
community of the gypsy moth larval midgut by using culturing and
culture-independent methods. Applied and Environmental Microbiology 70:293-300.

28. Hui XA, Wei GF, Jia SH, Huang JH, Miao XX, Zhou Z, Zhao LP, Huang YP. 2006. Microbial
communities in the larval midgut of laboratory and field populations of cotton
bollworm (*Helicoverpa armigera*). Canadian Journal of Microbiology 52:1085-1092.

29. Chen B, Du K, Sun C, Vimalanathan A, Liang X, Li Y, Wang B, Lu X, Li L, Shao Y. 2018. Gut
bacterial and fungal communities of the domesticated silkworm (*Bombyx mori*) and
wild mulberry-feeding relatives. The ISME Journal 12:2252-2262.

30. Colman DR, Toolson EC, Takacs-Vesbach CD. 2012. Do diet and taxonomy influence insect
gut bacterial communities? Molecular Ecology 21:5124-5137.

31. Xia XF, Zheng DD, Zhong HZ, Qin BC, Gurr GM, Vasseur L, Lin HL, Bai JL, He WY, You MS.
2013. DNA sequencing reveals the midgut microbiota of diamondback moth, *Plutella*
xylostella (L.) and a possible relationship with insecticide resistance. PLos ONE
8:e68852.

32. Meriweather M, Matthews S, Rio R, Baucom RS. 2013. A 454 survey reveals the
community composition and core microbiome of the common bed bug (*Cimex*
lectularius) across an urban landscape. PLoS ONE 8:e61465.

33. Chandler JA, Lang JM, Bhatnagar S, Eisen JA, Kopp A. 2011. Bacterial communities of
diverse *Drosophila* species: ecological context of a host-microbe model system. PLoS
Genetics 7:e1002272.

34. Werren JH. 1997. Biology of *Wolbachia*. Annual Review of Entomology 42:587-609.

35. Ahmed MZ, Araujo-Jnr EV, Welch JJ, Kawahara AY. 2015. *Wolbachia* in butterflies and
moths: geographic structure in infection frequency. Frontiers in Zoology 12:16.

36. Kageyama D, Nishimura G, Hoshizaki S, Ishikawa Y. 2002. Feminizing *Wolbachia* in an
insect, *Ostrinia furnacalis* (Lepidoptera: Crambidae). Heredity 88:444.

37. Berec L, Maxin D, Bernhauerová V, Rohr J. 2016. Male-killing bacteria as agents of insect
pest control. Journal of Applied Ecology 53:1270-1279.

38. Hurst GD, Johnson AP, Schulenburg JH, Fuyama Y. 2000. Male-killing *Wolbachia* in
Drosophila: a temperature-sensitive trait with a threshold bacterial density. Genetics

156:699-709.

39. Kaiser W, Huguet E, Casas J, Commin C, Giron D. 2010. Plant green-island phenotype
induced by leaf-miners is mediated by bacterial symbionts. Proceedings of the Royal
Society B-Biological Sciences 277:2311-2319.

40. Dittmer J, Bouchon D. 2018. Feminizing *Wolbachia* influence microbiota composition in
the terrestrial isopod *Armadillidium vulgare*. Scientific Reports 8:6998.

41. Simhadri RK, Fast EM, Guo R, Schultz MJ, Vaisman N, Ortiz L, Bybee J, Slatko BE, Frydman
HM. 2017. The gut commensal microbiome of *Drosophila melanogaster* is modified
by the endosymbiont *Wolbachia*. mSphere 2:e00287-17.

42. Shi M, White VL, Schlub T, Eden JS, Hoffmann AA, Holmes EC. 2018. No detectable effect
of Wolbachia wMel on the prevalence and abundance of the RNA virome of
Drosophila melanogaster. Proceedings of the Royal Society B-Biological Sciences
285:1883.

43. Bosch TJ, Welte CU. 2017. Detoxifying symbionts in agriculturally important pest insects.
Microbial Biotechnology 10:531-540.

44. Itoh H, Tago K, Hayatsu M, Kikuchi Y. 2018. Detoxifying symbiosis: microbe-mediated
detoxification of phytotoxins and pesticides in insects. Natural Products Reports
35:434-454.

45. Banerjee S, Kamila B, Barman S, Joshi SR, Mandal T, Haider G. 2019. Interlining Cr(VI)
remediation mechanism by a novel bacterium *Pseudomonas brenneri* isolated from
coalmine wastewater. Journal of Environmental Management 233:271-282.

46. He MY, Jiang JJ, Cheng DF. 2017. The plant pathogen Gluconobacter cerinus strain CDF1 is
beneficial to the fruit fly *Bactrocera dorsalis*. Amb Express 7:207.

47. Walterson AM, Stavrinides J. 2015. *Pantoea*: insights into a highly versatile and diverse
genus within the Enterobacteriaceae. FEMS Microbiology Reviews 39:968-984.

48. Bing X, Gerlach J, Loeb G, Buchon N. 2018. Nutrient-dependent impact of microbes on
Drosophila suzukii development. mBio 9:e02199-17.

49. Schultz J. 1989. Tannin-insect interactions. *In* Hemingway RW, Karchesy JJ, Branham SJ
(ed), Chemistry and Significance of Condensed Tannins. Springer, Boston, MA.

50. Ceja-Navarro JA, Vega FE, Karaoz U, Hao Z, Jenkins S, Lim HC, Kosina P, Infante F, Northen
TR, Brodie EL. 2015. Gut microbiota mediate caffeine detoxification in the primary
insect pest of coffee. Nature Communications 6:7618.

51. Vilanova C, Baixeras J, Latorre A, Porcar M. 2016. The generalist inside the specialist: gut
bacterial communities of two insect species feeding on toxic plants are dominated by
Enterococcus sp. Frontiers in Microbiology 7:1005.

52. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a
dual-index sequencing strategy and curation pipeline for analyzing amplicon
sequence data on the MiSeq Illumina sequencing platform. Appllied Environmental

Microbiology 79:5112-20.

53. Andrews S. 2013. FastQC: a quality control tool for high throughput sequence data.

Available online at: http://wwwbioinformaticsbabrahamacuk/projects/fastqc.

54. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30:2114-20.

55. Magoc T, Salzberg SL. 2011. FLASH: fast length adjustment of short reads to improve

genome assemblies. Bioinformatics 27:2957-63.

56. Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads.

Nature Methods 10:996-338.

57. Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST.

Bioinformatics 26:2460-1.

58. Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naive Bayesian classifier for rapid

assignment of rRNA sequences into the new bacterial taxonomy. Applied and

Environmental Microbiology 73:5261-7.

59. Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J, Ludwig W,

Glockner FO. 2014. The SILVA and "All-species Living Tree Project (LTP)" taxonomic

frameworks. Nucleic Acids Research 42:D643-8.

60. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,

Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ,

Weber CF. 2009. Introducing mothur: open-source, platform-independent,

community-supported software for describing and comparing microbial communities.

Applied and Environmental Microbiology 75:7537-41.

61. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N,

Pena AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE,

Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh

PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J, Knight R. 2010. QIIME allows

analysis of high-throughput community sequencing data. Nature Methods 7:335-336.

62. Langille MGI, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, Clemente JC,

Burkepile DE, Thurber RLV, Knight R, Beiko RG, Huttenhower C. 2013. Predictive

functional profiling of microbial communities using 16S rRNA marker gene sequences.

Nature Biotechnology 31:814-821.

63. DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P,

Andersen GL. 2006. Greengenes, a chimera-checked 16S rRNA gene database and

workbench compatible with ARB. Applied and Environmental Microbiology

72:5069-5072.

64. Huerta-Cepas J, Szklarczyk D, Heller D, Hernandez-Plaza A, Forslund SK, Cook H, Mende

DR, Letunic I, Rattei T, Jensen LJ, von Mering C, Bork P. 2019. eggNOG 5.0: a

hierarchical, functionally and phylogenetically annotated orthology resource based

on 5090 organisms and 2502 viruses. Nucleic Acids Research 47:D309-D314.

65. Kanehisa M, Goto S. 2000. KEGG: kyoto encyclopedia of genes and genomes. Nucleic

Acids Research 28:27-30.

66. Aßhauer KP, Wemheuer B, Daniel R, Meinicke P. 2015. Tax4Fun: predicting functional

profiles from metagenomic 16S rRNA data. Bioinformatics 31:2882-2884.

Table 1 Enrichment of KEGG pathways for gut bacterial microbiota of the peach fruit moth (PFM) *Carposina sasakii* and the oriental fruit moth (OFM) *Grapholita molesta*.

Pathway level 1	Pathway level 2	PFM	OFM
Metabolism	**Amino acid metabolism**	**9.20%**	**5.21%**
Metabolism	**Amino acid metabolism**	**9.11%**	**6.77%**
Metabolism	**Amino acid metabolism**	**8.55%**	**13.67%**
Metabolism	**Amino acid metabolism**	**8.37%**	**11.61%**
Metabolism	**Amino acid metabolism**	**8.15%**	**6.81%**
Metabolism	Lipid metabolism	2.92%	3.63%
Metabolism	Glycan biosynthesis and metabolism	2.47%	2.45%
Metabolism	Metabolism of terpenoids and polyketides	2.43%	2.94%
Metabolism	Metabolism of other amino acids	2.31%	2.69%
Metabolism	Xenobiotics biodegradation and metabolism	1.55%	3.86%
Metabolism	Biosynthesis of other secondary metabolites	0.28%	0.80%
Genetic Information Processing	**Translation**	**11.62%**	4.22%
Genetic Information Processing	Replication and repair	8.15%	4.10%
Genetic Information Processing	Folding, sorting and degradation	4.24%	2.29%
Genetic Information Processing	Transcription	0.58%	0.20%
Environmental Information Processing	**Membrane transport**	**10.24%**	**11.32%**
Environmental Information Processing	Signal transduction	2.86%	**8.70%**
Cellular Processes	Cell growth and death	2.83%	1.67%
Cellular Processes	Cell motility	0.36%	2.63%
Cellular Processes	Transport and catabolism	0.24%	0.26%
Cellular Processes	Cellular community - prokaryotes	0.09%	0.67%

The percentage is the proportion of the abundance of a pathway in the abundance of all pathways level 2. The top pathways at level 2 in each species as well as their proportion were bolded.

Table 2 Samples of the peach fruit moth (PFM) *Carposina sasakii* and the oriental fruit moth (OFM) *Grapholita molesta* used in the study

Code	Species	Collecting location	Host	Coordinate	NO.
PKPR	PFM	Kaosanji of Pinggu district (K)	Pear (PR)	40°12′N, 117°19′E	5
OKPR	OFM				5
PLPR	PFM	Lvfulong of Yanqing district (L)	Pear (PR)	40°32′N, 116°4′E	5
OLPR	OFM				7
PGAE	PFM	Liugou of Yanqing district (G)	Apple (AE)	40°27′N, 116°6′E	4
OGAE	OFM				6
PDAE	PFM	Dafengying of Yanqing district (D)	Apple (AE)	40°26′N, 115°54′E	5
OYPH	OFM	Linguosuo of Haidian district (Y)	Peach shoot (PH)	39°58′N, 116°13′E	5
OSPH	OFM	Shuangxin of Haidian district (S)	Peach shoot (PH)	39°57′N, 116°12′E	2

All samples were collected from the Beijing area, China. NO., the number of individuals used for 16S rRNA gene sequencing.

Fig. 1 Microbial composition identified in the peach fruit moth (PFM) *Carposina sasakii* and the oriental fruit moth (OFM) *Grapholita molesta*. Community composition of the microbiome on phylum (a), genus (b), and species (c) levels for the OFM and PFM. (d) The cooccurrence relation graph describes the abundance of correspondence between samples and species. Each unit was represented by one color.

Fig. 2 Core bacteria of the same species from different hosts and different orchards. (a) Venn diagram at the genus level of PFM in four orchards. (b) Composition of 24 core genera found in all four orchard samples. (c) Venn diagram at the genus level of OFM from 5 orchards. (d) Composition of 33 core genera found in all five orchard samples. See Table 1 for the codes.

Fig. 3 Comparison of gut bacterial microbiota between the peach fruit moth (PFM) *Carposina sasakii* and the oriental fruit moth (OFM) *Grapholita molesta* individuals for alpha and beta diversity. (a and c) Community richness and diversity by Ace and Shannon index for the OTU level between the two species. (b and d) Wilcoxon rank-sum test of the difference between OFM and PFM individuals for ACE and Shannon indices (*p* > 0.05 is marked as NA, 0.01 < *p* ≤ 0.05 is marked as *, 0.001 < *p* ≤ 0.01 is marked as **, and *p* ≤ 0.001 is marked as ***). (e) Beta diversity of the microbiome between two species estimated by PCoA analysis at the genus level. PCo1 and PCo2 are the first two principle components, while the values on the x- and y-axis are proportions explained by corresponding components, respectively (PERMANOVA test with 999 permutations, *p* = 0.001, see Table S5 for values).

Fig. 4 Beta diversity of gut bacterial microbiota between the peach fruit moth (PFM)

Carposina sasakii and the oriental fruit moth (OFM) *Grapholita molesta* from the same host

plant and orchard. PERMANOVA was performed to determine the differences among groups.

(a) Sampled were collected from apple orchard estimated by PCoA analysis on the genus

level. (b) Sampled were collected from pear orchard. (c) Sampled were collected from

another orchard of pear (PERMANOVA test with 999 permutations, see Table S5 for values).

Fig. 5 Effects of environmental differences on beta diversity of gut microbiota in the peach fruit moth (PFM) *Carposina sasakii* and the oriental fruit moth (OFM) *Grapholita molesta* between different orchards in the same insect species and the same host species (PERMANOVA test with 999 permutations, Table S6).

Fig. 6 Functional predictions of gut microbiota in the peach fruit moth (PFM) *Carposina sasakii* and the oriental fruit moth (OFM) *Grapholita molesta*. (a) COG function classification; (b) KEGG pathway abundance of top 10 KEGG pathways at level 3 and the statistical difference between PFM and OFM ($p \leq 0.0001$ is marked as ****).