Infection with novel *Bacteroides phage BV01* alters host transcriptome and bile acid

metabolism in a common human gut microbe

Danielle E. Campbell[a#], Lindsey K. Ly[b,c], Jason M. Ridlon[b,c,d], Ansel Hsiao[e], Rachel J.

Whitaker[a,d], Patrick H. Degnan[a,e]

[a]Department of Microbiology, University of Illinois, Urbana, Illinois, USA

[b]Division of Nutritional Sciences, University of Illinois, Urbana, Illinois, USA

[c]Department of Animal Sciences, University of Illinois, Urbana, Illinois, USA

[d]Carl R. Woese Institute for Genomic Biology, University of Illinois, Urbana, Illinois,

USA

[e]Department of Microbiology and Plant Pathology, University of California, Riverside,

California, USA

Running Head: Prophage BV01 alters transcriptome and bile metabolism

[#]Address correspondence to Danielle E. Campbell, campbe28@illinois.edu

Abstract Word Count: 233

Text Word Count: 6552

ABSTRACT

The bacterial genus *Bacteroides* is among the most abundant and common taxa in the human gut, yet little is known about the phages infecting the group. *Bacteroides phage BV01* (BV01) was identified as a prophage integrated on the chromosome of its host, *Bacteroides vulgatus* ATCC 8482. Phage BV01 is actively produced, and infects susceptible *B. vulgatus* hosts in the mouse gut. Infection with BV01 causes a generalized repression of the *B. vulgatus* transcriptome, downregulating 103 transcripts and upregulating only 12. Integration of BV01 disrupts the promoter sequence of a downstream gene encoding a putative tryptophan-rich sensory protein (*tspO*). Deletion of *tspO* and subsequent RNAseq analysis revealed that more than half of the differentially-regulated transcripts are shared with the BV01 lysogen, suggesting the transcriptomic response to BV01 is linked to *tspO*. Among these differentially-regulated transcripts are two encoding bile salt hydrolases. Bile acid deconjugation assays show that BV01 represses its host's ability to hydrolyze bile acids in a *tspO*-dependent manner. Analysis of 256 published healthy human gut metagenomes suggests that phage integration adjacent to *B. vulgatus*-like *tspO* genes is rare within an individual, but common among humans. Finally, this work proposes a novel phage family that includes BV01, the *Salyersviridae*, whose host range spans the *Bacteroides* and is detectable in human-associated samples. Together, these findings highlight the importance of phage-host interactions to our understanding of how gut microbes sense and interact with their environment.

IMPORTANCE

The links between human disease and the gut microbiome are numerous. Most

mechanisms by which most gut microbes and their activities change and impact human

health remain elusive. Phages, viruses that infect bacteria, are hypothesized to play a

central role in modulating both community dynamics and functional activities. Here we

have characterized an active prophage, BV01, which infects a pervasive and abundant

human gut-associated species. BV01 infection alters its host's transcriptional profile

including its metabolism of bile acids, molecules implicated in mediating health and

disease states in the gut. This highlights that prophages and other components of the

variable genome should not be overlooked in bacterial genomes because they may

dramatically alter host phenotypes. Furthermore, BV01 represents a new family of

phages infecting human gut symbionts, providing a foundation for future investigations

of phage-host interactions in these clinically-relevant but underexplored hosts.

INTRODUCTION

The human gut is colonized by a dense and diverse microbial community

comprised of bacterial, archaeal, and fungal cells, as well as the viruses that infect them.

This gut microbiome is vital to human health and development, and is linked to an

increasingly long list of disease states. Recent work has specifically implicated the gut

phageome in disease, including inflammatory bowel disease, malnutrition, AIDS,

colorectal cancers, and hypertension (1–5). Broadly, gut phages act as important

modulators of bacterial community structure (4, 6, 7) and metabolism (8). Despite their

apparent importance, little is known about how most gut-associated phages interact with

their bacterial hosts.

The *Bacteroides* is one of the most common and abundant bacterial genera in the

distal human gut. The genus is known to degrade a diversity of complex carbohydrates

(9, 10), and interact with host immune cells (11, 12). Within a single human host, many

Bacteroides species and strains coexist, competing for nutrients under changing

environmental conditions caused by host diet (13), host metabolites (14), host immune

system activities (15, 16), and phage predation (17). Moreover, horizontal gene transfer

plays an important role in shaping the evolution and function of *Bacteroides* genomes

(18, 19). How the diversity of *Bacteroides* strains in the human gut persists over time in

such a dense, dynamic, and competitive environment is likely multi-fold, and perhaps

afforded by their highly plastic genomes.

Phage diversity and phage-host interactions within most commensal gut-

associated bacteria, including *Bacteroides* species, is underexplored. Currently, the most

abundant gut-associated phages are the crAssphages (20, 21), a group of related lytic

phages that infect *Bacteroides intestinalis* and potentially other species. CrAssphages

demonstrate how traditional phage techniques (*e.g.,* agar overlay plaque assays) are not

reliable for *Bacteroides* hosts (22), likely due to heterogeneity in capsular polysaccharide

composition within isogenic cultures (23, 24). In fact, deletion of all capsular

polysaccharide synthesis loci allows for the isolation of many phages on the host

Bacteroides thetaiotaomicron VPI-5482 (17). Most phages isolated against *Bacteroides*

hosts thus far exhibit an obligately lytic lifestyle (17, 22, 25, 26), despite the potentially

large role of lysogeny in phage-host interactions in the gut, where at least 17% of the gut

phageome is predicted to be temperate (27, 28).

Prophage–host interactions have the potential to cause complex alterations to the

host phenotype by virtue of the temperate lifestyle having two distinct phases: lysogeny

and lysis. Unlike most strictly lytic phages, temperate phages may horizontally transfer

beneficial genes between hosts, such as antibiotic resistance genes (29) and auxiliary

metabolic genes (30, 31). Some phage regulatory machinery expressed from prophages

can modulate the transcription of host genes, resulting in altered phenotypes (32, 33).

Integration of prophages into the host genome may also disrupt or enhance the activity of

surrounding chromosomal genes (34–36). While much is known about how these

prophage–host interactions contribute to virulence in pathogens (37–41), exceedingly

little is about how temperate phages modulate the activities of commensals.

Here we have identified an active prophage, *Bacteroides phage BV01,* in a

genetically tractable host strain, *B. vulgatus* ATCC 8482, and characterized its effects on

the host's transcriptome and phenotype. Further we determine that BV01 represents a

larger group of *Bacteroides*-associated prophages comprising the proposed phage family

Salyersviridae, which are common members of the human gut phageome. This work

provides the first insights to how the *Bacteroides* react to temperate phage infection, and

establishes a model system for exploring complex phage–host interactions in an

important human gut symbiont.

RESULTS

***Bacteroides phage BV01* is a prophage in *B. vulgatus* ATCC 8482.** *Bacteroides*

phage BV01 was first partially predicted with ProPhinder and deposited in the ACLAME

database (42, 43). Through comparative genomics and annotation of this region, we

extended the predicted BV01 prophage to 58.9 Kb (NC_009614.1: 3,579,765..3,638,687),

comprised of 72 predicted ORFs (Fig. 1), a prediction that agrees with an observation of

prophage induction from *B. vulgatus* ATCC 8482 in a gnotobiotic mouse model (44).

BV01 encodes genes suggesting a temperate lifestyle, with a putative CI-like repressor

and a Cro-like anti-repressor, as well as a holing-lysin-spanin operon (Fig. 1).

BV01 is detectable outside of host cells in the supernatants of *in vitro* cultures as

a DNase-protected, dsDNA genome by both sequencing (Fig. 1) and PCR (Fig. 2A). An

isogenic cured lysogen (ΔBV01) strain constructed by replacement with the

corresponding chromosomal region of an uninfected *B. vulgatus* strain (*attB*) does not

release free BV01 (Fig. 1, 2A). Assembly of sequencing reads from free BV01 phage

DNA results in a circular contig that spans the phage attachment site (*attP*). The BV01

attP is identical to the left and right attachment sites (*attL* and *attR*), a pair of 25-bp direct

repeat sequences (5'-GTCTAGTTTAGTTTTTGTGTTGTAA-3'), suggesting BV01

enters a circular intermediate before replication.

To confirm that the release of DNase-protected BV01 genomes from host cells is

a phage-encoded and directed process, we sought to identify the phage integrase. BV01

encodes three genes with integrase domains (PF00589); we hypothesized that the gene

BVU_RS14130, adjacent to the phage attachment site, was the most likely candidate for

catalyzing integration and excision of BV01. A BVU_RS14130 deletion mutant (Δ*int*)

was constructed (Fig. 1) and its activity assayed by PCR of paired cell pellets and DNase-

treated culture supernatants (Fig. 2A). Phage DNA was not detected in the supernatants

of the Δ*int* strain. Furthermore, the Δ*int* mutant does not yield an amplicon for the

circularization of the BV01. Expression of the *int* gene *in trans* from a pNBU2 plasmid

complements both the circularization and release phenotypes (Fig. 2A). These results

demonstrate that the integrase encoded by BVU_RS14130 is necessary for phage

excision, circularization, and release from the host. Furthermore, they suggest that BV01

is an intact prophage capable of directing its own mobilization.

Despite numerous attempts, we have not identified *in vitro* inducing conditions

for BV01. This has included treatments with UV light and sub-inhibitory concentrations

of norfloxacin and mitomycin C (data not shown). BV01 supernatants never produced a

plaque on any of 10 *B. vulgatus* isolates tested, the *B. vulgatus* cured lysogen, *B.*

thetaiotaomicron VPI-5482, or *Bacteroides dorei* DSM 17855, on any of four media:

TYG, BHI-HB, BHI-HM, and TYG$_S$. To test for new lysogenic infections, BV01 was

tagged with a copy of *tetQ*, conferring tetracycline resistance, using allelic exchange.

Transduction of BV01 or BV01-*tetQ* via culture supernatants or in co-culture with

erythromycin resistant hosts tagged with pNBU2-*bla*-*ermG*$_b$ has never generated

transductants. We conclude that BV01 is a latent prophage in culture, which is further

supported by transcriptional data showing that BV01 exists in a largely repressed state

(Fig. S1).

The only infectious conditions that have been identified for BV01 are in a

gnotobiotic mouse model (Fig. 2B). Within a single day of gavage, BV01-*tetQ*

transductants were identified on doubly-selective media from mouse pellets from 4 of the

7 mice. Over the course of the 11-day experiment transductants were eventually observed

in all animals, in all cages. The average frequency of transduction ranged from 1.9 x 10^{-6}

to 3.6 x 10^{-9} per animal. These results support the hypothesis that an unknown

mammalian host factor is required for novel BV01 infection.

Lysogeny with BV01 alters the host transcriptome. It was hypothesized that

lysogeny with a prophage such as BV01 could alter the activities of the *B. vulgatus* host.

RNA sequencing (RNAseq) of the *B. vulgatus* wild-type lysogen and the isogenic cured

lysogeny was performed to identify transcripts differentially regulated in response to

BV01 lysogeny (Fig. 3).

Analysis of RNAseq data revealed 115 host transcripts differentially regulated in

response to lysogeny with BV01 (Fig. 3A), 103 of which (89%) are up-regulated in the

cured lysogen (Table S1). These transcriptional changes occur across the host genome

(Fig. 1B). Functional analysis of these transcripts revealed that most function in

metabolism and cellular processes and signaling (Fig. 3C). Pathway analysis using the

Kyoto Encyclopedia of Genes and Genomes Pathway Database (45), however, failed to

yield pathway-level differences, which is likely a reflection of the level of annotation of

the *B. vulgatus* genome. Taken together, these results indicate BV01 represses a diverse

array of its host's metabolic activities, suggesting it is acting through one or more

transcriptional regulators.

One possible explanation for the widespread transcriptomic response to BV01

lysogeny is that a phage product directly alters the transcriptional activity of host genes.

BV01 encodes two candidate genes that might act in this way: a predicted transcriptional

regulator (BVU_RS14475) and a predicted sigma factor-like protein (BVU_RS14235)

(Fig. S1). The transcriptional regulator encoded by BVU_RS14475 is the most highly

transcribed gene in the BV01 prophage, encoding the putative CI-like repressor protein,

which might directly interact with host promoters. Transcription of BVU_RS14235 is

very low, so it is less likely to play a major role in the observed transcriptional response

(Fig. S1). It is also possible that the observed transcriptional response to BV01 is the

result of a host response to infection. A universal stress protein (*uspA*) homolog

(BVU_RS16570) is upregulated in the BV01 lysogen, though it is not clear if that is a

primary or secondary effect of infection.

BV01 alters bile acid metabolism by disrupting the *tspO* promoter. Notably,

integration of BV01 at the *attB* is correlated with a 23-fold down-regulation of the

adjacent downstream transcript (BVU_RS14490), through an apparent disruption of the

gene's promoter (Fig. 4A). A low level of expression at this gene is observed in the wild-

type lysogen, perhaps a result of readthrough from phage transcripts. This gene encodes a

predicted tryptophan rich sensory protein (TspO) homolog (Fig. 4B), an intramembrane

protein whose endogenous ligand is unknown, but which is broadly implicated in

metabolic regulation and stress response in other bacteria (46–48). Although TspO is

conserved in many bacteria, archaea, and eukaryotes, it is considered an accessory

protein. Indeed, not all gut-associated members of the family Bacteroidales nor the genus

Bacteroides encode *tspO* (Fig. 4C). Within the *Bacteroides*, *tspO* is restricted to the clade

including *B. vulgatus* and *B. dorei*. Among *B. vulgatus* strains TspO is highly conserved

(Fig. 4D), suggesting it plays a specialized role in regulating the cellular activities unique

to *B. vulgatus*.

Given TspO's important role in regulating cellular activities in other bacterial

systems, it was hypothesized that it may be responsible for some of the differential

regulation observed in response to prophage BV01. A *tspO* deletion mutant was

constructed in the cured lysogen background (ΔBV01Δ*tspO*) and its transcriptome

sequenced alongside that of the wild-type and cured lysogen strains (Fig. 3). The

predicted *tspO* regulon extends far beyond the differential expression observed in the

cured lysogen (Fig. 3A, 3B), suggesting the small amount of *tspO* transcription in the

BV01 lysogen exerts effects on the rest of the genome. Transcripts differentially

regulated between the BV01 wild-type lysogen and cured lysogen which returned to wild

type-like levels upon further deletion of *tspO* were classified as *tspO*-dependent

transcripts (Fig. 3C). Of the 115 transcripts differentially regulated in response to BV01,

69 (60%) are *tspO*-dependent. Consistent with TspO's role in regulating stress, many

tspO-dependent transcripts fall into the COG category for post-translation modification,

protein turnover, and chaperones, including several thioredoxins, peroxidases, and

protein chaperones. *tspO*-dependent transcripts also account for the majority of metabolic

genes differentially regulated in response to BV01.

Two *tspO*-dependent transcripts that are down-regulated in response to lysogeny

with BV01 encode putative bile salt hydrolases (BVU_RS13575, 5.38-fold change, $q <$

10^{-100}; BVU_RS20010, 6.58-fold change, $q < 10^{-200}$). It was hypothesized these

transcriptional differences might reflect enzyme activities. To this end, *B. vulgatus* strains

were grown in the presence of bile acids and the deconjugation of those bile acids

measured by LC/MS (Fig. 5).

LC/MS results show that the wild-type *B. vulgatus* lysogen does not significantly

deconjugate glycocholic acid (GCA) to cholic acid (CA), and may exhibit modest

deconjugation of taurocholic acid (TCA) (Fig. 5). This agrees with a previous study that

showed *B. vulgatus* ATCC 8482 can deconjugate TCA but cannot deconjugate GCA over

a 48 hr incubation (49). Importantly, CA is clearly detectable only in the cured lysogen

background, consistent with our prediction based on RNAseq data. This bile acid

deconjugation phenotype is ablated with the deletion of *tspO*, further supporting the hypothesis that *tspO* activates transcription of bile salt hydrolases, resulting in their increased enzymatic activity.

To see if *tspO*-disrupted lysogens occur in natural human gut microbiomes, read mapping from 256 healthy human gut metagenomes was performed. Starting with reads which mapped to *tspO* in the reverse orientation, read mates were checked for mapping to Phage BV01 and its relatives (Fig. 6). All samples had reads mapping to *tspO*, with an average of 0.0004% of metagenome reads mapping, indicating the corresponding population of *B. vulgatus* and *B. dorei* encoding *tspO* is relatively abundant (Fig. S2A). Incidence of *tspO* associated with BV01 or a related phage is also common; 13.3% of samples (*n*=34) contained read pairs mapping to *tspO* and an adjacent prophage. Within an individual microbiome, incidence of *tspO*-disrupted lysogens appears rare, usually comprising 3% or less of the combined *B. vulgatus* and *B. dorei* population, although for some individuals this incidence rate can be more than 10% (Fig. S2B). In addition to these patterns, carriage of *tspO*-disrupted lysogens appears stable over time, as indicated by individuals sampled at multiple time points (Table S2). Together, these data suggest that BV01 and other phages' effects on downstream phenotypes via *tspO* are likely quite common among humans, though these lysogens comprise the minority of the overall microbiome.

BV01 represents the proposed viral family *Salyersviridae*. While searching for potential new hosts for BV01, its predicted 25-bp *attB* was queried against 154 gut-associated Bacteroidales genomes. It was found that all *Paraprevotella* and *Bacteroides* genomes had at least the first 21 bases of the attachment site conserved (Fig. 6A). In 20

genomes, two copies of the *att* site were found, and all are associated with putative

prophages. In all instances, the putative *attL* and *attR* sites are direct repeats, as is true for

BV01. Importantly, only *B. vulgatus* lysogens encode *tspO* (Fig. 4); other prophages and

attB sites occur in an alternative genomic context. Alignment of these putative prophage-

associated *att* sites finds that the first 22 basepairs are always conserved, and an

additional 3 basepairs are variable among lysogen genomes (Fig. 6B).

   Integration at the same *att* site suggests these prophages are genetically related.

To assess relatedness, VICTOR (50) was used to build a genome tree from all of the

identified prophages (Fig. 6C) and OPTSIL clustering (51) was used to predict taxonomic

groups, which we named *Salyersviridae*, *Salyersvirinae*, and *Salyersvirus* (Table 1).

Taxonomic clustering also defined phage species, four of which have more than one

member. Interestingly, the phage species BX01 has representatives in two distantly

related hosts, *Bacteroides xylanisolvens* XB1A (phage BX01) and *B. thetaiotaomicron*

1_1_6 (phage BT01), suggesting that at least some Salyersviruses have broad host ranges.

Phage BC01 was placed outside the subfamily *Salyersvirinae*, which is consistent with its

considerable sequence divergence (Fig. S3). Further examination of the BC01 *attL* and

attR sites found that the 22-bp consensus sequence is only a portion of the full 69-bp

repeat flanking the BC01 prophage, which further supports placement of BC01 outside

the *Salyersvirinae*. Also included in the analysis were three outgroup sequences from the

known *Bacteroides*-infecting lytic phages: B124-14, B40-8, and crAssphage.

Surprisingly, OPTSIL taxonomic clustering placed phages B124-14 and B40-8 in the

family *Salyersviridae*. This similarity is not detectable at the level of nucleotide sequence

(Fig. S3), but is likely driven by similarities in several proteins, including homologs to

the predicted lysin, excisionase, and ssDNA binding protein from BV01.

To check that the proposed *Salyersviridae* clade was comprised of active phages,

we searched for evidence of activity in culture and in wastewater samples. Sequencing

and PCR were used to test the activity of six additional Salyersviruses (Fig. S4). Phages

BV02, BV05, and BV06 were found to be released from host cells in culture, while BV03

is likely inactivated by genomic rearrangement. We did not observe activity for BV04 or

BO03 from *B. ovatus* ATCC 8483, both of which appear complete based on synteny with

intact Salyersviruses. This may be the result of inactivating mutations or not being

induced in the growth conditions tested (Fig. S4D).

Furthermore, we used read mapping to search for evidence of all 20

Salyersviridae phages in a wastewater metavirome (Fig. S5A). Reads mapped to all

Salyersviridae genomes, however it may be the result of sequence conservation within

the family, as reads often accumulate at the most conserved regions of each phage

genome. Further, very little read mapping occurred in the distal portion of the inactivated

prophage BV04, which resembles host chromosome more than phage sequence,

indicating that virome processing removed most cellular DNA prior to sequencing (Fig.

S5A). *De novo* assembly of the individual wastewater viromes finds contigs which align

with high identity but imperfectly to each *Salyersviridae* phage, supporting the findings

seen by read mapping, and suggesting the real diversity of the family *Salyersviridae* is far

greater than what has been observed integrated in cultured host genomes so far. This

analysis found that phages infecting *B. vulgatus* are more abundant than other

Salyersviridae phages based on maximum normalized read coverages. A similar

comparison concludes that most individual temperate *Salyersviridae* phages are

approximately equal in abundance to the lytic *Salyersviridae* phage B124-14 and B40-8,

and approximately ten-fold less abundant than crAssphage, in these wastewater viromes

(Fig. S5A).   Searches for *Salyersviridae* phages in individual healthy human fecal

metagenomes refine this conclusion, showing that in most human samples, *Salyersviridae*

phages are at least as abundant as crAssphage, though crAssphage can reach very high

abundances in a subset of individuals (Fig. S5B). Although confirmation of most

Salyersviridae activities will require better sampling and *in vitro* testing, these results

indicate the phage family is active in human-associated communities.

DISCUSSION

309   Here, we characterize a complex phage-host interaction between *Bacteroides*

phage BV01 and its host *B. vulgatus*. We first demonstrate that BV01 is an intact

prophage capable of directing its own excision, and it is transducible *in vivo* in a

gnotobiotic mouse model. Using a combination of genetics, RNAseq, and analytical

chemistry, we show that BV01 decreases its host's ability to deconjugate bile acids by

disrupting the transcription of the gene adjacent to the *attB* encoding a TspO homolog.

Furthermore, we show that *tspO* disruption by phage integration is common among, but

rare within, healthy human gut microbiomes, and can be mediated by BV01 or its

relatives. Together, these findings elucidate a complex mechanism by which a phage

alters its host's activities.

319   The repression of bile acid deconjugation as a consequence of BV01 integration is

particularly relevant in the context of the mammalian gut. Mammals secrete conjugated

primary bile acids into the small intestine, where they reach concentrations as high as 1

mM (52); though the majority of bile acids secreted into the small intestine are

readsorbed, they can still accumulate to concentrations of 0.2-1 mM in the colon (53).

While bile acids are broadly capable of damaging lipid membranes, generally

Bacteroides species are considered bile-resistant (54), the mechanism of which is

unknown. Bile acid deconjugation is a common activity encoded by gut-associated

microbes, though its direct benefit to those microbes is unclear. Microbial modification of

the bile acid pool can be linked to beneficial changes in the human host metabolism (49,

55) and varied epithelial susceptibility to viral pathogens (56). The link between BV01

and bile acid metabolism suggests a heretofore undescribed mechanism by which gut

phages might influence mammalian host phenotypes.

Here, bile acid deconjugation in *B. vulgatus* is dependent on a putative TspO.

Bacterial TspOs are important for regulating metabolic switches and stress regulation in

at least three diverse systems (46–48), though the mechanism of action for the protein is

unknown. The crystal structure of TspO shows a periplasm-facing binding pocket distinct

from the intramembrane cholesterol recognition consensus sequence, which may bind or

degrade porphyrins (57–59). Both the porphyrin degrading and cholesterol transporting

functions of TspO, however, have been disputed (60, 61). Despite this, it is notable that

cholesterol is structurally similar to bile acids, being their biosynthetic precursor. In at

least one other gut-associated microbe, *tspO* is up-regulated by bile acids, suggesting

TspO may be involved in bile acid metabolism in gut microbes more broadly (62). The

regulatory link described here between bile acid hydrolysis and TspO suggests a

hypothesis where the *B. vulgatus* TspO might be a sensor and regulator of bile acid

interactions.

Induction of BV01 from its integrated state and infection of new hosts remains

enigmatic. Prophage induction is canonically linked to stress-dependent pathways, as is

the case for lambdoid phages that respond to DNA damage via RecA-dependent cleavage

of the CI repressor protein (63). It is possible that prophages in *Bacteroides* hosts respond

to alternative stimuli, as is the case for CTnDOT, a well-studied *Bacteroides* conjugative

transposon, whose excision is inducible only by teteracycline (64). Neither DNA damage

nor antibiotics induce prophage BV01 *in vitro*, so all experiments here relied on an

apparently low rate of spontaneous prophage induction. Similarly, no infecting conditions

or susceptible hosts have been identified for BV01 *in vitro*. We demonstrate that BV01 is

transducible in a gnotobiotic mouse model, suggesting that an unknown mammalian host

factor is required for novel BV01 infection. Enigmatic infection dynamics may be the

result of the phase variable polysaccharide capsule, as recent work suggests heterogeneity

in capsular composition hinders phage infection on population-scales (17). Indeed, it has

long been observed that finding phages in the *Bacteroides* using traditional techniques is

difficult or impossible for most host strains (65, 66), making the host-first approach to

phage discovery used here especially appealing.

Finally, phage BV01 is the first representative of a broad family of phages that

spans an entire host genus, and includes lytic and temperate members. *Salyersviridae* is

common and diverse among natural human samples, but rare within individuals,

suggesting lysogenization may confer frequency-dependent advantages to the bacterial

host. The genetic context of non-*B. vulgatus Salyersviridae* lysogens remains unexplored,

providing ample opportunity for further discovery of novel phage-host interactions. The

absence of *tspO* in these other host systems may provide the ideal background for

studying more direct impacts of these phages on their hosts. Certainly, other interactions

between BV01 and its host remain to be studied, though they were overshadowed here by

the enormous effects of *tspO*. Future studies should also examine the role of

Salyersviridae phages on bacterial host fitness and evolution (67, 68), as these phages

directly impact their bacterial hosts and those interactions likely have important ripple

effects throughout the microbiome and on the mammalian host that remain to be

elucidated.

MATERIALS AND METHODS

Strains and culture conditions. All strains and plasmids used in the study are

listed in Table S3. *Escherichia coli* S17-1 *λ pir* was used for all routine recombinant

DNA cloning, and grown aerobically in Lysogeny Broth (LB) at 37°C. *B. vulgatus* strains

were cultured anaerobically in a vinyl anaerobic chamber using 70% N_2, 20% CO_2, and

10% H_2 gas mixture (Coy Laboratory Products, Grass Lake, MI). All *B. vulgatus* cultures

were grown on Difco Brain Heart Infusion (BHI) agar supplemented with 10%

defibrinated horse blood (BHI-HB; Quad Five, Ryegate, MT), or in tryptone-yeast

extract-glucose (TYG) broth (69) at 37°C. When necessary, ampicillin (100 μg/mL),

gentamicin (200 μg/mL), erythromycin (25 μg/mL), 5'-flourodeoxyuridine (FUdR; 20

μg/mL), or tetracycline (2 μg/mL) were supplemented in the media. Infection assays were

performed on BHI supplemented with 50 μg/mL hemin and 0.5 μg/mL menadione (BHI-

HM) and TYG_S (70).

Genetic manipulation. All primers used to construct genetic mutants are listed in

Table S3. Markerless deletion mutants in *B. vulgatus* were achieved by allelic exchange

using a system analogous to that developed in *B. thetaiotaomicron,* (71) and confirmed

by PCR and whole genome sequencing. The *tdk* gene (BVU_RS09305), encoding

thymidine kinase, was deleted from *B. vulgatus* ATCC 8482 by allelic exchange,

conferring resistance to the toxic nucleotide analog FUdR. Cloning was performed as

described by Degnan *et al.* (72). Briefly, the 3.5 Kb regions flanking either side of *tdk*

were amplified with Kapa HiFi Taq MasterMix (Kapa Biosystems, Wilmington, MA)

and joined by splicing overlap exchange (SOE) PCR. The SOE product was purified,

restriction digested and ligated into the suicide vector pKNOCK-*bla-ermG*b in *E. coli*,

and conjugated into *B. vulgatus.* Single recombinant merodiploids were selected for on

BHI-HB supplemented with gentamicin and erythromycin, and double recombinant

deletion mutants subsequently selected for on BHI-HB with FUdR. The counterselectable

suicide vector pExchange-*tdk*BV was constructed by amplifying *tdk* from *B. vulgatus* and

cloning the *tdk* amplicon into pKNOCK-bla-ermG$_b$ by the same methods used to clone

the SOE product above.

Subsequent deletions were accomplished similarly as described for *tdk*, except

using pExchange-*tdk*BV and flanking regions of ~1 Kb to create the deletion alleles

(*tspO, int*). For deletion of the entire BV01 provirus, an empty attachment site (*attB*) and

the flanking 800 bp were cloned from *B. vulgatus* VPI-4506, which has 99.9% nucleotide

identity to the analogous regions flanking BV01 in *B. vulgatus* ATCC 8482.

Complementation of the BV01 integrase (BVU_RS14130) was accomplished by

cloning the gene and its native promoter into the integrative plasmid pNBU2-*bla-ermG*b,

which has a single integration site in the *B. vulgatus* genome (*attN*;

NC_009614.1:3152550..3152572). This construct was conjugated into *B. vulgatus* and

transconjugants selected for on BHI-HB supplemented with gentamicin and erythromycin

as described elsewhere (72).

The BV01-*tetQ* strain was constructed by inserting *tetQ* from pNBU2-*bla-tetQ*

immediately downstream of the stop codon of BVU_RS14265, upstream of a predicted

transcriptional terminator. As was done for deletion constructs, the desired region was

constructed on the pExchange-*tdk*BV plasmid and moved into the wild-type *B. vulgatus*

strain by allelic exchange. First, a ~2 Kb region surrounding the BVU_RS14265 stop

codon was amplified in two pieces with SOE primers designed to insert adjacent SpeI and

BamHI cut sites downstream of the stop codon and ligated into pExchange-*tdk*BV. This

construct was confirmed by Sanger sequencing before *tetQ* and its promoter were

amplified from CTnDOT, and ligated into the SpeI and BamHI cut sites. Tetracycline

was used to select for mutants, and release of BV01-*tetQ* phages confirmed by PCR.

Select mutant strains were confirmed by whole genome sequencing and analyzed

with Breseq (73) aligned to the wild-type *B. vulgatus* ATCC 8482 genome

(NC_009614.1), and summarized in Table S4.

Genome sequencing. Cells were pelleted from 5 mL overnight culture in TYG by

centrifugation at 4,000 × *g* for 5 min at 4°C, resuspended in 0.5 mL TE buffer (10 mM

Tris, 1 mM EDTA), and lysed by adding sodium dodecyl sulfate (SDS) and proteinase K

(GoldBio, Olivette, MO) to final concentrations of 0.07% and 300 μg/mL, respectively,

and incubating for 2 hr at 55°C. Cellular material was removed by washing twice in an

equal volume of buffered phenol, phenol-chloroform-isoamyl alcohol (VWR, Radnor,

PA), and DNA precipitated with 100% ethanol in the presence of 0.3M sodium acetate at

-20°C overnight. DNA pellets were washed with 70% ethanol, dried, and resuspended in

TE buffer.

Phage DNA was prepared from overnight TYG culture supernatants collected

after centrifugation and concentrated by centrifugation with 30,000 MWCO Corning

Spin-X UF 20 Concentrators (Corning, NY) or by tangential flow filtration with a

Vivaflow 50R 30,000 MWCO Hydrosart filter (Sartorius, Gottingen, Germany).

Supernatants were treated with 200 μg/mL DNase I and 1 μg/mL RNase A for 1 hr at

room temperature to remove unprotected DNA and RNA. Virions were disrupted with

1% SDS and 1 mg/mL proteinase K for 2 hr at 55°C. DNA was further isolated using the

same phenol-chloroform-isoamyl alcohol extraction and ethanol precipitation procedures

as for cellular DNA.

DNA libraries were constructed with the Nextera XT Library Preparation Kit and

Index Kit (Illumina, San Diego, CA). DNA libraries were pooled and sequenced on both

the Illumina HiSeq5000 and HiSeq2500 and fastq files were generated from

demultiplexed reads with bcl2fastq Conversion Software (Illumina, San Diego, CA).

Reads were trimmed and assembled using the A5ud pipeline (74). Sequencing methods

and assembly data are summarized in Table S5.

Genome annotation. Annotation of cellular genomes was accomplished with a

custom Perl script that calls protein coding genes with Prodigal (75) and RNA coding

genes with tRNAscan-SE (77), Rnammer (78) and Infernal (78). Functional predictions

are assigned by searching against Kyoto Encyclopedia of Genes and Genomes (45),

Cluster of Orthologous Genes (80), Pfam (80), and TIGRFAM (81) databases, and

subCELlular LOcalization predictor (82) is used to predict cellular localization.

459   For phage genomes, genes were called by Prodigal and the gene calling tool

460   within Artemis (83). Functional predictions were made as above except with relaxed

461   search parameters (cut_tc in hmmscan), plus using Basic Local Search Alignment Tool

462   (84) with the Genbank virus database (85), Phyre2 (86) to identify conserved protein

463   folds, and iVireons (87) to predict structural proteins, and manually comparing and

464   combining results.

465   **Integrase activity assays.** Integrase activity was assayed through PCR of DNase-

treated supernatant DNA. Briefly, free phage DNA was prepared as for DNA sequencing,

and amplified with Kapa HiFi Taq MasterMix with primers specific to BV01

(BVU_RS14350) or spanning the circularized *attP* (Table S3). Free phage DNAs were

checked for the presence of contaminating cellular DNA by amplifying the 16S rRNA

gene with universal primers. Amplicons were cleaned with a Qiagen PCR Cleanup kit

(Hilden, Germany) and run on an agarose gel in 0.5X Tris-borate-EDTA buffer at 70V

alongside 1 Kb ladder (New England BioLabs, Ipswich, MA) or GeneRule Express DNA

Ladder (Thermo Scientific, Waltham, MA) and stained with GelRed (VWR, Radnor,

PA). Amplicons generated with *attP*-flanking primers were sequence confirmed by

Sanger sequencing performed by ACGT, Inc (Wheeling, IL).

476   **Gnotobiotic mice.** All experiments using mice were performed using protocols

approved by the University of California Riverside Institutional Animal Care and Use

Committee. Germfree C57BL/6J mice were maintained in flexible plastic gnotobiotic

isolators with a 12-hr light/dark cycle. Animals caged individually (*n*=1, female) or in

pairs of litter mates (*n*=6, males) were provided with standard, autoclaved mouse chow

(5K67 LabDiet, Purina) *ad libitum*. With no *a priori* reason to expect age to influence

transduction rates, animals ranged from 7 weeks to nearly 12 months old. Individually

antibiotic resistance marked bacterial strains were grown individually for ~20h in TYG

medium with appropriate antibiotics and frozen at -80ºC in anaerobic cryovials. Cell

viability was tested by plating and viable CFU counts were used to combine equal parts

of the wild-type *B. vulgatus* lysogen tagged with pNBU2-*bla-ermGb* and *B. vulgatus*

BV01-*tetQ* tagged with pNBU2-*bla-cfx*. Approximately 4 x 10^6 CFUs of the combined

strains were administered to each animal by oral gavage. Fecal samples were collected on

days 1, 3, 7 and 11 from each animal. Fecal pellets were processed by adding 500µl of

TYG+20% Glycerol to each tube and vigorously shaking in a beadbeater without beads

for 1m 30s. Fecal slurries were spun down at 2,000 x g for 1 s, followed by serial dilution

on selective media (BHI+Tet+Gn, BHI+Erm+Gn, BHI+Erm+Tet+Gn) to determine

CFUs. Animals were sacrificed on d11 following final fecal collection.

Transcriptomic response to lysogeny. *B. vulgatus* was grown overnight in 5 mL

TYG medium. Each culture was pelleted (4,000 × g for 5 min at 4°C), supernatant

decanted, and washed in an equal volume of TYG three times. Cells were normalized to

an OD_{600} of ~0.3 and used to inoculate cultures in 10 mL TYG at a final dilution of

1:1000 in biological triplicate. Cell growth was monitored and cells were harvested at an

OD_{600} of ~0.4. Total RNA was prepared with a Qiagen RNeasy kit (Hilden, Germany)

and treated on-column with RNase-free DNase (Qiagen, Hilden, Germany). RNA was

quantitated with a Qubit 2.0 fluorometer (Thermo Fisher, Waltham, MA) and stored at -

80°C.

RNA was submitted to the W. M. Keck Center for Comparative and Functional

Genomics at the University of Illinois at Urbana-Champaign for quality analysis, rRNA

depletion with the RiboZero Bacteria kit (Illumina, San Diego, CA), library construction

with the TruSeq Stranded mRNAseq Sample Prep kit (Illumina, San Diego, CA), and

sequencing on an Illumina NovaSeq 6000 with the NovaSeq S4 reagent kit. Fastq files of

demultiplexed reads were prepared with the bcl2fastq v2.20 Conversion Software

(Illumina, San Diego, CA).

RNAseq reads were quality filtered and trimmed with trim_galore v0.4.4

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/). Rockhopper (88)

was used to identify differentially expressed transcripts between isogenic mutants ($\geq$2-

fold change, $q \leq 0.01$).

Bile salt deconjugation assay & LC/MS. *B. vulgatus* strains were inoculated in

TYG liquid supplemented with 50 μM glycocholic acid (GCA) or 50 μM taurocholic

acid (TCA) and allowed to grow for 16 or 28 hr, respectively. Grown cultures were

brought to a pH 2.0-3.0 with 10 N hydrochloric acid, centrifuged for 5 min at 4,000 x *g*,

and the pellets discarded. Bile acids were isolated by solid phase extraction over Sep-Pak

tC18 500 mg cartridges (Waters Corp., Milford, MA). Cartridges were preconditioned by

serial washes with 6 mL hexane, 3 mL acetone, 6 mL methanol, and 6 mL water (pH =

3.0). Acidified supernatants were loaded before washing with 3 mL 40% methanol. The

column was allowed to dry, then bile acids eluted in 3 mL methanol. Samples were

evaporated under nitrogen, resuspended in 20 μL methanol, and centrifuged before

analysis by liquid chromatography-mass spectroscopy (LC/MS).

LC/MS for all samples was performed on a Waters Aquity UPLC coupled with a

Waters Synapt G2-Si ESI MS. Chromatography was performed using a Waters Cortecs

UPLC C18 column (1.6 μm particle size) (2.5 mm x 50 mm) with a column temperature

of 40° C. Samples were injected at 1 μl. Solvent A consisted of 95% water, 5%

acetonitrile, and 0.1% formic acid. Solvent B consisted of 95% acetonitrile, 5% water,

and 0.1% formic acid. The initial mobile phase was 90% Solvent A, 10% Solvent B and

increased linearly until the gradient reached 50% Solvent A and 50% Solvent B at 7.5

min. Solvent B was increased linearly again until it was briefly 100% at 8.0 min until

returning to the initial mobile phase (90% Solvent A, 10% Solvent B) over the next 2

min. The total run was 10 min with a flow rate of 10 μL/min. MS was performed in

negative ion mode. Nebulizer gas pressure was maintained at 400° C and gas flow was

800 L/hour. The capillary voltage was set at 2,000 V in negative mode. MassLynx was

used to analyze chromatographs and mass spectrometry data.

Taxonomic nomenclature. The family, subfamily, and generic names were

chosen to honor the microbiologist Abigail A. Salyers, who made significant

contributions to the understanding of function and genetics of human gut anaerobes and

the importance of their mobile genetic elements.

Wastewater collection, processing, and viromics. From the Urbana &

Champaign Sanitary District Northeast Plant (Urbana, IL), 1 L of unprocessed

wastewater was collected at each of three time points: May 25, 2016, June 23, 2016, and

October 3, 2016.

Wastewater samples were transported on ice, immediately centrifuged at 2,500 ×

g for 10 min at 4°C and filtered through a 0.4 μm polyethersulfone filter to remove large

particulate and cellular matter. The sample was split into three aliquots and processed

three ways. One aliquot was not processed further (F). Another aliquot was filtered a

second time through a 0.22 μm polyethersulfone filter (DF). The last aliquot was washed

three times with an equal volume of chloroform (FC). All aliquots were concentrated

100-fold and virome DNA was isolated from each as described for genome sequencing of

phages.

DNA libraries of virome DNA were prepared using the same methods as

described for genome sequencing and were sequenced on an Illumina HiSeq 2500

sequencer with a HiSeq v4 SBS sequencing kit (Illumina, San Diego, CA) producing

2×160-bp paired-end reads. Fastq files of demultiplexed reads were generated with the

bcl2fastq v2.17.1.14 Conversion Sotfware (Illumina, San Diego, CA). Reads were

trimmed and quality filtered using Trimmomatic 0.38 (89) and assembled with

metaSPAdes v3.13.0 using default parameters (90). Sequencing and assembly data for

wastewater viromes is summarized in Table S5. Read mapping to phage genomes was

performed with bwa (91).

Human Microbiome Project Healthy Human Subjects Study samples were

downloaded with portal_client (Table S2). Read mapping was performed with bwa (91).

Data availability. Trimmed RNAseq reads from this study are deposited in the

NCBI SRA under PRJNA622597; sample accession numbers are SAMN14522273 for

the wild-type lysogen (WT), SAMN14522274 for the cured lysogen (ΔBV01), and

SAMN14522275 for the cured lysogen *tspO* deletion (ΔBV01Δ*tspO*). Trimmed

wastewater virome reads are deposited in the NCBI SRA under PRJNA622299. Ten new

assembled *B. vulgatus* genomes are deposited in NCBI GenBank under PRJNA622758.

ACKNOWLEDGMENTS

We thank Nadja Shoemaker and Abigail Salyers for access to an impressive collection of *Bacteroides* isolates; Ken Ringwald for critical review of the manuscript; Jim Imlay for insightful discussion on metabolism and stress; Alvaro Hernandez and Chris Wright for DNA and RNA sequencing; Bruce Rabe for aid in wastewater collection; Jonathan Mitchell for maintenance and animal care at the UCR vivarium.

This research was supported by initial complement funding to PHD from UIUC and UCR and DC was supported by the Department of Microbiology from UIUC. LL was supported by the National Science Foundation Graduate Research Fellowship. Gnotobiotic mouse work and AH were supported by National Institute of General Medical Sciences grant R35GM124724. RJW is supported by the Allen Foundation with an Allen Distinguished Investigator Award.

REFERENCES

1. Norman JM, Handley SA, Baldridge MT, Droit L, Liu CY, Keller BC, Kambal A, Monaco CL, Zhao G, Fleshner P, Stappenbeck TS, McGovern DPB, Keshavarzian A, Mutlu EA, Sauk J, Gevers D, Xavier RJ, Wang D, Parkes M, Virgin HW. 2015. Disease-Specific Alterations in the Enteric Virome in Inflammatory Bowel Disease. Cell 160:447–460.

2. Reyes A, Blanton LV, Cao S, Zhao G, Manary M, Trehan I, Smith MI, Wang D, Virgin HW, Rohwer F, Gordon JI. 2015. Gut DNA viromes of Malawian twins discordant for severe acute malnutrition. Proc Nat Acad Sci USA 112:11941–11946.

3.   Monaco CL, Gootenberg DB, Zhao G, Handley SA, Ghebremichael MS, Lim ES,

Lankowski A, Baldridge MT, Wilen CB, Flagg M, Norman JM, Keller BC, Luévano

JM, Wang D, Boum Y, Martin JN, Hunt PW, Bangsberg DR, Siedner MJ, Kwon DS,

Virgin HW. 2016. Altered Virome and Bacterial Microbiome in Human

Immunodeficiency Virus-Associated Acquired Immunodeficiency Syndrome. Cell

Host & Microbe 19:311–322.

4.   Hannigan GD, Duhaime MB, Ruffin MT, Koumpouras CC, Schloss PD. 2018.

Diagnostic Potential and Interactive Dynamics of the Colorectal Cancer Virome.

mBio 9:e02248-18.

5.   Han M, Yang P, Zhong C, Ning K. 2018. The Human Gut Virome in Hypertension.

Front Microbiol 9.

6.   Moreno-Gallego JL, Chou S-P, Di Rienzi SC, Goodrich JK, Spector TD, Bell JT,

Youngblut ND, Hewson I, Reyes A, Ley RE. 2019. Virome Diversity Correlates with

Intestinal Microbiome Diversity in Adult Monozygotic Twins. Cell Host & Microbe

25:261-272.e5.

7.   Khan Mirzaei M, Khan MdAA, Ghosh P, Taranu ZE, Taguer M, Ru J, Chowdhury R,

Kabir MdM, Deng L, Mondal D, Maurice CF. 2020. Bacteriophages Isolated from

Stunted Children Can Regulate Gut Bacterial Communities in an Age-Specific

Manner. Cell Host Microbe 27:199-212.e5.

8.   Hsu BB, Gibson TE, Yeliseyev V, Liu Q, Lyon L, Bry L, Silver PA, Gerber GK.

2019. Dynamic Modulation of the Gut Microbiota and Metabolome by

Bacteriophages in a Mouse Model. Cell Host & Microbe 25:803-814.e5.

9.   Salyers AA, Vercellotti JR, West SE, Wilkins TD. 1977. Fermentation of mucin and

plant polysaccharides by strains of Bacteroides from the human colon. Appl Environ

Microbiol 33:319–322.

10. Kaoutari AE, Armougom F, Gordon JI, Raoult D, Henrissat B. 2013. The abundance

and variety of carbohydrate-active enzymes in the human gut microbiota. Nature Rev

Microbiol 11:497–504.

11. Hickey CA, Kuhn KA, Donermeyer DL, Porter NT, Jin C, Cameron EA, Jung H,

Kaiko GE, Wegorzewska M, Malvin NP, Glowacki RWP, Hansson GC, Allen PM,

Martens EC, Stappenbeck TS. 2015. Colitogenic Bacteroides thetaiotaomicron

Antigens Access Host Immune Cells in a Sulfatase-Dependent Manner via Outer

Membrane Vesicles. Cell Host & Microbe 17:672–680.

12. Shen Y, Torchia MLG, Lawson GW, Karp CL, Ashwell JD, Mazmanian SK. 2012.

Outer Membrane Vesicles of a Human Commensal Mediate Immune Regulation and

Disease Protection. Cell Host & Microbe 12:509–520.

13. Tuncil YE, Xiao Y, Porter NT, Reuhs BL, Martens EC, Hamaker BR. 2017.

Reciprocal Prioritization to Dietary Glycans by Gut Bacteria in a Competitive

Environment Promotes Stable Coexistence. mBio 8:e01068-17.

14. Ridlon JM, Harris SC, Bhowmik S, Kang D-J, Hylemon PB. 2016. Consequences of

bile salt biotransformations by intestinal bacteria. Gut Microbes 7:22–39.

15. Cullen TW, Schofield WB, Barry NA, Putnam EE, Rundell EA, Trent MS, Degnan

PH, Booth CJ, Yu H, Goodman AL. 2015. Antimicrobial peptide resistance mediates

resilience of prominent gut commensals during inflammation. Science 347:170–175.

16. Planer JD, Peng Y, Kau AL, Blanton LV, Ndao IM, Tarr PI, Warner BB, Gordon JI.

2016. Development of the gut microbiota and mucosal IgA responses in twins and

gnotobiotic mice. Nature 534:263–266.

17. Porter NT, Hryckowian AJ, Merrill BD, Gardner JO, Singh S, Sonnenburg JL,

Martens EC. 2019. Multiple phase-variable mechanisms, including capsular

polysaccharides, modify bacteriophage susceptibility in Bacteroides

thetaiotaomicron. bioRxiv 521070.

18. Coyne MJ, Zitomersky NL, McGuire AM, Earl AM, Comstock LE. 2014. Evidence

of Extensive DNA Transfer between Bacteroidales Species within the Human Gut.

mBio 5:e01305-14.

19. Lange A, Beier S, Steimle A, Autenrieth IB, Huson DH, Frick J-S. 2016. Extensive

Mobilome-Driven Genome Diversification in Mouse Gut-Associated Bacteroides

vulgatus mpk. Genome Biol Evol 8:1197–1207.

20. Yutin N, Makarova KS, Gussow AB, Krupovic M, Segall A, Edwards RA, Koonin

EV. 2018. Discovery of an expansive bacteriophage family that includes the most

abundant viruses from the human gut. Nat Microbiol 3:38–46.

21. Dutilh BE, Cassman N, McNair K, Sanchez SE, Silva GGZ, Boling L, Barr JJ, Speth

DR, Seguritan V, Aziz RK, Felts B, Dinsdale EA, Mokili JL, Edwards RA. 2014. A

highly abundant bacteriophage discovered in the unknown sequences of human faecal

metagenomes. Nat Commun 5:1–11.

22. Shkoporov AN, Khokhlova EV, Fitzgerald CB, Stockdale SR, Draper LA, Ross RP,

Hill C. 2018. ΦCrAss001 represents the most abundant bacteriophage family in the

human gut and infects Bacteroides intestinalis. Nat Commun 9:1–8.

23. Krinos CM, Coyne MJ, Weinacht KG, Tzianabos AO, Kasper DL, Comstock LE.

2001. Extensive surface diversity of a commensal microorganism by multiple DNA

inversions. Nature 414:555–558.

24. Patrick S, Blakely GW, Houston S, Moore J, Abratt VR, Bertalan M, Cerdeño-

Tárraga AM, Quail MA, Corton N, Corton C, Bignell A, Barron A, Clark L, Bentley

SD, Parkhill J. 2010. Twenty-eight divergent polysaccharide loci specifying within-

and amongst-strain capsule diversity in three strains of Bacteroides fragilis.

Microbiology, 156:3255–3269.

25. Tartera C, Jofre J. 1987. Bacteriophages active against Bacteroides fragilis in

sewage-polluted waters. Appl Environ Microbiol 53:1632–1637.

26. Jofre J, Blanch AR, Lucena F, Muniesa M. 2014. Bacteriophages infecting

Bacteroides as a marker for microbial source tracking. Water Research 55:1–11.

27. Ogilvie LA, Bowler LD, Caplin J, Dedi C, Diston D, Cheek E, Taylor H, Ebdon JE,

Jones BV. 2013. Genome signature-based dissection of human gut metagenomes to

extract subliminal viral sequences. Nat Commun 4:1–16.

28. Minot S, Sinha R, Chen J, Li H, Keilbaugh SA, Wu GD, Lewis JD, Bushman FD.

2011. The human gut virome: Inter-individual variation and dynamic response to

diet. Genome Res 21:1616–1625.

29. Abeles SR, Ly M, Santiago-Rodriguez TM, Pride DT. 2015. Effects of Long Term

Antibiotic Therapy on Human Oral and Fecal Viromes. PLoS One 10:e0134941.

30. Sullivan MB, Coleman ML, Weigele P, Rohwer F, Chisholm SW. 2005. Three

Prochlorococcus Cyanophage Genomes: Signature Features and Ecological

Interpretations. PLoS Biology 3:e144.

31. Anantharaman K, Duhaime MB, Breier JA, Wendt KA, Toner BM, Dick GJ. 2014.

Sulfur Oxidation Genes in Diverse Deep-Sea Viruses. Science 344:757–760.

32. Berger P, Kouzel IU, Berger M, Haarmann N, Dobrindt U, Koudelka GB, Mellmann

A. 2019. Carriage of Shiga toxin phage profoundly affects Escherichia coli gene

expression and carbon source utilization. BMC Genomics 20:504.

33. Hernandez-Doria JD, Sperandio V. 2018. Bacteriophage Transcription Factor Cro

Regulates Virulence Gene Expression in Enterohemorrhagic Escherichia coli. Cell

Host & Microbe 23:607-617.e6.

34. Rabinovich L, Sigal N, Borovok I, Nir-Paz R, Herskovits AA. 2012. Prophage

Excision Activates Listeria Competence Genes that Promote Phagosomal Escape and

Virulence. Cell 150:792–802.

35. Carey JN, Mettert EL, Fishman-Engel DR, Roggiani M, Kiley PJ, Goulian M. 2019.

Phage integration alters the respiratory strategy of its host. eLife 8:e49081.

36. Chen Y-Y, Wang J-T, Lin T-L, Gong Y-N, Li T-H, Huang Y-Y, Hsieh Y-C. 2019.

Prophage excision in streptococcus pneumoniaeserotype 19A ST320 promote

colonization: Insight into its evolution from the ancestral clone Taiwan 19F-14

(ST236). Frontiers in Microbiology 10:205.

37. Nakayama K, Kanaya S, Ohnishi M, Terawaki Y, Hayashi T. 1999. The complete

nucleotide sequence of φCTX, a cytotoxin-converting phage of Pseudomonas

aeruginosa: implications for phage evolution and horizontal gene transfer via

bacteriophages. Mol Microbiol 31:399–419.

38. Winter SE, Thiennimitr P, Winter MG, Butler BP, Huseby DL, Crawford RW,

Russell JM, Bevins CL, Adams LG, Tsolis RM, Roth JR, Bäumler AJ. 2010. Gut

inflammation provides a respiratory electron acceptor for Salmonella. Nature

467:426–429.

39. Bensing BA, Siboo IR, Sullam PM. 2001. Proteins PblA and PblB of Streptococcus

mitis, Which Promote Binding to Human Platelets, Are Encoded within a Lysogenic

Bacteriophage. Infection and Immunity 69:6186–6192.

40. Jermyn WS, Boyd EF. 2002. Characterization of a novel *Vibrio* pathogenicity island

(VPI-2) encoding neuraminidase (*nanH*) among toxigenic *Vibrio cholerae* isolates.

Microbiology, 148:3681–3693.

41. Sekulovic O, Fortier L-C. 2015. Global Transcriptional Response of *Clostridium*

difficile Carrying the φCD38-2 Prophage. Appl Environ Microbiol 81:1364–1374.

42. Leplae R, Lima-Mendez G, Toussaint A. 2010. ACLAME: A CLAssification of

Mobile genetic Elements, update 2010. Nucleic Acids Res 38:D57–D61.

43. Lima-Mendez G, Van Helden J, Toussaint A, Leplae R. 2008. Prophinder: a

computational tool for prophage prediction in prokaryotic genomes. Bioinformatics

24:863–865.

44. Reyes A, Wu M, McNulty NP, Rohwer FL, Gordon JI. 2013. Gnotobiotic mouse

model of phage–bacterial host dynamics in the human gut. Proc Nat Acad Sci USA

110:20236–20241.

45. Kanehisa M, Goto S. 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes.

Nucleic Acids Res 28:27–30.

46. Yeliseev AA, Kaplan S. 1999. A Novel Mechanism for the Regulation of

Photosynthesis Gene Expression by the TspO Outer Membrane Protein of

Rhodobacter sphaeroides 2.4.1. J Biol Chem 274:21234–21243.

47. Davey ME, Bruijn FJ de. 2000. A Homologue of the Tryptophan-Rich Sensory

Protein TspO and FixL Regulate a Novel Nutrient Deprivation-Induced

Sinorhizobium meliloti Locus. Appl Environ Microbiol 66:5353–5359.

48. Balsemão-Pires E, Jaillais Y, Olson BJ, Andrade LR, Umen JG, Chory J, Sachetto-

Martins G. 2011. The *Arabidopsis* translocator protein (AtTSPO) is regulated at

multiple levels in response to salt stress and perturbations in tetrapyrrole metabolism.

BMC Plant Biol 11:108.

49. Yao L, Seaton SC, Ndousse-Fetter S, Adhikari AA, DiBenedetto N, Mina AI, Banks

AS, Bry L, Devlin AS. 2018. A selective gut bacterial bile salt hydrolase alters host

metabolism. eLife.

50. Meier-Kolthoff JP, Göker M. 2017. VICTOR: genome-based phylogeny and

classification of prokaryotic viruses. Bioinformatics 33:3396–3404.

51. Göker M, García-Blázquez G, Voglmayr H, Tellería MT, Martín MP. 2009.

Molecular Taxonomy of Phytopathogenic Fungi: A Case Study in Peronospora. PLoS

One 4: e6319.

52. Northfield TC, McColl I. 1973. Postprandial concentrations of free and conjugated

bile acids down the length of the normal human small intestine. Gut 14:513–518.

53. Hamilton JP, Xie G, Raufman J-P, Hogan S, Griffin TL, Packard CA, Chatfield DA,

Hagey LR, Steinbach JH, Hofmann AF. 2007. Human cecal bile acids: concentration

and spectrum. Am J Phys-Gastr L 293:G256–G263.

54. Citron DM, Baron EJ, Finegold SM, Goldstein EJ. 1990. Short prereduced

anaerobically sterilized (PRAS) biochemical scheme for identification of clinical

isolates of bile-resistant Bacteroides species. J Clin Microbiol 28:2220–2223.

55. Joyce SA, MacSharry J, Casey PG, Kinsella M, Murphy EF, Shanahan F, Hill C,

Gahan CGM. 2014. Regulation of host weight gain and lipid metabolism by bacterial

bile acid modification in the gut. Proc Nat Acad Sci USA 111:7421–7426.

56. Grau KR, Zhu S, Peterson ST, Helm EW, Philip D, Phillips M, Hernandez A, Turula

H, Frasse P, Graziano VR, Wilen CB, Wobus CE, Baldridge MT, Karst SM. 2020.

The intestinal regionalization of acute norovirus infection is regulated by the

microbiota via bile acid-mediated priming of type III interferon. Nature Microbiol

5:84–92.

57. Li F, Liu J, Zheng Y, Garavito RM, Ferguson-Miller S. 2015. Crystal structures of

translocator protein (TSPO) and mutant mimic of a human polymorphism. Science

347:555–558.

58. Guo Y, Kalathur RC, Liu Q, Kloss B, Bruni R, Ginter C, Kloppmann E, Rost B,

Hendrickson WA. 2015. Structure and activity of tryptophan-rich TSPO proteins.

Science 347:551–555.

59. Zeno S, Veenman L, Katz Y, Bode J, Gavish M, Zaaroor M. 2012. The 18 kDa

mitochondrial translocator protein (TSPO) prevents accumulation of protoporphyrin

IX. Involvement of reactive oxygen species (ROS). Curr Mol Med 12:494–501.

60. Li F, Liu J, Garavito RM, Ferguson-Miller S. 2015. Evolving understanding of

translocator protein 18kDa (TSPO). Pharmacol Res 99:404–409.

61. Rone MB, Midzak AS, Issop L, Rammouz G, Jagannathan S, Fan J, Ye X, Blonder J,

Veenstra T, Papadopoulos V. 2012. Identification of a Dynamic Mitochondrial

Protein Complex Driving Cholesterol Import, Trafficking, and Metabolism to Steroid

Hormones. Mol Endocrinol 26:1868–1882.

62. Devendran S, Shrestha R, Alves JMP, Wolf PG, Ly L, Hernandez AG, Méndez-

García C, Inboden A, Wiley J, Paul O, Allen A, Springer E, Wright CL, Fields CJ,

Daniel SL, Ridlon JM. 2019. Clostridium scindens ATCC 35704: Integration of

Nutritional Requirements, the Complete Genome Sequence, and Global

Transcriptional Responses to Bile Acids. Appl Environ Microbiol 85.

63. Roberts JW, Roberts CW, Craig NL. 1978. *Escherichia coli recA* gene product

inactivates phage lambda repressor. Proc Nat Acad Sci USA 75:4714–4718.

64. Moon K, Shoemaker NB, Gardner JF, Salyers AA. 2005. Regulation of Excision

Genes of the *Bacteroides* Conjugative Transposon CTnDOT. Journal of Bacteriology

187:5732–5741.

65. Puig M, Gironés R. 1999. Genomic structure of phage B40-8 of *Bacteroides fragilis*.

Microbiol, 145:1661–1670.

66. Salyers AA, Bonheyo G, Shoemaker NB. 2000. Starting a New Genetic System:

Lessons from *Bacteroides*. Methods 20:35–46.

67. Oh J-H, Lin XB, Zhang S, Tollenaar SL, Özçam M, Dunphy C, Walter J, Pijkeren J-P

van. 2019. Prophages in *Lactobacillus reuteri* Are Associated with Fitness Trade-

Offs but Can Increase Competitiveness in the Gut Ecosystem. Appl Environ

Microbiol 86.

68. Zhao S, Lieberman TD, Poyet M, Kauffman KM, Gibbons SM, Groussin M, Xavier

RJ, Alm EJ. 2019. Adaptive Evolution within Gut Microbiomes of Healthy People.

Cell Host Microbe 25:656-667.e8.

69. Holdeman LV, Cato EP, Moore WEC. 1977. Anaerobic Laboratory Manual, 4th

Edition. Anaerobe Laboratory, Virginia Polytechnic Institute and State University,

Blacksburg.

70. Goodman AL, McNulty NP, Zhao Y, Leip D, Mitra RD, Lozupone CA, Knight R,

Gordon JI. 2009. Identifying Genetic Determinants Needed to Establish a Human Gut

Symbiont in Its Habitat. Cell Host & Microbe 6:279–289.

71. Koropatkin NM, Martens EC, Gordon JI, Smith TJ. 2008. Starch Catabolism by a

Prominent Human Gut Symbiont Is Directed by the Recognition of Amylose Helices.

Structure 16:1105–1115.

72. Degnan PH, Taga ME, Goodman AL. 2014. Vitamin B12 as a Modulator of Gut

Microbial Ecology. Cell Metab 20:769–778.

73. Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved

microbes from next-generation sequencing data using breseq. Methods Mol Biol

1151:165–188.

74. Riley AB, Kim D, Hansen AK. 2017. Genome Sequence of "*Candidatus Carsonella*

ruddii" Strain BC, a Nutritional Endosymbiont of Bactericera cockerelli. Genome

Announc 5:e00236-17.

75. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010.

Prodigal: prokaryotic gene recognition and translation initiation site identification.

BMC Bioinformatics 11:119.

76. Lowe TM, Eddy SR. 1997. tRNAscan-SE: A Program for Improved Detection of

Transfer RNA Genes in Genomic Sequence. Nucleic Acids Res 25:955–964.

77. Lagesen K, Hallin P, Rødland EA, Stærfeldt H-H, Rognes T, Ussery DW. 2007.

RNAmmer: consistent and rapid annotation of ribosomal RNA genes. Nucleic Acids

Res 35:3100–3108.

78. Nawrocki EP, Eddy SR. 2013. Infernal 1.1: 100-fold faster RNA homology searches.

Bioinformatics 29:2933–2935.

79. Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiryutin B, Koonin EV, Krylov

DM, Mazumder R, Mekhedov SL, Nikolskaya AN, Rao BS, Smirnov S, Sverdlov

AV, Vasudevan S, Wolf YI, Yin JJ, Natale DA. 2003. The COG database: an updated

version includes eukaryotes. BMC Bioinformatics 4:41.

80. Bateman A, Coin L, Durbin R, Finn RD, Hollich V, Griffiths-Jones S, Khanna A,

Marshall M, Moxon S, Sonnhammer ELL, Studholme DJ, Yeats C, Eddy SR. 2004.

The Pfam protein families database. Nucleic Acids Res 32:D138–D141.

81. Haft DH, Selengut JD, White O. 2003. The TIGRFAMs database of protein families.

Nucleic Acids Res 31:371–373.

82. Yu C-S, Cheng C-W, Su W-C, Chang K-C, Huang S-W, Hwang J-K, Lu C-H. 2014.

CELLO2GO: A Web Server for Protein subCELlular LOcalization Prediction with

Functional Gene Ontology Annotation. PLoS ONE 9:e99368.

83. Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. 2012. Artemis: an

integrated platform for visualization and analysis of high-throughput sequence-based

experimental data. Bioinformatics 28:464–469.

84. Mount DW. 2007. Using the Basic Local Alignment Search Tool (BLAST). Cold

Spring Harb Protoc 2007:pdb.top17.

85. Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. 2009. GenBank.

Nucleic Acids Res 37:D26–D31.

86. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 web

portal for protein modeling, prediction and analysis. Nat Protoc 10:845–858.

87. Seguritan V, Jr NA, Arnoult M, Raymond A, Lorimer D, Jr ABB, Salamon P, Segall

AM. 2012. Artificial Neural Networks Trained to Detect Viral and Phage Structural

Proteins. PLoS Comput Biol 8:e1002657.

88. McClure R, Balasubramanian D, Sun Y, Bobrovskyy M, Sumby P, Genco CA,

Vanderpool CK, Tjaden B. 2013. Computational analysis of bacterial RNA-Seq data.

Nucleic Acids Res 41:e140–e140.

89. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30:2114–2120.

90. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new

versatile metagenomic assembler. Genome Res 27:824–834.

91. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler

transform. Bioinformatics 25:1754–1760.

92. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and

high throughput. Nucleic Acids Res 32:1792–1797.

93. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 – Approximately Maximum-

Likelihood Trees for Large Alignments. PLOS ONE 5:e9490.

94. Noguchi H, Taniguchi T, Itoh T. 2008. MetaGeneAnnotator: Detecting Species-

Specific Patterns of Ribosomal Binding Site for Precise Gene Prediction in

Anonymous Prokaryotic and Phage Genomes. DNA Res 15:387–396.

95. Roux S, Enault F, Hurwitz BL, Sullivan MB. 2015. VirSorter: mining viral signal

from microbial genomic data. PeerJ 3:e985.

96. Simon R, Priefer U, Pühler A. 1983. A Broad Host Range Mobilization System for In

Vivo Genetic Engineering: Transposon Mutagenesis in Gram Negative Bacteria. Nat

Biotechnol 1:784–791.

97. Eggerth AH, Gagnon BH. 1933. The *Bacteroides* of Human Feces. Journal of

Bacteriology 25:389–413.

98. Johnson JL. 1978. Taxonomy of the *Bacteroides*. International Journal of Systematic

and Evolutionary Microbiology, 28:245–256.

99. Whittle G, Hund BD, Shoemaker NB, Salyers AA. 2001. Characterization of the 13-

Kilobase *ermF* Region of the *Bacteroides* Conjugative Transposon CTnDOT. Appl

Environ Microbiol 67:3488–3495.

100. Shoemaker NB, Vlamakis H, Hayes K, Salyers AA. 2001. Evidence for Extensive

Resistance Gene Transfer amongBacteroides spp. and among Bacteroides and Other

Genera in the Human Colon. Appl Environ Microbiol 67:561–568.

101. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh

PJ, Fierer N, Knight R. 2011. Global patterns of 16S rRNA diversity at a depth of

millions of sequences per sample. PNAS 108:4516–4522.

FIGURE LEGENDS

Figure 1. DNA sequencing reads mapped to the BV01 prophage region. Reads from

shotgun sequencing of *B. vulgatus* genomic DNA (WT) or isolated phage DNA (Free

BV01) were normalized to the total number of reads after trimming, and represented as a

coverage curve. A cured lysogen (ΔBV01) and integrase deletion mutant (Δint) of *B.*

vulgatus were confirmed by shotgun sequencing of genomic DNA. The discrete coverage

peak at position ~3,588,000 nt from ΔBV01 is attributed to a homologous sequence

elsewhere on the *B. vulgatus* chromosome. Putative functions of BV01 genes are

indicated by the colors in the legend.

Figure 2. Prophage BV01 is an intact prophage. (A) Excision and circularization

activities of the BV01 integrase are confirmed by PCR. The presence of phage DNA was

detected by amplification of a phage marker gene (BVU_RS14350). Amplification across

the phage attachment site (*attP*) indicates circularization of the BV01 genome; *attP*

amplicons from the integrase complement strain (Δ*int* + pNBU2-*int*) are ~1.2 Kb shorter

than wild-type amplicons due to deletion of the integrase gene. Supernatant fractions

were treated with DNase, eliminating all contaminating host genomic DNA, as

demonstrated by the amplification of a host marker gene (16S rRNA). Note that despite

apparent size shift of BVU_RS14350 amplicons from the pellets and supernatants,

Sanger sequencing validated that the products are in fact identical. PCR amplicons were

visualized by agarose gel electrophoresis alongside NEB 1 Kb DNA ladder

(BVU_RS14350, *attP*) or GeneRuler Express DNA ladder (16S rRNA); ladder band sizes

shown in Kb. (B) BV01 can transduce uninfected hosts in a gnotobiotic mouse. Germ-

free mice (*n*=7) were gavaged with an equal mixture of a BV01-*tetQ* lysogen and an

erythromycin-tagged cured lysogen (Day 0). Recipient, donor, and transductant cells

were identified by plating on Brain Heart Infusion (BHI) media with antibiotic selection:

erythromycin (Erm) or tetracycline (Tet).

Figure 3. Differential regulation of the host transcriptome in response to BV01

lysogeny. (A) Count of differentially regulated transcripts as compared to the wild-type

B. vulgatus lysogen (fold change $\geq$ 2, *q*-value $\leq$ 0.01). (B) Chromosomal localization of

the differentially expressed genes. Each dot represents a differentially expressed

transcript on a $\log_2$ scale; genes below the 2-fold change cutoff (yellow) and within the

BV01 prophage (blue) not shown. Positive fold change values correspond to increased

transcription in the second background listed. (C) General functional assignment of genes

differentially expressed between wild-type and cured lysogen strains was accomplished

using the Clusters of Orthologous Groups (COGs). Transcripts which are not

differentially expressed in other strain comparisons are shown in black. *tspO*-dependent

transcripts are marked on right (•). Positive fold change values correspond to increased

transcription in the second background listed. Letters correspond to COG categories: cell

wall/membrane/envelope biogenesis (M), post-translation modification, protein turnover,

and chaperones (O), signal transduction mechanisms (T), transcription (K), energy

production and conversion (C), amino acid transport and metabolism (E), carbohydrate

transport and metabolism (G), coenzyme transport and metabolism (H), lipid transport

and metabolism (I), inorganic ion transport and metabolism (P), secondary metabolite

biosynthesis, transport, and catabolism (Q), general function prediction only (R), function

unknown (S).

Figure 4. BV01 integration disrupts transcription of *tspO*. (A) Transcriptional activity

of the *tspO* gene region as it exists in the cured lysogen background. RNAseq reads from

wild-type (WT) and cured lysogen (ΔBV01) *B. vulgatus* were mapped to the region, and

coverage was normalized to the total number of reads mapping to the genome. The

average normalized read coverage for each genome is displayed as the y-axis maximum

(grey line). Maximum read coverage for the region is displayed as the grey dashed line.

(B) Amino acid alignment of *B. vulgatus* TspO with known TspO sequences was

generated with MUSCLE (92). Identical and similar residues are colored blue and gray,

respectively. Shown are TspO protein sequences from *B. vulgatus* (WP_005843416.1),

Bacillus cereus (GCF80909.1), *Rhodobacter sphaeroides* (AAF24291.1), *Sinorhizobium*

meliloti (AAF01195.1), human (NP_001243460.1), and rat (NP_036647.1). Secondary

structures, cholesterol recognition/interaction amino acid consensus (CRAC) sequence,

and critical residues (•) from *R. sphaeroides* TspO crystal structure are shown (57). (C)

The search for TspO homologs in the family *Bacteroidales* was accomplished with a

BLAST-based approach, using the *Bacillus cereus* copy of TspO (GCF80909.1) as a

query against a database of 154 gut-associated *Bacteroidales* genomes, 122 of which are

from the genus *Bacteroides*. Genome counts are indicated within categories. (D) Gene

tree estimated from TspO sequences across the *Bacteroidales*. All *B. vulgatus* and *B.*

dorei genomes included in the search encode *tspO*. Clade for *B. vulgatus* TspO sequences

is displayed as a polytomy; all *B. vulgatus* TspO sequences are at least 98% identical to

each other. Numbers above branches represent bootstrap values; only bootstraps over 50

shown. The gene tree was estimated using FastTree (93).

Figure 5. BV01 alters host interactions with bile acids in a *tspO*-dependent manner.

(A) Representative LC/MS traces showing *B. vulgatus* deconjugates taurocholic acid

(TCA) to cholic acid (CA) in the cured lysogen background (ΔBV01), but little or no

activity CA is detectable in the wild-type (WT) or cured lysogen *tspO* deletion

(ΔBV01Δ*tspO*) backgrounds. *B. vulgatus* cultures were incubated with 50 μM TCA for

16 hr prior to bile acid extraction. (B) Representative LC/MS traces showing *B. vulgatus*

deconjugates glycocholic acid (GCA) to CA in the ΔBV01 background, but not in the

WT or ΔBV01Δ*tspO* backgrounds. *B. vulgatus* cultures were incubated with 50 μM GCA

for 48 hr prior to bile acid extraction. Nordeoxycholic acid (norDCA) was added to a

final concentration of 15 μM as an internal standard after incubation. Peaks labeled for

their metabolites based on m/z; TCA = 514.29, GCA = 464.30, CA = 407.28, norDCA =

377.27.

Figure 6. *Salyersviridae* occur throughout the *Bacteroides* genus. (A) *Bacteroides*

phylogeny and occurrence of *Salyersviridae att* site. All duplications of the *att* site are

associated with a putative integrated prophage. Host phylogeny estimated by maximum

likelihood from concatenated alignment of 13 core genes. (B) Consensus *att* site for

Salyersvirinae. The *attP* is duplicated upon integration of a *Salyersvirinae* prophage,

resulting in direct repeats. Image made with the WebLogo online tool. (C) Phylogenomic

Genome-BLAST Distance Phylogeny implemented with the VCTOR online tool (50)

using amino acid data from all phage ORFs. For consistency, all phage genomes were

annotated with MetaGeneAnnotator (94) implemented via VirSorter (95). Support values

above branches are GBDP pseudo-bootstrap values from 100 replications. Family (F),

subfamily (Subf), genus (G), and species (Sp) assigned by OPTSIL clustering (51) (Table

1). Each leaf of the tree represents a unique phage species, except where indicated by

colored boxes. Active prophages confirmed by sequencing and/or PCR indicated with

"•"; prophages confirmed to have been inactivated by genome rearrangement indicated

with "x"; prophages which were tested for activity with inconclusive results indicated

with "*" (Fig. S4).

TABLES

 Table 1. *Bacteroides* phage taxonomy as determined by whole genome clustering.

Phage	Host	Family	Subfamily	Genus	Species
BV01	*B. vulgatus* ATCC 8482	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BV01*
BV02	*B. vulgatus* VPI-4245	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BV01*
BV03	*B. vulgatus* VPI-2365	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BV03*
BV04	*B. vulgatus* VPI-6186	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BV04*
BV05	*B. vulgatus* VPI-5710	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BV05*
BV06	*B. vulgatus* DH4096S	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BV06*
BF01	*B. finegoldii* CL09T03C10	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BF01*
BT01	*Bacteroides* sp. 1_1_6	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BX01*
BO01	*B. ovatus* ATCC 8483	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BO01*
BO02	*B. ovatus* CL02T12C04	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BO02*
BO03	*Bacteroides*	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BO03*

	sp. 2_2_4				
BO04	*Bacteroides* sp. D2	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BO04*
BX01	*B. xylanisolvens* XB1A	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BX01*
BX02	*Bacteroides* sp. D1	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BX02*
BX03	*Bacteroides* sp. 2_1_22	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BX02*
BX04	*B. ovatus* SD CC 2a	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BX02*
BX05	*B. xylanisolvens* SD CC 1b	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BX02*
BX06	*Bacteroides* sp. D22	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BX06*
BX07	*B. ovatus* SD CMC 3f	*Salyersviridae*	*Salyersvirinae*	*Salyersvirus*	*BX07*
BC01	*B. clarus* YIT 12056	*Salyersviridae*	?	?	*BC01*
B124-14	*B. fragilis*	*Salyersviridae*	?	?	*B124-14*

B40-8	*B. fragilis*	*Salyersviridae*	?	?	*B40-8*
crAssphage	*B. intestinalis*	?	?	?	?

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. Transcriptional activity of the BV01 prophage and surrounding

chromosome. RNAseq reads from wild-type lysogen (WT) and cured lysogen (ΔBV01)

strains were mapped to the region, and coverage was normalized to the total number of

reads mapping to the genome. One representative of three replicates shown for each. The

average normalized read coverage for each genome is displayed as the y-axis maximum

(grey line). Maximum read coverage for the region is indicated on the y-axis. Forward

reads (red) and reverse reads (blue) were plotted separately. Locations of two putative

BV01-encoded   transcriptional   regulators   are   indicated   (BVU_RS14235,

BVU_RS14475).

Figure S2. Prevalence of prophage insertion adjacent to *tspO* in human gut samples.

Reads from 256 healthy human gut metagenomes were obtained from the Human

Microbiome Project Healthy Human Subjects Study. (A) Reads were first mapped to

representative sequences of *tspO* from *B. vulgatus* and *B. dorei*. Percent abundance *tspO*

reads was calculated on a per sample basis as the number of reads mapping to *tspO*

divided by the total number of reads. Histogram shows counts of samples. (B) Reads

mapping to *tspO* were filtered to only include reads antisense to *tspO*, predicted to point

toward the *attB* based on the known genomic architecture. Mates to those reads were

subsequently mapped to BV01 and its *Salyersvirus* relatives (Fig. 6). Only samples with

read pairs bridging *tspO* and a phage sequence are shown (*n*=34). Percent *tspO* reads

mated with *Salyersvirus* reads was calculated as the read pairs bridging *tspO* and a phage

sequence divided by the total number of reads mapping antisense to *tspO*. Histogram

shows counts of samples.

1011

Figure S3. Paired whole genome tree and nucleotide alignment of *Salyersviridae*

phages. Phylogenomic Genome-BLAST Distance Phylogeny implemented with the

VCTOR online tool (50) using amino acid data from all phage ORFs. For consistency, all

phage genomes were annotated with MetaGeneAnnotator (94) implemented via VirSorter

(95). Support values above branches are GBDP pseudo-bootstrap values from 100

replications. Genome alignment of all phages made with MAUVE. One locally collinear

block (LCB) connects phages B124-14 and B40-8 to the *Salyersviridae* at the nucleotide

level (green). Other LCB connecting lines removed for clarity.

1020

Figure S4. Confirmation of activity of three additional Salyersviruses. DNase-treated

culture supernatants for predicted Salyersviruses BV02 (A), BV04 (B), and BV05 (C)

were sequenced. Assembly resulted in contigs corresponding to the free form of phages

BV02 and BV05; BV04 did not yield any contigs corresponding to the putative prophage

region, suggesting it is inactivated. Assembled free phage contigs were aligned to their

integrated prophage region with Mauve (A-C). Sequence reads were mapped back to

their free and integrated forms and represented as coverage curves (A-C). Vertical orange

lines indicate the location of *att* sequences; vertical dashed blue lines indicate the location

of contig breaks. (D) PCR amplification with phage-specific primers tests for phage

presence in pellet and supernatant fractions for 7 predicted Salyersviruses. Supernatant

fractions were treated with DNase, eliminating all contaminating host genomic DNA, as

demonstrated by the amplification of a host marker gene (16S rRNA). BV04 is not

detectable in supernatant, supporting the conclusion that it is an inactivated prophage.

PCR amplicons were visualized by agarose gel electrophoresis alongside GeneRuler

Express DNA ladder (16S rRNA); ladder band sizes shown in Kb.

1036

Figure S5. *Salyersviridae* sequence is detectable in human-associated samples. (A)

Wastewater viromes were collected, and processed in three ways prior to sequencing (see

Methods). Resulting reads were trimmed, pooled, and mapped to all *Salyersviridae*

genomes and crAssphage. Only *Salyersvirinae* genomes shown in alignment to better

demonstrate conservation, constructed with Mauve. Read coverages normalized to total

number of reads in the metavirome. The maximum normalized read coverage for BC01 is

4.1, B40-8 and B124-14 is 1.13, and crAssphage is 7.33. (B) Reads from 256 healthy

human gut metagenomes were obtained from the Human Microbiome Project Healthy

Human Subjects Study. Reads were mapped to all 20 temperate *Salyersviridae* phages

and crAssphage. Percent reads mapping was calculated on a per sample basis as the

number of reads mapping to any virus divided by the total number of reads. Histogram

shows counts of samples.

1049

1050   **SUPPLEMENTARY TABLE TITLES**

1051   **Table S1.** Rockhopper results for RNAseq from B. vulgatus WT (602), ΔBV01 (853) and

1052   ΔBV01ΔtspO (1662).

1053

1054   **Table S2.** Detection of *Salyersviridae* in Human Microbiome Project Healthy Human

1055   Subjects Study samples.

1056

1057   **Table S3.** Bacterial strains, plasmids, and primers used in this study.

1058

1059   **Table S4.** Breseq read mapping results of *B. vulgatus* WT and mutant strains used in this

1060   study.

1061

1062   **Table S5.** DNA sequencing data generated for B. vulgatus isolates, cell-free

1063   Salyersviridae phages, and wastewater viromes used in this study.

Figure 1. DNA sequencing reads mapped to the BV01 prophage region. Reads from shotgun sequencing of *B. vulgatus* genomic DNA (WT) or isolated phage DNA (Free BV01) were normalized to the total number of reads after trimming, and represented as a coverage curve. A cured lysogen (ΔBV01) and integrase deletion mutant (Δ*int*) of *B. vulgatus* were confirmed by shotgun sequencing of genomic DNA. The discrete coverage peak at position ~3,588,000 nt from ΔBV01 is attributed to a homologous sequence elsewhere on the *B. vulgatus* chromosome. Putative functions of BV01 genes are indicated by the colors in the legend.

Figure 2. Prophage BV01 is an intact prophage. (A) Excision and circularization activities of the BV01 integrase are confirmed by PCR. The presence of phage DNA was detected by amplification of a phage marker gene (BVU_RS14350). Amplification across the phage attachment site (*attP*) indicates circularization of the BV01 genome; *attP* amplicons from the integrase complement strain (Δ*int* + pNBU2-*int*) are ~1.2 Kb shorter than wild-type amplicons due to deletion of the integrase gene. Supernatant fractions were treated with DNase, eliminating all contaminating host genomic DNA, as demonstrated by the amplification of a host marker gene (16S rRNA). Note that despite apparent size shift of BVU_RS14350 amplicons from the pellets and supernatants, Sanger sequencing validated that the products are in fact identical. PCR amplicons were visualized by agarose gel electrophoresis alongside NEB 1 Kb DNA ladder (BVU_RS14350, *attP*) or GeneRuler Express DNA ladder (16S rRNA); ladder band sizes shown in Kb. (B) BV01 can transduce uninfected hosts in a gnotobiotic mouse. Germ-free mice (*n*=7) were gavaged with an equal mixture of a BV01-*tetQ* lysogen and an erythromycin-tagged cured

lysogen (Day 0). Recipient, donor, and transductant cells were identified by plating on Brain Heart

Infusion (BHI) media with antibiotic selection: erythromycin (Erm) or tetracycline (Tet).

Figure 3. Differential regulation of the host transcriptome in response to BV01 lysogeny. (A) Count of differentially regulated transcripts as compared to the wild-type *B. vulgatus* lysogen (fold change ≥ 2, *q*-value ≤ 0.01). (B) Chromosomal localization of the differentially expressed genes. Each dot represents a differentially expressed transcript on a log₂ scale; genes below the 2-fold change cutoff (yellow) and within the BV01 prophage (blue) not shown. Positive fold change values correspond to increased transcription in the second background listed. (C) General functional assignment of genes differentially expressed between wild-type and cured lysogen strains was accomplished using the Clusters of Orthologous Groups (COGs). Transcripts which are not differentially expressed in other strain comparisons are shown in black. *tspO*-dependent

transcripts are marked on right (•). Positive fold change values correspond to increased transcription in the second background listed. Letters correspond to COG categories: cell wall/membrane/envelope biogenesis (M), post-translation modification, protein turnover, and chaperones (O), signal transduction mechanisms (T), transcription (K), energy production and conversion (C), amino acid transport and metabolism (E), carbohydrate transport and metabolism (G), coenzyme transport and metabolism (H), lipid transport and metabolism (I), inorganic ion transport and metabolism (P), secondary metabolite biosynthesis, transport, and catabolism (Q), general function prediction only (R), function unknown (S).

Figure 4. BV01 integration disrupts transcription of *tspO*. (A) Transcriptional activity of the *tspO* gene region as it exists in the cured lysogen background. RNAseq reads from wild-type (WT) and cured lysogen (ΔBV01) *B. vulgatus* were mapped to the region, and coverage was normalized to the total number of reads mapping to the genome. The average normalized read coverage for each genome is displayed as the y-axis maximum (grey line). Maximum read coverage for the region is displayed as the grey dashed line. (B) Amino acid alignment of *B. vulgatus* TspO with known TspO sequences was generated with MUSCLE (49). Identical and similar residues are colored blue and gray, respectively. Shown are TspO protein sequences from *B. vulgatus* (WP_005843416.1), *Bacillus cereus* (GCF80909.1), *Rhodobacter sphaeroides* (AAF24291.1), *Sinorhizobium meliloti* (AAF01195.1), human (NP_001243460.1), and rat (NP_036647.1). Secondary structures, cholesterol recognition/interaction amino acid consensus (CRAC) sequence, and critical residues (•) from *R. sphaeroides* TspO crystal structure are shown (50). (C) The search

for TspO homologs in the family *Bacteroidales* was accomplished with a BLAST-based approach, using the *Bacillus cereus* copy of TspO (GCF80909.1) as a query against a database of 154 gut-associated *Bacteroidales* genomes, 122 of which are from the genus *Bacteroides*. Genome counts are indicated within categories. (D) Gene tree estimated from TspO sequences across the *Bacteroidales*. All *B. vulgatus* and *B. dorei* genomes included in the search encode *tspO*. Clade for *B. vulgatus* TspO sequences is displayed as a polytomy; all *B. vulgatus* TspO sequences are at least 98% identical to each other. Numbers above branches represent bootstrap values; only bootstraps over 50 shown. The gene tree was estimated using FastTree (51).

Figure 5. BV01 alters host interactions with bile acids in a *tspO*-dependent manner. (A) Representative LC/MS traces showing *B. vulgatus* deconjugates taurocholic acid (TCA) to cholic acid (CA) in the cured lysogen background (ΔBV01), but little or no activity CA is detectable in the wild-type (WT) or cured lysogen *tspO* deletion (ΔBV01Δ*tspO*) backgrounds. *B. vulgatus* cultures were incubated with 50 µM TCA for 16 hr prior to bile acid extraction. (B) Representative LC/MS traces showing *B. vulgatus* deconjugates glycocholic acid (GCA) to CA in the ΔBV01 background, but not in the WT or ΔBV01Δ*tspO* backgrounds. *B. vulgatus* cultures were incubated with 50 µM GCA for 48 hr prior to bile acid extraction. Nordeoxycholic acid (norDCA) was added to a final concentration of 15 µM as an internal standard after incubation. Peaks labeled for their metabolites based on m/z; TCA = 514.29, GCA = 464.30, CA = 407.28, norDCA = 377.27.

Figure 6. *Salyersviridae* occur throughout the *Bacteroides* genus. (A) *Bacteroides* phylogeny and occurrence of *Salyersviridae att* site. All duplications of the *att* site are associated with a putative integrated prophage. Host phylogeny estimated by maximum likelihood from concatenated alignment of 13 core genes. (B) Consensus *att* site for *Salyersvirinae*. The *attP* is duplicated upon integration of a *Salyersvirinae* prophage, resulting in direct repeats. Image made with the WebLogo online tool. (C) Phylogenomic Genome-BLAST Distance Phylogeny implemented with the VCTOR online tool (53) using amino acid data from all phage ORFs. For

consistency, all phage genomes were annotated with MetaGeneAnnotator (54) implemented via

VirSorter (55). Support values above branches are GBDP pseudo-bootstrap values from 100

replications. Family (F), subfamily (Subf), genus (G), and species (Sp) assigned by OPTSIL

clustering (56) (Table 1). Each leaf of the tree represents a unique phage species, except where

indicated by colored boxes. Active prophages confirmed by sequencing and/or PCR indicated with

"•"; prophages confirmed to have been inactivated by genome rearrangement indicated with "x";

prophages which were tested for activity with inconclusive results indicated with "*" (Fig. S4).