Structural and functional analysis of a potent sarbecovirus neutralizing antibody

Dora Pinto[1,7], Young-Jun Park[2,7], Martina Beltramello[1,7], Alexandra C. Walls[2,7], M.

Alejandra Tortorici[2,3], Siro Bianchi[1], Stefano Jaconi[1], Katja Culap[1], Fabrizia Zatta[1],

Anna De Marco[1], Alessia Peter[1], Barbara Guarino[1], Roberto Spreafico[4], Elisabetta

Cameroni[1], James Brett Case[6], Rita E. Chen[6], Colin Havenar-Daughton[4], Gyorgy

Snell[4], Amalio Telenti[4], Herbert W. Virgin[4], Antonio Lanzavecchia[1,5], Michael S.

Diamond[6], Katja Fink[1], David Veesler[2*] and Davide Corti[1*]

[1] Humabs Biomed SA, a subsidiary of Vir Biotechnology, 6500 Bellinzona, Switzerland.
[2] Department of Biochemistry, University of Washington, Seattle, Washington 98195, USA.
[3] Institut Pasteur & CNRS UMR 3569, Unité de Virologie Structurale, 75015, Paris, France.
[4] Vir Biotechnology, San Francisco, California 94158, USA
[5] Institute for Research in Biomedicine, Università della Svizzera Italiana, 6500 Bellinzona,
Switzerland.
[6] Departments of Medicine, Molecular Microbiology, Pathology & Immunology, Washington
University School of Medicine, St. Louis, MO 631110, USA
[7] These authors contributed equally

Correspondence: dveesler@uw.edu, dcorti@vir.bio

SARS-CoV-2 is a newly emerged coronavirus responsible for the current COVID-
19 pandemic that has resulted in more than one million infections and 73,000
deaths[1,2]**. Vaccine and therapeutic discovery efforts are paramount to curb the**
pandemic spread of this zoonotic virus. The SARS-CoV-2 spike (S) glycoprotein
promotes entry into host cells and is the main target of neutralizing antibodies.
Here we describe multiple monoclonal antibodies targeting SARS-CoV-2 S
identified from memory B cells of a SARS survivor infected in 2003. One
antibody, named S309, potently neutralizes SARS-CoV-2 and SARS-CoV
pseudoviruses as well as authentic SARS-CoV-2 by engaging the S receptor-
binding domain. Using cryo-electron microscopy and binding assays, we show
that S309 recognizes a glycan-containing epitope that is conserved within the
sarbecovirus subgenus, without competing with receptor attachment. Antibody
cocktails including S309 along with other antibodies identified here further
enhanced SARS-CoV-2 neutralization and may limit the emergence of
neutralization-escape mutants. These results pave the way for using S309 and
S309-containing antibody cocktails for prophylaxis in individuals at high risk of
exposure or as a post-exposure therapy to limit or treat severe disease.

Coronavirus entry into host cells is mediated by the transmembrane spike (S)

glycoprotein that forms homotrimers protruding from the viral surface[3]. The S

glycoprotein comprises two functional subunits: S_1 (divided into A, B, C and D domains)

that is responsible for binding to host cell receptors and S_2 that promotes fusion of the

viral and cellular membranes[4,5]. Both SARS-CoV-2 and SARS-CoV belong to the

sarbecovirus subgenus and their S glycoproteins share 80% amino acid sequence

identity[6]. SARS-CoV-2 S is closely related to the bat SARS-related CoV (SARSr-CoV)

RaTG13 with which it shares 97.2% amino acid sequence identity[1]. We and others

recently demonstrated that human angiotensin converting enzyme 2 (hACE2) is a

functional receptor for SARS-CoV-2, as is the case for SARS-CoV[1,6-8]. The S domain

B (S^B) is the receptor binding domain (RBD) and binds to hACE2 with high-affinity,

possibly contributing to the current rapid SARS-CoV-2 transmission in humans[6,9], as

previously proposed for SARS-CoV[10].

As the coronavirus S glycoprotein mediates entry into host cells, it is the main

target of neutralizing antibodies and the focus of therapeutic and vaccine design

efforts[3]. The S trimers are extensively decorated with N-linked glycans that are

important for protein folding[11] and modulate accessibility to host proteases and

neutralizing antibodies[12-15]. Cryo-electron microscopy (cryoEM) structures of SARS-

CoV-2 S in two distinct functional states[6,9] along with cryoEM and crystal structures of

SARS-CoV-2 S^B in complex with hACE2[16-18] revealed dynamic states of S^B domains,

providing a blueprint for the design of vaccines and inhibitors of viral entry.

Passive administration of monoclonal antibodies (mAbs) could have a major

impact on controlling the SARS-CoV-2 pandemic by providing immediate protection,

complementing the development of prophylactic vaccines. Accelerated development

of mAbs in a pandemic setting could be reduced to 5-6 months compared to the

traditional timeline of 10-12 months (Kelley B., Developing monoclonal antibodies at

pandemic speed, Nat Biotechnol, in press). The recent finding that ansuvimab

(mAb114) is a safe and effective treatment for symptomatic Ebola virus infection is a

striking example of the successful use of mAb therapy during an infectious disease

outbreak[19,20]. We previously isolated potently neutralizing human mAbs from memory

B cells of individuals infected with SARS-CoV[21] or MERS-CoV[22]. Passive transfer of

these mAbs protected animals challenged with various SARS-CoV isolates and SARS-

related CoV (SARSr-CoV)[21,23,24], as well as with MERS-CoV[22]. Structural

characterization of two of these mAbs in complex with SARS-CoV S and MERS-CoV

S provided molecular-level information on the mechanisms of viral neutralization[14]. In

particular, while both mAbs blocked S^B attachment to the host receptor, the SARS-

CoV-neutralizing S230 mAb acted by functionally mimicking receptor-attachment and

promoting S fusogenic conformational rearrangements[14]. Another mechanism of

SARS-CoV neutralization was recently described for mAb CR3022, which bound a

cryptic epitope only accessible when at least two out of the three S^B domains of a S

trimer were in the open conformation[25,26]. However, none of these mAbs neutralize

SARS-CoV-2.

Identification of a potent SARS-CoV-2 neutralizing mAb from a SARS survivor

We previously identified a set of human neutralizing mAbs from an individual

infected with SARS-CoV in 2003 that potently inhibited both human and zoonotic

SARS-CoV isolates[21,23,27]. To characterize the potential cross-reactivity of these

antibodies with SARS-CoV-2, we performed a new memory B cell screening using

peripheral blood mononuclear cells collected in 2013 from the same patient. We

describe here nineteen mAbs from the initial screen (2004 blood draw)[21,23] and six

mAbs from the new screen (2013 blood draw). The identified mAbs had a broad V

gene usage and were not clonally related (**Table 1).** Eight out of the twenty five mAbs

bound to SARS-CoV-2 S and SARS-CoV S transfected CHO cells with EC_{50} values

ranging between 1.4 and 6,100 ng/ml, and 0.8 and 254 ng/ml, respectively **(Fig. 1a-**

b). MAbs were further evaluated for binding to the SARS-CoV-2 and SARS-CoV S^B

domains as well as to the prefusion-stabilized OC43 S[28], MERS-CoV S[29,30], SARS-

CoV S[30] and SARS-CoV-2 S[6] ectodomain trimers. None of the mAbs studied bound to

prefusion OC43 S or MERS-CoV S ectodomain trimers, indicating a lack of cross-

reactivity outside the sarbecovirus subgenus **(Extended Data Fig.1)**. MAbs S303,

S304, S309 and S315 recognized the SARS-CoV-2 and SARS-CoV RBDs. In

particular, S309 bound with nanomolar affinity to both S^B domains, as determined by

biolayer interferometry **(Fig. 1c-d, Extended Data Fig. 2)**. Unexpectedly, S306 and

S310 stained cells expressing SARS-CoV-2 S at higher levels than those expressing

SARS-CoV S, yet it did not interact with SARS-CoV-2 or SARS-CoV S ectodomain

trimers and RBD constructs by ELISA. These results suggest that they may recognize

post-fusion SARS-CoV-2 S, which was recently proposed to be abundant on the

surface of authentic SARS-CoV-2 viruses[31] **(Fig. 1a-b and Extended Data Fig.3)**.

To evaluate the neutralization potency of the SARS-CoV-2 cross-reactive

mAbs, we carried out pseudovirus neutralization assays using a murine leukemia virus

(MLV) pseudotyping system[32]. S309 showed comparable neutralization potencies

against both SARS-CoV and SARS-CoV-2 pseudoviruses, whereas S303 neutralized

SARS-CoV-MLV but not SARS-CoV-2-MLV. S304 and S315 weakly neutralized

SARS-CoV-MLV and SARS-CoV-2-MLV (**Extended Data Fig.4**). In addition, S309

neutralized SARS-CoV-MLVs from isolates of the 3 phases of the 2002-2003 epidemic

with IC_{50} values comprised between 120 and 180 ng/ml and partially neutralized the

SARSr-CoV[33] WIV-1 (**Fig. 1e**). Finally, mAb S309 potently neutralized authentic

SARS-CoV-2 (2019n-CoV/USA_WA1/2020) with an IC_{50} of 69 ng/ml (**Fig. 1f**).

Structural basis of S309 cross-neutralization of SARS-CoV-2 and SARS-CoV

To study the mechanisms of S309-mediated neutralization, we characterized

the complex between the S309 Fab fragment and a prefusion stabilized SARS-CoV-2

S ectodomain trimer[6] using single-particle cryoEM. Similar to our previous study of apo

SARS-CoV-2 S[6], 3D classification of the cryoEM data enabled identification of two

structural states: a trimer with one S^B domain open and a closed trimer. We determined

3D reconstructions of the SARS-CoV-2 S ectodomain trimer with a single open S^B

domain and in a closed state (applying 3-fold symmetry), both with three S309 Fabs

bound, at 3.7 Å and 3.3 Å resolution, respectively (**Fig. 2a-c, Extended Data Fig. 5**

and Table 2). In parallel, we also determined a crystal structure of the S309 Fab at 3.3

Å resolution to assist model building (**Table 3**). The S309 Fab bound to the open S^B

domain is weakly resolved in the cryoEM map, due to marked conformational variability

of the upward pointing S^B domain, and was not modeled in density. The analysis below

is based on the closed state structure.

S309 recognizes a protein/glycan epitope on the SARS-CoV-2 S^B, distinct from

the receptor-binding motif. The epitope is accessible in both the open and closed S

states, explaining the stoichiometric binding of Fab to the S trimer (**Fig. 2a-c**). The

S309 paratope is composed of all six CDR loops that burie a surface area of ~1,050Å^2

at the interface with S^B through electrostatic interactions and hydrophobic contacts.

The 20-residue long CDRH3 sits atop the S^B helix comprising residues 337-344 and

also contacts the edge of the S^B five-stranded β-sheet (residues 356-361), overall

accounting for ~50% of the buried surface area (**Fig. 2d-e**). CDRL1 and CDRL2 extend

the epitope by interacting with the helix spanning residues 440-444 that is located near

the S 3-fold molecular axis. CDRH3 and CDRL2 sandwich the SARS-CoV-2 S glycan

at position N343 through contacts with the core fucose moiety (in agreement with the

detection of SARS-CoV-2 N343 core-fucosylated peptides by mass-spectrometry[34])

and to a lesser extent with the core N-acetyl-glucosamine (Fig. 2d). These latter

interactions bury an average surface of ~170 $Å^2$ and stabilize the N343 oligosaccharide

which is resolved to a much larger extent than in the apo SARS-CoV-2 S structures[6,9].

The structural data explain the S309 cross-reactivity between SARS-CoV-2 and

SARS-CoV as 19 out of 24 residues of the epitope are strictly conserved (**Fig. 2f** and

Extended Data Fig. 6a-b). $R346_{SARS-CoV-2}$, $R357_{SARS-CoV-2}$, $N354_{SARS-CoV-2}$ and

$L441_{SARS-CoV-2}$ are conservatively substituted to $K333_{SARS-CoV}$, $K344_{SARS-CoV}$ (except for

SARS-CoV isolate GZ02 where it is $R444_{SARS-CoV}$), $E341_{SARS-CoV}$ and $I428_{SARS-CoV}$

whereas $K444_{SARS-CoV-2}$ is semi-conservatively substituted to $T431_{SARS-CoV}$, in

agreement with the comparable binding affinities to SARS-CoV and SARS-CoV-2 S

(**Fig. 1c).** The oligosaccharide at position N343 is also conserved in both viruses and

corresponds to SARS-CoV N330, for which we previously detected core-fucosylated

glycopeptides by mass spectrometry[14] which would allow for similar interactions with

the S309 Fab. Analysis of the S glycoprotein sequences of the 2,229 SARS-CoV-2

isolates reported to date indicates that several mutations have occurred with variable

frequency on the SARS-CoV-2 S ectodomain (**Extended Data Fig. 7a-b**) but no

mutations arose within the epitope recognized by S309 mAb. Finally, S309 contact

residues showed a high degree of conservation across clade 1, 2 and 3 sarbecovirus

human and animal isolates[35] (**Extended Data Fig. 7c).** Collectively, the structural data

indicate that S309 could neutralize all SARS-CoV-2 isolates circulating to date and

possibly most other zoonotic sarbecoviruses.

Mechanism of S309-mediated neutralization of SARS-CoV-2 and SARS-CoV

The cryoEM structure of S309 bound to SARS-CoV-2 S presented here

combined with the structures of SARS-CoV-2 S^B and SARS-CoV S^B in complex with

$ACE2^{16-18,36}$ indicate that the Fab engages an epitope distinct from the receptor-binding

motif and would not clash with ACE2 upon binding to S **(Figure 3a-b)**. Biolayer

interferometry analysis of S309 Fab or IgG binding to the SARS-CoV-2 S^B domain or

the S ectodomain trimer confirmed the absence of competition between the mAb and

ACE2 for binding to SARS-CoV-2 S **(Figure 3c and Extended Data Fig. 8)**.

To further investigate the mechanism of S309-mediated neutralization, we

compared side-by-side transduction of SARS-CoV-2-MLV in the presence of either

S309 Fab or S309 IgG. Both experiments yielded comparable IC_{50} values (3.8 and 3.5

nM, respectively), indicating similar potencies for IgG and Fab **(Fig. 3d)**. However, The

S309 IgG reached 100% neutralization, whereas the S309 Fab plateaued at ~80%

neutralization **(Fig. 3d)**. This result indicates that one or more IgG-specific bivalent

mechanisms, such as S trimer cross-linking, steric hindrance or aggregation of

virions[37], may contribute to the ability to fully neutralize pseudovirions.

Fc-dependent effector mechanisms, such as NK-mediated antibody-dependent

cell cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis (ADCP) can

contribute to viral control in infected individuals. We observed efficient S309- and

S306-mediated ADCC of SARS-CoV-2 S-transfected cells, whereas the other mAbs

tested showed limited or no activity (**Fig. 3e and Extended Data Fig. 9a**). These

findings might be related to distinct binding orientations and/or positioning of the mAb

Fc fragment relative to the FcγRIIIa receptors. ADCC was observed only using NK

(effector) cells expressing the high-affinity FcγRIIIa variant (V158) but not the low-

affinity variant (F158) (**Fig. 3e**). These results, which we confirmed using a FcγRIIIa

cell reporter assay (**Fig. 3f**), suggest that S309 Fc engineering could potentially

enhance activation of NK cells with the low-affinity FcγRIIIa variant (F158)[38].

Macrophage or dendritic cell-mediated ADCP can contribute to viral control by clearing

virus and infected cells and by stimulating T cell response via presentation of viral

antigens[39,40]. Similar to the ADCC results, mAbs S309 and S306 showed the strongest

ADCP response (**Fig. 3g** and **Extended Data Fig. 8b)**. FcγRIIa signaling, however,

was only observed for S309 (**Fig. 3h**). These findings suggest that ADCP by

monocytes was dependent on both FcγRIIIa and FcγRIIa engagement. Collectively,

these results demonstrate that in addition to potent *in vitro* neutralization, S309 may

leverage additional protective mechanisms *in vivo*, as previously shown for other

antiviral antibodies[41,42].

MAb cocktails enhance SARS-CoV-2 neutralization

To gain more insight into the epitopes recognized by our panel of mAbs, we

used structural information, escape mutants analysis [23,27,30], and biolayer

inteferometry-based epitope binning to map the antigenic sites present on the SARS-

CoV and SARS-CoV-2 S^B domains (**Fig. 4a and Extended Data Fig.10**). This analysis

identified at least four antigenic sites within the S^B domain of SARS-CoV targeted by

our panel of mAbs. The receptor-binding motif, which is targeted by S230, S227 and

S110, is termed site I. Sites II and III are defined by S315 and S124, respectively, and

the two sites were bridged by mAb S304. Site IV is defined by S309, S109, and S303

mAbs. Given the lower number of mAbs cross-reacting with SARS-CoV-2, we were

able to identify sites IV targeted by S309 and S303, and site II-III targeted by S304 and

S315 (**Fig. 4b**).

Based on the above findings, we evaluated the neutralization potency of the site

IV S309 mAb in combination with either the site II S315 mAb or site II-III S304 mAb.

Although S304 and S315 alone were weakly neutralizing, the combination of either of

these mAbs with S309 resulted in an enhanced neutralization potency, compared to

single mAbs, against both SARS-CoV-2-MLV and authentic SARS-CoV-2 **(Fig. 4c-d**

and **Fig. 1e)**. A synergistic effect between two non-competing anti-RBD mAbs has

been already reported for SARS-CoV[43] and our data extend this observation to SARS-

CoV-2, providing a proof-of-concept for the use of mAbs combinations to prevent or

control SARS-CoV-2.

In summary, our study identifies S309 as a human mAb with broad neutralizing

activity against multiple sarbecoviruses, including SARS-CoV-2, via recognition of a

highly conserved epitope in the S^B domain comprising the N343-glycan (N330 in

SARS-CoV S). Furthermore, S309 can recruit effector mechanisms and synergizes

with weakly neutralizing mAbs, which may mitigate the risk of viral escape. Our data

indicate the potential to discover potently neutralizing pan-sarbecovirus mAbs,

highlight antigenic sites to include in vaccine design, and pave the way to support

preparedness for future sarbecovirus outbreaks. As S309 bears the promise to be an

effective countermeasure to curtail the COVID-19 pandemic caused by SARS-CoV-2,

Fc variants of S309 with increased half-life and effector functions have entered an

accelerated development path towards clinical trials.

ACKNOWLEDGEMENTS

This study was supported by the National Institute of General Medical Sciences (R01GM120553, D.V.), the National Institute of Allergy and Infectious Diseases (HHSN272201700059C to DV and 75N93019C00062 to MSD)), a Pew Biomedical Scholars Award (D.V.), an Investigators in the Pathogenesis of Infectious Disease Award from the Burroughs Wellcome Fund (D.V.), the University of Washington Arnold and Mabel Beckman cryoEM center, the Pasteur Institute (M.A.T.) and the beamline 5.0.1 at the Advanced Light Source at Lawrence Berkley National Laboratory. We gratefully acknowledge the authors, originating and submitting laboratories of the sequences from GISAID's EpiFlu™ Database upon which this research is based.

250
251

AUTHOR CONTRIBUTIONS

A.C.W., K.F., M.S.D., D.V. and D.C. designed the experiments. A.C.W., M.A.T., S.J., E.C. expressed and purified the proteins. K.C., F.Z., S.J., E.C. sequenced and expressed antibodies. D.P., M.B., A.C.W. and S.B. performed binding assays. D.P., M.B., A.C.W., A.P., A.D.M. carried out pseudovirus neutralization assays. J.B.C., R.E.C. performed neutralization assays with authentic SARS-CoV-2. B.G. performed effector function assays. Y.J.P. prepared samples for cryoEM and collected the data. Y.J.P. and D.V. processed the data, built and refined the atomic models. A.C.W. crystallized the S309 Fab. Y.J.P. collected and processed the X-ray diffraction data and built and refined the atomic model. R.S., A.T. and G.S. performed bioinformatic and conservation analysis. A.L. provided key reagents. A.C.W., K.F., C.H.D., H.W.V., A.L., D.V., D.C. analyzed the data and prepared the manuscript with input from all authors.

DECLARATION OF INTERESTS

D.P., S.B., K.C., E.C., C.H-D., G.S., M.B., A.K., K.F., A.P. F.Z., S.J., B.G., A.D.M., A.L., A.T., H.W.V, R.S. and D.C. are employees of Vir Biotechnology Inc. and may hold shares in Vir Biotechnology Inc. M.S.D. is a consultant for Inbios, Eli Lilly, Vir Biotechnology, NGM Biopharmaceuticals, and Emergent BioSolutions and on the Scientific Advisory Board of Moderna. The Diamond laboratory at Washington

MATERIALS AND METHODS

Ethics statement

Donors provided written informed consent for the use of blood and blood components

(such as sera), following approval by the Canton Ticino Ethics Committee, Switzerland.

Antibody discovery and expression

Monoclonal antibodies were isolated from EBV-immortalized memory B cells.

Recombinant antibodies were expressed in ExpiCHO cells transiently co-transfected

with plasmids expressing the heavy and light chain, as previously described[44]. Abs

S303, S304, S306, S309, S310 and S315 were expressed as rIgG-LS antibodies. The

LS mutation confers a longer half-life in vivo[45]. Antibodies S110 and S124 tested in

Fig. 1 and Extended Data Fig. 1 were purified mAbs produced from immortalized B

cells.

Transient expression of recombinant SARS-CoV-2 protein and flow cytometry

The full-length S gene of SARS-CoV-2 strain (SARS-CoV-2-S) isolate

BetaCoV/Wuhan-Hu-1/2019 (accession number MN908947) was codon optimized for

human cell expression and cloned into the phCMV1 expression vector (Genlantis).

Expi-CHO cells were transiently transfected with phCMV1-SARS-CoV-2-S, SARS-

spike_pcDNA.3 (strain SARS) or empty phCMV1 (Mock) using Expifectamine

CHO Enhancer. Two days after transfection, cells were collected for immunostaining

with mAbs. An Alexa647-labelled secondary antibody anti-human IgG Fc was used for

detection. Binding of mAbs to transfected cells was analyzed by flow-cytometry using a

ZE5 Cell Analyzer (Biorard) and FlowJo software (TreeStar). Positive binding was

defined by differential staining of CoV-S-transfectants versus mock-transfectants.

Affinity determination and competition experiments using Octet (BLI, biolayer

interferometry)

KD determination of full-length antibodies: Protein A biosensors (Pall ForteBio) were

used to immobilize recombinant antibodies at 2.7 μg/ml for 1min, after a hydration step

for 10 min with Kinetics Buffer (KB; 0.01% endotoxin-free BSA, 0.002% Tween-20,

0.005% NaN3 in PBS). Association curves were recorded for 5 minutes by incubating

the mAb-coated sensors with different concentration of SARS-CoV RBD (Sino

Biological) or SARS-CoV-2 RBD (produced in house; residues 331-550 of spike protein

from BetaCoV/Wuhan-Hu-1/2019, accession number MN908947). The highest RBD

concentration was 10 μg/ml, then serially diluted 1:2.5. Dissociation was recorded for

9 minutes by moving the sensors to wells containing KB. KD values were calculated

using a global fit model (Octet). Octet Red96 (ForteBio) equipment was used.

KD determination of full-length antibodies compared to Fab: His-tagged RBD of SARS-

CoV or SARS-CoV-2 were loaded at 3 µg/ml in KB for 15 minutes onto anti-HIS (HIS2)

biosensors (Molecular Devices, ForteBio). Association of mAb and Fab was performed

in KB at 15 μg/ml and 5 μg/ml respectively for 5 minutes. Dissociation in KB was

measured for 10 minutes.

MAbs competition experiments: His-tagged RBD of SARS-CoV or SARS-CoV-

2 was loaded for 5 minutes at 3 μg/ml in KB onto anti-Penta-HIS (HIS1K) biosensors

(Molecular Devices, ForteBio). Association of mAbs was performed in KB at 15 μg/ml.

ACE2 competition experiments: ACE2-His (Bio-Techne AG) was loaded for 30

minutes at 5 µg/ml in KB onto anti-HIS (HIS2) biosensors (Molecular Devices-

ForteBio).

SARS-CoV RBD-rabbitFc or SARS-CoV-2 RBD-mouseFc (Sino Biological Europe

GmbH) at 1 µg/ml was associated for 15 minutes, after a preincubation with or without

Ab (30 µg/ml, 30 minutes). Dissociation was monitored for 5 minutes.

ELISA

The following proteins were coated on 96 well ELISA plates at the following

concentrations: SARS-CoV RBD (Sino Biological, 40150-V08B1) at 1 µg/ml, SARS-

CoV-2 RBD (produced in house) at 10 µg/ml, ectodomains (stabilized prefusion trimer)

of SARS-CoV, SARS-CoV-2, OC43 and MERS, all at 1ug/ml. After blocking with 1%

BSA in PBS, antibodies es were added to the plates in a concentration range between

5 and 0.000028 μg/ml and incubated for 1 h at RT. Plates were washed and secondary

Ab Goat Anti Human IgG-AP (Southern Biotechnology: 2040-04) was added.

Substrate P-NitroPhenyl Phosphate (pNPP) (Sigma-Aldrich 71768) was used for
colour development. OD405 was read on an ELx808IU plate reader (Biotek).

Measurement of Fc-effector functions

ADCC assays were performed using ExpiCHO-S cells transient transfected with
SARS-CoV or SARS-CoV-2 S as targets. Target cells were incubated with titrated
concentrations of mAbs and after 10 minutes incubated with primary human NK cells
as effector cells at an effector:target ratio of 9:1. NK cells were isolated from fresh
blood of healthy donors using the MACSxpress NK Isolation Kit (Miltenyi Biotec, Cat.
Nr.: 130-098-185). ADCC was measured using LDH release assay (Cytotoxicity
Detection Kit (LDH) (Roche; Cat. Nr.: 11644793001) after 4 hours of incubation at
37°C.

ADCP assays were performed using ExpiCHO-S target cells transiently transfected
with SARS-CoV-2 S and fluorescently labeled with PKH67 Fluorescent Cell Linker Kits
(Sigma Aldrich, Cat. Nr.: MINI67) as targets. Target cells were incubated with titrated
concentrations of mAbs for 10minutes, followed by incubation with human PBMCs
isolated from healthy donors that were fluorescently labeled with Cell Trace Violet
(Invitrogen, Cat. Nr.: C34557) at an effector:target ratio of 20:1. After an overnight
incubation at 37°C, cells were stained with anti-human CD14-APC antibody (BD
Pharmingen, Cat. Nr.: 561708, Clone M5E2) to stain monocytes. Antibody-mediated
phagocytosis was determined by flow cytometry, gating on CD14$^+$ cells that were
double positive for cell trace violet and PKH67.

Determination of mAb-dependent activation of human FcγRIIIa or FcγRIIa was
performed using ExpiCHO cells transiently transfected with SARS-CoV-2 S
(BetaCoV/Wuhan-Hu-1/2019), incubated with titrated concentrations of mAbs for 10
minutes. ExpiCHO cells then were incubated with Jurkat cells expressing FcγRIIIa
receptor or FcγRIIa on their surface and stably transfected with NFAT-driven luciferase
gene (Promega, Cat. Nr.: G9798 and G7018) at an effector to target ratio of 6:1 for
FcγRIIIa and 5:1 for FcγRIIa. Activation of human FcγRs in this bioassay results in the
NFAT-mediated expression of the luciferase reporter gene. Luminescence was
measured after 21 hours of incubation at 37°C with 5% CO_2, using the Bio-Glo-TM
Luciferase Assay Reagent according to the manufacturer's instructions.

Pseudovirus neutralization assays

Murine leukemia virus (MLV)-based SARS-CoV S-pseudotyped viruses were prepared as previously described[6,32]. HEK293T cells were co-transfected with a SARS-CoV, SARS-CoV-2, CUHK, GZ02, or WiV1 S encoding-plasmid, an MLV Gag-Pol packaging construct and the MLV transfer vector encoding aluciferase reporter using the Lipofectamine 2000 transfection reagent (Life Technologies) according to the manufacturer's instructions. Cells were incubated for 5 hours at 37°C with 8% CO_2 with OPTIMEM transfection medium. DMEM containing 10% FBS was added for 72 hours. VeroE6 cells or DBT cells transfected with human ACE2 were cultured in DMEM containing 10% FBS, 1% PenStrep and plated into 96 well plates for 16-24 hours. Concentrated pseudovirus with or without serial dilution of antibodies was incubated for 1 hour and then added to the wells after washing 3X with DMEM. After 2-3 hours DMEM containing 20% FBS and 2% PenStrep was added to the cells for 48 hours. Following 48 hours of infection, One-Glo-EX (Promega) was added to the cells and incubated in the dark for 5-10 minutes prior to reading on a Varioskan LUX plate reader (ThermoFisher). Measurements were done in duplicate and relative luciferase units (RLU) were converted to percent neutralization and plotted with a non-linear regression curve fit in PRISM.

Live virus neutralization assay

SARS-CoV-2 strain 2019-nCoV/USA_WA1/2020 was obtained from the Centers for Disease Control and Prevention (gift of Natalie Thornburg). Virus was passaged once in Vero CCL81 cells (ATCC) and titrated by focus-forming assay on Vero E6 cells. Serial dilutions of indicated mAbs were incubated with 10^2 focus forming units (FFU) of SARS-CoV-2 for 1 hour at 37°C. MAb-virus complexes were added to Vero E6 cell monolayers in 96-well plates and incubated at 37°C for 1 hour. Subsequently, cells were overlaid with 1% (w/v) methylcellulose in MEM supplemented with 2% FBS. Plates were harvested 30 hours later by removing overlays and fixed with 4% PFA in PBS for 20 minutes at room temperature. Plates were washed and sequentially incubated with 1 µg/mL of CR3022[46] anti-S antibody and HRP-conjugated goat anti-human IgG in PBS supplemented with 0.1% saponin and 0.1% BSA. SARS-CoV-2-infected cell foci were visualized using TrueBlue peroxidase substrate (KPL) and

404   quantitated on an ImmunoSpot microanalyzer (Cellular Technologies). Data were
405   processed using Prism software (GraphPad Prism 8.0).
406
Recombinant Spike ectodomain production
The SARS-CoV-2 2P S (Genbank: YP_009724390.1) ectodomain was produced in
500mL cultures of HEK293F cells grown in suspension using FreeStyle 293 expression
medium (Life technologies) at 37°C in a humidified 8% CO2 incubator rotating at 130
r.p.m, as previously reported[6]. The culture was transfected using 293fectin
(ThermoFisher Scientific) with cells grown to a density of 10^6 cells per mL and
cultivated for three days. The supernatant was harvested and cells were resuspended
for another three days, yielding two harvests. Clarified supernatants were purified
using a 5mL Cobalt affinity column (Takara). Purified protein was filtered or
concentrated and flash frozen in a buffer containing 50 mM Tris pH 8.0 and 150 mM
NaCl prior to cryoEM analysis. The SARS-CoV S, HCoV-OC43 S and MERS-CoV S
constructs were previously described[14,28] and produced similarly to SARS-CoV-2 2P
S.
420
CryoEM sample preparation and data collection.
3 μL of SARS-CoV-2 S at 1.6 mg/mL was mixed with 0.45 μL of S309 Fab at 7.4 mg/mL
for 1 min at room temperature before application onto a freshly glow discharged 1.2/1.3
UltraFoil grid (300 mesh). Plunge freezing used a vitrobot MarkIV (ThermoFisher
Scientific) using a blot force of 0 and 6.5 second blot time at 100% humidity and 25°C.
Data were acquired using the Leginon software [47] to control an FEI Titan Krios
transmission electron microscope operated at 300 kV and equipped with a Gatan K2
Summit direct detector and Gatan Quantum GIF energy filter, operated in zero-loss
mode with a slit width of 20 eV. Automated data collection was carried out using
Leginon at a nominal magnification of 130,000x with a pixel size of 0.525 Å with tilt
angles ranging between 20° and 50°, as previously described[48]. The dose rate was
adjusted to 8 counts/pixel/s, and each movie was acquired in super-resolution mode
fractionated in 50 frames of 200 ms. 3,900 micrographs were collected in a single
session with a defocus range comprised between 1.0 and 3.0 μm.
435
436
437

CryoEM data processing

Movie frame alignment, estimation of the microscope contrast-transfer function parameters, particle picking and extraction were carried out using Warp [49]. Particle images were extracted with a box size of 800 binned to 400 yielding a pixel size of 1.05 Å. For each data set two rounds of reference-free 2D classification were performed using cryoSPARC [50] to select well-defined particle images. Subsequently, two rounds of 3D classification with 50 iterations each (angular sampling 7.5° for 25 iterations and 1.8° with local search for 25 iterations), using our previously reported closed SARS-CoV-2 S structure[6] as initial model, were carried out using Relion [51] without imposing symmetry to separate distinct SARS-CoV-2 S conformations. 3D refinements were carried out using non-uniform refinement along with per-particle defocus refinement in cryoSPARC[50]. Particle images were subjected to Bayesian polishing [52] before performing another round of non-uniform refinement in cryoSPARC [50] followed by per-particle defocus refinement and again non-uniform refinement. Reported resolutions are based on the gold-standard Fourier shell correlation (FSC) of 0.143 criterion and Fourier shell correlation curves were corrected for the effects of soft masking by high-resolution noise substitution[53].

CryoEM model building and analysis.

UCSF Chimera [54] and Coot were used to fit atomic models (PDB 6VXX and PDB 6VYB) into the cryoEM maps. The Fab was subsequently manually built using Coot[55,56]. N-linked glycans were hand-built into the density where visible and the models were refined and relaxed using Rosetta[57]. Glycan refinement relied on a dedicated Rosetta protocol, which uses physically realistic geometries based on prior knowledge of saccharide chemical properties [58], and was aided by using both sharpened and unsharpened maps. Models were analyzed using MolProbity [59], EMringer [60], Phenix [61] and privateer [62] to validate the stereochemistry of both the protein and glycan components. Figures were generated using UCSF ChimeraX [63].

Crystallization and X-ray structure determination of Fab S309

Fab S309 crystals were grown in hanging drop set up with a mosquito at 20°C using 150 nL protein solution in Tris HCl pH 8.0, 150 mM NaCl and 150nL mother liquor solution containing 1.1 M Sodium Malonate, 0.1 M HEPES, pH 7.0 and 0.5% (w/v) Jeffamine ED-2001. Crystals were cryo-protected using the mother liquor solution

supplemented with 30% glycerol. The dataset was collected at ALS beamline 5.0.2
and processed to 3.3 Å resolution in space group $P4_12_12$ using mosflm[64] and
Aimless[65]. The structure of Fab S309 was solved by molecular replacement using
Phaser[66] and homology models as search models. The coordinates were improved
and completed using Coot[55] and refined with REFMAC5[67]. Crystallographic data
collection and refinement statistics are shown in Table 3.
478

Sequence alignment
SARS-CoV-2 genomics sequences were downloaded from GISAID on March
29th 2020, using the "complete (>29,000 bp)" and "low coverage exclusion" filters. Bat
and pangolin sequences were removed to yield human-only sequences. The spike
ORF was localized by performing reference protein (YP_009724390.1)-genome
alignments with GeneWise2. Incomplete matches and indel-containing ORFs were
rescued and included in downstream analysis. Nucleotide sequences were
translated *in silico* using seqkit. Sequences with more than 10% undetermined
aminoacids (due to N basecalls) were removed. Multiple sequence alignment was
performed using MAFFT. Variants were determined by comparison of aligned
sequences (n=2,229) to the reference sequence using the R/Bioconductor package
Biostrings. A similar strategy was used to extract and translate spike protein
sequences from SARS-CoV genomes sourced from ViPR (search criteria: SARS-
related coronavirus, full-length genomes, human host, deposited before December
2019 to exclude SARS-CoV-2, n=53). Sourced SARS-CoV genome sequences
comprised all the major published strains, such as Urbani, Tor2, TW1, P2, Frankfurt1,
among others. Pangolin sequences as shown by Tsan-Yuk Lam et al[68] were sourced
from GISAID. Bat sequences from the three clades of sarbecoviruses as shown by Lu
et al[35] were sourced from Genbank. Civet and racoon dog sequences were similarly
sourced from Genbank.
499
500

501
References

1 Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of
probable bat origin. *Nature*, doi:10.1038/s41586-020-2012-7 (2020).
2 Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China,
2019. *N Engl J Med*, doi:10.1056/NEJMoa2001017 (2020).
3 Tortorici, M. A. & Veesler, D. Structural insights into coronavirus entry. *Adv Virus
Res* **105**, 93-116, doi:10.1016/bs.aivir.2019.08.002 (2019).
4 Walls, A. C. *et al.* Cryo-electron microscopy structure of a coronavirus spike
glycoprotein trimer. *Nature* **531**, 114-117, doi:10.1038/nature16988
10.1038/nature16988. Epub 2016 Feb 8. (2016).
5 Walls, A. C. *et al.* Tectonic conformational changes of a coronavirus spike
glycoprotein promote membrane fusion. *Proceedings of the National Academy
of Sciences* **114**, 11157-11162, doi:10.1073/pnas.1708727114 (2017).
6 Walls, A. C. *et al.* Structure, function and antigenicity of the SARS-CoV-2 spike
glycoprotein. *bioRxiv*, 2020.2002.2019.956581,
doi:10.1101/2020.02.19.956581 (2020).
7 Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2
and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*,
doi:10.1016/j.cell.2020.02.052 (2020).
8 Letko, M., Marzi, A. & Munster, V. Functional assessment of cell entry and
receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nature
Microbiology*, doi:10.1038/s41564-020-0688-y (2020).
9 Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion
conformation. *Science*, doi:10.1126/science.abb2507 (2020).
10 Li, W. *et al.* Receptor and viral determinants of SARS-coronavirus adaptation to
human ACE2. *EMBO J* **24**, 1634-1643, doi:10.1038/sj.emboj.7600640
10.1038/sj.emboj.7600640. Epub 2005 Mar 24. (2005).
11 Rossen, J. W. *et al.* The viral spike protein is not involved in the polarized sorting
of coronaviruses in epithelial cells. *J Virol* **72**, 497-503 (1998).
12 Walls, A. C. *et al.* Glycan shield and epitope masking of a coronavirus spike
protein observed by cryo-electron microscopy. *Nat Struct Mol Biol* **23**, 899-905,
doi:10.1038/nsmb.3293
10.1038/nsmb.3293. Epub 2016 Sep 12. (2016).
13 Xiong, X. *et al.* Glycan shield and fusion activation of a deltacoronavirus spike
glycoprotein fine-tuned for enteric infections. *J Virol*, doi:10.1128/JVI.01628-17
10.1128/JVI.01628-17. (2017).
14 Walls, A. C. *et al.* Unexpected Receptor Functional Mimicry Elucidates
Activation of Coronavirus Fusion. *Cell* **176**, 1026-1039 e1015,
doi:10.1016/j.cell.2018.12.028
10.1016/j.cell.2018.12.028. Epub 2019 Jan 31. (2019).
15 Yang, Y. *et al.* Two Mutations Were Critical for Bat-to-Human Transmission of
Middle East Respiratory Syndrome Coronavirus. *J Virol* **89**, 9119-9123,
doi:10.1128/JVI.01279-15
10.1128/JVI.01279-15. Epub 2015 Jun 10. (2015).
16 Yan, R. *et al.* Structural basis for the recognition of SARS-CoV-2 by full-length
human ACE2. *Science* **367**, 1444-1448, doi:10.1126/science.abb2762 (2020).
17 Shang, J. *et al.* Structural basis of receptor recognition by SARS-CoV-2. *Nature*,
doi:10.1038/s41586-020-2179-y (2020).

18   Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound
to the ACE2 receptor. *Nature*, doi:10.1038/s41586-020-2180-5 (2020).
19   Corti, D. *et al.* Protective monotherapy against lethal Ebola virus infection by a
potently   neutralizing   antibody.   *Science*   **351**,   1339-1342,
doi:10.1126/science.aad5224 (2016).
20   Levine, M. M. Monoclonal Antibody Therapy for Ebola Virus Disease. *N Engl J*
Med **381**, 2365-2366, doi:10.1056/NEJMe1915350 (2019).
21   Traggiai, E. *et al.* An efficient method to make human monoclonal antibodies
from memory B cells: potent neutralization of SARS coronavirus. *Nat Med* **10**,
871-875, doi:10.1038/nm1080
10.1038/nm1080. Epub 2004 Jul 11. (2004).
22   Corti, D. *et al.* Prophylactic and postexposure efficacy of a potent human
monoclonal antibody against MERS coronavirus. *Proc Natl Acad Sci U S A* **112**,
10473-10478, doi:10.1073/pnas.1510199112
10.1073/pnas.1510199112. Epub 2015 Jul 27. (2015).
23   Rockx, B. *et al.* Structural basis for potent cross-neutralizing human monoclonal
antibody protection against lethal human and zoonotic severe acute respiratory
syndrome   coronavirus   challenge.   *J   Virol*   **82**,   3220-3235,
doi:10.1128/JVI.02377-07
10.1128/JVI.02377-07. Epub 2008 Jan 16. (2008).
24   Menachery, V. D. *et al.* SARS-like WIV1-CoV poised for human emergence.
Proc Natl Acad Sci U S A **113**, 3048-3053, doi:10.1073/pnas.1517719113
10.1073/pnas.1517719113. Epub 2016 Mar 14. (2016).
25   Yuan, M. *et al.* A highly conserved cryptic epitope in the receptor-binding
domains   of   SARS-CoV-2   and   SARS-CoV.   *Science*,
doi:10.1126/science.abb7269 (2020).
26   Joyce, M. G. *et al.* A Cryptic Site of Vulnerability on the Receptor Binding
Domain   of   the   SARS-CoV-2   Spike   Glycoprotein.   *bioRxiv*,
2020.2003.2015.992883, doi:10.1101/2020.03.15.992883 (2020).
27   Rockx, B. *et al.* Escape from human monoclonal antibody neutralization affects
in vitro and in vivo fitness of severe acute respiratory syndrome coronavirus. *J*
Infect Dis **201**, 946-955, doi:10.1086/651022
10.1086/651022. (2010).
28   Tortorici, M. A. *et al.* Structural basis for human coronavirus attachment to sialic
acid receptors. *Nat Struct Mol Biol* **26**, 481-489, doi:10.1038/s41594-019-0233-
y (2019).
29   Park, Y. J. *et al.* Structures of MERS-CoV spike glycoprotein in complex with
sialoside   attachment   receptors.   *Nat   Struct   Mol   Biol*   **26**,   1151-1157,
doi:10.1038/s41594-019-0334-7 (2019).
30   Walls, A. C. *et al.* Unexpected Receptor Functional Mimicry Elucidates
Activation   of   Coronavirus   Fusion.   *Cell*   **176**,   1026-1039.e1015,
doi:10.1016/j.cell.2018.12.028 (2019).
31   Liu, C. *et al.* Viral Architecture of SARS-CoV-2 with Post-Fusion Spike Revealed
by   Cryo-EM.   *bioRxiv*,   2020.2003.2002.972927,
doi:10.1101/2020.03.02.972927 (2020).
32   Millet, J. K. & Whittaker, G. R. Murine Leukemia Virus (MLV)-based Coronavirus
Spike-pseudotyped Particle Production and Infection. *Bio Protoc* **6**,
doi:10.21769/BioProtoc.2035
10.21769/BioProtoc.2035. (2016).

33   Ge, X. Y. *et al.* Isolation and characterization of a bat SARS-like coronavirus
that uses the ACE2 receptor. *Nature* **503**, 535-538, doi:10.1038/nature12711
10.1038/nature12711. Epub 2013 Oct 30. (2013).
34   Watanabe, Y., Allen, J. D., Wrapp, D., McLellan, J. S. & Crispin, M. Site-specific
analysis of the SARS-CoV-2 glycan shield. *bioRxiv*, 2020.2003.2026.010322,
doi:10.1101/2020.03.26.010322 (2020).
35   Lu, R. *et al.* Genomic characterisation and epidemiology of 2019 novel
coronavirus: implications for virus origins and receptor binding. *Lancet* **395**,
565-574, doi:10.1016/S0140-6736(20)30251-8 (2020).
36   Li, F., Li, W., Farzan, M. & Harrison, S. C. Structure of SARS coronavirus spike
receptor-binding domain complexed with receptor. *Science* **309**, 1864-1868,
doi:10.1126/science.1116480 (2005).
37   Klasse, P. J. & Sattentau, Q. J. Occupancy and mechanism in antibody-
mediated neutralization of animal viruses. *J Gen Virol* **83**, 2091-2108,
doi:10.1099/0022-1317-83-9-2091 (2002).
38   Wang, X., Mathieu, M. & Brezski, R. J. IgG Fc engineering to modulate antibody
effector functions. *Protein Cell* **9**, 63-73, doi:10.1007/s13238-017-0473-8
(2018).
39   He, W. *et al.* Alveolar macrophages are critical for broadly-reactive antibody-
mediated protection against influenza A virus in mice. *Nat Commun* **8**, 846,
doi:10.1038/s41467-017-00928-3 (2017).
40   DiLillo, D. J. & Ravetch, J. V. Differential Fc-Receptor Engagement Drives an
Anti-tumor Vaccinal Effect. *Cell* **161**, 1035-1045, doi:10.1016/j.cell.2015.04.016
(2015).
41   Corti, D. *et al.* A neutralizing antibody selected from plasma cells that binds to
group 1 and group 2 influenza A hemagglutinins. *Science* **333**, 850-856,
doi:10.1126/science.1205669 (2011).
42   Hessell, A. J. *et al.* Fc receptor but not complement binding is important in
antibody protection against HIV. *Nature* **449**, 101-104,
doi:10.1038/nature06106 (2007).
43   ter Meulen, J. *et al.* Human monoclonal antibody combination against SARS
coronavirus: synergy and coverage of escape mutants. *PLoS Med* **3**, e237,
doi:10.1371/journal.pmed.0030237 (2006).
44   Stettler, K. *et al.* Specificity, cross-reactivity, and function of antibodies elicited
by Zika virus infection. *Science* **353**, 823-826, doi:10.1126/science.aaf8505
(2016).
45   Zalevsky, J. *et al.* Enhanced antibody half-life improves in vivo activity. *Nat
Biotechnol* **28**, 157-159, doi:10.1038/nbt.1601 (2010).
46   Tian, X. *et al.* Potent binding of 2019 novel coronavirus spike protein by a SARS
coronavirus-specific human monoclonal antibody. *Emerg Microbes Infect* **9**,
382-385, doi:10.1080/22221751.2020.1729069 (2020).
47   Suloway, C. *et al.* Automated molecular microscopy: the new Leginon system.
Journal of structural biology **151**, 41-60, doi:10.1016/j.jsb.2005.03.010 (2005).
48   Tan, Y. Z. *et al.* Addressing preferred specimen orientation in single-particle
cryo-EM through tilting. *Nat Methods* **14**, 793-796, doi:10.1038/nmeth.4347
(2017).
49   Tegunov, D. & Cramer, P. Real-time cryo-electron microscopy data
preprocessing with Warp. *Nature methods*, doi:10.1038/s41592-019-0580-y
(2019).

50   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC:
algorithms for rapid unsupervised cryo-EM structure determination. *Nature*
methods **14**, 290-296, doi:10.1038/nmeth.4169 (2017).
51   Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure
determination in RELION-3. *Elife* **7**, doi:10.7554/eLife.42166
10.7554/eLife.42166. (2018).
52   Zivanov, J., Nakane, T. & Scheres, S. H. W. A Bayesian approach to beam-
induced motion correction in cryo-EM single-particle analysis. *IUCrJ* **6**,
doi:doi:10.1107/S205225251801463X (2019).
53   Scheres, S. H. & Chen, S. Prevention of overfitting in cryo-EM structure
determination. *Nature methods* **9**, 853-854, doi:10.1038/nmeth.2115 (2012).
54   Goddard, T. D., Huang, C. C. & Ferrin, T. E. Visualizing density maps with UCSF
Chimera. *Journal of structural biology* **157**, 281-287,
doi:10.1016/j.jsb.2006.06.010 (2007).
55   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development
of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501,
doi:10.1107/S0907444910007493 (2010).
56   Brown, A. *et al.* Tools for macromolecular model building and refinement into
electron cryo-microscopy reconstructions. *Acta Crystallogr D Biol Crystallogr*
71, 136-153, doi:10.1107/S1399004714021683 (2015).
57   Wang, R. Y. *et al.* Automated structure refinement of macromolecular
assemblies from cryo-EM maps using Rosetta. *eLife* **5**, doi:10.7554/eLife.17219
(2016).
58   Frenz, B. D. R., S.; Borst A.J.; Walls, A.C.; Adolf-Bryfogle, J.; Schief, W.R.;
Veesler, D.; DiMaio, F. Automatically fixing errors in glycoprotein structures with
Rosetta. *Structure* **27**, 1-6, doi:https://doi.org/10.1016/j.str.2018.09.006 (2019).
59   Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12-21,
doi:10.1107/S0907444909042073 (2010).
60   Barad, B. A. *et al.* EMRinger: side chain-directed model and map validation for
3D cryo-electron microscopy. *Nat Methods* **12**, 943-946,
doi:10.1038/nmeth.3541
10.1038/nmeth.3541. Epub 2015 Aug 17. (2015).
61   Liebschner, D. *et al.* Macromolecular structure determination using X-rays,
neutrons and electrons: recent developments in Phenix. *Acta Crystallogr D*
Struct Biol **75**, 861-877, doi:10.1107/S2059798319011471 (2019).
62   Agirre, J. *et al.* Privateer: software for the conformational validation of
carbohydrate structures. *Nature structural & molecular biology* **22**, 833-834,
doi:10.1038/nsmb.3115 (2015).
63   Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in
visualization and analysis. *Protein Sci* **27**, 14-25, doi:10.1002/pro.3235 (2018).
64   Battye, T. G., Kontogiannis, L., Johnson, O., Powell, H. R. & Leslie, A. G.
iMOSFLM: a new graphical interface for diffraction-image processing with
MOSFLM. *Acta Crystallogr D Biol Crystallogr* **67**, 271-281,
doi:10.1107/S0907444910048675 (2011).
65   Evans, P. R. & Murshudov, G. N. How good are my data and what is the
resolution? *Acta Crystallogr D Biol Crystallogr* **69**, 1204-1214,
doi:10.1107/S0907444913000061 (2013).
66   McCoy, A. J. *et al.* Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-
674, doi:10.1107/S0021889807021206 (2007).

67   Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal
699   structures. *Acta Crystallogr D Biol Crystallogr* **67**, 355-367,
700   doi:10.1107/S0907444911001314 (2011).
701   68   Lam, T. T. *et al.* Identifying SARS-CoV-2 related coronaviruses in Malayan
702   pangolins. *Nature*, doi:10.1038/s41586-020-2169-0 (2020).
703

Figure 1

Figure 1: Identification of a potent SARS-CoV-2 neutralizing mAb from a SARS survivor. a-b, Binding of a panel of mAbs isolated from a SARS-immune patient to the SARS-CoV-2 (**a**) or SARS-CoV (**b**) S glycoproteins expressed at the surface of ExpiCHO cells (symbols are means of duplicates from one experiment). **c-d**, Affinity measurement of S309 full-length IgG1 and Fab for SARS-CoV-2 and SARS-CoV S^B domains measured using biolayer interferometry. **e**, Neutralization of SARS-CoV-2-MLV, SARS-CoV-MLV (bearing S from various isolates) and other sarbecovirus isolates by mAb S309. **f**, Neutralization of authentic SARS-CoV-2 (strain n-CoV/USA_WA1/2020) by mAbs as measured by a focus-forming assay on Vero E6 cells. (**e-f**) mean±SD (**e**) or means (**f**) of duplicates are shown. One representative out of two experiments is shown.

Figure 2

Figure 2: CryoEM structures of the SARS-CoV-2 S glycoprotein in complex with the S309 neutralizing antibody Fab fragment. a, Ribbon diagram of the partially open SARS-CoV-2 S trimer (one S^B domain is open) bound to three S309 Fabs. **b-c,** Ribbon diagrams of the closed SARS-CoV-2 S trimer bound to three S309 Fabs shown in two orthogonal orientations. **d,** Close-up view of the S309 epitope showing the contacts formed with the core fucose (labeled with a star) and the core N-acetyl-glucosamine of the oligosaccharide at position N343. **e,** Close-up view of the S309 epitope showing the 20-residue long CDRH3 siting atop the S^B helix comprising residues 337-344. The oligosaccharide at position N343 is omitted for clarity. In panels (**c-d**), selected residues involved in interactions between S309 and SARS-CoV-2 S are shown. **F,** Molecular surface representation of the SARS-CoV-2 S trimer showing the S309 footprint colored by residue conservation on one protomer among SARS-CoV-2 and SARS-CoV S glycoproteins. The other two protomers are colored pink and gold.

Figure 3

Figure 3: Mechanism of neutralization of S309 mAb. See next page for caption

Figure 3: Mechanism of neutralization of S309 mAb. **a-b**, Ribbon diagrams of S309 and ACE2 bound to SARS-CoV-2 S^B. This composite model was generated using the SARS-CoV-2 S/S309 cryoEM structure reported here and a crystal structure of SARS-CoV-2 S bound to ACE2[16]. **c**, Competition of S309 or S230 mAbs with ACE2 to bind to SARS-CoV S^B (left panel) and SARS-CoV-2 S^B (right panel). ACE2 was immobilized at the surface of biosensors before incubation with S^B domain alone or S^B precomplexed with mAbs. The vertical dashed line indicates the start of the association of mAb-complexed or free S^B to solid-phase ACE2. **d,** Neutralization of SARS-CoV-MLV by S309 IgG1 or S309 Fab, plotted in nM (means $\pm$SD is shown, one out of two experiments is shown). **e,** mAb-mediated ADCC using primary NK effector cells and SARS-CoV-2 S-expressing ExpiCHO as target cells. Bar graph shows the average area under the curve (AUC) for the responses of 3-4 donors genotyped for their FcγRIIIa (mean$\pm$SD, from two independent experiments). **f,** Activation of high affinity (V158) or low affinity (F158) FcγRIIIa was measured using Jurkat reporter cells and SARS-CoV-2 S-expressing ExpiCHO as target cells (one experiment, one or two measurements per mAb). **g,** mAb-mediated ADCP using Cell Trace Violet-labelled PBMCs as phagocytic cells and PKF67-labelled SARS-CoV-2 S-expressing ExpiCHO as target cells. Bar graph shows the average area under the curve (AUC) for the responses of four donor (mean$\pm$SD, from two independent experiments). **h,** Activation of FcγRIIa measured using Jurkat reporter cells and SARS-CoV-2 S-expressing ExpiCHO as target cells (one experiment, one or two measurements per mAb).

Figure 4

Figure 4: MAb cocktails enhance SARS-CoV-2 neutralization. a, Heat map showing the competition of mAb pairs for binding to the SARS-CoV S^B domain as measured by biolayer interferometry (as shown in Extended Data Fig. 9). **b,** Competition of mAb pairs for binding to the SARS-CoV-2 S^B domain . **c-d,** Neutralization of SARS-CoV-2-MLV by S309 combined with an equimolar amount of S304 or S315 mAbs. For mAb cocktails the concentration on the x axis is that of the individual mAbs.

Table 1: Characteristics of the antibodies described in this study. VH and VL % identity refers to V gene identity compared to germline (IMGT).

mAb	Blood sample date	VH (% identity)	HCDR3 Lenght	HCDR3 sequence	VL (% identity)	LCDR3 sequence	SARS-CoV	SARS-CoV-2	Specificity
S110	2004	VH3-30 (96.88)	18	AKDRFQFARSWYGDYFDY	VK2-30 (96.60)	MQGTHWPPT	+	+	RBD/non-RBD
S124	2004	VH2-26 (98.28)	17	ARINTAAYDYDSTTFDI	VK1-39 (98.57)	QQSYSTPPT	+	+	RBD
S109	2004	VH3-23 (93.75)	19	ARLESATQPLGYYFYGMDV	VL3-25 (97.85)	HSADISATSWV	+	-	RBD
S111	2004	VH3-30 (95.14)	16	ARDIRHLIVVVSDMDV	VK2-30 (98.30)	MQGTHWPPT	+	-	RBD
S127	2004	VH3-30 (96.53)	18	AKDLFGYCRSTSCESLDD	VK1-9 (98.92)	QQLNNYPLT	+	-	RBD
S215	2004	VH3-30 (90.28))	16	ARETRHYSHGLNWFDP	VK3-15 (98.92)	QQYNNWPTT	+	-	RBD
S217	2004	VH3-49 (95.58)	8	SWIHRIVS	VK1-33 (98.21)	QQYDNLPYT	+	-	RBD
S218	2004	VH3-30 (93.40)	16	ARDVKGHIVVMTSLDY	VK2-30 (97.62)	MQGTHWPPT	+	-	RBD
S219	2004	VH1-58(92.01)	18	AAEMATIQNYYYYYGMDV	VK1-39 (95.34)	QQSYSTPPT	+	-	RBD
S222	2006	VH1-2 (91.67)	15	ARGDVPVGTGWVFDF	VK1-39 (92.47)	QQSLSMVT	+	-	RBD
S223	2006	VH3-30 (95.14)	19	ATVSVEGYTSGWYLGTLDF	VK3-15 (98.21)	QQYNNWPGT	+	-	RBD
S224	2006	VH1-18 (90.97)	15	ARQSHSTRGGWHFSP	VK1-39 (95.70)	QQSYSVPYT	+	-	RBD
S225	2006	VH3-9 (96.18)	20	AKDISLVFWSVNPPRNGMDV	VK1-39 (98.57)	QQSYSSPLT	+	-	RBD
S226	2006	VH3-30 (89.61)	18	ARDSSWQSTGWPINWFDR	VK3-11 (96.11)	QQRSNWPPT	+	-	RBD
S227	2006	VH3-23 (95.14)	12	ASPLRNYGDLLY	VK1-5 (96.06)	QQYNSYSWT	+	-	RBD
S228	2006	VH3-30 (96.53)	16	ARDLQMRVVVVSNFDY	VK2D-30 (99.32)	MQATHWPPT	+	-	RBD
S230	2006	VH3-30 (90.97)	20	VTQRDNSRDYFPHYFHDMDV	VK2-30 (97.62)	MQGSHWPPT	+	-	RBD
S231	2006	VH3-30 (90.62)	17	ARDDNLDRHWPLRLGGY	VK2-30 (94.56)	MQGAHWPPT	+	-	RBD
S237	2006	VH3-21 (96.53)	11	ARGFERYYFDS	VL1-44 (96.84)	VAWDDILNAVV	+	-	RBD
S309	2013	VH1-18 (97.22)	20	ARDYTRGAWFGESLIGGFDN	VK3-20 (97.52)	QQHDTSLT	+	+	RBD
S315	2013	VH3-7 (97.92)	17	ARDLWWNDQAHYYGMDV	VL3-25 (97.57)	QSADSSGTV	+	+	RBD
S303	2013	VH3-23 (90.28)	17	ARERDDIFPMGLNAFDI	VK1-5 (97.49)	QQYDTYSWT	+	+	RBD
S304	2013	VH3-13 (97.89)	14	ARGDSSGYYYYFDY	VK1-39 (93.55)	QQSYVSPTYT	+	+	RBD
S306	2013	VH1-18 (95.49)	16	ASDYFDSSGYYHSFDY	VK3-11 (98.92)	QQRSNWPPGCS	+	+	non-RBD
S310	2013	VH1-69 (92.71)	19	ATRTYDSSGYRPYYYGLDV	VL2-23 (97.57)	CSYAGSDTVI	+	+	non-RBD

Table 2. CryoEM data collection and refinement statistics.

	SARS-CoV-2 S + S309 (closed)	SARS-CoV-2 S + S309 (one S^B open)
Data collection and processing		
Magnification	130,000	130,000
Voltage (kV)	300	300
Electron exposure (e$^-$/Å^2)	70	70
Defocus range (μm)	0.5-3.0	0.5-3.0
Pixel size (Å)	0.525	0.525
Symmetry imposed	C3	C1
Final particle images (no.)	168,449	119,608
Map resolution (Å)	3.3	3.7
FSC threshold	0.143	0.143
Map sharpening B factor (Å^2)	-91	-69
Validation		
MolProbity score	0.91	
Clashscore	0.9	
Poor rotamers (%)	0.1	
Ramachandran plot		
Favored (%)	97.24	
Allowed (%)	99.91	
Disallowed (%)	0.09	
EMRinger Score	2.58	

Table 3. X-ray data collection and refinement statistics.

	Fab S309
Data collection	
Space group	$P4_12_12$
Cell constants	
a,b,c (Å)	132.6, 132.6, 301.2
α,β,γ (˚)	90, 90, 90
Wavelength (Å)	0.9812
Resolution (Å)	68.6 - 3.3 (3.48 - 3.30)
Rmerge	18 (75)
I/σ(I)	13.2 (2)
CC(1/2)	99.0 (33)
Completeness (%)	99.4 (99.0)
Redundancy	12
Refinement	
Resolution (Å)	68.6 - 3.3
Unique reflections	41,395
Rwork/Rfree (%)	20.5 / 23.8
Number of protein atoms	8,347
Number of water atoms	0
R.m.s.d. bond lengths (Å)	0.06
R.m.s.d. bond angles (˚)	1.45
Favored Ramachandran residues (%)	96
Allowed Ramachandran residues (%)	3.54
Disallowed Ramachandran residues (%)	0.46

[1]Numbers in parentheses refer to outer resolution shell