Population connectivity predicts vulnerability to white-nose syndrome in the Chilean myotis (*Myotis chiloensis*) - a genomics approach

Thomas M. Lilley, *[1] Tiina M. Sävilammi TM, † Gonzalo Ossa, ‡§ Anna S Blomberg, † Anti Vasemägi, ** Veronica Yung, †† David L.J. Vendrami, ‡‡ Joseph S Johnson §§

* Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland.
† Department of Biology, University of Turku, Turku, Finland.
‡ ConserBat EIRL, San Fabian, Chile
§ Programa para la Conservación de los Murciélagos de Chile, Santiago, Chile
** Department of Aquatic Resources, Swedish University of Agricultural Sciences, Uppsala, Sweden
†† Sección Rabia, Subdepartamento de Enfermedades Virales, Instituto de Salud Pública, Santiago, Chile
‡‡ Department of Animal Behavior, University of Bielefeld, Bielefeld, Germany
§§ Department of Biological Sciences, Ohio University, Athens, Ohio

Corresponding author: Thomas M. Lilley

ORCiD:

Thomas M. Lilley: 0000-0001-5864-4958

Tiina M. Sävilammi: 0000-0001-9836-3843

Anna S. Blomberg: 0000-0002-6754-4948

David L.J. Vendrami: 0000-0001-9409-4084

Joseph S. Johnson: 0000-0003-2555-8142

Data availability

Running title: Population structure of *Myotis chiloensis*

Key words: Population genetics, population connectivity, population structure, chiroptera,

disease spread.

[1] Finnish Museum of Natural History, P. Rautatiekatu 13, University of Helsinki, PL 17, 00014

Helsinki, Finland

Phone: +358408337783

Email: thomas.lilley@helsinki.fi

ABSTRACT

Despite its peculiar distribution, the biology of the southernmost bat species in the world,
the Chilean myotis (*Myotis chiloensis*), has garnered little attention so far. The species has a
north-south distribution of c. 2800 km, mostly on the eastern side of the Andes mountain
range. Use of extended torpor occurs in the southernmost portion of the range, putting the
species at risk of bat white-nose syndrome (WNS), a fungal disease responsible for massive
population declines in North American bats. Here, we examined how geographic distance
and topology would be reflected in the population structure of *M. chiloensis* along the
majority of its range using a double digestion RAD-tag method. We sampled 66 individuals
across the species range and discovered pronounced isolation-by-distance. Furthermore,
and surprisingly, we found higher degrees of heterozygosity in the southernmost
populations compared to the north. A coalescence analysis revealed that our populations
may still not have reached secondary contact after the Last Glacial Maximum. As for the
potential spread of pathogens, such as the fungus causing WNS, connectivity among
populations was noticeably low, especially between the southern hibernatory populations in
the Magallanes and Tierra del Fuego, and more northerly populations. This suggests the
probability of geographic spread of the disease from the north through bat-to-bat contact to
susceptible populations is low. The study presents a rare case of defined population
structure in a bat species and warrants further research on the underlying factors
contributing to this.

Key words: population structure, Myotis, Chiroptera, opportunistic pathogen, disease
spread

          INTRODUCTION

Transmission of infectious diseases has garnered attention as one of the greatest risks to

human, agriculture and wildlife health over the last decade (Cangelosi *et al.* 2004; Semenza

and Menne 2009). Previous research demonstrates that the emergence of previously

unknown diseases often results from a change in the ecology of the host, pathogen, and/or

their environment (Scholthof 2007). An example of this is white-nose syndrome (hereafter

WNS), an epizootic disease that emerged in North America in 2006 (Blehert *et al.* 2009). The

disease is caused by the fungus, *Pseudogymnoascus destructans*, which infects insectivorous

bats during the hibernation period at latitudes where prey are not widely available during

winter (Lorch *et al.* 2011). Populations of highly susceptible species, especially from the

genus *Myotis*, have declined by >90% in areas affected by WNS (Frick *et al.* 2010). The

opportunistic pathogen can utilize alternative carbon sources (Raudabaugh and Miller 2013)

and can persist in the cold, humid environment within hibernacula in the absence of bat

hosts (Lorch *et al.* 2013; Hoyt *et al.* 2014).

P. destructans is native to Eurasia, where it has a large geographic range, with transmission

to North America likely facilitated by humans (Warnecke *et al.* 2012; Leopardi *et al.* 2015).

In North America, bats suffering from WNS were first detected in the state of New York

during the winter of 2006–2007 (Frick *et al.* 2010). The fungus has since spread across North

America, with records of prevalence in 33 U.S. states and 7 Canadian provinces. So far, *P.*

destructans has been detected on 17 species of bats, with more species likely to follow.

While human-assisted transmission of *P. destructans* likely has contributed to this spread,

the ecology and behaviour of cave-hibernating bats in North America also makes them

efficient vectors over large geographic areas (Wilder *et al.* 2015). Because WNS affects bats

during extended bouts of torpor, at low temperatures where it is able to grow and infect the

host, there has been speculation over how far into the southern North America the disease

will spread (Verant *et al.* 2012; Meierhofer *et al.* 2019). Although bats inhabiting lower

latitudes may suffer less from WNS, *P. destructans* conidia may be able to survive on the

body of bats for several months, even at temperatures up to 37 °C (Campbell *et al.* 2019).

This could facilitate the movement of WNS across Mesoamerica and the tropics, to arrive to

high southern latitudes where bats may be susceptible (Holz *et al.* 2019; Turbill and
Welbergen 2019).

Of species known to harbour the WNS fungus, *Tadarida brasiliensis* is of particular interest.
As a long-range migratory species, with movements spanning thousands of kilometres
(Cockrum 1969; Glass 1982), *T. brasiliensis* may be an important vector for spreading *P.*
destructans into the southern hemisphere (Ommundsen *et al.* 2017; McCracken *et al.* 2018).
Ecological niche models predict suitable habitat for the proliferation of *P. destructans* in
South America, highlighting the need to understand vectors such as *T. brasiliensis* as well as
human transmission (Escobar *et al.* 2014). However, once *P. destructans* arrives in South
America, its spread will not necessarily resemble that seen in North America, as it is likely to
be influenced by differing geology and species ecology.

The Chilean myotis (*Myotis chiloensis* [Waterhouse, 1840]) is the most Southerly distributed
species of bat in the world, together with the southern big-eared brown bat (*Histiotus*
magellanicus, Koopman 1967; Gardner 2007). *Myotis chiloensis* has a vast north-south
distribution that includes forested areas on both sides of the Andes from the northern shore
of Navarino Island to the southern border of the Atacama desert in Chile (Ossa and
Rodriguez-San Pedro 2015). The range of *M. chiloensis* overlaps with the distribution of *T.*
brasiliensis from the north, where *M. chiloensis* is not believed to hibernate, to 45°S
latitude, where *M. chiloensis* possibly hibernate and may therefore be susceptible to WNS
(Bozinovic *et al.* 1985). However, there is no information available on the population
structure of *M. chiloensis*, precluding an understanding of how *P. destructans* could be
transported from the northern edge of its range to more southern, and vulnerable,
populations. The connectedness of individuals across the range will determine the speed
and intensity of potential spread. Population structuring in bats is often relatively low
because of their efficient mode of dispersal, flight (Laine *et al.* 2013). An ability to disperse
more efficiently results in decreased population differentiation (Bohonak 1999) to the
extent that some bat species are panmictic across their range (Burland and Wilmer 2001;
Laine *et al.* 2013). Such high dispersal would likely result in rapid spread of *P. destructans*.
However, bats in the genus *Myotis* show instances of pronounced population structure that
may hinder spread of the fungus. For instance, the Gibraltar Strait, which separates the

Iberian Peninsula from the Maghreb in Morocco by a minimum gap of 14 km of the open

sea, represents a barrier for gene flow for *M. myotis* (Castella *et al.* 2000). Chile is littered

with such potential barriers to gene flow, such as the Atacama Desert, glaciers, ice fields,

the Andes Mountains, and the Magellan Strait, which in turn can hinder the potential spread

of *P. destructans*. Furthermore, populations may still be affected by the Last Glacial

Maximum, which covered a large part of Patagonia under ice until c. 10000 years bp (Sérsic

et al. 2011; Mansilla *et al.* 2018).

With tourism in southern Chile expected to increase (e.g. . http://www.conaf.cl/parques-

nacionales/visitanos/estadisticas-de-visitacion/), and migratory species such as *T.*

brasiliensis capable of carrying spores across large distances, there is a serious need to

better understand the population structure of Patagonian bat species before WNS spreads

to the region. The lack of knowledge on the extent of migration and mixing, and life history

traits in general, means that research in this area is now urgent and essential (Ossa and

Rodriguez-San Pedro 2015; Ossa 2016; Ossa *et al.* 2019). Studying population ecology

through molecular genetic methods allows for the identification of more accurate

population boundaries, which is important when assessing conservation in response to

threats of disease and dramatic declines in population size (Moritz 1994). This study will

therefore aim to describe population structure and isolation-by-distance in *M. chiloensis*

across the range of the species. By studying *M. chiloensis* along 2400 km of latitudinal

gradient using genome-wide SNP markers, we aimed to test if geography and the Last

Glacial Maximum influence genetic isolation patterns.

143   MATERIALS AND METHODS

Sample collection and DNA extraction

To describe population genetic structure in *M. chiloensis*, we obtained wing tissue samples

of 66 bats from two sources. A portion were obtained from live bats captured in the field

during November and December 2017 (i.e. austral spring) from two localities: Chicauma,

Metropolitana region (33 °S 70 °W) and Karukinka Reserve, Tierra del Fuego (64 °S 78 °W)

respectively (Capture permit: 4924_2017, Figure 1 A, Table S1). We used disposable biopsy

punches (5 mmm, MLT3335, Miltex Instrument Co, Plainsboro, New Jersey) to collect tissue

samples from the plagiopatagium of captured, live bats. The sampled bats were released at

the capture site. Additional samples were obtained from dead bats submitted to the Public

Health Institute of Chile for rabies testing (Figure 1 A, Table S1). Submitted bats included

latitude and longitude locations of origin. Tissue samples from the bats submitted for rabies

testing were obtained from the plagiopatium using sterile scalpels. To determine if *P.*

destructans had already spread to Chile, we swabbed the nose and wings of all bats in the

field and at the Public Health Institute of Chile bat with a sterile polyester swab (Puritan 25-

806 1PD, Guildford ME, USA) and stored at -20 °C until analysis.

We divided samples into four groups according to their geographic origin, with sub-locations

within each group to assist in further analyses. Sampling locations are presented in Figure 1

A and details of populations and samples are provided in Table 1 and Table S1. Tissue

samples were stored in 1.5 ml tubes with 95% EtOH and stored at -20 °C until further

analysis. Fungal spore samples were stored in 1.5 ml tubes and stored at -20 °C until further

analysis. We extracted DNA from tissue samples using QIAmp DNA Mini Kits (Qiagen, Hilden

Germany) and stored DNA at -80 °C. DNA from fungal swabs was extracted using Qlamp

DNA Micro Kits (Qiagen, Hilden Germany).

The amount of DNA in the final solution of each sample was tested and quantified using the

Thermo Scientific Nanodrop spectrophotometer, giving a result for the amount of DNA in

ng/ µL. Samples were frozen between DNA extraction and analysis.

Identification and quantification of *P. destructans*

Quantification of *P.destructans* load by qPCR was completed as described previously in

Johnson et al. (2015) with the exception of using 1 µl sample in the reaction, Roche Fast

Start Essential DNA Probe Master, and a Roche Lightcycler 480 instead of a BioRad iCycler.

RAD sequencing

We sequenced 66 individuals in total. Duplicate samples from 30 individuals were

additionally sequenced to estimate repeatability and error rate of the called genotypes.

DNA was prepared for genotyping-by-sequencing using a double digestion RAD-tag method

as described in Elshire et al. (2011). PstI-BamHI-digested libraries were prepared by the

Center of Evolutionary Applications (University of Turku; see Lemopoulos et al. 2017 and
references therein for further details) and sequenced in an Illumina HiSeq2500 run (100 bp
single end reads and pooling 96 samples on each lane) at Finnish Functional Genomics
Centre (Turku Bioscience).

Yield comparison samples

As the amount of DNA available may often be very limited in experiments where preserved
samples from the rabies laboratory are utilized, we wanted to estimate the effect of initial
DNA concentration to the resulting read coverage. We compared the total read coverages
of the replicate samples with Pearson's correlation.

The resulting fastq reads, separated by barcode for each sample, were quality-controlled
and low-quality bases trimmed with condetri v. 2.3 with parameter minlen=30 (min. length
of a trimmed read) followed by adapter-trim with cutadapt for Illumina universal adapters
from the end of the reads (identified with FastQC quality control in some of the samples).
Then, reads were mapped against *M. lucifugus* genome (mluc_2.0_assembly_supers.fasta)
using bwa mem with parameters -B 3 -O 5 -k 15. After mapping back with BWA mem, we
used SAMtools v. 1.4 and the associated bcftools for calling genotypes for SNPs and for
filtering SNPs based on minimum of 40% of the samples genotyped, at least eight
alternative alleles detected and SNP quality >=20 (bcftools filter). We further filtered the
SNPs based on exactly two alleles detected and excluded SNPs with particularly low (≤5) or
unusually high (≥125) mean coverage based on visual inspection of the mean coverage
distribution.

Population genetic analysis

For the final SNP dataset, which only included unduplicated samples, we performed
principal component analysis implemented in prcomp function of R stats package, and
calculated Euclidean genetic coordinates for each individual from PC1 and PC2. The four
pre-determined ancestral populations based on the sampling regions of each individual
were confirmed by hierarchical clustering of the Euclidean distanced calculated from PC1
and PC2. Individuals clustering to a neighbouring population were assumed to have
dispersed from their natal populations and were reassigned for the later analysis. We

further investigated the relative contributions of the ancestral populations inhabiting
regions 1-4 in the present-day nucleotide using ADMIXTURE (Zhou et al. 2009) analysis for
the A-samples, run with four expected populations (parameter K=4) based on prior
knowledge of the population structure, and with quasi-Newton convergence acceleration
method. We were particularly interested in identifying possible hybrid individuals.

After assigning each individual to the final regions, we calculated Nei's pairwise F_{ST} using
HierFSTat v. 0.04-22 pairwise. FST function for A-samples for SNP's with no missing values.
To assess the significance of genetic differentiation, we compared the actual obtained F_{ST}
estimates to the null distributions of F_{ST} values under panmixia, obtained from a hundred
random permutations of alleles. Latitude and longitude coordinates of the sampling
locations were used to calculate pairwise geographic distances between individuals in
kilometres using Haversine method assuming a spherical earth, implemented in function
distm in the R package geodist v. 1.5-10.

We estimated Isolation-by-distance with two methods: A Mantel test with complete
permutations and a linear model *geographic distance ~ genetic distance*. We used pairwise
F_{ST} as a measure for the genetic distances and mean of between-individual distances as
geographic coordinates for populations inhabiting each of the four regions. To study the
population structure in more detail, we repeated the Isolation-by-distance analysis for the
genetic vs. geographic distances from the most extreme individual (sample ID 700) using a
Mantel test with all possible permutations, and a linear model to identify individuals with
unusually high or low genetic differentiation. We then studied the systematic differentiation
of the individuals sampled in different regions by assessing the differences of the residual
distributions of each of the four populations from zero using t-tests and Bonferroni-
correction of the *P*-values.

We also calculated mean observed heterozygosity within the variable loci in Hardy-
Weinberg equilibrium (*FDR*>=0.05) in each of the four study regions and compared the
observed values to the mean expected heterozygosities using inbreeding coefficient *F,*
calculated as the difference between expected and observed heterozygosity divided by
expected heterozygosity (Serre 2006). The 95% confidence intervals for the heterozygosity
estimates were found by randomly sampling the variable loci for 1,000 times and extracting

the distributions of bootstrap means. The deviations of F statistics from zero were detected

using single-sample Wilcoxon tests. The significances of the regional differences in the

observed and expected heterozygosity distributions, and in the F statistics, were tested both

by inspecting the overlaps in the bootstrapped confidence intervals and using analysis of

variance and Tukey's post hoc tests. The significances of within-region differences between

the expected and observed heterozygosity were tested both by comparing the

bootstrapped confidence intervals and by pairwise t-tests and Bonferroni-corrected p-

values. Finally, the fraction of SNPs unique to any one region, and the number of SNPs

shared between all regions, were calculated from the observations of variable and non-

variable loci within regions.

Demographic modeling

To formally test whether any of the studied populations experienced secondary contact

following glaciation, we implemented a demographic analysis using the software

fastsimcoal2 (Excoffier *et al.* 2013) in combination with the folded site frequency spectra

(SFS) calculated from our data using easySFS.py utility (available from

https://github.com/isaacovercast/easySFS). Specifically, we evaluated support for two

alternative models (Supplementary Figure S6). The first model, representing our null

hypothesis of no secondary contact among populations, specified four distinct lineages

(region 1, region 2, region 3 and region 4) corresponding to the populations inhabiting the

four geographic regions sampled in the present study. These diverged from each other at

the time points T1, T2 and T3 as presented in Figure S6a, and exchanged no migrants.

Region 1 was used as the lineage from which the other three populations emerged, as this is

thought to be the population which is closest to the ancestral *Myotis chiloensis* population.

The second model, representing our alternative hypothesis of secondary contact among

populations, was identical with the exception that symmetric migration was present

between population pairs region 1- region 2 and region 3- region 4, and asymmetric

migration was implemented from region 2 to region 3 (Supplementary Figure S6b). In

addition to identify the model that was best supported by our data, we also estimated

divergence times (T1, T2 and T3) and effective population sizes (region 1, region 2, region 3

and region 4) for the four modeled lineages, as well as migration rates (Mig12, Mig21,
Mig34, Mig43 and Mig32).

We performed 50 independent *fastsimcoal2* runs for each model, with 100,000 simulations
and 40 cycles of the likelihood maximization algorithm. We then calculated Akaike's
Information Criteria (AIC) from the *fastsimcoal2* runs which yielded the highest maximum
likelihood for each model and used these values for model comparison. Finally, we
extracted parameter estimates from the best run of the most supported model and
calculated 95% confidence intervals based on 100 parametric bootstrap replicates, as
described in Excoffier et al. (2013).

Data availability

The RAD sequencing reads were deposited at NCBI SRA under BioProject ID PRJNA596389.

278   RESULTS

Filtering

Ninety-one of the 96 samples (66 individuals and 30 duplicates) had reads matched with a
barcode. Of the obtained genotypes, 54846 were biallelic and used in the later analysis,
while we excluded 88882 non-variable (homozygous to alternative) variants and 1708
variants with more than 2 alleles. After filtering, the mean SNP coverage ranged from 0.5 to
257.0 (Supplementary Figure S1). We excluded tags with < 5 or > 125 mean coverage,
leaving 47079 tags. The average rate of missing SNPs among the final unique samples was
6.2%, ranging from 0 to 24%.

Replicate samples

The correlation between input DNA concentration and the resulting mean per-sample read
coverage was only moderate (cor = 0.3394, t_{61} = 2.8178, *P* = 0.0065, Supplementary Figure
S2 A, Supplementary Figure 3 A-B). In contrast, we found strong and negative association
between mean read coverage after sequence assembly and the number of missing genotype
calls (cor = -0.9763, t_{61} = -35.223, *P* < 2.2e-16, Figure S2 B). Furthermore, read coverages
were very similar between the technical replicate samples (cor = 0.8736, t_{26} = 9.1542, *P* =

1.292e-09, Supplementary figure S2 C), allowing us to omit "B" samples (replicated). The
removal of the biological replicates was conducted to minimise the possible SNP calling bias
induced by some samples having approximately twice the amount of sequence data
compared to the others, if replicate samples had been combined. Identical genotype calls
ranged from 85.6% to 96.9% with an average of 94.2% identical genotype calls
(Supplementary figure S3 C), depending heavily on read coverage (cor = 0.9275, t_{26} = 12.641,
P = 1.312e-12 and cor = 0.8484, t_{26} = 8.1716, P = 1.186e-08 in "A" and "B" samples,
respectively; Supplementary figure 2 E-F). Although the correlation between the initial DNA
concentration and identical genotype calls between biological replicates was significant (cor
= 0.5990, t_{26} = 3.814, P = 0.0007579; Supplementary figure S2 D), this seemed to mainly be
due to two outlier observations with both very low concentration and repeatability.

Identification and quantification of *P. destructans*

Besides our control samples, no samples showed signs of amplification a portion of the
multicopy intergenic spacer region of the rRNA gene complex of *P. destructans* by 38 cycles,
which is generally considered as a cut-off for the presence of the pathogen DNA in the
samples (Muller *et al.* 2013). Therefore, we can conclude that the *M. chiloensis* individuals
sampled in this study did not carry *P. destructans*.

Population genetic analysis

Principal component analysis on 66 individuals and 5538 SNPs without any missing values
from non-duplicated samples (label including A, Figure 1 B) revealed a clear structuring of
individuals according to the sampling location indicative of strong population structure.
Based on hierarchical clustering of the two most important principal components
(Supplementary Figure 4), we confirmed the four pre-determined populations based on the
natural hierarchical structuring of the data. Based on the clustering, the sub-population
assignment of one individual, sample number 679, changed from Biobio to Maule (which
was the most common region assignment among the three nearest neighbors for that
individual). The estimation of ancestry of each sampled individual by examining the relative
contributions of ancestral populations inhabiting regions 1-4 in the present-day revealed

particularly pure ancestral lines in the northern and southern parts of the range, with

hybridization occurring in the central part of the range (Figure 2.).

Pairwise F_{ST}-value estimates between the populations sampled from the four geographical

locations ranged from 0.04 (between regions 2 and 3) to 0.113 (between the most distant

regions 1 and 4, Table 2). All estimated F_{ST} values were found significantly larger ($P<0.01$)

than the permuted F_{ST} distributions, with the 95 % confidence intervals of the null

distributions between 0 and 0.0205. Both the Mantel test approach and linear modelling

between genetic distances (Euclidean distances calculated from PC1 and PC2) and

geographical distances (latitude/longitude coordinates converted to distances in kilometres)

strongly suggested that we reject the null hypothesis of geographic and genetic distances

being unrelated. For the between-population comparisons with F_{ST}, we calculated Mantel

statistic r = 0.9497 (P = 0.041667) and R-squared estimate of 0.8773 (DF = 1,4, t = 6.063, P =

0.00374) (Supplementary figure 5. Also, for the between-individual distances, the observed

Mantel statistic r = 0.944 (P = 0.001), and R-squared estimate 0.943 (DF = 1,61; t = 31.994; P

< 2e-16) from the linear modelling suggesting that genetic and geographic distances are

strongly positively associated (Figure 3). More detailed exploration of the between-

individual linear model revealed, that the distribution of the residuals in regions 1 (DF = 20, t

= -3.0679, Bonferroni P = 0.024284) and 4 (DF = 11, t = -10.523, Bonferroni P = 1.7724e-06)

were marginally smaller than 0, while the residuals of individuals in regions 2 (DF = 15; t =

3.5896; adj. Bonferroni P = 0.002682) and 3 (DF = 13, t = 5.9325, Bonferroni P = 0.0001986)

were significantly larger than 0 (Figure 3).

Of the total of 47079 SNPs, 43903 were found to be in Hardy-Weinberg equilibrium.

Analyses of variance revealed statistically significant differences between populations in

observed heterozygosities [$F(3,106450)$=760.6, $P<2e-16$], in expected heterozygosities

[$F(3,106450)$=871.4, $P<2e-16$], and in the F statistics [F(3,106450)=55.29, P<2e-16]. Both

based on confidence intervals and pairwise comparisons, observed and expected

heterozygosity distributions were consistently higher (non-overlapping 95% confidence

intervals and adjusted $P<0.05$) in the southern regions than in north, except for the two

northernmost regions (region 1 vs. region 2), where a statistically significant difference was

not observed (Table 1, Table S2). Similarly, observed heterozygosities were consistently

lower than those estimated from allele frequencies (non-overlapping 95% confidence

intervals and adjusted *P*<0.05), which may be caused by within-region population structure

(Table 1, Table S2). Finally, pairwise comparisons of the inbreeding coefficient distributions

indicated that the excess of homozygotes was greater in the north than in the south when

compared to neutral expectation based on allele frequencies. This was supported with

statistically significant differences (non-overlapping confidence intervals and adjusted

P<0.05) observed in all comparisons except in region pairs 2 and 3; and 4 and 5 (Table 1,

Table S2). This indicated that the northern populations are inbreeding more than the

southern populations (Table 1, Table S2).

A large proportion of SNPs, 25.3%, were polymorphic in all four regions. The fraction of

SNPs unique only to one region decreased from going north to south: while 8.6% of the

SNPs were unique to region 1, only 2.5% unique SNPs were found in region 4.

We did not find support for secondary contact using demographic modelling with

Fastsimcoal2. Instead, the null model that included no migration among populations (Figure

S6a) received the highest AIC support (Table S3). Parameter priors and estimates, together

with their 95% confidence intervals, from the best model are reported in Table S4.

DISCUSSION

Our results present the first assessment of population structure in the widely distributed bat

species, *M. chiloensis*, using individual-based approach with genome-wide markers. We

found that geographic distance within the range of the species are reflected in its

population structure. Although we found a clear and robust population structure among

sampling sites, population structure is correlated with geographical distance, even though

populations are separated by ice fields, mountain ranges and stretches of open water. This

has implications for the protection of populations that may be susceptible to WNS. Our

results also show highest genetic variability in the species is at the southern extent of its

range.

Strong population structure is rarely seen in bats, even across large geographical scales in

genera such as *Myotis*, with shorter dispersal distances (Castella *et al.* 2000; Atterby *et al.*

2010; Laine *et al.* 2013). In fact, geographical distance often correlates significantly with

genetic distance in bats. This is partially due to autumn migration and swarming behaviour

in *Myotis* species, which brings together bats from broad geographic areas to breed and

promote recombination (Burns *et al.* 2014; Burns and Broders 2015). Furthermore, powered

flight allows effective dispersal, which is often male biased (Arnold 2007; Angell *et al.* 2013).

This is reflected in low fixation indices in widespread bat species, such as *M. daubentonii*,

where individuals separated by thousands of kilometres in Europe show low fixation indexes

(Laine *et al.* 2013). While fixation indices cannot be compared directly across species,

especially when different methodological approaches are used, they do give an indication of

the connectivity of populations.

Although geographic and genetic distances were found to have a strong positive correlation

in our study, we can concur that the southernmost population shows a higher degree of

isolation compared to the geographic distance to its closest comparative population to the

north. An F_{ST} of 0.075 between regions 3 and 4 using whole-genome data is already higher

than F_{ST} values recorded for *M. daubentonii* across Europe using microsatellites (Laine *et al.*

2013), and in our data, the geographic distance is only c. 1000 km. Our data, with tens of

thousands of SNP's also allowed a more precise estimate of fixation compared to a handful

of microsatellites. However, due to a limited number of individuals, it did not allow us to

examine dispersal as a function of sex, which in bats is often a male driven function (Arnold

2007; Laine *et al.* 2013; Angell *et al.* 2013). Our sampling may also have missed some

connecting populations in between regions 3 and 4, which could for instance be located on

the eastern slopes of the Andes. However, our test for relative contributions of ancestral

populations reveals bats in region 4, in the Magallanes and Tierra del Fuego, have no mixing

of ancestral populations with the other regions.

In the northern hemisphere, approximately 18000 years ago, at the end of the Late

Pleistocene, the ice sheets began to recede as the global climate became warmer. The biota

migrated northwards following their optimal environments (Huntley and Webb 1989). This

expansion of refugial populations has been associated with genetic variation decreasing

south-to-north in some species: a trend attributed to a series of bottlenecks when the biota

spread from the leading edge of the refugial population, leading to a loss of alleles and

decreasing genetic diversity (Hewitt 1996, 1999). Contrary to what one could expect based

on these latitudinal shifts in diversity in the northern latitude, genetic diversity in *M.*

chiloensis appears to increase with increasing latitude, from north to south. We presumed

this counterintuitive pattern of heterozygosity within the species may be related to the

glaciation history of South America, where populations isolated by glacial events could have

been able to hybridize. For instance, in some terrestrial European and Scandinavian

vertebrates, the intraspecific genealogical lineages, which formed in separate refugia, were

found to have come to secondary contact in the Fennoscandian area (Tegelström 1987;

Jaarola and Tegelström 1995; Nesbø *et al.* 1999; Knopp and Merilä 2009).

Myotis chiloensis is described as a vicariant species with respect to other closely related

Myotis species (*M. albescens, M. nigricans, M. levis*) from South America, Ruedi et al (2013)

estimated this isolation event at 5.5 My in late Miocene. This same time period saw the

beginning of a number of glaciations events in Patagonia with variable intensity and

duration (Rabassa *et al.* 2011). Glacial episodes isolated the Patagonian forest from around

middle Miocene well into the late Quaternary, the Last Glacial Maximum (Rabassa *et al.*

2011, Supplemental Figure S7). During the glaciations, the forests on the Pacific coast were

most like completely suppressed, possibly with isolated small refugia. On the Atlantic side,

the forest was fragmented from 36°S southwards (Sérsic *et al.* 2011). Finally, in Tierra del

Fuego the forest was probably displaced towards the current submarine shelf (Ponce *et al.*

2011). As the ice retreated refugial populations may have come into secondary contact in

the southern part of current range, which could explain the high heterozygosity as well as

the small fraction of unique SNP's of these populations. However, our coalescence analysis

rejected the secondary contact model, favouring the null model suggesting our study

populations are still largely separated after the Last Glacial Maximum. Indeed, the F_{ST} -

values are high, suggesting isolation of the populations. By contrast, our analysis for the

relative contributions of ancestral populations suggests mixing of populations. One potential

explanation for this apparent discrepancy is that we derived Site Frequency Spectrum from

a rather small number of individuals, which may carry a signal of migration that is not strong

enough to allow a model that includes secondary contact to be favoured over a simpler

model without migration.

The spread of *P. destructans* via one host to another was very rapid in North America
(Blehert *et al.* 2009; Frick *et al.* 2010). This was facilitated in part by the ecology of North
American cave-hibernating bats and the availability of suitable environment for the fungus
to propagate: limestone caves found throughout the Appalachian region in eastern North
America (Lorch *et al.* 2013). Furthermore, as a consequence of down-regulation of
metabolism during extended torpor bouts, attempted immune responses fall short, and
may even contribute to mortality in hosts infected with *P. destructans* (Field *et al.* 2015;
Lilley *et al.* 2017, 2019). In addition to these, the massive population declines associated
with WNS in affected species (Turner *et al.* 2011) were magnified by the panmictic
population structure across eastern North America in the most affected species, *M.*
lucifugus (Miller-Butterworth *et al.* 2014; Vonhof *et al.* 2015).

Our results for *M. chiloensis* from austral South America suggest the southernmost
population in region 4, may be less likely to be infected via their continental conspecifics,
because of reduced contact between the populations. Our results indicated no mixing of
ancestry in the southernmost individuals in our study, suggesting the *M. chiloensis* on Tierra
del Fuego are isolated from their mainland counterparts. To our knowledge, this is also the
only population to use extended torpor, a prerequisite for the propagation of *P. destructans*
and the onset of WNS (Ossa *et al.* Submitted for publication). Tierra del Fuego, the southern
tip of Patagonia and the continent of South America, experiences extended low winter
temperatures comparable to areas in North America where WNS is manifested. Our genetic
analysis for the presence of *P. destructans* on the sampled bats suggests the fungal
pathogen does not exist within the distribution range of our focal species at present. Even if
the fungus were to enter the region, variability in host behavior and environmental
characteristics may be the primary factors protecting hosts from the pathology related to
WNS (Zukal *et al.* 2014, 2016). Most strikingly large cave hibernacula with suitable, stable
environmental conditions favouring the environmental persistence of the pathogen in the
absence of the hosts, are very scarce and separated by hundreds of kilometers in most of
the southern range of *M. chiloensis*.

The observed distribution of *M. chiloensis* is vast, covering a range of forested habitats from
arid Sclerophyllous to sub-Antartic (Ossa and Rodriguez-San Pedro 2015). In this respect,

taking into consideration our results on clear population segregation begs to propose the
question on the species status of *M. chiloensis* as a whole. Indeed, *M. chiloensis* also
appears to vary phenotypically along its distribution range (Mann 1978; Ossa 2016). It has
been proposed that the species was composed of three sub-species: *M. ch. atacamensis*
which is presently known as *M. atacamensis; M. ch. arescens* from 29°S to 39°S; and *M. ch.*
chiloensis from 39°S to 53°S (Mann 1978). That classification was due according to their coat
colour changes in relation to exposure to solar radiation and ambient temperature, which is
correlated to latitude (Budyko 1969) and levels of precipitation in their habitat. They vary
from a lighter pelage to a dark brown colour on a gradient from the northern part of their
range to the south (Galaz *et al.* 2006). This promotes the theory, isolation by adaptation, as
a driver of population genetic structure. The genetic adaptations of an individual to their
local environment separates populations and leads to a reduced gene flow (Orsini *et al.*
2013). However, the isolation by adaptation theory negates the fact that there is a
possibility of a barrier so that gene flow is inhibited by climate or adaptations to the local
environment. It is possible to therefore state that isolation-by-dispersal limitation, and
moreover isolation-by- distance, are the more probable causes of the observed results in
the present study. Indeed, the reluctance of the species to cross barriers, for instance the
Andes, can clearly be seen by examining population 3, where six individuals sampled from
Puerto Aysén (437, 442, 443, 167, 256, 260) are visibly isolated on the PCA plot form
individuals on the other side of the Andes, under 150 km away. This also depicts the fine
scale resolution our individual-based SNP-based approach allows. However, further studies
should focus deeper on the taxon status of different population of the species currently
recognized as *M. chiloensis.*

The results highlight the importance to assess the population structure which may limit the
spread of white-nose syndrome disease. Whether *P. destructans* or another epizootic in the
future could spread depends largely on the population structure and connectedness of
hosts (Lilley *et al.* 2018).

497   ACKNOWLEDGEMENTS

We thank the Rufford Foundation (Rufford Small Grant 10502-1 and 23042-2), H2020 Marie
Skłodowska-Curie Actions (706196), and Ohio University Research Council for funding the

work. We thank Servicio Agricola y ganadero Diproren for the capture permits in Tierra del
Fuego (Res Ex: 1253/2016 and 4924/2017), Juan Carlos Aravena from the Instituto de la
Patagonia for his help, the personnel from WCS Chile for allowing us to conduct research at
Karukinka Natural Reserve as well as for their help with field work. We thank Michelle
Lineros and Tania Gatica from the National Health Institute for their help to obtain the
tissue samples. We thank Austin Waag for assistance with field work.

REFERENCES

Angell, R. L., R. K. Butlin, and J. D. Altringham, 2013 Sexual segregation and flexible mating

patterns in temperate bats. Plos One 8:.

Arnold, B. D., 2007 Population structure and sex-biased dispersal in the forest dwelling

vespertilionid bat, *Myotis septentrionalis*. Am. Midl. Nat. 157: 374–384.

Atterby, H., J. Aegerter, G. Smith, C. Conyers, T. Allnutt *et al.*, 2010 Population genetic

structure of the Daubenton's bat (*Myotis daubentonii*) in western Europe and the

associated occurrence of rabies. Eur. J. Wildl. Res. 56: 67–81.

Blehert, D. S., A. C. Hicks, M. Behr, C. U. Meteyer, B. M. Berlowski-Zier *et al.*, 2009 Bat white-

nose syndrome: an emerging fungal pathogen? Science 323: 227–227.

Bohonak, A. J., 1999 Dispersal, gene flow, and population structure. Q. Rev. Biol. 74: 21–45.

Bozinovic, F., L. Contreras, M. Rosenmann, and J. Torres-Mura, 1985 Bioenergetics of *Myotis*

chiloensis (Quiroptera: Vespertilionidae). Rev Chil Hist Nat 58: 39–45.

Budyko, M. I., 1969 The effect of solar radiation variations on the climate of the Earth. Tellus

21: 611–619.

Burland, T. M., and J. W. Wilmer, 2001 Seeing in the dark: molecular approaches to the

study of bat populations. Biol. Rev. 76: 389–409.

Burns, L. E., and H. G. Broders, 2015 Who swarms with whom? Group dynamics of *Myotis*

bats during autumn swarming. Behav. Ecol. 26: 866–876.

Burns, L. E., T. R. Frasier, and H. G. Broders, 2014 Genetic connectivity among swarming

sites in the wide ranging and recently declining little brown bat (*Myotis lucifugus*).

Ecol. Evol. 4: 4130–4149.

Campbell, L. J., D. P. Walsh, D. S. Blehert, and J. M. Lorch, 2019 Long-term survival of

Pseudogymnoascus destructans at elevated temperatures. J. Wildl. Dis.

Cangelosi, G., N. Freitag, and M. Riley-Buckley, 2004 From outside to inside: environmental

microorganisms as human pathogens. *American Society of Microbiology*.

Castella, V., M. Ruedi, L. Excoffier, C. Ibanez, R. Arlettaz *et al.*, 2000 Is the Gibraltar Strait a

barrier to gene flow for the bat *Myotis myotis* (Chiroptera: Vespertilionidae)? Mol.

Ecol. 9: 1761–1772.

Cockrum, E., 1969 Migration in the guano bat, *Tadarida brasiliensis*. Misc Publ Mus Nat Hist

Univ Kans. 51: 303–336.

Elshire, R. J., J. C. Glaubitz, Q. Sun, J. A. Poland, K. Kawamoto *et al.*, 2011 A robust, simple

genotyping-by-sequencing (GBS) approach for high diversity species (L. Orban, Ed.).

PLoS ONE 6: e19379.

Escobar, L. E., A. Lira-Noriega, G. Medina-Vogel, and A. T. Peterson, 2014 Potential for

spread of the white-nose fungus (*Pseudogymnoascus destructans*) in the Americas:

use of Maxent and NicheA to assure strict model transference. Geospatial Health

221–229.

Excoffier, L., I. Dupanloup, E. Huerta-Sánchez, V. C. Sousa, and M. Foll, 2013 Robust

Demographic Inference from Genomic and SNP Data. PLOS Genet. 9: e1003905.

Field, K. A., J. Johnson, T. Lilley, S. Reeder, E. Rogers *et al.*, 2015 The white-nose syndrome

transcriptome: activation of anti-fungal host responses in wing tissue of hibernating

bats. Plos Pathog. 11: e1005168.

Frick, W. F., J. F. Pollock, A. C. Hicks, K. E. Langwig, D. S. Reynolds *et al.*, 2010 An emerging

disease causes regional population collapse of a common North American bat

species. Science 329: 679–682.

Galaz, J., J. Yañez, A. Gantz, and D. Martínez, 2006 Orden Chiroptera. Pp. 67–89 in

Mamíferos de Chile (A. Muñoz-Pedreros and J.Yañez, eds.)., in *Mamíferos de Chile*,

Centro de Estudios Agrarios y Ambientales, Valdivia, Chile.

Gardner, A. (ed.), 2007 *Mammals of South America, Volume 1*. University of Chicago Press,

Chicago, Illinois.

Glass, B. P., 1982 Seasonal movements of Mexican freetail bats *Tadarida brasiliensis*

mexicana banded in the Great Plains. Southwest. Nat. 27: 127–133.

Hewitt, G. M., 1999 Post-glacial re-colonization of European biota. Biol. J. Linn. Soc. 68: 87–

112.

Hewitt, G. M., 1996 Some genetic consequences of ice ages, and their role in divergence and

speciation. Biol. J. Linn. Soc. 58: 247–276.

Holz, P., J. Hufschmid, W. Boardman, P. Cassey, S. Firestone *et al.*, 2019 Does the fungus

causing white-nose syndrome pose a significant risk to Australian bats? Wildl. Res.

46: 657–668.

Hoyt, J. R., K. E. Langwig, J. Okoniewski, W. F. Frick, W. B. Stone *et al.*, 2014 Long-term

persistence of *Pseudogymnoascus destructans*, the causative agent of white-nose

syndrome, in the absence of bats. EcoHealth.

Huntley, B., and T. Webb, 1989 Migration: species' response to climatic variations caused by

changes in the earth's orbit. J. Biogeogr. 5–19.

Jaarola, M., and H. Tegelström, 1995 Colonization history of north European field voles

(*Microtus agrestis*) revealed by mitochondrial DNA. Mol. Ecol. 4: 299–310.

Johnson, J. S., D. M. Reeder, T. M. Lilley, G. Á. Czirják, C. C. Voigt *et al.*, 2015 Antibodies to

Pseudogymnoascus destructans are not sufficient for protection against white-nose

syndrome. Ecol. Evol. 2203–2214.

Knopp, T., and J. Merilä, 2009 The postglacial recolonization of Northern Europe by *Rana*

arvalis as revealed by microsatellite and mitochondrial DNA analyses. Heredity 102:

174–181.

Koopman, K., 1967 The southernmost bats. J. Mammal. 48: 487–488.

Laine, V. N., T. M. Lilley, K. Norrdahl, and C. R. Primmer, 2013 Population genetics of

Daubenton's bat (*Myotis daubentonii*) in the Archipelago Sea, SW Finland. Ann. Zool.

Fenn. 50: 303–315.

Lemopoulos, A., S. Uusi-Heikkilä, A. Vasemägi, A. Huusko, H. Kokko *et al.*, 2017 Genome-
wide divergence patterns support fine-scaled genetic structuring associated with
migration tendency in brown trout. Can. J. Fish. Aquat. Sci. 75: 1680–1692.

Leopardi, S., D. Blake, and S. J. Puechmaille, 2015 White-nose syndrome fungus introduced
from Europe to North America. Curr. Biol. 25: R217–R219.

Lilley, T., J. Anttila, and L. Ruokolainen, 2018 Landscape structure and ecology influence the
spread of a bat fungal disease. Funct. Ecol. 32: 2483–2496.

Lilley, T. M., J. M. Prokkola, A. S. Blomberg, S. Paterson, J. S. Johnson *et al.*, 2019 Resistance
is futile: RNA-sequencing reveals differing responses to bat fungal pathogen in
Nearctic *Myotis lucifugus* and Palearctic *Myotis myotis*. Oecologia.

Lilley, T. M., J. M. Prokkola, J. S. Johnson, E. J. Rogers, S. Gronsky *et al.*, 2017 Immune
responses in hibernating little brown myotis (*Myotis lucifugus*) with white-nose
syndrome. Proc R Soc B 284: 20162232.

Lorch, J. M., C. U. Meteyer, M. J. Behr, J. G. Boyles, P. M. Cryan *et al.*, 2011 Experimental
infection of bats with *Geomyces destructans* causes white-nose syndrome. Nature
480: 376-U129.

Lorch, J. M., L. K. Muller, R. E. Russell, M. O'Connor, D. L. Lindner *et al.*, 2013 Distribution
and environmental persistence of the causative agent of white-nose syndrome,
Geomyces destructans, in bat hibernacula of the Eastern United States. Appl.
Environ. Microbiol. 79: 1293–1301.

Mann, G., 1978 Los pequeños mamiferos de Chile. Gayana Concepc. 40: 1–342.

Mansilla, C. A., R. D. McCulloch, and F. Morello, 2018 The vulnerability of the Nothofagus

forest-steppe ecotone to climate change: Palaeoecological evidence from Tierra del

Fuego (~53°S). Palaeogeogr. Palaeoclimatol. Palaeoecol. 508: 59–70.

McCracken, G. F., R. F. Bernard, M. Gamba-Rios, R. Wolfe, J. J. Krauel *et al.*, 2018 Rapid

range expansion of the Brazilian free-tailed bat in the southeastern United States,

2008–2016. J. Mammal. 99: 312–320.

Meierhofer, M. B., J. S. Johnson, S. J. Leivers, B. L. Pierce, J. E. Evans *et al.*, 2019 Winter

habitats of bats in Texas. PLOS ONE 14: e0220839.

Miller-Butterworth, C. M., M. J. Vonhof, J. Rosenstern, G. G. Turner, and A. L. Russell, 2014

Genetic structure of little brown bats (*Myotis lucifugus*) corresponds with spread of

white-nose syndrome among hibernacula. J. Hered. esu012.

Moritz, C., 1994 Defining 'Evolutionarily Significant Units' for conservation. Trends Ecol.

Evol. 9: 373–375.

Muller, L. K., J. M. Lorch, D. L. Lindner, M. O'Connor, A. Gargas *et al.*, 2013 Bat white-nose

syndrome: a real-time TaqMan polymerase chain reaction test targeting the

intergenic spacer region of *Geomyces destructans*. Mycologia 105: 253–259.

Nesbø, C. L., T. Fossheim, L. A. Vøllestad, and K. S. Jakobsen, 1999 Genetic divergence and

phylogeographic relationships among European perch (*Perca fluviatilis*) populations

reflect glacial refugia and postglacial colonization. Mol. Ecol. 8: 1387–1404.

Ommundsen, P., C. Lausen, and L. Matthias, 2017 First acoustic records of the Brazilian free-

tailed bat (*Tadarida brasiliensis*) in British Columbia. Northwest. Nat. 98: 132–136.

Orsini, L., J. Vanoverbeke, I. Swillen, J. Mergeay, and L. De Meester, 2013 Drivers of

population genetic differentiation in the wild: isolation by dispersal limitation,

isolation by adaptation and isolation by colonization. Mol. Ecol. 22: 5983–5999.

Ossa, G., 2016 Primer registro de la especie *Myotis chiloensis*, (Waterhouse, 1838)

(Chiroptera, Vespertilionidae) en el Parque Nacional Alberto de Agostini (Región de

Magallanes y Antártica Chilena). An. Inst. Patagon. 44: 85–88.

Ossa, G., J. S. Johnson, A. I. E. Puisto, V. Rinne, I. E. Sääksjärvi *et al.*, 2019 The Klingon

batbugs: Morphological adaptations in the primitive bat bugs, *Bucimex chilensis* and

Primicimex cavernis, including updated phylogeny of Cimicidae. Ecol. Evol. 9: 1736–

1749.

Ossa, G., T. M. Lilley, A. G. Waag, M. Meierhofer, and J. Johnson, Submitted for publication

Roosting ecology of the southernmost tree-dwelling bats, *Myotis chiloensis* and

Histiotus magellanicus, in southern Tierra del Fuego.

Ossa, G., and Rodriguez-San Pedro, 2015 *Myotis chiloensis* (Chiroptera: Vespertilionidae).

Mammal Species 47: 51–56.

Ponce, J. F., J. Rabassa, A. Coronato, and A. M. Borromei, 2011 Palaeogeographical evolution

of the Atlantic coast of Pampa and Patagonia from the last glacial maximum to the

Middle Holocene. Biol. J. Linn. Soc. 103: 363–379.

Rabassa, J., A. Coronato, and O. Martínez, 2011 Late Cenozoic glaciations in Patagonia and

Tierra del Fuego: an updated review. Biol. J. Linn. Soc. 103: 316–335.

Raudabaugh, D. B., and A. N. Miller, 2013 Nutritional capability of and substrate suitability

for *Pseudogymnoascus destructans*, the causal agent of wat white-nose syndrome.

PLoS ONE 8: e78300.

Ruedi, M., B. Stadelmann, Y. Gager, E. J. P. Douzery, C. M. Francis *et al.*, 2013 Molecular

phylogenetic reconstructions identify East Asia as the cradle for the evolution of the

cosmopolitan genus Myotis (Mammalia, Chiroptera). Mol. Phylogenet. Evol. 69: 437–

449.

Scholthof, K.-B. G., 2007 The disease triangle: pathogens, the environment and society. Nat.

Rev. Microbiol. 5: 152–156.

Semenza, J. C., and B. Menne, 2009 Climate change and infectious diseases in Europe.

Lancet Infect. Dis. 9: 365–375.

Serre, J., 2006 *Génétique des populations*. Dunod, Paris.

Sérsic, A. N., A. Cosacov, A. A. Cocucci, L. A. Johnson, R. Pozner *et al.*, 2011 Emerging

phylogeographical patterns of plants and terrestrial vertebrates from Patagonia.

Biol. J. Linn. Soc. 103: 475–494.

Tegelström, H., 1987 Transfer of mitochondrial DNA from the northern red-backed vole

(*Clethrionomys rutilus*) to the bank vole (*C. glareolus*). J. Mol. Evol. 24: 218–227.

Turbill, C., and J. A. Welbergen, 2019 Anticipating white-nose syndrome in the Southern

Hemisphere: Widespread conditions favourable to *Pseudogymnoascus destructans*

pose a serious risk to Australia's bat fauna. Austral Ecol. 45: 89–96.

Turner, G. G., D. M. Reeder, and J. T. H. Coleman, 2011 A five-year assessment of mortality

and geographic spread of white-nose syndrome in North American bats and a look to

the future. Bat Res. News 52: 13–27.

Verant, M. L., J. G. Boyles, W. Waldrep, G. Wibbelt, and D. S. Blehert, 2012 Temperature-

dependent growth of *Geomyces destructans*, the fungus that causes bat white-nose

syndrome. Plos One 7: e46280.

Vonhof, M. J., A. L. Russell, and C. M. Miller-Butterworth, 2015 Range-wide genetic analysis

of little brown bat (*Myotis lucifugus*) populations: estimating the risk of spread of

White-nose syndrome. PLoS ONE 10: e0128713.

Warnecke, L., J. M. Turner, T. K. Bollinger, J. M. Lorch, V. Misra *et al.*, 2012 Inoculation of

bats with European *Geomyces destructans* supports the novel pathogen hypothesis

for the origin of white-nose syndrome. Proc. Natl. Acad. Sci. U. S. A. 109: 6999–7003.

Wilder, A. P., T. H. Kunz, and M. D. Sorenson, 2015 Population genetic structure of a

common host predicts the spread of white-nose syndrome, an emerging infectious

disease in bats. Mol. Ecol. 24: 5495–5506.

Zukal, J., H. Bandouchova, T. Bartonicka, H. Berkova, V. Brack *et al.*, 2014 White-nose

syndrome fungus: a generalist pathogen of hibernating bats. Plos One 9: e97224.

Zukal, J., H. Bandouchova, J. Brichta, A. Cmokova, K. S. Jaron *et al.*, 2016 White-nose

syndrome without borders: *Pseudogymnoascus destructans* infection tolerated in

Europe and Palearctic Asia but not in North America. Sci. Rep. 6: 19829.

Figure captions:

Figure 1. Sampling locations and groupings for genetic sampling of *Myotis chiloensis* in Chile
(A). The two most important principal components calculated from allele frequencies
explain 15.8% of the total nucleotide variation (B).

Figure 2. Biogeographical ancestry (admixture) analysis based on nucleotide polymorphisms.
Each vertical bar represents an individual, ordered by latitude. Red, green, blue and purple
colours indicate the relative genetic contributions of ancestral populations inhabiting
regions 1-4, respectively.

Fig. 3. The correlation between geographic and genetic distances between individuals.
Geographic distances are measured in kilometres, and genetic distances as Euclidean
principal component distances from the reference individual (sample 700). Dashed lines
represent 95% confidence interval for the linear model. The violin plot highlights the
differences between model residuals for each study region. Residual distributions that differ
significantly from zero after Bonferroni correction for multiple testing are marked with * (P
< 0.05) and *** (P < 0.0001).

Numeric region	Number of individuals (A)	Number of duplicates (B)	male / female	Mean observed heterozygosity (95% confidence interval)	Mean expected heterozygosity (95% confidence interval)	Inbreeding coefficient F	Unique SNPs
1	20	6	16/4	0.2550 (0.2530-0.2570)	0.2793 (0.2777-0.2809)	0.0868 (0.0813 - 0.0925)	8.6%
2	19	6	9/10	0.2536 (0.2514-0.2557)	0.2765 (0.2749-0.2780)	0.08286 (0.0773- 0.0891)	4.4%
3	14	9	7/7	0.2872 (0.2848-0.2896)	0.2993 (0.29762-0.3009)	0.0404 (0.0343 − 0.0463)	3.6%
4	13	9	5/8	0.3248 (0.3221-0.3275)	0.3335 (0.3316-0.3352)	0.0261 (0.0197 − 0.0326)	2.5%
tot.	**66**	**30**					

Table 1. Individuals and samples per region.

F_{ST} and mean geographic distance (km)

	Region 4	Region 3	Region 2	Region 1
Region 4		*970.9*	*1550.3*	*2266.9*
Region 3	0.075***		*588.7*	*1326.6*
Region 2	0.089***	0.041***		*757.0*
Region 1	0.113***	0.072***	0.040***	

Table 2. Nei's pairwise F_{ST} and geographic distances (in italics) between populations inhabiting the four geographic regions. The significance level ($P<0.01$) of the F_{ST} statistics is denoted with ***.