Diminished adherence of *Biomphalaria glabrata* embryonic cell line to

sporocysts of *Schistosoma mansoni* following programmed knockout of the

allograft inflammatory factor

Fernanda Sales Coelho[1], Rutchanee Rodpai[2,3]André Miller[4], Shannon E. Karinshak[2,5], Victoria

H. Mann[2,5], Omar dos Santos Carvalho[1], Roberta Lima Caldeira[1], Marina de Moraes Mourão[1*],

Paul J. Brindley[2,5*], Wannaporn Ittiprasert[2,5*]

[1] Grupo de Pesquisa em Helmintologia e Malacologia Médica, Instituto René Rachou, Fundação

Oswaldo Cruz, Belo Horizonte, MG, Brazil.

[2] Department of Microbiology, Immunology and Tropical Medicine, School of Medicine and

Health Sciences, George Washington University, Washington, D.C., USA

[3] Department of Parasitology, Faculty of Medicine, Khon Kaen University, Khon Kaen, 40002

Thailand.

[4] Biomedical Research Institute , Schistosomiasis Resource Center, Rockville, MD, USA

[5] Research Center for Neglected Diseases of Poverty, School of Medicine and Health Sciences,

George Washington University, Washington, D.C., USA

*Correspondence; marina.mourao@fiocruz.br, pbrindley@gwu.edu, wannaporni@gwu.edu

E-mails:

FSC: fscoelho@aluno.fiocruz.br

RR: rutchanee5020@gmail.com

AM: amiller@afbr-bri.org

SHK: skspeight08@gmail.com

VHM: vmann@gwu.edu

OSC: omar.carvalho@fiocruz.br

RLC: roberta.caldeira@fiocruz.br

MMM: marina.mourao@fiocruz.br

PJB: pbrindley@gwu.edu

WI: wannaporni@gwu.edu

Abstract

Background: Larval development in an intermediate host gastropod snail of the genus

Biomphalaria is an obligatory component of the life cycle of *Schistosoma mansoni*.

Understanding of the mechanism(s) of host defense may hasten the development of tools that

block transmission of schistosomiasis. The allograft inflammatory factor 1, AIF, which is

evolutionarily conserved and expressed in phagocytes, is a marker of macrophage activation in

both mammals and invertebrates. AIF enhances cell proliferation and migration. The embryonic

cell line, termed Bge, from *Biomphalaria glabrata* is a versatile resource for investigation of the

snail-schistosome relationship since Bge exhibits a hemocyte-like phenotype. Hemocytes

perform central roles in innate and cellular immunity in gastropods and in some cases can kill the

parasite. However, the Bge cells do not kill the parasite *in vitro*.

Methods: Bge cells were transfected by electroporation with plasmid pCas-*Bg*AIFx4, encoding

the Cas9 nuclease and a guide RNA specific for exon 4 of the *B. glabrata* AIF (*Bg*AIF) gene.

Transcript levels for Cas9 and for *Bg*AIF were monitored by quantitative reverse-transcription-

PCR and, in parallel, adhesion of gene-edited Bge cells during co-culture with of schistosome

sporocysts was assessed.

Results: Gene knockout manipulation induced gene-disrupting indels, frequently 1-2 bp

insertions and/or 8-30 bp deletions, at the programmed target site; a range from 9 to 17% of the

copies of the *Bg*AIF gene in the Bge population of cells were mutated. Transcript levels for

*Bg*AIF were reduced by up to 73% (49.5±20.2% S.D, $P \leq 0.05$, n =12). Adherence by *Bg*AIF

gene-edited (Δ*Bg*AIF) Bge to sporocysts diminished in comparison to wild type cells, although

cell morphology did not change. Specifically, as scored by a semi-quantitative cell adherence

index (CAI), fewer ΔBgAIF than control wild type cells adhered to sporocysts; control CAI,

2.66±0.10, ΔBgAIF, 2.30±0.22 ($P \leq 0.01$).

Conclusion: The findings supported the hypothesis that BgAIF plays a role in the adherence of

B. glabrata hemocytes to sporocysts during schistosome infection *in vitro*. This demonstration

of the activity of programmed gene editing will enable functional genomics approaches using

CRISPR/Cas9 to investigate additional components of the snail-schistosome host-parasite

relationship.

Keywords: *Biomphalaria glabrata* embryonic cell line; Bge, CRISPR/Cas9; Gene editing;

Allograft inflammatory factor, Cell adhesion

Background

Evolution endowed the schistosomes with a complex life cycle that includes both a freshwater gastropod intermediate host and a definitive mammalian host. Several species of the freshwater snail genus *Biomphalaria* are the intermediate host for *Schistosoma mansoni*. The neotropical species, *Biomphalaria glabrata* has been studied extensively with respect to host-parasite relationship and coevolution with *S. mansoni* especially on mechanisms of susceptibility and/or resistance to the compatible parasites (1, 2). Genetic variation is evident among isolates and strains of *B. glabrata*, both in the laboratory and in the field, resulting in a spectrum of the susceptibility of infection with *S. mansoni* (3). Considerable advances have been made in the exploration and characterization of mechanisms of the internal defenses system (IDS) of the snail that determine susceptibility and resistance to schistosome (4-11). The resistance phenotype is underpinned by a complex genetic trait, where the schistosome larva fails to develop as the consequence of innate and cellular immune responses. Hemocytes of resistant snails encapsulate and destroy the sporocyst (11-18).

    B. glabrata embryonic cell line (Bge) (19) remains to date the only established cell line from any mollusk. The cell line originates from five-day-old embryos of *B. glabrata* susceptible to infection with *S. mansoni*. The Bge cell line has been studied extensively to interrogate the host-parasite relationship because the Bge cell exhibits a hemocyte-like behavior that includes encapsulation of the larval parasite, but does not kill the parasites (20-28).

    The genome sequence of *B. glabrata* has been reported (29), along with ongoing transcriptome and proteome catalogues that include factors participating in immunological surveillance, phagocytosis, cytokine responses, and pathogen recognition receptor elements including Toll-like receptors and fibrinogen-related proteins (30-36). An orthologue of the

evolutionary conserved allograft inflammatory factor (AIF) is an evolutionary conserved protein

typically expressed in phagocytes and granular leukocytes in both vertebrate and invertebrate.

Functions demonstrated for AIF include macrophage activation, enhancement of cellular

proliferation and of migration in mammalian and invertebrate cells; protostomes and

deuterostomes (37-41). AIF also plays a key role in the protective response by *B. glabrata* to

invasion by schistosomes (8, 9). *Bg*AIF, the orthologue in *B. glabrata* is expressed in hemocytes,

which participate in phagocytosis, cellular proliferation, and cellular migration. Elevated

expression of *Bg*AIF is a characteristic of the resistance of *B. glabrata* to schistosome infection

and has been considered as a marker of hemocyte activation (8, 9).

Expression of AIF also is seen during hemocyte activation in oysters (36, 38, 42, 43) and

during hepatic inflammation during murine schistosomiasis (44, 45). We hypothesized that

*Bg*AIF was involved in cell mediated immune response(s) by *B. glabrata* through activation of

hemocyte cell adhesion and/or migration after the schistosome miracidium has penetrated into

the tissues of the snail. We addressed this hypothesis by using CRISPR/Cas9-based

programmed genome editing to interrupt the *Bg*AIF gene of *B. glabrata* in the Bge cell line,

following reports that indicated the utility of using CRISPR-based programmed gene knockout

approach in other mollusks including the Pacific oyster, *Crassostrea gigas* and the slipper

limpet, *Crepidula fornicata* and the gastropod, *Lymnaea stagnalis* (46-48). As detailed below,

we demonstrated the activity of programmed genome editing in Bge cells, with gene knockout at

the *Bg*AIF locus.

Methods

Gene editing construct

The gene encoding the allograft inflammatory factor of *B. glabrata*, *Bg*AIF (2,226 bp; accession

number BGLB005061, https://www.vectorbase.org/) includes five exons interrupted by four

introns (Fig. 1a). A guide RNA (gRNA) for Cas9-catalyzed gene editing specific for the target *B.*

glabrata gene locus, *Bg*AIF, was identified in the BGLB005061 sequence using the

'CHOPCHOP' v3 tool, https://chopchop.cbu.uib.no/, with default parameters compatible for the

protospacer adjacent motif, NGG, of Cas9 from *Streptococcus pyogenes* (49-51) and screened

for off-target sites against the *Biomphalaria glabrata* genome (29). Based on the guidance from

the CHOPCHOP analysis, we chose the top ranked guide RNA (gRNA),

AGACTTTGTTAGGATGATGC, specific for exon 4 of the AIF gene, with predicted high

CRISPR/Cas9 efficiency for double-stranded cleavage in tandem with an absence of off-target

activity in the genome of *B. glabrata* (Fig. 1a). A CRISPR/Cas9 vector encoding the gRNA

targeting exon 4 of *Bg*AIF under the control of the mammalian U6 promoter and encoding Cas 9,

with nuclear localization signals 1 and 2, driven by the human cytomegalovirus (CMV)

immediate early enhancer and promoter was assembled using the GeneArt CRISPR Nuclease

Vector system (Thermo Fisher Scientific, Waltham, MA), according to the manufacturer's

protocol. Briefly, the 20 nt of either target (including 'gtttt' on the 3' end) or complementary to

target (including 'cggtg' on the 3'end) sequences were synthesized commercially, and annealed

according to manufacturer's protocol. The annealed double strand DNA (dsDNA) was ligated

into the linearized GeneArt® CRISPR Nuclease vector via *BamH*I and *BsmB*I restriction sites,

respectively, and the construct was termed pCas-*Bg*AIFx4 (Fig. 1b). (The sequence of GeneArt

CRISPR nuclease vector backbone is available at

https://www.thermofisher.com/order/catalog/product/A21174#/A21174). Chemically competent

TOP10, *E. coli* cells (Invitrogen, Thermo Fisher Scientific) were transformed with pCas-

*Bg*AIFx4 by the heat shock method and cultured on LB-agar supplemented with ampicillin at

100 µg/ml. Subsequently, the integrity of the recombinant plasmids from several single colonies

of ampicillin-resistant *E. coli* transformants was confirmed by amplicon PCR-based Sanger

direct nucleotide sequence analysis using a U6 gene-specific primer for gRNA ligation and

orientation (Fig. 1b).

***Biomphalaria glabrata* embryonic (Bge) cell line culture**

The Bge cell line was provided by the Schistosomiasis Resource Center (SRC), Biomedical

Research Institute (BRI), Rockville, MD. Historically, the Bge cell line was sourced by the SRC

from the American Type Culture Collection (Manassas, VA), catalog no. ATCC CRL 1494, and

thereafter maintained at BRI for >10 years. Bge cells were maintained at 26ºC in air in 'Bge

medium', which is comprised of 22% (v/v) Schneider's *Drosophila* medium, 0.13% galactose,

0.45% lactalbumin hydrolysate, 0.5% (v/v) phenol red solution, 20 µg/ml gentamycin, and

supplemented with 10% heat-inactivated fetal bovine serum (24, 52). Bge cells were grown to

80% confluence before transfection by electroporation with pCas-*Bg*AIFx4. The Bge cells were

free of contamination with *Mycoplasma*, as established with a PCR-based test (LookOut®

Mycoplasma PCR Detection kit, Sigma-Aldrich, St. Louis, MO).

Transfection of Bge cells by square wave electroporation

Bge cells were harvested using a cell scraper, washed twice in Bge medium, counted, and

resuspended at 20,000 cell/µl in Opti-MEM medium (Sigma-Aldrich, St. Louis, MO). Two

million cells were transferred into 0.2 mm path length electroporation cuvettes (BTX Harvard

Apparatus, Hollister, MA) containing 6 µg pCas-*Bg*AIFx4 in ~100 µl Opti-MEM. The cells were

subjected to electroporation using one pulse at 125 volts for 20 milliseconds, using a square

wave pulse generator (ECM 830, BTX Harvard Apparatus). Immediately thereafter, the Bge

cells were maintained in 12-well plates (Greiner Bio-One) at 26ºC. The mock control included

Opti-MEM only for electroporation. The presence of transcripts encoding the *B. glabrata* actin

and the Cas9 was monitored daily for nine days following transfection by electroporation (Fig.

1c).

Sequential isolation of total RNA and genomic DNA

To monitor the transfection of Bge cell by pCas9-*Bg*AIFx4, we investigated the expression of

Cas9 in Bge cells by reverse transcription PCR (RT-PCR). Both total RNA and genomic DNA

were extracted sequentially from cell pellets, as described (53, 54). In brief, each sample of total

RNA sample was extracted using the RNAzol® RT reagent (Molecular Research Center, Inc.,

Cincinnati, OH) according to the manufacturer's protocol. Subsequently, the DNA/protein pellet

retained after recovery of RNA was resuspend in DNAzol® solution (Molecular Research Center,

Inc), from which total DNA was recovered. The RNAs and DNAs were dissolved in nuclease-

free water and their concentration and purity established by spectrophotometry (Nanodrop 1000,

Thermo Fisher Scientific).

Expression of Cas9 in Bge cells

To investigate transcription from the pCas-*Bg*AIFx4 vector following transfection of Bge cells,

levels of transcribed Cas9 were investigated by semi-quantitative RT-PCR. The Cas9-specific

primers were Cas9-F, 5'- agcatcggccttgatatcgg-3' and Cas9-R, 5'- agaagctgtcgtccaccttg-3' (Fig.

1b). Total RNA from the non-transfected, mock (Opti-MEM electroporated-), and pCas-

*Bg*AIFx4 DNA electroporated-Bge cells were treated with DNase I (Ambion, Thermo Fisher

Scientific) to digest any residual vector pCas-*Bg*AIFx4 DNA and contaminating genomic DNAs.

The RNAs were reverse transcribed to cDNA using ProtoScript II reverse transcriptase with

oligo dT and random primers (First Strand cDNA Synthesis kit New England Biolabs, Ipswich,

MA). RT-PCRs specific for the Cas9 or actin gene of *B. glabrata, Bg*Actin (GenBank accession

number U53348.1) were undertaken, with BgActin serving as the positive control for RNA

integrity. The primer pairs used for the *Bg*Actin coding sequences were termed actin-F, 5'-

aagcgacgttttcttggtgc-3' and actin-R, 5'- acccataccaaccatcacacc-3'. Amplicons and molecular size

standards were separated by electrophoresis through Tris-acetate-EDTA-buffered agarose 1%

stained with ethidium bromide (Fig. 1c).

Analysis of programmed mutation of the allograft inflammatory factor gene of *B. glabrata*

Genomic DNA samples from the mock-transfected and pCas-*Bg*AIFx4-transfected cells were

amplified by PCR using AIF-F (5'-gcagatttgcaattcaacactta-3') and AIF-R (5'-

tgccagctagcttactgcat-3') primers that flank the CRISPR/Cas9 programmed double-stranded break

(DSB) site (Fig. 1a). Amplicons of 568 nt in length (from residues 489 to 1,056 of the

*Bg*AIF_BGLB0055061 gene) were obtained using the AIF-F and -R primer pair. Amplicons

were isolated from the agarose gel using the PCR cleanup and gel extraction kit (ClonTech,

Takara USA, Mountain View, CA) and the nucleotide sequence of amplicons determined by

Sanger direct sequencing (GENEWIZ, South Plainfield, NJ). Chromatograms of the sequence

reads from the control and experimental groups in each replicate experiment were subjected to

online analysis using the TIDE algorithm, https://tide.deskgen.com/ (55, 56) and also using the

Inference of CRISPR v2 Edits analysis (ICE) software, https://ice.synthego.com/#/ (Synthego

Corporation, Redwood City, CA) (57). Estimates of CRISPR efficiency, insertion-deletion

(INDEL)-substitution percentages, and the nucleotide sequence of mutant alleles were obtained

using both the TIDE and the ICE platforms (55, 56) (Fig. 2a, 2b).

Quantitative real time PCR analysis of transcription of *Bg*AIF

To evaluate the differential levels of the *Bg*AIF transcript among the control and experimental

groups, total RNAs were extracted and treated with DNase I, as above. DNase I treated-RNA

(200 ng) was reverse transcribed to cDNA, followed by quantitative RT-PCR, using the ViiA7

Real Time PCR System (Applied Biosystems, Scientific), and the SSoAdvanced Universal

SYBR Green Supermix reagents (Bio-Rad), according to the manufacturer's recommendations.

The following nucleotide primers used *Bg*AIF gene-specific primers amplify 119-257 nt of

*Bg*AIF GenBank accession number EX001601.1; *Bg*AIF-rt-F, 5'-cctgcttttaacccgacaga-3' and

*Bg*AIF-rt-R, 5'-tgaatgaaagctcctcgtca-3'. Differential *Bg*AIF gene expression were calculated

after normalizing with *Bg*Actin (primers as above) and comparison with the non-treated (control)

cells. The ΔΔCt method was used to calculate the differential gene expression (58), with

assistance of the GraphPad Prism 8 software (San Diego, CA) (Fig. 2c).

Schistosome sporocysts

Miracidia of the NMRI strain of *S. mansoni* were hatched from eggs recovered from livers of

schistosome infected mice (Schistosomiasis Resource Center, Biomedical Research Institute,

Rockville, MD) under axenic conditions (28), primary sporocysts were transformed from the

miracidia *in vitro*, as described (26). Briefly, miracidia were immobilized by chilling on ice for

25 min, following by pelleting using centrifugation, $500 \times g$ at 4°C, 60 sec. The miracidia were

washed with ice cold Chernin's balanced salt solution (28 mM NaCl, 0.5mM Na_2HPO_4, 2mM

KCl, 1.8mM $MgSO_4.7H_2O$, 0.6mM $NaHCO_3$, 3.6 mM $CaCl_2.2H_2O$) with 1 mg/ml glucose,

trehalose, and antibiotic, 10 µl/ml of 100·× penicillin/streptomycin (Thermo Fisher Scientific),

termed $CBSS^+$. Approximately 5,000 miracidia per well of 24-well plates were cultured in

CBSS+ at 26°C for 24 hrs, after which the sporocysts were washed to remove shed ciliated

epidermal plates and other debris, followed by transfer to a 1.5 ml microcentrifuge tube (26).

Sporocyst-Bge cell binding assay and cell adhesion index (CAI)

To investigate the if *Bg*AIF would affect the ability of cell adhesion to *S. mansoni* sporocyst, we

co-cultured the non-transfected Bge cell or non-selected-, transfected-pCas-*Bg*AIFx4 cells

(*Bg*AIF depleted- cells named 'Δ*Bg*AIF-Bge') with *in vitro* transformed sporocysts, then the cell

adhesion index (CAI) were calculated as described (26, 59). With the limitations in this study,

we were not be able to select or enrich for *Bg*AIF edited-cells, and hence the Δ*Bg*AIF-Bge cell

populations can be considered to be a population of gene mutated mixed with non-modified

(wild type) cells. CAI is a semi-quantitative method of cell adhesion to primary sporocysts using

the four categories of scores ranging from one to four - lower to higher numbers of cells adherent

to the parasite's surface. In brief, we mixed single cell suspensions of 500,000 Bge cells with

200 freshly prepared-sporocysts (total volume 200 µl of $CBSS^+$) in sterile, siliconized tubes (Bio

Plas, Thomas Scientific, Swedesboro, NJ). The Bge cell-sporocyst co-culture was maintained at

26°C for 24 hrs. Cellular morphology and adhesion of the cells to the surface of the sporocysts

was monitored and recorded using an inverted microscope, at 20× magnification (Zeiss Axio

Observer A1, Carl Zeiss LLC, White Plains, NY) after gently transferring the parasite-cell

suspension to the tissue culture plate (Greiner Bio-One). Scoring of the adherence index was

carried out in a blinded fashion to the investigator reading the score; ≥50 sporocysts from each

experimental group were counted each time, and triplicates of each treatment group were scored.

Seven independent biological replicates of this CAI-based sporocyst-Bge cell binding assay were

carried out. In total, ≥400 sporocysts were examined from each treatment and control group.

Averages for the CAI values were calculated from the cell adhesion scores ranging from 1 to 4

(examples presented in Fig. 3a) according to the formula, CAI = total binding value per number

of sporocysts (26).

Results

Cas9 nuclease transcribed in transfected Bge cells

Total RNA was extracted from non-transfected cell (wild type; WT), mock control and pCas-

*Bg*AIFx4-transfected Bge cells to assess the expression of Cas9 (Fig. 1b). The cDNAs from

either controls or pCas-*Bg*AIFx4-transfected Bge cells were employed as the template in PCRs

using two primer pairs, one specific for Cas9 and the other for *Bg*Actin, the actin gene of *B.*

glabrata that served as the reference gene (Fig. 1b, c). Transcripts encoding Cas9 in transient

pCas-*Bg*AIFx4 transfected-Bge cells were detected at 24 hrs after transfection and expression

was maintained for the nine days of the assay. The specific amplicon of Cas9 mRNA (231 bp)

was observed from pCas-BgAIFx4 transfected cells, but was absent from the non-transfected

cells (Fig. 1c). Our findings supported previous findings that revealed s CMV promoter driven

luciferase in Bge cells (60). Expression of the control reference *Bg*Actin was observed at 214 bp

amplicon in both controls and experimental samples (Fig. 1c).

Programmed mutation of *Bg*AIF confirmed functional CRISPR/Cas9 activity in Bge cells

Genomic DNAs from wild type Bge, medium-transfected (mock) and pCas-*Bg*AIFx4-transfected

cells were used as the template for PCRs with the primer pair, AIF-F and AIF-R, flanking the

programmed Cas9 cleavage site on *Bg*AIF, exon 4 (Fig. 1a, green arrows; amplicon size, ~200

nt). The red arrow indicates the predicted site of the Cas9-catalyzed double-strand break (DSB)

within the *Bg*AIF locus (Fig 1a). The nucleotide sequence of the amplicons was determined by

Sanger direct sequencing using the same primers. Both forward and reverse Sanger direct

sequencing reads from the same amplicon were estimated for insertion-deletion (INDELs) by the

ICE and the TIDE algorithms (55, 56). The reads from the Bge cells transfected with the pCas9-

*Bg*AIFx4 contained INDELs at or around the programmed CRISPR/Cas9 cleavage site. The

percentage of reads that included INDELs ranged from 8.9% to 17.1%, in the 12 biological

replicates that were carried out (Fig. 2a, b). Notably, the mutation profile in the vicinity of the

predicted DSB in *Bg*AIF was similar among these 12 replicates, which were undertaken

independently. Commonly observed+ INDELs at the DSBs site as revealed by the ICE analysis

included deletions of 8 to 30 bp and insertions of 1 or 2 bp (Fig. 2a). These mutations were

predicted to result in frameshift mutations, the consequent loss of the open reading frame, and

hence and permanent knockout of *Bg*AIF in the gene-edited Bge cell. The profile of the

frequency of mutations observed in each biological replicate was used to plot the curve (Prism 8

software) presented in figure 2b. These findings demonstrated that programmed genome editing

using CRISPR/Cas9 was active in Bge cells, and that the non-homologous end-joining (NHEJ)

pathway (61) was active in *B. glabrata* for the repair of programmed double-stranded breaks,

leading to targeted gene knockout.

Programmed mutation interrupted expression of *Bg*AIF

The aims of the study included the investigation of the activity or not CRISPR/Cas9 gene editing

in the Bge cell line and addressing the hypothesis that AIF functions in the activation of a

macrophage like phenotype by the Bge cell. Accordingly, Bge cells were transfected with pCas9-

*Bg*AIFx4 plasmid DNA. The experimental approach did not include drug resistance and/or

reporter gene markers in order to enrich for transfected Bge cells. However, even without

enrichment for transfected cells, there was a highly statistically significant reduction in levels of

*Bg*AIF transcripts in the transfected Bge cell population. Expression of *Bg*AIF transcripts were

assessed using RNAs from the cells at nine days post transfection. Comparison of the

experimental and control groups revealed significantly reduced levels of the *Bg*AIF in the pCas-

*Bg*AIFx4-transfected cells, mean 49.55±20.22%, range 28.1 to 86.3% (n =12) compared to the

wild type Bge (normalized sample, 100% expression), mock control cells (unpaired *t*-test and *F*

test to compare variances; F, DFn, Dfd=294.7, 11, 11, *P* < 0.0001) (Fig. 1b). An inverse

correlation between the percentage of INDELs and reduction in transcript levels was not

apparent (not shown).

Programmed knockout of *Bg*AIF interfered with adherence of Bge cells to schistosome

sporocysts

Single cell suspensions of Bge cells in the mock-treated and Δ*Bg*AIF groups were co-cultured

for 24 hrs in siliconized tubes with primary *S. mansoni* sporocysts. At that point, the numbers of

cells that had adhered to each sporocyst were scored. This was accomplished by examination of

at least five discrete sites of the well of the 24-well plate with ≥50 sporocysts of each group. The

cell adhesion index (CAI) were scored from 1 to 4, with a score of 1 indicating few or no

adherent cells and a score of 4 indicating that cells or clumps of cells covered more than half the

tegumental surface of the sporocyst, as defined in earlier reports (26) (Fig. 3a). Cells from

mock-transfected control group mostly adhered in clumps or singly to the surface of the parasite

(representative images in the upper panels of Fig. 3B), with CAI values that ranged from 2 to 4.

By contrast, fewer cells adhered to the surface of the sporocysts in the ΔBgAIF-Bge group

(representative images, lower panels in Fig. 3B), with CAI values ranging from 2 to 3. Only

~20% of the ΔBgAIF-Bge cells adhered to the surface of the sporocysts and most of the cells

retained remained spread singly on the surface of the well of tissue culture plate (Fig. 3b). More

specifically, the mean CAI values ascertained from the seven biological replicates (≥50 parasites

in each replicate (≥400 parasites scored), mean 2.66±0.10, range, 2.53 to 2.78 in the mock-

treated, transfection control group was significantly higher than the ΔBgAIF group, mean

2.25±0.22, range, 2.08 to 2.55 (Fig 3c) (unpaired t-test: t=3.661 df = 12, P = 0.0033). More

specifically, the CAI category-specific CAI values for mock treated cells averaged from the cell

adherence to single sporocysts, were 2.5±1.58 SE, 29.28±4.06, 20.85±2.15, and 13.14±2.55 for

categories 1, 2, 3 and 4, respectively. For the ΔBgAIF-Bge cells, the CAI values were

8.86±1.57, 31.42±4.07, 13.28±2.51, and 13.13±2.55 for categories 1, 2, 3 and 4, respectively.

Although we observed CAI scores of 1 to 4 in both mock control cells and ΔBgAIF cells,

nonetheless there were statistically significant higher numbers of sporocysts with the lowest

adherence, scored as '1', in the ΔBgAIF group compared with the mock control group as

confirmed using a multiple t-test (t ratio = 3.98, df = 12, P = 0.001). By contrast, there were

significantly higher numbers of sporocysts scored as '3' in the control compared with the

ΔBgAIF group (t ratio = 3.52032, df = 12, P = 0.004) (Fig. 3d). Last, morphological changes

were not apparent between the ΔBgAIF and the control group Bge cells.

Discussion

This report describes a novel use of programmed genome editing by the CRISPR/Cas9 approach

in the embryonic cell line from the gastropod snail, *B. glabrata*, an intermediate host snail of the

human blood fluke, *S. mansoni.* The Bge cell line is an informative tool in investigation of snail-

schistosome, host-parasite interactions. A key attribute of the Bge cell is its hemocyte-like

phenotype, given the central role of the snail hemocyte in innate and cellular immunity.

However, even though Bge cells adhere to the schistosome, the parasite is not killed by these

cells *in vitro*. The allograft inflammatory factor 1 (AIF) is a conserved calcium-binding protein

typically expressed in phagocytic and granular leukocytes and is a marker of macrophage

activation (38, 41, 45, 62-65). An orthologue, termed *Bg*AIF, is highly expressed in isolates of

B. glabrata that are resistant to infection with *S. mansoni* and this gene may be linked to

hemocyte activation (8, 9). Here, we targeted the *AIF* gene of *B. glabrata* embryonic cell line

using programmed gene knockout to further interrogate its role in the intermediate host-

schistosome interaction. We constructed a plasmid vector encoding the CRISPR/Cas9 nuclease

and a guide RNA targeting exon 4 of *Bg*AIF gene and the Cas9 nuclease from *Streptococcus*

pyogenes. Bge cells were transfected with the gene-editing construct by square wave

electroporation. Transcript levels of *Bg*AIF were significantly reduced by up to 71.9% following

transformation. In parallel, sequence reads of amplicons spanning the locus targeted for

programmed gene knock-out revealed on-target mutation on the *Bg*AIF gene, that had been

repaired by non-homologous end joining leading to gene-inactivating insertions and deletions. In

addition, the adherence of gene-edited Bge cells to sporocysts was significantly impeded in

comparison to control cells, as ascertained using a semi-quantitative, cell adherence index. In

our study, the %INDELs (8.9-17.1%) resulting from NHEJ after CRISPR/Cas9 gene editing on

*Bg*AIF exon 4 locus did not correlate with its transcript reduction (~50%) in all experimental

samples. Nonetheless, alternative mechanisms could be used as the fusion of suppressors with a

'dead' Cas9 which enables gene regulation and increase the level of repression of the target gene

(66).

The *B. glabrata* IDS comprises hemocytes and soluble proteins found in the hemolymph,

among them the *Bg*AIF (67-69). The response of resistant mollusks is given by the adherence

and encapsulation of sporocysts by hemocytes, leading to the parasite destruction (70). The AIF-

1 was demonstrated to be a pro-inflammatory cytokine that regulates immune-related genes of

the oyster, *Crassostrea ariakensis* (36). An orthologue in the leech *Hirudo medicinalis* promotes

macrophage-like migration by a chemotactic activity, in addition to being involved in the innate

immune responses as also seen in other species (41). The adherence of the mixed populations of

*Bg*AIF gene-edited/non edited-Bge cells to sporocysts was significantly impeded in comparison

to control cells, as ascertained using a semi-quantitative cell adherence index. These cells, albeit

in a low percentage, are less responsive to the *S. mansoni* parasite. These data suggested that, in

the presence of *S. mansoni*, the Bge cells need to secrete *Bg*AIF for activating the recruitment of

more adherent Bge cells. Thus, the *Bg*AIF protein appears to play a role in cell recognition,

migration, and/or adhesion, and to participate in the early immune response to the parasite. The

AIF gene is conserved broadly among protostomes and deuterostomes, including vertebrates, and

also in prebilaterian including sponges, where it likely performs similar functional roles in

macrophage activation and migration (71). In humans, the *Hm*AIF1 is an NF-κB pathway

regulator, a pathway that comprises a family of evolutionarily conserved proteins, important to

the immune system by participating in the expression of other proteins related to the immune

system (72, 73). Although more studies will be required to decipher the regulation of these

pathways in *B. glabrata*, after the pathogen invasion, the *Bg*AIF possibly acts throughout the

activation of the NF-κB pathway, leading to the recruitment of hemocytes and consequent

pathogen elimination (72, 74).

These findings confirmed the tractability of transfection of Bge cells by electroporation

with the genome-editing construct, pCas-*Bg*AIFx4, and that the CMV promoter drove

transcription of Cas9 in this snail species. Whereas transformation by plasmid DNA of Bge cells

by square wave electroporation appears to be novel, Bge cells have been transformed using DNA

complexed with cationic lipid-based transfection reagents and with polyethyleneimine (23)

Nevertheless, our study has some limitations. Thus far we have yet to enrich the transfected cells

from wild type cells. Future studies using a drug selectable marker can be designed to address

this issue. Other approaches to deliver the CRISPR/Cas gene-editing cargo can be tried

including repeated inoculation with ribonuclear protein complexes (75), titration of the

transfection chemicals (76), titration of electroporation parameters (77), and/or transduction by

lentiviral virions encoding the gRNA and *S. pyogenes* Cas9 nuclease as we have demonstrated

with eggs *of S. mansoni* (53, 54). Moreover, CIRCLE-Seq and like approaches can be employed

to investigate the off-target mutations (78).

Conclusions

Here we demonstrated CRISPR/Cas-based gene editing in a cell line from a medically important

taxon of freshwater gastropods that are vectors for the transmission of schistosomiasis. We

showed the functional role of a *B. glabrata* allograft inflammatory factor in the

recognition/attachment of *S. mansoni* sporocysts *in vitro*. The demonstration of the activity of

CRISPR/Cas9 gene editing in Bge cells suggests that genome editing in the germline and

somatic tissues of intact *B. glabrata* snails will also be functional. Whereas improvements can be

anticipated in these approaches, an obvious next step will be to gene edit the intact snail *B.*

glabrata. Transfection of germline cells within the snail using microinjection can be considered

(79). These findings, together with the first application of the CRISPR/Cas technique in the

genetic edition of *Lymnaea stagnalis* mollusk (48) are a step-change since they can favor the

creation of a genetically modified *Biomphalaria* line to study the biology and physiology of the

snail as well the schistosome-intermediate host relationship. Functional genomics using

CRISPR/Cas-based genome editing in schistosomes and other trematodes responsible for major

neglected tropical diseases has been reported (53, 54). The establishment of a functional genomic

protocols involving programmed gene editing to address fundamental questions in this host-

parasite relationship using genetically modified snails and schistosomes now seems to be

feasible.

Abbreviations

AIF: allograft inflammatory factor 1; Bge: *Biomphalaria glabrata* embryonic cell line; Cas9:

CRISPR associated protein 9; RNA: ribonucleic acid; PCR: polymerase chain reaction; CAI: cell

adherence index; CRISPR: Clustered Regularly-Interspaced Short Palindromic Repeats; gRNA:

guide RNA; CMV: cytomegalovirus; DBS: double strand break; SRC: Schistosomiasis Resource

Center; BRI: Biomedical Research Institute; RT-PCR: reverse transcription PCR; DNA:

deoxyribonucleic acid; cDNA: complementary deoxyribonucleic acid; IDS: internal defense

system; EDTA: ethylenediaminetetraacetic acid; INDEL: insertion–deletion mutation; TIDE:

Tracking of Indels by Decomposition; ICE: Inference of CRISPR Edits analysis; DNase I:

deoxyribonuclease I, CBSS: Chernin's balanced salt solution; WT: wild type; NHEJ: non-

homologous end-joining; NF-κB: nuclear factor kappa-B.

Acknowledgments

415 We thank Dr. Mathilde Knight for informative discussions on the snail-schistosome

416 immunobiology and Dr. Margaret Mentink-Kane for support with the Bge cell line. Bge cells

417 were provided by the NIAID Schistosomiasis Resource Center of the Biomedical Research

418 Institute, Rockville, Maryland through NIH-NIAID contract HHSN272201000005I for

419 distribution through BEI Resources.

Ethics approval and consent to participate

421 The protocols and procedures to maintain *S. mansoni* life cycle in mice performed at the NIAID

422 Schistosomiasis Resource Center of the Biomedical Research Institute, Rockville were approved

423 by the Institutional Animal Care and Use Committee (IACUC) (protocol number 18-04) and

424 followed by United States Animal Welfare Act and George Washington University IACUC

425 policies (assurance number A3205-01)

Consent for publication

427 Not applicable

Availability of data and materials

Data supporting the conclusions of this article are included within the article. The raw datasets

used and analyzed during the present study are available from the corresponding authors upon

reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior

(CAPES) - finance code 001, Productivity fellowship from Conselho Nacional de

Desenvolvimento Científico e Tecnológico (CNPq) granted to MMM (302518/2018-5) and RLQ

(308869/2017-6) and Fapemig (APQ-01766-15), Brazil. This research was supported in part by

the Wellcome Trust (strategic award number 107475/Z/15/Z, Functional Genomics Flatworms

Initiative, Hoffmann, K.H., principal investigator, Brindley, P.J., co-investigator). For the

purpose of Open Access, the authors have applied a CC BY public copyright license to any

Author Accepted Manuscript version arising from this submission.

Authors' contributions

WI, MM and PJB designed the study. FSC and WI wrote the protocol. MMM, VMH and PJB

reviewed the protocol. WI served as study team leader and director. VHM and AM obtained

ethical approval for vertebrate animal use. FSC, RR, AM and SEK conducted the gene editing

and cell culture experiments. FSC, RR and WI performed gene mutation analysis. FSC, OSC

and RCL performed CAI analysis. WI, MMM and PJB wrote the manuscript. All authors read

and approved the final manuscript.

Author details

[1] Grupo de Helmintologia e Malacologia Médica, Instituto René Rachou, Fundação Oswaldo

Cruz, Belo Horizonte, MG, Brazil. [2] Department of Microbiology, Immunology and Tropical

Medicine, School of Medicine and Health Sciences, George Washington University,

Washington, D.C., 20037, USA. [3] Department of Parasitology, Faculty of Medicine, Khon Kaen

University, Khon Kaen, 40002 Thailand. [4] Biomedical Research Institute, Schistosomiasis

Resource Center, Rockville, MD, USA. [5] Research Center for Neglected Diseases of Poverty,

School of Medicine and Health Sciences, George Washington University, Washington, D.C.,

20037, USA

References

1.   Webster JP, Davies CM. Coevolution and compatibility in the snail-schistosome system.

Parasitology. 2001;123 Suppl:S41-56.

2.   El-Ansary A, Al-Daihan S. Important aspects of *Biomphalaria* snail-schistosome

464 interactions as targets for antischistosome drug. Med Sci Monit. 2006;12(12):RA282-92.

3.   Richards CS, Shade PC. The genetic variation of compatibility in *Biomphalaria glabrata*

466 and *Schistosoma mansoni*. J Parasitol. 1987;73(6):1146-51.

4.   Knight M, Miller A, Liu Y, Scaria P, Woodle M, Ittiprasert W. Polyethyleneimine (PEI)

468 mediated siRNA gene silencing in the *Schistosoma mansoni* snail host, *Biomphalaria glabrata*.

469 PLoS Negl Trop Dis. 2011;5(7):e1212.

5.   Jannotti-Passos LK, Andrade HM, Caldeira RL, Romanha AJ, Murta SM, Chapeaurouge

471 DA*, et al.* Proteome analysis of the cardiac and pericardial tissue of *Biomphalaria tenagophila*

472 populations susceptible and resistant to *Schistosoma mansoni* infection. Acta Trop.

473 2008;105(3):229-34.

6.   Hanington PC, Lun CM, Adema CM, Loker ES. Time series analysis of the

475 transcriptional responses of *Biomphalaria glabrata* throughout the course of intramolluscan

476 development of *Schistosoma mansoni* and *Echinostoma paraensei*. Int J Parasitol.

477 2010;40(7):819-31.

7.   Pila EA, Sullivan JT, Wu XZ, Fang J, Rudko SP, Gordy MA*, et al.* Haematopoiesis in

molluscs: A review of haemocyte development and function in gastropods, cephalopods and

bivalves. Dev Comp Immunol. 2016;58:119-28.

8. Lockyer AE, Emery AM, Kane RA, Walker AJ, Mayer CD, Mitta G*, et al.* Early

differential gene expression in haemocytes from resistant and susceptible *Biomphalaria glabrata*

strains in response to *Schistosoma mansoni*. PLoS One. 2012;7(12):e51102.

9. Larson MK, Bender RC, Bayne CJ. Resistance of *Biomphalaria glabrata* 13-16-R1 snails

to *Schistosoma mansoni* PR1 is a function of haemocyte abundance and constitutive levels of

specific transcripts in haemocytes. Int J Parasitol. 2014;44(6):343-53.

10. Pinaud S, Portela J, Duval D, Nowacki FC, Olive MA, Allienne JF*, et al.* A shift from

cellular to humoral responses contributes to innate immune memory in the vector snail

Biomphalaria glabrata. PLoS Pathog. 2016;12(1):e1005361.

11. Barbosa L, Caldeira RL, Carvalho OS, Vidigal TH, Jannotti-Passos LK, Coelho PM.

Resistance to *Schistosoma mansoni* by transplantation of APO *Biomphalaria tenagophila*.

Parasite Immunol. 2006;28(5):209-12.

12. Adema CM, Loker ES. Digenean-gastropod host associations inform on aspects of

specific immunity in snails. Dev Comp Immunol. 2015;48(2):275-83.

13. Coustau C, Gourbal B, Duval D, Yoshino TP, Adema CM, Mitta G. Advances in

gastropod immunity from the study of the interaction between the snail *Biomphalaria glabrata*

and its parasites: A review of research progress over the last decade. Fish Shellfish Immunol.

2015;46(1):5-16.

14. Mitta G, Adema CM, Gourbal B, Loker ES, Theron A. Compatibility polymorphism in

snail/schistosome interactions: From field to theory to molecular mechanisms. Dev Comp

Immunol. 2012;37(1):1-8.

15. Mitta G, Gourbal B, Grunau C, Knight M, Bridger JM, Theron A. The Compatibility

between *Biomphalaria glabrata* snails and *Schistosoma mansoni*: an increasingly complex

puzzle. Adv Parasitol. 2017;97:111-45.

16. Pila EA, Li H, Hambrook JR, Wu X, Hanington PC. Schistosomiasis from a snail's

perspective: Advances in snail immunity. Trends Parasitol. 2017;33(11):845-57.

17. Famakinde DO. Molecular context of *Schistosoma mansoni* transmission in the

molluscan environments: A mini-review. Acta Trop. 2017;176:98-104.

18. Knight M, Arican-Goktas HD, Ittiprasert W, Odoemelam EC, Miller AN, Bridger JM.

Schistosomes and snails: a molecular encounter. Front Genet. 2014;5:230.

19. Hansen EL. A cell line from embryos of *Biomphalaria glabrata* (Pulmonata):

Establishment and characteristics. K M, editor. New York: Academic Press; 1976. 75-97 p.

20. Laursen JR, Yoshino TP. *Biomphalaria glabrata* embryonic (Bge) cell line supports in

vitro miracidial transformation and early larval development of the deer liver fluke, *Fascioloides*

magna. Parasitology. 1999;118 (Pt 2):187-94.

21. Wheeler NJ, Dinguirard N, Marquez J, Gonzalez A, Zamanian M, Yoshino TP*, et al.*

Correction to: Sequence and structural variation in the genome of the *Biomphalaria glabrata*

embryonic (Bge) cell line. Parasit Vectors. 2018;11(1):566.

22. Wheeler NJ, Dinguirard N, Marquez J, Gonzalez A, Zamanian M, Yoshino TP*, et al.*

Sequence and structural variation in the genome of the *Biomphalaria glabrata* embryonic (Bge)

cell line. Parasit Vectors. 2018;11(1):496.

23. Rinaldi G, Yan H, Nacif-Pimenta R, Matchimakul P, Bridger J, Mann VH*, et al.*

Cytometric analysis, genetic manipulation and antibiotic selection of the snail embryonic cell

line Bge from *Biomphalaria glabrata*, the intermediate host of *Schistosoma mansoni*. Int J Parasitol. 2015;45(8):527-35.

24.   Odoemelam E, Raghavan N, Miller A, Bridger JM, Knight M. Revised karyotyping and gene mapping of the *Biomphalaria glabrata* embryonic (Bge) cell line. Int J Parasitol. 2009;39(6):675-81.

25.   Vermeire JJ, Boyle JP, Yoshino TP. Differential gene expression and the effects of *Biomphalaria glabrata* embryonic (Bge) cell factors during larval *Schistosoma mansoni* development. Mol Biochem Parasitol. 2004;135(1):153-7.

26.   Castillo MG, Yoshino TP. Carbohydrate inhibition of *Biomphalaria glabrata* embryonic (Bge) cell adhesion to primary sporocysts of *Schistosoma mansoni*. Parasitology. 2002;125(Pt 6):513-25.

27.   Humphries JE, Elizondo L, Yoshino TP. Protein kinase C regulation of cell spreading in the molluscan *Biomphalaria glabrata* embryonic (Bge) cell line. Biochim Biophys Acta. 2001;1540(3):243-52.

28.   Yoshino TP, Laursen JR. Production of *Schistosoma mansoni* daughter sporocysts from mother sporocysts maintained in synxenic culture with *Biomphalaria glabrata* embryonic (Bge) cells. J Parasitol. 1995;81(5):714-22.

29.   Adema CM, Hillier LW, Jones CS, Loker ES, Knight M, Minx P, *et al.* Whole genome analysis of a schistosomiasis-transmitting freshwater snail. Nat Commun. 2017;8:15451.

30.   Dheilly NM, Duval D, Mouahid G, Emans R, Allienne JF, Galinier R, *et al.* A family of variable immunoglobulin and lectin domain containing molecules in the snail *Biomphalaria glabrata*. Dev Comp Immunol. 2015;48(1):234-43.

31. Zhang SM, Loker ES. Representation of an immune responsive gene family encoding

fibrinogen-related proteins in the freshwater mollusc *Biomphalaria glabrata*, an intermediate

host for *Schistosoma mansoni*. Gene. 2004;341:255-66.

32. Portet A, Pinaud S, Tetreau G, Galinier R, Cosseau C, Duval D*, et al.* Integrated multi-

omic analyses in *Biomphalaria-Schistosoma* dialogue reveal the immunobiological significance

of FREP-*Sm*PoMuc interaction. Dev Comp Immunol. 2017;75:16-27.

33. Hanington PC, Zhang SM. The primary role of fibrinogen-related proteins in

invertebrates is defense, not coagulation. J Innate Immun. 2011;3(1):17-27.

34. Adema CM. Fibrinogen-Related Proteins (FREPs) in Mollusks. Results Probl Cell Differ.

2015;57:111-29.

35. Gordy MA, Pila EA, Hanington PC. The role of fibrinogen-related proteins in the

gastropod immune response. Fish Shellfish Immunol. 2015;46(1):39-49.

36. Xu T, Xie J, Zhu B, Liu X, Wu X. Allograft inflammatory factor 1 functions as a pro-

inflammatory cytokine in the oyster, *Crassostrea ariakensis*. PLoS One. 2014;9(4):e95859.

37. Ai XL, Yao F, Wang XJ, Duan DB, Li K, Hu ZY*, et al.* Role of allograft inflammatory

factor-1 in regulating the proliferation, migration and apoptosis of colorectal cancer cells. Nan

Fang Yi Ke Da Xue Xue Bao. 2018;38(5):511-9.

38. Zhang Y, Li J, Yu F, He X, Yu Z. Allograft inflammatory factor-1 stimulates hemocyte

immune activation by enhancing phagocytosis and expression of inflammatory cytokines in

Crassostrea gigas. Fish Shellfish Immunol. 2013;34(5):1071-7.

39. Jia J, Cai Y, Wang R, Fu K, Zhao YF. Overexpression of allograft inflammatory factor-1

promotes the proliferation and migration of human endothelial cells (HUV-EC-C) probably by

up-regulation of basic fibroblast growth factor. Pediatr Res. 2010;67(1):29-34.

40. Autieri MV, Carbone CM. Overexpression of allograft inflammatory factor-1 promotes

proliferation of vascular smooth muscle cells by cell cycle deregulation. Arterioscler Thromb

Vasc Biol. 2001;21(9):1421-6.

41. Schorn T, Drago F, Tettamanti G, Valvassori R, de Eguileor M, Vizioli J*, et al.* Homolog

of allograft inflammatory factor-1 induces macrophage migration during innate immune response

in leech. Cell Tissue Res. 2015;359(3):853-64.

42. Martin-Gomez L, Villalba A, Carballal MJ, Abollo E. Molecular characterisation of TNF,

AIF, dermatopontin and VAMP genes of the flat oyster *Ostrea edulis* and analysis of their

modulation by diseases. Gene. 2014;533(1):208-17.

43. Li J, Chen J, Zhang Y, Yu Z. Expression of allograft inflammatory factor-1 (AIF-1) in

response to bacterial challenge and tissue injury in the pearl oyster, *Pinctada martensii*. Fish

Shellfish Immunol. 2013;34(1):365-71.

44. Chen QR, Guan F, Song SM, Jin JK, Lei DS, Chen CM*, et al.* Allograft inflammatory

factor-1 alleviates liver disease of BALB/c mice infected with *Schistosoma japonicum*. Parasitol

Res. 2014;113(7):2629-39.

45. Chen QR, Guan F, Yan DJ, Lei DS, Fu L, Xia HS*, et al.* The dynamic expression of

allograft inflammatory factor-1 in hepatic tissues and splenic cells of BALB/c mice with

Schistosoma japonicum infection. Tissue Antigens. 2012;79(1):33-41.

46. Yu H, Li H, Li Q, Xu R, Yue C, Du S. Targeted gene disruption in *Pacific oyster* based

on CRISPR/Cas9 ribonucleoproteincomplexes. Mar Biotechnol (NY). 2019;21(3):301-9.

47. Perry KJ, Henry JQ. CRISPR/Cas9-mediated genome modification in the mollusc,

Crepidula fornicata. Genesis. 2015;53(2):237-44.

48.   Abe M, Kuroda R. The development of CRISPR for a mollusc establishes the formin

*Lsdia*1 as the long-sought gene for snail dextral/sinistral coiling. Development. 2019;146(9).

49.   Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. CHOPCHOP v2: a web tool

for the next generation of CRISPR genome engineering. Nucleic Acids Res.

2016;44(W1):W272-6.

50.   Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E.

CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. Nucleic Acids

Res. 2019;47(W1):W171-W4.

51.   Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E. CHOPCHOP: a

CRISPR/Cas9 and TALEN web tool for genome editing. Nucleic Acids Res. 2014;42(Web

Server issue):W401-7.

52.   Knight M, Ittiprasert W, Odoemelam EC, Adema CM, Miller A, Raghavan N, *et al.* Non-

random organization of the *Biomphalaria glabrata* genome in interphase Bge cells and the

spatial repositioning of activated genes in cells co-cultured with *Schistosoma mansoni*. Int J

Parasitol. 2011;41(1):61-70.

53.   Arunsan P, Ittiprasert W, Smout MJ, Cochran CJ, Mann VH, Chaiyadet S, *et al.*

Programmed knockout mutation of liver fluke granulin attenuates virulence of infection-induced

hepatobiliary morbidity. Elife. 2019;8.

54.   Ittiprasert W, Mann VH, Karinshak SE, Coghlan A, Rinaldi G, Sankaranarayanan G, *et*

al. Programmed genome editing of the omega-1 ribonuclease of the blood fluke, *Schistosoma*

mansoni. Elife. 2019;8.

55.   Brinkman EK, Kousholt AN, Harmsen T, Leemans C, Chen T, Jonkers J *, et al.* Easy

quantification of template-directed CRISPR/Cas9 editing. Nucleic Acids Res. 2018;46(10):e58.

56.   Brinkman EK, van Steensel B. Rapid quantitative evaluation of CRISPR Genome editing

by TIDE and TIDER. Methods Mol Biol. 2019;1961:29-44.

57.   Hsiau T, Maures T, Waite K, Yang J, Kelso R, Holden K*, et al.* Inference of CRISPR

edits from Sanger Trace Data. bioRxiv. 2018.

58.   Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods. 2001;25(4):402-8.

59.   Bayne CJ, Loker ES, Yui MA, Stephens JA. Immune-recognition of *Schistosoma*

mansoni primary sporocysts may require specific receptors on *Biomphalaria glabrata*

hemocytes. Parasite Immunol. 1984;6(6):519-28.

60.   Lardans V, Boulo V, Duclermortier P, Serra E, Mialhe E, Capron A*, et al.* DNA transfer

in a *Biomphalaria glabrata* embryonic cell line by DOTAP lipofection. Parasitol Res.

1996;82(6):574-6.

61.   Symington LS, Gautier J. Double-strand break end resection and repair pathway choice.

Annu Rev Genet. 2011;45:247-71.

62.   Donovan KM, Leidinger MR, McQuillen LP, Goeken JA, Hogan CM, Harwani SC*, et al.*

Allograft Inflammatory Factor 1 as an immunohistochemical marker for macrophages in

multiple tissues and laboratoryanimal species. Comp Med. 2018;68(5):341-8.

63.   De Zoysa M, Nikapitiya C, Kim Y, Oh C, Kang DH, Whang I*, et al.* Allograft

inflammatory factor-1 in disk abalone (*Haliotis discus discus*): molecular cloning, transcriptional

regulation against immune challenge and tissue injury. Fish Shellfish Immunol. 2010;29(2):319-

26.

64.   Li Q, Bai Z, Zhao L, Li J. Characterization of allograft inflammatory factor-1 in

Hyriopsis cumingii and its expression in response to immune challenge and pearl sac formation.

Fish Shellfish Immunol. 2016;59:241-9.

65.   Liu G, Ma H, Jiang L, Zhao Y. Allograft inflammatory factor-1 and its immune

regulation. Autoimmunity. 2007;40(2):95-102.

66.   Roman E, Coman I, Prieto D, Alonso-Monge R, Pla J. Implementation of a CRISPR-

based system for gene regulation in *Candida albicans*. mSphere. 2019;4(1).

67.   van der Knaap WP, Loker ES. Immune mechanisms in trematode-snail interactions.

Parasitol Today. 1990;6(6):175-82.

68.   Vergote D, Bouchut A, Sautiere PE, Roger E, Galinier R, Rognon A*, et al.*

Characterisation of proteins differentially present in the plasma of *Biomphalaria glabrata*

susceptible or resistant to *Echinostoma caproni*. Int J Parasitol. 2005;35(2):215-24.

69.   Negrao-Correa D, Mattos AC, Pereira CA, Martins-Souza RL, Coelho PM. Interaction of

Schistosoma mansoni sporocysts and hemocytes of *Biomphalaria*. J Parasitol Res.

2012;2012:743920.

70.   Guaraldo AM, Magalhaes LA, Rangel Hde A, Pareja G. Evolution of spopocysts of

Schistosoma mansoni Sambon 1907 in *Biomphalaria glabrata* (Say 1818) and *Biomphalaria*

tenagophila (d'Orbigny 1835). Rev Saude Publica. 1981;15(4):436-48.

71.   Watano K, Iwabuchi K, Fujii S, Ishimori N, Mitsuhashi S, Ato M*, et al.* Allograft

inflammatory factor-1 augments production of interleukin-6, -10 and -12 by a mouse

macrophage line. Immunology. 2001;104(3):307-16.

72.   Symeonidou I, Kourelis A, Frydas I, Karagouni E, Anogeianaki A, Hatzistilianou M, *et*

al. Modulation of NF-kappa beta signalling pathways by parasites. J Biol Regul Homeost

Agents. 2010;24(4):471-9.

73.   Tato CM, Hunter CA. Host-pathogen interactions: subversion and utilization of the NF-

kappa B pathway during infection. Infect Immun. 2002;70(7):3311-7.

74.   Egana-Gorrono L, Chinnasamy P, Casimiro I, Almonte VM, Parikh D, Oliveira-Paula

GH, *et al.* Allograft inflammatory factor-1 supports macrophage survival and efferocytosis and

limits necrosis in atherosclerotic plaques. Atherosclerosis. 2019;289:184-94.

75.   Lin S, Staahl BT, Alla RK, Doudna JA. Enhanced homology-directed human genome

engineering by controlled timing of CRISPR/Cas9 delivery. Elife. 2014;3:e04766.

76.   Lino CA, Harper JC, Carney JP, Timlin JA. Delivering CRISPR: a review of the

challenges and approaches. Drug Deliv. 2018;25(1):1234-57.

77.   Fajrial AK, He QQ, Wirusanti NI, Slansky JE, Ding X. A review of emerging physical

transfection methods for CRISPR/Cas9-mediated gene editing. Theranostics. 2020;10(12):5532-

49.

78.   Tsai SQ, Nguyen NT, Malagon-Lopez J, Topkar VV, Aryee MJ, Joung JK. CIRCLE-seq:

a highly sensitive *in vitro* screen for genome-wide CRISPR-Cas9 nuclease off-targets. Nat

Methods. 2017;14(6):607-14.

79.   Huang J, You W, Xu Z, Yan Q, Shi C, Tang B, *et al.* An effective microinjection method

and TALEN-Mediated genome editing in *Pacific abalone*. Mar Biotechnol (NY).

2019;21(4):441-7.

Legends to figures

Fig. 1 Schematic diagram of *Bg*AIF gene structure, CRISPR/Cas9 vector and expression in Bge

cell. **a** Gene structure of *B. glabrata* allograft inflammatory factor (*Bg*AIF), accession number

BGLB005061 and gene editing target locus (red box) on exon 4. *Bg*AIF gene composed of 5

exons and 4 introns. The green arrows indicate the location of primers flanking expected double

strand breaks (DSB) which were used in PCR to generate the on-target amplicon for INDELs

estimation. **b** Map of the pCas-*Bg*AIFx4 vector which includes the Pol III-dependent

mammalian U6 gene promoter (red arrow) to drive transcription of the guide RNA targeting

exon 4 of *Bg*AIF gene (red arrow) and the CMV promoter to drive expression of the *S. pyogenes*

Cas9 nuclease (blue arrow). Primer pairs specific for the guide RNA and for Cas9 are indicated

(green arrows). **c** Expression of Cas9 and of *Bg*Actin (as the reference gene) transcripts as

established by semi-quantitative RT-PCR in pCas-*Bg*AIF-transfected (right) and control (left)

Bge cells from days one to nine following transfection. The amplicons of the expected sizes are

as indicated: 23 bp for Cas9 and 214 bp for *Bg*Actin. All RNA samples were positive for the

*Bg*Actin reference gene; the 214 bp band.

Fig. 2 Establishment of *Bg*AIF-knockout lines of Bge cells. **a** Representative examples of

frequent gene insertions-deletions (1-2 bp insertions and 8-30 bp deletions, straddling the

programmed CRISPR/Cas9-induced double-stranded break in exon 4, as determined by ICE

software-based analysis. **b** TIDE algorithm-based violin plot of insertion-deletion percentages

(%INDEL) computed using the amplicon sequence traces from the 12 biological replicates of

pCas-*Bg*AIF-transfected Bge cell populations. **c** Reduction of *Bg*AIF transcription by about 50%

following programmed genome editing of Bge cells (Δ*Bg*AIF-Bge) in comparison to control Bge

cells. Mean transcript reduction, 49.55± 20.22 (S.D.) percent, $P \le 0.0001$ (****), n =12

(unpaired Student's *t*-test).

Fig. 3 Programmed knockout of *Bg*AIF in Bge cells caused reduced adherence to primary

sporocysts. **a** Representative micrographs of primary sporocysts co-cultured with Bge cells in

our laboratory to profile the semi-quantitative scoring of the cell adhesion index (CAI); CAI

value = 1; no cells adhering to the surface of the sporocyst; value = 2; ≤10 cells adhering to the

sporocyst; value = 3; > 10 cells < half of the sporocyst surface covered by cells or clumps of

cells; value = 4; > half the sporocyst surface covered by Bge cells. **b** Representative

micrographs indicate the reduced levels of Δ*Bg*AIF-Bge cells adherence (right panel) in

comparison to control, mock-transfected Bge cells (left panel) to the co-cultured sporocysts. **c**

Bar chart to present the CAI values from control (mock-transfected) Δ*Bg*AIF-Bge cells during

co-culture with primary sporocysts at a co-culture ratio of one sporocyst to 100 Bge cells; CAI

value = 2.66±0.10, mean ±SD (476 sporocysts in total scored) for the mock-transfected Bge and

2.31±0.23 for the Δ*Bg*AIF-Bge cells (424 sporocysts in total scored); $P = 0.0033$, unpaired

Student's *t* test; n =7 biological replicates. **d.** The cell adhesion score ranging from 1-4 of

individual sporocyst. There were 2.5±SE 1.58, 29.28±4.06, 20.85±2.15 and 13.14±2.55 for cell

adhesion scores 1, 2, 3 and 4 from mock, respectively. While in Δ*Bg*AIF-Bge cell shown

8.86±1.57, 31.42±4.07, 13.28±2.51 and 13.13±2.55 of cell adhesion scores 1, 2, 3 and 4,

respectively. The higher amount of sporocyst from *Bg*AIF edited-mixed cells population shown

statistic significantly lower than control cell by *t*-test (t ratio = 3.98, P = 0.001). Also the lower

amount of sporocysts scored as '3' in ΔBgAIF-Bge cells comparing with control cell (t ratio =

2.40, P = 0.004).

a. *BG*LB005061

100 bp

AIF-F → ← AIF-R

exon 4

DSB

AGACTTTGTTAGGATGATGC TGG

target DNA on *Bg*AIF, exon 4

exon 5

b.

U6-F → U6 gRNA tracrRNA Amp^R Amp^R promoter CMV promoter SV40 NLS Cas9-F → ← Cas9-R Cas9 SV40 NLS

pCas-*Bg*AIFx4
9.22 kb

c.

control

pCas-*Bg*AIFx4-transfected

M ← → M ← →

← actin, 214 bp

← Cas9, 277 bp

day: 1 2 3 4 5 6 7 8 9   day: 1 2 3 4 5 6 7 8 9

a

CAI value = 1 CAI value = 2 CAI value = 3 CAI value = 4

b

WT Bge ΔBgAIF-Bge

c

CAI vaule

mock ΔBgAIF-Bge

d

● mock ■ ΔBgAIF-Bge

number of sporocyst

Cell adhesion score