Tocilizumab treatment in severe COVID-19 patients attenuates the inflammatory storm incited by monocyte centric immune interactions revealed by single-cell analysis

Chuang Guo[1,7], Bin Li[1,7], Huan Ma[1], Xiaofang Wang[2], Pengfei Cai[1], Qiaoni Yu[1], Lin Zhu[1], Liying Jin[1], Chen Jiang[1], Jingwen Fang[3], Qian Liu[1], Dandan Zong[1], Wen Zhang[1], Yichen Lu[1], Kun Li[1], Xuyuan Gao[1], Binqing Fu[1,4], Lianxin Liu[2], Xiaoling Ma[5], Jianping Weng[6], Haiming Wei[1,4], Tengchuan Jin[1,4],†, Jun Lin[1,4],†, Kun Qu[1,4],†

[1]Department of oncology, The First Affiliated Hospital of USTC, Division of Molecular Medicine, Hefei National Laboratory for Physical Sciences at Microscale, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, Anhui, 230021, China.
[2]Department of Hepatobiliary Surgery, the First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, Anhui, 230021, China.
[3]HanGene Biotech, Xiaoshan Innovation Polis, Hangzhou, Zhejiang, China
[4]CAS Center for Excellence in Molecular Cell Sciences, the CAS Key Laboratory of Innate Immunity and Chronic Disease, University of Science and Technology of China, Hefei, Anhui, 230027, China.
[5]Department of Laboratory Medicine, The First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, Anhui, 230001, China
[6]Department of Endocrinology and Metabolism, The First Affiliated Hospital of USTC, Division of Life Sciences of Medicine, University of Science and Technology of China, Hefei 230026, China.
[7]These authors contributed equally to this work.
†Corresponding should be addressed to Kun Qu (qukun@ustc.edu.cn).
Jun Lin (linjun7@ustc.edu.cn); Tengchuan Jin (jint@ustc.edu.cn)

Contact Information:
Kun Qu, Ph.D.
Division of Molecular Medicine, Hefei National Laboratory for Physical Sciences at Microscale, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, Anhui, 230027, China.
Email: qukun@ustc.edu.cn
Phone: +86-551-63606257

ABSTRACT

Coronavirus disease 2019 (COVID-19) has caused more than 40,000 deaths

worldwide[1]. Approximately 14% of patients with COVID-19 experienced severe

disease and 5% were critically ill[2]. Studies have shown that dysregulation of the

COVID-19 patients' immune system may lead to inflammatory storm and cause severe

illness and even death[3,4]. Tocilizumab treatment targeting interleukin 6 receptor has

shown inspiring clinical results of severe COVID-19 patients[5]. However, the immune

network with Tocilizumab treatment at single cell resolution has not been uncovered.

Here, we profiled the single-cell transcriptomes of 13,289 peripheral blood

mononuclear cells isolated at three longitudinal stages from two severe COVID-19

patients treated with Tocilizumab. We identified a severe stage-specific monocyte

subpopulation and these cells centric immune cell interaction network connected by the

inflammatory cytokines and their receptors. The over-activated inflammatory immune

response was attenuated after Tocilizumab treatment, yet immune cells including

plasma B cells and CD8[+] T cells still exhibited an intense humoral and cell-mediated

anti-virus immune response in recovered COVID-19 patients. These results provided

critical insights into the immunopathogenesis of severe COVID-19 and revealed

fundamentals of effectiveness in Tocilizumab treatment.

Keywords : Coronavirus disease 2019 (COVID-19); Severe acute respiratory
syndrome coronavirus 2 (SARS-CoV-2); Tocilizumab; Single-cell RNA sequencing
(scRNA-seq); Inflammatory storm; Monocyte

Main

As of Apr 1, 2020, WHO reported 40,598 deaths out of 823,626 confirmed cases infected by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), and these numbers are still growing rapidly[1]. Approximately 14% of patients with COVID-19 experienced severe disease, and 5% were critically ill suffered from 49% fatality rate[2], which may be caused by patients' abnormal immune system activation[3,4,6]. Hence, there is an urgent need for researchers to understand how the immune system respond to the viral infections at severe stage and thereby provide effective treatment strategies.

Studies have shown that the inflammatory storm caused by excessive immune responses was associated with the crucial cause for mortality in COVID-19[7,8]. Plasma concentrations of a series of inflammatory cytokines, such as granulocyte-macrophage colony-stimulating factor (GM-CSF), interleukin (IL)-6[4], tumor necrosis factor α (TNF-α), IL-2, 7, 10 and granulocyte colony-stimulating factor (G-CSF)[9] were increased after SARS-CoV-2 infections. Further investigation demonstrated peripheral inflammatory monocytes and pathogenic T cells may incite cytokine storm in severe COVID-19 patients[4,7]. To calming inflammatory storm, Tocilizumab, which targeting IL-6 receptors and has proved its effectiveness in the treatment of cytokine release syndrome that is severe or life-threatening[10,11], was used in the treatment of severe COVID-19. After receiving Tocilizumab, the body temperature of the patients returned to normal after 24 hours. The concentration of oxygen inhalation was significantly decreased on the 5th day[5]. However, the immune network arousing the inflammatory storm in severe or recovery stage during Tocilizumab therapy at single cell level has not been uncovered.

Here, we profiled the peripheral immune cells of COVID-19 patients using single-cell transcriptome sequencing. We obtained 5 peripheral blood samples from 2 severe COVID-19 patients at 3 consecutive time-points from the severe to recovery stages during Tocilizumab treatment (Fig. 1a). Specifically, the blood samples at severe stage were collected within 12 hours of Tocilizumab was given. The blood samples at recovery stage were collected at the 5th and 7th day after Tocilizumab treatment. The

patients at severe stage had decreased number of lymphocytes, increased percentage of

neutrophil and concentrations of C-reaction protein, and increased expression of IL-6

(Supplementary Table 1). Peripheral blood mononuclear cells (PBMCs) were isolated

and subjected to single-cell mRNA sequencing (scRNA-seq) using the 10X platform

(Fig. 1a, Supplementary Table 2). After filtering low quality cells, we retained a total

of 13,289 single transcriptomes of PBMCs. Of these, 4,364 cells were from severe stage

and 8,925 were from patients at recovery stage.

To investigate the heterogeneity and differences of PBMCs between COVID-19

patients and healthy controls, we integrated our COVID-19 single cell transcriptomes

with the published single-cell profiles of healthy PBMC from the 10X official website[12]

and obtained a total of 69,237 cells (See **Methods**) (Fig. 1b-d). We then applied Seurat[13]

to normalize and cluster the gene expression matrix, and identified 20 unique cell

subsets, which were visualized via uniform manifold approximation and projection

(UMAP) (Fig. 1b-d). Cell lineages, including monocytes, $CD4^+$ and $CD8^+$ T, $\gamma\delta$T, NK,

B, plasma B and myeloid dendritic cells (mDC), plasmacytoid dendritic cells (pDC),

platelet and $CD34^+$ progenitor cells were identified based on the expression of known

marker genes (Fig. 1e, Extended Data Fig.1a). With that, we delineated the landscape

of circulating immune cells in severe COVID-19 patients.

We next explored the distribution of immune cells from the severe, recovery and

healthy stage in each cell subpopulation (Fig. 1f, Extended Data Fig.1b). We observed

that a monocyte subpopulation (cluster 9) existed only in patients at severe stage.

Plasma B cells (cluster 11), effector $CD8^+$ T (cluster 6), proliferative $MKI67^+CD8^+$ T

cells (cluster 12) and NK cells (cluster 7) were significantly enriched in patients versus

control. However, a number of subpopulations, such as $\gamma\delta$ T cells (cluster 8), pDCs

(cluster 15) and mDCs (cluster 10 and 19), most monocytes (cluster 2, 13 and 14)

existed only in patients at recovery stage and healthy controls, indicating that these cell

types gradually become normal after the treatment. No significant differences were

observed in $CD4^+$ T (cluster 1 and 4), naïve $CD8^+$ T (cluster 3) and B cells (cluster 5)

in patients versus control.

Monocytes were reported to play a vital role in CAR-T induced cytokine-release

syndrome[14] and SARS-CoV-2 infection caused inflammatory storm[4], therefore we

explored the features and functions of the monocyte from COVID-19 patients. We

detected 1,737 monocytes in patients, 927 from severe stage and 810 from recovery

stage, and integrated 9,787 monocytes from health control. The UMAP plot displayed

two main clouds of monocytes that were clearly segregated (Fig 2a). One monocyte

subpopulation (cluster 9) was almost exclusively consisted of cells from severe stage

and others (cluster 2, 13, 14, 17) were dominated by cells from the recovery and healthy

stages (Fig 2b), suggesting a severe-stage specific monocyte subpopulation.

We then investigated the expressions of several selected inflammatory cytokines

and observed that these genes were all significantly enriched in severe stage-specific

monocytes (Fig. 2c, $P < 0.001$, Wilcoxon rank-sum test). To further explore the

transcriptional differences among the monocytes' subtypes, we performed a pairwise

comparison of the gene expressions in severe, recovery and healthy stages. We obtained

2,335 differentially expressed genes (DEGs) enriched in each stage, within which

reported cytokine storm related genes, such as *TNF*[9], *IL10*[9], *CCL3*[9] and *IL6*[4] were found

significantly higher expressed in severe stage-specific monocytes (Fig 2d,

Supplementary Table 3). In addition, we also discovered a large number of significant

and inflammatory related genes that were less reported (Fig 2d, fold change > 2, $P <$

10^{-3}), including chemokine genes *CCL4*, *CCL20*, *CXCL2*, *CXCL8* and *CXCL9*,

inflammasome activation associated genes *NLRP3* and *IL1B*, and complement pathway

genes *C1QA*, *C1QB* and *C1QC* (Extended Data Fig.2a-c, Supplementary Table 4).

These results indicated that this monocyte subpopulation may contribute to the

inflammatory storm in severe COVID-19 patients.

We also observed that genes involved in "acute inflammatory response" and

"leukocyte chemotaxis" were significantly decreased at recovery and healthy stage

(Fig.2e, f, Supplementary Table 5), suggesting that the inflammatory storm caused by

this monocyte subpopulation was suppressed after the Tocilizumab treatment.

Next, we explored the transcription factors (TFs) that may be involved in the

regulation of inflammatory storm in monocytes. We used SCENIC[15] and predicted 15

TFs that may regulate genes which were enriched in severe stage-specific monocyte

(Fig 2g). By constructing a gene regulatory network, we found 3 of them, namely ETS2,

NFIL3 and PHLDA2 were regulating the cytokine storm relevant genes (Extended Data

Fig.2d). In addition, we found the expressions of *ETS2*, *NFIL3* and *PHLDA2* and their

target genes were enhanced in severe-specific monocyte subpopulation (Fig 2h),

suggesting these 3 TFs may regulate the inflammatory storm in monocytes.

Given that monocytes in the severe stage may be involved in the regulation of a

variety of immune cell types, we used the accumulated ligand/receptor interaction

database[16] CellPhoneDB (www.cellphonedb.org) to identify alterations of molecular

interactions between monocytes and all the immune cell subsets (Supplementary Table

6). We found 15 pairs of cytokines and their receptors whose interaction were

significantly altered in severe versus recovery and healthy stages (Fig 3a). Consistent

with previous study[4], we found monocytes interacted with CD4[+] T cells and plasma B

cells in patients at severe stage through the ligand/receptor pairs of IL-6/IL-6R. This

interaction, together with many other cytokine storm relevant cell-cell communications

were then extensively attenuated after the treatment of Tocilizumab (Fig 3b), suggesting

that this drug may functioning by effectively blocking the inflammatory storm in severe

COVID-19 patients.

In addition, we also discovered many other ligand/receptor pairs involved in a

broader spectrum of immune cell communications enriched at the severe stage. For

example, TNF-α and its receptors, which connected monocytes with many other

immune cells. Others like IL-1β and its receptor, which connected monocytes with

CD8[+] T cells. Chemokines, such as CCL4L2, CCL3 and CCL4 and their receptors were

also found enriched at severe stage. These cytokines and their receptors may serve as

potential drug targets to treat COVID-19 patients at severe stage, and some of their

inhibitors are undergoing clinical trials in multiple places around the world

(Supplementary Table 7). Collectively, these findings illustrated the molecular basis of

cell-cell interactions at the peripheral blood of COVID-19 patients, leading to a better

understanding of the mechanisms of inflammatory storm of the disease.

Robust multi-factorial immune responses can be elicited against viral infection,

such as avian H7N9 disease[17,18]. A recent report has found effective immune responses

from a non-severe COVID-19 patient[19]. However, it is not clear whether the anti-virus

immune response would be affected after Tocilizumab treatment. Therefore, the anti-

virus immune responses from the humoral and cell-mediated immunity in COVID-19

patients at severe stage were compared with recovery stage and healthy controls. As

expected, we found plasma B cells were barely existing in healthy controls (Fig. 4a).

By contrast, significantly higher proportion of plasma B cells was exclusively increased

in both severe and recovery stages (Fig. 4a, b), suggesting powerful anti-virus humoral

immune responses during Tocilizumab treatment.

CD8$^+$ T cells are a critical component of cell-mediated immunity against viral

infections by killing infected cells and secreting proinflammatory cytokines. To identify

the anti-virus immune responses from the cell-mediated immunity during Tocilizumab

treatment, we detected 12,121 CD8$^+$ T cells from our analysis. Clustering these cells

revealed 3 subtypes: naïve CD8$^+$ T cells (cluster 3), effector CD8$^+$ T cells (cluster 6)

and a subset of CD8$^+$ T cells with proliferation characteristics (cluster 12) (Fig. 4c, d).

Among them, the cells from the severe patients were mainly distributed in the effector

CD8$^+$ T cell cluster (Fig. 4c, d). We then conducted pairwise comparisons to identify

DEGs of CD8$^+$ T cells between the severe, recovery and healthy stages (Fig. 4e,

Supplementary Table 8). We found that genes enriched in severe stage showed

"regulation of cell activation" signatures (Fig. 4f, Supplementary Table 9; $P < 10^{-10}$).

Meanwhile, genes involved in "leukocyte mediated cytotoxicity" and "positive

regulation of cytokine production" were highly enriched in CD8$^+$ T cells from COVID-

19 patients at both severe and recovery stage (Fig. 4g, Supplementary Table 9; $P < 10^-$

[5]). We further detected elevated expression of the 108 and 449 genes involved in these

GO terms (Fig. 4h, i, Supplementary Table 10). Together, these results demonstrated

the critical evidence that robust adaptive immune responses against SARS-CoV-2

infection exist in severe stage and remain after Tocilizumab treatment.

The immune system is crucial to fight off viral infection[20,21]. Recent studies have

illustrated that monocytes may be the main cause of exacerbation and even death of

COVID-19 patients through inflammatory storms[4]. In this study, we discovered a

specific monocyte subpopulation that may lead to the inflammatory storm in patients

at severe stage through single-cell mRNA sequencing. By analyzing the monocyte-

centric ligand/receptor interactions, we revealed a severe stage-specific landscape of

peripheral immune cell communication that may drive the inflammatory storm in

COVID-19 patients. With that we obtained a list of cytokine storm relevant

ligand/receptors that can serve as candidate drug targets to treat the disease, and

provided mechanistic insights of the immunopathogenesis of COVID-19.

There are always questions about whether Tocilizumab treatment may affect the

antiviral effect of the body[22,23]. Our single cell profiles illustrated a sustained humoral

and cell-mediated anti-virus immune response of COVID-19 patients at both severe and

recovery stage. For example, the proportion of plasma B cells with antibody-secreting

function were keeping at high levels and the cytotoxicity and cytokine production of

effector CD8$^+$ T cells were also remained stable in severe COVID-19 patients after

Tocilizumab treatment.

The distributions of NK cells were significantly different in the two patients

(Extended Data Fig.1b), and the analysis of gene expression differences did not enrich

significant biological functions, therefore we did not discuss them in depth. The normal

functions of other cell types, such as γδT cells and DCs, were almost lost under severe

conditions, and the contribution of these cells to the progression of the disease requires

further investigation.

Methods

Human samples

Peripheral blood samples were obtained from two severe COVID-19 patients. The patient severity was defined by the "Diagnosis and Treatment of COVID-19 (Trial Version 6)" which was released by The General Office of the National Health Commission and the Office of the National Administration of Traditional Chinese Medicine. Patient PZ was defined as a severe patient for his peripheral capillary oxygen saturation (SPO2) <93%. Patient PZ provided 2 blood samples at severe stage (Day 1) and recovery stage (Day 5). Patient PW was defined as critical ill for respiratory failure, multiple organ dysfunction (MOD) and SPO2 <93% under high flow oxygen (50 L/min, FIO2 50%). Patient PW provided 3 blood samples at severe stage (Day 1) and recovery stage (Day 5 and Day 7). All samples were collected from the First Affiliated Hospital of University of Science and Technology of China. Before blood draw, informed consent was obtained from each patient. Ethical approvals were obtained from the ethics committee of the First Affiliated Hospital of the University of Science and Technology of China (No. 2020-XG(H)-020).

Cell Isolation

We collected 2ml peripheral blood each time from the COVID-19 patients. Peripheral blood mononuclear cells (PBMC) were freshly isolated from the whole blood by using a density gradient centrifugation using Ficoll-Paque and cryopreserved for subsequent generation of single-cell RNA library.

Single-cell RNA-seq

We generated single-cell transcriptome library following the instructions of single-cell 3' solution v2 reagent kit (10x Genomics). Briefly, after thawing, washing and counting cells, we loaded the cell suspensions onto a chromium single-cell chip along with partitioning oil, reverse transcription (RT) reagents, and a collection of gel beads that

contain 3,500,000 unique 10X Barcodes. After generation of single-cell gel bead-in-

emulsions (GEMs), RT was performed using a C1000 Touch[TM] Thermal Cycler (Bio-

Rad). The amplified cDNA was purified with SPRIselect beads (Beckman Coulter).

Single-cell libraries were then constructed following fragmentation, end repair, polyA-

tailing, adaptor ligation, and size selection based on the manufacturer's standard

parameters. Each sequencing library was generated with unique sample index. Libraries

were sequenced on the Illumina NovaSeq 6000 system.

Single-cell RNA-seq data processing

The raw sequencing data of patients and health donors were processed using Cell

Ranger (version 3.1.0) against the GRCh38 human reference genome with default

parameters, and data from different patients and disease stages were combined by the

Cell Ranger 'aggr' function. We are uploading the scRNA-seq data of PBMCs from the

2 severe COVID-19 patients to the Genome Sequence Archive at BIG Data Center and

the accession number will be available upon request. We also used the scRNA-seq data

of PBMCs from 2 healthy donors, which can be downloaded from the 10X genomics

official website. Firstly, we filtered low quality cells using Seurat[13] (version 3.1.4). For

cells from COVID-19 patients (PZ and PW), we retained cells with detected gene

numbers between 500 and 6,000 and mitochondrial UMIs less than 10%. For cells from

healthy donors, we retained cells with detected gene numbers between 300 and 5,000

and mitochondrial UMIs less than 10%. Subsequently we normalized gene counts for

each cell using the "NormalizeData" function of Seurat with default parameters.

In downstream data processing, we used canonical correlation analysis (CCA) and

the top 40 canonical components to find the "anchor" cells between patients and healthy

controls. We then used the "IntegrateData" function in Seurat to integrate all the cells.

We clustered cells based on the integrated dataset using Seurat with parameter

"resolution=0.3" and generated 20 clusters. To display cells in a 2-dimensional space,

we ran the principal component analysis (PCA) on the integrated dataset and adopted

the first 50 principal components for the uniform manifold approximation and

projection (UMAP) analysis.

Differential expression analysis

To search for the differentially expressed genes (DEGs), we first assign the negative

elements in the integrated expression matrix to zero. We used Wilcoxon rank-sum test

to search for the DEGs between each pair of the 3 stages of cells (i.e. severe stage,

recovery stage and healthy control). We applied multiple thresholds to screen for DEGs,

including mean fold-change >2, P value <0.001, and were detected in >10% of cells in

at least one stage.

We defined stage A specific-DEGs as the intersections between the DEGs in stage

A versus stage B and the DEGs in stage A versus stage C. We defined stage A and B

common-DEGs as the intersections of the DEGs in stage A versus stage C and the

DEGs in the stage B versus stage C, minus the DEGs between stage A and B. In this

way, we obtained the specific-DEGs for each stage, and the common-DEGs for each

pair of the 3 stages. We then uploaded these DEG groups to the Metascape[24] website

(https://metascape.org/gp/index.html#/main/step1), and used the default parameters

perform Gene Ontology (GO) analysis for each stage.

Motif enrichment and regulatory network

We adopted SCENIC[15] (version 1.1.2) and RcisTarget database to build the gene

regulatory network of CD14$^+$ monocytes. Since the number of CD14$^+$ monocytes from

healthy control (N = 9,618) was more than those from the severe and recovery stages

(N = 1,607), to balance their contributions in the motif analysis, we randomly sampled

2,000 CD14$^+$ monocytes from the healthy control for calculation. We selected 13,344

genes that were detected in at least 100 monocytes or included in the DEGs of the 3

stages as the input features for SCENIC. With default parameters, SCENIC generated

the enrichment scores of 427 motifs. We used the student's t-test to calculate the P

values of these motifs between severe stage and healthy control, and selected severe-

specific enriched motifs with fold change >1.5 and *P* value < 10^{-100}.

We then applied the enrichment scores of the severe-specific enriched motifs and

the expressions of their targeted genes to Cytoscape[25] to construct a connection map for

the gene regulatory network, as shown in Extended Data Fig. 2b. The thickness of line

connecting TFs and target genes represented the weight of regulatory link predicted by

SCENIC.

Ligand/receptor interaction analysis

To identify potential cellular communications between monocytes and other cell types

(CD4[+] T, CD8[+] T, B, plasma B and NK cells), we applied the CellphoneDB[16] algorithm

to the scRNA-seq profiles from the severe, recovery and healthy stages. CellphoneDB

evaluated the impact of a ligand/receptor interactions based on the ligand expression in

one cell type and its corresponding receptor expression in another cell type. To identify

the enriched ligand/receptor interactions in patients at severe stage, we selected the

ligand/receptor interactions with more significant (*P* value < 0.05) cell-cell interaction

pairs in the severe stage than that in the recovery and healthy stages. We also included

ligand/receptor pairs which were highly expressed in severe stage.

Data Availability

We are uploading the scRNA-seq data of PBMCs from the 2 severe COVID-19 patients

to the Genome Sequence Archive at BIG Data Center and the accession number will be

available upon request. We also used the scRNA-seq data of PBMCs from 2 healthy

donors, which can be downloaded from the 10X genomics official website.

Acknowledgements

Funding: This work was supported by the National Key R&D Program of China

(2017YFA0102900 to K.Q.), the National Natural Science Foundation of China grants

(81788101, 31970858, 31771428, 91940306 and 91640113 to K.Q., 31700796 to C.G.

and 81871479 to J.L.), the Fundamental Research Funds for the Central Universities

(to K.Q.). We thank the USTC supercomputing center and the School of Life Science

Bioinformatics Center for providing supercomputing resources for this project.

Author Contributions

K.Q. conceived and supervised the project; K.Q., C.G. and J.L. designed the

experiments; C.G. and J.L. performed the experiments and conducted all the sample

preparation for NGS with the help from H.M. and T.C.; B.L. performed the data

analysis with the help from P.C., Q.Y., L.Z., L.J., C.J., Q.L., D.Z., W.Z., Y.L., K.L.,

X.G. and J.F; T.C., X.W., L.L. and X.M. provided COVID-19 blood samples and

clinical information. K.Q., C.G., J.L. and B.L. wrote the manuscript with the help of

B.F., H.W. and all the other authors.

Competing interests

Jingwen Fang is the chief executive officer of HanGen Biotech.

Figure Legends

Figure 1 | An atlas of peripheral immune cells in severe COVID-19 patients. a,

Flowchart depicting the overall design of the study. Blood draws from patient PZ were

performed at 2 time points (Day1 and Day5), and PW at 3 time points (Day1, Day5,

and Day7). Patients at Day 1 were at severe stage and Day 5 and Day 7 were at recovery

stages. Samples were collected within 12 hours of Tocilizumab was given at Day 1. **b-**

d, UMAP representations of single-cell transcriptomes of 13,289 PBMCs. Cells are

color-coded by clusters **(b)**, disease stages **(c)**, and the corresponding patient or healthy

control **(d)**. Dotted circles represented cell types with > 5% proportion of PBMCs in **(b)**

and clusters significantly enriched in patients versus control in **(c, d)**. Mono, monocyte;

NK, natural killer cells; mDC, myeloid dendritic cells; pDC, plasmacytoid dendritic

cells. **e**, Violin plots of selected marker genes (upper row) for multiple cell

subpopulations. The left column are cell subtypes identified by combination of marker

genes. **f**, Bar chart showing the proportion of immune cells from the severe, recovery

and healthy stage in each cell subpopulation.

Figure 2 | A unique monocyte subpopulation contributes to the inflammatory

storm in COVID-19 patients at severe stage. a, UMAP plot showing 4 clusters of

$CD14^+$ monocytes and 1 cluster of $CD16^+$ monocyte. Cells are color-coded by clusters.

b, Bar plot of the proportion of monocytes in cluster 9 at the severe, recovery and

healthy stages. **c**, UMAP plots showing the expressions of selected cytokines in all

monocytes clusters. **d**, Heatmap of differentially expressed genes (DEGs) in monocytes

from pairwise comparison between severe, recovery and healthy stages. **e,f**, Box plot

of the average expressions of genes involved in the signaling pathway "Acute

inflammatory response" and "Leukocyte chemotaxis" in monocytes from severe,

recovery and healthy stages. Center line, median; box limits, upper and lower quartiles;

whiskers, 1.5x interquartile range; points, outliers; **** represents P value $< 10^{-100}$,

student's t-test. **g**, Heatmap of the area under the curve (AUC) scores of expression

regulation by transcription factors (TFs) estimated using SCENIC. Shown are the top

differential TFs. **h**, UMAP plots showing the expressions of genes *ETS2*, *NFIL3* and

PHLDA2 in monocytes (top) and the AUC of the estimated regulon activity of the

corresponding TFs, indicating the degree of expression regulation of their target genes

(bottom).

Figure 3 | The monocyte-centric molecular interactions of peripheral immune cells

in COVID-19 patients at severe stage. a, Dot plot of predicted interactions between

monocytes and indicated immune cell types in the severe, recovery and healthy stages.

P values were measured by circle sizes. The expression levels of the interacted genes

were indicated by colors, scales on the right. **b**, Summary illustration depicting the

cytokine/receptor interactions between monocytes and other types of peripheral

immune cells in severe, recovery and healthy stages. Bolder lines indicated predicted

enriched ligand/receptor interactions between monocytes and other immune cell types.

Figure 4 | Enhanced humoral and cell-mediated immunity in severe COVID-19

patients. a, UMAP representations of B and plasma B cell clusters from the severe,

recovery and healthy stages. **b**, Bar plot of the proportions of plasma B cells in B cell

lineage from severe, recovery and healthy stages. **c**, UMAP representations of CD8$^+$ T

cell subtypes (left) and the distribution of cells from severe, recovery and healthy stages

in each subtype (right). **d**, Dot plot of the expression of *CCR7, PRDM1* and *MKI67* in

all CD8$^+$ T cell subtypes. **e**, Heatmap of differentially expressed genes in effector CD8$^+$

T cells from pairwise comparison between the severe, recovery and healthy stages. **f, g,**

Bar plots of GO terms enriched in effector CD8$^+$ T cells from the severe stage **(f)** or

severe and recovery stages **(g)**. **h, i,** Box plots of the average expressions of genes

involved in the signaling pathway "Leukocyte mediated cytotoxicity" and "Positive

regulation of cytokine production" in the effector CD8$^+$ T cells from severe stage,

recovery stage and healthy controls. Center line, median; box limits, upper and lower

quartiles; whiskers, 1.5x interquartile range; points, outliers; **** represents *P* value <

10^{-30}.

Extended Data Figure Legends and Supplementary Tables

Extended Data Figure 1 | Identification of single-cell subpopulations. a, UMAP

plots showing the expressions of selected marker genes in all identified cells. **b**, Bar

chart showing the percentage of cell subpopulations in different clinical stages of

patients and healthy controls.

Extended Data Figure 2 | Features of monocyte subpopulations. a-c, Bar plots of

enriched GO terms of genes highly expressed in monocytes at severe stage **(a)**, severe

and recovery stages **(b)**, and healthy controls **(c)**. **d,** Severe stage specific monocyte

regulatory network predicted by SCENIC. Transcription factors were shown in

rectangles and their target genes in circles.

Supplementary Table 1 | Baseline characteristcs and laboratory findings of

COVID-19 patients in this study.

Supplementary Table 2 | Sequencing data quality.

Supplementary Table 3 | DEGs of different stages of CD14 monocytes.

Supplementary Table 4 | GOterms of DEGs of CD14 monocytes.

Supplementary Table 5 | Gene sets of GO terms in Figure 2e and 2f.

Supplementary Table 6 | Interactions of cytokines and receptors in different stages.

Supplementary Table 7 | Drugs for targeting cytokines or their receptors.

Supplementary Table 8 | DEGs of different stages of effector CD8 T cells.

Supplementary Table 9 | GOterms of DEGs of effector CD8 T cells.

Supplementary Table 10 | Gene sets of GO terms in Figure 4h and 4i.

Reference:
450

451 1 WHO. Coronavirus disease 2019 (COVID-19) Situation Report - 72. (2020).

452 2 Wu, Z. & McGoogan, J. M. Characteristics of and Important Lessons From the
453  Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a
454  Report of 72314 Cases From the Chinese Center for Disease Control and
455  Prevention. *JAMA*, doi:10.1001/jama.2020.2648 (2020).

456 3 Mehta, P. *et al.* COVID-19: consider cytokine storm syndromes and
457  immunosuppression. *Lancet* **395**, 1033-1034, doi:10.1016/S0140-
458  6736(20)30628-0 (2020).

459 4 Zhou, Y. *et al.* Pathogenic T cells and inflammatory monocytes incite
460  inflammatory storm in severe COVID-19 patients. *National Science Review*,
461  doi:10.1093/nsr/nwaa041 (2020).

462 5 Xu, X., Han, Mingfeng, Li, Tiantian, Sun, Wei, Wang, Dongsheng, Fu, Binqing,
463  Zhou, Yonggang, Zheng, Xiaohu, Yang, Yun, Li, Xiuyong, Zhang, Xiaohua, Pan,
464  Aijun, Wei, Haiming. Effective Treatment of Severe COVID-19 Patients with
465  Tocilizumab. *[ChinaXiv:202003.00026]* (2020).

6 Zumla, A., Hui, D. S., Azhar, E. I., Memish, Z. A. & Maeurer, M. Reducing

mortality from 2019-nCoV: host-directed therapies should be an option. *Lancet*
395, e35-e36, doi:10.1016/S0140-6736(20)30305-6 (2020).

7   Chaofu Wang, J. X., Lei Zhao et al. Aveolar Macrophage Activation and
Cytokine Storm in the Pathogenesis of Severe COVID-19. *PREPRINT (Version*
*1) available at Research Square [+*https://doi.org/10.21203/rs.3.rs-19346/v1*+]*
(2020).

8   Li, G. *et al.* Coronavirus infections and immune responses. *J Med Virol* **92**, 424-
432, doi:10.1002/jmv.25685 (2020).

9   Huang, C. *et al.* Clinical features of patients infected with 2019 novel
coronavirus in Wuhan, China. *Lancet* **395**, 497-506, doi:10.1016/S0140-
6736(20)30183-5 (2020).

10   Kotch, C., Barrett, D. & Teachey, D. T. Tocilizumab for the treatment of
chimeric antigen receptor T cell-induced cytokine release syndrome. *Expert Rev*
Clin Immunol **15**, 813-822, doi:10.1080/1744666X.2019.1629904 (2019).

11   Le, R. Q. *et al.* FDA Approval Summary: Tocilizumab for Treatment of
Chimeric Antigen Receptor T Cell-Induced Severe or Life-Threatening
Cytokine Release Syndrome. *Oncologist* **23**, 943-947,
doi:10.1634/theoncologist.2018-0028 (2018).

12   Zheng, G. X. *et al.* Massively parallel digital transcriptional profiling of single
cells. *Nat Commun* **8**, 14049, doi:10.1038/ncomms14049 (2017).

13   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-
cell transcriptomic data across different conditions, technologies, and species.
Nat Biotechnol **36**, 411-420, doi:10.1038/nbt.4096 (2018).

14   Norelli, M. *et al.* Monocyte-derived IL-1 and IL-6 are differentially required for
cytokine-release syndrome and neurotoxicity due to CAR T cells. *Nat Med* **24**,
739-748, doi:10.1038/s41591-018-0036-4 (2018).

15   Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering.
Nat Methods **14**, 1083-1086, doi:10.1038/nmeth.4463 (2017).

16   Vento-Tormo, R. *et al.* Single-cell reconstruction of the early maternal-fetal
interface in humans. *Nature* **563**, 347-353, doi:10.1038/s41586-018-0698-6
(2018).

17   Guan, W. *et al.* Clinical Correlations of Transcriptional Profile in Patients
Infected With Avian Influenza H7N9 Virus. *J Infect Dis* **218**, 1238-1248,
doi:10.1093/infdis/jiy317 (2018).

18   Wang, Z. *et al.* Recovery from severe H7N9 disease is associated with diverse
response mechanisms dominated by CD8(+) T cells. *Nat Commun* **6**, 6833,
doi:10.1038/ncomms7833 (2015).

19   Thevarajan, I. *et al.* Breadth of concomitant immune responses prior to patient
recovery: a case report of non-severe COVID-19. *Nature Medicine*,
doi:10.1038/s41591-020-0819-2 (2020).

20   Braciale, T. J., Sun, J. & Kim, T. S. Regulating the adaptive immune response
to respiratory virus infection. *Nat Rev Immunol* **12**, 295-305,

doi:10.1038/nri3166 (2012).

21   Rouse, B. T. & Sehrawat, S. Immunity and immunopathology to viruses: what
decides the outcome? *Nat Rev Immunol* **10**, 514-526, doi:10.1038/nri2802
(2010).

22   Ahn, S. S. *et al.* Safety of Tocilizumab in Rheumatoid Arthritis Patients with
Resolved Hepatitis B Virus Infection: Data from Real-World Experience. *Yonsei*
Med J **59**, 452-456, doi:10.3349/ymj.2018.59.3.452 (2018).

23   Bersanelli, M. Controversies about COVID-19 and anticancer treatment with
immune checkpoint inhibitors. *Immunotherapy*, doi:10.2217/imt-2020-0067
(2020).

24   Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis
of systems-level datasets. *Nat Commun* **10**, 1523, doi:10.1038/s41467-019-
09234-6 (2019).

25   Cline, M. S. *et al.* Integration of biological networks and gene expression data
using Cytoscape. *Nat Protoc* **2**, 2366-2382, doi:10.1038/nprot.2007.324 (2007).

Figure1

a Tocilizumab / Severe COVID-19 Patients / PBMC / Single-cell RNA-seq / Data Analysis

PZ — Day 1, Day 5
PW — Day 1, Day 5, Day 7
Severe stage / Recovery stage
Droplets
Healthy PBMC from 10X — Integration

b Cell types
CD4+ T / Mono / Plasma B / CD8+ T / NK / B
UMAP2 / UMAP1

c Disease stages
UMAP2 / UMAP1

d Patients
UMAP2 / UMAP1

Cell type legend:
2, 9, 13, 17, 14 — Mono
1, 4 — CD4+ T
3, 6, 12 — CD8+ T
8 — γδT
7 — NK
5 — B
11 — Plasma B
10 — mDC
15 — pDC
16 — Platelet
18 — Progenitor
19, 20 — Unknown

Disease stages legend:
● Severe stage
● Recovery stage
● Healthy controls

Patients legend:
● Patient PW
● Patient PZ
● Healty controls

e
Markers: PTPRC, CD14, FCG3A, CD3D, IL7R, CCR7, CD8A, MKI67, TRGC1, NKG7, CD79A, CD38, CD1C, CLEC4C, PPBP, CD34

CD14+ Mono: 2, 9, 13, 17
CD16+ Mono: 14
Naive CD4+ T: 1
Effector CD4+ T: 4
Naive CD8+ T: 3
Effector CD8+ T: 6
MKI67+ CD8+ T: 12
γδT: 8
NK: 7
B: 5
Plasma B: 11
mDC: 10
pDC: 15
Platelet: 16
Progenitor: 18
Unknown: 19, 20

f Proportion of stages in each cell cluster
Percentage of cells
Clusters → 2, 9, 13, 17 (CD14+ Mono); 14 (CD16+ Mono); 1 (Naive CD4+ T); 4 (Effector CD4+ T); 3 (Naive CD8+ T); 6 (Effector CD8+ T); 12 (MKI67+ CD8+ T); 8 (γδT); 7 (NK); 5 (B); 11 (Plasma B); 10 (mDC); 15 (pDC); 16 (Platelet); 18 (Progenitor); 19, 20 (Unknown)

a

		IL1B_ADRB2			CCL20_CXCR3			FLT3LG_FLT3			CCL4L2_FFAR2			CCL4L2_VSIR			CCL4L2_PGRMC2			CCL3L1_CCR1			CCL4_CNR2			CCL4_SLC7A1			CCL3_CCR1			TNF_FFAR2			TNF_TNFRSF1A			TNF_TNFRSF1B			TNF_VSIR			IL6_IL6R		

c: healthy control   r: recovery stage   s: severe stage

−log₁₀(p-value)   0   3

log₁₀(expression)   low ▬▬▬ high

b

severe stage

recovery stage

healthy control

a

Severe stage — Plasma B, B

Recovery stage — Plasma B, B

Healthy control — Plasma B, B

UMAP2 / UMAP1

b

% of plasma B cells in B lineage

Severe, Recovery, Healthy

c

CD8⁺ T subtypes

3, 6, 12

Disease stages

UMAP1 / UMAP2

3 Naive CD8⁺ T
6 Effector CD8⁺ T
12 MKI67⁺ CD8⁺ T

Severe stage
Recovery stage
Healthy control

d

CCR7   *PRDM1*

MKI67

UMAP2 / UMAP1

high / low

e

DEGs between stages

Severe   Recovery   Healthy

568 genes

CD83   TNFRSF18
IGHG1   MAP3K8
IGHM   CLEC7A
IGKC   PIK3R6
IL1B
IL18

893 genes

FCGR2B   NCR1
FCGR3A   NCR3
KIR2DL1 FASLG
KIR2DL4 GZMB
KIR3DL1 PRF1
KIR3DL2 TNF
KLRB1   PIK3R3
KLRC1
KLRD1
KLRF1

322 genes

Expression
low / high

f

GO terms

regulation of leukocyte activation
regulation of cell activation
positive regulation of leukocyte activation
regulation of lymphocyte activation

-log₁₀(p-value)

g

GO terms

leukocyte mediated cytotoxicity
positive regulation of cytokine production
regulation of leukocyte mediated cytotoxicity
regulation of cytokine production

-log₁₀(p-value)

h

Leukocyte
mediated cytotoxicity

Average gene expression

****   ****

Severe   Recovery   Healthy

i

Positive regulation
of cytokine production

Average gene expression

****   ****

Severe   Recovery   Healthy