Repeated parallel losses of inflexed stamens in Moraceae: phylogenomics and generic

revision of the tribe Moreae and the reinstatement of the tribe Olmedieae (Moraceae)

Elliot M. Gardner[1,2,3,*], Mira Garner[1,4], Robyn Cowan[5], Steven Dodsworth[5,6], Niroshini

Epitawalage[5], Deby Arifiani[7], S. Sahromi[8], William J. Baker[5], Felix Forest[5], Olivier

Maurin[5], Nyree J.C. Zerega[9,10], Alexandre Monro[5], and Andrew L. Hipp[1,11]

[1] The Morton Arboretum, 4100 IL-53, Lisle, Illinois 60532, USA

[2] Case Western Reserve University, Department of Biology, 2080 Adelbert Road,

Cleveland, Ohio 44106, USA (current affiliation)

[3] Singapore Botanic Gardens, National Parks Board, 1 Cluny Road, 259569, Singapore

(current affiliation)

[4] University of British Columbia, Faculty of Forestry, 2424 Main Mall, Vancouver, British

Columbia, V6T 1Z4, Canada (current affiliation)

[5] Royal Botanic Gardens, Kew, Richmond, TW9 3AE, United Kingdom

[6] School of Life Sciences, University of Bedfordshire, Luton LU1 3JU, United Kingdom

(current affiliation)

[7] Herbarium Bogoriense, Research Center for Biology, Indonesian Institute of Sciences,

Cibinong, Jawa Barat, Indonesia

[8] Center for Plant Conservation Botanic Gardens, Indonesian Institute of Sciences, Bogor,

Jawa Barat, Indonesia

[9] Chicago Botanic Garden, Plant Science and Conservation, 1000 Lake Cook Road,

Glencoe, IL, 60022, USA

[10] Northwestern University, Plant Biology and Conservation Program, 2205 Tech Dr.,

Evanston, IL, 60208, USA

[11] The Field Museum, 1400 S Lake Shore Dr., Chicago, IL 60605, USA

[a] For correspondence, email elliot.gardner@case.edu

Abstract

We present a densely-sampled phylogenomic study of the mulberry tribe (Moreae,

Moraceae), an economically important clade with a global distribution, revealing multiple

losses of inflexed stamens, a character traditionally used to circumscribe Moreae. Inflexed

stamens facilitate ballistic pollen release and are associated with wind pollination, and the

results presented here suggest that losses of this character state may have evolved

repeatedly in Moraceae. Neither Moreae nor several of its major genera (*Morus*, *Streblus*,

Trophis) were found to be monophyletic. A revised system for a monophyletic Moreae is

presented, including the reinstatement of the genera *Ampalis, Maillardia, Taxotrophis,* and

Paratrophis, and the recognition of the new genus *Afromorus*, based on *Morus* subgenus

Afromorus. *Pseudostreblus* is reinstated and transferred to the Parartocarpeae, and

Sloetiopsis is reinstated and transferred to the Dorstenieae. The tribe Olmediae is reinstated,

replacing the Castilleae, owing to the reinstatement of the type genus *Olmedia,* and its

exclusion from Moreae. *Streblus* s.s. is excluded from Moreae and transferred to the

Olmediae, which is characterized primarily by involucrate inflorescences without regard to

stamen position. Eight new combinations are made.

Keywords: Moraceae, mulberry family; Moreae, Olmedieae, Castilleae, Parartocarpeae,

Afromorus, Ampalis, Bagassa, Maillardia, Milicia, Morus, Olmedia; Pachytrophe,

Paratrophis, Pseudostreblus, Sloetiopsis; Sorocea, Streblus, Taxotrophis, Trophis.

Introduction

The preservation of plesiomorphic (ancestral) characters can result in species that

are similar in appearance but distantly related, connected only by a remote common

ancestor. The mulberry family (Moraceae Gaudich., seven tribes, ca. 39 genera and 1,200

species) illustrates this principle well. Inflexed stamens in bud—an adaptation to wind

pollination that allows explosive pollen dispersal when flowers open—were traditionally

used to define a tribe of the family, the Moreae (mulberries and their allies). Yet

phylogenetic analyses have revealed that inflexed stamens, an ancestral feature of both

Moraceae and their sister family Urticaceae (nettles), have been lost repeatedly (Clement &

Weiblen, 2009). Thus, for example, Cecropiaceae C.C. Berg, traditionally distinguished

from the nettle family by the absence of inflexed stamens, is in fact embedded within the

Urticaceae Juss. (Berg, 1978; Clement & Weiblen, 2009). Likewise, while mulberries

(*Morus* L.), paper mulberries (*Broussonetia* L'Hér. ex Vent.), and osage oranges (*Maclura*

Nutt.) were once treated as tribe Moreae Gaudich. on account of their inflexed stamens,

phylogenetic analyses reveal them to belong to three distinct clades, each containing an

assemblage of genera with and without inflexed stamens (Clement & Weiblen, 2009;

Zerega & Gardner, 2019).

This study focuses on the tribe Moreae, a clade of six genera and an estimated 66

species (Clement & Weiblen, 2009). This widespread group of plants contains species of

ecological and cultural importance, as well as the economically important white mulberry

(*Morus alba* L.), whose leaves sustain *Bombyx mori* L. (Bombycidae), the invertebrate

proletariat of the silk industry.

Morphological basis for higher classification in Moraceae

By the end of the nineteenth century, Engler (1889) had circumscribed a Moraceae

that is quite similar to the modern concept of the family, with two subfamilies: the

Moroideae, comprising tribes Dorstenieae Gaudich., Broussonetieae Gaudich., Fatouae

Engler, Moreae, and Strebleae Bureau, characterized by stamens inflexed in bud (broadly

construed, including *Dorstenia* where they straighten gradually rather than spring outward

suddenly); and the Artocarpoideae, composed of tribes Brosimae Trécul, Euartocarpeae

Trécul, Ficeae Dumort., and Olmedieae Trécul and characterized by stamens straight in

bud. The two most influential scholars of Moraceae classification in the 20th century were

E.J.H. Corner and C.C. Berg. Corner considered inflorescence architecture to be the most

important character for higher-rank taxonomy within the family and inflexed stamens

consequently of secondary importance(Corner, 1962). Berg by contrast questioned the

utility of inflorescence architecture and took into account a variety of characters, especially

the presence of inflexed stamens (Berg, 1977b, 2001).

The taxonomic history of the family has leaned heavily on this stamen character.

Engler's Moreae (1899), unchanged in substance from Bureau's (1873), contained six

genera, *Ampalis* Bojer*, Pachytrophe* Bureau*, Paratrophis* Blume*, Pseudomoru*s Bureau,

and *Morus*, all with inflexed stamens that spring out suddenly at anthesis (Table 1).

Corner's expanded Moreae consisted of seven genera with either straight or inflexed

stamens, but with pistillate inflorescences never condensed into a head: *Fatoua* Gaudich,

Morus, *Sorocea* A. St.-Hil. (apparently including *Paraclarisia* Ducke), *Clarisia* Ruiz &

Pav., *Ampalis*, *Pachytrophe*, and *Streblus* Lour. (including *Bleekrodea* Blume*, Paratrophis,*

Pseudomorus, *Taxotrophis* Blume, *Sloetiopsis* Engl. and *Neosloetiopsis* Engl.). Corner

himself, however, found the diversity of his Moreae unsatisfactory, noting that "[t]oo many

genera on insufficient and invalid grounds trouble this small tribe" (Corner, 1962). Perhaps

in response, Berg's Moreae comprised all of the genera with inflexed stamens and none

without, including *Broussonetia* and *Maclura* but excluding *Sorocea* and *Clarisia*,

providing a simple character with which to delimit the tribe(Berg, 2001; Berg & al., 2006).

Recent phylogenetic work has supported a third approach, with the Moreae

comprising six genera, including genera with both straight (*Sorocea*, *Bagassa* Aubl.) and

inflexed (*Broussonetia*, *Maclura*) stamens (Clement & Weiblen, 2009). These studies also

suggest that the character state of inflexed stamens is plesiomorphic and is the ancestral

state for both Moraceae and Urticaceae (Datwyler & Weiblen, 2004; Zerega & al., 2005;

Clement & Weiblen, 2009). Inflexed stamens, which are precursors to ballistic pollen

release, are associated with wind pollination (Bawa & Crisp, 1980; Berg, 2001), while the

loss of inflexed stamens is associated with animal pollination. Although dominant in the

Moreae, inflexed stamens also occur in two other tribes: Maclureae W.L. Clement & G.

Weiblen and Dorstenieae. Genera lacking inflexed stamens occur in all seven tribes of

Moraceae (Clement & Weiblen, 2009; Zerega & Gardner, 2019).

Taxonomic summary of Moreae

Following Clement & Weiblen (2009), the Moreae comprise six genera and an

estimated 66 species: *Bagassa* (1), *Milicia* Sim (2), *Morus* (16), *Sorocea* (16), *Streblus*

(23), and *Trophis* P. Browne. (8) (Berg, 1977a, 2001; Berg & al., 2006; Clement &

Weiblen, 2009; Filho & al., 2009; Machado & al., 2013; Santos & Neto, 2015). Here, we

present an overview of these genera.

Morus L., the true mulberries, is characterized by leaves with crenate margins and
trinerved bases, stamens inflexed in bud, and many-flowered pistillate spikes whose four-
parted perianths become fleshy in fruit, the aggregations superficially resembling a
blackberry. *Morus* comprises approximately 16 species whose delimitation requires further
research. There are three subgenera: *Morus* (ca. 14 species) which is found in temperate to
tropical Asia and from North America to Mexico; *Gomphomorus* Leroy, a single species
restricted to tropical South America; and *Afromorus* (Bureau ex Leroy), a single species
restricted to tropical Africa. Previous phylogenetic work has suggested that these three
subgenera may not form a monophyletic clade (Nepal, 2012). *Milicia* (2 spp., Africa) has
inflorescences which somewhat resemble those of *Morus*, but the leaves of *Milicia*, with
entire margins and pinnate venation, prevent any confusion of the two genera.

Streblus Lour., with 23 species from India to Southeast Asia and Oceania, is
morphologically heterogeneous, but its species are all characterized by stamens inflexed in
bud and pistillate flowers with more or less free tepals that enclose the fruit loosely or not
at all. Initially described by Loureiro based on the widespread *S. asper* Lour.—notable for
its discoid-capitate staminate inflorescences with the rudiments of an involucre—*Streblus*
was broadened by Corner (1962), bringing in as sections *Taxotrophis*, *Phyllochlamys*
Bureau, *Paratrophis* (including *Pseudomorus*), *Pseudostreblus* Bureau, *Bleekrodea*, and
Sloetia Teijsm. & Binn. (apparently including *Sloetiopsis* but without making any
combinations). None of these have discoid-capitate inflorescences; they mostly have
spicate staminate inflorescences, and the latter two can have bisexual inflorescences.
Corner viewed these sections as fragments of an ancient lineage preserving ancestral
characters (Corner, 1975). Following additional work by Corner (1970, 1975), Berg
included the genera *Ampalis* and *Pachytrophe* as section *Ampalis* (Bojer) C.C. Berg,
subsumed *Sloetiopsis* and section *Taxotrophis* into section *Streblus*, and excluded section
Bleekrodea, reinstating it as a genus (Berg, 1988). Berg later reinstated section *Taxotrophis*,
whose species are unique in bearing axillary spines (Berg, 2005; Berg & al., 2006). In
2009, Clement and Weiblen reinstated *Sloetia* at generic rank and transferred it and
Bleekrodea to Dorstenieae based on phylogenetic evidence (Clement & Weiblen, 2009).

Trophis P. Browne is characterized by stamens inflexed in bud, spicate staminate

inflorescences, and tubular pistillate perianths, becoming fleshy and enclosing the fruit,

except for the monotypic section *Olmedia* (Ruiz & Pav.) C.C. Berg (*T. caucana*), which has

discoid-capitate staminate inflorescences with an involucre. *Trophis* (regrettably conserved

over Linnaeus's *Bucephalon* L.) as recognized by Berg (1988, 2001) had five sections:

Trophis, restricted to Latin America; *Calpidochlamys* (Diels) Corner (previously included

in *Paratrophis* and *Uromorus*), restricted to Southeast Asia (Corner, 1962); *Maillardia*

(Frapp. ex Duch.) C.C. Berg, restricted to Africa; *Malaisia* (Blanco) C.C. Berg, restricted to

Southeast Asia; and *Olmedia*, restricted to Latin America. The sinking of *Olmedia* into

Trophis by Berg (1988), requiring the re-typification of the tribe Olmedieae Trécul, which

became Castilleae C.C. Berg. Recently, *Malaisia* Blanco was reinstated as a genus and

transferred to Dorstenieae based on phylogenetic evidence (Clement & Weiblen, 2009),

reducing the current number of sections to four.

Two Neotropical genera included in Moreae by Clement and Weiblen (2009) but

not by Berg (2001) have straight stamens. While the monotypic *Bagassa* Aubl., with its

long staminate catkins, is wind pollinated (Bawa & Crisp, 1980), evidence suggests that

Sorocea A. St.-Hil. (19 spp.), which produces racemose staminate inflorescences that do

not have as many flowers as those of *Bagassa*, likely contains both wind- and insect-

pollinated species (Zapata & Arroyo, 1978; Bawa & al., 1985; Lewis, 1986). The pistillate

perianths are subtended by pluricellular hairs, believed to serve as a substrate for a fungus

which, in turn attracts pollinators (Berg, 2001). The staminate flowers of *Sorocea affinis*

Helmsl. are apparently fragrant (fide *S. Zona 778*, 21 Nov. 1997, FTBG♂♀), suggesting

insect pollination for that species. If insect pollination is confirmed in *Sorocea*, it would

likely represent another transition from ancestral wind to derived insect pollination in

Moraceae.

Disagreement between the two principal recent monographers of the Moraceae,

Corner (1962) and Berg (2001, 2005, 2006), over the delimitation of the Moreae has been

compounded by a series of phylogenies based on the analysis of molecular (DNA

sequence) data that have redefined genera and their relationships. This has resulted in a

poorly delimited tribe, both with respect to diagnostic morphological characters, the genera

it comprises, and the rank of several taxa (see above). Ensuring that the tribe and its genera

are monophyletic will result in a classification that better reflects evolutionary history,

providing a framework for answering broader scientific questions. Our aim, therefore, was

to generate a comprehensive phylogeny of the Moreae (Fig. 1) and its near allies by

sampling all of the potential genera and species in the tribe sensu Berg, and sensu Clement

& Weiblen (2009) and use this to redelimit the tribe and its genera. We aimed to do so

through increased taxon and genome sampling compared to previous studies, including a

nearly comprehensive sample of the taxa in Moreae (56/66 species) and the allied tribes

Artocarpeae (76/83 taxa) and Maclureae (12/12). Our data set combines phylogenomic data

generated using two largely non-overlapping sets of enrichment baits, one developed

specifically for the Moraceae (Gardner et al., 2016), the other for the whole of the

Angiosperms (Johnson & al., 2019), allowing us to explore the possibilities and challenges

of combining samples based on largely non-overlapping loci. The resulting generic revision

lays the groundwork for species-level revisionary work and provides clarity to this

economically important clade.

195   We also set out to reconstruct the evolutionary history of inflexed stamens within

Moraceae using ancestral state reconstruction. The loss of inflexed stamens in

Castilleae+Ficeae and Artocarpeae have been associated with transitions from wind to

animal pollination (Momose & al., 1998; Sakai & al., 2000; Datwyler & Weiblen, 2004;

Gardner & al., 2018). A more complete picture of evolutionary transitions between inflexed

and straight stamens may help focus further research on transitions in pollination biology

with Moraceae.

Materials and Methods

204   We used target enrichment sequencing (HybSeq) (Weitemier & al., 2014) to capture

333 genes previously developed for phylogenetic work in Moraceae (the "Moraceae333")

(Gardner & al., 2016; Johnson & al., 2016). This method allows for efficient capture of

hundreds of loci and is suitable for both fresh material and degraded DNA from herbarium

material (Villaverde & al., 2018; Brewer & al., 2019), which comprises much of the

material employed in this study.

Taxon sampling, *library preparation, and sequencing*

We sampled 56 out of 66 species (and all genera) in Moreae as well as select outgroup

taxa from Artocarpeae using DNA from leaf tissue preserved on silica gel or—in most

cases—samples from herbarium specimens (Table S1). In either case, DNA was extracted

using a modified CTAB method, usually with increased incubation times to maximize yield

from herbarium tissue(Doyle & Doyle, 1987; Hale & al., 2020). Samples were quantified

using a Qubit fluorometer (Invitrogen, Life Technologies, California, USA), and herbarium

samples were run on a gel to test for degradation. For most samples 200 ng of DNA was

used for library preparation; for some low-yield samples, as little as 50 ng was used, and for

very degraded samples, as much input as possible was used, up to 400 ng. Undegraded

DNA was fragmented either using NEB DNA Fragmentase (New England Biolabs,

Ipswich, Massachusetts, USA) or on a Covaris M220 (Covaris, Wobum, Massachusetts,

USA). DNA samples with an average fragment size less than 500 bp were not fragmented

at all, and partially-degraded samples with an average fragment size of over 500 bp were

fragmented on the Covaris M220. TruSeq-style library preparation was carried out using

either the KAPA Hyper Prep kit (Kapa Biosystems, Wilmington, MA) or the NEB DNA

Ultra 2 kit following the manufacturer's protocols, except that end repair, A-tailing, and

adapter ligation were carried out in reduced-volume reactions (0.25x for KAPA and 0.5x

for NEB) to reduce costs. Final products were quantified on the Qubit and combined in

equal molecular weights into pools of 16–20 samples. The pools totaled 1,200 µg each if

enough library preparation was available. Pools were hybridized for 16–24 hours to custom

Moraceae probes (Gardner et al, 2016) using a MYBaits kit (Arbor Biosciences, Ann

Arbor, Michigan, USA) following the manufacturer's protocol, except that the probes were

diluted 1:1 with nuclease-free water. Hybridization products were reamplified using KAPA

Hot Start PCR reagents following the MYbaits protocol, quantified on the Qubit, and

quality-checked on an Agilent BioAnalyzer (Agilent Technologies, Palo Alto, California,

USA). Samples with adapter dimer peaks were cleaned using 0.7x SPRI beads and re-run

on the Qubit and BioAnalyzer. An initial sequencing run took place on a MiSeq 2 x 300bp

run (v3) (Illumina, San Diego, California, USA) at the Field Museum of Natural History,

and then additional samples were sequenced on a HiSeq 4000 2 x 150bp run at the

Northwestern University Genomics Core.

We also included 47 Moraceae and Urticaceae samples enriched for the

Angiosperm353 probes and sequenced as part of the Plant and Fungal Trees of Life project

(PAFTOL, RBG Kew; https://www.kew.org/science/our-science/projects/plant-and-fungal-

trees-of-life; Table S1). Sample preparation and sequencing followed Johnson et al. (2019).

The PAFTOL samples were enriched with a universal probe set developed for angiosperms

(the "Angiosperms353").

Finally, we used samples sequenced for previous phylogenetics projects in *Artocarpus*

J.R. Forst. & G. Forst. and Parartocarpeae Zerega & E.M. Gardner to complete our

sampling (Johnson & al., 2016; Gardner, 2017; Kates & al., 2018; Zerega & Gardner,

2019). The final dataset contained 247 samples.

Assembly of reads

We trimmed reads using Trimmomatic (ILLUMINACLIP: TruSeq3-PE.fa:2:30:10

HEADCROP:3 LEADING:30 TRAILING:25 SLIDINGWINDOW:4:25 MINLEN:20)

(Bolger & al., 2014) and assembled them with HybPiper, which produces gene-by-gene,

reference-guided, *de novo* assemblies (Johnson & al., 2016). For the samples enriched with

the Angiosperms353 baits, we used the reference described by Johnson et al. (2019), and

for the samples enriched with the Moraceae333 baits, we used the reference described in

Zerega & Gardner (2019). To increase overlap between the two data sets beyond the 5

genes they inherently have in common, we used HybPiper to assemble the

Angiosperms353-enriched reads using the Moraceae333 targets and vice-versa, in order to

capture any additional genes found in off-target reads. The exception was most of the

Artocarpus data set, which was previously assembled for another study using only the

Moraceae333 targets. We used the HybPiper script "intronerate.py" to build "supercontig"

sequences for each gene, consisting of exons as well as any assembled flanking non-coding

sequences (intronic or intergenic). To these new assemblies, we added 76 Moraceae333

assemblies from Gardner (2017) to complete sampling for the Artocarpeae.

Main phylogenetic analyses

The full data set was analyzed using the "exon" sequences to ensure good alignment

across the entire family. To maximize phylogenetic resolution within Moreae, a smaller

data set consisting only of Moreae taxa and a single outgroup taxon (*Artocarpus*

heterophyllus) was analyzed using the "supercontig" sequences but following the same

methodology. For each of the 686 genes assembled, we discarded sequences shorter than

25% of the average length for the gene and discarded genes containing sequences for less

than 30 samples. We aligned each gene with MAFFT (Katoh & Standley, 2013)and used

Trimal (Capella-Gutiérrez & al., 2009) to discard sequences with an average pairwise

identity of less than 0.5 to all other sequences in the alignment (indicative of poor a poor

quality sequence that could not be properly aligned) as well as columns containing more

than 75% gaps. We used RAxML 8.2.4 (Stamatakis, 2014) to estimate a maximum-

likelihood tree for a partitioned supermatrix of all genes as well as for each gene

individually (GTRCAT model, 200 bootstrap replicates). We then used ASTRAL-III

(Zhang & al., 2017), a summary-coalescent method, to estimate a species tree from all gene

trees, estimating node support using both bootstrap (160 replicates, resampling across and

within genes) and local posterior probability (normalized quartet scores, representing gene

tree concordance). Finally, we used ASTRAL-III to test whether, for each node, the null

hypothesis of a polytomy could be rejected (Sayyari & Mirarab, 2018). Alignment,

trimming, and estimation of gene trees were parallelized using GNU Parallel (Tange,

2018).

Whole chloroplast phylogenetic tree

We also built a whole-chloroplast phylogenetic tree as follows. Rather than

assembling full-length genomes, which can be extremely slow to align, we assembled and

aligned the genomes in sections, dramatically speeding up the process. We created

HybPiper targets using the chloroplast genome of *Morus indica* L. (NCBI RefSeq accession

no. NC_008359.1) and the associated gene annotations. Each target consisted of a gene

feature concatenated with any internal or subsequent non-coding sequence, terminating one

base before the next gene feature began. These intervals were generated by manually

editing the NCBI gff3 file in Excel (Microsoft Corp., Redmond, Washington, USA) and

then extracted using BedTools (Quinlan & Hall, 2010). Assemblies for all Moraceae

samples and *Boehmeria nivea* (L.) Gaudich. were carried out in HybPiper using a coverage

cutoff of 2 and otherwise with default parameters. Within each target, sequencing less than

25% of the average length were discarded, and after alignment with MAFFT, sequences

with an average pairwise identity of less than 0.7 were discarded (the higher cutoff

reflecting the conserved nature of chloroplast DNA). Alignments were then concatenated

into a supermatrix, and samples with more than 50% undetermined characters were

discarded. Alignments were then rebuilt, filtered, and concatenated, and columns in the

final supermatrix with more than 75% missing characters were discarded. A maximum-

likelihood tree was generated using RAxML 8.2.4 under the GTRCAT model, with 200

rapid bootstrap replicates.

Phylogenetic network analysis

To further investigate relationships within the *Paratrophis* clade, including the proper

placement of *Morus insignis* and *Trophis philippinensis* (Bureau) Corner, we constructed

phylogenetic networks based on two reduced 8-taxon datasets ("exon" and "supercontig")

consisting of those taxa plus *Streblus anthropophagorum* (Seem.) Corner, *S. glaber* (Merr.)

Corner, *S. glaber* subsp. *australianus* (C.T. White) C.C. Berg, and *S. heterophyllus* (Blume)

Corner, with *S. mauritianus* (Jacq.) Blume as the outgroup. Alignment preparation followed

the workflow outline above except that only genes with sequences for all eight taxa were

retained. Rooted gene trees were used to infer the network in PhyloNet using the

"InferNetwork_MPL" command, allowing a maximum of four hybridization events and

collapsing gene tree nodes with less than 30% bootstrap support (Yu & Nakhleh, 2015;

Wen & al., 2018) (Wen et al., 2018; Yu and Nakhleh, 2015); AIC was scored for the best

five networks from 10 runs, using the number of branch lengths and hybridization events

calculated as the number of parameters (Kamneva & al., 2017).

ITS and rbcL phylogenetic trees

While we did not have our own sequences for *Streblus tonkinensis* (Eberh. & Dubard)

Corner, *S. ascendens* Corner, *S. banksii* (Cheeseman) C.J. Webb, and *S. smithii*

(Cheeseman) Corner, either *ITS* or *rbcL* sequences existed in NCBI GenBank for these taxa

(Fig. S2). We were not able to examine the underlying specimens ourselves, but we

considered the chances of misidentification low because those species are morphologically

and/or geographically distinctive. We constructed *ITS* and *rbcL* data sets from our own

samples using HybPiper, using *Streblus* sequences for these loci (obtained from GenBank)

as targets and reducing the coverage cutoff at 2 to recover the loci from off-target reads. To

these sequences, we added GenBank sequences of *Streblus* taxa as well as *Trophis caucana*

(Pittier) C.C. Berg. For each locus, we aligned sequences with MAFFT, discarded short

sequences (<490 bp), trimmed alignments to remove columns with over 75% gaps, and

built maximum-likelihood trees using RAxML (GTRGAMMA, 1000 rapid bootstraps).

Divergence time estimation

The supermatrix maximum-likelihood tree was time calibrated using ape v 5.3

(Paradis & Schliep, 2019) in R v 3.5.1 (Team, 2019). First, the tree was pruned of duplicate

taxa and non-Moraceae outgroup taxa, and edge lengths (in substitutions per site) were

multiplied by the number of sites in the alignment (to convert them to substitutions). The

following stem nodes were constrained with minimum ages (in million years, Ma) based on

fossil data, following(Zhang & al., 2019): *Ficus,* 56 Ma; *Broussonetia*, 33.9 Ma; *Morus*

(subg. *Morus*, based on U.S.S.R. locality of the fossil), 33.9 Ma; and *Artocarpus*, 64 Ma.

The crown node of Moraceae was constrained to a minimum age of 73.2 Ma and a

maximum age of 84.7 Ma.(Zhang & al., 2019), which had a more extensive outgroup

sampling that we have here. The tree was then time-calibrated using the chronos function

under two different models ("relaxed" and "correlated") (Kim & Sanderson, 2008; Paradis,

2013). The smoothing parameter (λ) was chosen using the cross-validation method in the

chronopl function (testing $\lambda = 0$ and 0_{-5} through 10_{15}), selecting the value of λ that

minimized the cross-validation statistic (Sanderson, 2002). The resulting trees were

visualized using the densiTree function in phangorn 2.4.0 (Schliep, 2011), and the time

calibrated tree representing the central tendency of these analyses was selected for use in all

further analyses, also taking into account the results of past family-wide studies. As the

penalized likelihood approach used does not integrate over model uncertainty or

uncertainty in calibration placement and timing, confidence intervals on node ages are not

provided in this study. A geologic timescale based on the strat2012 dataset added to tree

figures using PHYLOCH v 1.5-3 (Heibl, 2008).

Ancestral state reconstruction for inflexed stamens

All taxa were coded for presence (1) or absence (0) of inflexed stamens in bud.

Taxa such as *Dorsteniae* with stamens that are inflexed in bud but gradually straighten were

coded as 0. We reconstructed ancestral character states on the entire phylogeny by

stochastic mapping using the make.simmap function in the phytools v 0.6-99 (Revell,

2012). We also tested for trait-associated shifts in diversification rates using BAMM v

2.5.0 (Rabosky & al., 2013, 2014a; Rabosky, 2014), specifying the amount of missing taxa

per clade to account for the high proportion of missing taxa in Ficeae, Castilleae, and

Dorstenieae. Parameters were optimized using BAMMtools v 2.1.7 (Rabosky & al.,

2014b). We also tested for trait-dependent diversification using diversitree v 0.9-13

(FitzJohn, 2012). We evaluated BiSSE and trait-independent models on the entire tree and

on a pruned tree consisting only of the densely-sampled Maclureae+Moreae+Artocarpeae

clade.

Reads from have been deposited in Genbank (Non-PAFTOL samples: BioProject

PRJNA322184; PAFTOL samples: BioProject PRJEB35285, subprojects PRJEB37667 for

Moraceae and PRJEB37665 for Urticaceae). Tree files and scripts outlining the

phylogenetic analyses have been deposited in the Dryad Data Repository (###TBA###).

Results

The final data set contained 247 samples, 47 enriched with the Angiosperms353

baits, 196 enriched with the Moraceae333 baits, and 4 extracted from whole genomes) and

619 genes (286 Angiosperms353 and all of the Moraceae333) (Table S1). The "exon"

supermatrix (all taxa) contained 613,126 characters, and the "supercontig" supermatrix

(Moreae and select outgroup taxa only) contained 799,926 characters. The chloroplast data

set contained 113 loci.

On average, we assembled 39 Moraceae333 genes from the Angiosperms353-

enriched samples and 20 Angiosperms353 genes from the Moraceae333-enriched samples,

and the average pairwise overlap in assembled loci was 210 (Tables S1, S2). Nineteen taxa

were represented in both sets of samples, and 15 of these were always monophyletic (Figs.

2, 3). Twelve resolved as sister pairs: *Antiaropsis decipiens* (31 loci overlapping), *Bagassa*

guianensis Aubl. (38), *Batocarpus amazonicus* (Ducke) Fosberg (22), *Batocarpus*

orinocensis H. Karst. (34), *Brosimum alicastrum* Sw. (42), *Clarisia racemosa* Ruiz & Pav.

(59), *Maclura africana* (Bureau) Corner (213), *Milicia excelsa* (Welw.) C.C. Berg (45),

Streblus asper (Retz.) Lour. (75), *Streblus mauritianus* (Jacq.) Blume (40), *Streblus*

usambarensis (Engl.) C.C. Berg (43), and *Trophis caucana* (32). Three more represented

by more than two samples always resolved as a clade: *Sorocea bonplandii* (Baillon) W.C.

Burger, Lanj. & de Boer (3 samples; 19–35 loci overlapping), *Maclura tinctoria* (L.) D.

Don ex Steud. (4 samples; 37–43 loci overlapping), and *Malaisia scandens* (Lour.) Planch.

(3 samples; 27–43 loci overlapping). Three additional taxa resolved as sister pairs in the

ASTRAL analysis but as a grade in the supermatrix analysis: *Maclura cochinchinensis*

(Lour.) Corner (28 loci overlapping), *Streblus heterophyllus* (Blume) Corner (50), and

Trophis montana (Leandri) C.C. Berg (56). Only one species, *Utsetela gabonensis* Pellegr.

(18 loci overlapping) was never monophyletic, always forming a grade with *U. neglecta*

Jongkind.

The supermatrix and species trees were broadly concordant except within

Artocarpus, where the species tree was much more consistent with previous analyses within

that genus (Gardner et al., in review) (Figs. 2, 3). Otherwise, most differences were at

shallow phylogenetic depths, such as relationships within *Streblus* section *Paratrophis*. For

both data sets, the polytomy hypothesis was rejected ($P < 0.05$) for Moraceae and all tribe

and genus-level clades (following the revised classification presented below) (Figs. 2, 3).

Of the genera in Moreae sensu Clement & Weiblen, only *Milicia* and *Sorocea* were

monophyletic; *Morus*, *Streblus*, and *Trophis* were not; *Bagassa* is monotypic and resolved

as sister to Sorocea, with which it shares straight stamens. Four Moreae species resolved

with other tribes: *Trophis caucana* was nested within Castilleae, *Streblus asper* (Retz.)

Lour. was sister to Castilleae, *Streblus indicus* (Bureau) Corner was sister to

Parartocarpeae, and *Streblus usambarensis* was nested within Dorstenieae. Moreae was

otherwise monophyletic, comprising the following subclades: (1) *Streblus* section

Taxotrophis, sister to all other Moreae; (2) (a) *Trophis* section *Maillardia*, (b) *Milicia* +

Morus subgenus *Afromorus*, (c) *Streblus* section *Ampalis*, (d) *Streblus* section *Paratrophis*

in part + *Morus* subgenus *Gomphomorus*, (e) *Streblus* section *Paratrophis* in part + *Trophis*

section *Calpidochlamys*; (3) *Bagassa* + *Sorocea*; (4) *Trophis* sections *Trophis* and

Echinocarpa C.C. Berg + *Morus*. These are roughly geographic clades: (1) Southeast Asia;

(2) (a) Madagascar, (b) Southeast Africa, (c) Madagascar, (d) Pacific + South America, (e)

Southeast Asia + Pacific; (3) South America; (4) South America.

While *Trophis philippinensis* was inside the *Paratrophis* clade in all analyses, the

position of *Morus insignis* was not stable. In the "exon" analyses, it was always in

Paratrophis (Fig. 2), although the polotomy hypothesis could not be rejected for its

position as sister to *S. anthropophagorum*+*S. heterophyllus*. In the "supercontig" and

chloroplast analyses, it was sister to *Paratrophis*, and the polotomy test was rejected for

that node (Figs. 3, S1). In the five phylogenetic networks reported (Fig. 4, Table S3), *M.*

insignis was inside *Paratrophis* in three of them, had a hybrid origin involving *Paratrophis*

in one, and was only unambiguously sister to *Paratrophis* in one network, although even in

that one it was involved in a hybridization involving *Paratrophis*.

The chloroplast phylogenetic tree (Fig. S1) was generally in agreement with the

nuclear phylogenetic trees, with three notable differences. *Streblus indicus* and the

Parartocarpeae formed a grade paraphyletic to Dorstenieae, Castilleae, and Ficeae, rather

than a clade. *Streblus zeylanicus* (Thwaites) Kurz was sister to *S. taxoides* (B. Heyne ex

Roth) Kurz (instead of sister to all of section *Taxotrophis*), although with low support

(62%), and *S. glaber* subsp. *australianus* was not sister to subsp. *glaber*, although its

placement was not strongly supported (80%). Finally, *Morus insignis* was sister to

Paratrophis, agreeing with the "supercontig" analysis but not the "exon" analysis.

The ITS and *rbc*L phylogenetic trees based on few characters, were not very well

resolved, with only few nodes attaining 100% bootstrap support (Fig. S2). Nevertheless,

with the exception of a few stray taxa (one *Dorstenia* sample in ITS and one *Streblus*

indicus samples in *rbc*L), both phylogenetic trees reconstructed monophyletic tribes with at

least moderate support. In the ITS phylogenetic tree, *Streblus smithii* and *S. banksii* were

both in a well-supported clade comprising section *Paratrophis*, and *S. tonkinensis* was part

of the Castilleae clade, which also contained *S. asper* and *Trophis caucana*. In the *rbc*L

phylogenetic tree, the two *Streblus ascendens* samples, the four samples of the

undetermined *Streblus* from Papua New Guinea, and *S. smithii* were part of a well-

supported clade comprising section *Paratrophis*.

Time-calibration and ancestral state reconstruction

458   The cross-validation criterion was minimized by a smoothing parameter (λ) value of

10_6. The tree calibrated under the correlated model (logLik = -17; p-logLik = -17; ΦIC =

1108) was less sensitive to changes in λ and generally had younger ages than the tree

calibrated under the relaxed model (logLik = -56.7; p-logLik = -53129706, ΦIC =

4209013), the latter of which had perhaps implausibly long terminal branches (Table 2, Fig.

S3). The crown age of Moreae was Paleocene (59.1 Ma) under the correlated model and

late Cretaceous (75.4 Ma) under the relaxed model. Because the correlated tree was more

consistent with past family-wide studies (Zerega & al., 2005; Zhang & al., 2019), we used

that tree for all further analyses (Fig. 5). The BAMM analysis found a single credible rate

shift, not surprisingly at the crown node of *Ficus* (Fig. S4); that rate shift was—also not

surprisingly—associated with a loss of inflexed stamens, which are never found in *Ficus* (*P*

= 0.035); no rate shifts were found within Moreae. Ancestral reconstruction on the entire

tree found that ancestral Moraceae had stamens inflexed in bud; these were lost nine times

(in Ficeae, Olmedieae [=Castilleae], Parartocarpeae, twice in Dorstenieae, *Maclura* section

Cudrania (Trécul) Corner, Artocarpeae, *Bagassa,* and *Sorocea*) and regained once (in

Trophis caucaua) (Fig. 6). Model testing on both the whole tree and on the

Moreae+Artocarpeae+Maclureae subtree indicated that a BiSSE model was not

substantially better than a trait-independent model (Moreae only: AIC for BiSSE = 1110.4;

trait-independent = 1112.8; ΔAIC = 2.4; whole family: AIC for BiSSE = 1488.3; trait-

independent = 1490.4; ΔAIC = 2.1). In any event, the reconstructions were identical (Figs.

6).

Discussion

Sequencing and combination of data sets

This study was materially improved by our ability to combine samples enriched

with two largely non-overlapping bait sets. Despite minimal by-design overlap, we were

able to assemble many overlapping loci for samples with moderate to deep coverage (Table

S2), adding taxa that would not otherwise have been included in the study and replicating

taxa to confirm unexpected phylogenetic placement (e.g, *Streblus indicus*). Taxa replicated

across the two data sets performed well in phylogenetic analyses, with 15 out of 19 always

resolving as monophyletic in supermatrix analyses and 18/19 so resolving in ASTRAL

analyses, with only *Utsetela gabonensis* always forming a grade instead (with a difficult-to-

distinguish congener). Our results should embolden others to combine and repurpose data

sets in similar ways.

Higher taxonomy in Moraceae and the delimitation of the tribe Moreae

Our results support Corner's overall approach to the classification of Moraceae, if

not all of its details, including the primacy of inflorescence architecture and the

unreliability of inflexed stamens for higher taxonomy (Corner, 1962). Inflexed stamens are

a plesiomorphy that was lost nine times in Moraceae (Fig. 6), a preserved ancestral

character whose past taxonomic importance accounts for the rather extreme non-

monophyly of the Moreae. *Streblus* s.l. provides the best illustration of this principle,

appearing in four out of seven tribe-level clades within Moraceae. We may justly call its

disparate sections, as Corner did, "fragments of an ancestral *Streblus*." (Corner, 1975)—or

in modern parlance, a paraphyletic remnant preserving plesiomorphic staminate flowers

similar to those likely to have occurred in the ancestor of all Moraceae, with four free tepals

and four inflexed stamens (Clement & Weiblen, 2009). If one of the goals of modern

systematics is to establish taxonomic frameworks that reflect as far as possible real

evolutionary relationships, our results may serve as a warning to carefully investigate

whether characters used for taxonomy are derived (synapomorphic) or ancestral

(symplesiomorphic).

Broadly speaking, a tribal delimitation based on inflorescence architecture

(supplemented with other characters in some cases) agrees best with the phylogenetic trees

presented here. Moreae (as revised below) have unisexual spicate or racemose

inflorescences (the pistillate ones sometimes uniflorous), with the globose-capitate pistillate

inflorescences of the monotypic *Bagassa* as the sole exception. Stamens may be either

inflexed or straight. Elsewhere in the family, racemes and spikes are rare, with the former

found in *Maclura* (in part) and the latter found in the related genera *Broussonetia*,

Allaeanthus Thwaites, and *Malaisia* and arguably in *Batocarpus* H. Karst. and *Clarisia*; all

of these have capitate pistillate inflorescences. The taxa of *Streblus* s.l. and *Trophis* s.l. that

must be excluded from Moreae all have inflorescences that do not fit our general rule:

discoid-capitate (*Streblus asper* and *Trophis caucana*), cymose (*Streblus indicus*), or

bisexual (*Streblus usambarensis*). In these cases, perhaps Corner did not take his emphasis

on inflorescence architecture quite far enough, including too wide of a variety in this one

tribe. In critiquing the utility of inflorescence architecture for classification, Berg (1977b)

noted the similarity in the inflorescence structure of *Bleekrodea* (then part of Moreae, and

included in *Streblus* by Corner) to that of *Utsetela* and *Helianthostylis* (Dorstenieae), an

observation that proved prescient when *Bleekrodea* was found to belong to Dorstenieae

(Clement & Weiblen, 2009).

Olmedieae (as revised below, including Castilleae) can be defined entirely based

upon the presence of a discoid inflorescence subtended by an involucre of imbricate bracts.

Two species with such involucres previously classified as Moreae, *Streblus asper* and

Trophis caucana (=*Olmedia aspera*) always appeared in the Olmedieae (=Castilleae) clade

in our analyses (Figs. 2–3). *Olmedia aspera* (=*Trophis caucana*)—the nomenclatural type

of the tribe Olmedieae—was transferred to Moreae because of its inflexed stamens and lack

of self-pruning branches (Berg, 1977b). In inflorescence morphology, however, *T. caucana*

closely resembles other Castilleae, always subtended by an involucre of imbricate bracts

(Berg, 1977b, 2001). The staminate inflorescences of *Streblus asper* are strikingly similar
to those of *T. caucana*, and it is remarkable that the affinity between *S. asper* and the
Olmedieae has not been seriously considered until now. Previous barcoding or phylogenetic
studies have placed *Trophis caucana* (Kress & al., 2009) and *Streblus tonkinensis* (Chen &
al., 2016) (closely allied to *S. asper*) in the Castilleae clade, but those results went
unremarked upon, perhaps because of the broad scale of the studies (respectively, forest
community phylogenetics and the vascular plants of China).

The remaining five tribes can all be broadly defined based on inflorescence
architecture as Corner argued, sometimes supplemented by other characters as Berg
preferred, allowing of course for the exceptions made inevitable by the vicissitudes of
evolution. Ficeae of course is defined by the syconium (essentially an urceolate disc that
has been closed at the top). Maclureae have densely-packed globose infructescences and
can be distinguished from Artocarpeae by their four stamens (inflexed in most sections) and
armature. Artocarpeae also have densely-packed globose infructescences but only one
straight stamen and no armature. Parartocarpeae have a few connate involucral bracts and
(in large part) flowers embedded in fleshy receptacles. Dorstenieae are perhaps the most
heterogenous group, but in large part they have bisexual inflorescences, often capitate or
discoid, and often with ballistically-ejected endocarps.

The role of inflexed stamens in the evolution of Moraceae
The repeated losses of inflexed stamens (Fig. 6), which are associated with wind
pollination (Bawa & Crisp, 1980; Berg, 2001), raise the possibility that transitions from
wind to animal pollination, which have already been documented in Moraceae (Momose &
al., 1998; Sakai & al., 2000; Datwyler & Weiblen, 2004; Gardner & al., 2018), are even
more common within the family. Generally considered rare (Culley & al., 2002), the shift
from wind to animal pollination may be a repeated feature of Moraceae deserving of further
investigation. Further investigation of *Sorocea*, with its straight stamens and sometimes-
scented inflorescences, may reveal that, like *Artocarpus*, it contains both wind and animal
pollination. And while little is known about pollination in the Dorstenieae, the presence of
unisexual inflorescences and inflexed stamens (e.g., *Broussonetia*) as well as bisexual

inflorescences with straight stamens (e.g., *Brosimum, Dorstenia*) raises the possibility of

transitions within that clade as well; taxa with inflexed stamens but bisexual inflorescences

such as *Bleekrodea* and *Sloetia* (which is visited by bees, EMG pers. obs.) may represent

remnants of an intermediate state. Finally, the conclusion that a single transition to insect

pollination preceded the split between the Ficeae and Olmedieae (Castilleae) should be

reevaluated in light of the position of *Streblus asper* as sister to the latter (Figs. 2, 3). The

only shift in diversification rates our analyses recovered was on the branch leading to the

very diverse *Ficus*, suggesting that the shift away from wind pollination may not by itself

lead to increased diversification.

Explanation of taxonomic revisions

578   We present a generic revision of Moreae (Fig. 7) based on the present phylogenetic

study as well as morphological characters. Arranging monophyletic and morphologically

coherent genera requires one change of rank and 8 new combinations, but no entirely new

names. These revisions provide a framework for within-genus revisionary work, which will

require more intensive sampling and review of specimens within the genera circumscribed

here. We provide an explanation of the taxonomic changes, followed by a formal

presentation of the affected tribes and genera, with complete species lists and synonymy.

585   *Trophis* and the reinstatement of the Olmedieae — To make *Trophis* monophyletic,

we propose that it be applied strictly to the Neotropical clade that includes the type species

Trophis americana L. (=*Trophis racemosa* (L.) Urb.) . To accomplish this, we reinstate the

genus *Maillardia* and transfer *Trophis philippinensis* to *Paratrophis* (discussed below),

reinstating its former name *Paratrophis philippinensis* (Bureau) Fern.-Vill. Because

Trophis caucana does not belong in *Trophis* or Moreae but rather with the members of

Castilleae, we reinstate its former name *Olmedia aspera* Ruiz & Pav. and transfer it to that

tribe, whose former name Olmedieae must now be reinstated on account of its priority.

593   *Streblus* — We restrict the genus *Streblus* to three species comprising most of

section *Streblus*—*Streblus asper, Streblus tonkinensis*, and *Streblus celebensis* C.C. Berg—

and transfer the genus to Olmedieae. Although *S. celebensis* was not included in our

phylogeny, the sub-involucrate inflorescences are similar to those of *S. asper*, with which *S.*

celebensis differs primarily in vegetative characters, the latter having broadly-toothed

margins in the distal half of the leaf. Both occur in Sulawesi, where at least one specimen

with intermediate leaf morphology has been collected (Sulawesi, Kendari: *Kjellberg 452*,

24 Feb. 1929, L, det. *S. asper* by E.J.H. Corner). We therefore retain its taxonomic position

in *Streblus*. The remaining member of section *Streblus*, *S. usambarensis*, belongs with

Dorstenieae, and we therefore transfer it to that tribe, reinstating its former name

Sloetiopsis usambarensis. *Streblus indicus* is sister to Parartocarpeae, and we therefore

reinstate its former name *Pseudostreblus indicus* and transfer it to Parartocarpeae.

The remaining species of *Streblus* s.l. are properly placed in Moreae but are still

paraphyletic. We therefore reinstate the genera *Ampalis*, *Paratrophis*, and *Taxotrophis*,

largely corresponding to the former sections but requiring some new combinations.

Paratrophis as presented here is united by spicate male inflorescences (usually) with peltate

or reniform bracts (except for *Paratrophis philippinensis* (=*Trophis philippinensis*), which

does not have fleshy tepals in fruit). Within *Paratrophis* we include *Paratrophis ascendens*

(Corner) E.M. Gardner (=*Streblus ascendens* Corner) based on its inflorescence

morphology and phylogenetic position in the *rbc*L tree (Fig. S2b). Its previous position

within the monotypic section *Protostreblus* was due to the type specimen's spiral

phyllotaxy, but the latter may be atypical, as a more recent collection *Womersley NGF*

24791 (K, L, BO), has distichous leaves. Berg (1988), recognizing the close affinity

between *Pachytrophe dimepate* Bureau and *Ampalis mauritiana*, included both in *Streblus*

section *Ampalis*. Our phylogenetic results support this grouping, which we maintain in the

reinstated genus *Ampalis*, requiring one new combination. We follow Baillon in

maintaining *Pachytrophe* as a section of *Ampalis* in order to recognize the differences

between them in phyllotaxy, stipule amplexicaulity, and embryo characters. Further

intensive study of that species complex may of course warrant a different approach,

including potentially reducing both to a section of *Paratrophis*. Our species lists within the

former components of *Streblus* follow Berg's approach (1988, 2006), with two exceptions.

We provisionally recognize *Taxotrophis zeylanica* (Thwaites) Thwaites as distinct from

Taxotrophis taxoides (B. Heyne ex Roth) W.L. Chew ex E.M. Gardner (=*Streblus taxoides*

(B. Heyne ex Roth) Kurz) based on our phylogenetic work and following the *Flora of*

China, which recognizes *Streblus zeylanicus* as distinct from *S. taxoides* based on its

clustered pistillate inflorescences. In addition, we provisionally recognize *Paratrophis*

australiana (=*S. glaber* subsp. *australianus*) as distinct from *Paratrophis glabra* based on

its geographic and consistent morphological distinctiveness. *Taxotrophis* and *Paratrophis*

warrant further investigation to refine species limits.

Morus — Morus has never been revised, and the species concepts are often based on

minor morphological differences (Berg et al., 2006; Berg 2001), and the paraphyly of

several species in our analyses suggests that a broad *M. alba* similar to Bureau's (1873)

may be worth a second look. The monotypic subgenus *Afromorus* is sister to *Milicia*, but

the leaf morphology is markedly different, instead resembling other *Morus* species in its

trinerved based and crenate margins. We therefore raise *Afromorus* to genus level, requiring

one new combination. *Morus insignis*, from western Central and South America, bears a

remarkable resemblance to *Paratrophis*, in particular *P. pendulina*, especially in leaf

morphology, which in *M. insignis* is not consistently trinerved as in other mulberries. The

infructescence appears superficially like a mulberry because of its basally fleshy tepals,

although with more loosely-packed flowers, but closer inspection places it firmly within

Paratrophis, with drupes protruding from the persistent tepals, peltate bracts, and a sterile

groove. Because analyses differ as to whether *M. insignis* is sister to *Paratrophis* or part of

it, it seems best to treat them as congeners.

These changes result in ten monophyletic genera of Moreae, providing a framework

for revisionary work within the genera. Below, we present a complete genus and species list

for Moreae with brief descriptions for all genera and synonymies for new combinations and

reinstated or recircumscribed taxa. All cited protologues were reviewed, and dates for

works published piecemeal were confirmed by reference to Taxonomic Literature 2 online

(https://www.sil.si.edu/DigitalCollections/tl-2/index.cfm).

Taxonomic Treatment

Key to the tribes of Moraceae

Key to the tribes of Moraceae

This key follows the tribal circumscription of Clement & Weiblen (2009), as subsequently

modified by Zerega & al. (2010), Chung & al. (2017), Zerega & Gardner (2019), and this

study.

1. Inflorescence a syconium (urceolate with the opening entirely closed by ostiolar

663   bracts, flowers enclosed at all stages of development) – Ficeae (*Ficus*)

664   1. Inflorescence not a syconium (capitate, spicate, discoid, or urceolate, but

665   flowers not entirely enclosed at all developmental stages) – 2

667   2. Inflorescences (at least staminate) with an involucre of imbricate bracts; often

668   with self-pruning horizontal branches (except *Olmedia*, *Poulsenia, Streblus*) –

669   Olmedieae (*Antiaris, Antiaropsis, Castilla, Helicostylis, Maquira, Mesogyne,*

670   *Naucleopsis, Olmedia, Perebea, Poulsenia, Pseudolmedia, Sparattosyce*,

671   *Streblus*)

672   2. Inflorescences not involucrate; plants without self-pruning branches – 3

674   3. Plants woody; dioecious; pistillate inflorescences globose-capitate; spines axillary

675   or terminating short shoots – Chlorophoreae (*Maclura*)

676   3. Plants woody, herbaceous, or succulent; dioecious or monoecious; pistillate

677   inflorescences various; spines absent or if present, then pistillate inflorescences

678   not globose capitate – 4

680   4. Trees or shrubs; monoecious; inflorescences unisexual; staminate flowers with

681   one stamen (rarely two) – Artocarpeae (*Artocarpus, Batocarpus, Clarisia*)

682   4. Trees, shrubs, lianas, herbaceous, or succulent; monoecious or dioecious;

683   inflorescences unisexual or bisexual; staminate flowers with more than one

684   stamen (or if one stamen then dioecious) – 5

5. Trees or shrubs; monoecious; inflorescences unisexual; stamens straight in bud or

staminate flowers 5-parted and inflexed in bud – Parartocarpeae (*Hullettia,*

Parartocarpus, Pseudostreblus)

5. Trees, shrubs, lianas, herbaceous, or succulent; monoecious or dioecious;

inflorescences bisexual or unisexual; stamens straight or inflexed in bud but

staminate flowers never 5-parted – 6

6. Trees, shrubs, lianas, herbaceous, or succulent; inflorescences bisexual (or if

unisexual then a climber or herbaceous); endocarp body often ballistically ejected

from infructescence – Dorstenieae in part (*Allaeanthus* in part, *Bleekrodea,*

Bosqueiopsis, Brosimum, Broussonetia in part, *Dorstenia, Fatoua,*

Helianthostylis, Malaisia, Scyphosyce, Sloetia, Sloetiopsis, Treculia, Trilepsium,

Trymatococcus, Utsetela)

6. Trees or shrubs; inflorescences unisexual; endocarp body never ballistically

ejected – 7

7. Trees; stamens inflexed in bud; pistillate inflorescences globose-capitate –

Dorstenieae in part (*Allaeanthus* in part, *Broussonetia* in part)

7. Trees or shrubs; stamens inflexed or straight in bud; pistillate inflorescences

various, not globose-capitate if stamens are inflexed in bud – Moreae (*Afromorus,*

Ampalis, Bagassa, Maillardia, Milicia, Morus, Paratrophis, Taxotrophis,

Sorocea, Trophis)

Revisions

712   **Tribe MOREAE**

Moreae Gaudich. in Freyc., Voy. Uranie, Bot. (1830) 509.

Moreae subtribe Soroceae Miq in Martius, Fl. Bras. 4, 1, fasc. 12: 111 (1853), p.t. –

Soroceae (Miq.) C.C. Berg, Blumea 50: 537 (2005), p.p.

Strebleae Bureau in DC., Prodr. 17: 215 (1873), p.p.

Tree or shrubs, monoecious or dioecious. *Leaves* alternate or opposite, distichous or

spirally arranged; stipules lateral to amplexicaul. *Inflorescences* unisexual, uniflorous or

racemose, spicate, capitate, or globose; bracteate; tepals 4, free to connate; staminate

flowers with 4 stamens, filaments straight or inflexed in bud, pistillode usually present;

pistillate flowers with (mostly) free ovary, 2 stigmas. *Fruits* drupaceous or achene-like

with a fleshy persistent perianth, dehiscent or not. *Seeds* with or without endosperm, testa

usually with a thick vascularized part below the hilum, cotyledons equal or unequal,

straight or folded.

Genera and distribution: ten genera and 63 species with a worldwide distribution.

731   *Afromorus*

Afromorus **E.M. Gardner, gen. nov.** — based on *Morus* L. *subg. Afromorus* [A. Chev.,

Rev. Bot. Appl. Agr. Trop. 29 (315–316): 70 (1949), invalidly published]; J.-F. Leroy,

Rev. Bot. Appl. Agr. Trop. 29 (323–324): 482 (1949) & Bull. Mus. Hist. Nat. Paris,

ser. 2, 21: 732 (1949). TYPE: *Afromorus mesozygia* (Stapf ex A. Chev.) E.M.

Gardner.

Dioecious trees, shoot apices deciduous. *Leaves* distichous, triplinerved or at least trinerved

at the base. *Stipules* free, more or less lateral. *Inflorescences* solitary or paired, bracts

of varying shapes. Staminate inflorescences spicate, to 2.5mm long, flower 4-parted,

tepals imbricate, ciliolate, pistillode small, apiculate. Pistillate inflorescences

subglobose, ca. 5 mm across, flowers 4-parted, tepals ciliolate, stigma bifid, equal or

unequal, arms filiform to 5 mm long. *Infructescences* subglobose or less often slightly

elongate, ca. 1 mm across, tepals fleshy, yellowish to greenish, drupes ca. 5 x 3–5 mm.

Seeds ca. 4.5 x 2.5–4.5 mm.

Species and distribution: one species, in tropical Africa.

Note: The leaves of *Afromorus*—with crenate margins and trinerved bases—bear a striking

resemblance to those of *Morus*, and it is thus not surprising that the former was heretofore

included within the latter. However, leaves of *Afromorus* are distinct in being usually

completely triplinerved, without an upper pinnately-veined portion as in most species of

Morus. As the genus is monotypic, the type must be its only species, *Afromorus mesozygia*

(Stapf ex A. Chev.) E.M. Gardner.

The genus name is based on Leroy's invalidly published *Morus* subgenus

Afromorus, which was published only in French without a Latin diagnosis or description as

required under the Code.

1. *Afromorus mesozygia* (Stapf ex A. Chev.) E.M. Gardner, comb. nov. — based on *Morus*

mesozygia Stapf ex A. Chev. [Végétaux utiles de l'Afrique tropicale française 5: 263

(1909), nomen solum], J. Bot. (Morot) ser. 2, t. 2: 99 (1909).

Celtis lactea Sim, For. Fl. Port. E. Afr. : 97, t. 96 (1909), probably post-dating *Morus*

mesozygia (fide Berg. 1977) – *Morus lactea* (Sim) Mildbr., Notizbl. Bot. Gart.

Berlin 8: 243 (1922) – *Morus mesozygia* var. *lactea* (Sim) A. Chev., Rev. Bot.

Appl. Agr. Trop. 29: 72 (1949).

Morus mesozygia var. *sanda* A. Chev., Rev. Bot. Appl. Agr. Trop. 29: 71 (1949),

invalidly published (under Art. 39.1, Latin description or diagnosis lacking).

Morus mesozygia var. *colossea* A. Chev., Rev. Bot. Appl. Agr. Trop. 29: 71 (1949),

invalidly published (under Art. 39.1, Latin description or diagnosis lacking).

Ampalis

Ampalis Bojer, Hort. Maurit. 291 (1837)

Streblus Lour. subgen. *Parastreblus* Blume, Mus. Bot. Ludg.-Bat. 2: 89 (1856)

Streblus Lous. sect. *Ampalis* (Bojer) C.C. Berg, Proc. Kon. Ned. Akad. Wetensch. C

91(4): 358 (1988).

Pachytrophe Bureau, Prodr. [A.P. de Candolle] 17: 234 (1873). – *Ampalis* Boj. sect.

Pachytrophe (Bur.) Baillon, Hist. Pl. 6: 191 (1875–76)

Dioecious trees or shrubs. ***Leaves*** distichous to spirally arranged, pinnately veined. ***Stipules***

free, nearly lateral. ***Inflorescences*** solitary or paired in the leaf axils, spicate, with an

abaxial sterile groove, flowers in longitudinal rows, bracts basalt attached to subpeltate.

Staminate inflorescences to 9 cm long, flowers 4-parted, decussate-imbricate, stamens 4,

inflexed in bud, pistillode present. Pistillate inflorescences to 12 cm long, tepals 4, separate,

decussate-imbricate, ovary free, stigmas 2, equal. ***Infructescences*** with enlarged fleshy

perianths ca. 6–8 mm long, surrounding drupaceous fruits, the latter ca. 5–6 mm long.

Seeds ca. 4 × 4 mm, testa thickened and not distinctly vascularized, cotyledons equal.

Species and distribution: Two species, native to Madagascar and Comoros.

Note: The two sections differ in their phyllotaxy (usually spiral in *Ampalis* and ditichous in

Pachytrophe) and stipules (free in *Ampalis*, connate in *Pachytrophe*). Sect. *Ampalis*

usually has somewhat larger inflorescences.

Ampalis* sect. *Ampalis

1. *Ampalis mauritiana* (Jacq.) Urb., Symb. Antill. 8: 165 (1920) – *Morus mauritiana* Jacq.,

Collect. 3: 206 ("1789", 1791) – *Streblus mauritianus* (Jacq.) Blume, Mus. Bot.

Lugd.-Bat. 2: 80 (1856).

Streblus maritimus Palaky, Catal. Pl. Madag. 2: 31 (1907)

Morus nitida Willem. In Useri, Ann. Bot. 18: 56 (1796)

M. ampalis Poir. In Lam., Encycl. Bot. 4: 380 (1797)

Trophis cylindrica Roxb., Fl. Indica 3: 599 (1832), pro syn. *M. mauritianae.*

Ampalis madagascariensis, Bojer, Hort. Maurit. 291 (1837) nom. illeg., pro syn.

Ampalis madagascariensis var. *occidentalis* Léandri, Mém. Inst. Sci. Madag., ser. B,

1: 12, pl. (1948).

Morus rigida Hassk., Cat. Hort. Bog. 74 (1844).

Ampalis Boj. sect. *Pachytrophe* (Bureau) Baillon, Hist. Pl. 6: 191 (1875–76) ≡

Pachytrophe Bureau in A.DC, Prodr. 17: 234 (1873). – Type: *Pachytrophe dimepate*

Bureau.

2. *Ampalis dimepate* (Bureau) E.M.Gardner, comb. nov., based on *Pachytrophe dimepate*

Bureau in A.DC Prodr. 17: 234 (1873) – *Streblus dimepate* (Bureau) C.C.Berg, Proc.

Kon. Ned. Akad. Wetensch. C 91(4): 358 (1988).

Pachytrophe obovata Bureau, Prodr. [A.P. de Candolle] 17: 235 (1873).

Plecospermum bureaui A.G. Richt., Term. Füzetek 18: 296 (1895), nomen nudum et

superfl., pro. syn. nom. ined. "Plecospermum obovatum Bureau" (*Boivin 1717*, P).

Plecospermum (?) laurifolium Baill. in Grandidier, Hist. Madag. Vol. 35, Hist. Nat.

Pl., Tome 5, Atlas 3, 2e partie: pl. 294a (1895) – *Pachytrophe obovata* var.

laurifolia (Baill.) Léandri, Mém. Inst. Sci. Madag., ser. B, 1: 16 (1948).

Pachytrophe obovata var. *montana* Léandri, Mém. Inst. Sci. Madag., ser. B, 1: 16

(1948).

825   ***Bagassa***

Bagassa Aubl., Pl. Guiane. Suppl. 15 (1775).

Dioecious trees. ***Leaves*** opposite and decussate, lamina triplinerved, 3-lobed to entire.

Stipules free, lateral. ***Inflorescences*** solitary or paired in the leaf axils, bracteate. Staminate

inflorescences spicate with an abaxial sterile groove, to 12 cm long, flowers in longitudinal

rows with, tepals 4, stamens 2, straight in bud, pistillode present. Pistillate inflorescences

globose-capitate, ca. 1–1.5 cm across, flowers 4-lobed to 4-parted, stigmas 2, filiform.

Infructescences globose, ca. 2.5–3.5 cm across, green, tepals fleshy, yellowish to greenish,

drupes ca. 7–8 mm long. *Seeds* ca. 3 x 2 mm, testa thin and not vascularized, cotyledons

equal.

Species and distribution: One, in tropical South America.

1. *Bagassa guianensis* Aubl., Pl. Guiane. Supple. 15 (1775)

843   *Maillardia*

Maillardia Frapp. ex Duch. in Maillard, Notes sur l'île de la Réunion, Annex P. 3 (1862)

Trophis P. Br. sect. *Maillardia* (Duch.) Corner, Gard. Bull. Singapore 19: 230 (1962).

Dioecious trees or shrubs. *Leaves* distichous, pinnately veined. *Stipules* free, semi-

amplexicaul. *Inflorescences* solitary or paired in the leaf axils, bracts subpeltate. Staminate

inflorescences spicate with an abaxial sterile groove, flowers 4-parted, decussate-imbricate,

stamens 4, inflexed in bud, pistillode present. Pistillate inflorescences up to three together,

uni- or bi-florous, perianth tubular 4-lobed, ovary adnate to the perianth, stigmas 2, equal.

Infructescences with enlarged fleshy perianths surrounding drupaceous fruits, the latter to

18 mm long. *Seeds* to 13 mm long, testa thin, with a thickened vascularized part below the

hilum, cotyledons unequal.

Species and distribution: Two species, in Madagascar, Comoros, and the Seychelles.

1. *Maillardia borbonica* Duch., Ann. Notes Réunion, Bot. 1: 148 (1862) – *Trophis*

borbonica (Duch.) C.C.Berg, Proc. Kon. Ned. Akad. Wetensch. C 91(4): 355 (1988)

Maillardia lancifolia Frapp. ex Duch., Ann. Notes Réunion, Bot. 1: 148 (1862).

Trophis borbonica (Duch.) C.C. Berg, Proc. Kon. Ned. Akad. Wetensch. C 91: 355

(1988).

2. *Maillardia montana* Leandri, Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 1: 25
(1948) – *Trophis montana* (Leandri) C.C.Berg, Proc. Kon. Ned. Akad. Wetensch. C
91(4): 355 (1988).
Maillardia occidentalis Leandri, Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 1: 26
(1948)
Maillardia orientalis Léandri, Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 1: 27
(1948)
Maillardia mandrarensis Léandri, Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 1:
28 (1948).
Maillardia pendula Fosberg, Kew Bull. 29: 266, t. 2 (1974).
875
876
877   ***Milicia***
878 *Milicia* Sim, For. Fl. Port. E. Afr. 97 (1909).
879
Dioecious trees. ***Leaves*** distichous, lamina pinnately veined. ***Stipules*** free, not fully
amplexicaul. ***Inflorescences*** spicate, usually solitary in the leaf axils or on leafless nodes,
spicate, flowers in longitudinal rows alternating with rows of bracts, bracts mostly basally
attached, abaxial sterile groove present. Staminate inflorescences to 20 cm long, flowers 4-
parted, imbricate, stamens 4, inflexed in bud, pistillode present. Pistillate inflorescences to
4.5 cm long, flowers 4-parted, decussate-imbricate, ovary free, stigmas 2, unequal.
Infructescences with enlarged fleshy perianths, surrounding slightly flattened drupaceous
fruits, the latter to ca. 3 mm long. ***Seeds*** ca. 2 mm long, testa thin with a slightly thickened
vascularized part below the hilum, cotyledons equal.
889
890 Species and distribution: Two species in tropical Africa.
891
1. *Milicia excelsa* (Welw.) C.C. Berg, Bull. Jard. Bot. Natl. Belg. 52(1-2): 227 (1982) –
Morus excelsa Welw., Trans. Linn. Soc. Lond. (Bot.) 27: 69, t. 23 (1869) – *Maclura*

excelsa (Welw.) Bur. In DC., Prodr. 17: 231 (1873) – *Chlorophora excelsa* (Welw.)

Benth. & Hook., Gen. Pl. 3(1): 363 (1880).

Chlorophora tenuifolia Engl., Bot. Jahrb. 20: 139 (1894).

Milicia africana Sim, For. Fl. Port. E. Afr. 97, t. 122 (1909).

Chlorophora alba A. Chev., Bull. Soc. Bot. Fr. 58 (Mém. 8d): 209 (1912).

2. *Milicia regia* (A. Chev.) C.C. Berg, Bull. Jard. Bot. Natl. Belg. 52(1-2): 227 (1982) –

Chlorophora regia A. Chev., Bull. Soc. Bot. Fr. 58 (Mém. 8d): 209 (1912) – *Maclura*

regia (A. Chev.) Corner, Gard. Bull. Singapore 19: 237 (1962).

Morus

Morus L., Sp. Pl. 986 (1753).

Dioecious trees or shrubs. ***Leaves*** trinerved (to five-nerved) at the base. ***Stipules*** free,

nearly lateral. ***Inflorescences*** solitary or paired in the leaf axils, without an obvious sterile

groove, interfloral bracts absent. Staminate inflorescences spicate or racemose, to 8 cm

long, flowers 4-parted, imbricate, stamens 4, inflexed in bud, pistillode present. Pistillate

inflorescences subcapitate to spicate, up to 16 cm long, flowers tepals 4, separate,

decussate-imbricate, ovary free, stigmas 2, equal. ***Infructescences*** with enlarged fleshy

perianths enclosing achene-like fruits, the latter ca. 1 mm long. ***Seeds*** less than 1 mm long,

cotyledons equal.

Species and distribution: Sixteen species, Asia and North to Central America; introduced

worldwide.

1. *Morus alba* L., Sp. Pl. 2: 986 (1753).

var. *alba*

var. *multicaulis* (Perrottet) Loudon, Arbor. Frutic. Brit. 3: 1348 (1838).

2. *Morus australis* Poir. in Desrousseaux et al., Encycl. 4: 380 (1797).

3. *Morus boninensis* Koidz., Bot. Mag. (Tokyo) 31: 38 (1917)

4. *Morus cathayana* Hemsl., J. Linn. Soc., Bot. 26: 456 (1894)

var. *cathayana*

var. gongshanensis (Z. Y. Cao) Z. Y. Cao, Acta Bot. Yunnan. 17: 154 (1995).

5. *Morus celtidifolia* Kunth, Nov. Gen. Sp. [H.B.K.] 2: 33 (1817).

6. *Morus liboensis* S. S. Chang, Acta Phytotax. Sin. 22: 66 (1984).

7. *Morus macroura* Miq., Pl. Jungh. 1: 42 (1851).

var. *macroura*

var. *laxiflora* G.K.Upadhyay & A.A.Ansari, Rheedea 20: 44 (2010).

8. *Morus koordersiana* J.-F.Leroy, Bull. Mus. Natl. Hist. Nat. sér. 2, 21: 729 (1949)

(endemic to Sumatra and possibly synonymous with *M. macroura*. However, it was

not cited by Berg et al. (2006) either as a good species or as a synonym of the latter).

9. *Morus microphylla* Buckley, Acad. Nat. Sci. Philadelphia 1862: 8 (1863) (recognized in

the *Flora of North America* (Flora of North America Editorial Committee, 1993) but

likely conspecific with *M. celtidifolia* and considered so by Berg (2001)).

10. *Morus mongolica* (Bureau) C. K. Schneid. in Sargent, Pl. Wilson. 3: 296 (1916).

11. *Morus nigra* L. Sp. Pl. 2: 986 (1753).

12. *Morus notabilis* C. K. Schneid. in Sargent, Pl. Wilson. 3: 293 (1916).

13. *Morus rubra* L., Sp. Pl. 986 (1753).

var. *rubra*

var. *murrayana* (Saar & Galla) Saar, Phytologia 91: 106 (2009).

14. *Morus serrata* Roxb., Fl. Ind., ed. 1832, 3: 596 (1832).

15. *Morus trilobata* (S. S. Chang) Z. Y. Cao, Acta Phytotax. Sin. 29: 265 (1991).

16. *Morus wittiorum* Hand.-Mazz., Anz. Akad. Wiss. Wien, Math.-Naturwiss. Kl. 58: 88

(1921)

951   *Paratrophis*

Paratrophis Blume, Ann. Mus. Bot. Lugduno-Batavi 2 (1856) 81.

Pseudomorus Bureau, Ann. Sci. Nat., Bot. sér. 5, 11: 371 (1869).

Uromorus Bureau in A.DC., Prodr. 17: 236 (1873).

Calpidochlamys Diels, Bot. Jahrb. Syst. 67: 172 (1935); — *Trophis* sect.

Calpidochlamys Corner, Gard. Bull. Singapore 19: 230 (1962).

Chevalierodendron J.-F. Leroy, Compt. Rend. Hebd. Séances Acad. Sci. 227: 146

(1948) .

Streblus Lour. sect. *Paratrophis* (Blume) Corner, Gard. Bull. Singapore 19: 216

(1962).

Streblus Lour. sect. *Protostreblus* Corner, Blumea 18: 393 (1970).

Morus L. *subg. Gomphomorus* J.-F. Leroy, Bull. Mus. Hist. Nat. (Paris), Sér. 2, 21:

732 (1949)

Dioecious trees or shrubs. **Leaves** distichous (or spiral in *P. ascendens*), pinnately veined,

sometimes trinerved at the base. Stipules free, lateral. **Inflorescences** axillary, solitary or up

to 5 together, spicate, with an abaxial sterile groove, interfloral bracts mostly peltate,

flowers sessile in longitudinal rows, tepals 4, valvate, ciliolate. Staminate inflorescences up

to at least 20 cm. long, flowers with filaments inflexed in bud, pistillode present. Pistillate

inflorescences up to at least 10 cm long, flowers usually at least 2 (to many), tepals free

(except in *P. philippinensis*), stigma bifid, arms equal. **Fruits** drupaceous, red to black, with

tepals persistent but usually not enlarged or fleshy (except in *P. philippinensis* and *P.*

insignis), up to ca. 1 cm long. **Seeds** up to ca. 8 x 6 mm, cotyledons equal.

Species and distribution: Twelve species from the Malesian region to Australia, New

Zealand, and Oceania, Central and western South America.

1. *Paratrophis anthropophagorum* (Seem.) Benth. & Hook. f. ex Drake, Ill. Fl. Ins. Pacif.

296 1892

Trophis anthropophagorum Seem. [Bonplandia 9 (17–18): 259 (1861), nomen nudum]

Fl. Vit. 258, t. 68 (1868); — *Uromorus anthropophagorum* (Seem.) Bureau in de

Candolle, Prodr. 17 (1873) 236; — *Streblus anthropophagorum* (Seem.) Corner,

Gard. Bull. Sing.19 (1962) 221.

Caturus oblongatus Seem., Fl. Vit. 254 (1868).

Paratrophis ostermeyri Rech., Fedd. Rep. 5: 130 (1908).

Paratrophis viridissima Rech., Fedd. Rep. 5: 130 (1908).

Paratrophis zahlbruckneri Rech., Fedd. Rep. 5: 130 (1908).

Pseudomorus brunoniana (Endl.) Bureau var. *tahitensis* J. Nadeaud, Énum. Pl. Tahiti

43 (1873) – *Uromorus tahitensis* (J. Nadeaud) Bureau in de Candolle, Prodr. 17: 237

(1873) – *Paratrophis tahitensis* (Bureau) Benth. & Hook.f. ex Drake, Ill. Ins. Mar.

Pacif. Fasc. 7: 296 (1892); et Fl. Polyn. Franc. 193 (1892). – *Streblus tahitensis* (J.

Nadeaud) E.J.H. Corner, Gard. Bull. Singapore 19: 225 (1962).

2. *Paratrophis ascendens* (Corner) E.M. Gardner comb. nov. — based on *Streblus*

ascendens Corner, Blumea 18: 395, t.1 (1970).

3. *Paratrophis australiana* C.T. White, Contr. Arnold Arbor. 4: 15 (1933) — *Streblus*

glaber (Merr.) Corner var. *australianus* (C.T. White) Corner, Gard. Bull. Singapore

19: 221 (1962) — *Streblus glaber* (Merr.) Corner subsp. *australianus* (C.T. White)

C.C. Berg, Blumea 50: 548 (2005).

1002

1003   **4. *Paratrophis banksii*** Cheeseman, Man. N.Z. Fl. 633 (1906) – *Streblus banksii*

1004   (Cheeseman) C.J. Webb, in Connor & Edgar, New Zealand J. of Bot. 25 136 (1987).

1005   *Paratrophis heterophylla* var. *elliptica* Kirk T.N.Z.L 29: 500, t. 46 (1897); — *Streblus*

1006   *heterophyllus* var. *ellipticus* (Kirk) Corner, Gard. Bull. Singapore 19: 222 (1962).

1007   Note:— Morphologically very close to and not always distinguishable from *P.*

1008   *microphylla*.

1009

5. *Paratrophis glabra* (Merr.) Steenis**,** J. Bot. 72: 8 (1934) – *Gironniera glabra* Merr.,

Philipp. J. Sci., 1, Suppl. 42 (1906); Enum. Philipp. Flow. Pl. 2: 35 (1923) —

Chevalierodendron glabrum (Merr.) J.-F. Leroy, Compt. Rend. Hebd. Séances Acad. Sci.

227: 146 (1948) — *Streblus glaber* (Merr.) Corner, Gard. Bull. Singapore 19: 221 (1962).

Aphananthe negrosensis Elmer, Leafl. Philipp. Bot. 2: 575 (1909).

Pseudostreblus caudatus Ridl., J. Fed. Malay States Mus. 6: 54 (1915).

Streblus laevifolius Diels, Bot. Jahrb. Syst. 67: 171 (1935).

Streblus urophyllus Diels, Bot. Jahrb. Syst. 67: 172 (1935) — *Streblus glaber* (Merr.)

Corner subsp. *urophyllus* (Diels) C.C. Berg, Blumea 50: 548 (2005);

Streblus urophyllus Diels var. *salicifolius* Corner, Gard. Bull. Singapore. 19: 225 (1962).

1020

Note: Berg et al. (2006) treated *P. australiana* and *Streblus urophyllus* as subspecies of

Streblus glaber. *Paratrophis australiana*, endemic to Australia, has crenate leaf margins

and somewhat larger staminate inflorescences with more interfloral bracts than *P. glabra*;

these morphological differences are consistent and geographically confined, and we

therefore reinstate *P. australiana*. This stands in contrast to *Streblus urophyllus*, which we

provisionally maintain in synonomy under *P. glabra*. Although collections from Mt.

Wilhelm in New Guinea are remarkable for their thick coriaceous leaves and spinose

margins, similar toothed margins can be found at higher elevations in Borneo and Sulawesi,

suggesting that the striking leaf morphology of *S. urophyllus* is an alpine effect not

indicative of speciation. We reserve judgment on the status of *Streblus urophyllus* var.

salicifolius Corner, applied by Corner (1962) to specimens with linear leaves, provisionally

maintaining it in synonomy following Berg et al. (2006).

1033

6. *Paratrophis insignis* (Bureau) E.M. Gardner comb. nov. — based on *Morus insignis*

Bureau, in De Candolle, Prodr. 17: 247 (1873).

Morus peruviana Planch. ex Koidz., Fl. Symb. Orient.-Asiat. 88 (1930).

Morus trianae J.-F.Leroy, Bull. Mus. Hist. Nat. (Paris) Sér. 2, 21: 731 (1949).

Morus marmolii Legname, Lilloa 33: 334 (1973)

Note: The infructescences differ from most other *Paratrophis* in their fleshy free tepals and

denser aggregation of flowers, but the staminate inflorescences are typical of the genus.

1041

1042   **7. *Paratrophis microphylla*** (Raoul) Cockayne, Bot. Notes Kennedy's Bush & Scenic Res.

1043   Port Hills, Lyttelton (Rep. Scenery Preserv.) 3 (1915) – *Epicarpurus microphyllus*

1044   Raoul, Choix Pl. Nouv.-Zel. 14. (1846) – *Taxotrophis microphylla* (Raoul) F. Muell.,

1045   Fragm. (Mueller) 6(47): 193 (1868).

1046   *Paratrophis heterophylla* Blume, Mus. Bot. 2(1-8): 81. (1856).

1047

8. *Paratrophis pendulina* (Endl.) E.M. Gardner, comb. nov. — based on *Morus*

pendulina Endl., Prodromus Florae Norfolkicae 40 (1833) – *Pseudomorus pendulina*

(Endl.) Stearn, J. Arnold Arbor. 28: 427 (1947) – *Pseudomorus brunoniana* var.

pendulina (Endl.) Bureau, Ann. Sci. Nat. Bot. sér. 5, 11: 372 (1869) – *Streblus*

pendulinus (Endl.) F. Muell., Fragmenta Phytographiae Australiae (1868).

Morus brunoniana Endl., Atakta Bot. t. 32 (1835) – *Streblus brunonianus* (Endl.) F.

Muell., Fragm. Phyt. Australiae 6: 192 (1868) – *Pseudomorus brunoniana* (Endl.)

Bureau, Ann. Sci. Nat., Bot. sér. 5, 11: 372 (1869)

Pseudomorus brunoniana var. *australiana* Bureau, Ann. Sci. Nat., Bot. sér. 5, 11: 373

(1869) – *Pseudomorus pendulina* var. *australiana* (Bureau) Stearn, J. Arnold Arbor.

28: 427 (1947).

Pseudomorus brunoniana var. *australiana* subvar. *castaneaefolia* Bureau, Ann. Sci.

Nat., Bot. sér. 5, 11: 372 (1869).

Pseudomorus brunoniana var. *obtusa* Bureau, Ann. Sci. Nat., Bot. sér. 5, 11: 373

(1869) – *Pseudomorus pendulina var. obtusa* (Bureau) Stearn, J. Arnold Arbor. 28:

428 (1947), as var. *obtusata*.

Pseudomorus sandwicensis O. Deg., Fl. Hawaiiensis fam. 38 (1938) – *Pseudomorus*

brunoniana var. *sandwicensis* (O. Deg.) Skottsb., Acta Horti Gothob. 15: 347

(1944) – *Pseuomorus pendulina* var. *sandwicensis* (O. Degener) Stearn, J. Arnold

Arbor. 28: 428 (1947) – *Streblus sandwicensis* (O. Deg.) H. St. John, Pacific Trop.

Bot. Gard. Mem. 1: 374. 1973.

Note: This complex requires further investigation. Although the types of *Morus pendulina*,

Morus brunoniana, *Pseudomorus sandwicensis*, and *Pseudomorus brunoniana* var. *obtusa*

differ from one another, it is difficult to discern clear geographic patterns that warrant

maintaining separate species, and we therefore provisionally treat them all as one

widespread and variable species.

1074

9. *Paratrophis philippinensis* (Bureau) Fern.-Vill., Nov. App. 98 (1880) — *Uromorus*

philippinensis Bureau in A.DC., Prodr. 17: 237 (1873) — *Trophis philippinensis* (Bureau)

Corner, Gard. Bull. Singapore 19: 231 (1962).

Sloetia minahassae Koord., Versl. Minahasa 645 (1898).

Paratrophis grandifolia Elmer, Leafl. Philipp. Bot. 5: 1814 (1913).

Calpidochlamys branderhorstii Diels, Bot. Jahrb. Syst. 67: 173 (1935) — *Trophis*

branderhorstii (Diels) Corner, Gard. Bull. Singapore 19: 231 (1962).

Calpidochlamys drupacea Diels, Bot. Jahrb. Syst. 67: 173 (1935) — *Trophis drupacea*

(Diels) Corner, Gard. Bull. Singapore 19: 231 (1962).

Note: This species is the only *Paratrophis* with fused tepals and one of only two whose

tepals develop into a fleshy accessory fruit. The staminate inflorescences, however, are

typical of the genus.

1087

1088 **10. *Paratrophis sclerophylla* (Corner) E.M. Gardner, comb. nov.** — based on *Streblus*

1089 *sclerophyllus* Corner, Blumea 18: 399 (1970).

1090

1091 **11. *Paratrophis smithii*** Cheeseman T.N.Z.L 20:148 (1888) – *Streblus smithii* (Cheeseman)

1092 Corner Gard. Bull. Singapore 19: 224 (1962).

1093 Note: Morphologically very close to *P. anthropophagorum*.

1094

1095 **12. *Paratrophis solomonensis* (Corner) E.M. Gardner, comb. nov.** — based on *Streblus*

1096 *solomonensis* Corner, Gard. Bull. Singapore 19: 224 (1962).

1097 Note:— Morphologically very close to *P. anthropophagorum*.

1098

1099

1100 ***Sorocea***

1101

1102   *Sorocea* A. St-Hil., Mém. Mus. Hist. Nat. 7: 473 (1821).

1103   *Balanostreblus* Kurz, J. Asiat. Soc. Bengal., Pt. 2, Nat. Hist., 42: 247 (1873), p.p.

1104   *Pseudosorocea* Baill., Hist. Pl. 6: 210 (1875).

1105   *Trophisomia* Rojas Acosta, Bull. Acad. Inst. Geogr. Bot. 24: 211 (1914).

1106   *Paraclarisia* Ducke, Arq. Serv. Florest. 1(1): 2 (1939).

1107

Dioecious trees. **Leaves** alternate, distichous, lamina pinnately veined, 3-lobed to entire.

Stipules free, lateral. **Inflorescences** solitary or paired in the leaf axils or below the leaves,

racemose to spicate to subcapitate or uniflorous, bracts basally attached to peltate.

Staminate flowers 4-lobed to 4-parted, tepals decussate-imbricate, stamens (3–)4, straight

in bud, pistillode usually absent. Pistillate flowers tubular, 4-lobed to sub-entire, ovary

basally adnate to the perianth, stigmas 2, short, usually tongue-shaped. **Infructescences**

with drupaceous fruits enclosed by fleshy enlarged perianths, the latter red to orange,

turning black at maturity. **Seeds** large, testa thin, embryo green, cotyledons very unequal,

the smaller one minute and enclosed by the larger.

1117

1118   Species and distribution: 19 species in Central and South America from Mexico to

1119   Argentina.

1120

1121   Note: This species list follows the *Flora Neotropica* monograph (Berg 2001) with the

1122   addition of four new species and one status change published later. Names synonymized by

1123   Berg (2001) but recognized by Burger et al. (1962) are noted in parentheses. Complete

1124   synonymies can be found in those treatments.

1125

1126   *Sorocea* subg. *Sorocea*

1127   **Sorocea affinis** Hemsl., Biol. Cent.-Amer., Bot. 3: 150 (1883).

1128   **Sorocea angustifolia** Al.Santos & Romaniuc, Novon 24: 199 (2015).

1129   **Sorocea bonplandii** (Baill.) W.C. Burger, Lanj. & de Boer, Acta Bot. Neerl. 11: 465

1130   (1962).

1131   *Sorocea briquetii* J.F.Macbr., Publ. Field Columb. Mus., Bot. Ser. 11: 16 (1931) (incl.

1132   *S. pileate* W.C. Burger).

1133   *Sorocea carautana* M.D.M.Vianna, Carrijo & Romaniuc, Novon 19: 549 (2009).

1134   *Sorocea ganevii* R.M.Castro, Neodiversity 1: 18 (2006).

1135   *Sorocea guilleminiana* Gaudich., Voy. Bonite, Bot. 3: t. 74 (1843) (incl. *S.*

1136   *klotzschiana* Baill. and *S. macrogyna* Lanj. & Wess. Boer).

1137   *Sorocea hilarii* Gaudich., Voy. Bonite, Bot. 3: t. 71 (1843) (incl. *S. racemosa*

1138   Gaudich.).

1139   *Sorocea jaramilloi* C.C.Berg, Novon 6: 241 (1996).

1140   *Sorocea longipedicellata* A.F.P. Machado, M.D.M. Vianna & Romaniuc, Syst. Bot.

1141   38: 687 (2013).

1142   *Sorocea muriculata* Miq., C.F.P. von Martius & auct. suc. (eds.), Fl. Bras. 4(1): 113

1143   (1853).

1144   **subsp.** *muriculata* (incl. *S. amazonica* Miq.)

1145   **subsp.** *uaupensis* (Baill.) C.C. Berg, Proc. Kon. Ned. Akad. Wetensch. C 88: 387

1146   (1985) (incl. *S. guayanensis* W.C. Burger).

1147   *Sorocea pubivena* Hemsl., Biol. Cent.-Amer., Bot. 3: 150 (1883).

1148   **subsp.** *pubivena* (incl. *S. cufodontii* W.C. Burger)

1149   **subsp.** *hirtella* (Mildbr.) C.C. Berg, Novon 6: 243 (1996) (incl. *S. opima* J.F.

1150   Macbr.)

1151   **subsp.** *oligotricha* (Akkermans & C.C. Berg) C.C. Berg, Novon 6: 243 (1996).

1152   *Sorocea ruminata* C.C.Berg, Novon 6: 244 (1996).

1153   *Sorocea sarcocarpa* Lanj. & Wess. Boer, Acta Bot. Neerl. 11: 452 (1962).

1154   *Sorocea steinbachii* C.C. Berg, Proc. Kon. Ned. Akad. Wetensch. C 88: 385 (1985).

1155   *Sorocea trophoides* W.C. Burger, Acta Bot. Neerl. 11: 450 (1962).

1156

Sorocea subg. *Paraclarisia* (Ducke) W.C. Burger, Lanj. & Wess. Boer, Acta Bot. Neerl.

11: 468 (1962).

Sorocea duckei W.C.Burger, Acta Bot. Neerl. 11: 473 (1962).

Sorocea sprucei (Baill.) J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 11: 16

(1931).

subsp. *sprucei* (incl. *S. arnoldoi* Lanj. & Wess. Boer).

subsp. *saxicola* (Hassl.) C.C.Berg, Proc. Kon. Ned. Akad. Wetensch., Ser. C.,

Biol. Med. Sci. 88: 391 (1985).

Sorocea subumbellata (C.C. Berg) Cornejo, Novon 19: 297 (2009).

1166

1167

1168   *Taxotrophis*

1169 *Taxotrophis* Blume, Ann. Mus. Bot. Lugduno-Batavi 2 77 (1856); Hutch., Bull. Misc.

1170   Inform. Kew 147 (1918).

1171 *Streblus* Lour. sect. *Taxotrophis* (Blume) Corner, Gard. Bull. Singapore 19: 218 (1962);

1172   Berg et al., Fl. Males. Ser. 1, Vol. 17, Pt. 2 (2006).

1173

Monoecious or dioecious trees or shrubs, usually with lateral or terminal thorns. *Leaves*

distichous, pinnately veined, petioles adaxially pubescent. Stipules free, lateral.

Inflorescences axillary, solitary or paired, with an abaxial sterile groove, interfloral bracts

basally attached, flowers with 4 free tepals, valvate. Staminate inflorescences spicate to

sub-capitate, flowers with filaments inflexed in bud, pistillode present. Pistillate

inflorescences uniflorous or sub-spicate, flowers usually pedicellate, stigma bifid, arms

equal. *Fruits* drupaceous, up to ca. 1 cm long, usually loosely enclosed by the enlarged

persistent tepals. *Seeds* up to ca. 8 x 6 mm, cotyledons subequal to unequal.

1182

1183 Species and distribution: Six species, ranging from Sri Lanka to New Guinea.

1184

1. *Taxotrophis ilicifolia* (Kurz) S.Vidal, Revis. Pl. Vasc. Filip. 249 (1886) –

Balanostreblus ilicifolia Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 42(4): 248

(1874) – *Streblus ilicifolius* (Kurz) Corner, Gard. Bull. Singapore 19: 227 (1962).

Pseudotrophis laxiflora Warb., Bot. Jahrb. Syst. 13 294 (1891).

Taxotrophis obtusa Elmer, Leafl. Philipp. Bot. 5 1813 (1913).

Taxotrophis laxiflora Hutch., Bull. Misc. Inform. Kew 151 (1918). – *Streblus*

laxiflorus (Hutch.) Corner, Gard. Bull. Singapore 19 229 (1962).

Taxotrophis triapiculata Gamble, Bull. Misc. Inform. Kew 188 (1913).

Taxotrophis eberhardtii Gagnep., Fl. Indo-Chine 5 700 (1928).

Taxotrophis macrophylla auct. non Boerl.: Burkill, Dict. Econ. Prod. Malay Penins.

2126 (1935).

1196

2. *Taxotrophis macrophylla* (Blume) Boerl., Handl. Fl. Ned. Ind. 3: 359 (1900).

Streblus macrophyllus Blume, Ann. Mus. Bot. Lugduno-Batavi 2: 80 (1856) —

Diplocos ? macrophyllus (Blume) Bureau in A.DC., Prodr. 17: 216 (1873).

Pseudotrophis mindanaensis Warb. in Perkins, Fragm. Fl. Philipp. 1: 165 (1905);

Elmer, Leafl. Philipp. Bot. 5: 1815 (1913), '*Taxatrophis mindanaensis*' in nota.

Paratrophis caudata Merr., Philipp. J. Sci., 1, Suppl. 183 (1906).

Taxotrophis balansae Hutch., Bull. Misc. Inform. Kew 151 (1918).

Dimerocarpus brenieri Gagnep., Bull. Mus. Hist. Nat. (Paris) 27: 441 (1921).

1205

1206 **3. *Taxotrophis perakensis* (Corner) E.M. Gardner, comb. nov.**, based on *Streblus*

1207   *perakensis* Corner, Gard. Bull. Singapore 19: 223 (1962).

1208

1209 Note: Although Corner (1962) considered *S. perakensis* species part of section *Paratrophis*,

1210 albeit with some hesitation, Berg et al. (2006), whom we follow, placed it in *Taxotrophis*

1211 based on the spines that appear on some specimens. This is a rather variable species that

1212 requires further investigation to properly elucidate its limits and affinities.

1213

1214 **4. *Taxotrophis spinosa*** Steenis in Backer & Bakh.f., Fl. Java 2 (1965) 16.

1215   *Urtica spinosa* Blume, Bijdr. (1825) 507. — *Streblus spinosus* (Blume) Corner, Gard.

1216   Bull. Singapore 19: 229 (1962).

1217   *Taxotrophis javanica* Blume, Ann. Mus. Bot. Lugduno-Batavi 2: 77, t. 26 (1856).

1218

1219 **5. *Taxotrophis taxoides*** (B. Heyne ex Roth) W.L. Chew ex E.M. Gardner, comb. nov.

1220 *Trophis taxoides* B. Heyne ex Roth, Nov. Pl. Sp. 368 (1821). — *Trophis taxiformis*

1221   Spreng., Syst. Veg. 3 902 (1826), nom. nov. illeg. — *Streblus taxoides* (B. Heyne

1222   ex Roth) Kurz, Forest Fl. Burma 2: 465 (1877) — *Phyllochlamys taxoides* (B.

1223   Heyne ex Roth) Koord., Exkurs.-Fl. Java 2: 89 (1912).

1224 *Trophis spinosa* Roxb., Fl. Ind., ed. Carey 3: 762 (1832), non Willd. 1806, nec Blume

1225   1826. — *Epicarpurus spinosus* (Roxb.) Wight, Icon. Pl. Ind. Orient. 6: 7, t. 1962

1226   (1853), p.p. — *Phyllochlamys spinosa* (Roxb.) Bureau in A.DC., Prodr. 17: 218

1227   (1873);

1228 *Epicarpurus timorensis* Decne., Nouv. Ann. Mus. Hist. Nat. 3: 499, t. 21 (1834).

1229 *Taxotrophis roxburghii* Blume, Ann. Mus. Bot. Lugduno-Batavi 2: 78 (1856)

1230 *Streblus microphyllus* Kurz, Prelim. Rep. Forest Pegu App. A, cxviii; App. B, 84

1231   (1875); — *Streblus taxoides* (B. Heyne ex Roth) Kurz var. *microphylla* (Kurz)

1232   Kurz, Forest Fl. Burma 2: 465 (1877).

1233 *Phyllochlamys wallichii* King ex Hook.f., Fl. Brit. India 5: 489 (1888);

1234 *Phyllochlamys taxoides* (B. Heyne ex Roth) Koord. var. *parvifolia* Merr., Enum.

1235   Philipp. Flow. Pl. 2: 38 (1923).

1236 *Taxotrophis poilanei* Gagnep., Fl. Indo-Chine 5: 701 (1928).

1237 *Taxotrophis crenata* Gagnep., Fl. Indo-chine 5: 702, t. 82 (1928). — *Streblus crenatus*

1238   (Gagnep.) Corner, Gard. Bull. Singapore 19: 226 (1962).

1239 *Phyllochlamys tridentata* Gagnep., Fl. Indo-Chine 5: 714 (1928).

1240

1241 Note: In the 1950s, Dr. Chew Wee Lek annotated quite a lot of specimens at K and SING

1242 with the new combination *Taxotrophis taxoides*. However, the combination was never

1243 published, probably because the need for the combination was obviated in 1962 when

1244 Corner, his doctoral supervisor, reduced *Taxotrophis* to a section of *Streblus*.

1245

1246 **6. *Taxotrophis zeylanica*** (Thwaites) Thwaites, Enum. Pl. Zeyl. [Thwaites] 264 (1861).

1247   *Epicarpurus zeylanicus* Thwaites, Hooker's J. Bot. Kew Gard. Misc. 4: 1 (1852) —

1248   *Diplocos zeylanica* (Thwaites) Bureau, Prodr. [A. P. de Candolle] 17: 215 (1873)

1249   — *Streblus zeylanicus* (Thwaites) Kurz, Forest Fl. Burma 2: 464 (1877).

1250   *Taxotrophis caudata* Hutchinson, Bull. Misc. Inform. Kew 1918(4): 149 (1918).

1251

1252

1253   **Trophis**

1254

1255 *Trophis* Lour., P. Browne, Civ. Nat. Hist. Jamaica 357 (1756), nom. cons.

1256   *Bucephalon* L. Sp. Pl. 1190 (1753), nom. rejic.

1257   *Skutchia* Pax & K. Hoffm. ex C.V. Morton, J. Wash. Acad. Sci. 27: 306 (1937).

1258

Dioecious trees or shrubs. ***Leaves*** distichous, pinnately veined. Stipules free, lateral.

Inflorescences axillary or just below the leaves, solitary or paired, interfloral bracts basally

attached. Staminate inflorescences spicate to racemose with an abaxial sterile groove, tepals

4, basally connate, stamens 4, filaments inflexed in bud, pistillode present. Pistillate

inflorescences spicate to racemose or subcapitate, tepals 4, connate, forming a tubular

perianth, ovary adnate to the perianth or not, stigma bifid, arms equal. ***Fruits*** drupaceous,

up to ca. 1.5 cm long, adnate to the perianth or not, the perianth enlarged and fleshy or not.

Seeds up to ca. 1 cm long, cotyledons equal.

1267

1268 Species and distribution: Five species in the neotropics.

1269

1270 *Trophis* P. Browne sect. *Trophis*

1271

1272 **1. *Trophis cuspidata*** Lundell, Amer. Midl. Naturalist 19: 427 (1938).

1273 **2. *Trophis mexicana*** (Liebm.) Bureau, Prodr. [A. P. de Candolle] 17: 253 (1873).

1274 **3. *Trophis noraminervae*** Cuevas & Carvajal, Acta Bot. Mex. 47: 2 (1-7, fig.) (1999).

1275 **4. *Trophis racemosa*** Urb., Symb. Antill. (Urban). 4(2): 195 (1905).

1276

1277 *Trophis* P. Browne sect. *Echinocarpa* C.C. Berg, Proc. Kon. Ned. Akad. Wetensch., Ser. C,

1278 Biol. Med. Sci. 91: 353 (1988).

1279

1280   **5.** *Trophis involucrata* W.C. Burger, Phytologia 26: 432 (1973).

1281

1282

1283   **Tribe OLMEDIEAE**

1284

1285   Olmedieae Trécul, Ann. Sci. Nat., Bot. sér. 3, 8 (1847) 126

1286   Castilleae C.C. Berg, Acta Bot. Neerl. 26 (1977) 78

1287   Strebleae Bureau in DC., Prodr. 17: 215 (1873), p.p.

1288

Tree or shrubs, monoecious or dioecious (or androdioecious), mostly with self-pruning

branches. *Leaves* alternate or opposite, distichous or spirally arranged; stipules lateral to

amplexicaul. *Inflorescences* mostly unisexual, capitate, mostly discoid to urceolate,

involucrate, tepals mostly 4, connate or not. Staminate inflorescences usually many-

flowered; stamens 4 or fewer, with filaments straight or less often inflexed in bud, pistillode

mostly absent. Pistillate inflorescences one to many-flowered, ovary free or not, stigmas 2,

filiform. *Fruits* mostly drupaceous, mostly enclosed by a fleshy perianth or embedded in a

fleshy receptacle. *Seeds* with or without endosperm, testa thin, vascularized, cotyledons

mostly equal.

1298

1299   Genera and distribution: 13 genera with 63 species. Eight neotropical genera: *Castilla* (3

1300   spp.), *Helicostylis* (7), *Maquira* (4), *Naucleopsis* (22), *Olmedia* (1), *Perebea* (9),

1301   *Poulsenia* (1), and *Pseudolmedia* (9); and five Paleotropical genera: the widespread

1302   *Antiaris* (1); *Antiaropsis* (2) in New Guinea; *Mesogyne* (1) in Africa; *Sparattosyce* (1)

1303   in New Caledonia; and *Streblus* (3) in South to Southeast Asia.

1304

1305   *Olmedia*

1306

1307   *Olmedia* Ruiz & Pav., Syst. Veg. Fl. Peruv. Chil.1:257.1798. Type — *Olmedia aspera* Ruiz

1308   & Pav.

1309    *Trophis* section *Olmedia* (Ruiz & Pav.) Berg, Proc. Kon. Ned. Acad. Wetensch. C. 91:

1310     354. 1988.

1311

1312  Dioecious trees or shrubs. ***Leaves*** distichous, lamina pinnately veined. ***Stipules*** free, not

1313  fully amplexicaul. ***Inflorescences*** unisexual, with a well-developed involucre. Staminate

1314  inflorescences discoid, multiflorous; tepals 4, valvate, stamens 4, inflexed in bud, pistillode

1315  absent. Pistillate inflorescences usually uniflorous, perianth tubular, 4-dentate, ovary free,

1316  stigmas 2, equal. ***Fruits*** drupaceous, surrounded by fleshy persistent perianth and

1317  subtended by spreading, fleshy involucral bracts. ***Seeds*** ca. 5 mm long, cotyledons equal.

1318

1319  Species and distribution: One species, in the Neotropics.

1320

1321  **1.** ***Olmedia aspera*** Ruiz & Pav., Syst. Veg. Fl. Peruv. Chil.1:257.1798

1322    *Olmedia caucana* Pittier, Contr. U.S. Natl. Herb. 13:434. 1912. — *Trophis caucana*

1323     (Pitttier) C.C. Berg, Proc. Kon. Ned. Acad. Wetensch. C. 91: 354. 1988

1324    *Olmedia poeppigiana* Klotzsch, Linnaea 20: 525. 1847, as a synonym of *O. aspera*

1325     Poeppig & Endlicher, Nov. Gen. 2: 31. 1838, based on Poeppig s.n. or 1267, non

1326     *O.poeppigiana* Martius Flora (or Bot. Zeit) 24 (Beibl. 2): 93. 1841 (= *Helicostylis*

1327     *tomentosa* (Poeppig & Endlicher) Rusby).

1328    *Olmedia falcifolia* Pittier, Contr. U.S. Natl. Herb. 13: 435. 1912.

1329    *Trophis aurantiaca* Herzog, Repert. Spec. Nov. 7: 51. 1909

1330

1331

1332           ***Streblus***

1333

1334  *Streblus* Lour., Fl. Cochinch. (1790) 615.

1335  *Achymus* Juss., Dict. Sci. Nat. 1, Suppl. 31 (1816) .

1336  *Epicarpurus* Blume, Bijdr. 488 (1825).

1337  *Albrandia* Gaudich. in Freyc., Voy. Uranie, Bot. 509 (1830).

1338  *Calius* Blanco, Fl. Filip. 698 (1837).

1339   *Teonongia* Stapf, Hooker's Icon. Pl. 30: t. 2947 (1911).

1340   *Diplothorax* Gagnep., Bull. Soc. Bot. France 75 98 (1928).

1341

Trees or shrubs, dioecious or monoecious. **Leaves** distichous, lamina pinnately veined.

Stipules free, lateral. **Inflorescences** bisexual or unisexual, capitate, with a rudimentary

involucre, bracts basally attached. Staminate inflorescences discoid capitate, multiflorous;

tepals 4, imbricate, stamens 4, inflexed in bud, pistillode present but small. Pistillate

inflorescences usually uniflorous, tepals 4, ovary free, stigmas 2, equal. **Fruits** drupaceous,

up to ca. 8 mm long, initially enclosed by enlarged but not fleshy tepals, which may open

later. **Seeds** ca. 5 mm long, cotyledons equal or very unequal.

1349

1350   Species and distribution: three species from India to South China and from mainland

1351   Southeast Asia to the Philippines and the Moluccas.

1352

1353   **1.** *Streblus asper* (Retz.) Lour., Fl. Cochinch. 2: 615 (1790).

1354   *Trophis aspera* Retz., Observ. Bot. 5 (1788) – *Epicarpurus asper* (Retz.) Steud.

1355   Nomencl. Bot. ed. 2, 1: 556 (1840).

1356   *Trophis cochinchinensis* Poir., Encycl. 8 (1808) 123.

1357   *Epicarpurus orientalis* Blume, Bijdr. (1825) 488

1358   *Calius lactescens* Blanco, Fl. Filip. (1837) 698; ed. 3, 3 (1879) 1103, t. 171. —

1359   *Streblus lactescens* (Blanco) Blume, Ann. Mus. Bot. Lugduno-Batavi 2 (1856) 80.

1360   *Achymus pallens* Sol. ex Blume, Ann. Mus. Bot. Lugduno-Batavi 2 (1856) 79.

1361   *Cudrania crenata* C.H. Wright, J. Linn. Soc., Bot. 26 (1899) 469. — *Vanieria crenata*

1362   (C.H. Wright) Chun, J. Arnold Arbor. 8 (1927) 21.

1363   *Diplothorax tonkinensis* Gagnep., Bull. Soc. Bot. France 75 (1928) 98.

1364

1365   Note: Retzius's *Trophis aspera* predated Loureiro's *Streblus asper* by two years, making

1366   the latter an implied new combination under Article 41.1 of the Code (cf. Ex. 10).

1367   Synonymies of the *S. asper* have sometimes erroneously included *Trophis aculeata* Roth,

1368   which is actually *Maclura spinosa* (Roxb. ex Willd.) C.C. Berg.

1369

1370 **2.** *Streblus celebensis* C.C. Berg, Blumea 50 547 (2005).

1371

1372 **3.** *Streblus tonkinensis* (Eberh. & Dubard) Corner, Gard. Bull. Singapore 19 (1962): 228.

1373   *Bleeokrodea tonkinensis* Eberh. & Dubard, Compt. Rend. Hebd. SÈances Acad. Sci.

1374   145 (1907): 632. — *Teonongia tonkinensis* (Eberh. & Dubard) Stapf.

1375

1376

1377   **Tribe DORSTENIEAE**

1378

1379 Dorstenieae Gaudich. in Freyc., Voy. Uranie, Bot. (1830).

1380   Broussonetieae Gaudich. in Freyc., Voy. Uranie, Bot. 508 (1830).

1381 Brosimeae Trécul, Ann. Sci. Nat., Bot. sér. 3, 8: 146 (1847).

1382 Fatoueae Engl., Nat. Pflanzenfam. 3, 1: 71 (1888).

1383

Tree, shrubs, lianas, and herbs, monoecious or less often dioecious. *Leaves* alternate or less

commonly (sub)opposite, distichous or spirally arranged; stipules lateral to fully

amplexicaul. *Inflorescences* unisexual or bisexual, cymose, spicate, globose, or discoid to

turbinate or cup-shaped, multiflorous or uniflorous (the latter in pistillate inflorescences

only), bracteate or not, interfloral bracts mostly peltate. Staminate flowers with tepals (1–

)2–4 or absent, stamens 1–4 with filaments straight or inflexed in bud, pistillode present or

(more often) absent. Pistillate flowers free or connate or embedded in the receptacle, tepals

2–4, ovary free or not, stigmas 1 or 1, equal or unequal. *Fruits* drupaceous or drupe-like

due to a persistent fleshy perianth and/or receptacle, the whole endocarp unit often

ballistically ejected from the infructescence. *Seeds* with or without endosperm, large or

small; cotyledons equal or unequal.

1395

Genera and distribution: eleven genera and 63 species with a worldwide distribution:

Bleekrodea (Africa and Southeast Asia)*, Bosqueiopsis* (Africa)*, Brosimum* (Neotropics)*,

Broussonetia (Asia to Oceania; introduced worldwide)*, Dorstenia* (Africa and South

America, with one species in India)*, Fatoua* (Madagascar and Japan to New Caledonia;

introduced worldwide)*, Helianthostylis* (Neotropics)*, Malaisia* (Southeast Asia to New

Caledonia)*, Scyphosyce* (Africa)*, Sloetia* (Southeast Asia)*, Sloetiopsis* (Africa), *Trilepisium*

(Africa), *Trymatococcus* (Neotropics), *Utsetela* (Africa).

1403

1404

1405   *Sloetiopsis*

1406

1407   *Sloetiopsis* Engl., Bot. Jahrb. 39: 573 (1907).

1408   *Neosloetiopsis* Engl., Bot. Jahrb. 51: 426 (1914)

1409   *Streblus* auct. non Lour., C.C. Berg, Proc. Kon. Ned. Acad. Wetensch. C. 91: 357.

1410   1988.

1411

Tree or shrubs, dioecious (or monoecious). ***Leaves*** distichous, pinnately veined, cystoliths

present; stipules free, nearly amplexicaul. ***Inflorescences*** unisexual (or bisexual); staminate

inflorescences spicate with an abaxial sterile groove, bracts mostly peltate, tepals 4, free or

basally connate filaments inflexed in bud, pistillode small; pistillate inflorescences

uniflorous, bracts mostly basally attached, tepals 4, free, imbricate, 2 stigmas. ***Fruits***

drupaceous, fleshy endocarp dehiscent, tepals enlarged and persistent but not fleshy. ***Seeds***

globose, ca. 4 mm, endocarp coriaceous with a hard disc against the hilum, testa

vascularized with a thick apical cap, cotyledons equal.

1420

1421   **1. *Sloetiopsis usambarensis*** Engl., Bot. Jahrb. 39: 573 (1907) — *Streblus usambarensis*

1422   (Engl.) C.C. Berg, Proc. Kon. Ned. Acad. Wetensch. C. 91: 357. 1988.

1423   *Neosloetiopsis kamerunensis* Engl., Bot. Jahrb. 51: 426 (1914).

1424

1425   Note. The nearly amplexicaul stipules and occasional bisexual inflorescences reflect an

1426   affinity with the Southeast Asian *Sloetia*, also a member of the Dorstenieae.

1427

1428

1429 **Tribe PARARTOCARPEAE**

1430

Parartocarpeae N.J.C. Zerega & E.M. Gardner Zerega, Phytotaxa 388, 253–265 (2019).

Shrubs to large trees, monoecious or dioecious; abundant white exudate. *Leaves* distichous

or spirally arranged; simple; entire; pinnately veined; thin to thick coriaceous; glabrous,

pubescent, scabrid, or hispid pubescent. *Stipules* axillary, simple or paired, lateral.

Inflorescences solitary or paired in leaf axils, unisexual (or bisexual with a single apical

pistillate flower), uniflorous, cymose, or capitate with stamens or ovaries sunken into the

receptacle; pedunculate; involucre of 3–8 triangular bracts, basally connate. Staminate

inflorescences with numerous flowers, tepals 4–5, stamens 5 and normally positioned or 1–

3 in cavities in the receptacle with the anthers exerted through perforations in the upper

surface of the receptacle, filaments free or united. Pistillate inflorescences uniflorous or

(sub)globose with ovaries solitary in each cavity, unilocular, the style apical with a short

exerted stigma. *Fruits* drupaceous, enclosed by persistant tepals, or aggregated into

syncarps formed by the enlargement of the entire inflorescence.

1444

1445 Genera and distribution: Three genera and five species, from Southern China to the

1446 Solomon Islands: *Hullettia*, *Parartocarpus*, and *Pseudostreblus*.

1447

1448

1449 *Pseudostreblus*

1450

1451 *Pseudostreblus* Bureau, in A.P. de Candolle, Prodr. 17: 220 (1873).

1452 *Streblus* Lour. sect. *Pseudostreblus* (Bureau) Corner, Gard. Bull. Singapore 19: 217 (1962).

1453

Monoecious trees. *Leaves* distichous, pinnately veined. Stipules free, lateral.

Inflorescences axillary. Staminate inflorescences cymose, tepals 5, imbricate, filaments

inflexed in bud but apparently straightening gradually upon anthesis, pistillode minute,

conical, pubescent, interfloral bracts few, basally attached. Pistillate inflorescences

uniflorous, pedunculate, involucral bracts 3, ±connate, tepals 4, imbricate, stigma bifid,

arms equal. **Fruits** drupaceous, ca. 1 cm long, with tepals enlarged and loosely enclosing

the fruit. **Seeds** ellipsoid, ca. 6 x 8 mm, cotyledons unequal.

1461

1462   Species and distribution: One, in South to Southeast Asia and Southern China.

1463

1464   Note: *Pseudostreblus* is remarkable for its five-parted staminate flowers.

1465

1466   **1. *Pseudostreblus indicus* Bureau**, in A.P. de Candolle, Prodr. 17: 220 (1873).

1467   *Streblus indicus* (Bureau) Corner, Gard. Bull. Singapore 19: 226 (1962).

1468

1469

1470   **Conclusion**

1471   The revisions presented here, based on the best phylogenetic evidence available to date,

1472   provide for seven monophyletic tribes of Moraceae and ten monophyletic genera within

1473   Moreae. We hope that the resulting taxonomic and phylogenetic framework will provide a

1474   solid foundation for further research into the systematics and evolution of Moraceae.

1475

1476

Author contributions

EMG led the writing of the manuscript, did field, lab, and herbarium work, conducted all

phylogenetic analyses, and was responsible for the taxonomic treatment. MG performed lab

work for the Moraceae333 samples. RC, SD, NE, and OM performed lab work for the

PAFTOL samples, and OM curated the PAFTOL data. DA collaborated on all herbarium

work at BO. Sahromi facilitated access and participated in field collections at the Bogor

Botanical Gardens. NJCZ provided overall guidance on Moraceae and facilitated

sequencing at Northwestern University. WJB and FF acquired funding for and supervised

the PAFTOL project. AM performed herbarium sampling at K and supervised the

Moraceae portion of the PAFTOL project. EMG designed the overall study with input from

ALH and NJCZ. ALH supervised the overall study. All authors commented on and

contributed to the manuscript.

1489

1490

Acknowledgements

This work was supported by the United States National Science Foundation (DBI award

number 1711391, and DEB award numbers 0919119 and 1501373); the SYNTHESYS

Project http://www.synthesys.info/ which is financed by European Community Research

Infrastructure Action under the FP7 "Capacities" Program"; the National Parks Board of

Singapore through a *Flora of Singapore* Fellowship and a Research Fellowship; grants

from the Calleva Foundation and the Sackler Trust to the Plant & Fungal Trees of Life

Project at the Royal Botanic Gardens, Kew. We thank the Pritzker Laboratory for

Molecular Systematics at the Field Museum of Natural History (K. Feldheim) for access to

laboratory and sequencing facilities; S. Knapp and J. Wajer (BM) for helpful consultation

on nomenclatural issues; W. Clement for helpful comments on the circumscription of the

Olmedieae; the Bogor Botanical Gardens, Singapore Botanic Gardens, and The Morton

Arboretum for access to living collections; and the following herbaria for access to

collections for examination and/or sampling: BM, BKF, BO, CLM, F, IBSC, K, FTBG, L,

MO, NY, SING, SAN, SAR, SNP, and U. Finally, we gratefully acknowledge the careful

work of past scholars of Moraceae, in particular the late C.C. Berg and E.J.H. Corner.

1507

1508 **Conflicts of interest**

1509 The authors are unaware of any conflicts of interest affecting this research.

1510

1511 **Literature cited**

1512

Bawa, K., & Crisp, J. 1980. Wind-pollination in the understorey of a rain forest in Costa

Rica. *J. Ecol.*, 68: 871–876. Retrieved from

http://www.jstor.org/stable/10.2307/2259462

Bawa, K. S., Bullock, S. H., Perry, D. R., Coville, R. E., & Grayum, M. H. 1985.

Reproductive biology of tropical lowland rain forest trees. II. Pollination systems. *Am.*

J. Bot., 72: 346–356. https://doi.org/10.2307/2443527

Berg, C. C. 1977a. Revisions of African Moraceae (excluding Dorstenia, Ficus, Musanga
and Myrianthus). *Bull. Du Jard. Bot. Natl. Belgique*, 47: 267–407. Retrieved from
http://www.jstor.org/stable/10.2307/3667908

Berg, C. C. 1977b. The Castilleae, a tribe of the Moraceae, rename and redefined due tothe
exclusion of the type genus Olmedia from the 'Olmedieae'. *Acta Bot. Neerl.*, 26: 73–
82.

Berg, C. C. 1978. Cecropiaceae: A new family of the Urticales. *Taxon*, 27: 39–44.
https://doi.org/10.2307/1220477

Berg, C. C. 1988. The genera Trophis and Streblus (Moraceae) remodelled. *Proc. K. Ned.*
Akad. van Wet. Ser. C, Biol. Med. Sci., 91: 345–362.

Berg, C. C. 2001. *Flora Neotropica Monograph 83: Moreae, Artocarpeae, and Dorstenia*
(Moraceae), with Introductions to the Family and Ficus and with Additions and
Corrections to Flora Neotropica Monograph 7 (Vol. 83).

Berg, C. C. 2005. Flora Malesiana precursor for the treatment of Moraceae 8: other genera
than Ficus. *Blumea*, 50: 535–550. Retrieved from
http://www.ingentaconnect.com/content/nhn/blumea/2005/00000050/00000003/art000
16

Berg, C. C., Corner, E. J. H., & Jarrett, F. M. 2006. Moraceae (genera other than Ficus).
Flora Malesiana, Ser. 1, 17: 1–154.

Bolger, A. M., Lohse, M., & Usadel, B. 2014. Trimmomatic: A flexible trimmer for
Illumina Sequence Data. *Bioinformatics*, 1–7.
https://doi.org/10.1093/bioinformatics/btu170

Brewer, G. E., Clarkson, J. J., Maurin, O., Zuntini, A. R., Barber, V., Bellot, S., Biggs,
N., Cowan, R. S., Davies, N. M. J., Dodsworth, S., & others. 2019. Factors affecting
targeted sequencing of 353 nuclear genes from herbarium specimens spanning the
diversity of angiosperms. *Front. Plant Sci.*, 10: 1102.

Bureau, E. 1873. Moraceae and Artocarpace. in: *A. P. de Candolle, Prodromussystematis*
naturali 17. Paris: Treuttel & Würtz.

Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. 2009. trimAl: A tool for
automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*,

25: 1972–1973. https://doi.org/10.1093/bioinformatics/btp348

Chen, Z., Yang, T., & Lu, L. 2016. Tree of life for the genera of Chinese vascular plants.

J. Syst. Evol., 54: 273–276. https://doi.org/10.1111/jse.12219

Chung, K. F., Kuo, W. H., Hsu, Y. H., Li, Y. H., Rubite, R. R., & Xu, W. Bin. 2017.

Molecular recircumscription of Broussonetia (Moraceae) and the identity and

taxonomic status of B. kaempferi var. australis. *Bot. Stud.*

https://doi.org/10.1186/s40529-017-0165-y

Clement, W. L., & Weiblen, G. D. 2009. Morphological Evolution in the Mulberry

Family (Moraceae). *Syst. Bot.*, 34: 530–552.

https://doi.org/10.1600/036364409789271155

Corner, E. J. H. 1962. The Classification of Moraceae. *Gard. Bull.*, 19: 187–252.

Corner, E. J. H. 1970. New Species of Streblus and Ficus. *Blumea*, 18: 393–411.

Corner, E. J. H. 1975. The evolution of StrebIus Lour. (Moraceae): with a new species of

sect. Bleekrodea. *Phytomorphology*, 25: 1–12.

Culley, T. M., Weller, S. G., & Sakai, A. K. 2002. The evolution of wind pollination in

angiosperms. *Trends Ecol. Evol.*, 17: 361–369. https://doi.org/10.1016/S0169-

5347(02)02540-5

Datwyler, S. L., & Weiblen, G. D. 2004. THE ORIGIN OF THE FIG : PHYLOGENETIC

NDH F SEQUENCES. *Am. J. Bot.*, 91: 767–777.

Doyle, J., & Doyle, J. 1987. Genomic plant DNA preparation from fresh tissue-CTAB

method. *Phytochem. Bull.*, 19: 11–15.

Engler, G. H. A. 1889. Ulmaceae, Moraceae, and Urticaceae. in: *Engler & Prantl, Die*

natürlichen Pflanzenfamilien 3(1). Leipzig: W. Engelmann.

Filho, M. D. M. V., Carrijo, T. T., & Neto, S. R. 2009. Sorocea carautana (Moraceae): A

New Species from Southeastern Brazil. *Novon A J. Bot. Nomencl.*

https://doi.org/10.3417/2007143

FitzJohn, R. G. 2012. Diversitree : comparative phylogenetic analyses of diversification in

R. *Methods Ecol. Evol.*, 3: 1084–1092. https://doi.org/10.1111/j.2041-

210X.2012.00234.x

Gardner, E. M. 2017. *Dissertation: Evolutionary Transitions: Phylogenomics and*

Pollination of Artocarpus (Moraceae).

https://doi.org/https://doi.org/10.21985/N2CF59

Gardner, E. M., Gagné, R. J., Kendra, P. E., Montgomery, W. S., Raguso, R. A., McNeil, T. T., & Zerega, N. J. C. 2018. A flower in fruit's clothing: Pollination of jackfruit (artocarpus heterophyllus, moraceae) by a new species of gall midge, clinodiplosis ultracrepidata sp. Nov. (Diptera: Cecidomyiidae). *Int. J. Plant Sci.*, 179: 350–367. https://doi.org/10.1086/697115

Gardner, E. M., Johnson, M. G., Ragone, D., Wickett, N. J., & Zerega, N. J. C. 2016. Low-Coverage, Whole-Genome Sequencing of Artocarpus camansi (Moraceae) for Phylogenetic Marker Development and Gene Discovery. *Appl. Plant Sci.*, 4: 1600017. https://doi.org/10.3732/apps.1600017

Hale, H., Gardner, E. M ., Viruel, J., Pokorny, L. ., & Johnson, M. G. 2020. Strategies for reducing per-sample costs in target capture sequencing for phylogenomics and population genomics in plants. *Appl. Plant Sci.*, in press.

Heibl, C. 2008. PHYLOCH: R language tree plotting tools and interfaces to diverse phylogenetic software packages. http://www.christophheibl.de/Rpackages.html.

Johnson, M. G., Gardner, E. M., Liu, Y., Medina, R., Goffinet, B., Shaw, A. J., Zerega, N. J. C., & Wickett, N. J. 2016. HybPiper: Extracting Coding Sequence and Introns for Phylogenetics from High-Throughput Sequencing Reads Using Target Enrichment. *Appl. Plant Sci.*, 4: 1600016. https://doi.org/10.3732/apps.1600016

Johnson, M. G., Pokorny, L., Dodsworth, S., Botigué, L. R., Cowan, R. S., Devault, A., Eiserhardt, W. L., Epitawalage, N., Forest, F., Kim, J. T., Leebens-Mack, J. H., Leitch, I. J., Maurin, O., Soltis, D. E., Soltis, P. S., Wong, G. K., Baker, W. J., & Wickett, N. J. 2019. A Universal Probe Set for Targeted Sequencing of 353 Nuclear Genes from Any Flowering Plant Designed Using k-Medoids Clustering. *Syst. Biol.*, 68: 594–606. https://doi.org/10.1093/sysbio/syy086

Kamneva, O. K., Syring, J., Liston, A., & Rosenberg, N. A. 2017. Evaluating allopolyploid origins in strawberries (Fragaria) using haplotypes generated from target capture sequencing. *BMC Evol. Biol.*, 17: 1–19. https://doi.org/10.1186/s12862-017-1019-7

Kates, H. R., Johnson, M. G., Gardner, E. M., Zerega, N. J. C., & Wickett, N. J. 2018.

Allele phasing has minimal impact on phylogenetic reconstruction from targeted

nuclear gene sequences in a case study of Artocarpus. *Am. J. Bot.*

https://doi.org/10.1002/ajb2.1068

Katoh, K., & Standley, D. M. 2013. MAFFT multiple sequence alignment software

version 7: improvements in performance and usability. *Mol. Biol. Evol.*, 30: 772–80.

https://doi.org/10.1093/molbev/mst010

Kim, J., & Sanderson, M. J. 2008. Penalized likelihood phylogenetic inference: Bridging

the parsimony-likelihood gap. *Syst. Biol.* https://doi.org/10.1080/10635150802422274

Kress, W. J., Erickson, D. L., Jones, F. A., Swenson, N. G., Perez, R., Sanjur, O., &

Bermingham, E. 2009. Plant DNA barcodes and a community phylogeny of a tropical

forest dynamics plot in Panama. *Proc. Natl. Acad. Sci. U. S. A.*, 106: 18621–18626.

https://doi.org/10.1073/pnas.0909820106

Lewis, W. H. 1986. Airborne pollen of the Neotropics:Potential roles in pollination and

pollinosis. *Grana*, 25: 75–83. https://doi.org/10.1080/00173138609429936

Machado, A. F. P., Vianna Filho, M. D. M., Amorim, A. M., & Romaniuc Neto, S.

2013. A New *Sorocea* (Moraceae) From The Atlantic Rainforest, Brazil. *Syst.*

Bot. https://doi.org/10.1600/036364413x670340

Momose, K., Yamaoka, R., & Inoue, T. 1998. Pollination Biology of the Genus

Artocarpus, Moraceae. *Tropics*, 7: 165–172. https://doi.org/10.3759/tropics.7.165

Nepal, M. 2012. Phylogenetics of Morus (Moraceae) Inferred from ITS and trnL-trnF

Sequence Data. *Syst. Bot.*, 37: 442–450. https://doi.org/10.1600/036364412X635485

Paradis, E. 2013. Molecular dating of phylogenies by likelihood methods: A comparison

of models and a new information criterion. *Mol. Phylogenet. Evol.*

https://doi.org/10.1016/j.ympev.2013.02.008

Paradis, E., & Schliep, K. 2019. Ape 5.0: An environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics*, 35: 526–528.

https://doi.org/10.1093/bioinformatics/bty633

Quinlan, A. R., & Hall, I. M. 2010. BEDTools: A flexible suite of utilities for comparing

genomic features. *Bioinformatics*, 26: 841–842.

https://doi.org/10.1093/bioinformatics/btq033

Rabosky, D. L. 2014. Automatic detection of key innovations, rate shifts, and diversity-

dependence on phylogenetic trees. *PLoS One*, 9.

https://doi.org/10.1371/journal.pone.0089543

Rabosky, D. L., Donnellan, S. C., Grundler, M., & Lovette, I. J. 2014a. Analysis and

visualization of complex Macroevolutionary dynamics: An example from Australian

Scincid lizards. *Syst. Biol.*, 63: 610–627. https://doi.org/10.1093/sysbio/syu025

Rabosky, D. L., Grundler, M., Anderson, C., Title, P., Shi, J. J., Brown, J. W., Huang,

H., & Larson, J. G. 2014b. BAMMtools: An R package for the analysis of

evolutionary dynamics on phylogenetic trees. *Methods Ecol. Evol.*, 5: 701–707.

https://doi.org/10.1111/2041-210X.12199

Rabosky, D. L., Santini, F., Eastman, J., Smith, S. A., Sidlauskas, B., Chang, J., &

Alfaro, M. E. 2013. Rates of speciation and morphological evolution are correlated

across the largest vertebrate radiation. *Nat. Commun.*, 4: 1958.

https://doi.org/10.1038/ncomms2958

Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and

other things). *Methods Ecol. Evol.*, 3: 217–223. https://doi.org/10.1111/j.2041-

210X.2011.00169.x

Sakai, S., Kato, M., & Nagamasu, H. 2000. Artocarpus (Moraceae)–gall midge

pollination mutualism mediated by a male-flower parasitic fungus. *Am. J. Bot.*, 87:

440–445. Retrieved from http://www.amjbot.org/content/87/3/440.short

Sanderson, M. J. 2002. Estimating absolute rates of molecular evolution and divergence

times: A penalized likelihood approach. *Mol. Biol. Evol.*

https://doi.org/10.1093/oxfordjournals.molbev.a003974

Santos, A. dos, & Neto, S. R. 2015. Sorocea angustifolia (Moraceae), a New Species of

Sorocea from the Atlantic Forest in Brazil . *Novon A J. Bot. Nomencl.*

https://doi.org/10.3417/2013022

Sayyari, E., & Mirarab, S. 2018. Testing for polytomies in phylogenetic species trees

using quartet frequencies. *Genes (Basel).* https://doi.org/10.3390/genes9030132

Schliep, K. P. 2011. phangorn: Phylogenetic analysis in R. *Bioinformatics*, 27: 592–593.

https://doi.org/10.1093/bioinformatics/btq706

Stamatakis, A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-

analysis of large phylogenies. *Bioinformatics*, 30: 1312–1313.

https://doi.org/10.1093/bioinformatics/btu033

Tange, O. 2018. GNU Parallel 2018. *Https://Doi.Org/10.5281/Zenodo.1146014*.

https://doi.org/https://doi.org/10.5281/zenodo.1146014

Team, R. C. 2019. R: A Language and Environment for Statistical Computing. *Vienna,*

Austria.

Villaverde, T., Pokorny, L., Olsson, S., Rincón-Barrado, M., Johnson, M. G.,

Gardner, E. M., Wickett, N. J., Molero, J., Riina, R., & Sanmartín, I. 2018.

Bridging the micro- and macroevolutionary levels in phylogenomics: Hyb-Seq solves

relationships from populations to species and above. *New Phytol.*

https://doi.org/10.1111/nph.15312

Weitemier, K., Straub, S. C. K., Cronn, R. C., Fishbein, M., Schmickl, R., McDonnell,

A., & Liston, A. 2014. Hyb-Seq: Combining Target Enrichment and Genome

Skimming for Plant Phylogenomics. *Appl. Plant Sci.*, 2: 1400042.

https://doi.org/10.3732/apps.1400042

Wen, D., Yu, Y., Zhu, J., & Nakhleh, L. 2018. Inferring phylogenetic networks using

PhyloNet. *Syst. Biol.*, 67: 735–740. https://doi.org/10.1093/sysbio/syy015

Yu, Y., & Nakhleh, L. 2015. A maximum pseudo-likelihood approach for phylogenetic

networks. *BMC Genomics*, 16: 1–10. https://doi.org/10.1186/1471-2164-16-S10-S10

Zapata, T. R., & Arroyo, M. T. K. 1978. Plant Reproductive Ecology of a Secondary

Deciduous Tropical Forest in Venezuela. *Biotropica*, 10: 221.

https://doi.org/10.2307/2387907

Zerega, N. J. C., Clement, W. L., Datwyler, S. L., & Weiblen, G. D. 2005.

Biogeography and divergence times in the mulberry family (Moraceae). *Mol.*

Phylogenet. Evol., 37: 402–16. https://doi.org/10.1016/j.ympev.2005.07.004

Zerega, N. J. C., & Gardner, E. M. 2019. Delimitation of the new tribe Parartocarpeae

(Moraceae) is supported by a 333-gene phylogeny and resolves tribal level Moraceae

taxonomy. *Phytotaxa*, 388: 253–265.

1699 **Zerega, N. J. C., Nur Supardi, M. N., & Motley, T. J.** 2010. Phylogeny and

1700  Recircumscription of Artocarpeae (Moraceae) with a Focus on Artocarpus. *Syst. Bot.*,

1701  35: 766–782. https://doi.org/10.1600/036364410X539853

1702 **Zhang, C., Sayyari, E., & Mirarab, S.** 2017. ASTRAL-III: Increased Scalability and

1703  Impacts of Contracting Low Support Branches BT - Comparative Genomics: 15th

1704  International Workshop, RECOMB CG 2017, Barcelona, Spain, October 4-6, 2017,

1705  Proceedings. in: J. Meidanis & L. Nakhleh (eds.). Cham: Springer International

1706  Publishing. https://doi.org/10.1007/978-3-319-67979-2_4

1707 **Zhang, Q., Onstein, R. E., Little, S. A., & Sauquet, H.** 2019. Estimating divergence

1708  times and ancestral breeding systems in Ficus and Moraceae. *Ann. Bot.*, 123: 191–204.

1709  https://doi.org/10.1093/aob/mcy159

1710

1711

1712

1713 **Tables**

1714

1715 Table 1. Overview of the taxonomic history of Moreae.

Engler (1899)	Corner (1962)	Berg (2001)	Clement & Weiblen (2009)	Gardner et al.
				*Afromorus*c
Ampalis	*Ampalis*	(*Ampalis*)a	(*Ampalis*)a	*Ampalis*
			Bagassa	*Bagassa*
	(*Bleekrodea*)a	*Bleekrodea*		
		Broussonetia		
	Clarisia			
	Fatoua	*Fatoua*		
		Maclura		
	(*Maillardia*)b	(*Maillardia*)b	(*Maillardia*)b	*Maillardia*
		(*Malaisia*)b		
		Milicia	*Milicia*	*Milicia*
Morus	*Morus*	*Morus*	*Morus*	*Morus*
Pachytrophe	*Pachytrophe*	(*Pachytrophe*)a	(*Pachytrophe*)a	(*Pachytrophe*)d
Paratrophis	(*Paratrophis*)a	(*Paratrophis*)a	(*Paratrophis*)a	*Paratrophis*
Pseudomorus	(*Pseudomorus*)a	(*Pseudomorus*)a	(*Pseudomorus*)a	(*Pseudomorus*)e
	(*Pseudostreblus*)a	(*Pseudostreblus*)a	(*Pseudostreblus*)a	
	(*Sloetia*)	(*Sloetia*)a		
	(*Sloetiopsis*)a	(*Sloetiopsis*)a	(*Sloetiopsis*)a	
	Sorocea		*Sorocea*	*Sorocea*
	Streblus	*Streblus*	*Streblus*	
	(*Taxotrophis*)a	(*Taxotrophis*)a	(*Taxotrophis*)a	*Taxotrophis*
Trophis	*Trophis*	*Trophis*	*Trophis*	*Trophis*

1716 a Included in *Streblus*.

1717 b Included in *Trophis*.

1718 c Newly separated from *Morus*.

1719 d Included in *Ampalis*

1720   e Included in *Paratrophis*.

1721

1722

1723   Table 2. Divergence times (in Ma) estimated using penalized likelihood under correlated

1724   and relaxed models. Nomenclature follows the revisions proposed in this study.

Clade	Correlated	Relaxed
Moraceae	84.7	84.7
Parartocarpeae	75.3	74.7
Ficeae	24	34.5
Olmedieae	36.4	43.4
Dorstenieae	72.5	76.9
Maclureae	44.9	52.1
Artocarpeae	64	64
Moreae	59.1	75.4

1725

1726

1727

1728

1729

1730

1731   **Figures**

1732

Figure 1. Representatives of Moreae: *Paratrophis pendulina* (=*Streblus pendulinus*) (a)

infructescences and (b) staminate inflorescences, the latter showing the characteristic sterile

groove; *Sorocea racemosa* (c) infructescences and (d) staminate inflorescence; *Morus nigra*

(e) pistillate inflorescences with elaborate stigmas and (f) staminate inflorescences with

stamens that, having been inflexed in bud, are substantially longer than the perianth itself;

and a newly-placed member of Olmedieae (=Castilleae), *Streblus asper:* (g) infructescences

and (h) discoid-capitate staminate inflorescences with the rudiments of an involucre visible

below the unopened flowers. Photo credits: (a) F. & Kim Starr, under a CC-BY-3.0-US licence

(https://commons.wikimedia.org/wiki/File:Starr-051029-5102-Streblus_pendulinus-fruit-Auwahi-

Maui_(24481336159).jpg); (b) M. Marathon, CC-BY-SA-4.0

(https://commons.wikimedia.org/wiki/File:Streblus_brunonianus_flowers.jpg); (c)-(d) A. Popovkin, CC-BY-

2.0 (https://commons.wikimedia.org/wiki/File:Sorocea_racemosa_Gaudich._-_Flickr_-

_Alex_Popovkin,_Bahia,_Brazil_(3).jpg and

https://commons.wikimedia.org/wiki/File:Sorocea_racemosa_Gaudich._-_Flickr_-

_Alex_Popovkin,_Bahia,_Brazil_(9).jpg); (e) E. Gardner; (f) Schurdl, CC-BY-SA-4.0

(https://commons.wikimedia.org/wiki/File:Morus_nigra_100525_02.jpg); (g) D. Valke, CC-BY-SA-2.0

(https://commons.wikimedia.org/wiki/File:Bekar_(Konkani-_बेकर)_(4533519363).jpg); (h) Vinayaraj, CC

BY-SA 4.0 (https://commons.wikimedia.org/wiki/File:Streblus_asper_at_Panamaram_(5).jpg)

1751

1752

1768   Figure 2. Phylogenetic trees from the "exon" dataset. Maximum-likelihood based on a

1769   supermatrix of all loci, with bootstrap support and previous nomenclature (left), and a

1770   species tree based on gene trees from all loci with bootstrap/LPP support and revised

1771   nomenclature (right). Discordant branches are colored in red, and tribal classifications are

1772   shaded without (left) and with (right) the revised classifications presented here.

1773

1774

supermatrix

species tree

1775   Figure 3. Phylogenetic trees of the Moreae clade from the "supercontig" dataset.

1776   Maximum-likelihood based on a supermatrix of all loci, with bootstrap support and

1777   previous nomenclature (left), and a species tree based on gene trees from all loci with

1778   bootstrap/LPP support and revised nomenclature (right). Discordant branches are colored in

1779   red.

1780

1781

supermatrix

species tree

1782   Figure 4. The best five maximum-pseudo-likelihood phylogenetic networks for the

1783   *Paratrophis* clade.

1784

1785

1786   Figure 5. Time-calibrated phylogenetic tree, with revised nomenclature.

1787

1788

1789   Figure 6. Ancestral reconstruction of stamen position, with revised nomenclature. The

1790   reconstruction was identical under a trait-dependent (BiSSE) or a trait-independent model.

1791   Blue = inflexed in bud; yellow = straight in bud.

1792

1793

Pseudostreblus indica NewmanLA0518
Hulletia dumosa NZ242
Hullettia griffithiana Kerr16886
Pararartocarpus venenosus NZ874
Pararartocarpus bracteatus NZ730
Ficus sagittifolia ChaseMW19852
Ficus racemosa SRR1405699 SRR1405700
Ficus macrophylla EG30
Streblus asper EG696
Antiaropsis decipiens NZ281
Sparattosyce dioica WeiblenWW1223
Poulsenia armata Pennington14600
Helicostylis tomentosa Gomes511
Castilla elastica ChaseMW19850
Olmedia aspera Fuentes5323 S41
Pseudolmedia spuria Lobo263
Naucleopsis macrophylla BergP18534
Antiaris toxicaria 5595
Fatoua pilosa TaylorP218
Malaisia scandens EG122
Broussonetia papyrifera SRR1477753
Broussonetia cf papyrifera Heng14289
Broussonetia kazinoki ChaseMW17827
Bleekrodea madagascariensis DavisRakotonasolo3117
Sloetiopsis usambarensis Faden77 417
Brosimum alicastrum EG23
Helianthostylis sprucei Ribeiro1204
Trymatococcus amazonicus Prance13461
Bosqueiopsis gilletii Fronties243
Trilepisium madagascariense Richard403
Utsetela gabonensis Breteler14086
Utsetela neglecta Bissiengen923
Dorstenia hildebrandtii NZ311
Dorstenia barteri ChaseMW18153
Maclura tinctoria tinctoria Campos2679
Maclura africana GreenwayKanuri15269
Maclura andamanica Maxwell89−436
Maclura spinosa Perumal RHT22554
Maclura pomifera EG139
Maclura brasiliensis NeeVargas45025
Maclura thorellii Pierre4709
Maclura cochinchinensis SAGBE887
Maclura aff pubsecens Averyanov1198
Maclura tricuspidata MOR 68−7917
Maclura fruticosa NMCuong1193
Maclura cf cochinchinensis NZ757
Maclura amboinensis Takeuchi16578
Taxotrophis zeylanica Meijer701
Taxotrophis macrophylla DDS10673
Taxotrophis ilicifolius EG720
Taxotrophis spinosa EG702
Taxotrophis taxoides EG701
Maillardia borbonica Cadet4518
Maillardia montana Andrianantoanina 1023
Afromorus mesozygia Buckner309
Milicia excelsa McPherson16087
Milicia regia Cooper332
Pachytrophe dimepate CasmirZ405
Ampalis mauritiana Hoffman
Paratrophis microphylla Beecher sn
Paratrophis anthropophagorum Degeer4698
Paratrophis insignis Vasquez28068
Paratrophis pendulina Gray3277
Paratrophis glabra EG78
Paratrophis philippinensis EG430
Trophis involucrata Aguilar4679
Trophis racemosa Stevens37196
Trophis mexicana Stevens27939
Trophis cuspidata Gomez−Dominuez774
Morus notabilis SRR8138828
Morus rubra EG141
Morus microphylla Fishbein1058
Morus celtidifolia Sandoval637
Morus macroura EG28
Morus serrata Koelz4788
Morus australis MOR 241−716
Morus kagayamai AA20187−A
Morus cathayana Nie92144
Morus nigra EG29
Morus mongolica MOR 55−951
Morus alba MOR 920−261
Bagassa guianensis GW1677
Sorocea subumbellata Dodson9601
Sorocea duckei Stein 3931
Sorocea sprucei saxicola Nee35729
Sorocea trophoides Fuentes304
Sorocea affinis Espinosa706
Sorocea pubivena pubivena Galdames4477
Sorocea pubivena oligotricha Kennedy2124
Sorocea jaramilloi Madison4772
Sorocea muriculata muriculata Schunke16489
Sorocea pubivena hirtella Vasquez19236
Sorocea briquetii Valenzula954
Sorocea muriculata uaupensis Plowman sn
Sorocea steinbachii GW1501
Sorocea boplandii Tressens6347
Sorocea hilarii Carauta795
Sorocea guilleminiana Moya112 S38
Clarisia ilicifolia Carranza63386
Batocarpus amazonicus Salidas1233
Batocarpus costaricensis GW1463
Batocarpus oriniceros Vasquez27440
Clarisia racemosa Assuncao690
Clarisia biflora GW1460
Artocarpus scandens EG411
Artocarpus papuanus NZ61
Artocarpus limpato NZ609
Artocarpus altissimus EG441
Artocarpus sepicanus GW1701
Artocarpus longifolius ssp adpressus Nangkat15511
Artocarpus fulvicortex EG711
Artocarpus nitidus ssp lingnanensis NZ911
Artocarpus thailandicus NZ402
Artocarpus lacucha NZ420
Artocarpus gomezianus ssp zeylanicus NZ956
Artocarpus nitidus ssp humilis EG258
Artocarpus nitidus ssp griffithii NZ216
Artocarpus dadah NZ694
Artocarpus tonkinensis EG174
Artocarpus styracifolius EG176
Artocarpus hypargyraeus EG170
Artocarpus nanchuanensis EGL238
Artocarpus petelotii DDS14435
Artocarpus pithecogallus LiJianwu3200
Artocarpus gongshanensis HAST137747
Artocarpus gomezianus ssp gomezianus NZ533
Artocarpus cf lacucha NZ517
Artocarpus glaucus NZ852
Artocarpus nitidus ssp borneensis NZ686
Artocarpus tomentosulus NZ617
Artocarpus primackii NZ687
Artocarpus xanthocarpus Yang15648
Artocarpus vriesianus var vrieseanus GW1229
Artocarpus vrieseanus refractus Hoogland4822
Artocarpus albobrunneus AA2243
Artocarpus fretessii NZ929
Artocarpus ovatus NZ202
Artocarpus nitidus ssp nitidus PPI10374
Artocarpus rubrovenius JSB84
Artocarpus subrotundifolius Wenzel1576
Artocarpus heterophyllus EG98
Artocarpus annulatus NZ985
Artocarpus integer NZ918
Artocarpus sarawakensis EG527
Artocarpus brevipedunculatus NZ814
Artocarpus anisophyllus NZ606
Artocarpus lanceifolius ssp clementis NZ739
Artocarpus hirsutus NZ953
Artocarpus nobilis AHJ3283
Artocarpus chama NZ354
Artocarpus melinoxylus JLS2924
Artocarpus odoratissimus NZ618
Artocarpus rigidus ssp asperulus NZ507
Artocarpus hispidus NZ258
Artocarpus rigidus NZ728
Artocarpus treculianus nigrescens Ramos2107
Artocarpus pinnatisectus Escritor
Artocarpus multifidus PPI3911
Artocarpus treculianus Elmer12468
Artocarpus treculianus ovatifolius NZ203
Artocarpus blancoi Ramos42018
Artocarpus aff excelsus AveryanovVH1819
Artocarpus camansi EG149
Artocarpus mariannensis DD4
Artocarpus altilis K7 UluFiti
Artocarpus horridus EG437
Artocarpus cf horridus RM160
Artocarpus excelsus NZ780
Artocarpus obtusus NZ729
Artocarpus lowii MWL2
Artocarpus teijsmannii NZ946
Artocarpus sumatranus AA2766
Artocarpus kemando NZ612
Artocarpus maingayi NZ257
Artocarpus tamaran EG92
Artocarpus sericicarpus NZ771
Artocarpus elasticus EG87
Artocarpus scortechinii NZ209
Artocarpus corneri Fuchs21347
Artocarpus jarrettiae SAN120933

1794   Figure 7. Revised classification of Moreae on a maximum-likelihood phylogenetic tree of

1795   Moreae based on "supercontig" sequences.

1796

1797

1779   Figure 7. Revised classification of Moreae on a strict consensus tree of all four main

1780   phylogenomic analyses.

1781

1782