**Repeated parallel losses of inflexed stamens in Moraceae: phylogenomics and generic**

**revision of the tribe Moreae and the reinstatement of the tribe Olmedieae (Moraceae)**

Elliot M. Gardner[1,2,3,*], Mira Garner[1,4], Robyn Cowan[5], Steven Dodsworth[5,6], Niroshini

Epitawalage[5], Olivier Maurin[5], Deby Arifiani[7], S. Sahromi[8], William J. Baker[5], Felix

Forest[5], Nyree J.C. Zerega[9,10], Alexandre Monro[5], and Andrew L. Hipp[1,11]

[1] The Morton Arboretum, 4100 IL-53, Lisle, Illinois 60532, USA

[2] Case Western Reserve University, Department of Biology, 2080 Adelbert Road,

Cleveland, Ohio 44106, USA (current affiliation)

[3] Singapore Botanic Gardens, National Parks Board, 1 Cluny Road, 259569, Singapore

(current affiliation)

[4] University of British Columbia, Faculty of Forestry, 2424 Main Mall, Vancouver, British

Columbia, V6T 1Z4, Canada (current affiliation)

[5] Royal Botanic Gardens, Kew, Richmond, TW9 3AE, United Kingdom

[6] School of Life Sciences, University of Bedfordshire, Luton LU1 3JU, United Kingdom

(current affiliation)

[7] Herbarium Bogoriense, Research Center for Biology, Indonesian Institute of Sciences,

Cibinong, Jawa Barat, Indonesia

[8] Center for Plant Conservation Botanic Gardens, Indonesian Institute of Sciences, Bogor,

Jawa Barat, Indonesia

[9] Chicago Botanic Garden, Plant Science and Conservation, 1000 Lake Cook Road,

Glencoe, IL, 60022, USA

[10] Northwestern University, Plant Biology and Conservation Program, 2205 Tech Dr.,

Evanston, IL, 60208, USA

[11] The Field Museum, 1400 S Lake Shore Dr., Chicago, IL 60605, USA

[a] For correspondence, email elliot.gardner@case.edu

**Abstract**

We present a densely-sampled phylogenomic study of the mulberry tribe (Moreae,

Moraceae), an economically important clade with a global distribution, revealing multiple

losses of inflexed stamens, a character traditionally used to circumscribe Moreae. Inflexed

stamens facilitate ballistic pollen release and are associated with wind pollination, and the

results presented here suggest that losses of this character state may have evolved

repeatedly in Moraceae. Neither Moreae nor several of its major genera (*Morus*, *Streblus*,

*Trophis*) were found to be monophyletic. A revised system for a monophyletic Moreae is

presented, including the reinstatement of the genera *Ampalis, Maillardia, Pachytrophe,*

*Taxotrophis,* and *Paratrophis*, and the recognition of the new genus *Afromorus*, based on

*Morus* subgenus *Afromorus*. *Pseudostreblus* is reinstated and transferred to the

Parartocarpeae, and *Sloetiopsis* is reinstated and transferred to the Dorstenieae. The tribe

Olmediae is reinstated, replacing the Castilleae, owing to the reinstatement of the type

genus *Olmedia,* and its exclusion from Moreae. *Streblus* s.s. is excluded from Moreae and

transferred to the Olmediae, which is characterized primarily by involucrate inflorescences

without regard to stamen position. Eight new combinations are made.

**Keywords:** Moraceae, mulberry family; Moreae, Olmedieae, Castilleae, Parartocarpeae,

*Afromorus, Ampalis, Bagassa, Maillardia, Milicia, Morus, Olmedia; Pachytrophe,*

*Paratrophis, Pseudostreblus, Sloetiopsis; Sorocea, Streblus, Taxotrophis, Trophis*.

**Introduction**

The preservation of plesiomorphic (ancestral) characters can result in species that

are similar in appearance but distantly related, connected only by a remote common

ancestor. The mulberry family (Moraceae Gaudich., seven tribes, ca. 39 genera and 1,200

species) illustrates this principle well. Inflexed stamens in bud—an adaptation to wind

pollination that allows explosive pollen dispersal when flowers open—were traditionally

used to define a tribe of the family, the Moreae (mulberries and their allies). Yet

phylogenetic analyses have revealed that inflexed stamens, an ancestral feature of both

Moraceae and their sister family Urticaceae (nettles), have been lost repeatedly (Clement &

Weiblen, 2009). Thus, for example, Cecropiaceae C.C. Berg, traditionally distinguished

from the nettle family by the absence of inflexed stamens, is in fact embedded within the

Urticaceae Juss. (Berg, 1978; Clement & Weiblen, 2009). Likewise, while mulberries

(*Morus* L.), paper mulberries (*Broussonetia* L'Hér. ex Vent.), and osage oranges (*Maclura*

Nutt.) were once treated as tribe Moreae Gaudich. on account of their inflexed stamens,

phylogenetic analyses reveal them to belong to three distinct clades, each containing an

assemblage of genera with and without inflexed stamens (Clement & Weiblen, 2009;

Zerega & Gardner, 2019).

This study focuses on the tribe Moreae, a clade of six genera and an estimated 66

species (Clement & Weiblen, 2009). This widespread group of plants contains species of

ecological and cultural importance, as well as the economically important white mulberry

(*Morus alba* L.), whose leaves sustain *Bombyx mori* L. (Bombycidae), the invertebrate

proletariat of the silk industry.

*Morphological basis for higher classification in Moraceae*

By the end of the nineteenth century, Engler (1889) had circumscribed a Moraceae

that is quite similar to the modern concept of the family, with two subfamilies: the

Moroideae, comprising tribes Dorstenieae Gaudich., Broussonetieae Gaudich., Fatouae

Engler, Moreae, and Strebleae Bureau, characterized by stamens inflexed in bud (broadly

construed, including *Dorstenia* where they straighten gradually rather than spring outward

suddenly); and the Artocarpoideae, composed of tribes Brosimae Trécul, Euartocarpeae

Trécul, Ficeae Dumort., and Olmedieae Trécul and characterized by stamens straight in

bud. The two most influential scholars of Moraceae classification in the 20th century were

E.J.H. Corner and C.C. Berg. Corner considered inflorescence architecture to be the most

important character for higher-rank taxonomy within the family and inflexed stamens

consequently of secondary importance(Corner, 1962). Berg by contrast questioned the

utility of inflorescence architecture and took into account a variety of characters, especially

the presence of inflexed stamens (Berg, 1977b, 2001).

The taxonomic history of the family has leaned heavily on this stamen character.

Engler's Moreae (1899), unchanged in substance from Bureau's (1873), contained six

genera, *Ampalis* Bojer*, Pachytrophe* Bureau*, Paratrophis* Blume*, Pseudomoru*s Bureau,

and *Morus*, all with inflexed stamens that spring out suddenly at anthesis (Table 1).

Corner's expanded Moreae consisted of seven genera with either straight or inflexed

stamens, but with pistillate inflorescences never condensed into a head: *Fatoua* Gaudich,

*Morus*, *Sorocea* A. St.-Hil. (apparently including *Paraclarisia* Ducke), *Clarisia* Ruiz &

Pav., *Ampalis*, *Pachytrophe*, and *Streblus* Lour. (including *Bleekrodea* Blume*, Paratrophis,*

*Pseudomorus*, *Taxotrophis* Blume, *Sloetiopsis* Engl. and *Neosloetiopsis* Engl.). Corner

himself, however, found the diversity of his Moreae unsatisfactory, noting that "[t]oo many

genera on insufficient and invalid grounds trouble this small tribe" (Corner, 1962). Perhaps

in response, Berg's Moreae comprised all of the genera with inflexed stamens and none

without, including *Broussonetia* and *Maclura* but excluding *Sorocea* and *Clarisia*,

providing a simple character with which to delimit the tribe(Berg, 2001; Berg & al., 2006).

Recent phylogenetic work has supported a third approach, with the Moreae

comprising six genera, including genera with both straight (*Sorocea*, *Bagassa* Aubl.) and

inflexed (*Broussonetia*, *Maclura*) stamens (Clement & Weiblen, 2009). These studies also

suggest that the character state of inflexed stamens is plesiomorphic and is the ancestral

state for both Moraceae and Urticaceae (Datwyler & Weiblen, 2004; Zerega & al., 2005;

Clement & Weiblen, 2009). Inflexed stamens, which are precursors to ballistic pollen

release, are associated with wind pollination (Bawa & Crisp, 1980; Berg, 2001), while the

loss of inflexed stamens is associated with animal pollination. Although dominant in the

Moreae, inflexed stamens also occur in two other tribes: Maclureae W.L. Clement & G.

Weiblen and Dorstenieae. Genera lacking inflexed stamens occur in all seven tribes of

Moraceae (Clement & Weiblen, 2009; Zerega & Gardner, 2019).

*Taxonomic summary of Moreae*

Following Clement & Weiblen (2009), the Moreae comprise six genera and an

estimated 66 species: *Bagassa* (1), *Milicia* Sim (2), *Morus* (16), *Sorocea* (16), *Streblus*

(23), and *Trophis* P. Browne. (8) (Berg, 1977a, 2001; Berg & al., 2006; Clement &

Weiblen, 2009; Filho & al., 2009; Machado & al., 2013; Santos & Neto, 2015). Here, we

present an overview of these genera.

*Morus* L., the true mulberries, is characterized by leaves with crenate margins and
trinerved bases, stamens inflexed in bud, and many-flowered pistillate spikes whose four-
parted perianths become fleshy in fruit, the aggregations superficially resembling a
blackberry. *Morus* comprises approximately 16 species whose delimitation requires further
research. There are three subgenera: *Morus* (ca. 14 species) which is found in temperate to
tropical Asia and from North America to Mexico; *Gomphomorus* Leroy, a single species
restricted to tropical South America; and *Afromorus* (Bureau ex Leroy), a single species
restricted to tropical Africa. Previous phylogenetic work has suggested that these three
subgenera may not form a monophyletic clade (Nepal, 2012). *Milicia* (2 spp., Africa) has
inflorescences which somewhat resemble those of *Morus*, but the leaves of *Milicia*, with
entire margins and pinnate venation, prevent any confusion of the two genera.

*Streblus* Lour., with 23 species from India to Southeast Asia and Oceania, is
morphologically heterogeneous, but its species are all characterized by stamens inflexed in
bud and pistillate flowers with more or less free tepals that enclose the fruit loosely or not
at all. Initially described by Loureiro based on the widespread *S. asper* Lour.—notable for
its discoid-capitate staminate inflorescences with the rudiments of an involucre—*Streblus*
was broadened by Corner (1962), bringing in as sections *Taxotrophis*, *Phyllochlamys*
Bureau, *Paratrophis* (including *Pseudomorus*), *Pseudostreblus* Bureau, *Bleekrodea*, and
*Sloetia* Teijsm. & Binn. (apparently including *Sloetiopsis* but without making any
combinations). None of these have discoid-capitate inflorescences; they mostly have
spicate staminate inflorescences, and the latter two can have bisexual inflorescences.
Corner viewed these sections as fragments of an ancient lineage preserving ancestral
characters (Corner, 1975). Following additional work by Corner (1970, 1975), Berg
included the genera *Ampalis* and *Pachytrophe* as section *Ampalis* (Bojer) C.C. Berg,
subsumed *Sloetiopsis* and section *Taxotrophis* into section *Streblus*, and excluded section
*Bleekrodea*, reinstating it as a genus (Berg, 1988). Berg later reinstated section *Taxotrophis*,
whose species are unique in bearing axillary spines (Berg, 2005; Berg & al., 2006). In
2009, Clement and Weiblen reinstated *Sloetia* at generic rank and transferred it and
*Bleekrodea* to Dorstenieae based on phylogenetic evidence (Clement & Weiblen, 2009).

*Trophis* P. Browne is characterized by stamens inflexed in bud, spicate staminate

inflorescences, and tubular pistillate perianths, becoming fleshy and enclosing the fruit,

except for the monotypic section *Olmedia* (Ruiz & Pav.) C.C. Berg (*T. caucana*), which has

discoid-capitate staminate inflorescences with an involucre. *Trophis* (regrettably conserved

over Linnaeus's *Bucephalon* L.) as recognized by Berg (1988, 2001) had five sections:

*Trophis*, restricted to Latin America; *Calpidochlamys* (Diels) Corner (previously included

in *Paratrophis* and *Uromorus*), restricted to Southeast Asia (Corner, 1962); *Maillardia*

(Frapp. ex Duch.) C.C. Berg, restricted to Africa; *Malaisia* (Blanco) C.C. Berg, restricted to

Southeast Asia; and *Olmedia*, restricted to Latin America. The sinking of *Olmedia* into

*Trophis* by Berg (1988), requiring the re-typification of the tribe Olmedieae Trécul, which

became Castilleae C.C. Berg. Recently, *Malaisia* Blanco was reinstated as a genus and

transferred to Dorstenieae based on phylogenetic evidence (Clement & Weiblen, 2009),

reducing the current number of sections to four.

Two Neotropical genera included in Moreae by Clement and Weiblen (2009) but

not by Berg (2001) have straight stamens. While the monotypic *Bagassa* Aubl., with its

long staminate catkins, is wind pollinated (Bawa & Crisp, 1980), evidence suggests that

*Sorocea* A. St.-Hil. (19 spp.), which produces racemose staminate inflorescences that do

not have as many flowers as those of *Bagassa*, likely contains both wind- and insect-

pollinated species (Zapata & Arroyo, 1978; Bawa & al., 1985; Lewis, 1986). The pistillate

perianths are subtended by pluricellular hairs, believed to serve as a substrate for a fungus

which, in turn attracts pollinators (Berg, 2001). The staminate flowers of *Sorocea affinis*

Helmsl. are apparently fragrant (fide *S. Zona 778*, 21 Nov. 1997, FTBG♂♀), suggesting

insect pollination for that species. If insect pollination is confirmed in *Sorocea*, it would

likely represent another transition from ancestral wind to derived insect pollination in

Moraceae.

Disagreement between the two principal recent monographers of the Moraceae,

Corner (1962) and Berg (2001, 2005, 2006), over the delimitation of the Moreae has been

compounded by a series of phylogenies based on the analysis of molecular (DNA

sequence) data that have redefined genera and their relationships. This has resulted in a

poorly delimited tribe, both with respect to diagnostic morphological characters, the genera

it comprises, and the rank of several taxa (see above). Ensuring that the tribe and its genera

are monophyletic will result in a classification that better reflects evolutionary history,

providing a framework for answering broader scientific questions. Our aim, therefore, was

to generate a comprehensive phylogeny of the Moreae (Fig. 1) and its near allies by

sampling all of the potential genera and species in the tribe sensu Berg, and sensu Clement

& Weiblen (2009) and use this to redelimit the tribe and its genera. We aimed to do so

through increased taxon and genome sampling compared to previous studies, including a

nearly comprehensive sample of the taxa in Moreae (56/66 species) and the allied tribes

Artocarpeae (76/83 taxa) and Maclureae (12/12). Our data set combines phylogenomic data

generated using two largely non-overlapping sets of enrichment baits, one developed

specifically for the Moraceae (Gardner et al., 2016), the other for the whole of the

Angiosperms (Johnson & al., 2019), allowing us to explore the possibilities and challenges

of combining samples based on largely non-overlapping loci. The resulting generic revision

lays the groundwork for species-level revisionary work and provides clarity to this

economically important clade.

We also set out to reconstruct the evolutionary history of inflexed stamens within

Moraceae using ancestral state reconstruction. The loss of inflexed stamens in

Castilleae+Ficeae and Artocarpeae have been associated with transitions from wind to

animal pollination (Momose & al., 1998; Sakai & al., 2000; Datwyler & Weiblen, 2004;

Gardner & al., 2018). A more complete picture of evolutionary transitions between inflexed

and straight stamens may help focus further research on transitions in pollination biology

with Moraceae.

**Materials and Methods**

We used target enrichment sequencing (HybSeq) (Weitemier & al., 2014) to capture

333 genes previously developed for phylogenetic work in Moraceae (the "Moraceae333")

(Gardner & al., 2016; Johnson & al., 2016). This method allows for efficient capture of

hundreds of loci and is suitable for both fresh material and degraded DNA from herbarium

material (Villaverde & al., 2018; Brewer & al., 2019), which comprises much of the

material employed in this study.

*Taxon sampling*, *library preparation, and sequencing*

We sampled 56 out of 66 species (and all genera) in Moreae as well as select outgroup

taxa from Artocarpeae using DNA from leaf tissue preserved on silica gel or—in most

cases—samples from herbarium specimens (Table S1). In either case, DNA was extracted

using a modified CTAB method, usually with increased incubation times to maximize yield

from herbarium tissue(Doyle & Doyle, 1987; Hale & al., 2020). Samples were quantified

using a Qubit fluorometer (Invitrogen, Life Technologies, California, USA), and herbarium

samples were run on a gel to test for degradation. For most samples 200 ng of DNA was

used for library preparation; for some low-yield samples, as little as 50 ng was used, and for

very degraded samples, as much input as possible was used, up to 400 ng. Undegraded

DNA was fragmented either using NEB DNA Fragmentase (New England Biolabs,

Ipswich, Massachusetts, USA) or on a Covaris M220 (Covaris, Wobum, Massachusetts,

USA). DNA samples with an average fragment size less than 500 bp were not fragmented

at all, and partially-degraded samples with an average fragment size of over 500 bp were

fragmented on the Covaris M220. TruSeq-style library preparation was carried out using

either the KAPA Hyper Prep kit (Kapa Biosystems, Wilmington, MA) or the NEB DNA

Ultra 2 kit following the manufacturer's protocols, except that end repair, A-tailing, and

adapter ligation were carried out in reduced-volume reactions (0.25x for KAPA and 0.5x

for NEB) to reduce costs. Final products were quantified on the Qubit and combined in

equal molecular weights into pools of 16–20 samples. The pools totaled 1,200 µg each if

enough library preparation was available. Pools were hybridized for 16–24 hours to custom

Moraceae probes (Gardner et al, 2016) using a MYBaits kit (Arbor Biosciences, Ann

Arbor, Michigan, USA) following the manufacturer's protocol, except that the probes were

diluted 1:1 with nuclease-free water. Hybridization products were reamplified using KAPA

Hot Start PCR reagents following the MYbaits protocol, quantified on the Qubit, and

quality-checked on an Agilent BioAnalyzer (Agilent Technologies, Palo Alto, California,

USA). Samples with adapter dimer peaks were cleaned using 0.7x SPRI beads and re-run

on the Qubit and BioAnalyzer. An initial sequencing run took place on a MiSeq 2 x 300bp

run (v3) (Illumina, San Diego, California, USA) at the Field Museum of Natural History,

and then additional samples were sequenced on a HiSeq 4000 2 x 150bp run at the

Northwestern University Genomics Core.

We also included 47 Moraceae and Urticaceae samples enriched for the

Angiosperm353 probes and sequenced as part of the Plant and Fungal Trees of Life project

(PAFTOL, RBG Kew; https://www.kew.org/science/our-science/projects/plant-and-fungal-

trees-of-life; Table S1). Sample preparation and sequencing followed Johnson et al. (2019).

The PAFTOL samples were enriched with a universal probe set developed for angiosperms

(the "Angiosperms353").

Finally, we used samples sequenced for previous phylogenetics projects in *Artocarpus*

J.R. Forst. & G. Forst. and Parartocarpeae Zerega & E.M. Gardner to complete our

sampling (Johnson & al., 2016; Gardner, 2017; Kates & al., 2018; Zerega & Gardner,

2019). The final dataset contained 247 samples.

*Assembly of reads*

We trimmed reads using Trimmomatic (ILLUMINACLIP: TruSeq3-PE.fa:2:30:10

HEADCROP:3 LEADING:30 TRAILING:25 SLIDINGWINDOW:4:25 MINLEN:20)

(Bolger & al., 2014) and assembled them with HybPiper, which produces gene-by-gene,

reference-guided, *de novo* assemblies (Johnson & al., 2016). For the samples enriched with

the Angiosperms353 baits, we used the reference described by Johnson et al. (2019), and

for the samples enriched with the Moraceae333 baits, we used the reference described in

Zerega & Gardner (2019). To increase overlap between the two data sets beyond the 5

genes they inherently have in common, we used HybPiper to assemble the

Angiosperms353-enriched reads using the Moraceae333 targets and vice-versa, in order to

capture any additional genes found in off-target reads. The exception was most of the

*Artocarpus* data set, which was previously assembled for another study using only the

Moraceae333 targets. We used the HybPiper script "intronerate.py" to build "supercontig"

sequences for each gene, consisting of exons as well as any assembled flanking non-coding

sequences (intronic or intergenic). To these new assemblies, we added 76 Moraceae333

assemblies from Gardner (2017) to complete sampling for the Artocarpeae.

*Main phylogenetic analyses*

The full data set was analyzed using the "exon" sequences to ensure good alignment

across the entire family. To maximize phylogenetic resolution within Moreae, a smaller

data set consisting only of Moreae taxa and a single outgroup taxon (*Artocarpus*

*heterophyllus*) was analyzed using the "supercontig" sequences but following the same

methodology. For each of the 686 genes assembled, we discarded sequences shorter than

25% of the average length for the gene and discarded genes containing sequences for less

than 30 samples. We aligned each gene with MAFFT (Katoh & Standley, 2013)and used

Trimal (Capella-Gutiérrez & al., 2009) to discard sequences with an average pairwise

identity of less than 0.5 to all other sequences in the alignment (indicative of poor a poor

quality sequence that could not be properly aligned) as well as columns containing more

than 75% gaps. We used RAxML 8.2.4 (Stamatakis, 2014) to estimate a maximum-

likelihood tree for a partitioned supermatrix of all genes as well as for each gene

individually (GTRCAT model, 200 bootstrap replicates). We then used ASTRAL-III

(Zhang & al., 2017), a summary-coalescent method, to estimate a species tree from all gene

trees, estimating node support using both bootstrap (160 replicates, resampling across and

within genes) and local posterior probability (normalized quartet scores, representing gene

tree concordance). Finally, we used ASTRAL-III to test whether, for each node, the null

hypothesis of a polytomy could be rejected (Sayyari & Mirarab, 2018). Alignment,

trimming, and estimation of gene trees were parallelized using GNU Parallel (Tange,

2018).

*Whole chloroplast phylogenetic tree*

We also built a whole-chloroplast phylogenetic tree as follows. Rather than

assembling full-length genomes, which can be extremely slow to align, we assembled and

aligned the genomes in sections, dramatically speeding up the process. We created

HybPiper targets using the chloroplast genome of *Morus indica* L. (NCBI RefSeq accession

no. NC_008359.1) and the associated gene annotations. Each target consisted of a gene

feature concatenated with any internal or subsequent non-coding sequence, terminating one

base before the next gene feature began. These intervals were generated by manually

editing the NCBI gff3 file in Excel (Microsoft Corp., Redmond, Washington, USA) and

then extracted using BedTools (Quinlan & Hall, 2010). Assemblies for all Moraceae

samples and *Boehmeria nivea* (L.) Gaudich. were carried out in HybPiper using a coverage

cutoff of 2 and otherwise with default parameters. Within each target, sequencing less than

25% of the average length were discarded, and after alignment with MAFFT, sequences

with an average pairwise identity of less than 0.7 were discarded (the higher cutoff

reflecting the conserved nature of chloroplast DNA). Alignments were then concatenated

into a supermatrix, and samples with more than 50% undetermined characters were

discarded. Alignments were then rebuilt, filtered, and concatenated, and columns in the

final supermatrix with more than 75% missing characters were discarded. A maximum-

likelihood tree was generated using RAxML 8.2.4 under the GTRCAT model, with 200

rapid bootstrap replicates.

*Phylogenetic network analysis*

To further investigate relationships within the *Paratrophis* clade, including the proper

placement of *Morus insignis* and *Trophis philippinensis* (Bureau) Corner, we constructed

phylogenetic networks based on two reduced 8-taxon datasets ("exon" and "supercontig")

consisting of those taxa plus *Streblus anthropophagorum* (Seem.) Corner*, S. glaber* (Merr.)

Corner*, S. glaber* subsp. *australianus* (C.T. White) C.C. Berg, and *S. heterophyllus* (Blume)

Corner*, with *S. mauritianus* (Jacq.) Blume as the outgroup. Alignment preparation followed

the workflow outline above except that only genes with sequences for all eight taxa were

retained. Rooted gene trees were used to infer the network in PhyloNet using the

"InferNetwork_MPL" command, allowing a maximum of four hybridization events and

collapsing gene tree nodes with less than 30% bootstrap support (Yu & Nakhleh, 2015;

Wen & al., 2018) (Wen et al., 2018; Yu and Nakhleh, 2015); AIC was scored for the best

five networks from 10 runs, using the number of branch lengths and hybridization events

calculated as the number of parameters (Kamneva & al., 2017).

*ITS and rbcL phylogenetic trees*

While we did not have our own sequences for *Streblus tonkinensis* (Eberh. & Dubard)

Corner, *S. ascendens* Corner, *S. banksii* (Cheeseman) C.J. Webb, and *S. smithii*

(Cheeseman) Corner, either *ITS* or *rbcL* sequences existed in NCBI GenBank for these taxa

(Fig. S2). We were not able to examine the underlying specimens ourselves, but we

considered the chances of misidentification low because those species are morphologically

and/or geographically distinctive. We constructed *ITS* and *rbcL* data sets from our own

samples using HybPiper, using *Streblus* sequences for these loci (obtained from GenBank)

as targets and reducing the coverage cutoff at 2 to recover the loci from off-target reads. To

these sequences, we added GenBank sequences of *Streblus* taxa as well as *Trophis caucana*

(Pittier) C.C. Berg. For each locus, we aligned sequences with MAFFT, discarded short

sequences (<490 bp), trimmed alignments to remove columns with over 75% gaps, and

built maximum-likelihood trees using RAxML (GTRGAMMA, 1000 rapid bootstraps).

*Divergence time estimation*

The supermatrix maximum-likelihood tree was time calibrated using ape v 5.3

(Paradis & Schliep, 2019) in R v 3.5.1 (Team, 2019). First, the tree was pruned of duplicate

taxa and non-Moraceae outgroup taxa, and edge lengths (in substitutions per site) were

multiplied by the number of sites in the alignment (to convert them to substitutions). The

following stem nodes were constrained with minimum ages (in million years, Ma) based on

fossil data, following(Zhang & al., 2019): *Ficus,* 56 Ma; *Broussonetia*, 33.9 Ma; *Morus*

(subg. *Morus*, based on U.S.S.R. locality of the fossil), 33.9 Ma; and *Artocarpus*, 64 Ma.

The crown node of Moraceae was constrained to a minimum age of 73.2 Ma and a

maximum age of 84.7 Ma.(Zhang & al., 2019), which had a more extensive outgroup

sampling that we have here. The tree was then time-calibrated using the chronos function

under two different models ("relaxed" and "correlated") (Kim & Sanderson, 2008; Paradis,

2013). The smoothing parameter (λ) was chosen using the cross-validation method in the

chronopl function (testing $\lambda = 0$ and $0_{-5}$ through $10_{15}$), selecting the value of λ that

minimized the cross-validation statistic (Sanderson, 2002). The resulting trees were

visualized using the densiTree function in phangorn 2.4.0 (Schliep, 2011), and the time

calibrated tree representing the central tendency of these analyses was selected for use in all

further analyses, also taking into account the results of past family-wide studies. As the

penalized likelihood approach used does not integrate over model uncertainty or

uncertainty in calibration placement and timing, confidence intervals on node ages are not

provided in this study. A geologic timescale based on the strat2012 dataset added to tree

figures using PHYLOCH v 1.5-3 (Heibl, 2008).

*Ancestral state reconstruction for inflexed stamens*

All taxa were coded for presence (1) or absence (0) of inflexed stamens in bud.

Taxa such as *Dorsteniae* with stamens that are inflexed in bud but gradually straighten were

coded as 0. We reconstructed ancestral character states on the entire phylogeny by

stochastic mapping using the make.simmap function in the phytools v 0.6-99 (Revell,

2012). We also tested for trait-associated shifts in diversification rates using BAMM v

2.5.0 (Rabosky & al., 2013, 2014a; Rabosky, 2014), specifying the amount of missing taxa

per clade to account for the high proportion of missing taxa in Ficeae, Castilleae, and

Dorstenieae. Parameters were optimized using BAMMtools v 2.1.7 (Rabosky & al.,

2014b). We also tested for trait-dependent diversification using diversitree v 0.9-13

(FitzJohn, 2012). We evaluated BiSSE and trait-independent models on the entire tree and

on a pruned tree consisting only of the densely-sampled Maclureae+Moreae+Artocarpeae

clade.

Reads from have been deposited in Genbank (Non-PAFTOL samples: BioProject

PRJNA322184; PAFTOL samples: BioProject PRJEB35285, subprojects PRJEB37667 for

Moraceae and PRJEB37665 for Urticaceae). Tree files and scripts outlining the

phylogenetic analyses have been deposited in the Dryad Data Repository (###TBA###).

**Results**

The final data set contained 247 samples, 47 enriched with the Angiosperms353

baits, 196 enriched with the Moraceae333 baits, and 4 extracted from whole genomes) and

619 genes (286 Angiosperms353 and all of the Moraceae333) (Table S1). The "exon"

supermatrix (all taxa) contained 613,126 characters, and the "supercontig" supermatrix

(Moreae and select outgroup taxa only) contained 799,926 characters. The chloroplast data

set contained 113 loci.

On average, we assembled 39 Moraceae333 genes from the Angiosperms353-

enriched samples and 20 Angiosperms353 genes from the Moraceae333-enriched samples,

and the average pairwise overlap in assembled loci was 210 (Tables S1, S2). Nineteen taxa

were represented in both sets of samples, and 15 of these were always monophyletic (Figs.

2, 3). Twelve resolved as sister pairs: *Antiaropsis decipiens* (31 loci overlapping), *Bagassa*

*guianensis* Aubl. (38), *Batocarpus amazonicus* (Ducke) Fosberg (22),  *Batocarpus*

*orinocensis* H. Karst. (34), *Brosimum alicastrum* Sw. (42), *Clarisia racemosa* Ruiz & Pav.

(59), *Maclura africana* (Bureau) Corner (213), *Milicia excelsa* (Welw.) C.C. Berg (45),

*Streblus asper* (Retz.) Lour. (75), *Streblus mauritianus* (Jacq.) Blume (40), *Streblus*

*usambarensis* (Engl.) C.C. Berg (43), and *Trophis caucana* (32). Three more represented

by more than two samples always resolved as a clade: *Sorocea bonplandii* (Baillon) W.C.

Burger, Lanj. & de Boer (3 samples; 19–35 loci overlapping), *Maclura tinctoria* (L.) D.

Don ex Steud. (4 samples; 37–43 loci overlapping), and *Malaisia scandens* (Lour.) Planch.

(3 samples; 27–43 loci overlapping). Three additional taxa resolved as sister pairs in the

ASTRAL analysis but as a grade in the supermatrix analysis: *Maclura cochinchinensis*

(Lour.) Corner (28 loci overlapping), *Streblus heterophyllus* (Blume) Corner (50), and

*Trophis montana* (Leandri) C.C. Berg (56). Only one species, *Utsetela gabonensis* Pellegr.

(18 loci overlapping) was never monophyletic, always forming a grade with *U. neglecta*

Jongkind.

The supermatrix and species trees were broadly concordant except within

*Artocarpus*, where the species tree was much more consistent with previous analyses within

that genus (Gardner et al., in review) (Figs. 2, 3). Otherwise, most differences were at

shallow phylogenetic depths, such as relationships within *Streblus* section *Paratrophis*. For

both data sets, the polytomy hypothesis was rejected ($P < 0.05$) for Moraceae and all tribe

and genus-level clades (following the revised classification presented below) (Figs. 2, 3).

Of the genera in Moreae sensu Clement & Weiblen, only *Milicia* and *Sorocea* were

monophyletic; *Morus*, *Streblus*, and *Trophis* were not; *Bagassa* is monotypic and resolved

as sister to Sorocea, with which it shares straight stamens. Four Moreae species resolved

with other tribes: *Trophis caucana* was nested within Castilleae, *Streblus asper* (Retz.)

Lour. was sister to Castilleae, *Streblus indicus* (Bureau) Corner was sister to

Parartocarpeae, and *Streblus usambarensis* was nested within Dorstenieae. Moreae was

otherwise monophyletic, comprising the following subclades: (1) *Streblus* section

*Taxotrophis*, sister to all other Moreae; (2) (a) *Trophis* section *Maillardia*, (b) *Milicia* +

*Morus* subgenus *Afromorus*, (c) *Streblus* section *Ampalis*, (d) *Streblus* section *Paratrophis*

in part + *Morus* subgenus *Gomphomorus*, (e) *Streblus* section *Paratrophis* in part + *Trophis*

section *Calpidochlamys*; (3) *Bagassa* + *Sorocea*; (4) *Trophis* sections *Trophis* and

*Echinocarpa* C.C. Berg + *Morus*. These are roughly geographic clades: (1) Southeast Asia;

(2) (a) Madagascar, (b) Southeast Africa, (c) Madagascar, (d) Pacific + South America, (e)

Southeast Asia + Pacific; (3) South America; (4) South America.

While *Trophis philippinensis* was inside the *Paratrophis* clade in all analyses, the

position of *Morus insignis* was not stable. In the "exon" analyses, it was always in

*Paratrophis* (Fig. 2), although the polotomy hypothesis could not be rejected for its

position as sister to *S. anthropophagorum+S. heterophyllus*. In the "supercontig" and

chloroplast analyses, it was sister to *Paratrophis*, and the polotomy test was rejected for

that node (Figs. 3, S1). In the five phylogenetic networks reported (Fig. 4, Table S3), *M.*

*insignis* was inside *Paratrophis* in three of them, had a hybrid origin involving *Paratrophis*

in one, and was only unambiguously sister to *Paratrophis* in one network, although even in

that one it was involved in a hybridization involving *Paratrophis*.

The chloroplast phylogenetic tree (Fig. S1) was generally in agreement with the

nuclear phylogenetic trees, with three notable differences. *Streblus indicus* and the

Parartocarpeae formed a grade paraphyletic to Dorstenieae, Castilleae, and Ficeae, rather

than a clade. *Streblus zeylanicus* (Thwaites) Kurz was sister to *S. taxoides* (B. Heyne ex

Roth) Kurz (instead of sister to all of section *Taxotrophis*), although with low support

(62%), and *S. glaber* subsp. *australianus* was not sister to subsp. *glaber*, although its

placement was not strongly supported (80%). Finally, *Morus insignis* was sister to

*Paratrophis*, agreeing with the "supercontig" analysis but not the "exon" analysis.

The ITS and *rbc*L phylogenetic trees based on few characters, were not very well

resolved, with only few nodes attaining 100% bootstrap support (Fig. S2). Nevertheless,

with the exception of a few stray taxa (one *Dorstenia* sample in ITS and one *Streblus*

*indicus* samples in *rbc*L), both phylogenetic trees reconstructed monophyletic tribes with at

least moderate support. In the ITS phylogenetic tree, *Streblus smithii* and *S. banksii* were

both in a well-supported clade comprising section *Paratrophis*, and *S. tonkinensis* was part

of the Castilleae clade, which also contained *S. asper* and *Trophis caucana*. In the *rbc*L

phylogenetic tree, the two *Streblus ascendens* samples, the four samples of the

undetermined *Streblus* from Papua New Guinea, and *S. smithii* were part of a well-

supported clade comprising section *Paratrophis*.

*Time-calibration and ancestral state reconstruction*

The cross-validation criterion was minimized by a smoothing parameter ($\lambda$) value of

$10_6$. The tree calibrated under the correlated model (logLik = -17; p-logLik = -17; $\Phi$IC =

1108) was less sensitive to changes in $\lambda$ and generally had younger ages than the tree

calibrated under the relaxed model (logLik = -56.7; p-logLik = -53129706,  $\Phi$IC =

4209013), the latter of which had perhaps implausibly long terminal branches (Table 2, Fig.

S3). The crown age of Moreae was Paleocene (59.1 Ma) under the correlated model and

late Cretaceous (75.4 Ma) under the relaxed model. Because the correlated tree was more

consistent with past family-wide studies (Zerega & al., 2005; Zhang & al., 2019), we used

that tree for all further analyses (Fig. 5). The BAMM analysis found a single credible rate

shift, not surprisingly at the crown node of *Ficus* (Fig. S4); that rate shift was—also not

surprisingly—associated with a loss of inflexed stamens, which are never found in *Ficus* (*P*

= 0.035); no rate shifts were found within Moreae. Ancestral reconstruction on the entire

tree found that ancestral Moraceae had stamens inflexed in bud; these were lost nine times

(in Ficeae, Olmedieae [=Castilleae], Parartocarpeae, twice in Dorstenieae, *Maclura* section

*Cudrania* (Trécul) Corner, Artocarpeae, *Bagassa,* and *Sorocea*) and regained once (in

*Trophis caucaua*) (Fig. 6). Model testing on both the whole tree and on the

Moreae+Artocarpeae+Maclureae subtree indicated that a BiSSE model was not

substantially better than a trait-independent model (Moreae only: AIC for BiSSE = 1110.4;

trait-independent = 1112.8; $\Delta$AIC = 2.4; whole family: AIC for BiSSE = 1488.3; trait-

independent = 1490.4; ΔAIC = 2.1). In any event, the reconstructions were identical (Figs.

6).

**Discussion**

*Sequencing and combination of data sets*

This study was materially improved by our ability to combine samples enriched

with two largely non-overlapping bait sets. Despite minimal by-design overlap, we were

able to assemble many overlapping loci for samples with moderate to deep coverage (Table

S2), adding taxa that would not otherwise have been included in the study and replicating

taxa to confirm unexpected phylogenetic placement (e.g, *Streblus indicus*). Taxa replicated

across the two data sets performed well in phylogenetic analyses, with 15 out of 19 always

resolving as monophyletic in supermatrix analyses and 18/19 so resolving in ASTRAL

analyses, with only *Utsetela gabonensis* always forming a grade instead (with a difficult-to-

distinguish congener). Our results should embolden others to combine and repurpose data

sets in similar ways.

*Higher taxonomy in Moraceae and the delimitation of the tribe Moreae*

Our results support Corner's overall approach to the classification of Moraceae, if

not all of its details, including the primacy of inflorescence architecture and the

unreliability of inflexed stamens for higher taxonomy (Corner, 1962). Inflexed stamens are

a plesiomorphy that was lost nine times in Moraceae (Fig. 6), a preserved ancestral

character whose past taxonomic importance accounts for the rather extreme non-

monophyly of the Moreae. *Streblus* s.l. provides the best illustration of this principle,

appearing in four out of seven tribe-level clades within Moraceae. We may justly call its

disparate sections, as Corner did, "fragments of an ancestral *Streblus*." (Corner, 1975)—or

in modern parlance, a paraphyletic remnant preserving plesiomorphic staminate flowers

similar to those likely to have occurred in the ancestor of all Moraceae, with four free tepals

and four inflexed stamens (Clement & Weiblen, 2009). If one of the goals of modern

systematics is to establish taxonomic frameworks that reflect as far as possible real

evolutionary relationships, our results may serve as a warning to carefully investigate

whether characters used for taxonomy are derived (synapomorphic) or ancestral

(symplesiomorphic).

Broadly speaking, a tribal delimitation based on inflorescence architecture

(supplemented with other characters in some cases) agrees best with the phylogenetic trees

presented here. Moreae (as revised below) have unisexual spicate or racemose

inflorescences (the pistillate ones sometimes uniflorous), with the globose-capitate pistillate

inflorescences of the monotypic *Bagassa* as the sole exception. Stamens may be either

inflexed or straight. Elsewhere in the family, racemes and spikes are rare, with the former

found in *Maclura* (in part) and the latter found in the related genera *Broussonetia*,

*Allaeanthus* Thwaites, and *Malaisia* and arguably in *Batocarpus* H. Karst. and *Clarisia*; all

of these have capitate pistillate inflorescences. The taxa of *Streblus* s.l. and *Trophis* s.l. that

must be excluded from Moreae all have inflorescences that do not fit our general rule:

discoid-capitate (*Streblus asper* and *Trophis caucana*), cymose (*Streblus indicus*), or

bisexual (*Streblus usambarensis*). In these cases, perhaps Corner did not take his emphasis

on inflorescence architecture quite far enough, including too wide of a variety in this one

tribe. In critiquing the utility of inflorescence architecture for classification, Berg (1977b)

noted the similarity in the inflorescence structure of *Bleekrodea* (then part of Moreae, and

included in *Streblus* by Corner) to that of *Utsetela* and *Helianthostylis* (Dorstenieae), an

observation that proved prescient when *Bleekrodea* was found to belong to Dorstenieae

(Clement & Weiblen, 2009).

Olmedieae (as revised below, including Castilleae) can be defined entirely based

upon the presence of a discoid inflorescence subtended by an involucre of imbricate bracts.

Two species with such involucres previously classified as Moreae, *Streblus asper* and

*Trophis caucana* (=*Olmedia aspera*) always appeared in the Olmedieae (=Castilleae) clade

in our analyses (Figs. 2–3). *Olmedia aspera* (=*Trophis caucana*)—the nomenclatural type

of the tribe Olmedieae—was transferred to Moreae because of its inflexed stamens and lack

of self-pruning branches (Berg, 1977b). In inflorescence morphology, however, *T. caucana*

closely resembles other Castilleae, always subtended by an involucre of imbricate bracts

(Berg, 1977b, 2001). The staminate inflorescences of *Streblus asper* are strikingly similar

to those of *T. caucana*, and it is remarkable that the affinity between *S. asper* and the

Olmedieae has not been seriously considered until now. Previous barcoding or phylogenetic

studies have placed *Trophis caucana* (Kress & al., 2009) and *Streblus tonkinensis* (Chen &

al., 2016) (closely allied to *S. asper*) in the Castilleae clade, but those results went

unremarked upon, perhaps because of the broad scale of the studies (respectively, forest

community phylogenetics and the vascular plants of China).

The remaining five tribes can all be broadly defined based on inflorescence

architecture as Corner argued, sometimes supplemented by other characters as Berg

preferred, allowing of course for the exceptions made inevitable by the vicissitudes of

evolution. Ficeae of course is defined by the syconium (essentially an urceolate disc that

has been closed at the top). Maclureae have densely-packed globose infructescences and

can be distinguished from Artocarpeae by their four stamens (inflexed in most sections) and

armature. Artocarpeae also have densely-packed globose infructescences but only one

straight stamen and no armature. Parartocarpeae have a few connate involucral bracts and

(in large part) flowers embedded in fleshy receptacles. Dorstenieae are perhaps the most

heterogenous group, but in large part they have bisexual inflorescences, often capitate or

discoid, and often with ballistically-ejected endocarps.

*The role of inflexed stamens in the evolution of Moraceae*

The repeated losses of inflexed stamens (Fig. 6), which are associated with wind

pollination (Bawa & Crisp, 1980; Berg, 2001), raise the possibility that transitions from

wind to animal pollination, which have already been documented in Moraceae (Momose &

al., 1998; Sakai & al., 2000; Datwyler & Weiblen, 2004; Gardner & al., 2018), are even

more common within the family. Generally considered rare (Culley & al., 2002), the shift

from wind to animal pollination may be a repeated feature of Moraceae deserving of further

investigation. Further investigation of *Sorocea*, with its straight stamens and sometimes-

scented inflorescences, may reveal that, like *Artocarpus*, it contains both wind and animal

pollination. And while little is known about pollination in the Dorstenieae, the presence of

unisexual inflorescences and inflexed stamens (e.g., *Broussonetia*) as well as bisexual

inflorescences with straight stamens (e.g., *Brosimum, Dorstenia*) raises the possibility of

transitions within that clade as well; taxa with inflexed stamens but bisexual inflorescences

such as *Bleekrodea* and *Sloetia* (which is visited by bees, EMG pers. obs.) may represent

remnants of an intermediate state. Finally, the conclusion that a single transition to insect

pollination preceded the split between the Ficeae and Olmedieae (Castilleae) should be

reevaluated in light of the position of *Streblus asper* as sister to the latter (Figs. 2, 3). The

only shift in diversification rates our analyses recovered was on the branch leading to the

very diverse *Ficus*, suggesting that the shift away from wind pollination may not by itself

lead to increased diversification.

*Explanation of taxonomic revisions*

We present a generic revision of Moreae (Fig. 7) based on the present phylogenetic

study as well as morphological characters. Arranging monophyletic and morphologically

coherent genera requires one change of rank and 8 new combinations, but no entirely new

names. These revisions provide a framework for within-genus revisionary work, which will

require more intensive sampling and review of specimens within the genera circumscribed

here. We provide an explanation of the taxonomic changes, followed by a formal

presentation of the affected tribes and genera, with complete species lists and synonymy.

*Trophis* and the reinstatement of the Olmedieae — To make *Trophis* monophyletic,

we propose that it be applied strictly to the Neotropical clade that includes the type species

*Trophis americana* L. (=*Trophis racemosa* (L.) Urb.) . To accomplish this, we reinstate the

genus *Maillardia* and transfer *Trophis philippinensis* to *Paratrophis* (discussed below),

reinstating its former name *Paratrophis philippinensis* (Bureau) Fern.-Vill. Because

*Trophis caucana* does not belong in *Trophis* or Moreae but rather with the members of

Castilleae, we reinstate its former name *Olmedia aspera* Ruiz & Pav. and transfer it to that

tribe, whose former name Olmedieae must now be reinstated on account of its priority.

*Streblus* — We restrict the genus *Streblus* to three species comprising most of

section *Streblus*—*Streblus asper, Streblus tonkinensis*, and *Streblus celebensis* C.C. Berg—

and transfer the genus to Olmedieae. Although *S. celebensis* was not included in our

phylogeny, the sub-involucrate inflorescences are similar to those of *S. asper*, with which *S.*

*celebensis* differs primarily in vegetative characters, the latter having broadly-toothed

margins in the distal half of the leaf. Both occur in Sulawesi, where at least one specimen

with intermediate leaf morphology has been collected (Sulawesi, Kendari: *Kjellberg 452*,

24 Feb. 1929, L, det. *S. asper* by E.J.H. Corner). We therefore retain its taxonomic position

in *Streblus*. The remaining member of section *Streblus*, *S. usambarensis*, belongs with

Dorstenieae, and we therefore transfer it to that tribe, reinstating its former name

*Sloetiopsis usambarensis*. *Streblus indicus* is sister to Parartocarpeae, and we therefore

reinstate its former name *Pseudostreblus indicus* and transfer it to Parartocarpeae.

The remaining species of *Streblus* s.l. are properly placed in Moreae but are still

paraphyletic. We therefore reinstate the genera *Ampalis*, *Pachytrophe*, *Paratrophis*, and

*Taxotrophis*, largely corresponding to the former sections but requiring some new

combinations. *Paratrophis* as presented here is united by spicate male inflorescences

(usually) with peltate or reniform bracts (except for *Paratrophis philippinensis* (=*Trophis*

*philippinensis*), which does not have fleshy tepals in fruit). Within *Paratrophis* we include

*Paratrophis ascendens* (Corner) E.M. Gardner (=*Streblus ascendens* Corner) based on its

inflorescence morphology and phylogenetic position in the *rbc*L tree (Fig. S2b). Its

previous position within the monotypic section *Protostreblus* was due to the type

specimen's spiral phyllotaxy, but the latter may be atypical, as a more recent collection

*Womersley NGF 24791* (K, L, BO), has distichous leaves. Berg (1988), recognizing the

affinities between *Pachytrophe dimepate* Bureau and *Ampalis mauritiana*, included both in

*Streblus* section *Ampalis*, noting that in light of the diversity found in *Streblus* s.l.,

separation at the sectional level was no longer warranted; although our phylogenetic results

support this grouping, we provisionally retain them as separate genera because the

differences between them in phyllotaxy, stipule amplexicauli, and embryo characters

equal or exceed the distinctions between other genera in Moreae following our atomization

of *Streblus* s.l. Further intensive study of that species complex may of course warrant a

different approach, including potentially reducing both to a section of *Paratrophis*. Our

species lists within the former components of *Streblus* follow Berg's approach (1988,

2006), with two exceptions. We provisionally recognize *Taxotrophis zeylanica* (Thwaites)

Thwaites as distinct from *Taxotrophis taxoides* (B. Heyne ex Roth) W.L. Chew ex E.M.

Gardner (=*Streblus taxoides* (B. Heyne ex Roth) Kurz) based on our phylogenetic work and

following the *Flora of China*, which recognizes *Streblus zeylanicus* as distinct from *S.*

*taxoides* based on its clustered pistillate inflorescences. In addition, we provisionally

recognize *Paratrophis australiana* (=*S. glaber* subsp. *australianus*) as distinct from

*Paratrophis glabra* based on its geographic and consistent morphological distinctiveness.

*Taxotrophis* and *Paratrophis* warrant further investigation to refine species limits.

*Morus — Morus* has never been revised, and the species concepts are often based on

minor morphological differences (Berg et al., 2006; Berg 2001), and the paraphyly of

several species in our analyses suggests that a broad *M. alba* similar to Bureau's (1873)

may be worth a second look. The monotypic subgenus *Afromorus* is sister to *Milicia*, but

the leaf morphology is markedly different, instead resembling other *Morus* species in its

trinerved based and crenate margins. We therefore raise *Afromorus* to genus level, requiring

one new combination. *Morus insignis*, from western Central and South America, bears a

remarkable resemblance to *Paratrophis*, in particular *P. pendulina*, especially in leaf

morphology, which in *M. insignis* is not consistently trinerved as in other mulberries. The

infructescence appears superficially like a mulberry because of its basally fleshy tepals,

although with more loosely-packed flowers, but closer inspection places it firmly within

*Paratrophis*, with drupes protruding from the persistent tepals, peltate bracts, and a sterile

groove. Because analyses differ as to whether *M. insignis* is sister to *Paratrophis* or part of

it, it seems best to treat them as congeners.

These changes result in eleven monophyletic genera of Moreae, providing a

framework for revisionary work within the genera. Below, we present a complete genus

and species list for Moreae with brief descriptions for all genera and synonymies for new

combinations and reinstated or recircumscribed taxa. All cited protologues were reviewed,

and dates for works published piecemeal were confirmed by reference to Taxonomic

Literature 2 online (https://www.sil.si.edu/DigitalCollections/tl-2/index.cfm).

**Taxonomic Treatment**

*Key to the tribes of Moraceae*

This key follows the tribal circumscription of Clement & Weiblen (2009), as subsequently
modified by Zerega & al. (2010), Chung & al. (2017), Zerega & Gardner (2019), and this
study.

1.  Inflorescences urceolate with the opening entirely closed by ostiolar bracts (i.e.
syconium) such that the flowers are enclosed at all stages of development; lamina
with waxy glandular spots at the base of the midrib or in the axils of the basal lateral
veins beneath – Ficeae (*Ficus*)
1.  Inflorescences capitate, spicate, discoid, or urceolate, but flowers are not
entirely enclosed at all developmental stages (i.e. mature anthers or stigmas are
open to the surrounding environment); lamina without waxy glandular spots. – 2

2. Inflorescences (at least staminate) with an involucre of imbricate bracts; often
with self-pruning horizontal branches (except *Olmedia*, *Poulsenia, Streblus*) –
Olmedieae (*Antiaris, Antiaropsis, Castilla, Helicostylis, Maquira, Mesogyne,*
*Naucleopsis, Olmedia, Perebea, Poulsenia, Pseudolmedia, Sparattosyce*,
*Streblus*)
2. Inflorescences not involucrate; plants without self-pruning branches – 3

3. Plants woody; dioecious; pistillate inflorescences globose-capitate; spines axillary
or terminating short shoots; dioecious – Maclureae (*Maclura*)
3. Plants woody, herbaceous, or succulent; dioecious or monoecious; pistillate
inflorescences various; spines absent or if present, then pistillate inflorescences
not globose capitate – 4

4. Trees or shrubs; monoecious; inflorescences unisexual; staminate flowers with
one stamen (rarely two);– Artocarpeae (*Artocarpus, Batocarpus, Clarisia*)

4. Trees, shrubs, lianas, herbaceous, or succulent; monoecious or dioecious;

inflorescences unisexual or bisexual; staminate flowers with more than one

stamen (or if one stamen then dioecious) – 5

5. Trees or shrubs; monoecious; inflorescences unisexual; stamens straight in bud or

staminate flowers 5-parted and inflexed in bud;– Parartocarpeae (*Hullettia,*

*Parartocarpus, Pseudostreblus*)

5. Trees, shrubs, lianas, herbaceous, or succulent; not as above but otherwise

monoecious or dioecious; bisexual or unisexual inflorescences, stamens straight or

inflexed in bud but staminate flowers never 5-parted – 6

6. Trees, shrubs, lianas, herbaceous, or succulent; bisexual inflorescences (or if

unisexual then a climber or herbaceous) – Dorstenieae (*Allaeanthus* in part*,*

*Bleekrodea, Bosqueiopsis, Brosimum, Broussonetia* in part, *Dorstenia, Fatoua,*

*Helianthostylis, Malaisia, Scyphosyce, Sloetia, Sloetiopsis, Treculia, Trilepsium,*

*Trymatococcus, Utsetela*)

6. Trees or shrubs; unisexual inflorescences – 7

7. Trees; stamens inflexed in bud; pistillate inflorescences globose-capitate –

Dorstenieae (*Allaeanthus* in part, *Broussonetia* in part)

7. Trees or shrubs; stamens inflexed or straight in bud; if stamens inflexed in bud,

pistillate inflorescence not globose-capitate – Moreae (*Afromorus, Ampalis,*

*Bagassa, Maillardia, Milicia, Morus, Pachytrophe, Paratrophis, Taxotrophis,*

*Sorocea, Trophis*)

*Revisions*

713                        **Tribe MOREAE**

**Moreae** Gaudich. in Freyc., Voy. Uranie, Bot. (1830) 509.

Moreae subtribe Soroceae Miq in Martius, Fl. Bras. 4, 1, fasc. 12: 111 (1853), p.t. –

Soroceae (Miq.) C.C. Berg, Blumea 50: 537 (2005), p.p.

Strebleae Bureau in DC., Prodr. 17: 215 (1873), p.p.

Tree or shrubs, monoecious or dioecious. *Leaves* alternate or opposite, distichous or

spirally arranged; stipules lateral to amplexicaul. *Inflorescences* unisexual, uniflorous or

racemose, spicate, capitate, or globose; bracteate; tepals 4, free to connate; staminate

flowers with 4 stamens, filaments straight or inflexed in bud, pistillode usually present;

pistillate flowers with (mostly) free ovary, 2 stigmas. *Fruits* drupaceous or achene-like

with a fleshy persistent perianth, dehiscent or not. *Seeds* with or without endosperm, testa

usually with a thick vascularized part below the hilum, cotyledons equal or unequal,

straight or folded.

Genera and distribution: eleven genera and 63 species with a worldwide distribution.

*Afromorus*

*Afromorus* (**A. Chev. ex J.-F. Leroy) E.M. Gardner, stat. nov.** — based on *Morus* L.

*subg. Afromorus* [A. Chev., Rev. Bot. Appl. Agr. Trop. 29 (315–316): 70 (1949),

invalidly published]; J.-F. Leroy, Rev. Bot. Appl. Agr. Trop. 29 (323–324): 482

(1949) & Bull. Mus. Hist. Nat. Paris, ser. 2, 21: 732 (1949). TYPE: *Afromorus*

*mesozygia* (Stapf ex A. Chev.) E.M. Gardner.

Dioecious trees, shoot apices deciduous. *Leaves* distichous, triplinerved or at least trinerved

at the base. *Stipules* free, more or less lateral. *Inflorescences* solitary or paired, bracts

of varying shapes. Staminate inflorescences spicate, to 2.5mm long, flower 4-parted,

tepals imbricate, ciliolate, pistillode small, apiculate. Pistillate inflorescences

subglobose, ca. 5 mm across, flowers 4-parted, tepals ciliolate, stigma bifid, equal or

unequal, arms filiform to 5 mm long. **Infructescences** subglobose or less often slightly

elongate, ca. 1 mm across, tepals fleshy, yellowish to greenish, drupes ca. 5 x 3–5 mm.

**Seeds** ca. 4.5 x 2.5–4.5 mm.

Species and distribution: one species, in tropical Africa.

Note: The leaves of *Afromorus*—with crenate margins and trinerved bases—bear a striking

resemblance to those of *Morus*, and it is thus not surprising that the former was heretofore

included within the latter. However, leaves of *Afromorus* are distinct in being usually

completely triplinerved, without an upper pinnately-veined portion as in most species of

*Morus*. As the genus is monotypic, the type must be its only species, *Afromorus mesozygia*

(Stapf ex A. Chev.) E.M. Gardner.

Leroy's protologue of *Morus* subg. *Afromorus* was published only in French and

was therefore not a valid diagnosis or description in 1949. However, because the subgenus

was monotypic, the protologue referred to a single name based on a validly-published

description, *Morus lactea* (Sim.) Mildbr. (=*Celtis lactea* Sim.), and was therefore validly

published by reference to the latter.

**1. *Afromorus mesozygia*** (Stapf ex A. Chev.) E.M. Gardner, comb. nov. — based on *Morus*

*mesozygia* Stapf ex A. Chev. [Végétaux utiles de l'Afrique tropicale française 5: 263

(1909), nomen solum], J. Bot. (Morot) ser. 2, t. 2: 99 (1909).

*Celtis lactea* Sim, For. Fl. Port. E. Afr. : 97, t. 96 (1909), probably post-dating *Morus*

*mesozygia* (fide Berg. 1977) – *Morus lactea* (Sim) Mildbr., Notizbl. Bot. Gart.

Berlin 8: 243 (1922) – *Morus mesozygia* var. *lactea* (Sim) A. Chev., Rev. Bot.

Appl. Agr. Trop. 29: 72 (1949).

*Morus mesozygia* var. *sanda* A. Chev., Rev. Bot. Appl. Agr. Trop. 29: 71 (1949),

invalidly published (under Art. 39.1, Latin description or diagnosis lacking).

*Morus mesozygia* var. *colossea* A. Chev., Rev. Bot. Appl. Agr. Trop. 29: 71 (1949),

invalidly published (under Art. 39.1, Latin description or diagnosis lacking).

775                                            ***Ampalis***

*Ampalis* Bojer, Hort. Maurit. 291 (1837)

*Streblus* Lour. subgen. *Parastreblus* Blume, Mus. Bot. Ludg.-Bat. 2: 89 (1856)

*Streblus* Lous. sect. *Ampalis* (Bojer) C.C. Berg, Proc. Kon. Ned. Akad. Wetensch. C

91(4): 358 (1988), p.p.

Dioecious trees or shrubs. ***Leaves*** spirally arranged to nearly distichous, pinnately veined.

***Stipules*** free, nearly lateral. ***Inflorescences*** solitary or paired in the leaf axils, spicate, with

flowers in longitudinal rows, bracts present but sparse, usually subpeltate. Staminate

inflorescences to 9 cm long, flowers 4-parted, decussate-imbricate, stamens 4, inflexed in

bud, pistillode present. Pistillate inflorescences to 12 cm long, flowers tepals 4, separate,

decussate-imbricate, ovary free, stigmas 2, equal. ***Infructescences*** with enlarged fleshy

perianths, surrounding drupaceous fruits, the latter ca. 5–6 mm long. ***Seeds*** ca. 4 x 4 mm,

testa thickened and not distinctly vascularized, cotyledons equal.

Species and distribution: One species, native to Madagascar and Comoros.

**1.** *Ampalis mauritiana* (Jacq.) Urb., Symb. Antill. 8: 165 (1920) – *Morus mauritiana* Jacq.,

Collect. 3: 206 ("1789", 1791) – *Streblus mauritianus* (Jacq.) Blume, Mus. Bot.

Lugd.-Bat. 2: 80 (1856).

*Streblus maritimus* Palaky, Catal. Pl. Madag. 2: 31 (1907)

*Morus nitida* Willem. In Useri, Ann. Bot. 18: 56 (1796)

*M. ampalis* Poir. In Lam., Encycl. Bot. 4: 380 (1797)

*Trophis cylindrica* Roxb., Fl. Indica 3: 599 (1832), pro syn. *M. mauritianae.*

*Ampalis madagascariensis*, Bojer, Hort. Maurit. 291 (1837) nom. illeg., pro syn.

*Ampalis madagascariensis* var. *occidentalis* Léandri, Mém. Inst. Sci. Madag., ser. B,

1: 12, pl. (1948).

*Morus rigida* Hassk., Cat. Hort. Bog. 74 (1844).

Note: May be considered congeneric with *Pachytrophe*, with which it shares inflorescence

and infructescence morphology. They are distinguished on the basis of phyllotaxy,

which is (usually) spiral in *Ampalis* and distichous in *Pachytrophe*.

809                                     ***Bagassa***

*Bagassa* Aubl., Pl. Guiane. Suppl. 15 (1775).

Dioecious trees. ***Leaves*** opposite and decussate, lamina triplinerved, 3-lobed to entire.

***Stipules*** free, lateral. ***Inflorescences*** solitary or paired in the leaf axils, bracteate. Staminate

inflorescences spicate with an abaxial sterile groove, to 12 cm long, flowers in longitudinal

rows with, tepals 4, stamens 2, straight in bud, pistillode present. Pistillate inflorescences

globose-capitate, ca. 1–1.5 cm across, flowers 4-lobed to 4-parted, stigmas 2, filiform.

***Infructescences*** globose, ca. 2.5–3.5 cm across, green, tepals fleshy, yellowish to greenish,

drupes ca. 7–8 mm long. ***Seeds*** ca. 3 x 2 mm, testa thin and not vascularized, cotyledons

equal.

Species and distribution: One, in tropical South America.

**1.** ***Bagassa guianensis*** Aubl., Pl. Guiane. Supple. 15 (1775)

827                                       ***Maillardia***

*Maillardia* Frapp. ex Duch. in Maillard, Notes sur l'île de la Réunion, Annex P. 3 (1862)

*Trophis* P. Br. sect. *Maillardia* (Duch.) Corner, Gard. Bull. Singapore 19: 230 (1962).

Dioecious trees or shrubs. ***Leaves*** distichous, pinnately veined. ***Stipules*** free, semi-

amplexicaul. ***Inflorescences*** solitary or paired in the leaf axils, bracts subpeltate. Staminate

inflorescences spicate with an abaxial sterile groove, flowers 4-parted, decussate-imbricate,

stamens 4, inflexed in bud, pistillode present. Pistillate inflorescences up to three together,

uni- or bi-florous, perianth tubular 4-lobed, ovary adnate to the perianth, stigmas 2, equal.

*Infructescences* with enlarged fleshy perianths surrounding drupaceous fruits, the latter to

18 mm long. *Seeds* to 13 mm long, testa thin, with a thickened vascularized part below the

hilum, cotyledons unequal.

Species and distribution: Two species, in Madagascar, Comoros, and the Seychelles.

**1. *Maillardia borbonica*** Duch., Ann. Notes Réunion, Bot. 1: 148 (1862) – *Trophis*

*borbonica* (Duch.) C.C.Berg, Proc. Kon. Ned. Akad. Wetensch. C 91(4): 355 (1988)

*Maillardia lancifolia* Frapp. ex Duch., Ann. Notes Réunion, Bot. 1: 148 (1862).

*Trophis borbonica* (Duch.) C.C. Berg, Proc. Kon. Ned. Akad. Wetensch. C 91: 355

(1988).

**2. *Maillardia montana*** Leandri, Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 1: 25

(1948) – *Trophis montana* (Leandri) C.C.Berg, Proc. Kon. Ned. Akad. Wetensch. C

91(4): 355 (1988).

*Maillardia occidentalis* Leandri, Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 1: 26

(1948)

*Maillardia orientalis* Léandri, Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 1: 27

(1948)

*Maillardia mandrarensis* Léandri, Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 1:

28 (1948).

*Maillardia pendula* Fosberg, Kew Bull. 29: 266, t. 2 (1974).

861                                        *Milicia*

*Milicia* Sim, For. Fl. Port. E. Afr. 97 (1909).

Dioecious trees. *Leaves* distichous, lamina pinnately veined. *Stipules* free, not fully

amplexicaul. *Inflorescences* spicate, usually solitary in the leaf axils or on leafless nodes,

spicate, flowers in longitudinal rows alternating with rows of bracts, bracts mostly basally

attached, abaxial sterile groove present. Staminate inflorescences to 20 cm long, flowers 4-

parted, imbricate, stamens 4, inflexed in bud, pistillode present. Pistillate inflorescences to

4.5 cm long, flowers 4-parted, decussate-imbricate, ovary free, stigmas 2, unequal.

*Infructescences* with enlarged fleshy perianths, surrounding slightly flattened drupaceous

fruits, the latter to ca. 3 mm long. *Seeds* ca. 2 mm long, testa thin with a slightly thickened

vascularized part below the hilum, cotyledons equal.

Species and distribution: Two species in tropical Africa.

**1.** *Milicia excelsa* (Welw.) C.C. Berg, Bull. Jard. Bot. Natl. Belg. 52(1-2): 227 (1982) –

*Morus excelsa* Welw., Trans. Linn. Soc. Lond. (Bot.) 27: 69, t. 23 (1869) – *Maclura*

*excelsa* (Welw.) Bur. In DC., Prodr. 17: 231 (1873) – *Chlorophora excelsa* (Welw.)

Benth. & Hook., Gen. Pl. 3(1): 363 (1880).

*Chlorophora tenuifolia* Engl., Bot. Jahrb. 20: 139 (1894).

*Milicia africana* Sim, For. Fl. Port. E. Afr. 97, t. 122 (1909).

*Chlorophora alba* A. Chev., Bull. Soc. Bot. Fr. 58 (Mém. 8d): 209 (1912).

**2.** *Milicia regia* (A. Chev.) C.C. Berg, Bull. Jard. Bot. Natl. Belg. 52(1-2): 227 (1982) –

*Chlorophora regia* A. Chev., Bull. Soc. Bot. Fr. 58 (Mém. 8d): 209 (1912) – *Maclura*

*regia* (A. Chev.) Corner, Gard. Bull. Singapore 19: 237 (1962).

889                                                     *Morus*

*Morus* L., Sp. Pl. 986 (1753).

Dioecious trees or shrubs. *Leaves* trinerved (to five-nerved) at the base. *Stipules* free,

nearly lateral. *Inflorescences* solitary or paired in the leaf axils, without an obvious sterile

groove, interfloral bracts absent. Staminate inflorescences spicate or racemose, to 8 cm

long, flowers 4-parted, imbricate, stamens 4, inflexed in bud, pistillode present. Pistillate

inflorescences subcapitate to spicate, up to 16 cm long, flowers tepals 4, separate,

decussate-imbricate, ovary free, stigmas 2, equal. ***Infructescences*** with enlarged fleshy

perianths enclosing achene-like fruits, the latter ca. 1 mm long. ***Seeds*** less than 1 mm long,

cotyledons equal.

Species and distribution: Sixteen species, Asia and North to Central America; introduced

worldwide.

**1.** *Morus alba* L., Sp. Pl. 2: 986 (1753).

**var.** *alba*

**var.** *multicaulis* (Perrottet) Loudon, Arbor. Frutic. Brit. 3: 1348 (1838).

**2.** *Morus australis* Poir. in Desrousseaux et al., Encycl. 4: 380 (1797).

**3.** *Morus boninensis* Koidz., Bot. Mag. (Tokyo) 31: 38 (1917)

**4.** *Morus cathayana* Hemsl., J. Linn. Soc., Bot. 26: 456 (1894)

**var.** *cathayana*

**var.** *gongshanensis* (Z. Y. Cao) Z. Y. Cao, Acta Bot. Yunnan. 17: 154 (1995).

**5.** *Morus celtidifolia* Kunth, Nov. Gen. Sp. [H.B.K.] 2: 33 (1817).

**6.** *Morus liboensis* S. S. Chang, Acta Phytotax. Sin. 22: 66 (1984).

**7.** *Morus macroura* Miq., Pl. Jungh. 1: 42 (1851).

**var.** *macroura*

**var.** *laxiflora* G.K.Upadhyay & A.A.Ansari, Rheedea 20: 44 (2010).

**8.** *Morus koordersiana* J.-F.Leroy, Bull. Mus. Natl. Hist. Nat. sér. 2, 21: 729 (1949)

(endemic to Sumatra and possibly synonymous with *M. macroura*. However, it was

not cited by Berg et al. (2006) either as a good species or as a synonym of the latter).

**9.** *Morus microphylla* Buckley, Acad. Nat. Sci. Philadelphia 1862: 8 (1863) (recognized in

the *Flora of North America* (Flora of North America Editorial Committee, 1993) but

likely conspecific with *M. celtidifolia* and considered so by Berg (2001)).

**10.** *Morus mongolica* (Bureau) C. K. Schneid. in Sargent, Pl. Wilson. 3: 296 (1916).

**11. *Morus nigra*** L. Sp. Pl. 2: 986 (1753).

**12. *Morus notabilis*** C. K. Schneid. in Sargent, Pl. Wilson. 3: 293 (1916).

**13. *Morus rubra*** L., Sp. Pl. 986 (1753).

**var. *rubra***

**var. *murrayana*** (Saar & Galla) Saar, Phytologia 91: 106 (2009).

**14. *Morus serrata*** Roxb., Fl. Ind., ed. 1832, 3: 596 (1832).

**15. *Morus trilobata*** (S. S. Chang) Z. Y. Cao, Acta Phytotax. Sin. 29: 265 (1991).

**16. *Morus wittiorum*** Hand.-Mazz., Anz. Akad. Wiss. Wien, Math.-Naturwiss. Kl. 58: 88

(1921)

*Pachytrophe*

*Pachytrophe* Bureau, Prodr. [A.P. de Candolle] 17: 234 (1873). – *Ampalis* Boj. sect.

*Pachytrophe* (Bur.) Baillon, Hist. Pl. 6: 191 (1875–76)

*Streblus* Lour. sect. *Ampalis* (Boj.) C.C. Berg, Proc. Kon. Ned. Akad. Wetensch. C 91(4):

358 (1988), p.p.

Dioecious trees. ***Leaves*** distichous, pinnately veined. Stipules connate, semi-amplexicaul.

***Inflorescences*** usually paired in the leaf axils, spicate, with an abaxial sterile groove,

flowers sessile in longitudinal rows, tepals 4, decussate-imbricate, ciliolate. Staminate

inflorescences up to 5 cm. long, bracts basally attached, flowers with filaments inflexed in

bud, anthers introrse, pistillode quadrangular, ca. 0.5 mm long. Pistillate inflorescences up

to 2.5 cm long, bracts (sub)peltate, flowers 2–14, tepals free, stigma bifid, arms equal.

***Infructescences*** reddish, with enlarged fleshy perianths ca. 6–8 mm long, drupes thinly

fleshy. ***Seeds*** ca. 4.5 mm long.

Species and distribution: One, in tropical Africa (Madagascar).

Note: *Pachytrophe* could be considered congeneric with *Ampalis*, within which Baillon

treated it as a section (although without making any new specific combinations).

**1.** ***Pachytrophe dimepate*** Bureau, Prodr. [A.P. de Candolle] 17: 234 (1873).

*Pachytrophe obovata* Bureau, Prodr. [A.P. de Candolle] 17: 235 (1873).

*Plecospermum bureaui* A.G. Richt., Term. Füzetek 18: 296 (1895), nomen nudum et

958             superfl., pro. syn. nom. ined. "Plecospermum obovatum Bureau" (*Boivin 1717*, P).

959       *Plecospermum (?) laurifolium* Baill. in Grandidier, Hist. Madag. Vol. 35, Hist. Nat.

960             Pl., Tome 5, Atlas 3, 2e partie: pl. 294a (1895) – *Pachytrophe obovata* var.

961             *laurifolia* (Baill.) Léandri, Mém. Inst. Sci. Madag., ser. B, 1: 16 (1948).

962       *Pachytrophe obovata* var. *montana* Léandri, Mém. Inst. Sci. Madag., ser. B, 1: 16

963             (1948).

966                                         ***Paratrophis***

968 *Paratrophis* Blume, Ann. Mus. Bot. Lugduno-Batavi 2 (1856) 81.

969       *Pseudomorus* Bureau, Ann. Sci. Nat., Bot. sér. 5, 11: 371 (1869).

970       *Uromorus* Bureau in A.DC., Prodr. 17: 236 (1873).

971       *Calpidochlamys* Diels, Bot. Jahrb. Syst. 67: 172 (1935); — *Trophis* sect.

972             *Calpidochlamys* Corner, Gard. Bull. Singapore 19: 230 (1962).

973       *Chevalierodendron* J.-F. Leroy, Compt. Rend. Hebd. Séances Acad. Sci. 227: 146

974             (1948) .

975       *Streblus* Lour. sect. *Paratrophis* (Blume) Corner, Gard. Bull. Singapore 19: 216

976             (1962).

*Streblus* Lour. sect. *Protostreblus* Corner, Blumea 18: 393 (1970).

*Morus* L. *subg. Gomphomorus* J.-F. Leroy, Bull. Mus. Hist. Nat. (Paris), Sér. 2, 21:

732 (1949)

Dioecious trees or shrubs. ***Leaves*** distichous (or spiral in *P. ascendens*), pinnately veined,

sometimes trinerved at the base. Stipules free, lateral. ***Inflorescences*** axillary, solitary or up

to 5 together, spicate, with an abaxial sterile groove, interfloral bracts mostly peltate,

flowers sessile in longitudinal rows, tepals 4, valvate, ciliolate. Staminate inflorescences up

to at least 20 cm. long, flowers with filaments inflexed in bud, pistillode present. Pistillate

inflorescences up to at least 10 cm long, flowers usually at least 2 (to many), tepals free

(except in *P. philippinensis*), stigma bifid, arms equal. ***Fruits*** drupaceous, red to black, with

tepals persistent but usually not enlarged or fleshy (except in *P. philippinensis* and *P.*

*insignis*), up to ca. 1 cm long. ***Seeds*** up to ca. 8 x 6 mm, cotyledons equal.

Species and distribution: Twelve species from the Malesian region to Australia, New

Zealand, and Oceania, Central and western South America.

**1. *Paratrophis anthropophagorum*** (Seem.) Benth. & Hook. f. ex Drake, Ill. Fl. Ins. Pacif.

296 1892

*Trophis anthropophagorum* Seem. [Bonplandia 9 (17–18): 259 (1861), nomen nudum]

Fl. Vit. 258, t. 68 (1868); — *Uromorus anthropophagorum* (Seem.) Bureau in de

Candolle, Prodr. 17 (1873) 236; — *Streblus anthropophagorum* (Seem.) Corner,

Gard. Bull. Sing.19 (1962) 221.

*Caturus oblongatus* Seem., Fl. Vit. 254 (1868).

*Paratrophis ostermeyri* Rech., Fedd. Rep. 5: 130 (1908).

*Paratrophis viridissima* Rech., Fedd. Rep. 5: 130 (1908).

*Paratrophis zahlbruckneri* Rech., Fedd. Rep. 5: 130 (1908).

*Pseudomorus brunoniana* (Endl.) Bureau var. *tahitensis* J. Nadeaud, Énum. Pl. Tahiti

43 (1873) – *Uromorus tahitensis* (J. Nadeaud) Bureau in de Candolle, Prodr. 17: 237

(1873) – *Paratrophis tahitensis* (Bureau) Benth. & Hook.f. ex Drake, Ill. Ins. Mar.

Pacif. Fasc. 7: 296 (1892); et Fl. Polyn. Franc. 193 (1892). – *Streblus tahitensis* (J.

Nadeaud) E.J.H. Corner, Gard. Bull. Singapore 19: 225 (1962).

1009

1010 **2. *Paratrophis ascendens* (Corner) E.M. Gardner comb. nov.** — based on *Streblus*

1011 *ascendens* Corner, Blumea 18: 395, t.1 (1970).

1012

1013    **3. *Paratrophis australiana*** C.T. White, Contr. Arnold Arbor. 4: 15 (1933) — *Streblus*

1014      *glaber* (Merr.) Corner var. *australianus* (C.T. White) Corner, Gard. Bull. Singapore

1015      19: 221 (1962) — *Streblus glaber* (Merr.) Corner subsp. *australianus* (C.T. White)

1016      C.C. Berg, Blumea 50: 548 (2005).

1017

**4. *Paratrophis banksii*** Cheeseman, Man. N.Z. Fl. 633 (1906) – *Streblus banksii*

(Cheeseman) C.J. Webb, in Connor & Edgar, New Zealand J. of Bot. 25 136 (1987).

*Paratrophis heterophylla* var. *elliptica* Kirk T.N.Z.L 29: 500, t. 46 (1897); — *Streblus*

*heterophyllus* var. *ellipticus* (Kirk) Corner, Gard. Bull. Singapore 19: 222 (1962).

Note:— Morphologically very close to and not always distinguishable from *P.*

*microphylla*.

1024

**5. *Paratrophis glabra*** (Merr.) Steenis**,** J. Bot. 72: 8 (1934) – *Gironniera glabra* Merr.,

Philipp. J. Sci., 1, Suppl. 42 (1906); Enum. Philipp. Flow. Pl. 2: 35 (1923)  —

*Chevalierodendron glabrum* (Merr.) J.-F. Leroy, Compt. Rend. Hebd. Séances Acad. Sci.

227: 146 (1948) — *Streblus glaber* (Merr.) Corner, Gard. Bull. Singapore 19: 221 (1962).

*Aphananthe negrosensis* Elmer, Leafl. Philipp. Bot. 2: 575 (1909).

*Pseudostreblus caudatus* Ridl., J. Fed. Malay States Mus. 6: 54 (1915).

*Streblus laevifolius* Diels, Bot. Jahrb. Syst. 67: 171 (1935).

*Streblus urophyllus* Diels, Bot. Jahrb. Syst. 67: 172 (1935) — *Streblus glaber* (Merr.)

Corner subsp. *urophyllus* (Diels) C.C. Berg, Blumea 50: 548 (2005);

*Streblus urophyllus* Diels var. *salicifolius* Corner, Gard. Bull. Singapore. 19: 225 (1962).

1035

Note: Berg et al. (2006) treated *P. australiana* and *Streblus urophyllus* as subspecies of

*Streblus glaber*. *Paratrophis australiana*, endemic to Australia, has crenate leaf margins

and somewhat larger staminate inflorescences with more interfloral bracts than *P. glabra*;

these morphological differences are consistent and geographically confined, and we

therefore reinstate *P. australiana*. This stands in contrast to *Streblus urophyllus*, which we

provisionally maintain in synonomy under *P. glabra*. Although collections from Mt.

Wilhelm in New Guinea are remarkable for their thick coriaceous leaves and spinose

margins, similar toothed margins can be found at higher elevations in Borneo and Sulawesi,

suggesting that the striking leaf morphology of *S. urophyllus* is an alpine effect not

indicative of speciation. We reserve judgment on the status of *Streblus urophyllus* var.

*salicifolius* Corner, applied by Corner (1962) to specimens with linear leaves, provisionally

maintaining it in synonomy following Berg et al. (2006).

1048

**6. *Paratrophis insignis* (Bureau) E.M. Gardner comb. nov.** — based on *Morus insignis*

Bureau, in De Candolle, Prodr. 17: 247 (1873).

*Morus peruviana* Planch. ex Koidz., Fl. Symb. Orient.-Asiat. 88 (1930).

*Morus trianae* J.-F.Leroy, Bull. Mus. Hist. Nat. (Paris) Sér. 2, 21: 731 (1949).

*Morus marmolii* Legname, Lilloa 33: 334 (1973)

Note: The infructescences differ from most other *Paratrophis* in their fleshy free tepals and

denser aggregation of flowers, but the staminate inflorescences are typical of the genus.

1056

1057    **7. *Paratrophis microphylla* (Raoul) Cockayne, Bot. Notes Kennedy's Bush & Scenic Res.**

1058         Port Hills, Lyttelton (Rep. Scenery Preserv.) 3 (1915) – *Epicarpurus microphyllus*

1059         Raoul, Choix Pl. Nouv.-Zel. 14. (1846) – *Taxotrophis microphylla* (Raoul) F. Muell.,

1060         Fragm. (Mueller) 6(47): 193 (1868).

1061          *Paratrophis heterophylla* Blume, Mus. Bot. 2(1-8): 81. (1856).

1062

**8. *Paratrophis pendulina* (Endl.) E.M. Gardner, comb. nov.** — based on *Morus*

*pendulina* Endl., Prodromus Florae Norfolkicae 40 (1833) – *Pseudomorus pendulina*

(Endl.) Stearn, J. Arnold Arbor. 28: 427 (1947) – *Pseudomorus brunoniana* var.

*pendulina* (Endl.) Bureau, Ann. Sci. Nat. Bot. sér. 5, 11: 372 (1869) – *Streblus*

*pendulinus* (Endl.) F. Muell., Fragmenta Phytographiae Australiae (1868).

*Morus brunoniana* Endl., Atakta Bot. t. 32 (1835) – *Streblus brunonianus* (Endl.) F.

Muell., Fragm. Phyt. Australiae 6: 192 (1868) – *Pseudomorus brunoniana* (Endl.)

Bureau, Ann. Sci. Nat., Bot. sér. 5, 11: 372 (1869)

*Pseudomorus brunoniana* var. *australiana* Bureau, Ann. Sci. Nat., Bot. sér. 5, 11: 373

(1869) – *Pseudomorus pendulina* var. *australiana* (Bureau) Stearn, J. Arnold Arbor.

28: 427 (1947).

*Pseudomorus brunoniana* var. *australiana* subvar. *castaneaefolia* Bureau, Ann. Sci.

Nat., Bot. sér. 5, 11: 372 (1869).

*Pseudomorus brunoniana* var. *obtusa* Bureau, Ann. Sci. Nat., Bot. sér. 5, 11: 373

(1869) – *Pseudomorus pendulina var. obtusa* (Bureau) Stearn, J. Arnold Arbor. 28:

428 (1947), as var. *obtusata*.

*Pseudomorus sandwicensis* O. Deg., Fl. Hawaiiensis fam. 38 (1938) – *Pseudomorus*

*brunoniana* var. *sandwicensis* (O. Deg.) Skottsb., Acta Horti Gothob. 15: 347

(1944) – *Pseuomorus pendulina* var. *sandwicensis* (O. Degener) Stearn, J. Arnold

Arbor. 28: 428 (1947) – *Streblus sandwicensis* (O. Deg.) H. St. John, Pacific Trop.

Bot. Gard. Mem. 1: 374. 1973.

Note: This complex requires further investigation. Although the types of *Morus pendulina*,

*Morus brunoniana*, *Pseudomorus sandwicensis*, and *Pseudomorus brunoniana* var. *obtusa*

differ from one another, it is difficult to discern clear geographic patterns that warrant

maintaining separate species, and we therefore provisionally treat them all as one

widespread and variable species.

1089

**9. *Paratrophis philippinensis*** (Bureau) Fern.-Vill., Nov. App. 98 (1880) — *Uromorus*

*philippinensis* Bureau in A.DC., Prodr. 17: 237 (1873) — *Trophis philippinensis* (Bureau)

Corner, Gard. Bull. Singapore 19: 231 (1962).

*Sloetia minahassae* Koord., Versl. Minahasa 645 (1898).

*Paratrophis grandifolia* Elmer, Leafl. Philipp. Bot. 5: 1814 (1913).

*Calpidochlamys branderhorstii* Diels, Bot. Jahrb. Syst. 67: 173 (1935) — *Trophis*

*branderhorstii* (Diels) Corner, Gard. Bull. Singapore 19: 231 (1962).

*Calpidochlamys drupacea* Diels, Bot. Jahrb. Syst. 67: 173 (1935) — *Trophis drupacea*

(Diels) Corner, Gard. Bull. Singapore 19: 231 (1962).

Note: This species is the only *Paratrophis* with fused tepals and one of only two whose

tepals develop into a fleshy accessory fruit. The staminate inflorescences, however, are

typical of the genus.

1102

1103    **10. *Paratrophis sclerophyllus* (Corner) E.M. Gardner, comb. nov.** — based on *Streblus*

1104        *sclerophyllus* Corner, Blumea 18: 399 (1970).

1105

1106    **11. *Paratrophis smithii*** Cheeseman T.N.Z.L 20:148 (1888) – *Streblus smithii* (Cheeseman)

1107        Corner Gard. Bull. Singapore 19: 224 (1962).

1108    Note: Morphologically very close to *P. anthropophagorum*.

1109

1110    **12. *Paratrophis solomonensis* (Corner) E.M. Gardner, comb. nov.** — based on *Streblus*

1111        *solomonensis* Corner, Gard. Bull. Singapore 19: 224 (1962).

1112        Note:— Morphologically very close to *P. anthropophagorum*.

1113

1114

1115                                            ***Sorocea***

1116

1117    *Sorocea* A. St-Hil., Mém. Mus. Hist. Nat. 7: 473 (1821).

1118        *Balanostreblus* Kurz, J. Asiat. Soc. Bengal., Pt. 2, Nat. Hist., 42: 247 (1873), p.p.

1119        *Pseudosorocea* Baill., Hist. Pl. 6: 210 (1875).

1120        *Trophisomia* Rojas Acosta, Bull. Acad. Inst. Geogr. Bot. 24: 211 (1914).

1121        *Paraclarisia* Ducke, Arq. Serv. Florest. 1(1): 2 (1939).

1122

Dioecious trees. ***Leaves*** alternate, distichous, lamina pinnately veined, 3-lobed to entire.

***Stipules*** free, lateral. ***Inflorescences*** solitary or paired in the leaf axils or below the leaves,

racemose to spicate to subcapitate or uniflorous, bracts basally attached to peltate.

Staminate flowers 4-lobed to 4-parted, tepals decussate-imbricate, stamens (3–)4, straight

in bud, pistillode usually absent. Pistillate flowers tubular, 4-lobed to sub-entire, ovary

basally adnate to the perianth, stigmas 2, short, usually tongue-shaped. ***Infructescences***

with drupaceous fruits enclosed by fleshy enlarged perianths, the latter red to orange,

turning black at maturity. *Seeds* large, testa thin, embryo green, cotyledons very unequal,

the smaller one minute and enclosed by the larger.

1132

1133 Species and distribution: 19 species in Central and South America from Mexico to

1134 　　Argentina.

1135

1136 Note: This species list follows the *Flora Neotropica* monograph (Berg 2001) with the

1137 addition of four new species and one status change published later. Names synonymized by

1138 Berg (2001) but recognized by Burger et al. (1962) are noted in parentheses. Complete

1139 synonymies can be found in those treatments.

1140

*Sorocea* subg. *Sorocea*

*Sorocea affinis* Hemsl., Biol. Cent.-Amer., Bot. 3: 150 (1883).

*Sorocea angustifolia* Al.Santos & Romaniuc, Novon 24: 199 (2015).

*Sorocea bonplandii* (Baill.) W.C. Burger, Lanj. & de Boer, Acta Bot. Neerl. 11: 465

(1962).

*Sorocea briquetii* J.F.Macbr., Publ. Field Columb. Mus., Bot. Ser. 11: 16 (1931) (incl.

*S. pileate* W.C. Burger).

*Sorocea carautana* M.D.M.Vianna, Carrijo & Romaniuc, Novon 19: 549 (2009).

*Sorocea ganevii* R.M.Castro, Neodiversity 1: 18 (2006).

*Sorocea guilleminiana* Gaudich., Voy. Bonite, Bot. 3: t. 74 (1843) (incl. *S.*

*klotzschiana* Baill. and *S. macrogyna* Lanj. & Wess. Boer).

*Sorocea hilarii* Gaudich., Voy. Bonite, Bot. 3: t. 71 (1843) (incl. *S. racemosa*

Gaudich.).

*Sorocea jaramilloi* C.C.Berg, Novon 6: 241 (1996).

*Sorocea longipedicellata* A.F.P. Machado, M.D.M. Vianna & Romaniuc, Syst. Bot.

38: 687 (2013).

*Sorocea muriculata* Miq., C.F.P. von Martius & auct. suc. (eds.), Fl. Bras. 4(1): 113

(1853).

**subsp. *muriculata*** (incl. *S. amazonica* Miq.)

**subsp. *uaupensis*** (Baill.) C.C. Berg, Proc. Kon. Ned. Akad. Wetensch. C 88: 387

1161           (1985) (incl. *S. guayanensis* W.C. Burger).

1162      *Sorocea pubivena* Hemsl., Biol. Cent.-Amer., Bot. 3: 150 (1883).

1163      **subsp. *pubivena*** (incl. *S. cufodontii* W.C. Burger)

1164      **subsp. *hirtella*** (Mildbr.) C.C. Berg, Novon 6: 243 (1996) (incl. *S. opima* J.F.

1165           Macbr.)

1166      **subsp. *oligotricha*** (Akkermans & C.C. Berg) C.C. Berg, Novon 6: 243 (1996).

1167      *Sorocea ruminata* C.C.Berg, Novon 6: 244 (1996).

1168      *Sorocea sarcocarpa* Lanj. & Wess. Boer, Acta Bot. Neerl. 11: 452 (1962).

1169      *Sorocea steinbachii* C.C. Berg, Proc. Kon. Ned. Akad. Wetensch. C 88: 385 (1985).

1170      *Sorocea trophoides* W.C. Burger, Acta Bot. Neerl. 11: 450 (1962).

1171

1172  *Sorocea* subg. *Paraclarisia* (Ducke) W.C. Burger, Lanj. & Wess. Boer, Acta Bot. Neerl.

1173      11: 468 (1962).

1174      *Sorocea duckei* W.C.Burger, Acta Bot. Neerl. 11: 473 (1962).

1175      *Sorocea sprucei* (Baill.) J.F.Macbr., Publ. Field Mus. Nat. Hist., Bot. Ser. 11: 16

1176           (1931).

1177      **subsp. *sprucei*** (incl. *S. arnoldoi* Lanj. & Wess. Boer).

1178      **subsp. *saxicola*** (Hassl.) C.C.Berg, Proc. Kon. Ned. Akad. Wetensch., Ser. C.,

1179           Biol. Med. Sci. 88: 391 (1985).

1180      *Sorocea subumbellata* (C.C. Berg) Cornejo, Novon 19: 297 (2009).

1181

1182

1183                                    ***Taxotrophis***

1184  *Taxotrophis* Blume, Ann. Mus. Bot. Lugduno-Batavi 2 77 (1856); Hutch., Bull. Misc.

1185      Inform. Kew 147  (1918).

1186  *Streblus* Lour. sect. *Taxotrophis* (Blume) Corner, Gard. Bull. Singapore 19: 218 (1962);

1187      Berg et al., Fl. Males. Ser. 1, Vol. 17, Pt. 2 (2006).

1188

Monoecious or dioecious trees or shrubs, usually with lateral or terminal thorns. **Leaves** distichous, pinnately veined, petioles adaxially pubescent. Stipules free, lateral. **Inflorescences** axillary, solitary or paired, with an abaxial sterile groove, interfloral bracts basally attached, flowers with 4 free tepals, valvate. Staminate inflorescences spicate to sub-capitate, flowers with filaments inflexed in bud, pistillode present. Pistillate inflorescences uniflorous or sub-spicate, flowers usually pedicellate, stigma bifid, arms equal. **Fruits** drupaceous, up to ca. 1 cm long, usually loosely enclosed by the enlarged persistent tepals. **Seeds** up to ca. 8 x 6 mm, cotyledons subequal to unequal.

Species and distribution: Six species, ranging from Sri Lanka to New Guinea.

**1. *Taxotrophis ilicifolia*** S. Vidal, Revis. Pl. Vasc. Filip. 249 (1886).

    *Balanostreblus ilicifolia* Kurz, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 42(4): 248 (1874), p.p.

    *Pseudotrophis laxiflora* Warb., Bot. Jahrb. Syst. 13 294 (1891).

    *Taxotrophis obtusa* Elmer, Leafl. Philipp. Bot. 5 1813 (1913).

    *Taxotrophis laxiflora* Hutch., Bull. Misc. Inform. Kew 151 (1918). – *Streblus laxiflorus* (Hutch.) Corner, Gard. Bull. Singapore 19 229 (1962).

    *Taxotrophis triapiculata* Gamble, Bull. Misc. Inform. Kew 188 (1913).

    *Taxotrophis eberhardtii* Gagnep., Fl. Indo-Chine 5 700 (1928).

    *Taxotrophis macrophylla* auct. non Boerl.: Burkill, Dict. Econ. Prod. Malay Penins. 2126 (1935).

    *Streblus ilicifolius* (S. Vidal) Corner, Gard. Bull. Singapore 19: 227 (1962).

**2. *Taxotrophis macrophylla*** (Blume) Boerl., Handl. Fl. Ned. Ind. 3: 359 (1900).

    *Streblus macrophyllus* Blume, Ann. Mus. Bot. Lugduno-Batavi 2: 80 (1856) — *Diplocos ? macrophyllus* (Blume) Bureau in A.DC., Prodr. 17: 216 (1873).

    *Pseudotrophis mindanaensis* Warb. in Perkins, Fragm. Fl. Philipp. 1: 165 (1905); Elmer, Leafl. Philipp. Bot. 5: 1815 (1913), '*Taxatrophis mindanaensis*' in nota.

    *Paratrophis caudata* Merr., Philipp. J. Sci., 1, Suppl. 183 (1906).

1219    *Taxotrophis balansae* Hutch., Bull. Misc. Inform. Kew 151 (1918).

1220    *Dimerocarpus brenieri* Gagnep., Bull. Mus. Hist. Nat. (Paris) 27: 441 (1921).

1221

1222    **3. *Taxotrophis perakensis* (Corner) E.M. Gardner, comb. nov.**, based on *Streblus*

1223        *perakensis* Corner, Gard. Bull. Singapore 19: 223 (1962).

1224

1225    Note: Although Corner (1962) considered *S. perakensis* species part of section *Paratrophis*,

1226    albeit with some hesitation, Berg et al. (2006), whom we follow, placed it in *Taxotrophis*

1227    based on the spines that appear on some specimens. This is a rather variable species that

1228    requires further investigation to properly elucidate its limits and affinities.

1229

1230    **4. *Taxotrophis spinosa*** Steenis in Backer & Bakh.f., Fl. Java 2 (1965) 16.

1231        *Urtica spinosa* Blume, Bijdr. (1825) 507. — *Streblus spinosus* (Blume) Corner, Gard.

1232            Bull. Singapore 19: 229 (1962).

1233        *Taxotrophis javanica* Blume, Ann. Mus. Bot. Lugduno-Batavi 2: 77, t. 26 (1856).

1234

1235    **5. *Taxotrophis taxoides*** (B. Heyne ex Roth) W.L. Chew ex E.M. Gardner, comb. nov.

1236        *Trophis taxoides* B. Heyne ex Roth, Nov. Pl. Sp. 368 (1821). — *Trophis taxiformis*

1237            Spreng., Syst. Veg. 3 902 (1826), nom. nov. illeg. — *Streblus taxoides* (B. Heyne

1238            ex Roth) Kurz, Forest Fl. Burma 2: 465 (1877) — *Phyllochlamys taxoides* (B.

1239            Heyne ex Roth) Koord., Exkurs.-Fl. Java 2: 89 (1912).

1240        *Trophis spinosa* Roxb., Fl. Ind., ed. Carey 3: 762 (1832), non Willd. 1806, nec Blume

1241            1826. — *Epicarpurus spinosus* (Roxb.) Wight, Icon. Pl. Ind. Orient. 6: 7, t. 1962

1242            (1853), p.p. — *Phyllochlamys spinosa* (Roxb.) Bureau in A.DC., Prodr. 17: 218

1243            (1873);

1244        *Epicarpurus timorensis* Decne., Nouv. Ann. Mus. Hist. Nat. 3: 499, t. 21 (1834).

1245        *Taxotrophis roxburghii* Blume, Ann. Mus. Bot. Lugduno-Batavi 2: 78 (1856)

1246        *Streblus microphyllus* Kurz, Prelim. Rep. Forest Pegu App. A, cxviii; App. B, 84

1247            (1875); — *Streblus taxoides* (B. Heyne ex Roth) Kurz var. *microphylla* (Kurz)

1248            Kurz, Forest Fl. Burma 2: 465 (1877).

1249         *Phyllochlamys wallichii* King ex Hook.f., Fl. Brit. India 5: 489 (1888);

1250         *Phyllochlamys taxoides* (B. Heyne ex Roth) Koord. var. *parvifolia* Merr., Enum.

1251             Philipp. Flow. Pl. 2: 38 (1923).

1252         *Taxotrophis poilanei* Gagnep., Fl. Indo-Chine 5: 701 (1928).

1253         *Taxotrophis crenata* Gagnep., Fl. Indo-chine 5: 702, t. 82 (1928). — *Streblus crenatus*

1254             (Gagnep.) Corner, Gard. Bull. Singapore 19: 226 (1962).

1255         *Phyllochlamys tridentata* Gagnep., Fl. Indo-Chine 5: 714 (1928).

1256

1257 Note: In the 1950s, Dr. Chew Wee Lek annotated quite a lot of specimens at K and SING

1258 with the new combination *Taxotrophis taxoides*. However, the combination was never

1259 published, probably because the need for the combination was obviated in 1962 when

1260 Corner, his doctoral supervisor, reduced *Taxotrophis* to a section of *Streblus*.

1261

1262 **6. *Taxotrophis zeylanica*** (Thwaites) Thwaites, Enum. Pl. Zeyl. [Thwaites] 264 (1861).

1263         *Epicarpurus zeylanicus* Thwaites, Hooker's J. Bot. Kew Gard. Misc. 4: 1 (1852) —

1264             *Diplocos zeylanica* (Thwaites) Bureau, Prodr. [A. P. de Candolle] 17: 215 (1873)

1265             — *Streblus zeylanicus* (Thwaites) Kurz, Forest Fl. Burma 2: 464 (1877).

1266         *Taxotrophis caudata* Hutchinson, Bull. Misc. Inform. Kew 1918(4): 149 (1918).

1267

1268

1269                                     *Trophis*

1270

1271 *Trophis* Lour., P. Browne, Civ. Nat. Hist. Jamaica 357 (1756), nom. cons.

1272         *Bucephalon* L. Sp. Pl. 1190 (1753), nom. rejic.

1273         *Skutchia* Pax & K. Hoffm. ex C.V. Morton, J. Wash. Acad. Sci. 27: 306 (1937).

1274

Dioecious trees or shrubs. ***Leaves*** distichous, pinnately veined. Stipules free, lateral.

***Inflorescences*** axillary or just below the leaves, solitary or paired, interfloral bracts basally

attached. Staminate inflorescences spicate to racemose with an abaxial sterile groove, tepals

4, basally connate, stamens 4, filaments inflexed in bud, pistillode present. Pistillate

inflorescences spicate to racemose or subcapitate, tepals 4, connate, forming a tubular

perianth, ovary adnate to the perianth or not, stigma bifid, arms equal. *Fruits* drupaceous,

up to ca. 1.5 cm long, adnate to the perianth or not, the perianth enlarged and fleshy or not.

*Seeds* up to ca. 1 cm long, cotyledons equal.

1283

1284    Species and distribution: Five species in the neotropics.

1285

1286    *Trophis* P. Browne sect. *Trophis*

1287

1288    **1. *Trophis cuspidata*** Lundell, Amer. Midl. Naturalist 19: 427 (1938).

1289    **2. *Trophis mexicana*** (Liebm.) Bureau, Prodr. [A. P. de Candolle] 17: 253 (1873).

1290    **3. *Trophis noraminervae*** Cuevas & Carvajal, Acta Bot. Mex. 47: 2 (1-7, fig.) (1999).

1291    **4. *Trophis racemosa*** Urb., Symb. Antill. (Urban). 4(2): 195 (1905).

1292

1293    *Trophis* P. Browne sect. *Echinocarpa* C.C. Berg, Proc. Kon. Ned. Akad. Wetensch., Ser. C,

1294    Biol. Med. Sci. 91: 353 (1988).

1295

1296    **5. *Trophis involucrata*** W.C. Burger, Phytologia 26: 432 (1973).

1297

1298

1299                              **Tribe OLMEDIEAE**

1300

1301    Olmedieae Trécul, Ann. Sci. Nat., Bot. sér. 3, 8 (1847) 126

1302          Castilleae C.C. Berg, Acta Bot. Neerl. 26 (1977) 78

1303          Strebleae Bureau in DC., Prodr. 17: 215 (1873), p.p.

1304

Tree or shrubs, monoecious or dioecious (or androdioecious), mostly with self-pruning

branches. *Leaves* alternate or opposite, distichous or spirally arranged; stipules lateral to

amplexicaul. *Inflorescences* mostly unisexual, capitate, mostly discoid to urceolate,

involucrate, tepals mostly 4, connate or not. Staminate inflorescences usually many-

flowered; stamens 4 or fewer, with filaments straight or less often inflexed in bud, pistillode

mostly absent. Pistillate inflorescences one to many-flowered, ovary free or not, stigmas 2,

filiform. **Fruits** mostly drupaceous, mostly enclosed by a fleshy perianth or embedded in a

fleshy receptacle. **Seeds** with or without endosperm, testa thin, vascularized, cotyledons

mostly equal.

1314

1315  Genera and distribution: 13 genera with 63 species. Eight neotropical genera: *Castilla* (3

1316  spp.), *Helicostylis* (7), *Maquira* (4), *Naucleopsis* (22), *Olmedia* (1), *Perebea* (9),

1317  *Poulsenia* (1), and *Pseudolmedia* (9); and five Paleotropical genera: the widespread

1318  *Antiaris* (1); *Antiaropsis* (2) in New Guinea; *Mesogyne* (1) in Africa; *Sparattosyce* (1)

1319  in New Caledonia; and *Streblus* (3) in South to Southeast Asia.

1320

1321  *Olmedia*

1322

1323  *Olmedia* Ruiz & Pav., Syst. Veg. Fl. Peruv. Chil.1:257.1798. Type — *Olmedia aspera* Ruiz

1324  & Pav.

1325  *Trophis* section *Olmedia* (Ruiz & Pav.) Berg, Proc. Kon. Ned. Acad. Wetensch. C. 91:

1326  354. 1988.

1327

Dioecious trees or shrubs. **Leaves** distichous, lamina pinnately veined. **Stipules** free, not

fully amplexicaul. **Inflorescences** unisexual, with a well-developed involucre. Staminate

inflorescences discoid, multiflorous; tepals 4, valvate, stamens 4, inflexed in bud, pistillode

absent. Pistillate inflorescences usually uniflorous, perianth tubular, 4-dentate,  ovary free,

stigmas 2, equal. **Fruits** drupaceous, surrounded by fleshy persistent perianth and

subtended by spreading, fleshy involucral bracts. **Seeds** ca. 5 mm long, cotyledons equal.

1334

1335  Species and distribution: One species, in the Neotropics.

1336

1337  **1. Olmedia aspera** Ruiz & Pav., Syst. Veg. Fl. Peruv. Chil.1:257.1798

1338        *Olmedia caucana* Pittier, Contr. U.S. Natl. Herb. 13:434. 1912. — *Trophis caucana*

1339            (Pitttier) C.C. Berg, Proc. Kon. Ned. Acad. Wetensch. C. 91: 354. 1988

1340        *Olmedia poeppigiana* Klotzsch, Linnaea 20: 525. 1847, as a synonym of *O. aspera*

1341            Poeppig & Endlicher, Nov. Gen. 2: 31. 1838, based on Poeppig s.n. or 1267, non

1342            *O.poeppigiana* Martius Flora (or Bot. Zeit) 24 (Beibl. 2): 93. 1841 (= *Helicostylis*

1343            *tomentosa* (Poeppig & Endlicher) Rusby).

1344        *Olmedia falcifolia* Pittier, Contr. U.S. Natl. Herb. 13: 435. 1912.

1345        *Trophis aurantiaca* Herzog, Repert. Spec. Nov. 7: 51. 1909

1346

1347

1348                            ***Streblus***

1349

*Streblus* Lour., Fl. Cochinch. (1790) 615.

*Achymus* Juss., Dict. Sci. Nat. 1, Suppl. 31 (1816) .

*Epicarpurus* Blume, Bijdr. 488 (1825).

*Albrandia* Gaudich. in Freyc., Voy. Uranie, Bot. 509 (1830).

*Calius* Blanco, Fl. Filip. 698 (1837).

*Teonongia* Stapf, Hooker's Icon. Pl. 30: t. 2947 (1911).

*Diplothorax* Gagnep., Bull. Soc. Bot. France 75 98 (1928).

1357

Trees or shrubs, dioecious or monoecious. ***Leaves*** distichous, lamina pinnately veined.

***Stipules*** free, lateral. ***Inflorescences*** bisexual or unisexual, capitate, with a rudimentary

involucre, bracts basally attached. Staminate inflorescences discoid capitate, multiflorous;

tepals 4, imbricate, stamens 4, inflexed in bud, pistillode present but small. Pistillate

inflorescences usually uniflorous, tepals 4, ovary free, stigmas 2, equal. ***Fruits*** drupaceous,

up to ca. 8 mm long, initially enclosed by enlarged but not fleshy tepals, which may open

later. ***Seeds*** ca. 5 mm long, cotyledons equal or very unequal.

1365

1366   Species and distribution: three species from India to South China and from mainland

1367   Southeast Asia to the Philippines and the Moluccas.

1368

**1.** *Streblus asper* (Retz.) Lour., Fl. Cochinch. 2: 615 (1790).

    *Trophis aspera* Retz., Observ. Bot. 5 (1788) – *Epicarpurus asper* (Retz.) Steud.

      Nomencl. Bot. ed. 2, 1: 556 (1840).

    *Trophis cochinchinensis* Poir., Encycl. 8 (1808) 123.

    *Epicarpurus orientalis* Blume, Bijdr. (1825) 488

    *Calius lactescens* Blanco, Fl. Filip. (1837) 698; ed. 3, 3 (1879) 1103, t. 171. —

      *Streblus lactescens* (Blanco) Blume, Ann. Mus. Bot. Lugduno-Batavi 2 (1856) 80.

    *Achymus pallens* Sol. ex Blume, Ann. Mus. Bot. Lugduno-Batavi 2 (1856) 79.

    *Cudrania crenata* C.H. Wright, J. Linn. Soc., Bot. 26 (1899) 469. — *Vanieria crenata*

      (C.H. Wright) Chun, J. Arnold Arbor. 8 (1927) 21.

    *Diplothorax tonkinensis* Gagnep., Bull. Soc. Bot. France 75 (1928) 98.

Note: Retzius's *Trophis aspera* predated Loureiro's *Streblus asper* by two years, making

the latter an implied new combination under Article 41.1 of the Code (cf. Ex. 10).

Synonymies of the *S. asper* have sometimes erroneously included *Trophis aculeata* Roth,

which is actually *Maclura spinosa* (Roxb. ex Willd.) C.C. Berg.

**2.** *Streblus celebensis* C.C. Berg, Blumea 50 547 (2005).

**3.** *Streblus tonkinensis* (Eberh. & Dubard) Corner, Gard. Bull. Singapore 19 (1962): 228.

    *Bleeokrodea tonkinensis* Eberh. & Dubard, Compt. Rend. Hebd. SÈances Acad. Sci.

      145 (1907): 632. — *Teonongia tonkinensis* (Eberh. & Dubard) Stapf.

**Tribe DORSTENIEAE**

Dorstenieae Gaudich. in Freyc., Voy. Uranie, Bot. (1830).

    Broussonetieae Gaudich. in Freyc., Voy. Uranie, Bot. 508 (1830).

Brosimeae Trécul, Ann. Sci. Nat., Bot. sér. 3, 8: 146 (1847).

1398   Fatoueae Engl., Nat. Pflanzenfam. 3, 1: 71 (1888).

1399

Tree, shrubs, lianas, and herbs, monoecious or less often dioecious. *Leaves* alternate or less

commonly (sub)opposite, distichous or spirally arranged; stipules lateral to fully

amplexicaul. *Inflorescences* unisexual or bisexual, cymose, spicate, globose, or discoid to

turbinate or cup-shaped, multiflorous or uniflorous (the latter in pistillate inflorescences

only), bracteate or not, interfloral bracts mostly peltate. Staminate flowers with tepals (1–

)2–4 or absent, stamens 1–4 with filaments straight or inflexed in bud, pistillode present or

(more often) absent. Pistillate flowers free or connate or embedded in the receptacle, tepals

2–4, ovary free or not, stigmas 1 or 1, equal or unequal. *Fruits* drupaceous or drupe-like

due to a persistent fleshy perianth and/or receptacle, the whole endocarp unit often

ballistically ejected from the infructescence. *Seeds* with or without endosperm, large or

small; cotyledons equal or unequal.

1411

Genera and distribution: eleven genera and 63 species with a worldwide distribution:

*Bleekrodea* (Africa and Southeast Asia)*, Bosqueiopsis* (Africa)*, Brosimum* (Neotropics)*,*

*Broussonetia* (Asia to Oceania; introduced worldwide)*, Dorstenia* (Africa and South

America, with one species in India)*, Fatoua* (Madagascar and Japan to New Caledonia;

introduced worldwide), *Helianthostylis* (Neotropics)*, Malaisia* (Southeast Asia to New

Caledonia)*, Scyphosyce* (Africa)*, Sloetia* (Southeast Asia)*, Sloetiopsis* (Africa)*, Trilepisium*

(Africa)*, Trymatococcus* (Neotropics), *Utsetela* (Africa).

1419

1420

1421                                  ***Sloetiopsis***

1422

1423   *Sloetiopsis* Engl., Bot. Jahrb. 39: 573 (1907).

1424        *Neosloetiopsis* Engl., Bot. Jahrb. 51: 426 (1914)

1425        *Streblus* auct. non Lour., C.C. Berg, Proc. Kon. Ned. Acad. Wetensch. C. 91: 357.

1426        1988.

1427

Tree or shrubs, dioecious (or monoecious). *Leaves* distichous, pinnately veined, cystoliths

present; stipules free, nearly amplexicaul. *Inflorescences* unisexual (or bisexual); staminate

inflorescences spicate with an abaxial sterile groove, bracts mostly peltate, tepals 4, free or

basally connate filaments inflexed in bud, pistillode small; pistillate inflorescences

uniflorous, bracts mostly basally attached, tepals 4, free, imbricate, 2 stigmas. *Fruits*

drupaceous, fleshy endocarp dehiscent, tepals enlarged and persistent but not fleshy. *Seeds*

globose, ca. 4 mm, endocarp coriaceous with a hard disc against the hilum, testa

vascularized with a thick apical cap, cotyledons equal.

1436

1437 **1.** *Sloetiopsis usambarensis* Engl., Bot. Jahrb. 39: 573 (1907) — *Streblus usambarensis*

1438    (Engl.) C.C. Berg, Proc. Kon. Ned. Acad. Wetensch. C. 91: 357. 1988.

1439    *Neosloetiopsis kamerunensis* Engl., Bot. Jahrb. 51: 426 (1914).

1440

1441 Note. The nearly amplexicaul stipules and occasional bisexual inflorescences reflect an

1442 affinity with the Southeast Asian *Sloetia*, also a member of the Dorstenieae.

1443

1444

1445                          **Tribe PARARTOCARPEAE**

1446

Parartocarpeae N.J.C. Zerega & E.M. Gardner Zerega, Phytotaxa 388, 253–265 (2019).

Shrubs to large trees, monoecious or dioecious; abundant white exudate. *Leaves* distichous

or spirally arranged; simple; entire; pinnately veined; thin to thick coriaceous; glabrous,

pubescent, scabrid, or hispid pubescent. *Stipules* axillary, simple or paired, lateral.

*Inflorescences* solitary or paired in leaf axils, unisexual (or bisexual with a single apical

pistillate flower), uniflorous, cymose, or capitate with stamens or ovaries sunken into the

receptacle; pedunculate; involucre of 3–8 triangular bracts, basally connate. Staminate

inflorescences with numerous flowers, tepals 4–5, stamens 5 and normally positioned or 1–

3 in cavities in the receptacle with the anthers exerted through perforations in the upper

surface of the receptacle, filaments free or united. Pistillate inflorescences uniflorous or

(sub)globose with ovaries solitary in each cavity, unilocular, the style apical with a short

exerted stigma. ***Fruits*** drupaceous, enclosed by persistant tepals, or aggregated into

syncarps formed by the enlargement of the entire inflorescence.

1460

1461     Genera and distribution: Three genera and five species, from Southern China to the

1462     Solomon Islands: *Hullettia*, *Parartocarpus*, and *Pseudostreblus*.

1463

1464

1465                                     *Pseudostreblus*

1466

1467     *Pseudostreblus* Bureau, in A.P. de Candolle, Prodr. 17: 220 (1873).

1468     *Streblus* Lour. sect. *Pseudostreblus* (Bureau) Corner, Gard. Bull. Singapore 19: 217 (1962).

1469

Monoecious trees. ***Leaves*** distichous, pinnately veined. Stipules free, lateral.

***Inflorescences*** axillary. Staminate inflorescences cymose, tepals 5, imbricate, filaments

inflexed in bud but apparently straightening gradually upon anthesis, pistillode minute,

conical, pubescent, interfloral bracts few, basally attached. Pistillate inflorescences

uniflorous, pedunculate, involucral bracts 3, ±connate, tepals 4, imbricate, stigma bifid,

arms equal. ***Fruits*** drupaceous, ca. 1 cm long, with tepals enlarged and loosely enclosing

the fruit. ***Seeds*** ellipsoid, ca. 6 x 8 mm, cotyledons unequal.

1477

1478     Species and distribution: One, in South to Southeast Asia and Southern China.

1479

1480     Note: *Pseudostreblus* is remarkable for its five-parted staminate flowers.

1481

1482     **1.** ***Pseudostreblus indicus*** **Bureau**, in A.P. de Candolle, Prodr. 17: 220 (1873).

1483        *Streblus indicus* (Bureau) Corner, Gard. Bull. Singapore 19: 226 (1962).

1484

1485

1486     **Conclusion**

1487     The revisions presented here, based on the best phylogenetic evidence available to date,

1488     provide for seven monophyletic tribes of Moraceae and eleven monophyletic genera within

1489     Moreae. We hope that the resulting taxonomic and phylogenetic framework will provide a

1490     solid foundation for further research into the systematics and evolution of Moraceae.

1491

1492

**Author contributions**

EMG led the writing of the manuscript, did field, lab, and herbarium work, conducted all

phylogenetic analyses, and was responsible for the taxonomic treatment. MG performed lab

work for the Moraceae333 samples. RC, SD, NE, and OM performed lab work for the

PAFTOL samples, and OM curated the PAFTOL data. DA collaborated on all herbarium

work at BO. Sahromi facilitated access and participated in field collections at the Bogor

Botanical Gardens. NJCZ provided overall guidance on Moraceae and facilitated

sequencing at Northwestern University. WJB and FF acquired funding for and supervised

the PAFTOL project. AM performed herbarium sampling at K and supervised the

Moraceae portion of the PAFTOL project. EMG designed the overall study with input from

ALH and NJCZ. ALH supervised the overall study. All authors commented on and

contributed to the manuscript.

1505

1506

**Acknowledgements**

This work was supported by the United States National Science Foundation (DBI award

number 1711391, and DEB award numbers 0919119 and 1501373); the SYNTHESYS

Project http://www.synthesys.info/ which is financed by European Community Research

Infrastructure Action under the FP7 "Capacities" Program"; the National Parks Board of

Singapore through a *Flora of Singapore* Fellowship and a Research Fellowship; grants

from the Calleva Foundation and the Sackler Trust to the Plant & Fungal Trees of Life

Project at the Royal Botanic Gardens, Kew. We thank the Pritzker Laboratory for

Molecular Systematics at the Field Museum of Natural History (K. Feldheim) for access to

laboratory and sequencing facilities; S. Knapp and J. Wajer (BM) for helpful consultation

on nomenclatural issues; W. Clement for helpful comments on the circumscription of the

Olmedieae; the Bogor Botanical Gardens, Singapore Botanic Gardens, and The Morton

Arboretum for access to living collections; and the following herbaria for access to

collections for examination and/or sampling: BM, BKF, BO, F, IBSC, K, FTBG, L, MO,

NY, SING, SAN, SAR, SNP, and U. Finally, we gratefully acknowledge the careful work

of past scholars of Moraceae, in particular the late C.C. Berg and E.J.H. Corner.

1523

1524 **Conflicts of interest**

1525 The authors are unaware of any conflicts of interest affecting this research.

1526

1527 **Literature cited**

1528

**Bawa, K., & Crisp, J.** 1980. Wind-pollination in the understorey of a rain forest in Costa

Rica. *J. Ecol.*, 68: 871–876. Retrieved from

http://www.jstor.org/stable/10.2307/2259462

**Bawa, K. S., Bullock, S. H., Perry, D. R., Coville, R. E., & Grayum, M. H.** 1985.

Reproductive biology of tropical lowland rain forest trees. II. Pollination systems. *Am.*

*J. Bot.*, 72: 346–356. https://doi.org/10.2307/2443527

**Berg, C. C.** 1977a. Revisions of African Moraceae (excluding Dorstenia, Ficus, Musanga

and Myrianthus). *Bull. Du Jard. Bot. Natl. Belgique*, 47: 267–407. Retrieved from

http://www.jstor.org/stable/10.2307/3667908

**Berg, C. C.** 1977b. The Castilleae, a tribe of the Moraceae, rename and redefined due tothe

exclusion of the type genus Olmedia from the 'Olmedieae'. *Acta Bot. Neerl.*, 26: 73–

82.

**Berg, C. C.** 1978. Cecropiaceae: A new family of the Urticales. *Taxon*, 27: 39–44.

https://doi.org/10.2307/1220477

**Berg, C. C.** 1988. The genera Trophis and Streblus (Moraceae) remodelled. *Proc. K. Ned.*

*Akad. van Wet. Ser. C, Biol. Med. Sci.*, 91: 345–362.

**Berg, C. C.** 2001. *Flora Neotropica Monograph 83: Moreae, Artocarpeae, and Dorstenia*

*(Moraceae), with Introductions to the Family and Ficus and with Additions and*

*Corrections to Flora Neotropica Monograph 7* (Vol. 83).

**Berg, C. C.** 2005. Flora Malesiana precursor for the treatment of Moraceae 8: other genera

than Ficus. *Blumea*, 50: 535–550. Retrieved from

http://www.ingentaconnect.com/content/nhn/blumea/2005/00000050/00000003/art000

16

**Berg, C. C., Corner, E. J. H., & Jarrett, F. M.** 2006. Moraceae (genera other than Ficus).

*Flora Malesiana, Ser. 1*, 17: 1–154.

**Bolger, A. M., Lohse, M., & Usadel, B.** 2014. Trimmomatic: A flexible trimmer for

Illumina Sequence Data. *Bioinformatics*, 1–7.

https://doi.org/10.1093/bioinformatics/btu170

**Brewer, G. E., Clarkson, J. J., Maurin, O., Zuntini, A. R., Barber, V., Bellot, S., Biggs,**

**N., Cowan, R. S., Davies, N. M. J., Dodsworth, S., & others**. 2019. Factors affecting

targeted sequencing of 353 nuclear genes from herbarium specimens spanning the

diversity of angiosperms. *Front. Plant Sci.*, 10: 1102.

**Bureau, E.** 1873. Moraceae and Artocarpace. in: *A. P. de Candolle, Prodromussystematis*

*naturali 17*. Paris: Treuttel & Würtz.

**Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T.** 2009. trimAl: A tool for

automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*,

25: 1972–1973. https://doi.org/10.1093/bioinformatics/btp348

**Chen, Z., Yang, T., & Lu, L.** 2016. Tree of life for the genera of Chinese vascular plants.

*J. Syst. Evol.*, 54: 273–276. https://doi.org/10.1111/jse.12219

**Chung, K. F., Kuo, W. H., Hsu, Y. H., Li, Y. H., Rubite, R. R., & Xu, W. Bin**. 2017.

Molecular recircumscription of Broussonetia (Moraceae) and the identity and

taxonomic status of B. kaempferi var. australis. *Bot. Stud.*

https://doi.org/10.1186/s40529-017-0165-y

**Clement, W. L., & Weiblen, G. D.** 2009. Morphological Evolution in the Mulberry

Family (Moraceae). *Syst. Bot.*, 34: 530–552.

https://doi.org/10.1600/036364409789271155

**Corner, E. J. H.** 1962. The Classification of Moraceae. *Gard. Bull.*, 19: 187–252.

**Corner, E. J. H.** 1970. New Species of Streblus and Ficus. *Blumea*, 18: 393–411.

**Corner, E. J. H.** 1975. The evolution of StrebIus Lour. (Moraceae): with a new species of
sect. Bleekrodea. *Phytomorphology*, 25: 1–12.

**Culley, T. M., Weller, S. G., & Sakai, A. K.** 2002. The evolution of wind pollination in
angiosperms. *Trends Ecol. Evol.*, 17: 361–369. https://doi.org/10.1016/S0169-
5347(02)02540-5

**Datwyler, S. L., & Weiblen, G. D.** 2004. THE ORIGIN OF THE FIG : PHYLOGENETIC
NDH F SEQUENCES. *Am. J. Bot.*, 91: 767–777.

**Doyle, J., & Doyle, J.** 1987. Genomic plant DNA preparation from fresh tissue-CTAB
method. *Phytochem. Bull.*, 19: 11–15.

**Engler, G. H. A.** 1889. Ulmaceae, Moraceae, and Urticaceae. in: *Engler & Prantl, Die*
*natürlichen Pflanzenfamilien 3(1)*. Leipzig: W. Engelmann.

**Filho, M. D. M. V., Carrijo, T. T., & Neto, S. R.** 2009. Sorocea carautana (Moraceae): A
New Species from Southeastern Brazil. *Novon A J. Bot. Nomencl.*
https://doi.org/10.3417/2007143

**FitzJohn, R. G.** 2012. Diversitree : comparative phylogenetic analyses of diversification in
R. *Methods Ecol. Evol.*, 3: 1084–1092. https://doi.org/10.1111/j.2041-
210X.2012.00234.x

**Gardner, E. M.** 2017. *Dissertation: Evolutionary Transitions: Phylogenomics and*
*Pollination of Artocarpus (Moraceae).*
https://doi.org/https://doi.org/10.21985/N2CF59

**Gardner, E. M., Gagné, R. J., Kendra, P. E., Montgomery, W. S., Raguso, R. A.,**
**McNeil, T. T., & Zerega, N. J. C.** 2018. A flower in fruit's clothing: Pollination of
jackfruit (artocarpus heterophyllus, moraceae) by a new species of gall midge,
clinodiplosis ultracrepidata sp. Nov. (Diptera: Cecidomyiidae). *Int. J. Plant Sci.*, 179:
350–367. https://doi.org/10.1086/697115

**Gardner, E. M., Johnson, M. G., Ragone, D., Wickett, N. J., & Zerega, N. J. C.** 2016.
Low-Coverage, Whole-Genome Sequencing of Artocarpus camansi (Moraceae) for
Phylogenetic Marker Development and Gene Discovery. *Appl. Plant Sci.*, 4: 1600017.
https://doi.org/10.3732/apps.1600017

**Hale, H., Gardner, E. M ., Viruel, J., Pokorny, L ., & Johnson, M. G.** 2020. Strategies

for reducing per-sample costs in target capture sequencing for phylogenomics and

population genomics in plants. *Appl. Plant Sci.*, in press.

**Heibl, C.** 2008. PHYLOCH: R language tree plotting tools and interfaces to diverse

phylogenetic software packages. http://www.christophheibl.de/Rpackages.html.

**Johnson, M. G., Gardner, E. M., Liu, Y., Medina, R., Goffinet, B., Shaw, A. J.,**

**Zerega, N. J. C., & Wickett, N. J.** 2016. HybPiper: Extracting Coding Sequence and

Introns for Phylogenetics from High-Throughput Sequencing Reads Using Target

Enrichment. *Appl. Plant Sci.*, 4: 1600016. https://doi.org/10.3732/apps.1600016

**Johnson, M. G., Pokorny, L., Dodsworth, S., Botigué, L. R., Cowan, R. S., Devault, A.,**

**Eiserhardt, W. L., Epitawalage, N., Forest, F., Kim, J. T., Leebens-Mack, J. H.,**

**Leitch, I. J., Maurin, O., Soltis, D. E., Soltis, P. S., Wong, G. K., Baker, W. J., &**

**Wickett, N. J.** 2019. A Universal Probe Set for Targeted Sequencing of 353 Nuclear

Genes from Any Flowering Plant Designed Using k-Medoids Clustering. *Syst. Biol.*,

68: 594–606. https://doi.org/10.1093/sysbio/syy086

**Kamneva, O. K., Syring, J., Liston, A., & Rosenberg, N. A.** 2017. Evaluating

allopolyploid origins in strawberries (Fragaria) using haplotypes generated from target

capture sequencing. *BMC Evol. Biol.*, 17: 1–19. https://doi.org/10.1186/s12862-017-

1019-7

**Kates, H. R., Johnson, M. G., Gardner, E. M., Zerega, N. J. C., & Wickett, N. J.** 2018.

Allele phasing has minimal impact on phylogenetic reconstruction from targeted

nuclear gene sequences in a case study of Artocarpus. *Am. J. Bot.*

https://doi.org/10.1002/ajb2.1068

**Katoh, K., & Standley, D. M.** 2013. MAFFT multiple sequence alignment software

version 7: improvements in performance and usability. *Mol. Biol. Evol.*, 30: 772–80.

https://doi.org/10.1093/molbev/mst010

**Kim, J., & Sanderson, M. J.** 2008. Penalized likelihood phylogenetic inference: Bridging

the parsimony-likelihood gap. *Syst. Biol.* https://doi.org/10.1080/10635150802422274

**Kress, W. J., Erickson, D. L., Jones, F. A., Swenson, N. G., Perez, R., Sanjur, O., &**

**Bermingham, E.** 2009. Plant DNA barcodes and a community phylogeny of a tropical

forest dynamics plot in Panama. *Proc. Natl. Acad. Sci. U. S. A.*, 106: 18621–18626.

https://doi.org/10.1073/pnas.0909820106

**Lewis, W. H.** 1986. Airborne pollen of the Neotropics:Potential roles in pollination and

pollinosis. *Grana*, 25: 75–83. https://doi.org/10.1080/00173138609429936

**Machado, A. F. P., Vianna Filho, M. D. M., Amorim, A. M., & Romaniuc Neto, S.**

2013. A New <I>Sorocea</I> (Moraceae) From The Atlantic Rainforest, Brazil. *Syst.*

*Bot.* https://doi.org/10.1600/036364413x670340

**Momose, K., Yamaoka, R., & Inoue, T.** 1998. Pollination Biology of the Genus

Artocarpus, Moraceae. *Tropics*, 7: 165–172. https://doi.org/10.3759/tropics.7.165

**Nepal, M.** 2012. Phylogenetics of Morus (Moraceae) Inferred from ITS and trnL-trnF

Sequence Data. *Syst. Bot.*, 37: 442–450. https://doi.org/10.1600/036364412X635485

**Paradis, E.** 2013. Molecular dating of phylogenies by likelihood methods: A comparison

of models and a new information criterion. *Mol. Phylogenet. Evol.*

https://doi.org/10.1016/j.ympev.2013.02.008

**Paradis, E., & Schliep, K.** 2019. Ape 5.0: An environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics*, 35: 526–528.

https://doi.org/10.1093/bioinformatics/bty633

**Quinlan, A. R., & Hall, I. M.** 2010. BEDTools: A flexible suite of utilities for comparing

genomic features. *Bioinformatics*, 26: 841–842.

https://doi.org/10.1093/bioinformatics/btq033

**Rabosky, D. L.** 2014. Automatic detection of key innovations, rate shifts, and diversity-

dependence on phylogenetic trees. *PLoS One*, 9.

https://doi.org/10.1371/journal.pone.0089543

**Rabosky, D. L., Donnellan, S. C., Grundler, M., & Lovette, I. J.** 2014a. Analysis and

visualization of complex Macroevolutionary dynamics: An example from Australian

Scincid lizards. *Syst. Biol.*, 63: 610–627. https://doi.org/10.1093/sysbio/syu025

**Rabosky, D. L., Grundler, M., Anderson, C., Title, P., Shi, J. J., Brown, J. W., Huang,**

**H., & Larson, J. G.** 2014b. BAMMtools: An R package for the analysis of

evolutionary dynamics on phylogenetic trees. *Methods Ecol. Evol.*, 5: 701–707.

https://doi.org/10.1111/2041-210X.12199

**Rabosky, D. L., Santini, F., Eastman, J., Smith, S. A., Sidlauskas, B., Chang, J., &**

**Alfaro, M. E.** 2013. Rates of speciation and morphological evolution are correlated

across the largest vertebrate radiation. *Nat. Commun.*, 4: 1958.

https://doi.org/10.1038/ncomms2958

**Revell, L. J.** 2012. phytools: An R package for phylogenetic comparative biology (and

other things). *Methods Ecol. Evol.*, 3: 217–223. https://doi.org/10.1111/j.2041-

210X.2011.00169.x

**Sakai, S., Kato, M., & Nagamasu, H.** 2000. Artocarpus (Moraceae)–gall midge

pollination mutualism mediated by a male-flower parasitic fungus. *Am. J. Bot.*, 87:

440–445. Retrieved from http://www.amjbot.org/content/87/3/440.short

**Sanderson, M. J.** 2002. Estimating absolute rates of molecular evolution and divergence

times: A penalized likelihood approach. *Mol. Biol. Evol.*

https://doi.org/10.1093/oxfordjournals.molbev.a003974

**Santos, A. dos, & Neto, S. R.** 2015. Sorocea angustifolia (Moraceae), a New Species of

Sorocea from the Atlantic Forest in Brazil . *Novon A J. Bot. Nomencl.*

https://doi.org/10.3417/2013022

**Sayyari, E., & Mirarab, S.** 2018. Testing for polytomies in phylogenetic species trees

using quartet frequencies. *Genes (Basel).* https://doi.org/10.3390/genes9030132

**Schliep, K. P.** 2011. phangorn: Phylogenetic analysis in R. *Bioinformatics*, 27: 592–593.

https://doi.org/10.1093/bioinformatics/btq706

**Stamatakis, A.** 2014. RAxML version 8: A tool for phylogenetic analysis and post-

analysis of large phylogenies. *Bioinformatics*, 30: 1312–1313.

https://doi.org/10.1093/bioinformatics/btu033

**Tange, O.** 2018. GNU Parallel 2018. *Https://Doi.Org/10.5281/Zenodo.1146014*.

https://doi.org/https://doi.org/10.5281/zenodo.1146014

**Team, R. C.** 2019. R: A Language and Environment for Statistical Computing. *Vienna,*

*Austria*.

**Villaverde, T., Pokorny, L., Olsson, S., Rincón-Barrado, M., Johnson, M. G.,**

**Gardner, E. M., Wickett, N. J., Molero, J., Riina, R., & Sanmartín, I.** 2018.

Bridging the micro- and macroevolutionary levels in phylogenomics: Hyb-Seq solves

relationships from populations to species and above. *New Phytol.*

https://doi.org/10.1111/nph.15312

**Weitemier, K., Straub, S. C. K., Cronn, R. C., Fishbein, M., Schmickl, R., McDonnell,**

**A., & Liston, A.** 2014. Hyb-Seq: Combining Target Enrichment and Genome

Skimming for Plant Phylogenomics. *Appl. Plant Sci.*, 2: 1400042.

https://doi.org/10.3732/apps.1400042

**Wen, D., Yu, Y., Zhu, J., & Nakhleh, L.** 2018. Inferring phylogenetic networks using

PhyloNet. *Syst. Biol.*, 67: 735–740. https://doi.org/10.1093/sysbio/syy015

**Yu, Y., & Nakhleh, L.** 2015. A maximum pseudo-likelihood approach for phylogenetic

networks. *BMC Genomics*, 16: 1–10. https://doi.org/10.1186/1471-2164-16-S10-S10

**Zapata, T. R., & Arroyo, M. T. K.** 1978. Plant Reproductive Ecology of a Secondary

Deciduous Tropical Forest in Venezuela. *Biotropica*, 10: 221.

https://doi.org/10.2307/2387907

**Zerega, N. J. C., Clement, W. L., Datwyler, S. L., & Weiblen, G. D.** 2005.

Biogeography and divergence times in the mulberry family (Moraceae). *Mol.*

*Phylogenet. Evol.*, 37: 402–16. https://doi.org/10.1016/j.ympev.2005.07.004

**Zerega, N. J. C., & Gardner, E. M.** 2019. Delimitation of the new tribe Parartocarpeae

(Moraceae) is supported by a 333-gene phylogeny and resolves tribal level Moraceae

taxonomy. *Phytotaxa*, 388: 253–265.

**Zerega, N. J. C., Nur Supardi, M. N., & Motley, T. J.** 2010. Phylogeny and

Recircumscription of Artocarpeae (Moraceae) with a Focus on Artocarpus. *Syst. Bot.*,

35: 766–782. https://doi.org/10.1600/036364410X539853

**Zhang, C., Sayyari, E., & Mirarab, S.** 2017. ASTRAL-III: Increased Scalability and

Impacts of Contracting Low Support Branches BT  - Comparative Genomics: 15th

International Workshop, RECOMB CG 2017, Barcelona, Spain, October 4-6, 2017,

Proceedings. in: J. Meidanis & L. Nakhleh (eds.). Cham: Springer International

Publishing. https://doi.org/10.1007/978-3-319-67979-2_4

**Zhang, Q., Onstein, R. E., Little, S. A., & Sauquet, H.** 2019. Estimating divergence

times and ancestral breeding systems in Ficus and Moraceae. *Ann. Bot.*, 123: 191–204.

https://doi.org/10.1093/aob/mcy159

1726

1729 **Tables**

1730

1731 Table 1. Overview of the taxonomic history of Moreae.

| Engler (1899) | Corner (1962) | Berg (2001) | Clement & Weiblen (2009) | Gardner et al. |
|---|---|---|---|---|
| | | | | *Afromorus*[c] |
| *Ampalis* | *Ampalis* | (*Ampalis*)[a] | (*Ampalis*)[a] | *Ampalis* |
| | | | *Bagassa* | *Bagassa* |
| | (*Bleekrodea*)[a] | *Bleekrodea* | | |
| | | *Broussonetia* | | |
| | *Clarisia* | | | |
| | *Fatoua* | *Fatoua* | | |
| | | *Maclura* | | |
| | (*Maillardia*)[b] | (*Maillardia*)[b] | (*Maillardia*)[b] | *Maillardia* |
| | | (*Malaisia*)[b] | | |
| | | *Milicia* | *Milicia* | *Milicia* |
| *Morus* | *Morus* | *Morus* | *Morus* | *Morus* |
| *Pachytrophe* | *Pachytrophe* | (*Pachytrophe*)[a] | (*Pachytrophe*)[a] | *Pachytrophe* |
| *Paratrophis* | (*Paratrophis*)[a] | (*Paratrophis*)[a] | (*Paratrophis*)[a] | *Paratrophis* |
| *Pseudomorus* | (*Pseudomorus*)[a] | (*Pseudomorus*)[a] | (*Pseudomorus*)[a] | (*Pseudomorus*)[d] |
| | (*Pseudostreblus*)[a] | (*Pseudostreblus*)[a] | (*Pseudostreblus*)[a] | |
| | (*Sloetia*) | (*Sloetia*)[a] | | |
| | (*Sloetiopsis*)[a] | (*Sloetiopsis*)[a] | (*Sloetiopsis*)[a] | |
| | *Sorocea* | | *Sorocea* | *Sorocea* |
| | *Streblus* | *Streblus* | *Streblus* | |
| | (*Taxotrophis*)[a] | (*Taxotrophis*)[a] | (*Taxotrophis*)[a] | *Taxotrophis* |
| *Trophis* | *Trophis* | *Trophis* | *Trophis* | *Trophis* |

1732 [a] Included in *Streblus*.

1733 [b] Included in *Trophis*.

1734 [c] Newly separated from *Morus*.

1735 [d] Included in *Paratrophis*.

1736

1737

1738    Table 2. Divergence times (in Ma) estimated using penalized likelihood under correlated

1739    and relaxed models. Nomenclature follows the revisions proposed in this study.

| Clade | Correlated | Relaxed |
|---|---|---|
| Moraceae | 84.7 | 84.7 |
| Parartocarpeae | 75.3 | 74.7 |
| Ficeae | 24 | 34.5 |
| Olmedieae | 36.4 | 43.4 |
| Dorstenieae | 72.5 | 76.9 |
| Maclureae | 44.9 | 52.1 |
| Artocarpeae | 64 | 64 |
| Moreae | 59.1 | 75.4 |

1740

1741

1742

1743

1744

1745

1746 **Figures**

1747

Figure 1. Representatives of Moreae: *Paratrophis pendulina* (=*Streblus pendulinus*) (a)

infructescences and (b) staminate inflorescences, the latter showing the characteristic sterile

groove; *Sorocea racemosa* (c) infructescences and (d) staminate inflorescence; *Morus nigra*

(e) pistillate inflorescences with elaborate stigmas and (f) staminate inflorescences with

stamens that, having been inflexed in bud, are substantially longer than the perianth itself;

and a newly-placed member of Olmedieae (=Castilleae), *Streblus asper:* (g) infructescences

and (h) discoid-capitate staminate inflorescences with the rudiments of an involucre visible

below the unopened flowers. Photo credits: (a) F. & Kim Starr, under a CC-BY-3.0-US licence

(https://commons.wikimedia.org/wiki/File:Starr-051029-5102-Streblus_pendulinus-fruit-Auwahi-

Maui_(24481336159).jpg); (b) M. Marathon, CC-BY-SA-4.0

(https://commons.wikimedia.org/wiki/File:Streblus_brunonianus_flowers.jpg); (c)-(d) A. Popovkin, CC-BY-

2.0 (https://commons.wikimedia.org/wiki/File:Sorocea_racemosa_Gaudich._-_Flickr_-

_Alex_Popovkin,_Bahia,_Brazil_(3).jpg and

https://commons.wikimedia.org/wiki/File:Sorocea_racemosa_Gaudich._-_Flickr_-

_Alex_Popovkin,_Bahia,_Brazil_(9).jpg); (e) E. Gardner; (f) Schurdl, CC-BY-SA-4.0

(https://commons.wikimedia.org/wiki/File:Morus_nigra_100525_02.jpg); (g) D. Valke, CC-BY-SA-2.0

(https://commons.wikimedia.org/wiki/File:Bekar_(Konkani-_बेकर)_(4533519363).jpg); (h) Vinayaraj, CC

BY-SA 4.0 (https://commons.wikimedia.org/wiki/File:Streblus_asper_at_Panamaram_(5).jpg)

1766

1767

1768    Figure 2. Phylogenetic trees from the "exon" dataset. Maximum-likelihood based on a

1769    supermatrix of all loci, with bootstrap support and previous nomenclature (left), and a

1770    species tree based on gene trees from all loci with bootstrap/LPP support and revised

1771    nomenclature (right). Discordant branches are colored in red, and tribal classifications are

1772    shaded without (left) and with (right) the revised classifications presented here.

1773

1774

Parartocarpeae

Ficeae

Castilleae | Olmedieae

Dorstenieae

Maclureae

Artocarpeae

Moreae

supermatrix

species tree

1775    Figure 3. Phylogenetic trees of the Moreae clade from the "supercontig" dataset.

1776    Maximum-likelihood based on a supermatrix of all loci, with bootstrap support and

1777    previous nomenclature (left), and a species tree based on gene trees from all loci with

1778    bootstrap/LPP support and revised nomenclature (right). Discordant branches are colored in

1779    red.

1780

1781

supermatrix

species tree

1782     Figure 4. The best five maximum-pseudo-likelihood phylogenetic networks for the

1783     *Paratrophis* clade.

1784

1785

1786    Figure 5. Time-calibrated phylogenetic tree, with revised nomenclature.

1787

1788

1789    Figure 6. Ancestral reconstruction of stamen position, with revised nomenclature. The

1790    reconstruction was identical under a trait-dependent (BiSSE) or a trait-independent model.

1791    Blue = inflexed in bud; yellow = straight in bud.

1792

1793

*Pseudostreblus indica* NewmanLA0518
*Hulletia dumosa* NZ242
*Hulletia griffithiana* Kerr16886
*Parartocarpus venenosus* NZ874
*Parartocarpus bracteatus* NZ730
*Ficus sagittifolia* ChaseMW19852
*Ficus racemosa* SRR1405699 SRR1405700
*Ficus macrophylla* EG30
*Streblus asper* EG696
*Antiaropsis decipiens* NZ281
*Sparattosyce dioica* WeiblenWW1223
*Poulsenia armata* Pennington14600
*Helicostylis tomentosa* Gomes511
*Castilla elastica* ChaseMW19850
*Olmedia aspera* Fuentes5323 S41
*Pseudolmedia spuria* Lobo263
*Naucleopsis macrophylla* BergP18534
*Antiaris toxicaria* 5595
*Fatoua pilosa* TaylorP218
*Malaisia scandens* EG122
*Broussonetia papyrifera* SRR1477753
*Broussonetia cf papyrifera* Heng14289
*Broussonetia kazinoki* ChaseMW17827
*Bleekrodea madagascariensis* DavisRakotonasolo3117
*Sloetiopsis usambarensis* Faden77 417
*Brosimum alicastrum* EG23
*Helianthostylis sprucei* Ribeiro1204
*Trymatococcus amazonicus* Prance13461
*Bosqueiopsis gilletii* Frontiers243
*Trilepisium madagascariense* Richard403
*Utsetela gabonensis* Breteler14086
*Utsetela neglecta* Bissiengen923
*Dorstenia hildebrandtii* NZ311
*Dorstenia barteri* ChaseMW18153
*Maclura tinctoria tinctoria* Campos2679
*Maclura africana* GreenwayKanuri15269
*Maclura andamanica* Maxwell89–436
*Maclura spinosa* Perumal RHT22554
*Maclura pomifera* EG139
*Maclura brasiliensis* NeeVargas45025
*Maclura thorellii* Pierre4709
*Maclura cochinchinensis* SAGBE887
*Maclura aff pubsecens* Averyanov1198
*Maclura tricuspidata* MOR 68–7917
*Maclura fruticosa* NMCuong1193
*Maclura cf cochinchinensis* NZ757
*Maclura amboinensis* Takeuchi16578
*Taxotrophis zeylanica* Meijer701
*Taxotrophis macrophylla* DDS10673
*Taxotrophis ilicifolius* EG720
*Taxotrophis spinosa* EG702
*Taxotrophis taxoides* EG701
*Maillardia borbonica* Cadet4518
*Maillardia montana* Andrianantoanina 1023
*Afromorus mesozygia* Buckner309
*Milicia excelsa* McPherson16087
*Milicia regia* Cooper332
*Pachytrophe dimepate* CasmirZ405
*Ampalis mauritiana* Hoffman
*Paratrophis microphylla* Beecher sn
*Paratrophis anthropophagorum* Degeer4698
*Paratrophis insignis* Vasquez28068
*Paratrophis pendulina* Gray3277
*Paratrophis glabra* EG78
*Paratrophis philippinensis* EG430
*Trophis involucrata* Aguilar4679
*Trophis racemosa* Stevens37196
*Trophis mexicana* Stevens27939
*Trophis cuspidata* Gomez–Dominuez774
*Morus notabilis* SRR8138828
*Morus rubra* EG141
*Morus microphylla* Fishbein1058
*Morus celtidifolia* Sandoval637
*Morus macroura* EG28
*Morus serrata* Koelz4788
*Morus australis* MOR 241–716
*Morus kagayamai* AA20187–A
*Morus cathayana* Nie92144
*Morus nigra* EG29
*Morus mongolica* MOR 55–951
*Morus alba* MOR 920–261
*Bagassa guianensis* GW1677
*Sorocea subumbellata* Dodson9601
*Sorocea duckei* Stein 3931
*Sorocea sprucei saxicola* Nee35729
*Sorocea trophoides* Fuentes304
*Sorocea affinis* Espinosa706
*Sorocea pubivena pubivena* Galdames4477
*Sorocea pubivena oligotricha* Kennedy2124
*Sorocea jaramilloi* Madison4772
*Sorocea muriculata muriculata* Schunke16489
*Sorocea pubivena hirtella* Vasquez19236
*Sorocea briquetii* Valenzula954
*Sorocea muriculata uaupensis* Plowman sn
*Sorocea steinbachii* GW1501
*Sorocea boplandii* Tressens6347
*Sorocea hilarii* Carauta795
*Sorocea guilleminiana* Moya112 S38
*Clarisia ilicifolia* Carranta6386
*Batocarpus amazonicus* Salidas1233
*Batocarpus costaricensis* GW1463
*Batocarpus oriniceros* Vasquez27440
*Clarisia racemosa* Assuncao690
*Clarisia biflora* GW1460
*Artocarpus scandens* EG411
*Artocarpus papuanus* NZ61
*Artocarpus limpato* NZ609
*Artocarpus altissimus* EG441
*Artocarpus sepicanus* GW1701
*Artocarpus longifolius ssp adpressus* Nangkat15511
*Artocarpus fulvicortex* EG711
*Artocarpus nitidus ssp lingnanensis* NZ911
*Artocarpus thailandicus* NZ402
*Artocarpus lacucha* NZ420
*Artocarpus gomezianus ssp zeylanicus* NZ956
*Artocarpus nitidus ssp humilis* EG258
*Artocarpus nitidus ssp griffithii* NZ216
*Artocarpus dadah* NZ694
*Artocarpus tonkinensis* EG174
*Artocarpus styracifolius* EG176
*Artocarpus hypargyraeus* EG170
*Artocarpus nanchuanensis* EGL238
*Artocarpus petelotii* DDS14435
*Artocarpus pithecogallus* LiJianwu3200
*Artocarpus gongshanensis* HAST137747
*Artocarpus gomezianus ssp gomezianus* NZ533
*Artocarpus cf lacucha* NZ517
*Artocarpus glaucus* NZ852
*Artocarpus nitidus ssp borneensis* NZ686
*Artocarpus tomentosulus* NZ617
*Artocarpus primackii* NZ687
*Artocarpus xanthocarpus* Yang15648
*Artocarpus vriesianus var vrieseanus* GW1229
*Artocarpus vrieseanus refractus* Hoogland4822
*Artocarpus albobrunneus* AA2243
*Artocarpus fretessii* NZ929
*Artocarpus ovatus* NZ202
*Artocarpus nitidus ssp nitidus* PPI10374
*Artocarpus rubrovenius* JSB84
*Artocarpus subrotundifolius* Wenzel1576
*Artocarpus heterophyllus* EG98
*Artocarpus annulatus* NZ985
*Artocarpus integer* NZ918
*Artocarpus sarawakensis* EG527
*Artocarpus brevipedunculatus* NZ814
*Artocarpus anisophyllus* NZ606
*Artocarpus lanceifolius ssp clementis* NZ739
*Artocarpus hirsutus* NZ953
*Artocarpus nobilis* AHJ3283
*Artocarpus chama* NZ354
*Artocarpus melinoxylus* JLS2924
*Artocarpus odoratissimus* NZ618
*Artocarpus rigidus ssp asperulus* NZ507
*Artocarpus hispidus* NZ258
*Artocarpus rigidus* NZ728
*Artocarpus treculianus nigrescens* Ramos2107
*Artocarpus pinnatisectus* Escritor
*Artocarpus multifidus* PPI3911
*Artocarpus treculianus* Elmer12468
*Artocarpus treculianus ovatifolius* NZ203
*Artocarpus blancoi* Ramos42018
*Artocarpus aff excelsus* AveryanovVH1819
*Artocarpus camansi* EG149
*Artocarpus mariannensis* DD4
*Artocarpus altilis* K7 UluFiti
*Artocarpus horridus* EG437
*Artocarpus cf horridus* RM160
*Artocarpus excelsus* NZ780
*Artocarpus obtusus* NZ729
*Artocarpus lowii* MWL2
*Artocarpus teijsmannii* NZ946
*Artocarpus sumatranus* AA2766
*Artocarpus kemando* NZ612
*Artocarpus maingayi* NZ257
*Artocarpus tamaran* EG92
*Artocarpus sericicarpus* NZ771
*Artocarpus elasticus* EG87
*Artocarpus scortechinii* NZ209
*Artocarpus corneri* Fuchs21347
*Artocarpus jarrettiae* SAN120933

1794      Figure 7. Revised classification of Moreae on a maximum-likelihood phylogenetic tree of

1795      Moreae based on "supercontig" sequences.

1796

1797

*Taxotrophis*

*Maillardia*

*Afromorus*

*Milicia*

*Pachytrophe*

*Ampalis*

*Paratrophis*

*Trophis*

*Morus*

*Bagassa*

*Sorocea*