Phylogenomic analyses of non-Dikarya fungi supports horizontal gene transfer driving

diversification of secondary metabolism in the amphibian gastrointestinal symbiont,

Basidiobolus

Javier F. Tabima[1]

Ian A. Trautman[1]

Ying Chang[1]

Yan Wang[2,3,4]

Stephen Mondo[5]

Alan Kuo[5]

Asaf Salamov[5]

Igor V. Grigoriev[5,6]

Jason E. Stajich[2,3]

Joseph W. Spatafora[1]

Affiliations:
1. Department of Botany and Plant Pathology, College of Agricultural Sciences, Oregon State
University, Corvallis, U.S.A.
2. Department of Microbiology and Plant Pathology, University of California—Riverside,
Riverside, California, USA
3. Institute for Integrative Genome Biology, University of California—Riverside, Riverside,
California, USA
4. Current address: Department of Biological Sciences, University of Toronto Scarborough,
Toronto, Ontario, Canada and the Department of Ecology and Evolutionary Biology, University
of Toronto, Toronto, Ontario, Canada

5. US Department of Energy Joint Genome Institute, Lawrence Berkeley National Laboratory,

Berkeley, California, USA

6. Department of Plant and Microbial Biology, University of California-Berkeley, Berkeley,

California, USA

Abstract

Research into secondary metabolism (SM) production by fungi has resulted in the discovery of

diverse, biologically active compounds with significant medicinal applications. However, the

fungi rich in SM production are taxonomically restricted to Dikarya, two phyla of Kingdom

Fungi, Ascomycota and Basidiomycota. Here, we explore the potential for SM production in

Mucoromycota and Zoopagomycota, two phyla of nonflagellated fungi that are not members of

Dikarya, by predicting and identifying core genes and gene clusters involved in SM. The

majority of non-Dikarya have few genes and gene clusters involved in SM production except for

the amphibian gut symbionts in the genus *Basidiobolus*. *Basidiobolus* genomes exhibit an

enrichment of SM genes involved in siderophore, surfactin-like, and terpene cyclase production,

all these with evidence of constitutive gene expression. Gene expression and chemical assays

confirm that *Basidiobolus* has significant siderophore activity. The expansion of SMs in

Basidiobolus are partially due to horizontal gene transfer from bacteria, likely as a consequence

of its ecology as an amphibian gut endosymbiont.

Introduction

Fungi produce a wealth of biologically active small molecules – secondary or specialized

metabolites – that function in interactions with other organisms, environmental sensing, growth

and development, and numerous other processes (Rokas et al. 2020). Several of these

compounds have led to the successful development of pharmaceuticals (e.g., antibiotics,

immunosuppressants, statins, etc.) that have had dramatic and positive impacts on human health.

Understanding the evolution of fungal secondary metabolites and linking them with their

ecological and physiological functions in nature can inform searches for compounds with

applications in human society.

Secondary metabolism (SM) is imprecisely defined but can be characterized generally as

the production of bioactive compounds that are not part of primary metabolism and that are not

required for growth and survival in the laboratory (Keller et al 2005, Brakhage 2013, Rokas et al.

2020). In fungi, the genes responsible for the synthesis of secondary metabolites are frequently

co-located in biosynthetic gene clusters (Smith et al. 1990, Brakhage 2013), which contain the

genes that control regulation of expression, biosynthesis, tailoring, and transport of these

compounds out of the cell (Smith et al. 1990, Keller et al 2005, Osbourn 2010). In the kingdom

Fungi, the diversity of products synthesized via SM is substantial and primarily includes

alkaloids, peptides, polyketides, and terpenes (Collemare et al. 2008, Helaly et al. 2018). Each

of these groups of compounds are synthesized by core genes that are characteristic of the

pathways and include, but are not limited to, dimethylallyl tryptophan synthases (DMAT), non-

ribosomal peptide synthetases (NRPS), polyketide synthetases (PKS), and terpene cyclases (TC).

These bioactive compounds fulfill various roles that are hypothesized to increase the fitness of

the fungus by promoting better recognition and adaptation to environmental cues.

Biosynthesis of secondary metabolites is heterogeneous across the fungal tree of life, but

the vast majority of discovered and predicted secondary metabolites are reported within the

fungal phyla Ascomycota and Basidiomycota of the subkingdom Dikarya. Filamentous

ascomycetes are the major producers of secondary metabolites (e.g., penicillin, cyclosporin, etc.),

with the majority of genes and gene clusters involved in fungal SM discovered in the subphylum

Pezizomycotina (Collemare et al. 2008, Helaly et al. 2018). Although less than Ascomycota,

Basidiomycota is also a prominent producer of SM, including some of the better-known

hallucinogens (e.g., psilocybin of *Psilocybe*; Reynolds et al. 2018) and compounds toxic to

humans (e.g., amanitin of *Amanita*; Luo et al. 2010).

For reasons that are unclear, the remainder of kingdom Fungi is characterized by a

paucity of secondary metabolites (Voight et al. 2016). This includes the zoosporic fungi and

relatives classified in Blastocladiomycota, Chytridiomycota and Rozellomycota, and the

nonflagellated, zygomycete fungi of Mucoromycota and Zoopagomycota. This pattern of SM

diversity supports the hypothesis that diversification of secondary metabolism is a characteristic

of Ascomycota and Basidiomycota (subkingdom Dikarya), which share a more recent common

ancestor relative to the other phyla. Recent genome sampling efforts have focused on increased

sequencing of non-Dikarya species (Nagy et al. 2014, Kohler et al. 2015, Spatafora et al. 2016,

Quandt et al. 2017, Ahrendt et al. 2018). These efforts have provided a better understanding of

the relationships of the phyla of kingdom Fungi (e.g., Spatafora et al. 2016), and processes and

patterns that shaped the evolution of morphologies (e.g., Nagy et al. 2014) and ecologies (e.g.,

Chang et al. 2019, Quandt et al. 2017) within the kingdom. The availability of a diversity of

these genomes provides an opportunity to characterize and focus on the secondary metabolism

composition of non-Dikarya taxa, which have remained relatively unexplored.

While the majority of non-Dikarya taxa have low SM diversity, genomic sequencing of

the genus *Basidiobolus* (Phylum Zoopagomycota) revealed that it possesses an unusually large

composition of SM gene clusters. Species of *Basidiobolus* have complex life cycles, which

produce multiple spore types that occur in multiple environmental niches. These species are

symbionts found in the digestive tracts of reptiles and amphibians, but their function and impact

on the host remains unknown. The fungus is dispersed with the feces where it sporulates

producing both forcibly discharged asexual spores (blastoconidia) and passively dispersed

asexual spores (capilloconidia) that adhere to exoskeletons of small insects. These insects are

consumed by insectivorous amphibians, completing the life cycle. *Basidiobolus* also reproduces

sexually through the production of zygospores (meiospores) either by selfing (homothallic) or

outcrossing (heterothallic) according to species. The fungus is also isolated from leaf litter and

can be maintained in pure culture, findings that are consistent with a saprobic (decomposition of

organic matter) phase to the life cycle. *Basidiobolus* must have adapted for survival in numerous

environmental niches including the amphibian digestive system, amphibian feces, insect

phoresis, and on decaying plant matter or leaf litter.

In this study we demonstrate that the genomes of *Basidiobolus* contain a larger number of

genes related to SM than predicted by phylogeny and that in several cases the evolution of many

of these SM genes is inconsistent with vertical evolution. Our objectives were to: i) characterize

the diversity of SM in *Basidiobolus*, ii) identify the phylogenetic sources of this diversity, and

iii) determine which classes of SM gene clusters are functional and may predict the secondary

metabolites produced by species of *Basidiobolus*. Finally, we propose a model in which the

amphibian gastrointestinal system is an environment that promotes noncanonical evolution of its

fungal inhabitants.

Results

Secondary metabolite gene cluster prediction. – A total of 38 secondary metabolism (SM) gene

clusters and 44 SM core genes were predicted in the *B. meristosporus* CBS 931.73 genome, 40

SM gene clusters and 44 SM core genes for *B. meristosporus* B9252, and 23 SM gene clusters

and 23 SM core genes for *B. heterosporus* B8920 (Table 1; Supplementary Table 2). Seventy-

eight percent of the SM gene models predicted were found to be shared across the *Basidiobolus*

isolates. Ten SM core genes were found to be unique to *B. meristosporus* CBS 931.73, 10 SM

genes unique to *B. meristosporus* B9252, and 5 SM genes unique to *B. heterosporus* B8920

(Additional file 1).

A total of 721 SM gene clusters were predicted for the 66 additional Mucoromycota and

Zoopagomycota genomes including 74 non-ribosomal peptide synthetases (NRPS), 167 NRPS-

like, 97 polyketide synthases (PKS), 91 PKS-like, 292 terpene cyclase (TC) gene models across

both phyla (Figure 1, Supplementary Table 2). For Zoopagomycota (including *Basidiobolus*),

284 SM gene models were predicted, including one NRPS-PKS hybrid, 71 NRPS, 56 NRPS-

Like, 54 PKS, 46 PKS-Like, and 56 TC gene models. In Mucoromycota, 563 total SM gene

models were predicted, including 45 NRPS, 142 NRPS-Like, 49 PKS, 51 PKS-Like, and 256 TC

gene models. The three isolates with the most numerous predicted SM gene clusters, not

including *Basidiobolus* genomes, were *Dimargaris cristalligena* RSA 468 with 33 predicted SM

proteins (21 NRPS, 7 NRPS-Like, 2 PKS-Like, 3 TC), *Linderina pennispora* ATCC 12442 V 1.0

with 23 SM predicted (1 NRPS, 15 PKS, 3 PKS-like, 4 TC), and *Martensiomyces pterosporus*

CBS 209.56 v1.0 with 23 SM proteins predicted (1 NRPS, 5 PKS, 14 PKS-Like, 3 TC). No

DMAT gene models were predicted for any member of Mucoromycota or Zoopagomycota.

Expression of core SM genes in Basidiobolus. – A total of 83.45% of the RNA sequenced reads

were mapped uniquely to the reference genome of *B. meristosporus* CBS 931.73, while 12.08%

of the reads were mapped in more than one location. Only 4.47% of the RNA sequenced reads

did not map to the reference genome. The majority of predicted SM for *B. meristosporus* CBS

931.73 were expressed at the same or higher levels than constitutive housekeeping genes, such as

Beta-tubulin, Elongation Factor 1, Actin, and Ubiquitin (Figure 2). The highest expressed SM

core genes per SM group were: NRPS – gene model 387529 (Cluster 5) with 74.03 transcripts

per million (TPM) mapped; NRPS-like – gene model 221915 (Cluster 20) with 45.58 TPM

mapped; PKS – gene model 290138 (Cluster 37) with 687.55 TPM mapped; PKS-like – gene

model 207695 (Cluster 38) with 26.15 TPM mapped, and Terpene cyclase – gene model 301341

(Cluster 13) with 78.26 TPM mapped.

Phylogenetic analysis of NRPS/NRPS-Like A-domains. – The phylogenetic reconstruction of A-

domains was performed with 951 A-domains from a combined dataset including the A-domain

dataset from Bushley and Turgeon (2010) and the predictions of NRPS/NRPS-like A-domains

from Mucoromycota and Zoopagomycota genome sequences (Additional file 2). A total of 395

NRPS A-domains were predicted for *Basidiobolus*, Mucoromycota and Zoopagomycota genome

sequences (Additional File 3). The phylogenetic analyses recovered the nine major families

reported by Bushley and Turgeon (2010) with the addition of the two new clades, surfactin-like

and ChNSP 12-11-like, reported here. The total number of Mucoromycota/Zoopagomycota A-

domains and their distribution across these clades are as follows: 74 to the AAR clade; 115 to the

major bacterial clade (MBC), including 104 A-domains with the surfactin-like clade with and an

additional 11 A-domains scattered elsewhere in the MBC; the CYCLO clade with eight A-

domains; 65 A-domains to the ChNSP 12-11-like clade; 25 A-domains to the ChNSP 12-11

clade; 11 A-domains to the PKS/NRPS clade; 16 A-domains to the SID clade; 76 A-domains to

the SIDE clade; and 5 A-domains to the EAS clade (Figure 3). No A-domains were clustered

into the ACV clade or the ChNSP 10 clade, and the remaining A-domains grouped within the

outgroup clade (Figures 3 and 4, Supplementary Figure 1). Similar phylogenetic origins for

multiple A-domains were found in a single NRPS core gene (Figure 4).

171   Of fungi classified in Mucoromycota or Zoopagomycota, *Basidiobolus* genomes

contained the most A-domains clustered in the EAS, SIDE, and CYCLO clades. The A-domains

clustered within EAS represent two domains from *B. meristosporus* CBS 931.73, and one A-

domain of *Rhizophagus irregularis* DAOM 197198 v2.0. The A-domains clustered within SIDE

represent 76 domains, 44 from *B. meristosporus* CBS 931.73, 19 from *B. meristosporus* B9252,

and 13 from *B. heterosporus.* The A-domains clustered within the CYCLO clade represent six

domains, four from *B. meristosporus* CBS 931.73 and two from *B. meristosporus* B9252. The

NRPS A-domains grouped in the CYCLO cluster correspond to cluster 5 NRPS from *B.*

meristosporus CBS 931.73 (Protein ID 387529) and cluster 9 of *B. meristosporus* B9252

(Protein ID N161_4477). Gene model 387529 was predicted as a tetra-modular protein, with two

N-methyltransferase domains and a thioesterase domain. In addition, AntiSMASH analyses

predict a six gene cluster (cluster 5) that includes gene model 387529, a zinc finger transcription

factor (312194), carrier protein (277549), transporter (277552), peptidase (277554), and a

SNARE associated protein (334759) (Supplementary Figure 2). All of these gene models have

evidence of gene expression (Figure 2).

The *Basidiobolus* NRPS A-domains clustered in the surfactin-like clade included 32 A-

domains from eight gene models from *B. meristosporus* CBS 931.73. These included gene

model 375475 (Cluster 19) with eight A-domains, gene models 307892 (Cluster 33) and 343011

(Cluster 35) with seven A-domains each; gene model 368581 (Cluster 8) with three A-domains;

gene models 337511 (Cluster 16), 372991 (Cluster 17), and 298977 (Cluster 7) each with two A-

domains; and gene model 322666 (Cluster 2) with one A-domain. *B. meristosporus* B9252

contained two surfactin-like A-domains from one gene model (N161_8304; Cluster 18). *B.*

heterosporus possessed 13 A-domains from seven gene models, including N168_07733 (Cluster

6), N168_06479 (Cluster 13), N168_02885 (Cluster 1) and N168_00034 (Cluster 14) with two

A-domains each, and gene models N168_05934 (Cluster 8), N168_04239 (Cluster 12) and

N168_07140 (Cluster 9) with one A-domain each.

Evolutionary relationships of predicted PKSs. – A total of 421 KS domains were included in the

phylogenetic reconstruction (Additional File 4). Two KS domains were predicted from

Chytridiomycota genomes, 21 from Neocallimastigomycota, 46 from Basidiomycota, 78 from

Ascomycota, 54 from Mucoromycota, and 76 from Zoopagomycota for a total of 310 predicted

fungal KS domains from genome sequences (Additional File 4). The additional KS domains

were obtained from Kroken et al. (2003). For Mucoromycota and Zygomycota, the genomes with

the highest number of KS domains were *Linderina pennispora* ATCC 12442 v1.0 (20 KS

domains), *Coemansia spiralis* RSA 1278 v1.0 (11 KS domains), *Martensiomyces pterosporus*

CBS 209.56 v1.0 (10 KS domains), *Coemansia reversa* NRRL 1564 v1.0 (8 KS domains),

Coemansia mojavensis RSA 71 v1.0 (7 KS domains), and the *Basidiobolus* genomes: *B.*

meristosporus CBS 931.73 (5 KS domains), *B. meristosporus* B9252 (4 KS domains), and *B.*

heterosporus (3 KS domains).

Five major clades of PKS proteins were reported by Kroken et al. (2013) and recovered

by this study, including the fungal reducing (R) PKS, animal fatty acid synthases (FAS), fungal

non-reducing (NR) PKS, bacterial PKS, and fungal FAS (Supplementary Figures 3 and 4). The

Fungal reducing PKS1 clade included domains from *B. meristosporus* CBS 931.73 (2 KS

domains) and *B. meristosporus* B9252 (2 KS domains), as well as a new sub-clade called

"reducing PKS clade V". This clade comprised KS domains from Basidiomycota,

Neocallimastigomycota and one Chytridiomycota representative. No KS domains of

Mucoromycota or Zoopagomycota genomes were clustered within the animal FAS clade. The

Fungal NR PKS clade contained a new clade, "non-reducing PKS proteins clade IV", comprising

KS domains from Basidiomycota, Neocallimastigomycota, and one KS domain of *B.*

meristosporus B9252. The Bacterial PKS clustered KS domains from *B. meristosporus* CBS

931.73 (1 KS domain) and *B. meristosporus* B9252 (1 KS domain) as sole representative of

Mucoromycota and Zoopagomycota. Finally, the fungal FAS clade comprised all the remaining

KS domains of Mucoromycota and Zoopagomycota genomes (Supplementary Figures 3 and 4).

To better understand the predicted PKSs of Mucoromycota and Zoopagomycota that

clustered in the fungal FAS clade, the patterns of domains that comprise fungal PKS protein

sequences were analyzed. Predicted PKS of Ascomycota contained AT, KS and PP domains in

all sequences (Supplementary Figure 5). Predicted PKS from Basidiomycota contained AT and

KS domains in all sequences, while KR and PP domains were present on 36% of the sequences.

All PKS that contain a KS domain in the FAS clade are missing the PP domain. For

Chytridiomycota, the predicted PKS with the KS domain clustered in the Fungal (R) clade

contained AT, KS and DH domains, but no PP domain. Chytridiomycota PKS with KS domains

associated to FAS clade only possessed AT and KS domains. For Neocallimastigomycota, 100%

of PKS clustered in fungal R and NR clades contained AT and KS domains, however, only 5

PKS contained the PP domain. All Neocallimastigomycota PKS that clustered within the FAS

clade contained only AT and KS domains. For Mucoromycota and Zoopagomycota, 100% of

PKS contained either AT and/or KS domain, but no other domains (Supplementary Figure 6).

Evolutionary relationships of predicted terpene cyclase gene models. – A total of 1,108 terpene

cyclase or terpene cyclase-like gene models were identified and used for phylogenetic analyses

(Additional file 5). These include 256 identified in the Mucoromycota species genomes, 56 in the

Zoopagomycota genomes, and 401 ortholog proteins from 58 fungal genomes of Basidiomycota

and Ascomycota. An additional 395 TC candidates were identified in bacterial genomes from

RefSeq via BLASTP. The phylogenetic reconstruction of TC resulted in two main clades: an

outgroup clade that comprised predicted TC annotated as tRNA threonylcarbamoyladenosine

dehydratase and phytoene synthases, and a main ingroup TC core clade (Figure 5,

Supplementary Figure 7). The tRNA threonylcarbamoyladenosine dehydratase subclade

contained mostly bacterial sequences plus one TC gene model of *B. meristosporus* B9252 and *B.*

meristosporus CBS 931.73 each. The phytoene synthases clade comprised two subclades, one

clade that grouped most bacterial gene models, and a second clade clustering bacterial and

Mucoromycota predicted TC gene models, but no *Basidiobolus* TC core gene models were found

in this clade.

The TC core clade (Figure 5 and Supplementary Figure 7) comprised a bacterial

exclusive clade (Bacteria I), followed by four highly supported sub-clades containing TC core

genes from Mucoromycota, Zoopagomycota, and Dikarya, and are referred to here as Fungi TC

clades I – IV. The majority of TC genes from Mucoromycota and Zoopagomycota clustered

within clade Fungi TC II, which included no other fungal or bacterial TC. These genes were

annotated as Squalene synthetases by the *Mycocosm* genome portal.

The Fungi TC clades I, III and IV included mostly Mucoromycota TCs. The Fungi TC I

clade contained TC genes from Mucoromycota isolates, as well as TC genes from the

ascomycete isolates *Fusarium verticillioides* 7600 and *Hypoxylon sp*. EC38 v1.0, and the

basidiomycete *Suillus luteus* UH-Slu-Lm8-n1 v2.0. These gene models were annotated as

associated with the Ubiquitin C-terminal hydrolase UCHL1 according to the *MycoCosm* portal.

Fungi TC III clade also comprised mostly Mucoromycota isolates, with additional TCs from

Zoopagomycota isolates *Piptocephalis cylindrospora* RSA 2659 single−cell v3.0 and

Syncephalis pseudoplumigaleata Benny S71−1 single−cell v1.0. Annotations of genes in Fungi

TC III clade in *Mycocosm* indicated that these proteins are part of the isoprenoid/propenyl

synthetases, responsible for synthesis of isoprenoids. Isoprenoids play a role on synthesis of

various compounds such as cholesterol, ergosterol, dolichol, ubiquinone or coenzyme Q (Finn et

al. 2009). Finally, Fungi TC IV clade followed a similar Mucoromycota-enriched pattern with

the exceptions of *Dimargaris cristalligena* RSA 468 single−cell v1.0, *Linderina pennispora*

ATCC 12442 v1.0, *M. pterosporus* CBS 209.56 v1.0, and *Syncephalis fuscata* S228 v1.0. Fungi

TC IV clade also included TC genes from Ascomycete sequenced genomes. The majority of

these genes were annotated as containing a terpene synthase family, metal binding domain, and a

polyprenyl synthetase domain in the *MycoCosm* genome portal.

Within the fungal clades, the NADH dehydrogrenase (ubiquinone) complex clade

represented a highly supported clade clustering bacterial, Mucoromycota and Zoopagomycota

TCs. Annotations associated with gene models found in this clade indicated that this clade

comprised TC associated to NADH dehydrogrenase (ubiquinone) complex.

The terminal clades of TC clusters include the Bacteria II + *Basidiobolus* clade that

included bacterial TC and six TC SM core genes predicted from *Basidiobolus* (two from each

genome), and the clade comprising TC core genes solely from Dikarya and one gene from

Mortierella verticillata NRRL 6337 and *Rhizopus microsporus ATCC11559 v1.0* (Figures 5 and

Supplementary Figure 7).

Signatures for HGT in Basidiobolus *genomes*. – The identity search for genes with evidence for

HGT identified 934 genes in *B. meristosporus* CBS 931.73, 620 genes of *B. meristosporus*

B9252, and 382 genes of and *B. heterosporus* B8920 with zero BLASTP hits to fungal proteins.

These genes were used for the coverage assay to identify gene model coverage deviation from

the harboring scaffold median coverage (Sup. Fig. 7). This assay resulted in 810 genes of *B.*

meristosporus CBS 931.73, 503 genes of *B. meristosporus* B9252, and 301 genes of and *B.*

heterosporus B8920 with z-scores under 2 standard deviations from the harboring scaffold

median coverage. These genes were considered candidate genes with evidence for HGT, and

represented 5%, 4% and 3% of the gene content of *B. meristosporus* CBS 931.73, *B.*

meristosporus B9252, and *B. heterosporus* B8920, respectively. The HGT candidates showed

significant differences in GC content when contrasted to genes considered of fungal origin

(Mean fungal GC content (expected) = 0.492, Mean HGT GC content (Observed) = 0.486, t =

3.4119, df = 891.77, p-value = 0.0006741), but no differences in codon usage or 5-mer

composition were detected between HGT candidate and fungal genes (Supplementary Figures 10

and 11). Differences were also found for intron number and normalized intron length between

genes HGT candidate genes and fungal genes, where the distribution of intron number shows a

median of 0 introns for HGT candidates and 2 for fungal genes (Kruskal-Wallis test; χ^2=272.88,

df = 1, p-value < 2.2e-16; Sup. Fig. 9). The majority of HGT candidate genes (61% of the genes)

have no introns, compared to 36% of genes from fungal origin with no introns (Sup. Fig. 9). The

candidate HGT genes have a significantly smaller normalized intron length than the genes with a

fungal origin (Kruskal-wallis test; χ^2=272.88, df = 1, p-value < 2.2e-16).

A large percentage of SM core genes appeared to be the product of HGT from bacterial

species into *Basidiobolus*. The SM core genes identified as candidate HGT genes is 61%, 44%

and 52% for *B. meristosporus* CBS 931.73, *B. meristosporus* B9252, and *B. heterosporus*

B8920, respectively (Table 2). NRPS/NRPS-like SM core genes represent the largest percentage

of HGT evidence, while TC have the lowest percentage of HGT evidence (Table 2, Table 3).

The identification of taxonomic sources for HGT into *Basidiobolus* indicated that most HGT

comes from bacteria in the phylum Proteobacteria with 234, 160 and 95 gene models in *B.*

meristosporus CBS 931.73, *B. meristosporus* B9252, and *B. heterosporus* B8920, respectively.

Firmicutes (112, 63, and 39 gene models, respectively) and Actinobacteria (127, 67, and 40 gene

models, respectively) (Figure 6, Supplementary Table 4) were the second and third most

abundant source of HGT from bacteria into *Basidiobolus*.

Most HGT candidates genes resulted in no gene ontology (GO) annotation (659 gene

models) or no InterPro domain annotation (140 gene models, Supplementary Table 5). The ten

top GO terms were oxidation-reduction process (59 gene models), protein binding (34 gene

models), phosphorelay sensor kinase activity (32 gene models), N-acetyltransferase activity (28

gene models), catalytic activity (26 gene models), extracellular space (25 gene models),

hydrolase activity (24 gene models), oxidoreductase activity (23 gene models), hydrolase activity

(24 gene models), hydrolyzing O-glycosyl compounds(23 gene models) , and NRPS (18 gene

models) (Figure 6, Supplementary Table 5).

Siderophore activity in Basidiobolus meristosporus. – The siderophore activity assay resulted in

visible activity for the three replicates for *B. meristosporus* and *Cladosporium sp.*, and a small

halo for one of the replicates of *C. thromboides*. No visible siderophore activity was detected for

the other two replicates of *C. thromboides* or for the empty AY-CAS plates (Figure 7,

Supplementary Figure 12). The analysis of variance for the area of siderophore activity measured

showed significant differences across isolates/empty plates (ANOVA, F value = 19.62, p =

0.0004). Post-hoc tests showed no differences between *B. meristosporus* and *Cladosporium sp.*

siderophore activity (Tukey HSD, p = 0.9801489) nor between *C. thromboides* and the empty

AY-CAS plates (Tukey HSD, p = 0.9396603), but significant differences were found for all

other comparisons (Tukey HSD, $p < 0.001$ for all remaining comparisons).

Discussion

Secondary, or specialized, metabolite (SM) production is an important element of fungal

metabolism. It has resulted in numerous natural products with human health implications, such

as mycotoxins in our food supply, and medical applications, including antibiotics,

immunosuppressants, and antitumor agents. The majority of the known genetic and chemical

diversity of fungal SM has been described from the phyla Ascomycota and Basidiomycota with

few SM gene clusters, and limited SM production, reported for fungi in Mucoromycota and

Zoopagomycota. This observation has led to the dogma that 'zygomycete' species are

depauperate of these chemical pathways (Voight et al. 2016). Here we report gene clusters

involved in SM production in the largest survey to date of Mucoromycota and Zoopagomycota

species using genomics approaches and estimate the SM potential of these fungi.

Genome sequencing of a total of 69 isolates across diverse lineages of Mucoromycota

and Zoopagomycota enabled detailed identification of SM clusters. These results support the

hypothesis that zygomycete fungi have a low abundance of secondary metabolism (Figure 1,

Supplementary Table 2) and agree with previous reports (Voight et al. 2016). Outliers to this

pattern exist, however, and are particularly true of the genus *Basidiobolus* (Zoopagomycota),

which possesses a large number of SM gene clusters predicted for the NRPS, PKS and TC

families. This discovery of abundant candidate genes for production of secondary metabolite in

Basidiobolus is novel and its presence is most consistent with a signal of horizontal gene transfer

from bacteria to fungi, a phenomenon we propose is facilitated by living in the amphibian gut

environment.

Distribution and Evolution of NRPS Genes Across Mucoromycota and Zoopagomycota. – A

deeper examination of *Basidiobolus* SM gene clusters indicates that this genus surpasses the

number of expected NRPS genes for zygomycete species (Figure 1). Most of these SM genes

also show evidence for transcription, indicating that the majority of these genes are expressed

constitutively under laboratory conditions (Figure 2). Several of these core genes, such as the

NRPS gene model 387529, appear to be expressed at a higher rate than several housekeeping

genes.

A more detailed census of core genes reveals unique patterns of evolution of NRPS genes

across Mucoromycota and Zoopagomycota when compared to Dikarya fungi (Figure 3, Table 1).

The only NRPS A-domains found throughout the two phyla are members of the AAR clade (with

the exception of *Rhizopus delemar* 99-80). These A-domains are from the genes that encode α-

aminoadipate reductases (AAR), an enzyme responsible for the reduction of alpha-aminoadipic

acid, which is essential for the lysine biosynthesis pathway and is present in all fungal phyla

(Bushley and Turgeon 2010). In contrast, the remaining NRPS genes, and their respective A-

domains, show discontinuous and patchy distributions across Mucoromycota and

Zoopagomycota (Figure 3).

376   The most pronounced NRPS diversification in *Basidiobolus* are for genes that are

predicted to encode for siderophores, iron chelating metabolites. A-domains for predicted

siderophores are distributed throughout four clades including the three clades of major bacterial

genes and the fungal SID and SIDE clades (Figure 3, Sup. Fig. 1). Major bacterial clades (MBC)

exclusively comprise bacterial siderophore synthases, such as pyoverdine, yersiniabactin and

pyochelin (Bushley and Turgeon 2010), with the exception of the surfactin-like clade. Our

results show that genomes of *Mortierella* and *Basidiobolus* contain A-domains that are members

of the MBC, and that they are the only fungal representatives of these clades. SID clade contains

all NRPS associated with siderophore production from Ascomycota and Basidiomycota species.

All *Basidiobolus* isolates contain one NRPS A-domain in this clade, as well as three A-domains

from three NRPS gene models of *Conidiobolus coronatus* NRRL28638. Finally, SIDE, a clade

comprising NRPS genes responsible for the production of siderophores in filamentous

ascomycetes is expanded in *Basidiobolus* (Figure 3), which is also the only zygomycete with A-

domains clustered in this clade. These findings are consistent with enrichment of both bacterial

and fungal siderophores and are suggestive of the importance of iron metabolism in

Basidiobolus.

The CYCLO clade contains the A-domains for core genes associated with biosynthesis of

cyclic peptides, such as beauvericin and cyclosporin (Bushley and Turgeon 2010, Bushley et al

2013). Sister to all other CYCLO clade A-domains are the A-domains that comprise the NRPS

core gene model 387529 from *B. meristosporus* (Supplementary Figure 2). 387529 is expressed

at the highest rate of any SM gene under laboratory conditions (Figure 2).   It is annotated as a

tetra-modular gene model, that includes two N-methylation domains, four adenylation domains,

four condensation domains, and a TE domain. When compared to *simA*, the NRPS responsible

for biosynthesis of cyclosporin, structural similarities can be found, such as the presence of the

N-methylation domains and the TE terminator domain. Its phylogenetic and structural

similarities to *simA* suggest that 387529 results in the synthesis of a cyclic peptide with

methylated amino acid residues.

The surfactin-like clade contains A-domains for bacterial core genes with similarities,

but not identical, to the *Bacillus subtilis* surfactin termination module (*srfA-C* gene; Peypoux et

al. 1999) including the third, fourth and fifth A-domains of the five A-domain gene model

NP_930489.1 of *Photorhabdus luminescens* gene, two domains of the gene PvdD (AAX16295.1;

pyoverdine synthetase) of *Pseudomonas aeruginosa*, and a single A-domain of the bimodular

NRPS dbhF protein of *Bacillus subtilis* (AAD56240.1) which is involved in the biosynthesis of

the siderophore bacillibactin. This surfactin-like clade contains eleven A-domains predicted from

Basidiobolus genomes including the gene models 298977, 368581 and 372991 from *B.*

meristosporus CBS 931.73. Gene model 298977 showed no evidence for gene expression, while

gene models 368581 and 372991 show high rates of expression (Fig. 2). This is the first report of

the prediction of a surfactin-like gene in fungi, but surfactant production was recently reported in

Mortierella alpina (Baldeweg et al. 2019). These include malpinins, amphiphilic acetylated

hexapeptides that function as natural emulsifiers during lipid secretion, and malpibaldins,

hydrophobic cyclopentapeptides. This finding is consistent with the genomic data and reveals

that *Mortierella*, in addition to *Basidiobolus*, possesses homologs in the surfactin-like clade that

are phylogenetically different from *B. subtilis* surfactins.

Surfactins, encoded by the *srfA* gene cluster in *Bacillus subtilis*, are functionally active as

surfactants, as well as toxins and antibiotics. However, the surfactin genes from *B. subtilis* (SrfA-

AA, SrfA-AB and SrfA-AC) were included in our analysis and clustered in a different clade

within the MBC. A-domains from single module NRPS-like protein from *B. meristosporus* CBS

931.73 (Gene model 146993) and from a NRPS-PKS hybrid A domain from of *B. meristosporus*

B9252 (Gene model N161_14278) clustered with the *B. subtilis* SrfA genes. The placement of

this NRPS-PKS A-domain is interesting because surfactins are lipopeptides, which contain a

hydrophobic fatty acid chain, whose biosynthesis is consistent with an NRPS-PKS hybrid. The

A-domains from the remaining Mucoromycota and Zoopagomycota NRPS-PKS hybrids

clustered as sister to the original clade of Dikarya NRPS-PKS hybrids, supporting the hypothesis

of a single origin of fungal NRPS-PKS hybrids (Bushley and Turgeon 2010).

Mucoromycota and Zoopagomycota lack PKS diversity. – Polyketide synthases (PKS) are

abundant SM of Ascomycota and Basidiomycota and are involved in antibiotic production,

carotenoid biosynthesis and other functional roles. In contrast, literature on PKS diversity for

Mucoromycota and Zoopagomycota is limited. Our analyses update the phylogenetic

reconstruction reported by Kroken et al. (2013) by adding genomic information from

Chytridiomycota, Mucoromycota and Zoopagomycota (Supplementary Figures 3 and 4). Overall,

we report the discovery of two new clades: A clade of non-reducing PKS proteins (clade IV)

comprising Neocastimastigomycota and Basidiomycota, and a reducing PKS clade (reducing

PKS clade V) consisting of Neocastimastigomycota, Chytridiomycota and Basidiomycota. The

results of our PKS prediction revealed a number of potential PKS core genes in zygomycete

species (Supplementary Figures 3 and 4), but the results of our phylogenetic reconstruction show

that the KS domain of the predicted PKS gene models are fungal fatty acid synthases (FAS).

Only *Basidiobolus meristosporus* genomes possessed KS domains associated with fungal PKS,

either reducing or non-reducing.

A domain-by-domain presence/absence analysis of PKS genes models (Supplementary

Figures 5 and Figure 6) shows that in addition to AT and KT domains, a third domain (either

KR, DH or PP) is found in the majority of fungal PKSs (Supplementary Figure 5). Conversely,

the zygomycete genes in the FAS clade only possess the AT and/or KT domains, which are

domains common between FAS and PKS, including the majority of PKSs predicted for

Basidiobolus. However, *Basidiobolus* KS domains are found in both fungal and bacterial PKS

clades. Gene models 292783 and 237744 from both isolates of *B. meristosporous* cluster with

fungal reducing PKS II and possess the DH domain (Supplementary Figure 4), and the

expression levels of 292783 is consistent with an actively transcribed gene (Figure 2).

Terpene cyclase-like genes are expanded in zygomycetes. – Terpene cyclases (TC) are the most

common SM predicted for zygomycete species (Figure 1, Supplementary Table 2). Phylogenetic

reconstruction shows that zygomycete TC core genes are clearly distinct from Ascomycota and

Basidiomycota TC, where at least 4 new clades of predominantly zygomycete TC are found

(Figure 5, Supplementary Figure 7). Fungi II TC clade comprises TC from all zygomycete

genomes analyzed with the exception of *Piptocephalis cylindrospora* RSA 2659 single−cell

v3.0, *Phycomyces blakesleeanus NRRL1555* v2.0, and five genomes of *Rhizophagus irregularis*,

which show no prediction of TC in their genomes. No Dikarya TC genes cluster within this

clade. Fungi I, III and IV group TC genes are found almost exclusively in Mucoromycota

species, as well as some Dikarya species. TC genes from Fungi I are present only in

Mucoromycotina genomes. Functional annotations of TC genes from this clade indicate that

these TC genes code for proteins associated with squalene and phytoene synthases and are part

of the synthesis of carotenoids. Carotenoids are important compounds for the synthesis pathway

of trisporic acid, the main molecule responsible for initiating sexual reproduction in zygomycetes

(Burmester et al. 2007). Finally, *Basidiobolus* is the only non-Dikarya genus with TC genes

clustered within the Bacteria II clade of TC. Both these SM core genes show evidence for

expression comparable to housekeeping genes or other SM core genes (Figure 2). The presence

of bacterial-like TC genes in *Basidiobolus* present more evidence on the plasticity of the genome

of *Basidiobolus* and its ability to integrate and possibly express foreign SM associated DNA.

Horizontal Gene Transfer of SM genes to Basidiobolus? – Basidiobolus is a genus with a

complex biology, alternating ecologies, and multiple spore types. This complex biology is also

reflected at the genomic scale. The sequenced genomes show a larger genome size than other

zygomycetes, as well as a higher number of genes than any other Zoopagomycota genomes

sequenced to date. Our SM prediction assay is concordant with these patterns in which

Basidiobolus has an excess of SM gene clusters when compared to other zygomycetes. Evidence

points to HGT as a main driver of SM diversity in *Basidiobolus* as supported by the phylogenetic

reconstructions of NRPS, PKS and TC gene clusters with bacterial homologs. *Basidiobolus* and

Mortierella are the only fungi with genes associated with the bacterial clades in each of these SM

phylogenetic reconstructions. Moreover, these HGT candidates are integrated into the

Basidiobolus genome assembly and do not show evidence of artifactual assembly as evidenced

by discontinuous coverage (Supplementary Figure 8). The most abundant functional group of

SM core genes overall are siderophores and their overall functionality is supported by both the

RNA expression analyses (Figure 2) and siderophore plate assays (Figure 7).

One stage of the *Basidiobolus* life cycle the fungus lives as a gut endosymbiont where it

co-occurs with bacteria and other organisms that comprise the gut microbiome. Animal gut

environments can facilitate HGT between bacteria and fungi, as previously reported for the

zoosporic species of Neocallimastigomycetes (Chytridiomycota), which live in the ruminant gut

environment and whose genomes exhibit a 2-3.5% frequency of genes with HGT evidence

(Wang et al. 2018; Murphy et al. 2019). Phylogenomic analyses of the *Basidiobolus* NRPS A-

domains support a phylogenetic affinity with A-domains from bacterial taxa and more rarely

other fungi, a pattern most consistent with HGT. Our HGT survey comprised an extensive

search of reference genomes across the tree of life available in NCBI RefSeq, as well as all of the

gene models predicted for the Mucoromyocta and Zoopagomycota genomes. We find that 3% to

5% of all predicted gene models present in *Basidiobolous* genomes are consistent with signatures

of HGT from bacteria. However, the percentages radically change to 41% to 66% of predicted

SM core gene models with bacterial HGT evidence. These SM gene models with bacterial

signatures are highly abundant, with NRPS and NRPS-like genes comprising the top 25 ontology

categories of HGT genes in *B. meristosporus* (Supplementary Table 5). These results are

consistent with the life history of *Basidiobolus*, where the fungus lives in close proximity with

other microorganisms associated with the amphibian gut environment.

The additional analyses to discover distinct genetic features for HGT candidates showed

that the largest differences found are in intron number and intron length, but not in nucleotide

composition or by codon usage. A significantly smaller number of introns and smaller

normalized intron length in HGT candidates provide more support to the HGT hypothesis, where

we expected that bacterially transferred genes would maintain a smaller number and length of

introns. Intronic expansion of transferred genes into fungal species after an HGT event appears

to be rapid in order to reflect the genetic makeup across the genome (Da Lage et al. 2013) and

can explain the introns in some of the HGT candidates. However, up to 60% of the HGT

candidates still maintain absence of introns as expected for genes of bacterial origin. Finally, the

nucleotide composition of HGT candidates and fungal genes were indistinguishable. Reports

show that foreign genes with similar codon usage are more likely to become fixed on the

receiving genome (Medrano-Soto et al. 2004, Amorós-Montoya et al. 2010, Tuller 2011). We

interpet these results to indicate that horizontally transfered genes are evolving towards a similar

nucleotide composition of the fungal genome based on the 5-mer/codon usage assay, but still

maintain high protein similarity to and group with donor lineage copy in phylogenetic

reconstructions.

The taxonomic survey of our HGT analysis shows that a diverse array of bacteria may

have consistently contributed genetic information into the *Basidiobolus* genomes (Figure 6 and

Supplementary Table 4). The most abundant bacterial taxonomic groups associated with HGT

are the Proteobacteria, Firmicutes, Actinobacteria/high GC gram positive bacteria, and

Bacteriodetes (Figure 6, Supplementary Table 4). The proportion of HGT for each taxonomic

group appears to be consistent among the three *Basidiobolus* genomes, and there are

consistencies between the most common taxonomic groups responsible for HGT in *Basidiobolus*

and the reported composition of bacteria associated with the gut microbiome in amphibians

(Bletz et al. 2016, Kohl et al. 2013) and reptiles (Colston and Jackson, 2016, Costello et al.

2010).

Conclusions

Our results confirm that the majority of zygomycete fungi classified in Mucoromycota

and Zoopagomycota do not possess a large genomic potential for secondary metabolism.

Significant departures from this pattern exist, however, as exemplified by *Basidiobolus*, a genus

with a complex genomic evolution and potential for considerable and diverse secondary

metabolite production. First, it possesses larger than average genome with less than 8% content

of repetitive regions, but a genetic plasticity to integrate and express extrinsic DNA. Second, the

incorporation of extrinsic DNA is consistent with selection for increased SM production,

especially gene models that are related to the capture of resources available in anaerobic

conditions (iron chelation by siderophores) and metabolites that may play roles in antibiosis

(surfactin-like genes) or host interaction. Third, the amphibian gut environment predisposes

Basidiobolus to the acquisition of these SM core genes via HGT from co-inhabiting bacterial

species. More information is needed to further test these hypotheses, including sequencing of

additional *Basidiobolus* species with long read technologies; more accurate characterization of

amphibian microbiomes that test positive for *Basidiobolus*; and LC-MSMS characterization of

the *Basidiobolus* metabolome.

Materials and Methods

Data collection. – Annotated genome and amino-acid translation of predicted gene model sequences for three isolates of two species within the genus *Basidiobolus* were used in this study*: Basidiobolus meristosporus* CBS 931.73 (Mondo et al. 2017), isolated from gecko dung in the locality of Lamco, Ivory Coast; *B. meristosporus* B9252 (Chibucos et al. 2016) isolated from human eye in Saudi Arabia; and *B. heterosporus* B8920 (Chibucos et al. 2016) isolated from plant debris in India. The genomic sequence of *B. meristosporus* CBS 931.73 was sequenced with PacBio and annotated by Mondo et al. (2017) and obtained from the US Department of Energy Joint Genome Institute *MycoCosm* genome portal (https://mycocosm.jgi.doe.gov; Grigoriev et al. 2014). Genomic sequences and annotation of *B. meristosporus* B9252 and *B. heterosporus* B8920 sequenced by Chibucos et al. (2016) were obtained directly from the authors. The raw reads for these two species are available in GenBank (Accession numbers GCA_000697375.1 and GCA_000697455.1). Data sources for the remaining genomes included in these analyses are available in Supplementary Table 1.

Secondary metabolite gene cluster prediction. – Secondary Metabolite gene clusters for *Basidiobolus* species were predicted with AntiSMASH v4.2.0 (Weber et al. 2017) and the Secondary Metabolite Unique Regions Finder (SMURF; Khaldi et al. 2010) from the annotated genomes of the three isolates used in this study. The AntiSMASH prediction was performed on local HPC, while SMURF predictions were obtained by submission of the genomes the SMURF web server (http://smurf.jcvi.org/). Predictions were contrasted manually to determine shared clusters of secondary metabolites. Predicted SM proteins were retrieved for the genomes of 66 additional Mucoromycota and Zoopagomycota species based on the SMURF predictions

available at *MycoCosm* and by local prediction with AntiSMASH. Orthologous sets of core SM

genes across *Basidiobolus* were identified using OrthoFinder (Emms et al. 2015).

Secondary metabolite expression analysis. – To assess the expression of predicted SM proteins

in *B. meristosporus,* we calculated summarized counts of RNA transcript per million (TPM) for

genes in *B. meristosporus* CBS 931.73 isolate by aligning RNA-Seq reads to the assembled *B.*

meristosporus CBS 931.73 genome with HiSat v2.1.0 (Kim et al. 2019). The aligned sequence

reads were processed with HTS-seq (Anders et al. 2014) to generate the counts of overlapping

reads found for each gene and the normalized TPM for the genes was calculated using the cpm

function in edgeR (Robinson et al. 2010). A distribution of RNA-Seq read counts per gene was

plotted using the ggplot2 package in the R statistical framework (R Core Team 2018).

Phylogenomic analyses. – Phylogenetic analyses were used to assess the evolutionary

relationships of the NRPS, PKS, and terpene cyclase/synthase predicted proteins. For the NRPS

genes, the adenylation domains (A-domains) were identified by hmmsearch from HMMer 3.0

suite (Eddy 2004), using the A-domain profile reported by Bushley and Turgeon (2010) as a

reference profile HMM. The predicted A domains were extracted from the resulting HMMER

table into a FASTA file using the esl-reformat program included in the HMMER suite. The

predicted A-domains for all *Basidiobolus*, Mucoromycota and Zoopagomycota species were

added to an A-domain amino-acid alignment reported by Bushley and Turgeon (2010) with

MAFFT v7 (Katoh et al. 2017) (Additional file 1). This reference alignment contains A-domains

from NRPS proteins from nine major subfamilies of fungal and bacterial NRPS proteins. The

phylogenetic domain tree was constructed using a maximum likelihood approach implemented in

RAxML v. 8.2.11 (Stamatakis et al. 2014) with the JTT amino acid substitution matrix, after

model selection using the PROTGAMMAAUTO option, and 1000 bootstrap replicates

(raxmlHPC-PTHREADS -T 12 -n NRPS -s infile.fasta -f a -x 12345 -p 12345 -m PROTCATJTT

–N 1000). A graphical representation of the A-domains from the NRPS/NRPS-like core gene

models (Fig. X) was constructed by coloring the A-domain position in the core gene according to

its phylogenetic origin using the ggplot2 R package (Wickham, 2016).

For the PKS genes, the KS domains of all predicted PKS proteins from the *Basidiobolus*,

Mucoromycota and Zoopagomycota species were identified using hmmsearch, using the KS

domain profile (PF001009) available in PFAM v31 (Finn et al. 2009). The predicted KS domains

for all *Basidiobolus*, Mucoromycota and Zoopagomycota species were added to an existing KS

domain amino-acid alignment reported by Kroken et al. (2003) using MAFFT v7 (Katoh et al.

2017) (Additional file 2). This existing alignment contained KS domains from PKS proteins

from reduced and unreduced PKS from bacterial and fungal species. The Kroken et al. (2003)

database was expanded by adding predicted KS domains from PKS proteins of additional

published fungal genomes in order to include more fungal diversity in the dataset: Eight

Ascomycota isolates (*Aspergillus nidulans* FGSC A4, *Beauveria bassiana* ARSEF 2860,

Capronia coronata CBS 617.96, *Capronia semiimmersa* CBS 27337, *Cladophialophora*

bantiana CBS 173.52, *Cochliobolus victoriae* FI3 v1.0, *Microsporum canis* CBS 113480,

Trichoderma atroviride v2.0), nine Basidiomycota isolates (*Acaromyces ingoldii* MCA 4198

v1.0, *Fibroporia radiculosa* TFFH 294, *Fomitiporia mediterranea* v1.0, *Gloeophyllum trabeum*

v1.0, *Gymnopus luxurians* v1.0, *Laccaria bicolor* v2.0, *Microbotryum lychnidis−dioicae* p1A1

Lamole, *Piloderma croceum* F 1598 v1.0, *Pisolithus tinctorius* Marx 270 v1.0), four

Neocallimastigomycota isolates (*Anaeromyces robustus* v1.0, *Orpinomyces sp.*, *Piromyces finnis*

v3.0, *Piromyces sp. E2* v1.0) and one Chytridiomycota species (*Spizellomyces punctatus* DAOM

BR117). The predicted PKS proteins from additional species were all obtained from the DOE-

JGI *MycoCosm* genome portal by searching for all genes with "PKS" on the SM annotation from

MycoCosm. The KS domains were identified for the subset of PKS proteins using a HMMER KS

profile as mentioned above. A phylogenetic tree was reconstructed using maximum likelihood

using all KS domains using similar parameters described in the NRPS step.

The terpene cyclase (TC) proteins were predicted in AntiSMASH for all 69 assembled

genome sequences from *Basidiobolus*, Mucoromycota and Zoopagomycota. To include

additional fungal TC proteins in the phylogenetic reconstruction, we identified TC proteins from

58 published genomes of Dikarya isolates (21 Basidiomycetes and 37 Ascomycetes) available in

DOE-JGI *MycoCosm*, each from a different family (Supplementary Table 3). One isolate per

family was randomly selected for the analysis. To screen for TC proteins in Dikarya,

OrthoFinder was used to build orthologous clusters of genes between *Basidiobolus*,

Mucoromycota and Zoopagomycota TC and the Dikarya proteome dataset. Dikarya proteins

clustered within orthologous groups that contain TC of *Basidiobolus*, Mucoromycota and

Zoopagomycota were considered valid TC orthologs and used in downstream analyses. Finally,

we identified bacterial TC to evaluate the potential for HGT into *Basidiobolus,* Mucoromycota or

Zoopagomycota species. Bacterial TC's were identified by screening each protein from the

RefSeq, release 87 (May 2018, O'Leary et al. 2016) FASTA dataset against a BLAST database

(Altschul et al. 1997), one from each of the zygomycete orthologous groups identified by

OrthoFinder in the previous step, for a total of four protein sequences in the database. The

BLASTP program was used to perform the searches, using an e-value of $1e^{-10}$ and the

BLOSUM62 substitution matrix. Positive matches from the BLASTP assay were used as

bacterial TC for subsequent analyses. A multi-sequence alignment containing all predicted TC

from *Basidiobolus*, Mucoromycota and Zoopagomycota species, the orthologous TC from

additional fungal species, and the TC proteins from reference bacterial predicted gene models

result of the BLASTP search was performed in MAFFT v7 using the G-INS-1 algorithm for

progressive global alignment (Katoh et al. 2017) (Additional file 3). A phylogenetic tree was

reconstructed using maximum likelihood in RAxML using similar parameters as mentioned

before.

Horizontal gene transfer in Basidiobolus *species.* – Identification of genic regions with evidence

for horizontal gene transfer (HGT) from bacteria was performed by searching all translated

proteins from the predicted gene models of the *Basidiobolus* isolates against a custom BLAST

protein database. This database included all amino-acid sequences available in the NCBI RefSeq

proteomics database and all available amino-acid sequences for Mucoromycota and

Zoopagomycota species at the DOE-JGI *MycoCosm* genome portal. The proteins were searched

with BLASTP against the combined RefSeq/Mucoromycota/Zoopagomycota custom database,

using an e-value cutoff of $1e^{-10}$ and the BLOSUM62 substitution matrix. A summary of the

taxonomy identifier for all best hits was obtained by identifying whether the best hit was a

Mucoromycota/Zoopagomycota protein, or a RefSeq protein. We used the rentrez package

(Winter, 2017) in the R statistical framework to extract the top-ten taxonomic identifiers from

the NCBI database when hits did not correspond to Mucoromycota/Zoopagomycota. Proteins

that had no hits to a fungal protein and only hits to a bacterial protein were considered candidate

HGT genes.

To increase the accuracy of the prediction of genes product of HGT, we tested whether

HGT candidate genes showed signs of errant assembly or in silica incorporation into the

Basidiobolus genomes. The mean genomic read coverage of each candidate gene was calculated

and compared to the mean coverage of the scaffold harboring the HGT candidate gene for all

three *Basidiobolus* isolates. A z-score was calculated in R to determine the number of standard

deviations of the HGT candidate from the mean coverage of the harboring scaffold. All HGT

candidates with standard deviations greater or less than two were removed from the analysis.

The genomic reads were mapped to each reference genome using BWA (Li and Durbin, 2009).

Coverage per HGT and harboring scaffold was estimated using samtools (Li et al. 2009). A

summary plot of the proportion of genes with bacterial best hits was constructed using the

ggplot2 package in R.

To identify differences in the GC content between HGT candidates and genes of fungal

origin, a paired t-test was performed between the mean GC content for HGT candidates and

fungal genes for the predicted genes of *Basidiobolus meristosporus* CBS 931.73 as this species

has the genome with longest contigs and best assemblies overall. In addition, to identify if the

HGT candidate genes had a reduced number of introns than the fungal genes, a summary of the

number of introns and normalized intron length (intron length divided by gene model length) was

performed in the GenomicRanges R package (Lawrence et al. 2013). A Kruskal-Wallis test was

performed in R to identify significant differences in the number of introns and the normalized

intron length for the HGT candidates and the fungal genes. A nucleotide composition analysis

based on 5-mers and codon usage was performed to observe differences between candidate HGT

genes and fungal genes for *Basidiobolus meristosporus* CBS 931.73. The 5-mer analysis was

conducted in all predicted coding sequences from the annotated gene models using the

oligonucleotideFrequency function of the Biostring R package (Pagès et al. 2019). Codon usage

was estimated using the uco function of the seqinr R package (Charif and Lobry 2007). Both

these indices were divided by gene length to normalize the nucleotide composition by the effect

of gene length. Principal component analyses were performed in R for both 5-mer and codon

usage analysis to compare the HGT candidates to the fungal candidate genes. Lastly, the putative

functions of the genes with evidence for HGT were summarized using the functional annotations

available in the gene format file (GFF) of *B. meristosporus* CBS 931.73.

Measuring siderophore activity in Basidiobolus. – To measure the siderophore activity (chelation

of ferric ions) of *Basidiobolus*, an assay of detection of siderophore activity based on the

universal chrome azurol S (CAS) assay (Andrews et al. 2016) was performed for the strain

Basidiobolus meristosporus CBS 931.73. This colorimetric assay uses a complex of Fe(III) –

CAS – DDAPS (Surfactant). When this complex is combined with acetate yeast agar (AY agar),

it results in a greenish-blue color, where the color changes to yellow upon the removal of the

iron. A total of 450 ml of AY agar (Andrews et al. 2016) was mixed with 50 ml of autoclaved

10X CAS assay (20 ml of 10 mM $Fe(NO_3)_3$, 40 ml of 10 mM chrome azurol S, and 100 ml of 10

mM DDAPS). A 10 cm layer of the AY-CAS media was poured in small petri dishes and cooled.

An upper layer of 10 cm of AY agar was poured after cooling, and the media was left to diffuse

overnight and stored at 4°C for 24 hours. *B. meristosporus*, *Conidiobolus thromboides* FSU 785

(negative control)*, and Cladosporium sp.* from the *herbarum* species complex PE-07 (positive

control) were transferred into the AY-CAS plates by transferring a small amount of mycelium

via a sterile toothpick and piercing the media in the center. The assay was performed in triplicate

for each isolate used. Cultures were grown at room temperature for 12 days. Siderophore activity

was measured as the area of the plate that has changed to yellow color, when compared to a

negative control and an empty petri dish with AY-CAS agar. Pictures of the plates were taken.

These images were imported into Adobe Photoshop CC 2019 and size-corrected to 5.5 cm

(diameter of the small petri dish). All images were concatenated in the same file. The Color

Range tool of Adobe Photoshop CC tool was used to measure the yellow area for all plates. The

area measurements were exported into R, where an analysis of variance was performed to

determine significant differences across the siderophore activity of the strains.

Data availability

All additional files included as supplementary materials in this manuscript and custom

scripts used for this analysis can be found in the ZyGoLife GitHub repository

(https://github.com/zygolife/Basidiobolus_SM_repo).

Acknowledgments
725

We would like to thank Dr. Vincent Bruno and Dr. Ashraf S. Ibrahim for kindly sharing the files

of the genome annotation, assembly and gene predictions of *B. meristosporus* B9252 and *B.*

heterosporus B8920; Dr. Marc Cubeta and Dr. Gregory Bonito for discussions and

recommendations for additional experiments regarding siderophore assays and surfactin-like

genes; Dr. Gregory Bonito for allowing us to use the unpublished genome sequences of

Mortierella humilis PMI_1414, *Mortierella sp.* GBAus27b; Dr. Olafur S. Andrésson for

allowing us to use the unpublished genome sequences of *Lobaria pulmonaria*; Dr. Paul Dryer for

allowing us to use the unpublished genome sequence of *Xanthoria parietina 46-1-SA22*; Dr.

Francis Martin and Dr. Pierre Gladieux for allowing us to use the unpublished genome sequence

of *Achaetomium strumarium CBS333.67*; Dr. Francis Martin for allowing us to use the

unpublished genome sequences of *Boletus edulis V1* and *Amylostereum chailletii DWAch2*; Dr.

Marie-Noëlle Rosso for allowing us to use the unpublished genome sequences of

Abortiporus biennis CIRM-BRFM1778; Dr. Francis Martin and Dr. Gregory Bonito for allowing

us to use the unpublished genome sequence of *Atractiellales rhizophila v2.0;* the 1000 Fungal

Genomes Project for the use of the dikarya unpublished genomes; and Sabrina Heitmann, Dr. Ed

Barge, Dr. Devin Leopold and Dr. Posy Busby for the *Cladosporium sp.* from the *herbarum*

species complex PE-07 sample. This work was supported by the National Science Foundation

(DEB-1441604 to JWS; DEB-1557110 and DEB-1441715 to JES). The work conducted by the

U.S. Department of Energy Joint Genome Institute, a DOE Office of Science User Facility, is

supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-

AC02-05CH11231. Any opinions, findings, and conclusions or recommendations expressed in

this material are those of the author(s) and do not necessarily reflect the views of the National

Science Foundation.

References

Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W. and

Lipman, D.J., 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database

search programs. *Nucleic Acids Research*, *25*(17), pp.3389-3402.

Amorós-Moya, D., Bedhomme, S., Hermann, M. and Bravo, I.G., 2010. Evolution in

regulatory regions rapidly compensates the cost of nonoptimal codon usage. *Molecular Biology*

and Evolution, 27(9), pp.2141-2151.

Anders, S., Pyl, P.T. and Huber, W., 2015. HTSeq—a Python framework to work with

high-throughput sequencing data. *Bioinformatics*, *31*(2), pp.166-169.

Andrews, M. Y., Santelli, C. M., & Duckworth, O. W. 2016. Layer plate CAS assay for

the quantitation of siderophore production and determination of exudation patterns for fungi.

Journal of Microbiological Methods, 121(C), pp. 41–43.

http://doi.org/10.1016/j.mimet.2015.12.012

Ahrendt, S.R., Quandt, C.A., Ciobanu, D., Clum, A., Salamov, A., Andreopoulos, B.,

Cheng, J.F., Woyke, T., Pelin, A., Henrissat, B. and Reynolds, N.K., 2018. Leveraging single-

cell genomics to expand the fungal tree of life. *Nature Microbiology, 3*(12), p.1417.

Armaleo, D., Müller, O., Lutzoni, F., Andrésson, Ó.S., Blanc, G., Bode, H.B., Collart,

F.R., Dal Grande, F., Dietrich, F., Grigoriev, I.V. and Joneson, S., 2019. The lichen symbiosis

re-viewed through the genomes of Cladonia grayi and its algal partner Asterochloris glomerata.

BMC genomics, 20(1), p.605

Baldeweg, F., Warncke, P., Fischer, D. and Gressler, M., 2019. Fungal Biosurfactants

from Mortierella alpina. *Organic Letters*, *21*(5), pp.1444-1448.

Bletz, M.C., Goedbloed, D.J., Sanchez, E., Reinhardt, T., Tebbe, C.C., Bhuju, S., Geffers,

R., Jarek, M., Vences, M. and Steinfartz, S., 2016. Amphibian gut microbiota shifts differentially

in community structure but converges on habitat-specific predicted functions. *Nature*

Communications, *7*, p.13699.

Blin, K., Wolf, T., Chevrette, M.G., Lu, X., Schwalen, C.J., Kautsar, S.A., Suarez Duran,

H.G., de Los Santos, E.L., Kim, H.U., Nave, M. and Dickschat, J.S., 2017. antiSMASH 4.0—

improvements in chemistry prediction and gene cluster boundary identification. *Nucleic Acids*

Research, *45*(W1), pp.W36-W41.

Brakhage, A.A., 2013. Regulation of fungal secondary metabolism. *Nature Reviews*

Microbiology, *11*(1), p.21.

Braun V., Hantke K. (2013) The Tricky Ways Bacteria Cope with Iron Limitation. In:

Chakraborty R., Braun V., Hantke K., Cornelis P. (eds) Iron Uptake in Bacteria with Emphasis

on E. coli and Pseudomonas. SpringerBriefs in Molecular Science. Springer, Dordrecht

Burmester, A., Richter, M., Schultze, K., Voelz, K., Schachtschabel, D., Boland, W.,

Wöstemeyer, J. and Schimek, C., 2007. Cleavage of β-carotene as the first step in sexual

hormone synthesis in zygomycetes is mediated by a trisporic acid regulated β-carotene

oxygenase. *Fungal Genetics and Biology*, *44*(11), pp.1096-1108.

Burmester, A., Shelest, E., Glöckner, G., Heddergott, C., Schindler, S., Staib, P., Heidel,

A., Felder, M., Petzold, A., Szafranski, K. and Feuermann, M., 2011. Comparative and

functional genomics provide insights into the pathogenicity of dermatophytic fungi. *Genome*

biology, 12(1), p.R7

Bushley, K.E. and Turgeon, B.G., 2010. Phylogenomics reveals subfamilies of fungal

nonribosomal peptide synthetases and their evolutionary relationships. *BMC Evolutionary*

Biology, *10*(1), p.26.

Bushley, K.E., Raja, R., Jaiswal, P., Cumbie, J.S., Nonogaki, M., Boyd, A.E., Owensby,

C.A., Knaus, B.J., Elser, J., Miller, D. and Di, Y., 2013. The genome of *Tolypocladium inflatum:*

evolution, organization, and expression of the cyclosporin biosynthetic gene cluster. *PLoS*

genetics, 9(6).

Capella-Gutiérrez, S., Silla-Martínez, J.M. and Gabaldón, T., 2009. trimAl: a tool for

automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, *25*(15),

pp.1972-1973.

Chang, Y., Wang, S., Sekimoto, S., Aerts, A.L., Choi, C., Clum, A., LaButti, K.M.,

Lindquist, E.A., Yee, Ngan C., Ohm, R.A., Salamov, A.A., Grigoriev, I.V., Spatafora, J.W., and

Berbee M.L. 2015. Phylogenomic Analyses Indicate that Early Fungi Evolved Digesting Cell

Walls of Algal Ancestors of Land Plants. *Genome Biology and Evolution*, 14;7(6):1590-601. doi:

10.1093/gbe/evv090.

Chang, Y., Desirò, A., Na, H., Sandor, L., Lipzen, A., Clum, A., Barry, K., Grigoriev,

I.V., Martin, F.M., Stajich, J.E. and Smith, M.E., 2019. Phylogenomics of Endogonaceae and

evolution of mycorrhizas within Mucoromycota. *New Phytologist*, 222(1), pp.511-525.

Charif, D. and Lobry, J.R., 2007. SeqinR 1.0-2: a contributed package to the R project for

statistical computing devoted to biological sequences retrieval and analysis. In Structural

approaches to sequence evolution (pp. 207-232). Springer, Berlin, Heidelberg.

Chen, E.C.H., Morin, E., Beaudet, D., Noel, J., Yildirir, G., Ndikumana, S., Charron, P.,

St-Onge, C., Giorgi J., Kruger, M., Marton, T., Ropars, J., Grigoriev, I.V., Hainaut, M.,

Henrissat, B., Roux, C., Martin, F. and Corradi, N. 2018. High intraspecific genome diversity in

the model arbuscular mycorrhizal symbiont Rhizophagus irregularis. *New Phytologist,*

220(4):1161-1171. doi: 10.1111/nph.14989.

Chibucos, M.C., Soliman, S., Gebremariam, T., Lee, H., Daugherty, S., Orvis, J., Shetty,

A.C., Crabtree, J., Hazen, T.H., Etienne, K.A. and Kumari, P., 2016. An integrated genomic and

transcriptomic survey of mucormycosis-causing fungi. *Nature Communications*, *7*, p.12218.

Collemare, J., Pianfetti, M., Houlle, A.E., Morin, D., Camborde, L., Gagey, M.J.,

Barbisan, C., Fudal, I., Lebrun, M.H. and Böhnert, H.U., 2008. Magnaporthe grisea avirulence

gene ACE1 belongs to an infection specific gene cluster involved in secondary metabolism.

New Phytologist, *179*(1), pp.196-208.

Colston, T.J. and Jackson, C.R., 2016. Microbiome evolution along divergent branches of

the vertebrate tree of life: what is known and unknown. *Molecular Ecology*, *25*(16), pp.3776-

3800.

Corrochano, L.M., Kuo, A., Marcet-Houben, M., Polaino, S., Salamov, A., Villalobos-

Escobedo, J.M., Grimwood, J., Alvarez, M.I., Avalos, J., Bauer, D., Benito, E.P., Benoit, I.,

Burger, G., Camino, L.P., Canovas, D., Cerda-Olmedo, E., Cheng, J.F., Dominguez, A., Elias,

M., Eslava, AP., Glaser, F., Gutierrez, G., Heitman, J., Henrissat, B., Iturriaga, E.A., Lang, B.F.,

Lavin, J.L., Lee, S.C., Li, W., Lindquist, E., Lopez-Garcia, S., Luque, E.M., Marcos, A.T.,

Martin, J., McCluskey, K., Medina, H.R., Miralles-Duran, A., Miyazaki, A., Munoz-Torres, E.,

Oguiza, J.A., Ohm, R.A., Olmedo, M., Oreja,s M., Ortiz-Castellanos, L., Pisabarro, A.G.,

Rodriguez-Romero, J., Ruiz-Herrera, J., Ruiz-Vazquez, R., Sanz, C., Schackwitz, W., Shahriari,

M., Shelest, E., Silva-Franco, F., Soanes, D., Syed, K., Tagua, V.G., Talbot, N.J., Thon, M.R.,

Tice, H., de Vries, R.P., Wiebenga, A., Yadav, J.S., Braun, E.L., Baker, S,E., Garre, V.,

Schmutz, J., Horwitz, B.A., Torres-Martinez, S., Idnurm, A., Herrera-Estrella, A., Gabaldon and

T., Grigoriev, I.V. 2016. Expansion of Signal Transduction Pathways in Fungi by Extensive

Genome Duplication. *Current Biology*, 26(12):1577-1584. doi: 10.1016/j.cub.2016.04.038.

Costello, E.K., Gordon, J.I., Secor, S.M. and Knight, R., 2010. Postprandial remodeling

of the gut microbiota in Burmese pythons. *The ISME journal*, *4*(11), p.1375.

Da Lage, J.L., Binder, M., Hua-Van, A., Janeček, Š. and Casane, D., 2013. Gene make-

up: rapid and massive intron gains after horizontal transfer of a bacterial α-amylase gene to

Basidiomycetes. *BMC evolutionary biology*, 13(1), p.40.

DiGuistini, S., Wang, Y., Liao, N.Y., Taylor, G., Tanguay, P., Feau, N., Henrissat, B.,

Chan, S.K., Hesse-Orce, U., Alamouti, S.M. and Tsui, C.K., 2011. Genome and transcriptome

analyses of the mountain pine beetle-fungal symbiont Grosmannia clavigera, a lodgepole pine

pathogen. *Proceedings of the National Academy of Sciences*, 108(6), pp.2504-2509.

Dimkpa, C., 2016. Microbial siderophores: Production, detection and application in

agriculture and environment. *Endocytobiosis & Cell Research*, *27*(2), pp. 7-16

Eddy, S.R., 2004. What is a hidden Markov model?. *Nature Biotechnology*, *22*(10),

p.1315.

Emms, D.M. and Kelly, S., 2015. OrthoFinder: solving fundamental biases in whole

genome comparisons dramatically improves orthogroup inference accuracy. *Genome Biology*,

16(1), p.157.

Finn, R.D., Mistry, J., Tate, J., Coggill, P., Heger, A., Pollington, J.E., Gavin, O.L.,

Gunasekaran, P., Ceric, G., Forslund, K. and Holm, L., 2009. The Pfam protein families

database. *Nucleic Acids Research*, *38*(suppl_1), pp.D211-D222.

Floudas, D., Binder, M., Riley, R., Barry, K., Blanchette, R.A., Henrissat, B., Martínez,

A.T., Otillar, R., Spatafora, J.W., Yadav, J.S. and Aerts, A., 2012. The Paleozoic origin of

enzymatic lignin decomposition reconstructed from 31 fungal genomes. *Science,* 336(6089),

pp.1715-1719.

Ffrench-Constant, R.H., Waterfield, N., Burland, V., Perna, N.T., Daborn, P.J., Bowen,

D. and Blattner, F.R., 2000. A genomic sample sequence of the entomopathogenic bacterium

Photorhabdus luminescens W14: potential implications for virulence. *Applied Environmental*

Microbiology, *66*(8), pp.3310-3329.

Gaillard DL (2014) Population genetics and microbial communities of the Gopher

Tortoise (Gopherus polyphemus). Unpublished MS Thesis, The University of Southern

Mississippi.

Gawad, C., Koh, W. and Quake, S.R., 2016. Single-cell genome sequencing: current state

of the science. *Nature Reviews Genetics*, *17*(3), p.175.

Gostinčar, C., Ohm, R.A., Kogej, T., Sonjak, S., Turk, M., Zajc, J., Zalar, P., Grube, M.,

Sun, H., Han, J. and Sharma, A., 2014. Genome sequencing of four Aureobasidium pullulans

varieties: biotechnological potential, stress tolerance, and description of new species. *BMC*

genomics, 15(1), p.549.

Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, Riley R, Salamov A, Zhao

X, Korzeniewski F, Smirnova T, Nordberg H, Dubchak I, Shabalov I., 2014. MycoCosm portal:

gearing up for 1000 fungal genomes. Nucleic Acids Res. 42(1):D699-704.

Haridas, S., Albert, R., Binder, M., Bloem, J., LaButti, K., Salamov, A., Andreopoulos,

B., Baker, S.E., Barry, K., Bills, G. and Bluhm, B.H., 2020. 101 Dothideomycetes genomes: a

test case for predicting lifestyles and emergence of pathogens. *Studies in Mycology*.

https://doi.org/10.1016/j.simyco.2020.01.003

Helaly, S.E., Thongbai, B. and Stadler, M., 2018. Diversity of biologically active

secondary metabolites from endophytic and saprotrophic fungi of the ascomycete order

Xylariales. *Natural Product Reports*, *35*(9), pp.992-1014.

Hong, P.Y., Wheeler, E., Cann, I.K. and Mackie, R.I., 2011. Phylogenetic analysis of the

fecal microbial community in herbivorous land and marine iguanas of the Galápagos Islands

using 16S rRNA-based pyrosequencing. *The ISME journal*, *5*(9), p.1461.

Katoh, K., Rozewicki, J. and Yamada, K.D., 2017. MAFFT online service: multiple

sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics*.

Karlsson, M., Atanasova, L., Jensen, D.F. and Zeilinger, S., 2016. Necrotrophic

mycoparasites and their genomes. *Spectrum*, *5*(2).

Keller, N.P., Turner, G. and Bennett, J.W., 2005. Fungal secondary metabolism—from

biochemistry to genomics. *Nature Reviews Microbiology*, *3*(12), p.937.

Khaldi, N., Seifuddin, F.T., Turner, G., Haft, D., Nierman, W.C., Wolfe, K.H. and

Fedorova, N.D., 2010. SMURF: genomic mapping of fungal secondary metabolite clusters.

Fungal Genetics and Biology, *47*(9), pp.736-741.

Kohl, K.D., Cary, T.L., Karasov, W.H. and Dearing, M.D., 2013. Restructuring of the

amphibian gut microbiota through metamorphosis. *Environmental Microbiology Reports*, *5*(6),

pp.899-903.

Kohler, A., Kuo, A., Nagy, L.G., Morin, E., Barry, K.W., Buscot, F., Canbäck, B., Choi,

C., Cichocki, N., Clum, A. and Colpaert, J., 2015. Convergent losses of decay mechanisms and

rapid turnover of symbiosis genes in mycorrhizal mutualists. *Nature Genetics*, *47*(4), p.410.

Kroken, S., Glass, N.L., Taylor, J.W., Yoder, O.C. and Turgeon, B.G., 2003.

Phylogenomic analysis of type I polyketide synthase genes in pathogenic and saprobic

ascomycetes. *Proceedings of the National Academy of Sciences*, *100*(26), pp.15670-15675.

Lawrence, M., Huber, W., Pages, H., Aboyoun, P., Carlson, M., Gentleman, R., Morgan,

M.T. and Carey, V.J., 2013. Software for computing and annotating genomic ranges. *PLoS*

Computational Biology, 9(8), p.e1003118.

Lastovetsky, O.A., Gaspar, M.L., Mondo, S.J., LaButti, K.M., Sandor, L., Grigoriev, IV.,

Henry, S.A. and Pawlowska, T.E. 2016. Lipid metabolic changes in an early divergent fungus

govern the establishment of a mutualistic symbiosis with endobacteria. *Proceedings of the*

National Academy of Sciences, 113(52):15102-15107. doi: 10.1073/pnas.1615148113.

Li, H. and Durbin, R., 2009. Fast and accurate short read alignment with Burrows–

Wheeler transform. *Bioinformatics*, *25*(14), pp.1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,

Abecasis, G. and Durbin, R., 2009. The sequence alignment/map format and SAMtools.

Bioinformatics, *25*(16), pp.2078-2079.

Luo, H., Hong, S.Y., Sgambelluri, R.M., Angelos, E., Li, X. and Walton, J.D., 2014.

Peptide macrocyclization catalyzed by a prolyl oligopeptidase involved in α-amanitin

biosynthesis. *Chemistry & Biology*, *21*(12), pp.1610-1617.

Ma, L.J., Ibrahim, A.S., Skory, C., Grabherr, M.G., Burger, G., Butler, M., Elias, M.,

Idnurm, A., Lang, B.F., Sone, T., Abe, A., Calvo, S.E., Corrochano, L.M., Engels, R., Fu, J.,

Hansberg, W., Kim, J.M., Kodira, C.D., Koehrsen, M.J., Liu, B., Miranda-Saavedra, D.,

O'Leary, S., Ortiz-Castellanos, L., Poulter, R., Rodriguez-Romero, J., Ruiz-Herrera, J., Shen,

Y.Q., Zeng, Q., Galagan, J., Birren, B.W., Cuomo, C.A. and Wickes B.L. 2009. Genomic

analysis of the basal lineage fungus Rhizopus oryzae reveals a whole-genome duplication. *PLoS*

Genetics, 5(7):e1000549. doi: 10.1371/journal.pgen.1000549.

Martino, E., Morin, E., Grelet, G.A., Kuo, A., Kohler, A., Daghino, S., Barry, K.W.,

Cichocki, N., Clum, A., Dockter, R.B. and Hainaut, M., 2018. Comparative genomics and

transcriptomics depict ericoid mycorrhizal fungi as versatile saprotrophs and plant mutualists.

New Phytologist, 217(3), pp.1213-1229.

Medrano-Soto, A., Moreno-Hagelsieb, G., Vinuesa, P., Christen, J.A. and Collado-Vides,

J., 2004. Successful lateral transfer requires codon usage compatibility between foreign genes

and recipient genomes. *Molecular Biology and Evolution, 21*(10), pp.1884-1894.

Mondo, S.J., Dannebaum, R.O., Kuo, R.C., Louie, K.B., Bewick, A.J., LaButti, K.,

Haridas, S., Kuo, A., Salamov, A., Ahrendt, S.R. and Lau, R., 2017. Widespread adenine N6-

methylation of active genes in fungi. *Nature Genetics*, *49*(6), p.964.

Morin, E., Kohler, A., Baker, A.R., Foulongne-Oriol, M., Lombard, V., Nagye, L.G.,

Ohm, R.A., Patyshakuliyeva, A., Brun, A., Aerts, A.L. and Bailey, A.M., 2012. Genome

sequence of the button mushroom Agaricus bisporus reveals mechanisms governing adaptation

to a humic-rich ecological niche. *Proceedings of the National Academy of Sciences*, 109(43),

pp.17501-17506.

Murphy, C., Youssef, N., Hanafy, R.A., Couger, M.B., Stajich, J.E., Wang, Y., Baker, K.,

Dagar, S., Griffith, G., Farag, I. and Callaghan, T.M., 2019. Horizontal gene transfer as an

indispensible driver for Neocallimastigomycota evolution into a distinct gut-dwelling fungal

lineage. *bioRxiv*, p.487215.

Nagy, G., Farkas, A., Csernetics, Á., Bencsik, O., Szekeres, A., Nyilasi, I., Vágvölgyi, C.

and Papp, T., 2014. Transcription of the three HMG-CoA reductase genes of Mucor

circinelloides. *BMC Microbiology*, *14*(1), p.93.

Nagy, L.G., Riley, R., Tritt, A., Adam, C., Daum, C., Floudas, D., Sun, H., Yadav, J.S.,

Pangilinan, J., Larsson, K.H. and Matsuura, K., 2016. Comparative genomics of early-diverging

mushroom-forming fungi provides insights into the origins of lignocellulose decay capabilities.

Molecular Biology and Evolution, 33(4), pp.959-970.

Nguyen, T.A., Cissé, O.H., Wong, J.Y., Zheng, P., Hewitt, D., Nowrousian, M., Stajich,

J.E. and Jedd, G., 2017. Innovation and constraint leading to complex multicellularity in the

Ascomycota. *Nature communications*, 8(1), pp.1-13.

Ohm, R.A., Feau, N., Henrissat, B., Schoch, C.L., Horwitz, B.A., Barry, K.W., Condon,

B.J., Copeland, A.C., Dhillon, B., Glaser, F. and Hesse, C.N., 2012. Diverse lifestyles and

strategies of plant pathogenesis encoded in the genomes of eighteen Dothideomycetes fungi.

PLoS Pathogens, 8(12).

Okagaki, L.H., Nunes, C.C., Sailsbery, J., Clay, B., Brown, D., John, T., Oh, Y., Young,

N., Fitzgerald, M., Haas, B.J. and Zeng, Q., 2015. Genome sequences of three phytopathogenic

species of the Magnaporthaceae family of fungi. *G3: Genes, Genomes, Genetics*, 5(12), pp.2539-

2545.

O'Leary, N.A., Wright, M.W., Brister, J.R., Ciufo, S., Haddad, D., McVeigh, R., Rajput,

B., Robbertse, B., Smith-White, B., Ako-Adjei, D. and Astashyn, A., 2015. Reference sequence

(RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation.

Nucleic Acids Research, *44*(D1), pp.D733-D745.

Osbourn, A., 2010. Gene clusters for secondary metabolic pathways: an emerging theme

in plant biology. *Plant Physiology*, *154*(2), pp.531-535.

Pagès, H., Aboyoun, P., Gentleman, R. and DebRoy, R. 2019. Biostrings: Efficient

manipulation of biological strings. R package version 2.52.0.

Padamsee, M., Kumar, T.A., Riley, R., Binder, M., Boyd, A., Calvo, A.M., Furukawa,

K., Hesse, C., Hohmann, S., James, T.Y. and LaButti, K., 2012. The genome of the xerotolerant

mold Wallemia sebi reveals adaptations to osmotic stress and suggests cryptic sexual

reproduction. *Fungal Genetics and Biology*, 49(3), pp.217-226.

Paterson, R.R.M., 2008. Cordyceps–a traditional Chinese medicine and another fungal

therapeutic biofactory?. *Phytochemistry*, *69*(7), pp.1469-1495.

Pendleton, A.L., Smith, K.E., Feau, N., Martin, F.M., Grigoriev, I.V., Hamelin, R.,

Nelson, C.D., Burleigh, J.G. and Davis, J.M., 2014. Duplications and losses in gene families of

rust pathogens highlight putative effectors. *Frontiers in plant science*, 5, p.299.

Peypoux, F., Bonmatin, J.M. and Wallach, J., 1999. Recent trends in the biochemistry of

surfactin. *Applied Microbiology and Biotechnology*, 51(5), pp.553-563.

Pusztahelyi, T., Holb, I.J. and Pócsi, I., 2015. Secondary metabolites in fungus-plant

interactions. *Frontiers in Plant Science*, *6*, p.573.

Quandt, C.A., Beaudet, D., Corsaro, D., Walochnik, J., Michel, R., Corradi, N. and

James, T.Y., 2017. The genome of an intranuclear parasite, Paramicrosporidium saccamoebae,

reveals alternative adaptations to obligate intracellular parasitism. *Elife*, *6*, p.e29594.

Reynolds, H.T., Vijayakumar, V., Gluck☐Thaler, E., Korotkin, H.B., Matheny, P.B. and

Slot, J.C., 2018. Horizontal gene cluster transfer increased hallucinogenic mushroom diversity.

Evolution Letters, *2*(2), pp.88-101.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. 2010. edgeR: a Bioconductor

package for differential expression analysis of digital gene expression data. *Bioinformatics* 26,

139-140

Rokas, A., Mead, M. E., Steenwyk, J. L., Raja, H. A., & Oberlies, N. H. 2020.

Biosynthetic gene clusters and the evolution of fungal chemodiversity. *Natural Product Reports*

17, pp. 167-178. http://doi.org/10.1039/c9np00045c

Riley, R., Salamov, A.A., Brown, D.W., Nagy, L.G., Floudas, D., Held, B.W., Levasseur,

A., Lombard, V., Morin, E., Otillar, R. and Lindquist, E.A., 2014. Extensive sampling of

basidiomycete genomes demonstrates inadequacy of the white-rot/brown-rot paradigm for wood

decay fungi. *Proceedings of the National Academy of Sciences*, 111(27), pp.9923-9928.

1006

Schwartze, V.U., Winter, S., Shelest, E., Marcet-Houben, M., Horn, F., Wehner, S.,

Linde, J., Valiante, V., Sammeth, M., Riege, K., Nowrousian, M., Kaerger, K., Jacobsen, I.D.,

Marz, M., Brakhage, A.A., Gabaldon, T., Bocker, S. and Voigt, K. 2014. Gene expansion shapes

genome architecture in the human pathogen Lichtheimia corymbifera: an evolutionary genomics

analysis in the ancient terrestrial mucorales (Mucoromycotina). *PLoS Genetics*, 10(8):e1004496.

doi: 10.1371/journal.pgen.1004496.

Smith, D.J., Burnham, M.K., Edwards, J., Earl, A.J. and Turner, G., 1990. Cloning and

heterologous expression of the penicillin biosynthetic gene cluster from Penicillium

chrysogenum. *Bio/technology*, *8*(1), p.39.

Spatafora, J.W., Chang, Y., Benny, G.L., Lazarus, K., Smith, M.E., Berbee, M.L.,

Bonito, G., Corradi, N., Grigoriev, I., Gryganskyi, A. and James, T.Y., 2016. A phylum-level

phylogenetic classification of zygomycete fungi based on genome-scale data. *Mycologia*, *108*(5),

pp.1028-1046.

Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. *Bioinformatics*, *30*(9), pp.1312-1313.

Team, R.C., 2018. R: A language and environment for statistical computing.

Teixeira, M.D.M., Moreno, L.F., Stielow, B.J., Muszewska, A., Hainaut, M., Gonzaga,

L., Abouelleil, A., Patané, J.S.L., Priest, M., Souza, R. and Young, S., 2017. Exploring the

genomic diversity of black yeasts and relatives (Chaetothyriales, Ascomycota). *Studies in*

mycology, 86, pp.1-28.

Terfehr, D., Dahlmann, T.A., Specht, T., Zadra, I., Kürnsteiner, H. and Kück, U., 2014.

Genome sequence and annotation of Acremonium chrysogenum, producer of the β-lactam

antibiotic cephalosporin C. *Genome Announcements*, 2(5), pp.e00948-14.

Tisserant, E., Malbreil, M., Kuo, A., Kohler, A., Symeonidi, A., Balestrini, R., Charron,

P., Duensing, N., Frei dit Frey, N., Gianinazzi-Pearson, V., Gilbert, L.B., Handa, Y., Herr, J.R.,

Hijri, M., Koul, R., Kawaguchi, M., Krajinski, F., Lammers, P.J., Masclaux, F.G., Murat, C.,

Morin, E., Ndikumana, S., Pagni, M., Petitpierre, D., Requena, N., Rosikiewicz, P., Riley, R.,

Saito, K., San Clemente, H., Shapiro, H., van Tuinen, D., Becard, G., Bonfante, P., Paszkowski,

U., Shachar-Hill, Y.Y., Tuskan, G.A., Young, J.P., Sanders, I.R., Henrissat, B., Rensing, S.A.,

Grigoriev, I.V., Corradi, N., Roux, C. and Martin, F. 2013. Genome of an arbuscular mycorrhizal

fungus provides insight into the oldest plant symbiosis. *Proceedings of the National Academy of*

Sciences, 10;110(50):20117-22. doi: 10.1073/pnas.1313452110.

Tuller, T. 2011. Codon bias, tRNA pools, and horizontal gene transfer. *Mobile Genetic*

Elements, 1(1), pp. 75-77, DOI: 10.4161/mge.1.1.15400

Uehling, J., Gryganskyi, A., Hameed, K., Tschaplinski, T., Misztal, PK., Wu S., Desiro,

A., Vande Pol, N., Du, Z., Zienkiewicz, A., Zienkiewicz, K., Morin, E., Tisserant, E., Splivallo,

R., Hainaut, M., Henrissat, B., Ohm, R., Kuo, A., Yan, J., Lipzen, A., Nolan, M., LaButti, K.,

Barry, K., Goldstein, A.H., Labbe, J., Schadt, C., Tuskan, G., Grigoriev, I., Martin, F., Vilgalys,

R. and Bonito, G. 2017. Comparative genomics of Mortierella elongata and its bacterial

endosymbiont Mycoavidus cysteinexigens. 2017. *Environmental Microbiology,* 19(8):2964-

2983. doi: 10.1111/1462-2920.13669

Vesth, T.C., Nybo, J.L., Theobald, S., Frisvad, J.C., Larsen, T.O., Nielsen, K.F., Hoof,

J.B., Brandl, J., Salamov, A., Riley, R. and Gladden, J.M., 2018. Investigation of inter-and

intraspecies variation through genome sequencing of Aspergillus section Nigri. *Nature Genetics*,

50(12), pp.1688-1695.

Voigt, K., Wolf, T., Ochsenreiter, K., Nagy, G., Kaerger, K., Shelest, E. and Papp, T.,

2016. 15 Genetic and Metabolic Aspects of Primary and Secondary Metabolism of the

Zygomycetes. In *Biochemistry and Molecular Biology* (pp. 361-385). Springer, Cham.

Wisecaver, J.H. and Rokas, A., 2015. Fungal metabolic gene clusters—caravans traveling

across genomes and environments. *Frontiers in Microbiology*, 6, p.161.

Wang, D., Wu, R., Xu, Y. and Li, M. 2013. Draft Genome Sequence of Rhizopus

chinensis CCTCCM201021., Used for Brewing Traditional Chinese Alcoholic Beverages.

Genome Announcements, 1(2):e0019512. doi: 10.1128/genomeA.00195-12.

Wang, Y., White, M.M., Kvist, S. and Moncalvo, J.M. 2016. Genome-Wide Survey of

Gut Fungi (Harpellales) Reveals the First Horizontally Transferred Ubiquitin Gene from a

Mosquito Host. *Molecular Biology and Evolution*, 33(10):2544-54. doi:

10.1093/molbev/msw126.

Wang, Y., Youssef, N., Couger, M.B., Hanafy, R., Elshahed, M. and Stajich, J.E., 2018.

Comparative genomics and divergence time estimation of the anaerobic fungi in herbivorous

mammals. *bioRxiv*, p.401869.

Wang, Y.Y., Liu, B., Zhang, X.Y., Zhou, Q.M., Zhang, T., Li, H., Yu, Y.F., Zhang, X.L.,

Hao, X.Y., Wang, M. and Wang, L., 2014. Genome characteristics reveal the impact of

lichenization on lichen-forming fungus Endocarpon pusillum Hedwig (Verrucariales,

Ascomycota). *BMC Genomics*, 15(1), p.34.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New

York.

Winter, D.J., 2017. rentrez: An R package for the NCBI eUtils API (No. e3179v1). *PeerJ*

Preprints.

Xiao, J.H. and Zhong, J.J., 2007. Secondary metabolites from Cordyceps species and

their antitumor activity studies. *Recent Patents on Biotechnology*, *1*(2), pp.123-137.

Xiao, G., Ying, S.H., Zheng, P., Wang, Z.L., Zhang, S., Xie, X.Q., Shang, Y., Leger,

R.J.S., Zhao, G.P., Wang, C. and Feng, M.G., 2012. Genomic perspectives on the evolution of

fungal entomopathogenicity in Beauveria bassiana. *Scientific reports*, 2, p.483.

Yang, J., Wang, L., Ji, X., Feng, Y., Li, X., Zou, C., Xu, J., Ren, Y., Mi, Q., Wu, J. and

Liu, S., 2011. Genomic and proteomic analyses of the fungus Arthrobotrys oligospora provide

insights into nematode-trap formation. *PLoS pathogens,* 7(9).

Zhang, X.Y., Bao, J., Wang, G.H., He, F., Xu, X.Y. and Qi, S.H., 2012. Diversity and

antimicrobial activity of culturable fungi isolated from six species of the South China Sea

gorgonians. *Microbial Ecology*, *64*(3), pp.617-627.

Zuccaro, A., Lahrmann, U., Gueldener, U., Langen, G., Pfiffi, S., Biedenkopf, D., Wong,

P., Samans, B., Grimm, C., Basiewicz, M. and Murat, C., 2011. Endophytic life strategies

decoded by genome and transcriptome analyses of the mutualistic root symbiont Piriformospora

indica. *PLoS pathogens*, 7(10).

1090

1091

Tables

1093

Table 1. Predicted secondary metabolite (SM) core genes for *Basidiobolus* isolates used in this
study. NRPS: Non-ribosomal peptide synthetases, PKS: Polyketide synthases, TC: Terpene
cyclases.

1097

Isolate	Total SM	NRPS/NRPS-like	PKS/PKS-Like	NRPS-PKS hybrids	TC
B. meristosporus CBS 931.73	44	30	4	0	10
B. meristosporus B9252	43	29	7	1	6
B. heterosporus B8920	23	18	1	0	4

1098
1099
1100 **Table 2.** Number of predicted secondary metabolite core genes with evidence for HGI in
Basidiobolus genomes. NRPS: Non-ribosomal peptide synthetases, PKS: Polyketide synthases,
TC: Terpene cyclases.

1103

Isolate	Total SM	NRPS/NRPS-like	PKS/PKS-Like	NRPS-PKS hybrids	TC
B. meristosporus CBS 931.73	26/44 (61%)	22/30 (60%)	2/4 (50%)	0/0 (0%)	2/10 (2%)
B. meristosporus B9252	19/43 (44%)	16/29 (55%)	1/7 (14%)	0/1 (0%)	2/6 (1.4%)
B. heterosporus B8920	12/23 (52%)	12/18 (66%)	0/1 (0%)	0/0 (0%)	0/4 (0%)

1104
1105

1106
1107   **Table 3.** Summary of secondary metabolite core genes with HGT evidence from *Basidiobolus*
1108   isolates.
1109
1110

Isolate	Gene	Z-score of gene coverage	SM class	Taxonomy for best NCBI hit
	N168_02885	-0.104	NRPS	b-proteobacteria
	N168_00992	0.028	NRPS	enterobacteria
	N168_05934	-0.183	NRPS	b-proteobacteria
	N168_07140	0.260	NRPS	b-proteobacteria
	N168_06479	0.328	NRPS	b-proteobacteria
	N168_00176	0.013	NRPS	cyanobacteria
	N168_08721	0.561	NRPS	cyanobacteria
B. heterosporus	N168_05966	0.672	NRPS	d-proteobacteria
	N168_05324	-0.038	NRPS-Like	cyanobacteria
	N168_08580	0.047	NRPS-Like	d-proteobacteria
	N168_03036	0.022	NRPS-Like	d-proteobacteria
	N168_04239	0.070	NRPS-Like	b-proteobacteria
	N161.mRNA.1431.1	-0.350	NRPS	cyanobacteria
	N161.mRNA.1485.1	-0.258	NRPS	d-proteobacteria
	N161.mRNA.2152.1	0.074	NRPS	cyanobacteria
	N161.mRNA.4115.1	0.158	NRPS	enterobacteria
	N161.mRNA.4211.1	0.243	NRPS	cyanobacteria
	N161.mRNA.4324.1	0.213	NRPS	CFB group bacteria
	N161.mRNA.8304.1	0.358	NRPS	b-proteobacteria
	N161.mRNA.9145.1	0.201	NRPS	cyanobacteria
	N161.mRNA.9317.1	0.677	NRPS	d-proteobacteria
	N161.mRNA.9639.1	-0.151	NRPS	d-proteobacteria
	N161.mRNA.11289.1	0.596	NRPS	cyanobacteria
B. meristosporus B9252	N161.mRNA.1486.1	0.662	NRPS-Like	firmicutes
	N161.mRNA.5862.1	1.131	NRPS-Like	firmicutes
	N161.mRNA.6846.1	0.760	NRPS-Like	firmicutes
	N161.mRNA.6935.1	0.624	NRPS-Like	d-proteobacteria
	N161.mRNA.8699.1	-0.201	NRPS-Like	firmicutes
	N161.mRNA.6146.1	0.999	PKS	high GC Gram+
	N161.mRNA.1413.1	0.481	Terpene	CFB group bacteria
	N161.mRNA.13969.1	1.099	Terpene	CFB group bacteria

	366903	0.043	NRPS	a-proteobacteria
	368581	-0.232	NRPS	b-proteobacteria
	349800	0.208	NRPS	high GC Gram+
	351909	-0.120	NRPS	cyanobacteria
	372749	-0.088	NRPS	firmicutes
	372991	-0.260	NRPS	b-proteobacteria
	373247	0.533	NRPS	firmicutes
	373940	-0.142	NRPS	firmicutes
	375475	0.522	NRPS	enterobacteria
	338397	0.033	NRPS	d-proteobacteria
	375580	0.033	NRPS	cyanobacteria
	375613	-0.369	NRPS	firmicutes
	377413	-0.245	NRPS	g-proteobacteria
	387529	-0.157	NRPS	b-proteobacteria
	307892	0.404	NRPS	a-proteobacteria
B. meristosporus CBS 931.73	298977	-0.201	NRPS	b-proteobacteria
	343011	0.147	NRPS	firmicutes
	363930	-0.358	NRPS	firmicutes
	300898	-0.165	NRPS-Like	firmicutes
	322666	-0.174	NRPS-Like	CFB group bacteria
	146993	-0.200	NRPS-Like	cyanobacteria
	382467	-0.264	NRPS-Like	a-proteobacteria
	340613	-0.136	NRPS-Like	GNS bacteria
	237744	-0.001	PKS	bacteria
	292783	-0.429	PKS	high GC Gram+
	301341	-0.210	Terpene	CFB group bacteria
	304520	0.389	Terpene	cyanobacteria

Figure 1. Genome size, number of genes, proportion of secondary metabolism (SM) gene clusters and number of predicted SM gene clusters for 66 Zoopagomycota and Mucoromycota sequenced genomes. Color code represents the category of SM predicted per isolate. NRPS: Non-ribosomal peptide synthetases. PKS: Polyketide synthases. Glom.: Glomeromycotina. Mort.: Mortierellomycotina. Ent.: Entomophtoromycotina. Kickx.: Kickxellomycotina. Zoo.: Zoopagomycotina

Figure 2. Distribution of number of RNAseq counts in transcripts per million (TPM) per genic feature from *B. meristosporus* CBS 931.73. Colors represent SM and genes of interest. X-axis represents the TPM count. Y-axis represents distribution of TPM. The scale is in log(TPM), and the values are in TPM absolute values. Histogram represents the distribution of mapped reads in TPM for all predicted gene models with non-zero TPM values across the *B. meristosporus* CBS 931.73 genome.

Figure 3. Phylogenetic sources and abundance of A-domains from NRPS predicted gene models for Zoopagomycota and Mucoromycota species. The maximum likelihood phylogenetic tree (left) represent a simplification of the reconstructed tree which includes the clades that include more than one A-domain from Mucoromycota/Zoopagomycota genomes. The heatmap (right) represents the abundances A-domains predicted for each domain clustered within each clade. *Basidiobolus* species are enriched in NRPS from the major bacterial clade (MBC), cyclosporin (CYCLO), surfactin-like, and siderophore (SIDE, SID) clades.

Figure 4. Graphical representation of the A-domains of each NRPS core gene predicted for *Basidiobolus* genomes. Horizontal grey lines represent the length of the predicted NRPS core gene. Arrows represent A-domains and are located in the position within the gene model. Colors represent the phylogenetic origin. Numbers represent the name of each gene model and predicted SM cluster for each genome.

Figure 5. Phylogenetic sources and abundance of terpene cyclase (TC) core genes predicted for Zoopagomycota and Mucoromycota species. The maximum likelihood phylogenetic tree is a simplification of the maximum likelihood tree reconstructed using the TC core genes. The heatmap (right) represents the abundances of predicted TC genes within each reconstructed clade. Glom.: Glomeromycotina. Mort.: Mortierellomycotina. Ent.: Entomophtoromycotina. Kickx.: Kickxellomycotina. Zoo.: Zoopagomycotina

Figure 6. Plausible taxonomic sources of HGT genes. Bar-plot represents the proportion of diversity of HGT candidates for each *Basidiobolus* genome. Colors represent the taxonomy term for the RefSeq best hit from BLAST. Overall, between 3% to 5% of the gene models predicted for Basidiobolus species appear to be product of HGT from taxonomic groups of bacteria associated to reptilian and amphibian gut tracts (Proteobacteria, firmicutes, and CFB/bacterioidetes).

Figure 7. Siderophore activity of *Basidiobolus meristosporus* CBS 931.73 in a universal CAS assay using layered AY-CAS plates after 12 days. Bars represent mean siderophore activity measured per strain as the yellow area in AY-CAS plates for three replicates. Error bars represent the standard deviation for each replicate. *Cladosporium sp.* PE-07 represents the positive control. *Conidiobolus thromboides* FSU 785 represents a zygomycete with no evidence for sidero-phore NRPS expansion.