Title: **Neural harmonics reflect grammaticality**

Authors: Alessandro Tavano[1][*], Stefan Blohm[2], Christine Knoop[2], R Muralikrishnan[1], Mathias Scharinger[2,3], Valentin Wagner[2], Dominik Thiele[1], Oded Ghitza[4], Nai Ding[5], Winfried Menninghaus[2], David Poeppel[1,6]

Affiliations:

1. Department of Neuroscience, Max Plank Institute for Empirical Aesthetics, 60322 - Frankfurt am Main, Germany
2. Department of Language and Literature, Max Plank Institute for Empirical Aesthetics, 60322 - Frankfurt am Main, Germany
3. University of Marburg, 35037 – Marburg, Germany
4. Department of Biomedical Engineering, Boston University, 02215 - Boston, USA
5. College of Biomedical Engineering and Instrument Sciences, Zhejiang University, 310027 - Hangzhou, Zhejiang Province, China
6. Department of Psychology, New York University, 10003 - New York City, New York, USA

* Corresponding author: alessandro.tavano@ae.mpg.de

Abstract

Can neural activity reveal syntactic structure building processes and their violations? To verify this, we recorded electroencephalographic and behavioral data as participants discriminated concatenated isochronous sentence chains containing only grammatical sentences (regular trials) from those containing ungrammatical sentences (irregular trials). We found that the repetition of abstract syntactic categories generates a harmonic structure of their period independently of stimulus rate, thereby separating endogenous from exogenous neural rhythms. Behavioral analyses confirmed this dissociation. Internal neural harmonics extracted from regular trials predicted participants' grammatical sensitivity better than harmonics extracted from irregular trials, suggesting a direct reflection of grammatical sensitivity. Instead, entraining to external stimulus rate scaled with task sensitivity only when extracted from irregular trials, reflecting attention-capture processing. Neural harmonics to repeated syntactic categories constitute the first behaviorally relevant, purely internal index of syntactic competence.

Keywords: Entrainment, brain rhythms, harmonics, syntax, syntactic competence, Grammaticality

Introduction

Speech and language are related but distinct dimensions of verbal communication [1–4]. While temporal regularities guide speech chunking in ways that are relevant for language production and comprehension [3, 5–13]), the core of language processing lies in the internal and tacit knowledge of syntactic rules, whose function is to determine how words combine into meaningful utterances [14]. Isolating a direct

neural correlate of such linguistic competence has proven an extremely difficult task.

A growing body of literature has linked brain rhythms to various aspects of language

comprehension. Recently, Ding and colleagues [15] tried to capture a neural

signature of structure building in the frequency domain by having native speakers of

English and Mandarin Chinese listen to continuous spoken sentences, concatenated

as continuous trains of monosyllabic words presented at a fixed rate of 4 Hz. They

tested sentences composed of two-word noun phrases (NP = a phrase headed by a

noun, such as "My shoe") followed by two-word verb phrases (VP = "is wet", see

Figure 1a). The lexical input therefore appeared at a rate of 4 Hz, phrasal units at 2

Hz, and sentences at 1 Hz. This design was selected to allow for 'frequency tagging'

of sentence building processes, such that the neural data could be decomposed into

linguistically relevant rhythmic response components. Remarkably, a Fast Fourier

Transform (FFT) analysis highlighted significant peaks of spectral energy emerging

not only at 4 Hz (word rate), but also at 2 Hz and 1 Hz. The authors deemed the

neural responses as a direct window into the linguistic hierarchy: 2 Hz rhythms (2-

word units) tracked the phrases, and 1 Hz rhythms (4-word units) tracked the

sentences. Such linguistic rhythms emerged only if participants listened to their own

language [16], and have been further interpreted by some as tracking the

incremental combination of words into phrases, and phrases into sentences [17].

The findings have led to a number of follow-up studies, across languages.

The appeal of the interpretation notwithstanding, linguistic considerations question

the isomorphic mapping of syntactic structure and brain rhythms. If a frequency

analysis (e.g., an FFT) simply captures the rhythmic repetition of abstract linguistic

categories, is it plausible that the spectral peak at 2 Hz reflects individual phrase

processing? In fact, in such a frequency tagging design, each of the phrasal nodes -

NP and VP - repeats every second, not every half a second, so the repetition of a

phrasal node, in and by itself, cannot generate a 2 Hz rhythm. For the 2 Hz phrasal

rhythm interpretation to hold, one would have to hypothesize that: a) the human

brain is insensitive to the difference between noun phrases and verb phrases; b)

and, conversely, it is sensitive to a superordinate phrase category (which we term

sPhrase), cycling periodically through all NPs and VPs (see Figures 1a,d, and 1b,e).

Since in fact the human brain distinguishes between abstract phrasal categories [18–

23], we suggest a simpler alternative model, based on categorical repetition, which

does not require a direct mapping of structure building onto individual rhythms. We

assume phrasal and sentence nodes repeat with the same period, so they jointly

contribute to the same categorical rhythm: 1 Hz, in the example at hand. The 2 Hz

peak would then be generated simply as a first harmonic consequence (see Figures

1a,d, and 1c,f). To test the 'sPhrase response' model against the 'categorical

harmonics' model by varying the size of phrasal units within sentences (see Figure

1a,d), ensuring that the period of phrases was not a multiple of sentence period, so

as to be able to detect *sPhrase* rhythms in an FFT analysis. We recorded

encephalographic (EEG) and behavioral data while 31 participants discriminated

isochronous trials of ten sentences, half of them containing only grammatical

sentences (*regular trials*), and half containing two randomly distributed

ungrammatical sentences (*irregular trials* as defined by word order violations, see

Figure 2).

The results are not consistent with the existence of a superordinate *sPhrase* rhythm,

suggesting instead that the brain tracks the repetition of abstract phrasal categories -

NPs, VPs and sentence nodes -, into a single neural rhythm, which generates its

own harmonic structure. The higher harmonics are mathematically independent from

stimulus (word) rate, leading to a clear-cut distinction between internally constructed

versus externally driven rhythms. Importantly, the median power of harmonics

structures for regular trials was larger than that for irregular trials, suggesting that

brain rhythms encode the distinction between grammatical and ungrammatical

sentences, i.e. grammaticality. This was confirmed by the emergence of a functional

double dissociation: (a) entraining to external stimuli was at ceiling for regular trials

but scaled with task sensitivity for irregular trials, likely indexing lower-level attentive

processes [24]; (b) neural harmonics extracted from regular trials better predicted

individual grammaticality judgments than those extracted from irregular trials, linking

for the first time brain rhythms to internalized linguistic knowledge governing

behavioral decisions.

Results

Behavior. Monosyllabic stimuli formed sentences of either four (Conditions 1 and 2,

see Figure 2a) or five words (Conditions 3 and 4, see Figure 2b). Disyllabic stimuli

formed sentences of five words (Conditions 5 and 6, Figure 2c). For each condition,

participants distinguished between trials containing solely grammatical sentences

(regular trials), or trials in which two neighboring sentences were rendered

ungrammatical by randomly shuffling word order (irregular trials).

Estimates of task sensitivity and response bias were obtained using a signal

detection approach [25]. Discriminability index (d') measures of task sensitivity,

showed that in all conditions participants successfully distinguished irregular from

regular trials, suggesting that the grammatical nature of the task was well

understood: all $ts_{(30)} \geq 7.94$, all $ps \leq 7.205^{-09}$, FDR-corrected (threshold p = 0.008.

See Figure 3, left panel). There was a significant difference among conditions:

$F_{(30,185)}$ = 4.16, p = 0.001, η^2 = 0.04. When FDR-corrected (threshold p = 0.006), only

a difference between condition 1 and condition 4 survived, $t_{(30)}$ = -3.34, p = 0.001,

suggesting that sensitivity for condition 1 (mean = 1.83, SD = 1.24) was lower than

for condition 4 (mean = 2.67, SD = 1.75). Response bias measures showed that

participants adopted a stricter decision criterion in distinguishing between regular

and irregular trials for conditions 1, 4, 5 and 6: all $ts_{(30)} \geq 3.65$, all ps $\leq 9.749^{-04}$, FDR-

corrected (threshold p = 0.021. See Figure 3, right panel). This means that they were

more likely to classify an irregular trial as regular. This suggests that the

impoverished acoustic quality of stimuli following the pitch-flattening procedure did

not drive participants' grammatical judgments. No significant difference in criterion

estimates among conditions was found, suggesting that the stimulus generation

procedure was bias-free: $F_{(30,185)}$ = 1.81, p = 0.115.

Entrainment to word rate. To avoid low-level confounds on syntactic chunking, we

first asked if speech sound statistics acted as implicit cues to phrasal boundaries.

There emerged no significant difference between phonemic transition probabilities

within and between phrases (see Supplemental Information, Figure S1a,b). We

found significant differences in word frequency (Figure S1c), reflecting the known

asymmetry in the distribution of monosyllabic and disyllabic words in natural

languages (for the German language, see [26]).

As for power peaks, we then used the same analysis approach for both acoustic

stimulus sequences and neural data, in order to make their results directly

comparable: A Fast Fourier Transform analysis was run on individual trials,

normalized power estimates were extracted using the complex modulus, and signal-

to-noise ratio estimates (SNR) were calculated for each subject and condition. This

approach highlights the concentration of auditory or neural activity across

frequencies (See Materials and Methods). Acoustic data showed a significant word

rate entrainment effect: all $ts_{(30)} \geq 55.04$, all $ps \leq 1.080^{-31}$, FDR-corrected (threshold p

= 0.001, Figure S2a,b,c). Neural frequency tagging effects in the EEG data were

analyzed separately for regular and irregular trials. We compared activity at target

frequencies to their own floor noise level, defined as the average of the samples two

positions to the left and right side of each target frequency. For regular trials, we

found significant peaks of spectral energy at word rhythm in all conditions, thereby

verifying that the brain robustly entrained to both monosyllabic and disyllabic

rhythms: all $ts_{(30)} \geq 7.97$, all $ps \leq 6.648^{-09}$, FDR-corrected (threshold p = 0.008). The

same held for irregular trials: all $ts_{(30)} \geq 5.95$, all $ps \leq 1.564^{-06}$, FDR-corrected (See

Figures 4, 5, and S3). We then tested the effect of grammaticality (irregular vs.

regular trials) and internal syntactic structure (Conditions 1 vs. 2, with four words

running at 4 Hz; conditions 3 vs. 4, with five words running at 4 Hz; conditions 5 vs.

6, with five words running at 2 Hz) in a series of two-way rmANOVAs, but found no

significant main effect or interaction: all $Fs_{(1,30)} \leq 6.35$, all $ps \geq 0.017$, FDR corrected

(threshold p = 0.0083). Hence, neural entrainment to word rate did not reflect the

judgments of grammaticality required to distinguish regular from irregular trials.

Internal linguistic rhythms. Next, we focused on the evidence for *sPhrase* vs

categorical models. First, there was no evidence for linguistic rhythms in the auditory

signal (see Figure S2, Supplemental Information). Second, we hypothesized that

either significant *sPhrase* rhythms emerge across conditions, or a robust harmonic

structure emerges across conditions. We tested these hypotheses on regular trial

data. Condition 1 replicates the main condition in Ding et al., 2016). As for *sPhrase*

rhythms, condition 1 is effectively uninformative, as a putative *sPhrase* rhythm would

overlap with the first harmonic to sentence rhythm. For condition 2, a *sPhrase*

rhythm for the VP should have had a peak at 1.33 Hz, but we could not detect it: $t_{(30)}$

= -1.08, p = 0.28 (Figure 4a). For conditions 3 and 4, which mirror each other in the

size of NPs and VPs, *sPhrase* rhythms should have emerged at 1.33 Hz for three-

word phrasal units, and 2 Hz for two-word phrasal units (Figure 2b). Again, no

evidence could be found for those rhythms being different from floor noise: all $ts_{(30)} \leq$

-1.71, all ps $\geq$ 0.09 (See Figure 4b). Finally, for conditions 5 and 6, two *sPhrase*

rhythms should emerge: 0.66 Hz for three-word units, and 1 Hz for two-word units

(Figure 2c). Again and conclusively, no evidence supporting their emergence from

noise could be found: all ts(30) $\leq$ 1.29, all ps $\geq$ 0.20 (See Figure 4c). Conversely, we

found strong evidence for the presence of categorical rhythms, contributed by both

sentence and phrasal nodes, and their harmonic structures. Categorical rhythms

were predicted at 1 Hz for conditions 1 and 2, 0.8 Hz for conditions 3 and 4, and 0.4

for conditions 5 and 6. Significant peaks for abstract category rhythms were found in

all conditions relative to floor noise, except condition 6: all $ts_{(30)} \geq$ 4.03, all ps $\leq 3.471^{-}$

^{04}, FDR-corrected (threshold p = 0.015). Condition 6 approached significance: $t_{(30)}$ =

1.88, p = 0.068. The first harmonic process was significant in all conditions: all $ts_{(30)} \geq$

2.57, all ps $\leq$ 0.015, FDR-corrected. The second harmonic process was significant

in all conditions: all $ts_{(30)} \geq$ 2.56, all ps $\leq$ 0.015, FDR-corrected. Importantly, our

experimental design for conditions 3 to 6 ensure that first and second harmonic

processes cannot be mathematically derived as subharmonics of stimulus rate [27]:

for example, in conditions 3 and 4 there exists no integer factor which when

multiplied by the first harmonic at 1.6 Hz would yield the 4 Hz word rate. This simple

fact grants complete functional independence to harmonic structures, suggesting the

existence of purely endogenous neural generators.

Grammaticality in the frequency domain. We then turned to irregular trials, in order to

assess the extent to which internalized language knowledge affects frequency

domain responses. Category rhythm peaks were largely suppressed when

participants attended to irregular trials: We found significant activity only in conditions

2, 3 and 4: all $ts_{(30)} \geq 2.68$, all $ps \leq 0.011$, FDR-corrected (threshold $p = 0.017$. Figure

5abc, left panel). Conditions 5 and 6 approached significance: $ts_{(30)} = 3.33$ and 1.91,

$ps = 0.017$ and 0.065, respectively. Importantly, no significant harmonic process was

found in any experimental condition: all $ts_{(30)} \leq 2.76$, all $ps \geq 0.009$, FDR-corrected

(threshold $p = 0.008$. See Figure 5abc, left panel). We directly compared regular

against irregular trials in an rmANOVA with factors Grammaticality (regular vs.

irregular) and Condition for each harmonic structure peak (all comparisons FDR-

corrected). There was no significant Grammaticality x Condition interaction (all

$Fs_{(1,30)} \leq 4.31$, all $ps \geq 0.0465$, FDR-corrected with threshold $p = 0.005$). There was a

significant Condition effect for the first harmonic of conditions 3 and 4: $F_{(1,30)} = 9.25$,

$p = 0.004$, $\eta^2 = 0.28$. Activity for condition 3 (mean = 1.20 μV^2, SD = 0.30) was

reduced relative to condition 4 (mean = 1.48 μV^2, SD = 0.44). More importantly, we

found a significant Grammaticality effect for all category, first harmonic and second

harmonic processes (all $Fs_{(1,30)} \geq 5.55$, all $ps \leq 0.025$, FDR-corrected with threshold

$p = 0.025$), except for category peaks in conditions 5 and 6, and first harmonics in

conditions 3 and 4 (Figure 5abc, right panel). This suggests that higher harmonics

encode perceived grammaticality more reliably than lower ones. More generally, we

conclude that task sensitivity, driven by perceived grammaticality, modulates the

strength of neural harmonic response.

A functional double dissociation in the frequency domain. An important test of the

function of neural responses in the frequency domain is the extent to which they

predict behavior. We created two simple neural predictors: an internal one, by

calculating for each participant the median power across harmonics and conditions,

separately for regular and irregular trials; and an external one, by averaging the

strength (in SNR) to which each participant entrained to external stimulus rate across

conditions, again separately for regular and irregular trials. We then correlated each

neural predictor with standardized task sensitivity values, averaged across

conditions. A functionally relevant double dissociation emerged. As for the internal

index, power estimates extracted from both regular and irregular trials significantly

predicted participants' ability to decide whether a trial was regular or irregular:

regular trials, $r_{(29)} = 0.69$, $p < 0.001$; irregular trials, $r_{(29)} = 0.42$, $p = 0.017$. A

comparison between correlation coefficients using the Steiger test [28] determined

that the fit for regular trials was superior to that for irregular trials: $Z = 1.77$, $p = 0.038$

(Figure 6a). This suggests that the extraction of abstract category rhythms was

boosted when linguistic knowledge was employed for grammaticality judgments. As

for the external index, we previously showed that entrainment to word rates did not

differ between conditions or by grammaticality (Figure S3). However, power

estimates extracted from irregular trials predicted behavior, with higher power

indicating increased attention to irregular stimuli: $r(29) = 0.40$, $p = 0.022$. Locking to

stimulus rate within regular trials was at ceiling regardless of behavioral

performance, confirming that entrainment per se does not reflect internal language

knowledge: $r(29) = 0.02$, $p = 0.89$ (Figure 6b). A Steiger test comparison confirmed

this conclusion: $Z = 1.95$, $p = 0.025$.

Discussion

We show that the perceived repetition, at constant rates, of linguistic categories

generates a harmonic structure of categorical rhythms, which directly reflects the

dynamics of syntactic structure building, and its violations. We varied the syntactic

complexity of phrases across six conditions, and demonstrated that linguistic

hierarchy, i.e. the assumption that words combine into phrases and phrases combine

to form sentences, is not isomorphically reflected in the hierarchy of energy peaks in

the frequency domain [15]. Rather, the iso-periodic repetition of nominal and verbal

phrases, as well as sentences, feeds into one categorical peak reflecting categorical

repetition, which in turn generates its own harmonic structure and correlates with

task sensitivity. Intriguingly, the categorical model closely reflects certain syntactic

theories which conceive of sentence level simply as a phrasal node, reflecting a

functional category on a pair with other phrases [29]. More broadly, because

frequency tagging is so widely used as a design, these results provide insights into

correctly interpreting the cognitive/linguistic mechanisms that underwrite rhythmic

brain activity.

Behavioral analyses document that participants not only were generally highly

proficient in distinguishing regular from irregular trials but that they also took a

conservative stance in deeming a trial as irregular, suggesting awareness about their

own grammaticality judgments. Importantly, our experimental manipulations

document that the harmonic structures generated by the category-driven syntactic

rhythms are independent from mere entrainment to external stimulus rate. For

example, in conditions 3 and 4, the first and second harmonics at 1.6 and 2.4 Hz

cannot be derived as sub-harmonic processes of the 4 hz word rate. Furthermore,

we document a double dissociation between internally generated and externally

elicited rhythms. The median power of internal harmonics extracted from regular

trials predicts grammatical judgments (i.e., task sensitivity) better than power

extracted from irregular trials. However, the median power of word rate extracted

from regular trials is at ceiling regardless of task sensitivity, suggesting that the two

measures are unrelated. Word rates extracted from irregular trials correlated with

task sensitivity. We interpret these findings as evidence that category-driven

harmonic structures reflect internal, top-down generators of syntactic knowledge,

whereas the strength of entrainment to word sequences mediates the deployment of

task-relevant attention [24], with more attention to stimuli leading to better detection

of irregular trials.

From a functional viewpoint, the linguistic representations and computations that can

be captured by our experimental setup cannot provide decisive evidence for

incremental buildup processes combining words into phrases, and phrases into

sentences [17], or for lexical buildup processes [30]. This is because the brain

rhythms - resulting from an FFT analysis - do not depict the temporal unfolding of

linguistic structure; rather, they depict the end result of linguistic parsing, i.e. the

effect of the repeated recognition of phrasal and sentential nodes within trials in the

form of a harmonic structure. The findings indicate that the strength of harmonic

structures predicts the degree to which native German speakers are sensitive to

word order violations, i.e., to the grammaticality manipulation we selected for our

experiment. The online comprehension of German relies substantially on word order

within and across phrases [31]. However, within the Germanic language family,

English, sensibly, assigns word order more weight for comprehension than German does [32]. We therefore predict that the sensitivity to word order violations, as reflected by the strength of neural harmonics, should be larger for languages which strongly depend on word order for comprehension, such as English, and smaller for morphologically richer languages, such as Italian, in which comprehension is locally bound by rules of declension. This perspective opens new, exciting avenues of cross-linguistic research for testing different types of grammaticality constraints in the frequency domain.

The data functionally dissociate between internal and external rhythmic neural modulations within and below the delta band range (all categorical rhythms ≤ 1 Hz). Delta band (0.5-4 Hz) and infra-slow (< 0.5 Hz) oscillations are natural spectral niches for linguistic operations because their extended time windows match the ranges of spontaneous language segment duration [33, 34], allowing for the incremental integration of statistically driven linguistic information [35]. In congruence with this stance, slow oscillations have been found to convey top-down knowledge across brain areas [36]. However, delta rhythms also robustly entrain to salient events in continuous speech [37], as well as to meaningless word lists [15]. Hence, rhythmic activity in the delta and sub-delta bands in response to spoken language likely reflects a mixture of neural processes pertaining to attentive speech tracking as well language knowledge, possibly running in a concurrent and interactive fashion. To partition their contribution, linguistically balanced and controlled designs are of the essence.

Of note, the strength of harmonic processes (first, second and higher order) is predictable only up to a point [27]. For example, it is unclear from our data why some harmonic processes are larger than others (see the difference between conditions 3

322 and 4 as far as the first harmonic is concerned). In our view, what is missing at

323 present is a good theory of signal-to-noise distribution for harmonic process. This

324 also suggests caution in interpreting the lack of a significant peak as an argument

325 against a harmonic interpretation.

327 Conclusions

328 By using carefully designed linguistic structures, we conclusively distinsguished

329 internally constructed neural rhythms from externally neural rhythms induced via

330 entrainment. The latter likely convey differences in stimulus-driven attention capture,

331 while the former reflect the perception of regular and violated syntactic structure

332 rules.

334 Materials and methods

Participants. Thirty-two young adults participated in the Experiment (Age range: 19-

30, 8 males). The EEG data from one participant were not faithfully recorded, and

were thus discarded. All analyses were run on the remaining participants (N = 31):

they were all German native speakers, right-handed (Oldfield Test), and self-

reported normal hearing, normal-to-corrected vision, no medical history of treatments

affecting the central nervous system, and no psychiatric disturbances. They were

compensated with 10 euros per hour (~3 hours, including instruction, EEG montage

and cleaning). All experimental procedures were approved by the Ethics Committee

of the Max Planck Society, and were undertaken with a written informed consent

signed by each participant.

Stimuli. Stimuli were words composing different sentence types. To reduce implicit

word- and phrase-level prosodic cues, stimuli were individually generated using the

MacinTalk Text-to-Speech Synthesizer (OSX El Capitan, voice Anna, f0 set at 220

Hz), and then pitch-flattened to 220 Hz across all time points (fundamental frequency

estimated using an autocorrelation approach, Praat, v. 6.0.16, www.praat.org). If an

original stimulus was longer than the relative SOA, its duration was compressed to fit

the stimulation window while preserving pitch.

Individual trials were composed of ten sentences, randomly selected from predefined

lists of 50 sentences, one list per sentence type condition (see Procedure, and

Appendix). There were two types of trial: regular trials, which contained only

grammatically correct sentences; and irregular trials, which contained also two

ungrammatical sentences, obtained by randomly shuffling word positions across two

successive sentences. There were 50 trials per condition, half were regular and half

irregular, thereby yielding unbiased estimates of accuracy in discriminating regular

from irregular trials, a task akin to classic grammaticality judgments.

Procedure. Participants sat comfortably in a IAC 40a sound attenuating and

electrically shielded recording booth (IAC Acoustics), approximately 1m from an LCD

computer screen. They were instructed to fixate a cross at the center of the screen

and listen attentively to each trial; When a trial ended, they were asked to determine

whether it was regular or irregular. They were asked to be as accurate as possible,

with no time limit. Participants used their right hand to press on two pre-defined

buttons (arrow up regular, arrow down irregular) on a keyboard. Stimuli were

delivered diotically at 75 dBs SPL via loudspeakers positioned at circa 1.2 m from

participants, 25 cm to the left and right of the LCD screen. On presentation, stimuli

371 were further attenuated using a fixed -20 dB SPL step, resulting in a comfortable and

372 perceptually controlled environment. Brief rest periods between blocks were self-

373 determined by each participant.

374 There were six sentence type conditions. Trial presentation was blocked by sentence

375 type. Stimuli within a trial were presented at a constant Stimulus Onset Asynchrony

376 (isochronous SOA): 250 ms for monosyllabic words, 500 ms for disyllabic words. As

377 in Ding et al. [15], there was no gap between sentences within a trial: Participants

378 listened to a continuous flow of words equally spaced in time [38], further reducing

379 prosodic cues at phrasal and sentence level.

380 The main experimental manipulation pertained to the size - in number of words - of

381 Noun Phrases (NP) and Verb Phrases (VP) composing each sentence. An NP is a

382 phrase with a noun as head (e.g., "The girls"), while a VP has a verb as head (e.g.,

383 "play rugby"). NPs usually perform the grammatical functions of verb subject or verb

384 object. In our experiments, all grammatical sentences contained an NP functioning

385 as verb subject, and a VP, which either included a second NP functioning as a verb

386 object or a different phrasal component. The Appendix lists all token sentences. See

387 Figure 2 for an exemplary analysis of a sentence used in each condition. In the main

388 text, NP always refers to a Noun Phrase functioning as subject, and VP refers to the

389 whole Verb Phrase, regardless of its internal composition.

390 Stimulus sequences were created using custom scripts written in Matlab (R2015b,

391 64 bit, mathworks.com). Sequence delivery was controlled by Psychophysics

392 Toolbox Version 3 (PTB-3, psychtoolbox.org) for Matlab, running on a Windows 7

393 computer (ASIO sound card for optimal stimulus latency control).

Data Recording and Analysis. Behavioral accuracy data were subject to a signal

detection theory approach, obtaining measures of task sensitivity (d' = zscore(Hits)

minus zscore(False Alarms)) and response bias (criterion = -

0.5*(zscore(Hits)+zscore(False Alarms))) for each participant and condition [25].

To record EEG data, we used an actiCAP 64-channel, active electrode set (10-10

system, Brain Vision Recorder, Brain Products, brainproducts.com) to record

electroencephalographic (EEG) activity at a sampling rate of 1KHz, with a 0.1 Hz

online filter (12 dB/octave roll-off). Additionally, electrocardiographic (ECG) and

electrooculographic (EOG) signals were recorded using a standard bipolar montage.

All impedances were kept below 10 kOhm. Data were recorded with an on-line

reference to FCz channel, and offline re-referenced to the average activity of all

scalp channels, and downsampled to 250 Hz. Using the EEGLAB toolbox for Matlab

[39] (sccn.ucsd.edu), continuous EEG data were first visually inspected to remove

large non-stereotypical artifacts (e.g., sudden head movements, chewing), digitally

filtered at 35 Hz low-pass (kaiser window, beta = 5.65326, filter order 93, transition

bandwidth 10 Hz), high-pass filtered at 1 Hz (filter order 455, transition bandwidth 2

Hz) to ensure data stationarity, submitted to an automatic artifact rejection based on

spectrum thresholding (threshold = 10 standard deviations, 1-35 Hz), and then

decomposed into Independent spatial components using the Infomax algorithm,

which allows for optimal source separation [40]. The resulting Independent

Components (ICs) were tested using the SASICA toolbox for EEGLAB [41]: ICs

reflecting blinks/vertical eye movements, lateral eye movements and heart-beat,

were detected by means of a correlation threshold (0.7) with bipolar Vertical,

Horizontal EOG, and ECG channels, respectively, and found to be present in all

participants (range: 1-3 ICs for vertical eye movements, 1-2 ICs for horizontal, 1 IC

for heart beat). To reduce inter-trial signal variability, ICs reflecting muscle artifacts

were identified via autocorrelation (threshold = 0, lag = 20 s), while ICs reflecting

focal topography was used for bad electrodes (= 7 standard deviations threshold

relative to the mean across electrodes). ICA results were copied back to the original

EEG datasets (0.1 Hz high-pass), and ICs marked as artifactual were rejected before

trial epoching. Each ten-sentence trial varied in duration between 10 seconds (4

words per sentence, running at 4 Hz) and 25 seconds (5 words per sentence,

running at 2 Hz). Individual trial power estimates were extracted for each electrode

from Hann-windowed and standardized data using the complex modulus of the Fast

Fourier Transform, correcting for the Hann window loss of power (sqrt(1.5)). Signal-

to-noise ratio estimates (SNR) were calculated for each peak of interest using the

average of two samples before and two samples after.

For all hypothesized peaks of energy, we created a surrogate noise floor condition

by averaging the SNR values of two samples below and two samples above the

peak sample. Each floor condition was used as the term of comparison in a T-test to

verify the peak's significance. To verify the significance of syntactic manipulations,

repeated measures analysis of variance (rmANOVAs) were run on each condition

pair with the same stimulation rate and number of words in a sentence. False-

discovery-rate (FDR) correction with Q value = 0.05 was applied in all cases of

multiple comparisons [42].

Appendices

See Table 1

Acknowledgements

The authors would like to thank Cornelius Abel, Jana Gessert, Freya Materne,

Claudia Lehr, Alexander Lindau, Georg-Friederich Paasch for help with stimulus

creation and data collection, and Johanna Rimmele, Andrea Martin and

Nina Kazanina for critical feedback on data analysis.

References

1. Hickok G, Poeppel D. Dorsal and ventral streams: a framework for understanding aspects of the functional anatomy of language. Cognition 2004; 92: 67–99.

2. Hickok G, Poeppel D. The cortical organization of speech processing. Nat Rev Neurosci. 2007;8: 393–402.

3. Giraud AL, Poeppel D. Cortical oscillations and speech processing: emerging computational principles and operations. Nat Neurosci. 2012;15(4) :511–517.

4. Tavano A, Scharinger M. Prediction in speech and language processing. Cortex 2015; 68: 1–7.

5. Ding N, Patel AD, Chene L, Butler H, Luo C, Poeppel D. Temporal modulations in speech and music. Neurosci Biobehav Rev. 2017;81: 181–187.

6. Ghitza O, Greenberg S. On the possible role of brain rhythms in speech perception: intelligibility of time-compressed speech with periodic and aperiodic insertions of silence. Phonetica 2009;66: 11326.

7. Ghitza O. Acoustic-driven delta rhythms as prosodic markers. Lang Cogn Neurosci. 2017;32: 54561.

8. Assaneo F, Poeppel D. The coupling between auditory and motor cortices is rate-restricted: Evidence for an intrinsic speech-motor rhythm. Sci Adv. 2018;4(2): eaao3842.

9. Steinhauer K, Alter K, Friederici AD. Brain potentials indicate immediate use of prosodic cues in natural speech processing. Nat Neurosci. 1999;2: 191—196.

10. Hwang H, Steinhauer K. Phrase length matters: The interplay between implicit prosody and syntax in Korean garden path sentences. J Cogn Neurosci. 2017;23(11): 295– 300.

11. Tavano A, Pesarin A, Murino V, Cristani M. Automatic conversational scene

analysis. low-level acoustic feature classification of speech activity in children with

Asperger syndrome and typically developing peers. PLoS One 2014;9(1): e85819.

12. Scharinger M, Steinberg J, Tavano A. Integrating speech in time depends on

temporal expectancies and attention. Cortex 2017;93: 28–40.

13. Keitel A, Gross J, Kayser C. Perceptually relevant speech tracking in auditory

and motor cortex reflects distinct linguistic features. PLoS Biology 2018;16(3):

e2004473.

14. Chomsky N. Language architecture and its import for evolution. Neurosci

Biobehav Rev. 2017;81: 295-300.

15. Ding N, Melloni L, Zhang H, Tian X, Poeppel D. Cortical tracking of hierarchical

linguistic structures in connected speech. Nat Neurosci. 2016;19:15864.

16. Ding N, Pan X, Luo C, Su N, Zhang W, Zhang J. Attention is required for

knowledge-based sequential grouping: Insights from the integration of syllables into

words. J Neurosci. 2017; 38(5): 1178-1188.

17. Martin AE, Doumas LA. A mechanism for the cortical computation of hierarchical

linguistic structure. PLoS Biology 2017;15(3): e2000663.

18. Pallier C, Devauchelle AD, Dehaene S. Cortical representation of the constituent

structure of sentences. PNAS USA 2011;108: 2522–2527.

19. Pylkknen, Liina B, Bemis DK, Elorrieta EB. Building phrases in language

production: an Meg study of simple composition. Cognition 2014;133(2): 371–384.

20. Fruchter J, Linzen T, Westerlund M, Marantz A. Lexical preactivation in basic

linguistic phrases. J Cogn Neurosci. 2015;27(10): 1912–1935.

21. Pylkknen L. The neural basis of combinatory syntax and semantics. Science

2019;366(6461): 62–66.

22. Zaccarella E, Mayer L, Makuuchi M, Friederici AD. Building by syntax: The

neural basis of minimal linguistic structures. Cereb Cortex 2017;27(1): 411–421.

23. Nelson MJ, El Karoui I, Giber K, Yang X, Cohen L, Koopman H et al.

Neurophysiological dynamics of phrase-structure building during sentence

processing. PNAS USA 2017;114: E3669–E3678.

24. Andersen SK, Müller MM. Driving steady-state visual evoked potentials at

arbitrary frequencies using temporal interpolation of stimulus presentation. BMC

Neurosci. 2015;16: 95.

25. Macmillan NA, Creelman CD. Detection theory: A user's guide. 2nd ed. New

York: Lawrence Erlbaum Associates Publishers; 2005.

26. Kupietz, M. BCKH, Witt A. The German reference corpus Dereko: A primordial

sample for linguistic research. In: Calzolari N, Choukri K, Maegaard B, Mariani J,

Odijk J, Piperidis S, et al., editors. Proceedings of the 7th conference on international

language resources and evaluation (Irec 2010). Valletta, Malta: European language

resources association (Elra); 2010. pp. 1848-1854.

27. Zhou H, Melloni L, Poeppel D, Ding N. Interpretations of frequency domain

analyses of neural entrainment: Periodicity, fundamental frequency, and harmonics.

Front Hum Neurosci. 2016;10: 274.

28. Steiger JH. Tests for comparing elements of a correlation matrix. Psychological

Bulletin 1980;87: 245–251.

29. Rizzi L, Cinque G. Functional categories and syntactic theory. Ann Rev Ling.

2016;2(1): 139–163.

30. Frank SL, Yang J. Lexical representation explains cortical entrainment during

speech comprehension. PLoS ONE 2018;13(5): e0197304.

31. Weyerts H, Penke M, Münte TF, Heinze HJ, Clahsen H. Word order in sentence

processing: An experimental study of verb placement in German. J Psycholing Res.

2002;31(3): 21168.

32. Koster J. Word order in sentence processing: An experimental study of verb

placement in German. In: Abraham W, editor. Groninger arbeiten zur

germanistischen linguistik. Groningen, The Netherlands: University of Groningen;

1999. pp. 1-42.

33. Bieber D. Variation across speech and writing. Cambridge: Cambridge University

Press; 1988.

34. Greenberg S, Carvey H, Hitchcock L, Chang S. Temporal properties of

spontaneous speech - a syllable-centric perspective. J Phonetics 2003;31(3): 465–

485.

35. Meyer L, Henry ML, Gaston P, Schmuck N, Friederici AD. Linguistic bias

modulates interpretation of speech via neural delta-band oscillations. Cereb Cortex

2017;27(9): 4293–4302.

36. Park H, Ince RAA, Schyns PG, Thut G, Gross J. Frontal top-down signals

increase coupling of auditory low-frequency oscillations to continuous speech in

human listeners. Curr Biol. 2015;25: 1649–1653.

37. Keitel A, Robin AA Ince, Joachim Gross CK. Auditory cortical delta-entrainment

interacts with oscillatory power in multiple fronto-parietal networks. Neuroimage

2017;147: 32–42.

38. Rossion B. Understanding face perception by means of human

electrophysiology. Trends Cogn Sci. 2014;18(6): 310–318.

39. Delorme A, Makeig S. Eeglab: an open source toolbox for analysis of single-trial

eeg dynamics. J Neurosci Methods 2004;134: 9–21.

40. Bell A, Sejnowski T. An information-maximization approach to blind separation

and blind deconvolution. Neural Computation 1995;7: 1129–1159.

41. Chaumon, M. BD, Busch N. A practical guide to the selection of independent

components of the electroencephalogram for artifact correction. J Neurosci Methods

2015;250: 47–63.

42. Genovese, Christopher R. LN, Nichols T. Thrsholding of statistical maps in

functional neuroimaging using the false discovery rate. Neuroimage 2002;15: 870–

878.

43. Baayen, H PR, Gulikers L. The Celex lexical database [cd-rom]; 1995.

Philadelphia, PA: Linguistic data consortium, University of Pennsylvania.

 Figure legends

Figure 1, Neural response models: a. A syntactic description of a sample sentence

from condition 1, replicating in German the structure used by Ding et al. [15] in

English and Mandarin Chinese. b. According to the *sPhrase* model, linguistic brain

rhythms emerge by cycling through all nodes of the same syntactic level, regardless

of their distinct categorical status. A Fast Fourier Transform (FFT) analysis captures

the *sPhrase* model results. Notice that the model predicts also the absence of a 3 Hz

higher harmonic. c. The categorical model holds that the brain distinguishes between

NPs and VPs as separate abstract categories: their individual period equals that of

the sentence node, as phrases also repeat every second. The FFT analysis of brain

data should therefore highlight a 1 Hz peak of energy and its harmonics, reflecting

the contribution from both phrase and sentence levels of syntactic competence. d. A

sample sentence from condition 2 with the same number of words, but different

phrasal organization. e. *sPhrase* model results: The NP - with one word running at 4

Hz - would generate a 4 Hz rhythm and thus be indistinguishable from external word

rhythm, while the VP - with three words running at 4 Hz - would yield a distinct peak

of energy at 1.33 Hz in the resulting FFT model. f. A categorical analysis of condition

2 assumes that each phrase type cycles independently, thus effectively overlapping

in predictions with condition 1. The resulting neural response should not differ from

that of condition 1. See text for specific model predictions in each condition.

Figure 2, Syntactic variability across conditions: a. Condition 1 and 2 have four

monosyllabic words running at a constant speed of 4 Hz, with sentences in condition

1 composed of a two-word NP and a two-word VP - replicating Ding et al. [15] -,

while sentences in condition 2 are composed by a one-word NP followed by a three-

word VP. b. Conditions 3 and 4 have five monosyllabic words running at a constant

speed of 4 Hz: Sentences in condition 3 are composed of a three-word NP followed

by a two-word VP, while condition 4 presents the reverse pattern. c. Conditions 5

and 6 replicate the structure in conditions 3 and 4, only with disyllabic words running

at 2 Hz. For each condition, a syntactic tree of an exemplary sentence and the

corresponding waveform are provided. Red and green color codes identify even and

odd conditions from each pair (1-2, 3-4, 5-6, respectively), here and throughout the

text. See Supplemental Information for a complete list of sentence materials.

Figure 3, Behavioral accuracy: a. In all conditions, participants were able to

discriminate between regular and irregular trials. A significant difference between

conditions 1 and 4 suggests that discriminating irregular from regular trials in

condition 1, the original condition of Ding et al. [15], was more difficult than in

condition 4. b. Positive response bias measures for conditions 1, 4, 5 and 6 suggest

that participants adopted a conservative criterion when operating grammatical

judgments: They were more likely to respond that a given trial was regular, rather

than irregular.

Figure 4, Neural linguistic rhythms in regular trials. The left panel plots the FFT

results for regular trials across condition pairs: 1-2, 3-4, 5-6. Word rate is highlighted,

along with the positional distribution of identifiable *sPhrase* rhythms (vertical

hyphenated bars). No significant peaks were found for any of the hypothesized

sPhrase rhythms. Conversely, a significant harmonic structure was found in all

conditions: the right panel reports individual peak values for category rhythms, as

well as first and second harmonic process, plotted against floor noise estimates.

Significance is marked as follows: ** $\leq$ 0.01, * $\leq$ 0.05, ns = non-significant. Notice

that all probability values, here and in the text, are objectively reported, and deemed

significant or not within an FDR approach.

Figure 5, Neural linguistic rhythms in irregular trials: The left panel plots the FFT

results for all irregular trials in all condition pairs. A generalized suppression of

category rhythms and their harmonic structure is clear. Right panel: for each

condition pair, SNR estimates are plotted - with horizontal bar indicating a sample's

median – according to whether data were extracted from regular trials (REG), or

irregular trials (IRR). The significance of each comparison is marked as follows: ** $\leq$

0.01, * $\leq$ 0.05, ns = non significant (FDR-corrected).

Figure 6, Brain/behavior fit: a. The median power of internal harmonics extracted

from regular trials across conditions predicts grammatical judgments (black thick

line) better than power extracted from irregular trial harmonics (ochre thick line); b.

Entrainment to stimulus rate from regular trials is at ceiling, unmodulated, and thus

not predictive of task sensitivity (black thick line). Entrainment values extracted from

irregular trials (ochre thick line) significantly increase with task sensitivity. We take

this finding to suggest that attention to stimuli modulated the ability to detect

ungrammatical sentences within irregular trials. Thin, dotted lines indicate each

correlation's 95% confidence interval. Significance: *** $\leq$ 0.001, * $\leq$ 0.05, ns = non

significant.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6